


NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering all
Aspects of Natural Products Research



Volume 6. Issue 12. Pages 1799-1968. 2011
ISSN 1934-578X (printed); ISSN 1555-9475 (online)
www.naturalproduct.us

EDITOR-IN-CHIEF**DR. PAWAN K AGRAWAL**

Natural Product Inc.
7963, Anderson Park Lane,
Westerville, Ohio 43081, USA
agrawal@naturalproduct.us

EDITORS**PROFESSOR ALESSANDRA BRACA**

Dipartimento di Chimica Bioorganica e Biofarmacia,
Università di Pisa,
via Bonanno 33, 56126 Pisa, Italy
braca@farm.unipi.it

PROFESSOR DEAN GUO

State Key Laboratory of Natural and Biomimetic Drugs,
School of Pharmaceutical Sciences,
Peking University,
Beijing 100083, China
gda5958@163.com

PROFESSOR YOSHIHIRO MIMAKI

School of Pharmacy,
Tokyo University of Pharmacy and Life Sciences,
Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan
mimaki@ps.toyaku.ac.jp

PROFESSOR STEPHEN G. PYNE

Department of Chemistry
University of Wollongong
Wollongong, New South Wales, 2522, Australia
spyne@uow.edu.au

PROFESSOR MANFRED G. REINECKE

Department of Chemistry,
Texas Christian University,
Forts Worth, TX 76129, USA
m.reinecke@tcu.edu

PROFESSOR WILLIAM N. SETZER

Department of Chemistry
The University of Alabama in Huntsville
Huntsville, AL 35809, USA
wsetzer@chemistry.uah.edu

PROFESSOR YASUHIRO TEZUKA

Institute of Natural Medicine
Institute of Natural Medicine, University of Toyama,
2630-Sugitani, Toyama 930-0194, Japan
tezuka@inn.u-toyama.ac.jp

PROFESSOR DAVID E. THURSTON

Department of Pharmaceutical and Biological Chemistry,
The School of Pharmacy,
University of London, 29-39 Brunswick Square,
London WC1N 1AX, UK
david.thurston@pharmacy.ac.uk

HONORARY EDITOR**PROFESSOR GERALD BLUNDEN**

The School of Pharmacy & Biomedical Sciences,
University of Portsmouth,
Portsmouth, PO1 2DT U.K.
axuf64@dsl.pipex.com

ADVISORY BOARD

Prof. Berhanu M. Abegaz
Gaborone, Botswana

Prof. Viqar Uddin Ahmad
Karachi, Pakistan

Prof. Øyvind M. Andersen
Bergen, Norway

Prof. Giovanni Appendino
Novara, Italy

Prof. Yoshinori Asakawa
Tokushima, Japan

Prof. Lee Banting
Portsmouth, U.K.

Prof. Julie Banerji
Kolkata, India

Prof. Alejandro F. Barrero
Granada, Spain

Prof. Anna R. Bilia
Florence, Italy

Prof. Maurizio Bruno
Palermo, Italy

Prof. César A. N. Catalán
Tucumán, Argentina

Prof. Josep Coll
Barcelona, Spain

Prof. Geoffrey Cordell
Chicago, IL, USA

Prof. Cristina Gracia-Viguera
Murcia, Spain

Prof. Duvvuru Gunasekar
Tirupati, India

Prof. A.A. Leslie Gunatilaka
Tucson, AZ, USA

Prof. Kurt Hostettmann
Lausanne, Switzerland

Prof. Martin A. Iglesias Arteaga
Mexico, D. F., Mexico

Prof. Jerzy Jaroszewski
Copenhagen, Denmark

Prof. Leopold Jirovetz
Vienna, Austria

Prof. Karsten Krohn
Paderborn, Germany

Prof. Hartmut Laatsch
Gottingen, Germany

Prof. Marie Lacaille-Dubois
Dijon, France

Prof. Shoen-Sheng Lee
Taipei, Taiwan

Prof. Francisco Macias
Cadiz, Spain

Prof. Imre Mathe
Szeged, Hungary

Prof. Joseph Michael
Johannesburg, South Africa

Prof. Ermino Murano
Trieste, Italy

Prof. M. Soledade C. Pedras
Saskatoon, Canada

Prof. Luc Pieters
Antwerp, Belgium

Prof. Peter Proksch
Düsseldorf, Germany

Prof. Phila Raharivelomanana
Tahiti, French Polynesia

Prof. Monique Simmonds
Richmond, UK

Prof. Valentin Stonik
Vladivostok, Russia

Prof. Winston F. Tinto
Barbados, West Indies

Prof. Karen Valant-Vetschera
Vienna, Austria

Prof. Peter G. Waterman
Lismore, Australia

INFORMATION FOR AUTHORS

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site <http://www.naturalproduct.us>.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the Natural Product Inc. (NPI). A photocopy license is available from the NPI for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

To Subscribe: Natural Product Communications is a journal published monthly. 2011 subscription price: US\$1,995 (Print, ISSN# 1934-578X); US\$1,995 (Web edition, ISSN# 1555-9475); US\$2,495 (Print + single site online); US\$595 (Personal online). Orders should be addressed to Subscription Department, Natural Product Communications, Natural Product Inc., 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

Natural Product Communications

2011

Volume 6, Number 12

Contents

<u>Original Paper</u>	<u>Page</u>
Sibiralactone: A New Monoterpene from <i>Sibiraea angustata</i> Guangbo Xie, Xianlong Wang, Tibor Kurtán, Attila Mándi and Tianzhi Wang	1799
Bioconversion of Proposed Precursors into Theobroxide and Related Compounds Peng Li, Kosaku Takahashi, Ahmed Elkhateeb, Hideyuki Matsuura, Teruhiko Yoshihara and Kensuke Nabeta	1801
Microbial Hydroxylation of <i>S</i>-(-)-Perillyl Alcohol by <i>Fusarium heterosporium</i> Ismail Kiran	1805
A Phytochemical Investigation of <i>Zanthoxylum setulosum</i> Tameka M. Walker, Bernhard Vogler, Debra M. Moriarity, William A. Haber and William N. Setzer	1807
Cytotoxic Cembranoids from the Red Sea Soft Coral <i>Sarcophyton glaucum</i> Mohamed-Elamir F. Hegazy, Ahmed A. El-Beih, Alaa Y. Moustafa, Abdelhamed A. Hamdy, Montaser A. Alhammady, Rehab M. Selim, Mohamed Abdel-Rehim and Paul W. Paré	1809
C-Lactam Derivatives of Oleanolic Acid. The synthesis of C-lactam by Beckmann rearrangement of C-oxime Barbara Bednarczyk – Cwynar	1813
Analysis of Native Carotenoid Composition of Sweet Bell Peppers by Serially Coupled C₃₀ Columns Daniele Giuffrida, Paola Dugo, Giacomo Dugo, Germana Torre and Luigi Mondello	1817
New Antifungal Cholestane and Aldehyde Derivatives from the Red Alga <i>Laurencia papillosa</i> Walied M. Alarif, Sultan S. Al-Lihaibi, Ahmed Abdel-Lateff and Seif-Eldin N. Ayyad	1821
Steroidal Saponins from the Fruits of <i>Cestrum ruizteranianum</i> Elier Galarraga M., Anne-Claire Mitaine-Offer, Juan Manuel Amaro-Luis, Tomofumi Miyamoto, Chiaki Tanaka, Laurent Pouységu, Stéphane Quideau, Luis B. Rojas and Marie-Aleth Lacaille-Dubois	1825
Isolation and Cholinesterase Activity of Amaryllidaceae Alkaloids from <i>Nerine bowdenii</i> Lucie Cahliková, Stanislav Zavadil, Kateřina Macáková, Irena Valterová, Andrea Kulhánková, Anna Hošťálková, Jiří Kuneš and Lubomír Opletal	1827
HPLC Determination of Majdine in <i>Vinca herbacea</i> Natia Gagua, Beatrice Baghdikian, Fathi Mabrouki, Riad Elias, Valentina Vachnadze, Aliosha Bakuridze and Evelyne Ollivier	1831
Pyridine Metabolism and Trigonelline Synthesis in Leaves of the Mangrove Legume trees <i>Derris indica</i> (<i>Millettia pinnata</i>) and <i>Caesalpinia crista</i> Yuling Yin, Hamako Sasamoto and Hiroshi Ashihara	1835
Anti-adipogenic Activity of <i>Cordyceps militaris</i> in 3T3-L1 Cells Qing Liu, In Pyo Hong, Mi-Jeong Ahn, Hwan-Soo Yoo, Sang-Bae Han, Bang Yeon Hwang and Mi Kyeong Lee	1839
Two New Cyclopeptides and One New Nonenolide from <i>Xylaria</i> sp. 101 Yao-Yao Li, Zhi-Yu Hu, and Yue-Mao Shen	1843
A Novel Flavonoid and Furoquinoline Alkaloids from <i>Vepris glomerata</i> and their Antioxidant Activity Joyce J. Kiplimo, Md. Shahidul Islam and Neil A. Koorbanally	1847
Flavonoid Constituents and Free Radical Scavenging Activity of <i>Alchemilla mollis</i> Antoaneta Trendafilova, Milka Todorova, Milena Nikolova, Anna Gavrilova and Antonina Vitkova	1851
Ultrasound-assisted Extraction of Total Phenols and Flavonoids from Dry Tobacco (<i>Nicotiana tabacum</i>) Leaves Ivana T. Karabegović, Vlada B. Veljković and Miodrag L. Lazić	1855
Characterization of Polyphenolic Compounds in Unripe Chinotto (<i>Citrus myrtifolia</i>) Fruit by HPLC/PDA/ESI/MS-MS Monica Scordino, Leonardo Sabatino, Adalgisa Belligno and Giacomo Gagliano	1857
Bioactive Compounds, RP-HPLC Analysis of Phenolics, and Antioxidant Activity of Some Portuguese Shrub Species Extracts Ângelo Luís, Fernanda Domingues and Ana Paula Duarte	1863
HPLC/PDA/ESI-MS Evaluation of Saffron (<i>Crocus sativus</i> L.) Adulteration Leonardo Sabatino, Monica Scordino, Maria Gargano, Adalgisa Belligno, Pasqualino Traulo and Giacomo Gagliano	1873

Continued Overleaf

Ferric Reducing, Antiradical and β-Carotene Bleaching Activities of Nicotinic Acid and Picolinic Acid Bioconjugates of Curcumin	
Archana Pandey, Kanti Bhooshan Pandey, Ravindra Kumar Gupta and Syed Ibrahim Rizvi	1877
Antiviral Activities of Diarylheptanoids Isolated from <i>Alpinia officinarum</i> against Respiratory Syncytial Virus, Poliovirus, Measles Virus, and Herpes Simplex Virus Type 1 <i>in vitro</i>	
Katsuhiko Konno, Rie Sawamura, Yi Sun, Ken Yasukawa, Tomomi Shimizu, Wataru Watanabe, Masahiko Kato, Ryuichi Yamamoto and Masahiko Kurokawa	1881
Isolation of C-glycosyl Xanthenes from <i>Coffea pseudozanguebariae</i> and Their Location	
Pascale Talamond, Geneviève Conejero, Jean-Luc Verdeil and Jean-Luc Poëssel	1885
Antifungal Activity and Isomerization of Octadecyl <i>p</i>-coumarates from <i>Ipomoea carnea</i> subsp. <i>fistulosa</i>	
Eugene Sebastian J. Nidiry, Girija Ganeshan and Ankanahalli N. Lokesh	1889
New Glucose Esters from the Fresh Leaves of <i>Jacaranda mimosaeifolia</i>	
Christianah A. Elusiyan and Tiwalade A. Olugbade	1893
Shamiminol: A New Aromatic Glycoside from the Stem Bark of <i>Bombax ceiba</i>	
Shaheen Faizi, Sadia Zikr-Ur-Rehman and Muhammad Ali Versiani	1897
Two New Phenolic Glycosides from <i>Viburnum plicatum</i> var. <i>plicatum</i> f. <i>plicatum</i>	
Saki Katagiri, Yoshiki Watanabe, Yasunori Yaoita, Masao Kikuchi and Koichi Machida	1901
Antimicrobial Chemical Constituents from the Endophytic Fungus <i>Phomopsis</i> sp. from <i>Notobasis syriaca</i>	
Hidayat Hussain, Michel Kenne Tchimine, Ishtiaq Ahmed, Kathrin Meier, Michael Steinert, Siegfried Draeger, Barbara Schulz and Karsten Krohn	1905
Phomosines H–J, Novel Highly Substituted Biaryl Ethers, Isolated from the Endophytic Fungus <i>Phomopsis</i> sp. from <i>Ligustrum vulgare</i>	
Karsten Krohn, Umar Farooq, Hidayat Hussain, Ishtiaq Ahmed, Joachim Rheinheimer, Siegfried Draeger, Barbara Schulz and Teunis van Ree	1907
Isolation and Characterization of a new Benzofuran from the Fungus <i>Alternaria</i> sp. (HS-3) Associated with a Sea Cucumber	
Xuekui Xia, Jun Qi, Fang Wei, Airong Jia, Wenpeng Yuan, Xiumei Meng, Miansong Zhang, Changheng Liu and Changyun Wang	1913
Potent Toxic Macrocytic Trichothecenes from the Marine-Derived Fungus <i>Myrothecium verrucaria</i> Hmp-F73	
Li Zhao, Li Liu, Nan Wang, Shu-Jin Wang, Jing-Chun Hu and Jin-Ming Gao	1915
Synthesis and Bioactivity of Novel Coumarin Derivatives	
Ai-Ying Guan, Chang-Ling Liu, Miao Li, Zhi-Nian Li, Ming-Xing Zhang and Hong Zhang	1917
Kinase Inhibitory, Haemolytic and Cytotoxic Activity of Three Deep-water Sponges from North Western Australia and their Fatty Acid Composition	
Ana Zivanovic, Natalie J. Pastro, Jane Fromont, Murray Thomson and Danielle Skropeta	1921
Antimicrobial and Cytotoxic Effects of Mexican Medicinal Plants	
Maria del Rosario Jacobo-Salcedo, Angel Josabad Alonso-Castro, Luis A. Salazar-Olivo, Candy Carranza-Alvarez, Luis Ángel González-Espíndola, Fabiola Domínguez, Sandra Patricia Maciel-Torres, Concepción García-Lujan, Marisela del Rocio González-Martínez, Maricela Gómez-Sánchez, Eduardo Estrada-Castillón, Rocio Zapata-Bustos, Pedro Medellín-Milán and Alejandro García-Carrancá	1925
Chemometrics Evaluation of the Herbal Drug <i>Andrographis paniculata</i>	
Shiv Narayan Sharma, Zenu Jha and D. K. Sharma	1929
<i>Garcinia cambogia</i> Leaf and Seawater for Tannase Production by Marine <i>Aspergillus awamori</i> BTMFW032 under Slurry State Fermentation	
Beena P. S, Soorej M. Basheer, Sarita G. Bhat and Chandrasekaran M	1933
Gas Chromatographic Quantitative Analysis of Methanol in Wine: Operative Conditions, Optimization and Calibration Model Choice	
Rosario Caruso, Grazia Laura Gambino, Monica Scordino, Leonardo Sabatino, Pasqualino Traulo and Giacomo Gagliano	1939
Composition and Biological Potential of Essential Oil from <i>Thelechitonina trilobata</i> Growing in South Africa	
Jamie Peebles, Ephraim Gwebu, Opeoluwa Oyediji, Sarah Nanyonga, Nokuthula Kunene, David Jackson, William Setzer and Adebola Oyediji	1945
Chemical Composition and Antibacterial Activity of Essential oil from <i>Salvia mukerjeei</i>	
Lalit Mohan, Anuradha Negi, Anand B. Melkani and Vasu Dev	1949
<u>Review/Account</u>	
Revealing Indigenous Indonesian Traditional Medicine: Anti-infective Agents	
Ari S. Nugraha and Paul A. Keller	1953

Manuscripts in Press

1967

Cummulative Index

Contents

Author Index

Keywords Index

i-xx

1-6

i-viii

LIST OF AUTHORS

Abdel-Lateff, A 1821	González-Martínez, MR 1925	Machida, K 1901	Skropeta, D 1921
Abdel-Rehim, M 1809	Guan, AY 1917	Maciel-Torres, SP 1925	Steinert, M 1907
Ahmed, I 1905	Gupta, RK 1877	Mándi, A 1799	Sun, Y 1881
Ahn, MJ 1839	Gwebu, E 1945	Matsuura, H 1801	Takahashi, K 1801
Alarif, WM 1821	Haber, WA 1807	Medellin-Milán, P 1925	Talamond, P 1885
Alhammady, MA 1809	Hamdy, AA 1809	Meier, K 1907	Tanaka, C 1825
Al-Lihaibi, SS 1821	Han, SB 1839	Melkani, AB 1949	Tchimene, MK 1905
Alonso-Castro, AJ 1925	Hegazy, MF 1809	Meng, X 1913	Thomson, M 1921
Amaro-Luis, JM 1825	Hong, IP 1839	Mitaine-Offier, A 1825	Todorova, M 1851
Ashihara, H 1835	Hošťáková, A 1827	Miyamoto, T 1825	Torre, G 1817
Ayyad, SN 1821	Hu, JC 1915	Mohan, L 1949	Traulo, P 1873
Baghdikian, B 1831	Hu, ZY 1843	Mondello, L 1817	Traulo, P 1939
Bakuridze, A 1831	Hussain, H 1905	Moriarity, DM 1807	Trendafilova, A 1851
Basheer, SM 1933	Hwang, BY 1839	Moustafa, AY 1809	
Bednarczyk – Cwynar, B 1813	Islam, MS 1847	Nabeta, K 1801	Vachnadze, V 1831
Beena PS 1933	Jackson, D 1945	Nanyonga, S 1945	Valterová, I 1827
Belligno, A 1857, 1873	Jacobo-Salcedo, MR 1925	Negi, A 1949	Veljković, VB 1855
Bhat, SG 1933	Jha, Z 1929	Nidiry, ESJ 1889	Verdeil, JL 1885
Cahliková, L 1827	Jia, A 1913	Nikolova, M 1851	Versiani, MA 1897
Carranza-Alvarez, C 1925	Karabegović, IT 1855	Nugraha, AS 1953	Vitkova, A 1851
Caruso, R 1939	Katagiri, S 1901	Ollivier, E 1831	Vogler, B 1807
Chandraksekaran M 1933	Kato, M 1881	Olugbade, TA 1893	Walker, TM 1807
Conejero, G 1885	Keller, PAQ 1953	Opletal, L 1827	Wang, C 1913
Dev, V 1949	Kikuchi, M 1901	Oyedéji, A 1945	Wang, N 1915
Domingues, F 1863, 1925	Kiplimo, JJ 1847	Oyedéji, O 1945	Wang, SJ 1915
Draeger, S 1907	Kiran, I 1805	Pandey, A 1877	Wang, T 1799
Duarte, AP 1863	Konno, K 1881	Pandey, KB 1877	Wang, X 1799
Dugo, G 1817	Koorbanally, NA 1847	Paré, PW 1809	Watanabe, W 1881
Dugo, P 1817	Krohn, K 1907	Pastro, NJ 1921	Watanabe, Y 1901
El-Beih, AA 1809	Kulhánková, A 1827	Peebles, J 1945	Wei, F 1913
Elias, R 1831	Kunene, N 1945	Poëssel, JL 1885	Xia, X 1913
Elier Galarraga M., E 1825	Kuneš, J 1827	Pouységu, L 1825	Xie, G 1799
Elkhateeb, A 1801	Kurokawa, M 1881	Qi, J 1913	Yamamoto, R 1881
Elusiyan, CA 1893	Kurtán, T 1799	Quideau, S 1825	Yaoita, Y 1901
Estrada-Castillón, E 1925	Lacaille-Dubois, M 1825	Ree, Tv 1907	Yasukawa, K 1881
Faizi, S 1897	Lazić, ML 1855	Rizvi, SI 1877	Yin, Y 1835
Fromont, J 1921	Lee, MK 1839	Rojas, LB 1825	Yoo, HS 1839
Gagliano, G 1857, 1873, 1939	Li, M 1917	Sabatino, L 1857, 1873, 1939	Yoshihara, T 1801
Gagua, N 1831	Li, P 1801	Salazar-Olivo, LA 1925	Yuan, W 1913
Gambino, GL 1939	Li, YY 1843	Sasamoto, H 1835	Zapata-Bustos, R 1925
Ganeshan, G 1889	Li, ZN 1917	Sawamura, R 1881	Zavadil, S 1827
Gao, JM 1915	Liu, C 1913	Schulz, B 1907	Zhang, HJ 1917
García-Carrancá, A 1925	Liu, CL 1917	Scordino, M 1857, 1939, 1973	Zhang, M 1913
García-Lujan, C 1925	Liu, L 1915	Selimi, RM 1809	Zhang, MX 1917
Gargano, M 1873	Liu, Q 1839	Setzer, W 1945	Zhao, L 1915
Gavrilova, A 1851	Lokesha, AN 1899	Setzer, WN 1807	Zikr-Ur-Rehman, S 1897
Giuffrida, D 1817	Luis, Á 1863	Sharma, DK 1929	Zivanovic, A 1921
Gómez-Sánchez, M 1925	Mabrouki, F 1831	Sharma, SN 1929	
González-Espíndola, LA 1925	Macáková, K 1827	Shen, YM 1843	
		Shimizu, T 1881	

Sibiralactone: A New Monoterpene from *Sibiraea angustata*Guangbo Xie^{a,*}, Xianlong Wang^a, Tibor Kurtán^b, Attila Mándi^b and Tianzhi Wang^c^aSchool of Life Science & Technology, University of Electronic Science & Technology of China, Chengdu 610054, People's Republic of China^bDepartment of Organic Chemistry, University of Debrecen, PO Box 20, H-4010 Debrecen, Hungary^cWest China School of Pharmacy, Sichuan University, Chengdu 610041, People's Republic of China

gbxie@uestc.edu.cn

Received: August 23rd, 2011; Accepted: October 26th, 2011

Sibiralactone (**1**), a new monoterpene derivative, was isolated from the leaves of *Sibiraea angustata*. The structure was determined by the analysis of its NMR data and the absolute configuration was established by TDDFT ECD calculation of the solution conformers.

Keywords: *Sibiraea angustata*, sibiralactone, monoterpene, TDDFT ECD calculation.

Sibiraea angustata (Rehd.) Hand.-Mazz. is indigenous of the bush flora and gravel areas found in Qinghai, Gansu, Sichuan and Yunnan provinces of China at an altitude of 3000–4000 m [1]. When it is fed to domestic animals, loss of weight was observed [2a]. Previous phytochemical investigations of the plant revealed the presence of several triterpenoids, phenolic acids, fatty alcohols [2b–2e] and a monoterpene lactone, 1-*O*- β -D-glucopyranosylgeraniol-10,5-olide exhibiting lipid metabolism-ameliorating and hypoglycemic activity [3]. Herein we report the structure elucidation of sibiralactone (**1**), the second monoterpene lactone from *S. angustata* leaves.

Sibiralactone (**1**) was determined as C₁₀H₁₆O₄ on the basis of a HRESI-MS peak of [M + Na]⁺ at 223.0942 (Calcd. 223.0946), indicating three degrees of unsaturation. The IR spectrum showed absorption bands characteristic of hydroxyl groups (3307 cm⁻¹) and a γ -lactone moiety (1780 cm⁻¹). The ¹H NMR spectrum showed proton signals for two isolated methyl groups [δ _H 1.80 (3H, s), 1.81 (3H, s)], two oxymethine groups [δ _H 3.88 (1H, t, *J* = 8.4 Hz), 4.85 (1H, t, *J* = 8.4 Hz)], one methine group [2.60 (1H, dt, *J* = 2.8, 9.6 Hz)], one olefinic H-atom [δ _H 5.21 (1H, br. d, *J* = 9.2 Hz)], one oxymethylene group [δ _H 3.72 (1H, dt, *J* = 2.4, 10.8 Hz), 4.00 (1H, dt, *J* = 4.0, 10.8 Hz)] and one methylene group [δ _H 1.73 (1H, m), 2.25 (1H, dq, *J* = 3.2, 6.4, 14.8 Hz)]. The proton spin system of (O)-CH₂-CH₂-CH-CH(O)-CH(O)-CH=C(CH₃)₂ could be deduced by an ¹H-¹H COSY measurement. The ¹³C NMR spectroscopic data, including the DEPT spectrum, showed two methyls (δ _C 18.6 and 25.9), two oxymethines (δ _C 78.6 and 79.7), one methine (δ _C 49.0), one oxymethylene (δ _C 61.8), one methylene (δ _C 31.2), one olefinic methine (δ _C 120.1), one quaternary olefinic carbon (δ _C 143.0), and a carbonyl (δ _C 176.0). The ester carbonyl at δ _C 176.0 showed HMBC correlation with the C-7 methylene




Figure 1: ¹H-¹H COSY, key HMBC and selected NOE correlations of **1**.




Figure 2: Experimental and B3LYP/TZVP ECD spectra of (4*S*, 5*S*, 6*R*)-**1** obtained as the Boltzmann-weighted average of the computed four solution conformers.

protons at δ _H 1.73 and 2.25, and the oxymethine at δ _H 4.85, which together with other correlations confirmed the γ -lactone moiety and the planar structure of **1** (Figure 1). The all *cis* relative configuration was deduced on the ground of NOE effects between H-4 (δ _H 4.85), H-5 (δ _H 3.88) and H-6 (δ _H 2.60) (Figure 1), implying (4*R**, 5*R**, 6*S**) relative configuration. The all *cis* relative configuration also corroborated well the observed ³*J*_{H,H} coupling constants of 4-H (*J* = 8.4 Hz), 5-H (*J* = 8.4 Hz), and 6-H (*J* = 2.8, 9.6 Hz).




Figure 3: DFT optimized geometries of the four lowest-energy conformers of (4S, 5S, 6R)-1.

For the determination of absolute configuration, the electronic circular dichroism (ECD) spectra were calculated with three functionals (B3LYP, BH&HLYP, PBE0) for the computed four lowest-energy conformers (with populations larger than 5%) of (4S, 5S, 6R)-1 and the Boltzmann-weighted ECD spectrum was compared with the experimental ECD curve. Since the calculated and experimental ECD curves were mirror images (Figure 2), the absolute configuration of sibiralactone (**1**) was determined as (+)-(4R, 5R, 6S).

Experimental

General: Optical rotation, Perkin-Elmer Model 341 polarimeter; IR, Nicolet NEXUS 670 FT-IR spectrometer; ECD, Applied Photophysics Ltd. Chirascan spectropolarimeter; HRESI-MS, Waters Q-TOF Premier; NMR (1D and 2D) on a Varian Unity 400/54 instrument.

Plant material: The leaves of *Sibiraea angustata* were collected at Songpan County, Sichuan Province in August, 2008. The plant was identified by Prof. Tiznzh Wang. A voucher specimen (No. SA0808) was deposited at the Herbarium of the Sichuan University, Chengdu, China.

Extraction and isolation: Powdered leaves of *S. angustata* (5 kg) were extracted with 80% aq. ethanol under reflux. The extracts were evaporated *in vacuo*, then partitioned between H₂O and EtOAc. The EtOAc extract (185 g) was chromatographed over a silica gel column (2000 g, 100-200 mesh), eluted with a gradient solvent system [CHCl₃-MeOH (90:1-2:1)] to give 12 fractions (Fr.1-12). Fr. 4 (2.9 g) was chromatographed over a silica gel column (200-300 mesh, 60 g)

eluted with solvent systems of light petroleum-acetone (5:1) and cyclohexane-EtOAc (4:1) to afford compound **1** (24 mg).

Sibiralactone (1)

colorless gum

[α]_D: +32.7 (*c* 1.15, CHCl₃).

CD (MeOH, λ [nm] ($\Delta\epsilon$), *c* = 0.028): 217 (-1.41), 196 (9.12).

IR ν_{\max} (ethanol): 3307, 2848, 1780, 1454, 1378, 1045, 879 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): 1.73 (1H, m, H-7a), 1.80 (3H, s, H-10), 1.81 (3H, s, H-1), 2.25 (1H, dq, *J* = 14.8, 6.4, 3.2 Hz, H-7b), 2.60 (1H, dt, *J* = 9.6, 2.8 Hz, H-6), 3.72 (1H, dt, *J* = 10.8, 2.4 Hz, H-8a), 3.88 (1H, t, *J* = 8.4 Hz, H-5), 4.00 (1H, dt, *J* = 10.8, 4.0 Hz, H-8b), 4.85 (1H, t, *J* = 8.4 Hz, H-4), 5.21 (1H, br.d, *J* = 9.2 Hz, H-3).

¹³C NMR (100 MHz CDCl₃): 18.6 (CH₃, C-10), 25.9 (CH₃, C-1), 31.2 (CH₂, C-7), 49.0 (CH, C-6), 61.8 (CH₂, C-8), 78.6 (CH, C-5), 79.7 (CH, C-4), 120.1 (CH, C-3), 143.0 (C, C-2), 176.0 (C, C-9).

HRESI-MS: *m/z* 223.0942 [M + Na⁺] (calcd. 223.0946 for C₁₀H₁₆N₄NaO₄).

Computational section: Conformational searches were carried out by means of the MacroModel 9.7.211 [4] software using Merck Molecular Force Field (MMFF) with implicit solvent model for chloroform. Geometry reoptimizations at B3LYP/6-31 G(d) level of theory followed by TDDFT calculations using various functionals (B3LYP, BH&HLYP, PBE0) and TZVP basis set were performed by the Gaussian 03 [5] package. Boltzmann distributions were estimated from the ZPVE corrected B3LYP/6-31 G(d) energies. ECD spectra were generated as the sum of Gaussians [6] with 3000 cm⁻¹ half-height width (corresponding to ca. 12 nm at 200 nm), using dipole-velocity computed rotational strengths for conformers above 5%. The MOLEKEL [7] software package was used for visualization of the results.

Acknowledgments - The authors are grateful to Shepo Shi (Institute of Biological Chemistry, Washington State University) for discussion on the structure elucidation. This work was supported financially by the Fundamental Research Funds for the Central Universities (ZYGX2010J102). T. K. thanks the National Office for Research and Technology (NKTH, K-68429), TÁMOP 4.2.1/B-09/1/KONV-2010-0007 and János Bolyai Foundation.

References

- [1] Yu DJ, Gu CZ. (1974) *Flora Republicae Popularis Sinicae*. Volume 36. Science Publishing House, Beijing. 69-70.
- [2] (a) Wang XL, Guo JM, Yang BX. (1995) Studies on narrowleaf *Sibiraea* (*Sibiraea angustata*) in regulating lipid metabolism. *Chinese Traditional and Herbal Drugs*, 26, 473-475, 504; (b) Zhang CZ, Feng SL, Li C, Zhao CX, Shang XY, Ma HP. (1992) Triterpenoid constituents from liucha (*Sibiraea angustata*). *China Journal of Chinese Materia Medica*, 17, 360, 373; (c) Zhang CZ, Li C, Feng SL, Zhao CL, Wang WH. (1993) Isolation and structure of sibirate from *Sibiraea angustata*. *Acta Pharmaceutica Sinica*, 28, 798-800; (d) Liu JX, Wu SJ, Wei XY, Yang RZ. (1999) Studies on the chemical constituents of *Sibiraea angustata*. *Journal of Tropical and Subtropical Botany*, 7, 248-251; (e) Tao TT, Sei Y, Wang TZ, Bai C, Chang YB. (2006) Chemical constituents of *Sibiraea angustata*. *Chinese Journal Natural Medicines*, 4, 257-259.
- [3] Sei Y, Kano Y, Takabori R. (2007) 1-O-beta-D-glucopyranosyl-raniol-10,5-olide and use thereof, US 2007/0059339.
- [4] MacroModel, Schrödinger LLC, 2009. <http://www.schrodinger.com/Products/macromodel.html>
- [5] Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR, Montgomery Jr JA, Vreven T, Kudin KN, Burant JC, et al. Gaussian 03, Revision C.02. 2004, Gaussian Inc., Wallingford CT.
- [6] Stephens PJ, Harada N. (2010) ECD cotton effect approximated by the Gaussian curve and other methods. *Chirality*, 22, 229-233.
- [7] Varetto U. MOLEKEL 5.4., 2009, Swiss National Supercomputing Centre: Manno, Switzerland.

Bioconversion of Proposed Precursors into Theobroxide and Related Compounds

Peng Li^{a,*}, Kosaku Takahashi^a, Ahmed Elkhateeb^{a,†}, Hideyuki Matsuura^a, Teruhiko Yoshihara^a and Kensuke Nabeta^a

^aLaboratory of Bioorganic Chemistry, Division of Applied Bioscience, Graduate School of Agriculture, Hokkaido University, Kita 9, Nishi 9, Kita ku, Sapporo 060-8589, Japan

[†]Permanent address: Phytochemistry and Plant Systematic Department, National Research Centre, Dokki, Giza, Egypt

Current address of corresponding author: Central Research Institute of Electric Power Industry, Nagasaki, Yokosuka 240-0196 Japan

yolipeng@criepi.denken.or.jp

Received: June 14th, 2011; Accepted: September 2nd, 2011

We have previously reported a tetraketide origin for theobroxide and its related compound. In the present study, bioconversion of natural and deuterium-labeled precursors of this proposed biosynthetic pathway by *Lasiodiplodia theobromae* was investigated. Theobroxide was quantified after bioconversion from each proposed precursor. The transformation of the isotopically labeled precursor to products was tracked by ²H NMR measurement.

Keywords: *Lasiodiplodia theobromae*, bioconversion, fungal biosynthesis, theobroxide, ²H NMR.

Fungi are lower eukaryotic microbes having important relationships with human beings not only as plant and animal pathogens, but also as major producers of enzymes, amino acids, and biologically active secondary metabolites. *Lasiodiplodia theobromae* (synonym *Botryodiplodia theobromae*) is a common pathogenic fungus that has been found throughout the tropics and the subtropics [1]. The culture filtrate of *L. theobromae* showed significant plant growth inhibitory activity [2]. For several decades, a lot of attention has been paid to this fungus. A variety of bioactive components have been isolated from it and been characterized, such as jasmonic acid (JA), a plant hormone widely distributed in plants [3] and micro-organisms [2]; mellein, a metabolite of *Aspergillus melleus* [4] and *A. ochraceus* [5], as a hair pencil component of the male oriental fruit moth *Grapholitha molesta* [6]; and theobroxide, a potato-tuber and flower bud inducing substance (1*S*,2*R*,5*S*,6*R*)-3-methyl-7-oxa-bicyclohept-3-en-2,5-diol (1), with its related compound (1*S*,4*R*,5*S*,6*R*)-7,9-dioxa-3-methyl-8-oxobicyclo[4.3.0]-2-nonene-4,5-diol (2), both showing inducing activity for different physiological phenomena. Research on some plant growth regulators from *L. theobromae* and their biosynthesis, translocation, functions, and modes of action has also been intensively undertaken by the authors' group [7–12].

In our previous study, we reported a tetraketide origin for biosynthesis of 1 and 2 by using ¹³C and ²H labeled




Figure 1: The proposed biosynthetic pathway of theobroxide 1 and compound 2.

acetates [7]. In the proposed biosynthetic pathway (Figure 1), we supposed that four acetyl-CoA are linked 'head-to-tail' to generate a triketide-CoA, and then one of the resulting keto groups is reduced to a hydroxyl group. Generation of a carbanion at the β -keto residue would allow an aldol condensation to form a six-membered carbocycle. Then a sequence of plausible dehydration and enolisation reactions would produce the 6-methylsalicylic acid (3, 6-MSA) as a natural precursor [13,14]. Decarboxylation of compound 3 gives 3-methylphenol (4), which is *para*-hydroxylated to 2-methylbenzene-1,4-diol (5). Precursor 5 is then oxidized to 2-methyl-[1,4]benzoquinone (6). Finally, theobroxide (1) is

produced through epoxidization and reduction of both hydroxyl groups, and is converted to its carbonyldioxy derivative **2** by an unidentified mechanism.

As a continuing study, we performed bioconversion experiments to confirm this proposed biosynthetic pathway by incorporating both natural and isotopic proposed precursors into these two target compounds.

In this study, the appropriate conditions and process for feeding of four proposed precursors to *L. theobromae* were investigated first. The mycelia grew well when the fungal culture was fed with these chemicals at a concentration of under 10^{-4} M. However, they did not grow when fed with compound **6** at 10^{-3} M, implying that a higher concentration of **6** may kill the fungus. No obvious difference in mycelial growth was observed when fed with the other three compounds at 10^{-3} M compared with the control group, which was fed only with 0.5 mL of DMSO (average weight: 3.98–4.35 g/200 mL culture). The average mycelial weight of the positive control group was 4.57 g/200 mL culture. The results indicated that these four compounds did not significantly suppress growth of *L. theobromae* OCS 71 at a concentration of 10^{-4} M. TLC monitoring also showed that theobroxide can be produced well in the presence of 10^{-4} M of **6**. Therefore, feeding experiments with *L. theobromae* using the proposed precursors **3–6** were carried out at a concentration of 10^{-4} M.

The results are presented in Figure 2. As can be seen, feeding of compounds **3** or **4** increased the amount of theobroxide in the fungal culture by two times compared with the control. In the case of fungal culture fed with compounds **5** or **6**, theobroxide was produced in almost three times higher amount compared with the control group. The higher content of theobroxide might be because compounds **5** and **6** are closer to theobroxide in the biosynthetic pathway. The above results provide evidence for compounds **3**, **4**, **5**, and **6** as intermediates in the biosynthetic pathway of theobroxide.




Figure 2: Content of theobroxide after bioconversion from proposed precursors. Values represent the means of three independent experiments with \pm SE which indicate significant differences ($p < 0.05$) with respect to the control.

The biosynthesis of theobroxide **1** and compound **2** from the proposed precursor of compound **5** in *L. theobromae*

Table 1: Incorporation of deuterium-labelled **5a** into compounds **1** and **2**.

Position	Compound 1		Compound 2	
	$^1\text{H}^a$	$^2\text{H}^a$	$^1\text{H}^b$	$^2\text{H}^b$
1	3.31 (1H, br s)	-	5.10 (1H, m)	-
2	4.20 (1H, br s)	-	5.56 (1H, m)	-
3	-	-	-	-
4	5.46 (1H, br d, $J = 5.4$ Hz)	-	3.88 (1H, br d, $J = 8.1$ Hz)	-
5	4.41 (1H, br s)	-	3.55 (1H, br t, $J = 8.1$ Hz)	-
6	3.25 (1H, br s)	-	4.59 (1H, br t, $J = 8.1$ Hz)	-
8	1.79 (3H, s)	1.79 (3D, s)	-	-
10	-	-	1.84 (3H, s)	1.84 (3D, s)

^aMeasured in CDCl_3 for ^1H NMR and in CHCl_3 for ^2H NMR.

^bMeasured in CD_3OD for ^1H NMR and in CH_3OH for ^2H NMR.

was further examined by means of ^2H NMR based isotopic tracing. As shown in Table 1, ^2H NMR spectroscopic data of the resulting compounds **1** and **2** revealed that the chemical shifts of the incorporated deuterium were δ 1.79 for **1** and δ 1.84 for **2**, respectively. These chemical shifts are consistent with those of the methyl proton of natural theobroxide **1** and compound **2**. These results indicated that retention of deuterium of the methyl group occurred during bioconversion from **5a** to **1** and **2** in *L. theobromae*. Accordingly, compound **5** was further proved to be an intermediate in the biosynthetic pathway of theobroxide.

In conclusion, in this study, some evidence was provided for the previously proposed biosynthetic pathway of theobroxide (**1**) and compound **2** as depicted in Figure 1. For future work, the biotransformation mechanism of theobroxide to its carbonyldioxy derivative **2** should be clarified experimentally.

Experimental

General experimental procedures: Spores of *L. theobromae* were maintained on agar (1.5%) in a 2%-potato D-glucose medium (Nissui Seiyaku) at 25°C and transferred at intervals of 6 months. All chemicals were purchased from either Wako or Aldrich Chemical Company. Compounds **3** and **5a** were prepared by published methods [15, 16]. Preparative and analytical TLC were performed on Merck Kieselgel 60F₂₅₄ precoated glass plates (0.5 mm and 0.25 mm, respectively). Compounds were visualized by either exposure to UV light or by staining with 5% molybdophosphoric acid *n*-hydrate in EtOH followed by heating on a hot plate. Open column chromatography was performed with silica gel (60N spherical neutral 63–210 μm , Kanto Chemical). ^1H NMR (270 MHz and 500 MHz) and ^2H NMR (76.5 MHz) spectra were recorded on a Bruker AMX-500 FT-NMR.

Determination of feeding conditions: Two different concentrations (10^{-3} M and 10^{-4} M) of each proposed precursor were tested for fungal growth. For one group, 3 500 mL Erlenmeyer flasks containing 200 mL of a potato D-glucose medium (1% glucose) were inoculated with a

piece (1 cm²) of agar bearing the slantwise culture of *L. theobromae* OCS 71, individually. Chemicals were first dissolved in 0.5 mL DMSO (dimethyl sulfoxide) and then fed into the culture medium. As a control group, 3 flasks of fungal culture were fed with 0.5 mL of DMSO without chemicals. As a positive control group, 3 flasks only contained fungal culture. All groups were statically cultured at 25°C in the dark for 7 days, and then the mycelia were harvested and dried upon filter paper under reduced pressure for 12 h. The fresh weight of mycelia was measured. The presence of theobroxide in the culture medium of each group was monitored by TLC with MeOH–CHCl₃ (5:95, v/v).

Quantification of theobroxide: After 7 days, the fungal culture in each flask was filtered using 3 layers of gauze. One mL of culture filtrate from each flask was added into 100 µL of internal standard, 2-methoxy-4-methylphenol (100 µg in EtOAc), and then extracted with 1 mL of EtOAc, 3 times. The EtOAc layer was again filtered using a cellulose acetate 0.45 µm filter, dried over Na₂SO₄ and then concentrated to dryness under vacuum. The concentrated sample was re-dissolved in 1 mL of acetonitrile (HPLC grade). Ten µL of sample solution was analyzed by HPLC (Waters model 600 system) equipped with an RP ODS column (GL Science, 4.6 mm ID × 250 mm) and a photodiode array detector (Waters 996). Acetonitrile-water (1:1, v/v) was used as the mobile phase at a flow rate of 1.0 mL/min. The detection wavelength was set at 210 nm and the column temperature was 25°C. The retention time of theobroxide was 2.7 min and that of the internal standard 6.1 min. The content of theobroxide

was estimated using the following equation: ($C_t = f \times A_t / A_s \times C_s$), where C_t , A_t , A_s , and C_s are the content of target compound, peak area of target compound, peak area of standard substance, and content of injected standard substance, respectively. The correction factor f is the average value (R.S.D. < 2.0%) of 4 tests, which is calculated as $f = (M_t / A_t) / (M_s / A_s)$ using the mass (M) and peak area (A) of the target compound (t) and standard substance (s). Statistical analysis of data was performed by ANOVA analysis and the Bonferoni test ($p < 0.05$ and $p < 0.01$) in order to compare different data in the experiments.

Feeding experiments: *L. theobromae* was grown by static culture on liquid potato-1%-D-glucose medium supplemented with 10⁻³ M [7,7-²H₃]-2-methyl-benzene-1,4-diol (**5a**). The culture broth from each feeding experiment of either 7-day-old or 14-day-old cultures was combined individually, filtered, and concentrated to 100 mL. The filtrate was extracted 3 times with an equal volume of EtOAc. The EtOAc extract was concentrated to dryness under reduced pressure and then chromatographed using a silica gel column (24 g silica gel for 7-day-old culture and 45 g for 14 day-old culture) eluting with MeOH–CHCl₃ (5:95, v/v). Compound **1** (0.9 mg/100 mL from 7-day-old culture) and compound **2** (0.36 mg/100mL from 14-day-old culture) were purified by recrystallization from EtOAc: *n*-hexane.

Acknowledgments - The authors thank Dr Eri Fukushima (GC-MS and NMR Laboratory, Hokkaido University, Japan) for NMR measurements.

References

- [1] Hirai T. (1938) Disease of the banana in transport from Formosa (Japanese). *Annual Meeting of the Phytopathological Society of Japan*, **8**, 145-166.
- [2] Aldridge DC, Galt S, Giles D, Turner WB. (1971) Metabolites of *Lasiodiplodia theobromae*. *Journal of Chemical Society*, **9**, 1623-1627.
- [3] Meyer A, Miersch O, Buttner C, Dathe W, Sembdner G. (1984) Occurrence of the plant growth regulator jasmonic acid in plants. *Journal of Plant Growth Regulation*, **3**, 1-8.
- [4] Nishikawa H, Kaishi NN. (1993) Metabolite from *Aspergillus melleus*. *Nippon Nogeikagaku Kaishi, Journal of the Agricultural Chimerical Society of Japan*, **9**, 772-774.
- [5] Yabuta T, Sumiki Y. (1993) Metabolite from *Aspergillus ochraceus*. *Nippon Nogeikagaku Kaishi, Journal of the Agricultural Chimerical Society of Japan*, **9**, 1264-1275.
- [6] Mori K, Gupta AK. (1985) Chiral synthesis of (*R*)-(-)-mellein and (3*R*, 4*aS*)-(+)-ramulosin. *Tetrahedron*, **41**, 5295-5299.
- [7] Peng L, Ryo T, Kosaku T, Kensuke N. (2006) Biosynthesis of theobroxide and its related compounds, metabolites of *Lasiodiplodia theobromae*. *Phytochemistry*, **68**, 819-823.
- [8] Peng L, Kosaku T, Hideyuki M, Yoshihara T. (2005) Novel potato micro-tuber-inducing compound, (3*R*,6*S*)-6-Hydroxylasiodiplodin, from a strain of *Lasiodiplodia theobromae*. *Bioscience, Biotechnology, and Biochemistry*, **69**, 1610-1612.
- [9] Nakamori K, Matsuura H, Yoshihara T, Ichihara A, Koda Y. (1994) Potato micro-tuber inducing substances from *Lasiodiplodia theobromae*. *Phytochemistry*, **35**, 835-839.
- [10] Yoshihara T, Amanuma M, Tsutsumi T, Okumura Y, Matsuura H, Ichihara A. (1996) Metabolism and transport of [2-¹⁴C]jasmonic acid in potato plant. *Plant Cell Physiology*, **37**, 586-590.
- [11] Kashima T, Takahashi K, Matsuura H, Nabeta K. (2009) Biosynthesis of resorcylic acid lactone lasiodiplodin in *Lasiodiplodia theobromae*. *Bioscience, Biotechnology, and Biochemistry*, **73**, 1118-1122.
- [12] Takei R, Takahashi K, Matsuura H, Nabeta K. (2009) Biosynthesis of resorcylic acid lactone. (5*S*)-5-Hydroxylasiodiplodin in *Lasiodiplodia theobromae*. *Bioscience, Biotechnology, and Biochemistry*, **72**, 2522-2524.
- [13] Birch AJ, Donovan FW. (1953) Studies in relation to biosynthesis. Some possible routes to derivatives of orcinol and phloroglucinol. *Australian Journal of Chemistry*, **6**, 360-368.

- [14] Birch AJ, Massy-Westropp RA, Moye CJ. (1955) Studies in relation to biosynthesis. 2-Hydroxy-6-methylbenzoic acid in *Penicillium griseofulvum* Dierckx. *Australian Journal of Chemistry*, **8**, 539-544.
- [15] Eliel EL, Rivard DE, Bergstahler AW. (1953) The synthesis of 6-formylsalicylic acid, esters of 6-bromomethylsalicylic acid, and related substances. *Journal of Organic Chemistry*, **18**, 1979-1988.
- [16] Francesco M, Salvadori P. (2005) An easy two-step reduction of salicylic acids and alcohols to 2-methylphenols. *Synthesis*, **15**, 2479-2481.

Microbial Hydroxylation of *S*-(-)-Perillyl Alcohol by *Fusarium heterosporium*

Ismail Kiran

Department of Chemistry, Faculty of Arts and Sciences, Eskisehir Osmangazi University, 26480, Eskişehir, Turkey

ikiran@ogu.edu.tr; ikiran1971@gmail.com

Received: March 23rd, 2011; Accepted: October 4th, 2011

S-(-)-Perillyl alcohol (*p*-mentha-1, 8-diene-7-ol) (**1**) (500 mg) was converted by *Fusarium heterosporium* ATCC 15625 over 10 days at 25°C to a new metabolite, 1,2-dihydroxyperillyl alcohol (*p*-mentha-8-en-1,2,7-triol) (**3**) in a yield of 13% (70 mg). The structure of **3** was established by IR and NMR spectroscopic, specific rotation, and mass spectral studies.

Keywords: Biotransformation, *Fusarium heterosporium*, *S*-(-)-perillyl alcohol.

Perillyl alcohol (POH) is a dietary monoterpene found in a variety of foods such as mints, cranberries and cherries [1,2]. It is a hydroxy derivative of limonene and possesses chemopreventive properties [3]. At present it is claimed that POH and limonene inhibit, in a dose-dependent manner, the development of various kinds of carcinoma [4-6]. They may act during the initiation phase of carcinogenesis, preventing interaction of carcinogens with DNA, or during the promotion phase, inhibiting cancer cell development and migration. These led to induction of cancer cell apoptosis, re-differentiation of tumor cells, and influence the molecular mechanisms which regulate their functions. It is believed that post-translational isoprenylation of proteins controlling the growth of cells is the most important mechanism that perillyl alcohol and limonene influence [3,7]. Therefore, the biological derivatization of perillyl alcohol and its structural analog limonene with the aim of producing new metabolites with a variety of biological properties, besides their use as flavoring agents, is an important field of xenobiochemistry, pharmacology and toxicology [8]. The aim of this study was to obtain derivatives of perillyl alcohol through microbial biotransformation.

A few studies are available for the biotransformation of *S*-(-)-perillyl alcohol by microorganisms, plant cell cultures and mammals. These include epoxidation by *Streptomyces ikutanensis* Ya-2-1 giving 8,9-epoxy perillyl alcohol and glycosylation by *Eucalyptus perriniana* suspension cells, which produced the monoglucoside [9,10]. In the present study, the microbiological transformation of *S*-(-)-perillyl alcohol (**1**) by the plant pathogenic fungus *Fusarium heterosporium* ATCC 15625 was examined. The microbial transformation was carried out on an orbital shaker for 10 days at 25°C. This resulted in the production




Figure 1: Biotransformation of *S*-(-)-perillyl alcohol by *F. heterosporium*.

of a complex mixture of products as detected by TLC. The pure metabolite, dihydroxyperillyl alcohol (**3**), was obtained from the column by eluting with 90%, v/v, ethyl acetate in light petroleum (60-80°C) (see Figure 1). The MS of the metabolite showed a molecular ion at *m/z* 186 corresponding to the molecular formula of C₁₀H₁₈O₃. The ¹³C NMR spectrum indicated a new lowfield hydroxyl-bearing carbon and methine signals at δ_C 69.8 and 72.2 ppm, respectively, whereas two carbon signals resonating at δ_C 137.2 and 122.1 disappeared. This indicated that the endocyclic double bond of *S*-(-)-perillyl alcohol (**1**) had been dihydroxylated.




Figure 2: HMBC correlations of 7-CH₂OH signals for *p*-mentha-8-en-1,2,7-triol.

The position of the newly introduced hydroxyl groups were established by the use of Heteronuclear Multi Bond Coherence (HMBC) spectra and comparing the ¹³C NMR and the DEPT spectra of the metabolite with that of **1** and its known metabolites. The DEPT spectrum displayed the presence of a methyl, a hydroxyl bearing methylene, four methylenes, one methine, one hydroxyl bearing methine and a hydroxyl bearing quaternary carbon signals,

indicating that hydroxylations had taken place on C-2 and C-3. The HMBC spectrum confirmed this and showed the key bond connectivity of proton signals of the hydroxy-bearing carbon atoms at 3.45 (1H, d, J 11.4 MHz, $7\text{-CH}_A\text{H}_B\text{OH}$) and 3.82 (1H, d, J 11.4 MHz, $7\text{-CH}_A\text{H}_B\text{OH}$) to the carbon signals at δ_C 72.2 (C-1), 69.8 (C-2) and 25.4 (C-6).

In summary, *F. heterosporium* ATCC 15625 efficiently dihydroxylated perillyl alcohol at the C-2 and C-3 positions, producing a dihydroxylated metabolite, which is recorded in the literature for the first time. The pure metabolite was found to be optically active.

Experimental

General experimental procedures: IR spectra were recorded in KBr pellets. Optical rotation was measured on a Krüss Optronic P8000-T series polarimeter. GC-MS and LCMS spectra were recorded on a 6890 N network GC system, a 5975 inert mass and Applied Biosystems API 3200 instruments, respectively. ^1H NMR and ^{13}C NMR spectra were determined on a Bruker DPX 500 spectrometer, while DEPT spectra were measured at 125 MHz in deuteriochloroform with tetramethylsilane as an internal standard reference. *S*-(-)-perillyl alcohol (*p*-mentha-1,8-diene-7-ol) (**1**) was purchased from Aldrich (218391). The solvents used for purification were distilled prior to use. Silica gel type 60 (Merck, 230–400 mesh) was used for CC. TLC was carried out on a 0.25 mm thick silica gel plate (Merck silica gel 60 GF₂₅₄) in *n*-hexane/ethyl acetate (1:1, v/v). Compounds were detected either under UV light (at 254 nm) or by spraying with vanillin/sulfuric acid (1:1).

Microorganism: *Fusarium heterosporium* ATCC 15625 was provided by Prof. Dr Fatih Demirci, Faculty of Pharmacy at Anadolu University, Eskişehir, Turkey.

References

- Belanger JT. (1998) Perillyl alcohol: applications in oncology. *Alternative Medicine Review*, **3**, 448–457.
- Paduch R, Kandefer-Szerszeń M, Trytek M, Fiedurek J. (2007) Terpenes: substances useful in human healthcare. *Archivum Immunologiae et Therapia Experimentalis*, **55**, 315–327.
- Chan NLS, Wang H, Wang Y, Leung HY, Leung LK. (2006) Polycyclic aromatic hydrocarbon-induced CYP1B1 activity is suppressed by perillyl alcohol in MCF-7 cells. *Toxicology and Applied Pharmacology*, **213**, 98–104.
- Crowell PL. (1999) Prevention and therapy of cancer by dietary monoterpenes. *Journal of Nutrition*, **129**, 775S–778S.
- Bardon S, Foussard V, Fournel S, Loubat A. (2002) Monoterpenes inhibit proliferation of human colon cancer cells by modulating cell cycle-related protein expression. *Cancer Letters*, **181**, 187–194.
- Gupta A, Myrdal PB. (2004) Development of a perillyl alcohol topical cream formulation. *International Journal of Pharmacology*, **269**, 373–383.
- Crowell PL. (1997) Monoterpenes in breast cancer chemoprevention. *Breast Cancer Research Treatment*, **46**, 191–197.
- Bhatti HN, Zubair M, Rasool N, Hassan Z, Ahmad VU. (2009) Transformation of sesquiterpenoids. *Natural Product Communications*, **4**, 1155–1168.
- Noma Y, Asakawa Y. (2010) Biotransformation of monoterpenoids by microorganisms, insects and mammals. In *Handbook of essential oils*, Baser KHC, Buchbauer G. (Eds), CRC Press, New York, USA, 645–646.
- Cardillo R, Nasini G, De Pava OV. (1992) Isolation of illudin M, illudine and shisoal from the mushroom *Clitocybe phosphorea*; biotransformation of perillyl alcohol and aldehyde. *Phytochemistry*, **31**, 2013–2014.
- Kiran I, Yildirim HN, Hanson JR, Hitchcock PB. (2004) The antifungal activity and biotransformation of diisophorone by the fungus *Aspergillus niger*. *Journal of Chemical Technology and Biotechnology*, **79**, 1366–1370.

General culture conditions: The cultures were maintained and precultured on potato dextrose agar (PDA) slants at 5°C and 25°C, respectively [11]. A medium for growing *F. heterosporium* was prepared by mixing glucose (20 g), yeast extract (5 g), polypeptone (5 g), NaCl (5 g) and Na_2HPO_4 (5 g) in distilled water (1 L). *S*-(-)-perillyl alcohol (**1**) (500 mg) was added into 20 Erlenmeyer flasks (250 mL) containing 100 mL of freshly prepared and autoclaved medium after 2 days' growth. The fermentations were continued for a further 10 days. The mycelium was filtered off and the broth was extracted with ethyl acetate. The extract was dried over Na_2SO_4 and the solvent evaporated to give a residue which was purified by means of CC (3 cm wide, 60 cm long) on silica eluting with increasing concentrations of ethyl acetate in light petroleum. A control experiment was run simultaneously.

Biotransformation by *F. heterosporium*: The product, 1,2-dihydroxyperillyl alcohol (**3**) (70 mg; 13 % yield), was eluted with 90%, v/v, ethyl acetate in light petroleum (60–80°C) as an oily compound.

$[\alpha]_D^{20}$: 96.0 (2.5 mg in 1 mL acetone, 20°C).

FT-IR ν_{max} cm^{-1} : 3404 (OH), 1641 (C=C);

^1H NMR (500 MHz, CDCl_3): 1.75 (3H, s, 10- CH_3), 3.45 (1H, d, J = 11.4 MHz, 7- $\text{CH}_A\text{H}_B\text{OH}$), 3.82 (1H, d, J = 11.4 MHz, 7- $\text{CH}_A\text{H}_B\text{OH}$), 3.91 (1H, s, 2-CH), and 4.75 (2H, br. s, 9- CH_2OH).

^{13}C NMR (125 MHz, CDCl_3): C-8 (149.3), C-9 (108.9), C-1 (72.2), C-2 (69.8), C-7 (68.0), C-4 (37.8), C-3 (33.9), C-5 (28.7), C-6 (25.4), C-10 (20.8).

LCMS m/z : $[\text{M}+1]$ 187.066, (calcd for $\text{C}_{10}\text{H}_{18}\text{O}_3$, $[\text{M}+1]$ 187.241).

Acknowledgements - Dr Hakan DAL and Dr Fatih Göger from Anadolu University, Eskişehir-TURKEY are kindly acknowledged for providing the NMR and LCMS analyses, respectively.

A Phytochemical Investigation of *Zanthoxylum setulosum*

Tameka M. Walker^a, Bernhard Vogler^a, Debra M. Moriarity^b, William A. Haber^c and William N. Setzer^{a,*}

^aDepartment of Chemistry, University of Alabama in Huntsville, Huntsville, Alabama 35899, USA

^bDepartment of Biological Sciences, University of Alabama in Huntsville, Huntsville, Alabama 35899, USA

^cMissouri Botanical Garden, St. Louis, Missouri 63166, USA
Apdo. 50-5655, Monteverde, Puntarenas, Costa Rica, Central America

wsetzer@chemistry.uah.edu

Received: September 19th, 2011; Accepted: September 28th, 2011

The crude bark extract of *Zanthoxylum setulosum* from Monteverde, Costa Rica was notably cytotoxic (100% kill at 100 µg/mL) to MCF-7, MDA-MB-231, and MDA-MB-468 cells *in vitro*. Phytochemical studies of the bark extract revealed the triterpenoid lupeol, the lignan sesamin, the sesquiterpene sesquichamaenol, and the xanthone lichexanthone. This is the first report of the isolation of sesquichamaenol and lichexanthone from the bark extract of *Z. setulosum*. All structures were determined using NMR spectroscopic techniques (¹H NMR and ¹³C NMR) and GC-MS and by comparison with literature data. Lupeol proved to be the cytotoxic component of *Z. setulosum* bark.

Keywords: *Zanthoxylum setulosum*, lupeol, sesamin, sesquichamaenol, lichexanthone, cytotoxicity.

Tropical rainforest plants have evolved chemical defensive mechanisms to avoid herbivory and resist pathogens [1]. Monteverde, Costa Rica, is home to numerous species of *Zanthoxylum* that contain common bioactive constituents and have various traditional uses [2a]. *Zanthoxylum*, a genus of about 250 species in Rutaceae, is a rich source of lignans, alkaloids, coumarins, amides, flavonoids, terpenes and other metabolites with pharmacological properties [2b,2c].

Z. setulosum P. Wilson is a medium to large tree [3]. Lupeol and sesamin have been previously identified in the leaves of *Z. setulosum* [4]. *Z. setulosum* bark extract had been previously shown to be cytotoxic [5] and in this work, the crude chloroform bark extract was cytotoxic to MCF-7, MDA-MB-231, and MDA-MB-468 breast tumor cells (100% kill at 100 µg/mL).

The crude chloroform bark extract of *Z. setulosum* was subjected to flash chromatography leading to the isolation of lupeol, sesamin, lichexanthone, and sesquichamaenol. The structures of the isolated components were determined using NMR and MS spectral data and by comparison of these with the literature [6-8].

Previous phytochemical investigation of *Z. setulosum* has shown the isolation from the leaf extract of sesamin, syringaresinol, the isopentyl ether of pluviatol, savinin, kusunokinin, lupeol, aurantiamide, and skimmianine [4]. In this work we report the isolation of sesquichamaenol

and lichexanthone that have not been previously reported from *Z. setulosum*. In addition, lupeol and sesamin have been isolated in larger quantities from the bark than previously isolated from the leaves of this tree [4].

Sesquichamaenol is a sesquiterpene that is reported to be a source of various metabolites with pharmacological activities [9]. Lichexanthone is recorded as having antibacterial, antimalarial, antioxidant, and antitumor activity [10]. Lichexanthone has been isolated from a variety of *Zanthoxylum* species, but has not been reported in *Z. setulosum* [11]. Cytotoxicity testing of isolated and purified compounds indicated lupeol to be the cytotoxic material in the crude bark extract (100% kill at 100 µg/mL). Lupeol had been previously shown to be cytotoxic to Hep-G2, A-431 and H-4IIE cells [12]. However, neither sesamin, sesquichamaenol, nor lichexanthone showed cytotoxic activity.

Experimental

Collection and extraction of plant material: *Zanthoxylum setulosum* bark was collected in June, 2004, in Provincia Puntarenas, Costa Rica, 6 km SSW of Santa Elena on the main road from Monteverde. *Z. setulosum* was identified and authenticated by W.A. Haber and a voucher specimen (Haber 9982) has been deposited in Missouri Botanical Garden Herbarium. The chopped, dried bark (952 g) was extracted with refluxing chloroform for 6 h. Evaporation of the solvent gave 31.9 g of crude chloroform extract.

Chromatographic separation of bark extract: The crude chloroform bark extract (25.02 g) was subjected to flash chromatography using a silica gel (200-400 mesh) column, 70 cm in length \times 5 cm in diameter. Elution was carried out using a solvent gradient (*n*-hexane with increasing proportions of ethyl acetate, followed by ethyl acetate with increasing proportions of ethanol). TLC was performed on silica gel precoated aluminum plates with a fluorescent indicator 254 nm for analysis of isolated fractions, and spots were detected by illumination with UV light. Lupeol was isolated from fraction 14 (yield 1.28 g). Sesamin crystallized from fraction 29 and 30, as colorless prisms (yield 2.05 g).

Fractions were checked by TLC and HPLC analysis and fractions having comparable TLC features were combined to form "superfractions." Combined fractions 5-11 (22 mg) and 20-23 (69 mg) were subjected to chromatographic separation on a silica gel (200-400 mesh) column, eluting with a *n*-hexane/EtOAc step gradient. Superfraction 5-11 was separated to give 10 mg of sesquichamaenol, and superfraction 20-23 to give 10 mg of lichexanthone.

Gas chromatography-mass spectrometry: Superfractions of *Z. setulosum* were subjected to gas chromatographic-mass spectral analysis on an Agilent system consisting of a 6890 Gas Chromatograph, a 5973 Mass selective detector (MSD), and an Agilent ChemStation data system. The GC

column was an HP-5ms fused silica capillary consisting of a (5% phenyl)-methylpolysiloxane stationary phase, a film thickness of 0.25 μ m, a length of 30 m, and an internal diameter of 0.25 mm. The GC carrier gas was helium and the column had a head pressure of 7.07 psi and a flow rate of 1.0 mL/min. The inlet temperature was 200°C and the MSD temperature 280°C. The GC oven temperature was programmed as follows: 40°C initial temperature held for 10 min; increased at 3°/min to 200°C; increased 2°/min to 220°C. The extracts were dissolved in dichloromethane and 1 μ L of the sample was injected using a split injection method. The MS of constituents were compared with those in the literature [7-9].

Cytotoxicity screening: *In-vitro* cytotoxic activity against MCF-7, MDA-MB-231, and MDA-MB-468 cells was determined using the MTT method for cell viability, as previously described [13].

Acknowledgments – We are grateful to the Monteverde Cloud Forest Preserve and the Tropical Science Center for granting permission to collect plant materials under a cooperative rights agreement and to the Commission for the Development of Biodiversity of Costa Rica's Ministry of the Environment, Energy, and Telecommunications for Research Permit R-001-2006-OT-CONAGEBIO.

References

- [1] Howe HF, Westley LC. (1990) *Ecological Relationships of Plants and Animals*, Oxford University Press, Oxford, UK.
- [2] (a) Haber WA, Zuchowski W, Bello E. (2000) *An Introduction to Cloud Forest Trees, Monteverde, Costa Rica*, Mountain Gem Publications, Monteverde de Puntarenas, Costa Rica; (b) Gentry AH. (1993) *A Field Guide to the Families and Genera of Woody Plants of Northwest South America*, University of Chicago Press, Chicago, Illinois; (c) Stermitz FR, Sharifi IA. (1977) Alkaloids of *Zanthoxylum monophyllum* and *Zanthoxylum punctatum*. *Phytochemistry*, **16**, 2003-2006.
- [3] Boehme AK, Noletto JA, Haber WA, Setzer WN. (2008) Bioactivity and chemical composition of the leaf essential oils of *Zanthoxylum rhoifolium* and *Zanthoxylum setulosum* from Monteverde, Costa Rica. *Natural Product Research*, **22**, 31-36.
- [4] Mora A, Castro V, Poveda L, Chavarria M, Murillo R. (2011) Chemical constituents from *Zanthoxylum setulosum* (Rutaceae). *The Latin American and Caribbean Bulletin of Medicinal and Aromatic Plants*, **10**, 155-158.
- [5] Setzer MC, Moriarity DM, Lawton RO, Setzer WN, Gentry GA, Haber WA. (2003) Phytomedicinal potential of tropical cloudforest plants from Monteverde, Costa Rica. *Revista de Biología Tropical*, **51**, 647-674.
- [6] (a) Baures PW, Miski M, Eggleston DS. (1992) Structure of sesamin. *Acta Crystallographica Section C Crystal Structure Communications*, **48**, 574-576; (b) Laggoune S, Brouard I, Leon F, Calliste CA, Duroux J, Bermejo J, Kabouche Z, Kabouche A. (2011) Lignans and an abundant flavone glycoside with free-radical scavenging activity from the roots of the endemic species *Stachys mialhesi* de Noe. *Records of Natural Products*, **5**, 238-241.
- [7] (a) Letcher RM. (1968) Chemistry of lichen constituents-VI: Mass spectra of usnic acid, lichexanthone and their derivatives. *Organic Mass Spectrometry*, **1**, 551-561; (b) Díaz AB, Vera JR, Cote V, Bruno-Colmenarez J, Diaz de Delgado G. (2010) NMR elucidation and crystal structure analysis of 1-hydroxy-3,6-dimethoxy-8-methyl-9h-xanthen-9-one (lichexanthone) isolated from *Vismia baccifera* (Guttiferae). *Latin American and Caribbean Bulletin of Medicinal and Aromatic Plants*, **9**, 470-474.
- [8] Koorbanally NA, Randrianarivelojosia M, Mulholland DA, van Ufford LQ, van den Berg AJ. (2002) Bioactive constituents of *Cedrelopsis microfoliata*. *Journal of Natural Products*, **65**, 1349-1352.
- [9] Warrior G, Lima EJS, Fernandes JB, da Silva MFGF, Vieira PC, Magellan CMI, Pagnocca CF, Bueno OC, Hebling MJA, Pirani JR. (2005) Sesquiterpenes from the stem of *Pilocarpus riedelianus* and activities of microorganisms. *Química Nova*, **40**, 986-990.
- [10] Pinto MM, Sousa ME, Nascimento MS. (2005) Xanthone derivatives: New insights in biological activity. *Current Medicinal Chemistry*, **12**, 2517-2538.
- [11] Jimenez C, Marcos M, Villaverde MC, Riguera R, Castedo L, Stermitz F. (1989) A chromone from *Zanthoxylum* species. *Phytochemistry*, **28**, 1992-1993.
- [12] Moriarity DM, Huang J, Yancey CA, Zhang P, Setzer WN, Lawton RO, Bates RB, Caldera S. (1998) Lupeol is the cytotoxic principle in the leaf extract of *Dendropanax cf. querceti*. *Planta Medica*, **64**, 370-372.
- [13] Moriarity DM, Bansal A, Cole RA, Takaku S, Haber WA, Setzer WN. (2007) Selective cytotoxic activities of leaf essential oils from Monteverde, Costa Rica. *Natural Product Communications*, **2**, 1263-1268.

Cytotoxic Cembranoids from the Red Sea Soft Coral *Sarcophyton glaucum*

Mohamed-Elamir F. Hegazy^{a,b}, Ahmed A. El-Beih^{b,c,*}, Alaa Y. Moustafa^d, Abdelhamed A. Hamdy^c,
Montaser A. Alhammady^e, Rehab M. Selim^c, Mohamed Abdel-Rehim^{b,f,g} and Paul W. Paré^h

^aChemistry of Medicinal Plants Department, National Research Centre, Dokki, Giza 12622, Egypt

^bPharmaceutical Research Laboratory, Center of Excellence for Advanced Sciences,
National Research Center, Dokki 12622, Cairo, Egypt

^cChemistry of Natural and Microbial Products Department, National Research Centre, Dokki,
Giza 12622, Egypt

^dZoology Department, Faculty of Science, Sohag University, Egypt

^eNational Institute of Oceanography and Fisheries, Red Sea Branch, Hurghada, Egypt

^fDepartment of Chemistry & Biomedical Sciences, Faculty of Technology and Science,
Karlstad University, SE-651 88 Karlstad, Sweden

^gDepartment of Clinical Pharmacology & DMPK, Astra Zeneca, R&D, Södertälje, 15185 Södertälje,
Sweden

^hDepartment of Chemistry and Biochemistry, Texas Tech University, Lubbock, Texas 79409-1061, USA


aae2eg@yahoo.com

Received: June 21st, 2011; Accepted: August 3rd, 2011

One new cembrane diterpene, 2*R*,7*R*,8*R*-dihydroxydeepoxysarcophine (**1**), together with three known compounds, 7*α*,8*β*-dihydroxy-deepoxysarcophine (**2**), 7*β*-acetoxy-8*α*-hydroxydeepoxysarcophine (**3**), and sarcophine (**4**), have been isolated from the Red Sea soft coral *Sarcophyton glaucum*. Their structures were determined using 1D and 2D NMR spectroscopy. 7*β*-Acetoxy-8*α*-hydroxydeepoxysarcophine (**3**) exhibits cytotoxic activity against HepG2, HCT-116, and HeLa cells with IC₅₀ values of 3.6, 2.3, and 6.7 µg/mL, respectively.

Keywords: *Sarcophyton glaucum*, cembrane diterpenes, dihydroxydeepoxysarcophine, sarcophine.

Marine organisms, comprising more than half of the total global diversity, offer an enormous source of novel and biologically active compounds. The soft corals (phylum Cnidaria, class Anthozoa, subclass Octocorallia, order Scleractinia) have proven to be a biochemical warehouse for terpenoids. Nearly 30 species of soft corals of the genus *Sarcophyton*, from different geographical areas, have been chemically examined. They contain antiproliferative sesquiterpenes [1] and an impressive series of cembranoid diterpenes [2]. Many cembrane-type compounds have been shown to exhibit cytotoxicity and anti-inflammatory activity [3-5]. Herein, we report the isolation of the new compound 2*R*,7*R*,8*R*-dihydroxydeepoxysarcophine (**1**), together with three known cembranoides, 7*α*,8*β*-dihydroxydeepoxysarcophine (**2**), 7*β*-acetoxy-8*α*-hydroxydeepoxysarcophine (**3**), and sarcophine (**4**), from the Red Sea soft coral *S. glaucum* and cytotoxic activity against HepG2, HCT-116, HeLa, and MCF-7 cells. Structures of these isolated metabolites were elucidated through extensive 1D and 2D spectroscopic analyses.




Compound **1** was obtained as colorless crystals, $[\alpha]_D^{25}$ -21.2 (*c* 1.86, MeOH). The HR-ESI-MS exhibited a $[M+Na]^+$ ion at *m/z* 357.20371 (calc. 357.20363), suggesting a molecular formula C₂₀H₂₉O₄Na and six degrees of unsaturation that was supported by NMR data. The IR spectrum indicated the presence of an α,β -unsaturated- γ -lactone group (1750 and 1686 cm⁻¹), a ketone group (1707 cm⁻¹), an olefinic group (1669 cm⁻¹), and broad absorption for OH stretching (3000–3353 cm⁻¹) functionalities. A literature survey and careful comparison of the ¹H and ¹³C NMR spectroscopic data of **1** (Table 1) revealed that it was in part close to the known compounds, 7*β*,8*α*-dihydroxydeepoxysarcophine (**1a**) and 7*α*,8*β*-dihydroxydeepoxysarcophine (**2**) [6a]. The ¹³C NMR and

Table 1: NMR spectroscopic data of compound **1**.

Position	δ_H (J in Hz)	δ_C	HMBC
1	--	163.1 C	
2	5.43 (dd, 10.0, 1.5)	80.1 CH	C-1, C-3, C-4
3	4.95 (dd, 10.0, 1.0)	118.7 CH	C-5, C-18
4	--	146.2 C	
5	2.24 (m)	36.5 CH ₂	C-3, C-4, C-6, C-7, C-18
6	1.60 (m)	29.1 CH ₂	C-4, C-5, C-7
	1.79 (m)		C-4, C-5, C-7, C-8
7	3.33 (dd, 10.0, 1.5)	73.8 CH	C-5, C-6, C-8, C-19
8	--	75.3 C	
9	1.68 (m)	38.4 CH ₂	C-7, C-8, C-10, C-11, C-19
10	2.10 (m)	22.4 CH ₂	C-12
11	5.24 (td, 7.5, 1.0)	126.6 CH	C-9, C-10, C-13, C-20
12	--	135.3 C	
13	2.37 (m)	36.7 CH ₂	C-1, C-11, C-12, C-14, C-20
14	2.10 (m)	26.1 CH ₂	C-1, C-2, C-12, C-13, C-15
	2.62 (m)		C-1, C-2, C-12, C-13, C-15
15	--	123.1 C	
16	--	175.1 C	
17	1.88 (brt, 1.5)	9.3 CH ₃	C-1, C-15, C-16
18	1.90 (d, 1.0)	17.7 CH ₃	C-3, C-4, C-5
19	1.20 (s)	24.3 CH ₃	C-7, C-8, C-9
20	1.64 (s)	16.1 CH ₃	C-11, C-12, C-13

Recorded in CDCl₃ and obtained at 500 and 125 MHz for ¹H and ¹³C NMR, respectively.

**Figure 1:** (A) ¹H-¹H COSY (bold lines) and key HMBC (arrows) correlations of **1**; (B) Selected NOESY correlations in compound **1**.

DEPT spectra (Table 1) revealed the presence of four olefinic functionalities at C-3 (δ 118.7), C-11 (δ 126.6), C-12 (δ 135.3), and C-4 (δ 146.2), two oxymethine carbons at C-7 (δ 73.8) and C-2 (δ 80.1), and one oxygenated quaternary carbon at C-8 (δ 75.3). The carbon signal resonances at C-15 (δ 123.1), C-1 (δ 163.1), and C-16 (δ 175.1) were assigned to the α,β -unsaturated- γ -lactone functionality that was confirmed by comparison of NMR data for **1** with **4**.


From the ¹H NMR and ¹H-¹H COSY spectra, the following proton correlations were assigned H-2/H-3, H₂-5/H₂-6/H-7, H₂-9/H₂-10/H-11, and H₂-13/H₂-14. The HMBC correlations H-2/C-1, H-2/C-4, H-3/C-5, H₂-9/C-7, H₂-10/C-12, H-11/C-13, H₂-13/C-1, and H₂-14/C-2 confirmed the connectivities of the 14-membered ring from C1–C14. The position of the methyl groups H₃-18 (δ 1.90), H₃-19 (δ 1.20), and H₃-20 (δ 1.64) was deduced from the HMBC correlations H-3/C-18, H₃-18/C-4, H₃-19/C-8, H₃-19/C-7, H₃-19/C-9, and H₃-20/C-12. The last methyl group, H₃-17 (δ 1.88), was connected to the α,β -unsaturated- γ -lactone ring from the HMBC correlations with the olefinic carbons C-1 (δ 163.1) and C-15 (δ 123.1) and the carbonyl carbon C-16 (δ 175.1). The most significant differences observed

Table 2: Comparison of ¹H and ¹³C NMR spectroscopic data of compounds **1**, **2**, and **1a**.

	1	2	1a		1	2	1a
H-2	5.43, dd (10.0, 1.5)	5.57, dq (13.6, 2.4)	5.55, dq (10.1, 1.3)	C-2	80.1	79.3	79.1
H-7	3.33, dd (10.0, 1.5)	3.47, d (13.5)	3.66, dd (9.6, 9.5)	C-7	73.8	72.5	72.3
H-11	5.24, td (7.5, 1.0)	4.98 (m)	5.10, dd (6.7, 6.6)	C-11	126.6	125.2	124.0
H ₃ -19	1.20 (s)	1.18 (s)	1.54 (s)	C-19	24.3	24.2	26.4

in the ¹H NMR spectrum of **1** are the downfield chemical shift of H-11, which resonates at δ 5.24 (td, J = 7.5, 1.0), and the upfield chemical shift of H-7 [δ 3.33 (dd, J = 10.0, 1.5)] rather than the values reported for compounds **2** and **1a** [6b] (Table 2). These data indicate differences in the stereochemistry of the three chiral carbons C-2, C-7, and C-8 compared with those of **1a** and **2**.

The relative configuration of **1** was determined on the basis of the study of the coupling constants and NOESY experiments. The H-2 and H-3 vicinal coupling constant (10.0 Hz), as well as a NOESY correlation of H-2 with H₃-18, establish a *trans* configuration between the γ -lactone (H-2) and the olefinic protons (H-3); the proton signal H-5 (2.24, m) showed clear correlations with the H₃-18 methyl protons (1.90, d, J = 1.0); the oxygenated proton at C-7 (δ_H 3.33, dd, J = 10.0, 1.5) showed a clear correlation with the H-6 (1.60, m) and H₃-19 (1.20, s) protons. From this information, we noted that H-2, H-7, and H₃-19 all faced in the same direction (Figure 1b).

**Figure 2:** Circular dichroism spectra of compounds **1** and **2** in MeOH.

The absolute stereochemistry of **1** at C-2 was determined via circular dichroism (CD) analysis (Figure 2). The positive and negative Cotton effect ($[\theta]_{248} +1.12$; $[\theta]_{225} -2.19$) observed in the CD spectrum for the electronic transitions of the 2(5*H*)-furanone moiety indicate a left-handed (*M*) helix for the five-membered α,β -unsaturated- γ -lactone ring [6c]. Consistent with this interpretation, the similar optical rotation for **1** $\{[\alpha]_D^{25} -21.2$ (c 1.86, MeOH)} and *ent*-sarcophine $\{[\alpha]_D^{25} = -16.0$ (c 1.19, MeOH)} [7], confirm the same *R* absolute configuration at C-2, C-7, and C-8 [6-8]. These data establish the isolation and identification of 7 β ,8 β -dihydroxydepoxy sarcophine, a newly identified natural product.

Compounds **2–4** were previously reported from the soft coral *S. glaucum* [6b,9] with no report of cytotoxicity. The

Table 3: Cytotoxic effect of different samples against different human cancer cell lines, using MTT assay ($n=4$); data expressed as the mean value of IC_{50} ($\mu\text{g/mL}$) \pm S.E. Sample results are compared with paclitaxel.

Compound	Cell line		
	Hep-G2	HCT-116	HeLa
1	38.8 ± 1.0	26.3 ± 2.7	> 50
2	23.5 ± 2.4	15.6 ± 1.9	44.1 ± 5.4
3	3.6 ± 1.0	2.3 ± 1.5	6.7 ± 0.8
Paclitaxel	0.49 ± 0.1	0.4 ± 0.1	0.6 ± 0.08

cytotoxic activity of compounds **1–3** was tested against HepG2, HCT-116, HeLa, and MCF-7 cells. **7 β -Acetoxy-8 α -hydroxydepoxy-sarcophine (3)** showed strong cytotoxic activity against HepG2, HCT-116, and HeLa cells with IC_{50} values of 3.6, 2.3, and 6.7 $\mu\text{g/mL}$, respectively, while no activity against MCF-7 cells was observed for compounds **1** and **2**.

Using MTT assay, the effect of isolated compounds on the growth of various cell lines was studied after 48 h of incubation. As shown in Table 3, compound **3** showed a potential cytotoxic activity (low IC_{50} values <10 $\mu\text{g/mL}$) against HepG2, HCT-116, and HeLa cells, but not against MCF-7 cells, while the other tested compounds were not cytotoxic (IC_{50} values >10 $\mu\text{g/mL}$). Paclitaxel, a known anti-cancer drug, was used as a positive control.




Figure 3: (A) Type of cell death was investigated in HepG2, HCT-116, and HeLa cells after treatment with compound **3**, using acridine orange/ethidium bromide staining to compare between the percentage of necrotic cells (grey segment) and apoptotic cells (black segment). (B) Effect of compound **3** (grey bars) on the percentage of DNA fragmentation in different cancer cell types compared with untreated cells (black bars). The fragmented DNA was determined with the diphenylamine reaction. Values are means of three measurements (mean \pm SD), where (*) represent the $p < 0.05$ in comparison with control.

To detect the type of cell death induced by compound **3** in different cancer cell lines, cells were treated with the appropriate IC_{50} value of compound **3** for 12 h and the apoptosis and necrosis percentages were recorded using acridine orange/ethidium bromide staining and analysis under a fluorescence microscope. As shown in Figure 3A, compound **3** led to an apoptosis-dependant cell death in all of the tested cell lines (59.7–83.2% of the total number of dead cell population), while the percentage of necrotic cells was 16.8–40.3% of the total number of the dead cell population. Comparing the untreated cells with paclitaxel-treated cells indicated that the latter significantly induced DNA fragmentation up to 57% ($p < 0.001$) in different cell lines, as shown in Figure 3B. Similarly, treatment of different cells with compound **3** resulted in high, to variable extents, DNA fragmentation levels that were

significantly different from the control ($p < 0.01$ – 0.001), in the following order: HCT-116 > HeLa > HepG2.

Experimental

General experimental procedures: 1D and 2D NMR, Varian Mercury Plus 300 MHz and Varian Unity INOVA 500 spectrometer; CI-MS, Finnigan LCQ Ion Trap mass spectrometer; HR-CI-MS, Fourier transform ion cyclotron mass spectrometer (Ion Spec, Varian); Optical rotations, Perkin–Elmer-341 MC digital polarimeter; CD, OLIS DSM-10UV/Vis CD; CC, Silica gel 60 (Merck, 230–400 mesh) and Sephadex LH-20 (Sigma); TLC, Pre-coated silica gel plates (Merck, Kieselgel 60 F₂₅₄, 0.25 mm).

Animal material: The *S. glaucum* was collected by SCUBA in June 2009 from Hurghada, on the Egyptian Red Sea coast. A voucher specimen (03RS24) was deposited in the Marine Biological Station, Hurghada, Red Sea, Egypt.

Extraction and isolation: The frozen soft coral was chopped into small pieces (4 kg, wet wt) and extracted with EtOAc at room temperature ($4\text{L} \times 5$). The combined EtOAc extracts were concentrated to a brown gum (20.0 g), which was subjected to CC on a silica gel column (6×120 cm) using *n*-hexane–EtOAc (gradient separation) to yield 8 fractions. Fraction 2, eluted with *n*-hexane–EtOAc (3:1), was separated by Sephadex LH-20 using *n*-hexane– CHCl_3 –MeOH (7:4:0.5) to give **1** (10 mg) and **3** (15 mg). Fraction 6 eluted with *n*-hexane–EtOAc (1:1) was separated on silica gel using *n*-hexane–EtOAc, followed by Sephadex LH-20 using *n*-hexane– CHCl_3 –MeOH (7:4:0.5), to give **2** (12 mg).

2R,7R,8R-Dihydroxydepoxy-sarcophine (1)

Colorless crystals.

$[\alpha]_D^{25}$: -21.2 (c 1.86, MeOH).

IR (film): 1750, 1686, 1707, 1669 cm^{-1} .

^1H and ^{13}C NMR: Table 1.

HRCIMS: m/z $[M + \text{Na}]^+$ calcd for $\text{C}_{20}\text{H}_{29}\text{O}_4\text{Na}$: 357.20363; found: 357.20371.

Cell culture: Human breast adenocarcinoma (MCF-7), hepatocellular carcinoma (HepG2), cervical carcinoma (HeLa), and colon carcinoma cells (HCT-116) were purchased from ATCC, USA, and used to evaluate the cytotoxic effect of the tested extracts. Cells were routinely cultured in Dulbecco's modified Eagle's medium (DMEM), while HCT-116 cells were grown in McCoy's medium. Media were supplemented with 10% fetal bovine serum (FBS), 2 mL glutamine, containing 100 units/mL penicillin G sodium, 100 units/mL streptomycin sulfate, and 250 ng/mL amphotericin B. Cells were maintained at sub-confluency at 37°C in humidified air containing 5% CO_2 . For sub-culturing, monolayer cells were harvested after trypsin/EDTA treatment at 37°C . Cells were used when confluence had reached 75%. Test samples were dissolved in DMSO, and then diluted 1000 times for the assay. All experiments were repeated 3 times, unless mentioned.

Cytotoxicity assay: The cytotoxic effect of the compounds on the growth of different human cancer cell lines was estimated by the 3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2*H*-tetrazolium bromide (MTT) assay [10]. Cells (5×10^4 cells/well) were incubated for 48 h with various concentrations of the compounds at 37°C in a FBS-free medium, before submitted to MTT assay. The absorbance was measured with a microplate reader (BioRad, München, Germany) at 570 nm. The relative cell viability was determined by the amount of MTT converted to the insoluble formazan salt. The data are expressed as the mean percentage of viable cells as compared with untreated cells. The relative cell viability was expressed as the mean percentage of viable cells compared with the respective untreated cells (control). The half maximal growth inhibitory concentration (IC_{50}) value was calculated from the line equation of the dose-dependent curve of each compound. The results were compared with paclitaxel.

Apoptosis and necrosis staining: The type of cell death was investigated in compound-treated and -untreated cells using acridine orange/ethidium bromide staining [11,12]. In brief, a mixture of 100 µg/mL acridine orange and 100 µg/mL ethidium bromide was prepared in PBS. The cell uptake of the stain was monitored under a fluorescence microscope, and the apoptotic, necrotic, and viable cells were counted. The early apoptotic cells had yellow

chromatin in the nuclei that were highly condensed or fragmented. Apoptotic cells also exhibited membrane blebbing. The late apoptotic cells had orange chromatin with nuclei that were highly condensed and fragmented. The necrotic cells had bright orange chromatin in round nuclei. Only cells with yellow, condensed, or fragmented nuclei were counted as apoptotic cells in a blinded, nonbiased manner.

DNA fragmentation: DNA fragmentation was essentially assayed as reported previously [13]. Briefly, compound-treated and -untreated cell pellets were re-suspended in 250 µL 10 mM Tris, 1 mM EDTA, pH 8.0 (TE-buffer), and incubated with an additional volume of lysis buffer (5 mM Tris, 20 mM EDTA, pH 8.0, 0.5% Triton X-100) for 30 min at 48°C. After lysis, the intact chromatin (pellet) was separated from DNA fragments (supernatant) by centrifugation. Pellets were re-suspended in TE-buffer and samples were precipitated by 10% trichloroacetic acid at 48°C. The sample pellets were added to 5% trichloroacetic acid and boiled. DNA contents were quantified using the diphenylamine reagent [14]. The percentage of DNA fragmented was calculated as the ratio of the DNA content in the supernatant to the amount in the pellet.

Acknowledgements - Financial assistance was provided in part by a grant from the Robert Welch Foundation (D-1478).

References

- [1] Sawant SS, Youssef DTA, Sylvester PW, Wali V, El Sayed KA. (2007) Antiproliferative sesquiterpenes from the Red Sea soft coral *Sarcophyton glaucum*. *Natural Product Communications*, **2**, 117-119.
- [2] Yan XH, Gavagnin M, Cimino G, Guo YW. (2007) Two new biscebranes with unprecedented carbon skeleton and their probable biogenetic precursor from the Hainan soft coral *Sarcophyton latum*. *Tetrahedron Letters*, **48**, 5313-5316.
- [3] Gross H, Wright AD, Beil W, König G. (2004) Two new bicyclic cembranoides from a new *Sarcophyton* species and determination of the absolute configuration of sarcoglaucol-16-one. *Organic & Biomolecular Chemistry*, **2**, 1133-1138.
- [4] Sawant S, Youssef D, Mayer A, Sylvester P, Wali V, Arant M, Sayed KE. (2006) Anticancer and anti-inflammatory sulfur-containing semisynthetic derivatives of sarcophine. *Chemical & Pharmaceutical Bulletin*, **54**, 1119-1123.
- [5] Sawant S, Youssef D, Reiland J, Ferniz M, Marchetti D, Sayed KE. (2006) Biocatalytic and antimetastatic studies of the marine cembranoid sarcophine and 2-*epi*-16-deoxysarcophine. *Journal of Natural Products*, **69**, 1010-1013.
- [6] (a) Yao LG, Liu HL, Guo YW, Mollo E. (2009) New cemranoids from the Hainan soft coral *Sarcophyton glaucum*. *Helvetica Chimica Acta*, **92**, 1085-1091; (b) El Sayed KA, Hamann MT, Waddling CA, Jensen C, Lee SK, Dunstan CA, Pezzuto JM. (1998) Structurally novel bioconversion products of the marine natural product sarcophin effectively inhibit JB6 cell transformation. *Journal of Organic Chemistry*, **63**, 7449-7455; (c) Kamel HN, Ferreira D, Garcia-Fernandez LF, Slattery M. (2007) Cytotoxic Diterpenoids from the hybrid soft coral *Sinularia maxima* × *Sinularia polydactyla*. *Journal of Natural Products*, **70**, 1223-1227.
- [7] Kashman Y, Zadock E, Neeman I. (1974) Some new cembrane derivatives of marine origin. *Tetrahedron*, **30**, 3615-3620.
- [8] Yin SW, Shi YP, Li XM, Wang BG. (2005) A novel hydroperoxyl substituted cembranoliditerpene from marine soft coral *Lobophytum crissum*. *Chinese Chemical Letters*, **16**, 1489-1491.
- [9] Grote D, Shaker KH, Soliman HSM, Hegazi MM, Seifert K. (2008) Cembranoid diterpenes from the soft corals *Sarcophyton* sp. and *Sarcophyton glaucum*. *Natural Product Communications*, **3**, 1473-1478.
- [10] Hansen MB, Nielsen SE, Berg K. (1989) Re-examination and further development of a precise and rapid dye method for measuring cell growth/cell kill. *Journal of Immunological Methods*, **119**, 203-210.
- [11] Coligan JE, Kruisbeek AM, Margulies DH, Shevach EM, Strober W. (1995) Current protocols in immunology. In *Related isolation procedures and functional assays*. Vol.1, Colico R. (Ed.). John Wiley & Sons, Inc., New York, NY, p 3.17.1.
- [12] Ribble D, Goldstein NB, Norris DA, Shellman YG. (2005) A simple technique for quantifying apoptosis in 96-well plates. *BMC Biotechnology*, **5**, 12, doi:10.1186/1472-6750-5-12 (<http://www.biomedcentral.com/1472-6750/5/12>).
- [13] Messmer UK, Reed UK, Brüne B. (1996) Bcl-2 protects macrophage from nitric oxide-induced apoptosis. *Journal of Biological Chemistry*, **271**, 20192-20197.
- [14] Burton KA. (1956) A study of the conditions and mechanism of the diphenylamine reaction for the colorimetric estimation of deoxyribonucleic acid. *Biochemical Journal*, **62**, 315-323.

C-Lactam Derivatives of Oleanolic Acid. The synthesis of C-lactam by Beckmann rearrangement of C-oxime

Barbara Bednarczyk – Cwynar

Department of Organic Chemistry, Poznan University of Medical Sciences, Grunwaldzka Str. 6,
60 – 780 Poznan, Poland

bcwynar@ump.edu.pl

Received: May 18th, 2011; Accepted: October 11th, 2011

Oleanolic acid, one of the most known triterpenes, was subjected to different chemical transformations within C-3 β -hydroxyl group, a double bond between C-12 and C-13, and a carboxyl function at C-17 in order to obtain new derivatives. The key compound consists of four six-membered rings (A, B, D, E) and one enlarged ring (C ring) containing a nitrogen atom and a carbonyl function – lactam. This type of derivative can be obtained by Beckmann rearrangement of the appropriate oxime. The lactam can be transformed into thiolactam with the use of Lavesson's reagent. The method is also presented for new derivatives synthesis, as well as their structure elucidation by spectroscopic means.

Keywords: triterpenes, oleanolic acid, azaderivatives of oleanolic acid, lactams, Beckmann rearrangement.

Oleanolic acid is a very-well tested compound that is broadly distributed in the plant kingdom. This acid is present, often together with other triterpenes, in a great number of plant sources [1]. The presence of a β -hydroxyl group at C-3, a double bond between carbons 12 and 13, and a carboxyl function at C-17 allows the possibility of performing a set of transformations that lead to many new derivatives.

For many years oleanolic acid has been subjected to numerous tests to determine its biological activity [eg. 2 and works cited within]. These efforts were focused on obtaining new types of derivatives with special pharmacological activity [eg. 3–7 and works cited therein].

Lactam derivatives are a special type of new compound within oleanolates. Lactams are internal amides. The nitrogen atom constitutes an element of a ring. The lactam system appears in numerous compounds with pharmacological activity, for example, psychotropic drugs, antiviral medicines and antibiotics.

There are only a few known derivatives of oleanolic acid with a lactam system. The first sythesis of this type of compound was published in 1971 [8]. The scientists developed a method of oleanolic acid azaderivatives synthesis with an expanded A-ring, pyridine as solvent, and tosyl chloride as a rearrangement activator. The reaction was carried out at 100°C for 3 hours. Two main products were separated, one with a nitrile function and cleaved A-ring (about 24%), and the second with a seven-membered lactam system (32%). Sundararamaiah *et al.* [9]

performed some experiments with the usage of phosphoryl chloride as a rearrangement agent in pyridine at either slightly elevated temperature (30 – 35°C) or at 130°C. Better results were obtained in the latter case and the yield of nitrile and lactam were 12% and 40%, respectively.

The next azaderivative of oleanolic acid was obtained in 1974 [10]. The new compounds contained two nitrogen atoms, one within a morpholide function, and the second within a lactam or nitrile system. The synthesis was performed in pyridine with the use of tosyl chloride as a rearrangement activator at room temperature. The nitrile was obtained in a yield of 74%, but the yield of lactam is unknown.

The synthesis of a six-membered lactam system within an oleanolic acid molecule is also known, but this compound is deprived of its C-23 and C-24 angular groups [11]. As a result of phosphorus pentachloride action on methyl oleanolate at -15°C, dehydration took place and a new bond between C-3 and C-5 was formed. Further transformation led to a product with a five-membered A-ring of the nor-type and with a C-3 carbonyl group. This compound, when treated with hydroxylamine hydrochloride, formed an oxime that by the action of phosphorus pentachloride was transformed into a lactam.

Oleanolic acid (**1**) used for transformations was isolated by us from a by-product residue obtained during production of mistletoe extract. The triterpenic acid was transformed into its methyl ester (**2**). Physico-chemical and spectral data of both compounds were in agreement with literature data [12].




Figure 1: Synthesis of C-lactam derivatives of oleanolic acid.

Reagents and conditions: I: $(\text{CH}_3)_2\text{SO}_4$, EtOH, NaOH, heating; II: $(\text{CH}_3\text{CO})_2\text{O}$, pyridine, rt.; III: *m*-CPBA, CH_2Cl_2 , rt; IV: $\text{NH}_2\text{OH} \times \text{HCl}$, CH_3COONa , EtOH, heating; V: POCl_3 , pyridine, rt; VI: Lavesson's reagent, C_6H_6 , heating.

Next, methyl oleanolate (**2**) was acetylated with 10 equiv. of acetic anhydride in anhydrous pyridine to give methyl 3-acetoxyoleanolate (**3**), the spectral data of which were in accordance with those in the literature [12]. On oxidation of **2** with 2 equiv. of 3-chloroperbenzoic acid in dry CHCl_3 at room temperature overnight, methyl 3-acetoxy-12,13-epoxyoleanolate was formed. The crude product was purified by column chromatography. Under acidic conditions, the 12,13-epoxide underwent hydrolysis and further oxidation to form the 12-oxo compound. Pure product (**4**) was obtained as oil that solidified after cooling. The crystallization from ethanol led to white needles, the physico-chemical and spectral data of which were in agreement with those in the literature [13].


The product of the above reaction was converted into the appropriate hydroxyimine compound by the action of hydroxylamine hydrochloride in ethanol. The total transformation of ketone **4** into oxime **5** finished after 3 hours of heating; the yield was about 80%. After crystallization from ethanol, 3 β -acetoxy-12-hydroxyiminoolean-28-oic acid methyl ester (**5**) formed white crystals.

The obtained 12-oxime (**5**) was subjected to Beckmann rearrangement in pyridine at room temperature with phosphoryl chloride as Lewis acid (rearrangement agent). As a result, a mixture of at least three compounds was obtained with one of them, the most polar, as the main product. The mixture was subjected to column chromatography and derivative **6** was isolated in about 70% yield.

In the ^1H NMR spectrum of the above compound the presence of a characteristic N-H signal, located at δ 5.53, was observed. The signal, a doublet, $J = 3.4$ Hz, indicated the coupling of the lactam proton with a single atom of hydrogen, which can be only that at C-13. It is stated on

the basis of this observation that the product of this Beckmann rearrangement of oxime **5** has the structure **6a**. If the product had the structure **6b**, the N-H proton would be coupled with two hydrogen atoms at C-11, and the signal for this lactam proton would be observed as a triplet. On the basis of the above evidence it can be stated that the starting oxime **5** was the *E* isomer. The structures of the less polar products was not established and so regiostructural tests were performed in order to establish them.

The lactam system in 3 β -acetoxy-12-oxo-12a-aza-C-homoolean-28-oic acid methyl ester (**6**) is not a reactive function. It forms the appropriate thioxoderivative, named thiolactam (**7**). The reaction is carried out in boiling benzene with Lavesson's reagent. After purification by column chromatography, the main product was obtained in a yield of about 88%.



In the IR spectrum of the newly obtained triterpene the alteration of the N-H signal was noticed. This was caused by the presence of a neighbouring thiocarbonyl group instead of the carbonyl one. The signal was observed at about 3395 cm^{-1} for thiolactam (for lactam: 3410 cm^{-1}). The signal indicating the presence of a thiocarbonyl function was found at 1445 cm^{-1} . In the ^1H NMR spectrum of product **7**, the thiolactam signal, observed at δ 7.77, had a higher value in comparison with the signal for a lactam system (δ 5.53). As in the case of the mother lactam, the signal for **7** had the form of a doublet. Comparison of the ^{13}C NMR spectra of thioazaderivative **7** and azaderivative

6 showed the change of the C-12 signal from δ 176.6 (lactam) to δ 208.2 (thiolactam). The molecular ion in the MS of the thiolactam was at m/z = 559.3, with an intensity of 100.00%, which confirmed the structure of the obtained product 7.

Experimental

General: The solvents used for experiments (pure or pure for analysis) were purchased from either Chempur or POCh; the other reagents were purchased from Sigma-Aldrich. TLC analysis (reactions progress and level of compounds purity) was conducted on HPTLC aluminum sheets (Merck, HPLC Alufolien Kieselgel 60 F245, Merck Art. 5547) with benzene: ethyl acetate mixtures in different ratios used as eluents. The chromatograms were visualised by spraying them with 10% ethanolic sulphuric acid solution and then heating at about 110°C for several minutes. Column chromatography was performed using Kieselgel 60; 0.063 – 0.200 mm (70 – 230 mesh), Merck. Melting points were determined using an open capillary method in a Kofler apparatus and are uncorrected. IR spectra were recorded using a Specord IR-75 spectrophotometer, for 0.5% mixtures of tested compounds and KBr. The position of absorption bands is given in cm^{-1} , exact to $\pm 5\text{cm}^{-1}$. NMR spectra for hydrogen (^1H) and carbon atoms (^{13}C) were recorded in CDCl_3 solutions using a Varian Gemini 300 VT apparatus, for frequencies of 300 MHz and 75 MHz, respectively, with Me_4Si as the internal standard. The values for chemical shifts are given in δ , exact to ± 0.01 ppm and ± 0.1 ppm, respectively; J values are given exact to ± 0.1 Hz. The multiplicity of signals in ^1H NMR spectra are marked as follows: **s** – singlet, **s/br/** – broad singlet, **d** – doublet, **t** – triplet, **dd** – doublet of doublets, **dt** – doublet of triplets. MS were recorded using an AMD 402 spectrometer with electroionisation. The m/z values are given exact to ± 0.1 . HREIMS were also recorded using an AMD 402 spectrometer; m/z values are given exact to ± 0.0001 . Elemental analyses (C, H, N) were performed with Perkin-Elmer 2400 CHN analyzer. The elucidation of the chemical structures was based on IR, ^1H NMR, ^{13}C NMR and MS analysis.

3 β -Acetoxy-12-oxoolean-28-oic acid methyl ester (4): To a solution of 3 β -acetoxyolean-12-en-28-oic acid methyl ester (**3**, 5.13 g, 10 mmol) in dry chloroform (150 cm^3), a dry solution of *m*-CPBA (3.45 g, 20 mmol) in chloroform (50 cm^3) was added and the resulting solution was left at room temperature in darkness for 2 days. The mixture was washed with 5% solutions of FeSO_4 , Na_2CO_3 , HCl and with water. The organic solution was dried with MgSO_4 , the solvent removed and the obtained solid (5.04 g) was subjected to column chromatography.

Yield: 3.84 g, 72.6%.

White needles (ethanol).

MP: 198 – 199°C.

R_f : 0.82 (C_6H_6 – AcOEt , 4:1), dark-yellow spot.

Spectral data and melting point agreed with literature data [14].

3 β -Acetoxy-12-hydroxyiminoolean-28-oic acid methyl ester (5): To a hot solution of 3 β -acetoxy-12-oxoolean-28-oic acid methyl ester (**4**, 5.28 g, 10 mmol) in ethanol (105 cm^3), hydroxylamine hydrochloride (3.47 g, 50 mmol) and anhydrous sodium acetate (6.56 g, 80 mmol) were added. The obtained mixture was refluxed for 3 h, cooled and poured into water (525 cm^3) slightly acidified with HCl. The resulted precipitate was filtered, washed with water and dried.

Yield: 4.24 g, 78.0%.

White needles (ethanol).

MP: 196 – 198.5°C.

R_f : 0.80 (C_6H_6 – AcOEt , 4:1), yellow-brown spot.

IR (KBr): 3430 (OH, N-OH), 1730 and 1710 ($\text{C}=\text{O}$, CH_3COO and COOCH_3), 905 (N-O, N-OH).

^1H NMR (300 MHz, CDCl_3): 8.05 (1H, s/br/, N-OH), 4.47 (1H, dt, J = 5.7 and 10.5 Hz, $\text{C}_3\text{-H}_a$), 3.67 (3H, s, COOCH_3), 2.85 (1H, dt, J = 3.3 and 11.0 Hz; $\text{C}_{18}\text{-H}_\beta$), 2.52 (1H, d, J = 3.7 Hz; $\text{C}_{13}\text{-H}$), 2.04 (3H, s, CH_3COO), 0.91, 0.91, 0.88, 0.87, 0.85, 0.85, 0.81 (7 x s, 7 x CH_3).

^{13}C NMR (75 MHz, CDCl_3): 178.5 (C-28), 170.8 (CH_3COO), 159.8 (C-12), 80.7 (C-3), 51.8 (COOCH_3), 48.1 (C-17), 43.3 (C-13), 21.3 (CH_3COO).

EIMS, m/z (%): 543.4 (49.6) M^{+} .

Anal. calcd for $\text{C}_{33}\text{H}_{53}\text{NO}_5$ (%): C = 72.89; H = 9.82; N = 2.58; found: C = 72.53; H = 9.86; N = 2.85.

3 β -Acetoxy-12-oxo-12a-aza-C-homoolean-28-oic acid methyl ester (6): To a solution of 3 β -acetoxy-12-hydroxyiminoolean-28-oic acid methyl ester (**5**, 5.44 g, 10 mmol) in dried pyridine (125 cm^3), phosphoryl chloride (2.8 cm^3 , 4.60 g, 30 mmol) was added dropwise with cooling. The obtained mixture was left at room temperature for one day and then poured into water (625 cm^3), slightly acidified with HCl. The resulting precipitate was filtered, washed with water, dried and subjected to column chromatography.

Yield: 3.94 g, 72.5%.

Long, colorless needles (ethanol).

MP: 289 – 290°C.

$[\alpha]_D$: -14.0 (c 0.58, CHCl_3).

R_f : 0.10 (C_6H_6 – AcOEt , 4:1), dark-pink spot.

IR (KBr): 3410 (N-H, lactam), 1730 and 1710 ($\text{C}=\text{O}$, CH_3COO and COOCH_3), 1670 ($\text{C}=\text{O}$, lactam).

^1H NMR (300 MHz, CDCl_3): 5.53 (1H, d, J = 3.4 Hz; N-H, lactam), 4.49 (1H, dd, J = 5.1 and 11.3 Hz; $\text{C}_3\text{-H}_a$), 4.01 (1H, t, J = 6.0 Hz; $\text{C}_{13}\text{-H}$), 3.73 (3H, s, COOCH_3), 2.04 (3H, s, CH_3COO), 1.00, 0.98, 0.96, 0.87, 0.86, 0.84, 0.82 (21H, 7 x s, 7 x CH_3).

^{13}C NMR (75 MHz, CDCl_3): 177.3 (COOCH_3), 176.6 (C-12), 170.6 (CH_3COO), 80.2 (C-3), 52.0 (COOCH_3), 50.4 (C-13), 47.3 (C-17), 21.3 (CH_3COO).

EIMS, m/z (%): 543.0 (27.7) M^{+} .

HREIMS: for M^{+} of $\text{C}_{33}\text{H}_{53}\text{NO}_5$ calcd.: 543.3924, found: 543.3939.

Anal. calcd. for $\text{C}_{33}\text{H}_{53}\text{NO}_5$ (%): C = 72.89; H = 9.82; N = 2.58, found: C = 73.14; H = 10.01; N = 2.72.

3 β -Acetoxy-12-thioxo-12a-aza-C-homoolean-28-oic acid methyl ester (7): To a hot solution of 3 β -acetoxy-12-oxo-12a-aza-C-homoolean-28-oic acid methyl ester (**6**, 2.72 g, 5 mmol) in dry benzene (130 cm³), Lavesson's reagent (2.02 g, 5 mmol) was added and the resulting solution was refluxed for 90 minutes. Next it was cooled, washed with a 5% solution of K₂CO₃ and water. The organic solution was dried with MgSO₄, the solvent removed and the obtained solid (2.63 g) subjected to column chromatography. Yield: 2.47 g, 88.3%. White, thick needles (ethanol). MP: 235 – 237°C. [α]_D: –53.6 (*c* 0.50, CHCl₃). R_f: 0.80 (C₆H₆ – AcOEt, 4:1), yellow-brown spot.

IR (KBr): 3395 (N-H, thiolactam), 1720 and 1710 (C=O, CH₃COO and COOCH₃), 1495 (C=S, thiolactam).

¹H NMR (300 MHz, CDCl₃): 7.77 (1H, d, *J* = 5.7 Hz; N-H, thiolactam), 4.49 (1H, dd, *J* = 4.7 and 11.7 Hz; C₃-H_a), 4.22 (1H, t, *J* = 6.3 Hz; C₁₃-H), 3.73 (3H, s, COOCH₃), 2.04 (3H, s, CH₃COO), 1.01, 0.99, 0.98, 0.90, 0.85, 0.84, 0.82 (21H, 7 x s, 7 x CH₃).

¹³C NMR (75 MHz, CDCl₃): 208.2 (C-12), 177.2 (C-28), 170.7 (CH₃COO), 80.2 (C-3), 54.4 (C-13), 52.0 (COOCH₃), 47.1 (C-17), 21.2 (CH₃COO).

EIMS, *m/z* (%): 559.3 (100.0) M⁺.

HREIMS: for M⁺ of C₃₃H₅₃NO₄S calcd.: 559.3695, found: 559.3696.

References

- [1] Yeung MF, Che ChT. (2009) A review on presence of oleanolic acid in natural products. *Natura Proda Medica*, **2**, 77–290.
- [2] Liu J. (2005) Pharmacology of oleanolic acid and ursolic acid. *Journal of Ethnopharmacology*, **100**, 92–94.
- [3] Couch RD, Browning RG, Honda T, Gribble GW, Wright DL, Sporn MB, Anderson AC. (2005) Studies on the reactivity of CDDO, a promising new chemopreventive and chemotherapeutic agent: Implications for a molecular mechanism of action. *Bioorganic and Medicinal Chemistry Letters*, **15**, 2215–2219.
- [4] García – Granados A, López PE, Melguizo E, Parra A, Simeo Y. (2004) Partial synthesis of C-ring derivatives from oleanolic and maslinic acids. Formation of several triene systems by chemical and photochemical isomerization processes. *Tetrahedron*, **60**, 1491–1503.
- [5] Hichri F, Jannet HB, Cheriaa J, Jegham S, Mighri Z. (2003) Antibacterial activities of a few prepared derivatives of oleanolic acid and of other natural triterpenic compounds. *Comptes Rendus Chimie*, **6**, 473–483.
- [6] Finlay HJ, Honda T, Gribble GW. (2002) Synthesis of novel [3,2-b]Indole fused oleanolic acids as potential inhibitors of cell proliferation. *ARKIVOC*, **12**, 38–46.
- [7] Honda T, Honda Y, Favaloro FGJr, Gribble GW, Suh N, Place EA, Rendi MH, Sporn MB. (2002) A novel dicyanotriterpenoid, 2-cyano-3,12-dioxooloana-1,9(11)-dien-28-onitrile, active at picomolar concentrations for inhibition of nitric oxide production. *Bioorganic and Medicinal Chemistry Letters*, **12**, 1027–1030.
- [8] Wrzeciono U, Turowska W, Kusnierkiewicz G. (1971) Azotowe pochodne trójtępenów. IV. 3,4-Seko- i A-homo-pochodne oleananu. *Roczniki Chemii (Polish Journal of Chemistry)*, **45**, 1233–1241.
- [9] Sundararamaiah T, Ramraj SK, Lakshman Rao K, Vimala Bai V. (1976) Synthesis of A-azatriterpenes-I: A-aza triterpenes from methyl oleanonate, methyl betulonate and lupenone. *Journal of Indian Chemical Society*, **53**, 664–665.
- [10] Yasue M, Sakakibara J, Kaiya T. (1974) Syntheses of nitrogen-containing triterpenes. III. Derivatives of betulinic acid and oleanolic acid. *Yakugaku Zasshi*, **94**, 1468–1474.
- [11] García – Granados A, López PE, Melguizo E, Moliz JN, Parra A, Simeó Y. (2003) Epoxides, cyclic sulfites, and sulfate from natural pentacyclic triterpenoids: theoretical calculations and chemical transformations. *Journal of Organic Chemistry*, **68**, 4833–4844.
- [12] Ahmad VU, Atta-ur-Rahman. (1994) *Handbook of Natural Products Data*. Volume 2. Pentacyclic Triterpenoids, Elsevier, Amsterdam – London – New York – Tokyo, 1–509.
- [13] Zhu YM, Shen JK, Wang HK, Cosentino LM, Lee KH. (2001) Synthesis and anti-HIV activity of oleanolic acid derivatives. *Bioorganic and Medicinal Chemistry Letters*, **11**, 3115–3118.
- [14] Okamoto I., Takeya T., Kagawa Y., Kotani E. (2000) Iron(III)picolinate-induced oxygenation and subsequent rearrangement of triterpenoid derivatives with hydrogen peroxide. *Chemical & Pharmaceutical Bulletin*, **48**, 120–125.

Analysis of Native Carotenoid Composition of Sweet Bell Peppers by Serially Coupled C₃₀ Columns

Daniele Giuffrida^{a*}, Paola Dugo^{b,c}, Giacomo Dugo^a, Germana Torre^b and Luigi Mondello^{b,c}

^aUniversità di Messina, Facoltà di Scienze MM. FF. NN., Dipartimento di Scienze degli Alimenti e dell'Ambiente, Contrada Papardo, Salita Sperone 31, 98166 Messina, Italy

^bUniversità di Messina, Facoltà di Farmacia, Dipartimento farmaco-chimico, Viale Annunziata, 98168, Messina, Italy

^cUniversity Campus Bio-Medico, Rome, Italy

dgiuffrida@unime.it

Received: June 29th, 2011; Accepted: September 26th, 2011

Serial coupled columns reversed-phase separations in high-performance liquid chromatography can be a useful tool for the analysis of complex real samples. The great difficulties found when analyzing complex carotenoid samples, due to the high natural variability of these compounds, as well as to the presence of carotenoid esters, are well documented. In the present contribution, the applicability of connecting two C₃₀ columns to increase significantly the separation power, resolution and peak capacity for the analysis of carotenoids in a complex carotenoid sample, like sweet bell peppers, has been shown for the first time. By using LC coupled to PDA/APCI-MS detectors, 56 different carotenoids have been detected in red sweet bell peppers. By using two serial coupled C₃₀ columns a peak capacity of 95.4 was obtained, compared with 73 achieved using a single column. Moreover, resolution greatly improved between different critical peaks when using two serial coupled C₃₀ columns, compared with a single column. Interestingly, free carotenoids, mono-esters and di-esters were quantitatively equally represented (around 33% for each different class) in red sweet bell pepper, showing, therefore, a value for the ratio of mono-esters/di-esters of around 1, which could be considered a parameter of typicality. Free β -carotene (12.6%), capsanthin-C14:0 (8.4%), and capsanthin-C12:0-C14:0 (8.9%) were the most abundant carotenoids in the three different classes of red sweet bell pepper. No carotenoid esters were detected in either yellow or green sweet bell peppers. The application of such methodology in the analysis of other complex carotenoid matrices could be a future objective of research.

Keywords: carotenoids, sweet bell peppers, liquid chromatography-DAD-APCI-MS.

Carotenoids are based on a C₄₀ tetraterpenoid skeleton which can undergo a high diversity of modifications, such as cyclization in one or both ends, hydrogenation, dehydrogenation, and addition of lateral groups, among others, resulting in an extremely wide group of compounds. Usually, these compounds are divided into two groups: hydrocarbons (commonly known as carotenes) and oxygenated compounds (generally named xanthophylls). To further increase the natural variability of these compounds, it has to be considered that the carotenoids can be present in nature as either free carotenoids or in a more stable form esterified with fatty acids, in the case of the oxygenated compounds. To simplify to some extent their analysis, a saponification procedure has been traditionally employed to release all the carotenoid esters and to analyze all these compounds in their free form. Although this saponification step acts also as a clean-up procedure, some drawbacks are found, mainly related to the formation of artefacts, as well as to the production of carotenoid degradation. Moreover, as a result of the saponification step, information on the native carotenoid composition of the studied samples is lost. Thus

a better approach to carotenoid content is through classifying plant materials depending on either a free or esterified xanthophylls profile. In fact, the carotenoid esters could be used as a marker of authenticity of food products and could be useful as a ripeness degree index; moreover, esters may enhance food product oxidative stability and may improve carotenoid bioavailability. Although esterification does not change the chromophore properties of the carotenoid molecules, it does modify the immediate molecular environment; thus, chemical activities may be altered depending on the kind of fatty acid bound to the xanthophylls. Carotenoids are an important kind of natural pigment that can be widely found in plant-derived food and products. Although these compounds have been traditionally used in the food industry as colorants, nowadays, they attract a great deal of attention since they have been described to possess several important functional properties, mainly antioxidant activity [1], as well as prevention of cardio vascular diseases [2,3], cancer [4] and macular degeneration [5]. These properties make these compounds ideal for the always increasing functional food industry, as well as promoting the

required and recommended amounts of carotenoids in the human diet. *Capsicum annuum* is a carotenogenic fruit: during ripening, the transformation of the chloroplast into chromoplast occurs. Chlorophylls disappear and more and new carotenoids are formed. *Capsicum* species uniquely have capsanthin-capsorubin synthase that synthesizes two red pigments, capsanthin and capsorubin. Moreover, esterification greatly increases during the fruit ripening process. *Capsicum* is one of the oldest and most popular vegetables and spices in the world. Considering the carotenoid profile among various *C. annuum* cultivars [6-16], the spicy ones have received greater attention than the non spicy cultivars; the native carotenoid profile in sweet bell peppers had not been previously investigated. Therefore, the aim of this study was to investigate the native carotenoid composition in sweet bell peppers by developing an analytical LC method which allowed the direct identification of the carotenoids in the samples, based on the use of serially connected C₃₀ columns coupled with DAD and APCI-MS detectors. High performance liquid chromatography has been selected as the analytical tool for a large number of applications, including carotenoid analysis. However, due to the great complexity of some natural samples containing this kind of compound, conventional LC could not have enough separation power. In this work, serial connection of two columns is proposed as an alternative to conventional LC. The applicability of connecting two C₃₀ columns to increase significantly the separation power, resolution and peak capacity for the analysis of carotenoids in sweet bell peppers has been demonstrated. Moreover, here we report the first investigation of the native carotenoid profile in sweet bell peppers at three different ripening stages: green, yellow and red (Figures 1, 2 and 3). As shown in Figure 3B, 56 different carotenoids have been detected in red sweet bell peppers, including many esters (for peak identification see Table 1). No carotenoid esters were detected in either yellow or green sweet bell peppers. The identification of these compounds was carried out combining the information provided by the two detectors employed (i.e., DAD and APCI-MS detectors) and the commercial standards available. As can be observed in Figure 3, the order of elution of the different compounds is highly dependent on the polarity and hydrophobicity of the molecules. Therefore, free xanthophylls elute before mono-esterified ones, and the di-esterified xanthophylls have longer retention times. By considering the different fragmentations in the APCI-MS producing different regioisomers, various xanthophyll di-esters were detected in red bell peppers.

As can be seen in Table 2, by using two serially coupled C₃₀ columns a peak capacity of 95.4 was obtained, compared with 73 using a single column. Moreover, the resolution values of some critical pairs (12-13, 29-30 and 32-33) were significantly improved by the coupling of the two C₃₀ columns.




Figure 1: Chromatograms (455 nm) of sweet bell green peppers using one C₃₀ column. Peak identification: 1. Luteoxanthin; 2. Lutein; 3. Zeaxanthin; 4. β-Carotene.




Figure 2: Chromatograms (455 nm) of sweet bell yellow peppers using one C₃₀ column. Peak identification: 1. Luteoxanthin; 2. Antheraxanthin; 3. Zeaxanthin; 4. Lutein; 5. Zeaxanthin; 6. β-cryptoxanthin; 7. β-carotene




Figure 3: Chromatograms (455 nm) of sweet bell red peppers using one C₃₀ column (A), and two C₃₀ columns (B). For peak identification see Table 1.

Interestingly, free carotenoids, mono-esters and di-esters were quantitatively equally represented (around 33% for each different class) in red sweet bell pepper, and the ratio of di-esters/mono-esters was around 1, which could be considered a typicality parameter of red peppers. β-Carotene (6%), lutein (20-30%) and zeaxanthin (2-5%) were the most abundant carotenoids present in yellow and green sweet bell peppers, whereas violaxanthin (2.2%) capsanthin (8.1%) and β-carotene (12.6%) characterized the free carotenoid fraction of sweet red bell peppers. Myristic acid (C14:0) was the most abundant fatty acid present as mono-ester with capsanthin (8.4%), capsorubin (3.1%) and antheraxanthin (2.7%) in red bell peppers. Lauric and myristic acids were mainly present in the di-esters of capsanthin and capsorubin in red bell peppers. In particular, the most abundant di-esters in red bell peppers were capsanthin-C12:0, C14:0 (8.9%), capsanthin-C14:0, C14:0 (7.1%) and capsorubin-C14:0, C14:0 (3.3%).

No carotenoid esters were detected in either green or yellow sweet bell peppers. Interestingly, Minguez-Mosquera [6]

Table 1: UV-Vis, MS data and identification of the different carotenoids separated in sweet bell red pepper using two serially connected C₃₀ columns.

No.	Peak identification	λ_{\max} (nm)	[M + H] ⁺
1	Neoxanthin	416,440,469	601,583
2	n.i.	419,446,467	601,583
3	β -Apo-10-carotenal	436	376,361
4	Violaxanthin	418,441,470	601,583
5	Luteoxanthin	400,422,448	601,583
6	n.i.	418,440,467	613,595
7	n.i.	400,424,448	601,583
8	n.i.	400,423,448	615,597
9	Capsanthin-5,6-epoxide	469,486	601,583,565
10	Anteraxanthin	424s,445,474	585,567
11	n.i.	357,448,469	585,567
12	Mutatoxanthin A	406,429,451	585,567,549
13	n.i.	457s,479,507	601,583,567,491
14	Mutatoxanthin B	406,429,452	585,567,549
15	Capsanthin	474	585,567,479
16	Zeaxanthin	427,451,477	569,551,476
17	Phytofluene	332,348,367	543
18	β -Cryptoxanthin	426,451,477	553,535
19	Anteraxanthin-laureate	422,447,474	767,749,567
20	n.i.	400,424,448	875,597
21	n.i.	424,443,469	875,597,565
22	n.i.	451,467	823,583
23	n.i.	425s,447,467	877,823,597
24	n.i.	466	879,599,583
25	n.i.	400,424,448	879,793,597
26	Anteraxanthin-myristate	424,446,474	795,777,567
27	n.i.	425,446,474	876,777
28	Mutatoxanthin-laureate	408,425,452	767,749,567
29	Capsorubin-myristate	479	811,583,565
30	β -carotene	452,478	537
31	Capsanthin-laureate	474	767,567,549
32	Anteraxanthin-myristate	425,447,474	795,777,567
33	n.i.	424s,448,471	853,795,567
34	Mutatoxanthin-myristate	406,429,452	795,777,567
35	Capsanthin-myristate	474	795,777,567
36	n.i.	460	795,567,549
37	n.i.	426,451,474	879,684,533
38	n.i.	424,443,469	961,821,547
39	n.i.	466,472	823,567,549
40	β -Cryptoxanthin-laureate	426,451,477	735,535
41	n.i.	469	765,549
42	β -Cryptoxanthin-myristate	425,452,479	765,535
43	Capsanthin-di-laureate (C12:0,C12:0)	473	949,749,549
44	Capsorubin-laureate-myristate (C12:0,C14:0)	479	993,793,765,565
45	n.i.	428,451,470	1021,793,565
46	Capsanthin-laureate-myristate (C12:0,C14:0)	474	977,777,749,549
47	Capsorubin-di-myristate (C14:0,C14:0)	481	1021,793,565
48	Mutatoxanthin-laureate-myristate (C12:0,C14:0)	406,429,454	977,793,765,565
49	Zeaxanthin-laureate-myristate (C12:0,C14:0)	425,451,478	961,761,733,533
50	Capsanthin-di-myristate (C14:0,C14:0)	474	1005,777,549
51	n.i.	469	1005,777,749,549
52	Capsorubin-myristate-palmitate (C14:0-C16:0)	479	1049,821,793,565
53	Zeaxanthin-di-myristate (C14:0,C14:0)	427,452,477	989,761,533
54	Capsanthin-myristate-palmitate (C14:0,C16:0)	474	1033,805,777,549
55	n.i.	469	805,777,549
56	n.i.	469	805,713,549,551

Table 2: Values of Peak capacity (Pc) and resolution (Rs) between selected peaks for the analysis of sweet bell red peppers using two different set ups.

Set Up	Pc	Rs (12-13)	Rs (29-30)	Rs (32-33)
1 x C30	73.0	0.23	0.66	0.33
2 x C30	95.4	0.85	1.28	0.87

reported that in the yellow spicy *C. annuum* cultivar Bola, 50% of the carotenoids were esterified, and that at the fully ripened red stage the percentages of the free carotenoid pigments and the partially and totally esterified forms of these were 21.3%, 35.6%, and 43.1%, respectively, and, therefore, different from the percentages reported in this work for red sweet bell peppers, where the three fractions were equally represented. The increase in xanthophyll esterification during ripening reported in this work is in agreement with the report by Hornero-Mendez [8] for various cultivars of spicy *C. annuum* cultivars. This process has been related to a phenomenon intimately linked with and inherent to the degeneration of chloroplast and formation of chromoplast. Such a phenomenon might be the result of hydrophobicity requirements on the part of the carotenoid, so that, with all its hydroxyl groups esterified, it will be included more readily in the lipid matrix of chromoplast membranes and organelles (plastoglobules) [8]. Schweiggert *et al.* [9] reported that capsanthin and β -carotene were the main free carotenoids in spicy red peppers, in agreement with the results reported in this work for sweet red bell peppers, and they reported also a similar esters profile for spicy red pepper pods, although the sweet bell red peppers analyzed in this work showed a higher degree of esterification with capsanthin rather than with capsorubin. Apart from a very early report by Gregory *et al.* [11] on the carotenoid esters in sweet red bell peppers carried out by gas chromatographic analysis of the transesterified fatty acids obtained after saponification and which reported some generic tentative identifications, the work here reported is the first direct study of the native carotenoid composition of sweet bell peppers using a liquid chromatographic methodology. In fact, previous works on the carotenoid composition of sweet bell peppers [12-16] were carried out after a saponification step.

Although the application of the methodology reported in this work is instrumentally quite simple, it has been demonstrated to be a valid way to further improve resolution and efficiency in LC.

Experimental

Samples and chemicals: Fresh samples of green, yellow and red sweet bell peppers were supplied by a local producer. Carotenoid standards, namely, β -carotene, lutein, β -cryptoxanthin, zeaxanthin, capsanthin and lutein-di-palmitate (C_{16:0}, C_{16:0}) were purchased from Extrasynthese (Genay, France). All the carotenoid standards were stored protected from light at -18°C. All the solvents used, namely, methanol, methyl tert-butyl ether (MTBE) and water, were HPLC grade and purchased from Sigma-Aldrich (Milan, Italy). BHT (butylatedhydroxy

toluene) and potassium hydroxide were obtained from Sigma-Aldrich (Milan, Italy).

Carotenoid extraction: The sweet bell pepper samples (200 g) were homogenized, and the carotenoids extracted 4 times with methanol/ethyl acetate/light petroleum (1:1:1). The upper phase was kept and ca. 2 mg of BHT was added prior to evaporation under vacuum until dryness. The dry residue was then resuspended in a given volume of MTBE/methanol (1:1) and stored protected from light at -18°C until use.

LC-DAD/APCI-MS analyses: To carry out the analyses a Shimadzu HPLC instrument (Shimadzu, Milan, Italy) was employed including a SCL-10A-VP system controller, two LC-10AD-VP pumps and a SPD-10Avp diode array detector. Besides a Shimadzu mass spectrometer LCMS-2010 equipped with an APCI interface in the positive ion mode was installed in parallel. The APCI parameters were set as follows: probe voltage (kV), 4.5; probe temperature, 400°C; block temperature, 200°C; CDL temperature, 250°C; Q array voltage, 20 and 80 V; gas flow, 2.5 L/min.

Data acquisition and system control was performed by the LCMSsolutionver 3.30 software (Shimadzu). The injections were made manually through a Rheodyne injection valve (injection volume 20 µL). The C₃₀ columns employed consisted of two YMC 30 analytical columns (YMC Europe, Schermbeck, Germany) with 5 µm C₃₀ reversed-phase material (250 × 4.6 mm I.D.), including a pre-column YMC 30 (S-5 µm, 10 × 4.0 mm I.D.). The HPLC solvent systems were (A) methanol/MTBE/water (83:15:2, v/v/v) and (B) methanol/MTBE/water (8:90:2, v/v/v) used following a linear gradient depending on the use of one or two serial coupled columns. Gradient for the separation with 1 x C₃₀ column: 0-20 min, 0% B; 20-160 min 100% B; then reconditioning. Gradient for the separation with 2 x C₃₀ columns: gradient times were doubled keeping unchanged the B%. The flow rate employed was 1 mL/min and the chromatograms were recorded at 450 nm and the UV-Vis spectra were recorded in the range from 250 to 600 nm (sampling rate: 12.5 Hz; time constant: 0.64 s). The column oven temperature was 40°C.

References

- [1] Beutner S, Bloedorn B, Frixel S, Hernandez Blanco I, Hoffman T, Martin HD, Mayer B, Noack P, Ruck C, Schmidt M, Schulke I, Sell S, Ernst H, Haremza S, Seybold G, Sies H, Stahl W, Walsh R. (2001) Quantitative assessment of antioxidant properties of natural colorants and phytochemicals: carotenoids, flavonoids, phenols and indigoids. The role of β-carotene in antioxidant functions. *Journal of Science Food & Agriculture*, **81**, 559-568.
- [2] Arab L, Steck S. (2000) Lycopene and cardiovascular disease. *American Journal of Clinical Nutrition*, **71**, 1691S-1695S.
- [3] Rao AV, Rao RG. (2007) Carotenoids and human health. *Pharmaceutical Research*, **55**, 207-216.
- [4] Nishino H, Tokuda H, Satomi Y, Masuda M, Bu P, Onozuka M, Yamaguchi S, Okuda Y, Takayasu J, Tsuruta J, Okuda M, Ichiishi E, Murakoshi M, Kato T, Misawa N, Narisawa T, Takasuka N, Yano M. (1999) Cancer prevention by carotenoids. *Pure & Applied Research*, **12**, 2273-2278.
- [5] Snodderly M D. (1995) Evidence for protection against age-related macular degeneration by carotenoids and antioxidant vitamins. *American Journal of Clinical Nutrition*, **62**, S1448-S1461.
- [6] Minguez-Mosquera MI, Hornero-Mendez D. (1994) Changes in carotenoid esterification during the fruit ripening of *Capsicum annum* Cv. Bola. *Journal of Agricultural and Food Chemistry*, **42**, 640-644.
- [7] Markus F, Daoood HG, Kapitany J, Biacs PA. (1999) Changes in the carotenoid and antioxidant content of spice red pepper (paprika) as a function of ripening and some technological factors. *Journal of Agricultural and Food Chemistry*, **47**, 100-107.
- [8] Hornero-Mendez D, Minguez-Mosquera MI. (2000) Xanthophyll esterification accompanying carotenoid overaccumulation in chromoplast of *Capsicum annum* ripening fruits is a constitutive process and useful for ripeness index. *Journal of Agricultural and Food Chemistry*, **48**, 1617-1622.
- [9] Schweiggert U, Kammerer DR, Carle R, Schieber A. (2005) Characterization of carotenoids and carotenoid esters in red pepper pods (*Capsicum annum* L.) by high-performance liquid chromatography/atmospheric pressure chemical ionization mass spectrometry. *Rapid Communications in Mass Spectrometry*, **19**, 2617-2628.
- [10] Hernandez-Hervert D, Sayago-Ayerdi SG, Goni I. (2010) Bioactive compounds of four hot pepper varieties (*Capsicum annum* L.), antioxidant capacity, and intestinal bioaccessibility. *Journal of Agricultural and Food Chemistry*, **58**, 3399-3406.
- [11] Gregory GK, Chen TS, Philip T. (1987) Quantitative analysis of carotenoids and carotenoid esters in fruits by HPLC: red bell peppers. *Journal of Food Science*, **52**, 1071-1073.
- [12] Sun T, Xu Z, Wu CT, Janes M, Prinyawattakul W, No HK. (2007) Antioxidant activities of different colored sweet bell peppers (*Capsicum annum* L.). *Journal of Food Science*, **72**, S98-S102.
- [13] de Azevedo-Meleiro CH, Rodriguez-Amaya DB. (2009) Qualitative and quantitative differences in the carotenoid composition of yellow and red peppers determined by HPLC-DAD-MS. *Journal of Separation Science*, **32**, 3652-3658.
- [14] Matsufuji H, Ishikawa K, Nunomura O, Chino M, Takeda M. (2007) Anti-oxidant content of different coloured sweet peppers, white, green, orange and red (*Capsicum annum* L.). *International Journal of Food Science and Technology*, **42**, 1482-1488.
- [15] Guil-Guerrero JL, Martinez-Guirado C, Rebolloso-Fuentes M, Carrique-Perez A. (2006) Nutrient composition and antioxidant activity of 10 pepper (*Capsicum annum*) varieties. *European Food Research and Technology*, **224**, 1-9.
- [16] Marin A, Ferreres F, Tomas-Barberan FA, Gil MI. (2004) Characterization and quantification of antioxidant constituents of sweet pepper (*Capsicum annum* L.). *Journal of Agricultural and Food Chemistry*, **52**, 3861-3869.

New Antifungal Cholestane and Aldehyde Derivatives from the Red Alga *Laurencia papillosa*Walied M. Alarif^{a*}, Sultan S. Al-Lihaibi^a, Ahmed Abdel-Lateff^{b#} and Seif-Eldin N. Ayyad^c^aDepartment of Marine Chemistry, Faculty of Marine Sciences, King AbdulAziz University, P.O. Box 80207, Jeddah 21589, Saudi Arabia^bDepartment of Health Information Technology, Jeddah Community College, King AbdulAziz University, P.O. Box 80283, Jeddah 21589, Saudi Arabia^cDepartment of Chemistry, Faculty of Science, King AbdulAziz University, PO. Box 80207, Jeddah 21589, Saudi Arabia[#]Permanent address: Department of Pharmacognosy, Faculty of Pharmacy, Minia University, Minia 61519, Egypt

walied1737@yahoo.com


Received: July 2nd, 2011; Accepted: October 20th, 2011

The chloroform/methanol extract of the red alga, *Laurencia papillosa*, collected from the Red Sea in Saudi Arabia, was found to contain two cholestane derivatives: 3 α , 6 α -dihydroxy-5 β -cholestan-12-one (**1**) and the known, 6 β -hydroxycholest-4-en-3-one (**2**), which was isolated separately in a pure form for the first time. In addition to these compounds, a new aldehyde derivative, (*E*)-2-[(*E*) tridec-2-en-2-yl] heptadec-2-enal (**3**), was isolated. The structures of all compounds were established based on extensive spectroscopic (1D and 2D NMR, UV, IR) and mass spectrometric studies. All compounds, except **2**, were tested for their antifungal activity. Significant activities were associated with **1** and **3** against *Candida albicans*, *Aspergillus fumigatus*, and *A. flavus*.

Keywords: Red alga, *Laurencia papillosa*, cholestane derivatives, aldehyde, antifungal activity.

Marine organisms have historically provided a rich source of structurally diverse, biologically active secondary metabolites [1a-1d]. The genus *Laurencia* (family Rhodomelaceae, order Ceramiales) possesses nearly 140 species worldwide are considered as a rich source of bioactive metabolites [2a-2c]. *L. papillosa* C. Agardh is a source of different substances such as steroids [3a-3c] and an unsaturated fatty acid aldehyde [4a]. A number of long chain unsaturated fatty acid aldehydes with antibiotic activity were reported from the fungus *Cladosporium* sp., obtained from intertidal marine sediment [4b]. It was clear from the literature that steroidal and aldehydic compounds have antimicrobial activities. On this basis, *L. papillosa*, collected from the water around El-Shuaiba, Saudi Arabia, was extracted with chloroform and methanol (1:1) and the extract fractionated by NP-silica gel column chromatography, followed by preparative thin-layer chromatography to yield a new cholestane derivative, 3 α ,6 α -dihydroxy-5 β -cholestan-12-one (**1**), and a known steroid, 6 β -hydroxycholest-4-en-3-one (**2**). In addition, a new aldehydic derivative {(2*E*)-2-(tridec-2-en-2-yl) heptadec-2-enal} (**3**) was also isolated.

The molecular formula of **1** was established as C₂₇H₄₆O₃ by EI-MS {*m/z*: 418 [C₂₇H₄₆O₃]⁺ and 400 [M- H₂O]⁺}. This result was validated by HRFABMS (*m/z* 441.3357

**Figure 1:** Structures of isolated compounds **1-3**.

[M+Na]⁺). The ¹³C NMR spectra (¹H decoupled and DEPT) of **1** showed 27 resonances attributable to 5 X CH₃, 10 X CH₂, 9 X CH, and 3 X C (Table 1). One of the five elements of unsaturation, as indicated by the molecular formula of **1**, deduced to be a carbonyl group, appeared at δ_c 212.8 with the absence of any olefinic proton or any double bond signal. The IR spectrum showed a ν (C-H)_{Aliphatic} vibration band at 2930 cm⁻¹. The spectrum also displayed two strong bands at 3450 and 1706 cm⁻¹ assigned to ν (OH) and a six membered ring ν (C=O) [4c], respectively; the molecule thus has four rings. As the ¹H and ¹³C NMR spectroscopic data enabled all but two of the hydrogen atoms within **1** to be accounted for, it was evident that the remaining two protons were present as part of a hydroxyl function. After association of all the protons with directly bonded carbons via 2D NMR (HMQC)

Table 1: ^1H (CDCl₃, 600 MHz) and ^{13}C NMR (CDCl₃, 150 MHz) spectral data of compound **1**^a.

P	^{13}C	H (m, J Hz)	P	^{13}C	H (m, J Hz)
1a	41.3	1.67 (1H, m) ^{b, c}	13	53.8	
1b		1.32 (1H, m)	14	56.4	1.26 (1H, m)
2a	31.7	1.77 (1H, m)	15	27.2	1.66 (2H, m)
2b		1.36 (1H, m)	16	27.6	1.76 (2H, m)
3	65.4	4.18 (1H, m)	17	53.1	1.22 (1H, m)
4a	36.1	2.02 (1H, m)	18	12.0	0.68 (3H, s)
4b		1.67 (1H, m)	19	12.3	0.74 (3H, s)
5	42.0	1.64 (1H, brs)	20	38.0	1.36 (1H, m)
6	73.9	3.61 (1H, dd, 12, 1.8)	21	21.1	0.93(3H, d, 6.6)
7a	39.5	1.71 (1H, dd, 12, 3.0)	22a	40.0	1.32 (1H, m)
7b		1.44 (1H, m)	22b		1.10 (1H, m)
8	38.0	1.51 (1H, m)	23a	24.0	1.33 (1H, m)
9	51.7	1.98 (1H, ddd, 13.3, 4.5, 1.8)	23b		1.13(1H, m)
10	35.9		24	29.7	1.12 (1H, m)
11a	46.9	2.31(1H, dd, 13.0, 4.5)	25	28.2	1.39 (1H, m)
11b		2.03(1H, d, 13.0)	26	23.0	0.90(3H, d, 6.6)
12	212.8		27	22.5	0.91(3H, d, 6.6)

All assignments are based on 1D and 2D NMR measurements (HMBC, HMQC, COSY). ^b Implied multiplicities as determined by DEPT (C= s, CH = d, CH₂ = t, CH₃ = q). ^c J in Hz.

spectral measurements, it was possible to deduce the structure of **1** by interpretation of the ^1H - ^1H COSY and ^1H - ^{13}C HMBC spectra. From the ^1H - ^1H COSY spectrum of **1**, ^1H - ^1H spin system between H₂-1 and H₂-2, H₂-2 and H-3, H-3 and H₂-4, H₂-4 and H-5, H-5 and H-6, H-6 and H₂-7, H₂-7 and H-8, and H-8 and H-9 were observed. Long-range C-H correlations observed between the resonances of H-9 and those of C-1, C-5, C-7, C-8 and C-10; between H₂-1 and C-2, C-3, C-5, C-9 and C-19; between H-3 and C-1 and C-5; between H-6 and C-8 and C-10 established that rings A and B were fused together. In the ^1H - ^1H COSY spectrum of **1**, a ^1H - ^1H spin system between H₂-11 and H-9 indicated that C-11 was attached to C-9. A CH long-range coupling between H-9 and C-8, C-10, C-11, C-14 and C-19 were observed. HMBC correlations, this time between H-14 and C-8, C-13, C-15, and C-18, between H₃-18 and C-12, C-13, C-14, and C-17 were observed. Long-range C-H correlations observed between the resonances of C-12 and H₂-11, H-14 were observed. Ring C was established by deduction through the previous data. From the ^1H - ^1H COSY spectrum of **1**, a ^1H - ^1H spin system between H₂-15 and H-14 and H₂-16, and between H₂-16 and H-17 were observed indicating a C-C bond between C-14 to C-15, C-15 to C-16, and C-16 to C-17. Ring D was established based on ^1H - ^1H COSY and ^1H - ^{13}C HMBC spectra. The main skeleton of **1** was established as a steroidal derivative. From the ^1H - ^1H COSY spectrum of **1**, a ^1H - ^1H spin system between H₃-26 and H-25 and between H-25 and H₃-27 indicated a forked tail isopropane derivative [4d]. The positions of the two hydroxyl groups were determined by examination of the ^1H and ^{13}C NMR, ^1H - ^1H COSY, HSQC and HMBC spectra. From the ^1H - ^1H COSY spectrum, correlations were observed between H-25 and H₂-24, between H₂-24 and H₂-23, between H₂-23 and H₂-22, and between H₂-22, H₂-20 and H₂-21, which led to establishing the side chain as an iso-octane derivative. This side chain is attached to the steroidal nucleus between C-17 and C-20 based on HMBC correlations between H₃-21 and C-17, C-20 and C-22. A proton at δ 4.18 appears as a multiplet at position

3, with equatorial orientation (hydroxyl group occupies α -position). The proton at δ 3.61, appearing as a doublet, with *J* values of 12 and 1.8 Hz, was located at position 6 with axial orientation (OH-6 occupies α -orientation). The weak or zero chemical shift effect of the hydroxyl group located at C-6 on Me-19 at δ 0.74 is further evidence for the α -position of OH-6. The proton at position 5 was found to have a β -orientation, owing to the signal at δ 1.64 (brs) which implies an equatorial orientation for H-5. The configuration of the OH at C-6 is equatorial downward, a deduction which led to the axial orientation of H-5. The *cis*-junction between rings A and B is responsible for the strain on both rings. On this basis, the chemical shifts of C-1 and C-11 appeared more downfield than the normal values. The configuration of 3-OH is alpha based on the ^{13}C chemical shift (δ 65.4) [5a]. The ^{13}C chemical shifts of C-1 and C-11 are downfield from the normal ones, based on the strain that appears to be due to the *cis*-junction between rings A and B. Also the α -OH located at C-6 was confirmed by the correlation between H-8 and C-6, which is in good agreement with the ^{13}C chemical shift of C-6 (δ 73.9) [5a,5b]. A literature survey indicated that **1** is a new cholestanone derivative and was named as 6 α -dihydroxy-5 β -cholestan-12-one.

The structure of **2** was based on the molecular formula of C₂₇H₄₄O₂, which was abstracted from the EI-MS: *m/z* 400 [C₂₇H₄₄O₂]⁺ and 382 [M-H₂O]⁺. After extensive study of the ^1H and ^{13}C NMR spectral data of **2** (see experimental part), it was clear that these matched those published for the steroidal derivative; 6 β -hydroxycholest-4-en-3-one, isolated from the red alga *Hypnea musciformis* as non solvable mixture with 6 β -hydroxycholest-4, 22-dien-3-one [6a]. A computer survey, including Science Finder, indicated that **2** is published here for the first time in a pure form, and the complete physical properties are presented.

Compound **3** had the molecular formula C₃₀H₅₆O, derived from the high-resolution mass measurement of the parent ion, which implied three degrees of unsaturation. The IR bands at 2730, 1680 and 1640 ν_{max} were characteristic of an α , β -unsaturated aldehyde. This conjugation was supported by the ^1H NMR spectrum, which showed an aldehyde proton at δ 9.32 and one vinyl proton at δ 6.44 (H-3, *t*, *J* = 7 Hz). ^1H NMR spectral data also revealed the presence of an olefinic proton at δ 5.33 (H-2', *m*), two primary methyls at δ 0.88 (6 H, *t*, *J* = 6.6 Hz), four methylene protons at δ 2.20-2.35 (4 H, *m*, 4, 3'), four methylene protons at δ 1.50 (4H, *m*, 5, 4'), olefinic methyl protons at δ 1.61 (3 H, *m*), and a huge methylene envelope at δ 1.27 (cf Experimental). The ^{13}C NMR and DEPT spectral data of **3** were supportive of the above analysis, showing an aldehyde group at δ_c 195.4 (C-1, *s*), two double bonds [δ_c 143.7 (C-2, *s*), 155.5 (C-3, *d*), 129.6 (C-1', *s*), 130.1 (C-2', *d*)], aliphatic methylene carbons at δ_c 23.0-31.0, two methyls at δ_c 14.2, and an olefinic methyl at δ_c 22.0. Thus, three degrees of unsaturation were accounted for as an aldehydic group and two double bonds, which are *trans* oriented owing to the values of the

chemical shifts of the allylic methylene at $\delta_c > 30$, and the J values, which are similar to those published [6b]. Hence **1**, had an acyclic α , β -unsaturated aldehyde with two fatty acid chains terminating at methyl groups, in which the length of both chains was determined by MS, and ^1H and ^{13}C NMR spectra. The structure connectivity of **3** was concluded by interpretation of its ^1H - ^1H COSY cross peaks, which showed that H-3 at δ 6.44 correlated with H₂-4 at δ 2.35; H-2' at δ 5.33 correlated with H₂-3' at δ 2.22; there was no observed correlation between H-3 and H-2'. The structure was unambiguously indicated as **3** by studying the HMBC spectrum in which H-1, resonating at δ 9.32, correlates with C-2 at δ_c 143.7, C-3 at δ_c 155.5 and C-1' at δ_c 129.6; also the olefinic proton H₁-3, resonating at δ_c 6.44, correlates with C-2 and C-4 at δ_c 31.0; the CH₃ protons resonating at δ_{H} 1.60 correlate with both C-1', C-2 and C-2' at δ_c 130.1. The above correlations denoted that the CHO group is connected to the quaternary carbon C-2, (δ_c 143.7, which in turn is connected to both C-3 and C-1', and the olefinic methyl is attached to C-1'; other correlations are shown in figure 2. From the above, compound **3** was assigned the structure (*E*)-2-((*E*) tridec-2-en-2-yl) heptadec-2-enal.




Figure 2: Selected HMBC correlations of **3**

The antifungal activity of all compounds, except **2**, was assessed using the microtiter broth dilution method for yeast susceptibility testing. Compound **1** had significant activity against *Candida albicans* with a MIC of 2000 $\mu\text{g/mL}$. Although **1** showed no activity against *A. fumigatus* and *A. flavus*, **3** had significant activity against these with MIC of 200 and 1000 $\mu\text{g/mL}$, respectively. The positive control, ketoconazole, showed potent activity against *C. albicans* and *A. fumigatus* with a MIC of 40 $\mu\text{g/mL}$, and 625 $\mu\text{g/mL}$ towards *A. flavus*.

Experimental

General: Optical rotations, ATAGO POLAX-L 2 polarimeter; GC/MS, Shimadzu-QP 2010 spectrometer; TLC was performed on silica gel (Kieselgel 60, F₂₅₄) of 0.25 mm layer thickness. Compounds were detected by using ethanol /sulfuric acid as spray reagent.

Algal sample: *Laurencia papillosa* was collected in June 2009, off the Saudi Arabian Red Sea Coast at Jeddah. A voucher sample (JAD 03050) is deposited at the Marine Chemistry Department, King Abdulaziz University, Jeddah, Saudi Arabia.

Extraction and isolation: The air dried algal material (200 g) was extracted with equal volumes of a mixture of light petroleum, chloroform and methanol (2 X 6 L, 24 h for each batch) at room temperature. The extract was

concentrated under reduced pressure to obtain 6 g residue. The residue (6 g) was homogenized with a small amount of silica gel (50 g) and poured onto the top of the column that was packed in light petroleum (40-60°C). The fractions were collected (50 mL each) employing a gradient elution from light petroleum to diethyl ether and from diethyl ether to ethyl acetate. The similar fractions were pooled together according to TLC pattern into three pools (P-A, P-B and P-C), employing 50%-sulfuric acid in methanol as spraying reagent. P-A was purified by PTLC-Silica gel, using light petroleum-diethyl ether (9:1), yielded **3**. P-B was purified by PTLC-Silica gel, using benzene: ethyl acetate (4:1) to give **2**. Finally, P-C was purified by PTLC-Silica gel, using benzene-ethyl acetate (5: 2) and re-chromatographed on Sephadex LH-20 using a mixture of MeOH-CHCl₃ (8:2), and finally purified by preparative TLC-silica gel employing benzene-ethyl acetate (6: 4), yielded **1**.

3 α , 6 α -Dihydroxy-5 β - cholestan-12-one (**1**)

White solid (7 mg, 0.0028 %).

MP: 170°C.

$[\alpha]_{\text{D}}^{18}$: +93 (*c* 0.1, CHCl₃).

R_f : 0.48 (benzene/EtOAc, 6:4).

IR (neat) ν_{max} : 3450, 2930, 1706 cm^{-1} .

^1H and ^{13}C NMR (CDCl₃): Table 1.

HRFAB MS: (m/z 441.3357 [M+Na]⁺). Calcd. 441.3345, EI-MS m/z : 418 [M]⁺, 400 [M- H₂O]⁺, 385 [M-CH₃-H₂O]⁺, 318 (100) [M-C₇H₁₆]⁺.

6 β -Hydroxycholest-4-en-3-one (**2**)

White solid (2 mg, 0.0008 %).

MP: 119-120°C.

$[\alpha]_{\text{D}}^{18}$: +29 (*c* 0.03, CHCl₃).

R_f : 0.35 (*n*-hexane/EtOAc, 7:3).

IR (KBr): 3635, 3520, 1685 cm^{-1} .

UV (MeOH): strong absorption at λ_{max} 237 nm.

^1H NMR (CDCl₃), δ (ppm): 0.85 (d, $J=6.6$ Hz, 3H, CH₃-26), 0.87 (d, $J=6.6$ Hz, 3H, CH₃-27) 0.92 (d, $J=6.6$ Hz, 3H, CH₃-21), 0.74 (s, 3H, CH₃-18), 1.38 (s, 3H, CH₃-19), 1.52 (m, 1H, H-25), 1.28 (m, 2H, H-24), 1.16 (m, 2H, H-23), 1.24 (m, 2H, H-22), 1.35 (m, 1H, H-20), 1.12 (m, 1H, H-17), 1.28 (m, 1H, H-16), 1.24 (m, 2H, H-15), 1.03 (m, 1H, H-14), 1.12 (m, 2H, H-12), 1.51 (m, 2H, H-11), 1.33 (m, 1H, H-9), 1.97 (m, 1H, H-8), 2.54-2.00 (m, 2H, H-7), 4.35 (t, $J=2.8$ Hz, 1H, H-6), 5.82 (s, 1H, H-4), 2.40 (m, 2H, H-2), 2.05-1.60 (m, 2H, H-1).

^{13}C NMR (CDCl₃), δ (ppm): 37.0 (C-1), 34.4 (C-2), 200.5 (C-3), 126.3 (C-4), 168.5 (C-5), 73.3 (C-6), 38.5 (C-7), 29.7 (C-8), 53.6 (C-9), 37.9 (C-10), 21.0 (C-11), 39.5 (C-12), 42.2 (C-13), 55.8 (C-14), 24.1 (C-15), 56.1 (C-17), 12.1 (C-18), 19.5 (C-19), 32.8 (C-20), 18.7 (C-21), 38.0 (C-22), 24.1 (C-23), 39.5 (C-24), 28.1 (C-25), 22.5 (C-26), 22.5 (C-27).

EIMS m/z : 400 [M, C₂₇H₄₄O₂]⁺, 385 [M-CH₃]⁺, 370 [M-CH₃-H₂O]⁺, 339 [M-C₃H₇-H₂O]⁺, 269 [M-C₈H₁₇-H₂O]⁺, 199 [M-C₁₃H₂₇-H₂O]⁺.

(E)-2-((E) Tridec-2-en-2-yl) heptadec-2-enal (3)

White residue (10 mg, 0.005%).

R_f: 0.70 (light petroleum/diethylether, 9:1).

IR (neat): ν_{\max} 2930, 1680, 1640 cm⁻¹.

UV (Et₂O): 259 nm.

¹H (600 MHz, CDCl₃) δ : 9.32 (1H, s, H-1), 6.44 (1H, t, *J* = 7 Hz, H-3), 2.20-2.35 (4H, m, H-4, 3'), δ 5.33 (1H, m, H-2'), 1.60 (3H, s, CH₃-1'), 1.50 (4H, m, H-5, 4'), 1.27 (36H) and 0.88 (6H, t, *J* = 6.6 Hz H-17 and H-13').

¹³C NMR (150 MHz, CDCl₃): 195.4 (C-1), 143.7 (C-2), 155.5 (C-3), 129.6 (C-1'), 130.1 (C-2'), 31 (C-4 and C-3'), 23-29 (19CH₂), 22.0 (CH₃-1') and 14.2 (C-17 and C-12').

GCMS *m/z*: 432 [M, C₃₀H₅₆O]⁺, 431 [M- H]⁺, 305 [M-C₉H₁₉]⁺, 249 [M-C₁₃H₂₇]⁺, 183, 127, 43 (100).

GCMS *m/z*: 432 [M, C₃₀H₅₆O]⁺, 431 [M- H]⁺, 305 [M-C₉H₁₉]⁺, 249 [M-C₁₃H₂₇]⁺, 183, 127, 43 (100).

HR-EIMS: 432.4319 requires C₃₀H₅₆O, calcd. 432.4331;

Microtiter broth dilution method for yeast susceptibility

testing: Sterile microtiter assay trays containing 96 round-bottom wells (Dynatech Laboratories, Inc., Alexandria, Va.) were employed. The stock solutions of antifungal agents were appropriately diluted in the various assay media to give the following working solution of ketoconazole (Janssen Pharmaceuticals, Piscataway, N.J.), starting from 20 to 1000 μ g/mL. Two fold serial dilutions were performed with respective assay broths and dispensed into appropriate wells. Each vertical column of wells contained a single antifungal agent in progressive dilutions and was inoculated with a single clinical isolate. The first

horizontal row of wells (A1-A12) contained no antifungal agent and served both as a growth and sterility control. The inoculation of 0.1 mL of yeast suspension created a final range of antifungal concentrations identical to that of the agar dilution plates. The final volume in each well was 0.2 mL.

Inoculum: *Aspergillus fumigatus* WT (Af293), *A. flavus* WT (NRRL3357), and *Candida albicans* were grown in YPD (1% yeast, 2% peptone and 2% dextrose) medium. After 24 h of incubation at 37°C in a 5% CO₂ atmosphere, the isolates were diluted 1:100 in normal saline, followed by a 1:100 dilution in yeast peptone dextrose medium. Prepared in this manner, a volume of 0.1 mL of the final broth dilution contained approximately 1000 CFU of the previously mentioned fungi. Microtiter broth dilution plates were incubated at 37°C in 5% CO₂ for 48 h. MIC was defined as that concentration of the antifungal agent contained in the microtiter well or in the agar dilution plate in which the absence of visual turbidity (colonies) was first observed.

Acknowledgment - This work was supported by grants from the Deanship of Scientific Research (DSR) No. 429/003-8, King Abdul-Aziz University (KAU). Dr Nessim, I. Rady, Assistant Prof. of algal flora, Faculty of Science, Umm Al-Qura University, is acknowledged for algal identification, and Dr Shaymaa Hassan, Faculty of Pharmacy, Mansoura University, for her help in measuring the antifungal activity.

References

- [1] (a) Kladi M, Xenaki H, Vagias C, Papazafiri P, Roussis V. (2006) New cytotoxic sesquiterpenes from the red algae *Laurencia obtusa* and *Laurencia microcladia*. *Tetrahedron*, **62**, 182-189; (b) Kaur K, Jain M, Kaur T, Jain R. (2009) Antimalarials from nature. *Bioorganic and Medicinal Chemistry*, **17**, 3229-3256; (c) Li H-Y, Matsunaga S, Fusetani N. (1998) Antifungal metabolites from marine sponges. *Current Organic Chemistry*, **2**, 2649-682; (d) Vairappan CS. (2003) Potent antibacterial activity of halogenated metabolites from Malaysian red alga, *Laurencia maiuscula* (Rhodomelaceae, Ceramiales). *Biomolecular Engineer*, **20**, 255-259.
- [2] (a) Bruno LA, Beatriz GF, Mutue TF, Valéria LT. (2008) Sesquiterpenes of the Brazilian marine red alga *Laurencia filiformis* (Rhodophyta, Ceramiales). *Natural Product Communications*, **3**, 1653-1654; (b) Dias DA, White JM, Urban S. (2009) *Laurencia filiformis*: Phytochemical profiling by conventional and HPLC-NMR approaches. *Natural Product Communications*, **4**, 157-172; (c) Kladi M, Vagias C, Papazafiri P, Furnari G, Serio D, Roussis V. (2007). New sesquiterpenes from the red alga *Laurencia microcladia*. *Tetrahedron*, **63**, 7606-7611.
- [3] (a) Al-lihaibi SS, Ayyad SN, Al-wessabi EO, Alarif WM. (2010) 3 β , 7 β -Dihydroxy-5 α -cholestan-11-one: A new oxidation pattern of cholestane skeleton from *Laurencia papillosa*. *Biochemical Systematics and Ecology*, **38**, 861-863; (b) Lisboa BP, Ganshow LI, Halket JM, Pinheiro-Joventio F. (1982) The trace sterols of the red alga *Laurencia papillosa* investigated by open tubular capillary column gas chromatography. *Compound Biochemical Physiology*, **73B**, 257-264; (c) Zornista K, Kamen S, Rosalina S, Stefka D-K, Simon P. (2006) Comparative investigation on sterols from some Black Sea red algae. *Natural Products Research*, **20**, 113-118.
- [4] (a) Suzuki M, Kurosawa E, Kurata K. (1987) (E)-2-Tridecyl-2-heptadecenal, an unusual metabolite from the red alga *Laurencia* species. *Bulletin of the Chemical Society of Japan*. **60**, 3793-3794; (b) Gallo ML, Seldes AM, Cabrera GM. (2004) Antibiotic long-chain and α,β -unsaturated aldehydes from the culture of the marine fungus *Cladosporium* sp., *Biochemical Systematics and Ecology*, **32**, 545-551; (c) Anderson IG, Haslewood AD. (1967) New evidence for the structure of myxinol. *Biochemistry Journal*, **104**, 1061-1063; (d) Goad JL, Akihisa T. (1977) Analysis of sterols 1st. ed. Blackie Academic & Professional, 197-217.
- [5] (a) Zhao K, Wang Y, Han L. (2007) 4,5-Epoxy cholestane-3,6- diols: templates for generating the full set of eight cholestane -3, 5, 6- triol stereoisomerisms in multigram scales, but not for cholestane- 3, 4, 6- triol. *Steroids*, **72**, 95-104; (b) Jiang B, Shi H-P, Xu M, Wang W-J, Zhao W-S. (2008) Stereoselective synthesis of certonardolsterol D3. *Tetrahedron*, **64**, 9738-9744.
- [6] (a) Bultel-Ponce V, Etahriri S, Guyot M. (2002) New ketosteroid from the red alga *Hypnea musciformis*. *Bioorganic Medicinal Chemistry Letter*, **12**, 1715-1718; (b) Su B-N, Takaishi Y, Morinins H-K. (1999) Four novel phenylpropanol ester lipid metabolites from *Morina chinensis*, *Journal of Natural Products*, **62**, 1325-1327.

Steroidal Saponins from the Fruits of *Cestrum ruizteranianum*

Elier Galarraga M.^{a,b}, Anne-Claire Mitaine-Offer^a, Juan Manuel Amaro-Luis^b,
Tomofumi Miyamoto^c, Chiaki Tanaka^c, Laurent Pouységu^d, Stéphane Quideau^d,
Luis B. Rojas^b and Marie-Aleth Lacaille-Dubois^{a,*}

^aLaboratoire de Pharmacognosie, UMIB, UPRES-EA 3660, Faculté de Pharmacie,
Université de Bourgogne, 7, Bd Jeanne d'Arc, BP 87900, 21079 Dijon Cedex, France

^bLaboratorio de Productos Naturales. Departamento de Química, Facultad de Ciencias.
Universidad de Los Andes. Mérida, Venezuela-5101

^cGraduate School of Pharmaceutical Sciences, Kyushu University, Fukuoka, 812-8582, Japan

^dInstitut des Sciences Moléculaires, CNRS-UMR 5255 & Institut Européen de Chimie et Biologie,
Université de Bordeaux, 2 rue Robert Escarpit, 33607 Pessac Cedex, France

m-a.lacaille-dubois@u-bourgogne.fr

Received: September 23rd, 2011; Accepted: September 29th, 2011

Seven spirostane and furostane-type glycosides were isolated from the aqueous methanolic extract of the fruits of *Cestrum ruizteranianum* and characterized mainly by 2D NMR spectroscopy and mass spectrometry. These known saponins belong to the Δ^5 -spirostene and Δ^5 -furostene series and are reported in this species for the first time.

Keywords: Solanaceae, *Cestrum ruizteranianum*, steroidal saponins.

Cestrum ruizteranianum Benítez & D'Arcy is a medium height shrub endemic of the Venezuelan Andes [1]. The genus *Cestrum* (Solanaceae) comprises more than 300 species, which are native to warm temperate to tropical regions, mainly distributed in South America, but known as ornamental plants in the whole world [2]. The species of this genus are rich in steroidal saponins and exhibit many potential medicinal uses as insecticide, herbicide, molluscicide, antimicrobial agent, and anticancer agents for instance [2]. We chose to study *C. ruizteranianum* for chemotaxonomic reasons, and because no work on its saponin composition had been reported.

Seven saponins were obtained by several chromatographic methods from the dried fruits of *C. ruizteranianum*. The isolated compounds (**1-7**) were unambiguously identified by 2D NMR spectroscopy and mass spectrometry (FAB) as: pennogenin 3-*O*- β -chacotrioside (**1**) [3], (25*R*,26*R*)-spirost-5-ene-3 β ,17 α ,26-triol 3-*O*- β -chacotrioside (**2**) [4], methyl protodioscin (**3**) [5], protodioscin (**4**) [5], 26-*O*- β -D-glucopyranosyl-(25*R*)-furost-5-ene-3 β ,17 α ,22,26-tetraol 3-*O*- β -chacotrioside (**5**) [4], 26-*O*- β -D-glucopyranosyl-22-methoxy-(25*R*)-furost-5-ene-3 β ,26-diol-3-*O*- α -L-rhamnopyranosyl-(1 \rightarrow 4)- α -L-rhamnopyranosyl-(1 \rightarrow 4)-[α -L-rhamnopyranosyl-(1 \rightarrow 2)]- β -D-glucopyranoside (Methyl proto Pb) (**6**) [6], and 26-*O*- β -D-glucopyranosyl-(25*R*)-furost-5-ene-3 β ,22,26-triol-3-*O*- α -L-rhamnopyranosyl-(1 \rightarrow 4)- α -L-rhamnopyranosyl-(1 \rightarrow 4)-[α -L-rhamnopyranosyl-(1 \rightarrow 2)]-

β -D-glucopyranoside (**7**) [7]. All spectroscopic data were in agreement with those previously reported.

Chemotaxonomic significance: Steroidal saponins with spirostane and furostane-type skeletons are very common in *Cestrum* species [2], like in *C. parqui* [8], *C. diurnum* [9], *C. nocturnum* [10], *C. sendtnerianum* [11], *C. axillare* [12], and *C. laevigatum* [13]. From a chemotaxonomic point of view, it is not surprising to find steroidal saponins in the fruits of *C. ruizteranianum*, but the isolated compounds (**1-7**) are found for the first time in this species.

Experimental

Plant material: Plant material was collected in Mucuchies-Gavidia (2950 m), Municipio Rangel, Mérida-Venezuela, in April 2010 and identified by Ing. Juan Carmona Arzola, Universidad de Los Andes (Mérida-Venezuela). A voucher specimen (J. M. Amaro *et al.*, No. 1862) was deposited in the MERF Herbarium, Faculty of Pharmacy, ULA.

Extraction and isolation: Eight g of the MeOH/H₂O (7:3) extract of the dried fruits was fractionated by isocratic vacuum liquid chromatography (VLC) on silica gel (40-63 μ m) (CHCl₃/MeOH/H₂O, 13:7:2, lower phase, 200 mL, each) yielding 3 fractions (A-C). Fraction A (360 mg) was then separated by medium pressure liquid chromatography (MPLC) on silica gel (15-40 μ m) (CHCl₃/MeOH/H₂O,

13:7:2, lower phase) to obtain compounds **1** (11 mg) and **2** (15 mg). Fraction B (4.4 g) was chromatographed on a Sephadex LH-20 column (MeOH/H₂O, 1:1) yielding 5 subfractions (I-V). Subfraction I (1.1 g) was then

submitted to successive MPLC on normal and RP-18 silica gel to afford compounds **3** (9.8 mg), **4** (10.0 mg), **5** (11.1 mg), **6** (11.6 mg) and **7** (10.8 mg). All saponins were isolated as white amorphous powders.

References

- [1] Jauregui D, de Rios NR, Benitez de Rojas CE. (2001) Leaf anatomic studies in Solanaceae of Venezuela. VI. Leaf anatomy of 10 species of *Cestrum* L. *Acta Cientifica Venezolana*, **52**, 248-260.
- [2] Begum AS, Goyal M. (2007) Research and medicinal potential of genus *Cestrum* (Solanaceae)-a review. *Pharmacognosy Reviews*, **1**, 320-332.
- [3] Nohara T, Miyahara K, Kawasaki T. (1975) Steroid saponins and sapogenins of underground parts of *Trillium kamschaticum* Pall. II. Pennogenin- and kryptogenin 3-*O*-glycosides and related compounds. *Chemical & Pharmaceutical Bulletin*, **23**, 872-885.
- [4] Ando J, Miyazono A, Zhu XH, Ikeda T, Nohara T. (1999) Studies on the constituents of Solanaceous plants, steroidal glycosides from *Solanum nodiflorum*. *Chemical & Pharmaceutical Bulletin*, **47**, 1794-1796.
- [5] Kawasaki T, Komori T, Miyahara K, Nohara T. (1974) Furostanol bisglycosides corresponding to dioscin and gracillin. *Chemical & Pharmaceutical Bulletin*, **9**, 2164-2175.
- [6] Miyamura M, Nakano K, Nohara T, Tomimatsu T, Kawasaki T. (1982) Steroid saponins from *Paris polyphylla* Sm.-Supplement. *Chemical & Pharmaceutical Bulletin*, **30**, 712-718.
- [7] Kawabe N, Togami Y, Uenishi N, Rii H, Chen S. (1996) Extraction of anticancer saponins from *Ypsilandra thibetica*. Jpn. Kokai Tokkyo Koho, JP 08040914 A.
- [8] Chaieb I, Tayeb W. (2009) Comparison of the molluscicidal activity of *Cestrum parqui* (Solanaceae) and *Quillaja saponaria* (Quillajaceae) saponins. *Tunisian Journal of Medicinal Plants and Natural Products*, **2**, 31-35.
- [9] Fouad MA, Mohamed KM, Kamel MS, Matsunami K, Otsuka H. (2008) Cesdiurins I-III, steroidal saponins from *Cestrum diurnum*. *Journal of Natural Medicines*, **62**, 168-173.
- [10] Mimaki Y, Watanabe K, Sakagami H, Sashida Y. (2002) Steroidal glycosides from the leaves of *Cestrum nocturnum*. *Journal of Natural Products*, **65**, 1863-1868.
- [11] Haraguchi M, Mimaki Y, Motidome M, Morita H, Takeya K, Itokawa H, Yokosuka A, Sashida Y. (2000) Steroidal saponins from the leaves of *Cestrum sendtnerianum*. *Phytochemistry*, **55**, 715-720.
- [12] De Amorim A, Borba HR, Pereira NA, Matsunami K, Otsuka H. (1992) Effect of the toxic fraction of *Cestrum axillare* on mitochondria. *Revista Brasileira de Farmacia*, **73**, 2-4.
- [13] Canham PAS, Warren FL (1950) Saponins. I. Isolation of gitogenin and digitogenin from *Cestrum laevigatum*. *Journal of the South African Chemical Institute*, **3**, 9-12.

Isolation and Cholinesterase Activity of Amaryllidaceae Alkaloids from *Nerine bowdenii*

Lucie Cahliková^a, Stanislav Zavadil^a, Kateřina Macáková^a, Irena Valterová^b,
Andrea Kulhánková^a, Anna Hošťálková^a, Jiří Kuneš^b and Lubomír Opletal^a

^aDepartment of Pharmaceutical Botany and Ecology, Faculty of Pharmacy, Charles University, Heyrovského 1203, 500 05 Hradec Králové, Czech Republic

^bInstitute of Organic Chemistry and Biochemistry of the AS CR, Flemingovo nám. 2, 166 10 Praha 6, Czech Republic

^cDepartment of Inorganic and Organic Chemistry, Faculty of Pharmacy, Charles University, Heyrovského 1203, 500 05 Hradec Králové, Czech Republic

cahlikova@faf.cuni.cz

Received: September 26th, 2011; Accepted: October 20th, 2011

Amaryllidaceae species are known as ornamental plants. Some contain galanthamine, an acetylcholinesterase inhibitor. The chemical composition of the alkaloid extract of bulbs of *Nerine bowdenii* Watson has been analyzed by means of GC/MS. Twenty-two compounds were detected and nineteen of them identified, one of which was belladine. The alkaloid extract showed promising cholinesterase inhibitory activities against human blood acetylcholinesterase (HuAChE; $IC_{50} = 87.9 \pm 3.5 \mu\text{g/mL}$) and human plasma butyrylcholinesterase (HuBuChE; $IC_{50} = 14.8 \pm 1.1 \mu\text{g/mL}$). Belladine inhibited HuAChE and HuBuChE in a dose-dependent manner with IC_{50} values of $781 \pm 12.5 \mu\text{M}$ and $284.8 \pm 4.2 \mu\text{M}$, respectively.

Keywords: *Nerine bowdenii*, GC/MS, belladine, acetylcholinesterase, butyrylcholinesterase.

Alzheimer's disease (AD) is the most common form of dementia. Epidemiological data indicate a considerable potential increase in the prevalence of the disease over the next two decades [1]. Although the pathogenesis of AD is complicated and not fully established, two major hypotheses (amyloid cascade and cholinergic) are currently under consideration regarding the molecular mechanism.

In AD patients, deficit of cholinergic functions, which results in decreased levels of the neurotransmitter acetylcholine (ACh) in the cortex, is responsible for the memory impairments [2a]. The principal role of acetylcholinesterase (AChE) is the termination of nerve impulse transmission at the cholinergic synapses by rapid hydrolysis of ACh. Inhibition of AChE serves as a strategy for the treatment of AD, senile dementia, ataxia, myasthenia gravis, and Parkinson's disease [2b]. In the healthy brain, AChE is the most important enzyme regulating the level of ACh, while another enzyme, butyrylcholinesterase (BuChE) plays only a minor role. Moreover, in late AD stages, levels of AChE have declined by up to 85% and BuChE represents the predominant cholinesterase in the brain. BuChE, primarily associated with glial cells, but also with specific neuronal pathways, cleaves ACh in a manner similar to that of AChE to terminate its physiological action [2c]. Galanthamine, which is used for the treatment of AD, has been shown to

be much less potent against BuChE than AChE [2d]. This fact has targeted BuChE as a new approach to intercede in the progression of AD and requires research into new inhibitors with dual enzymatic activity. Currently, cholinesterase inhibition is the most used therapeutic treatment for the symptoms of AD [2e, f].

Extracts from plants belonging to the Amaryllidaceae family have long been used as herbal remedies. These plants are known to synthesize a particular type of bioactive compound known as Amaryllidaceae alkaloids [3]. Some species of this family contain galanthamine, a long-acting, selective, reversible and competitive acetylcholinesterase inhibitor, as well as other alkaloids with interesting pharmacological activities, such as anticancer, antiviral, antimalarial, anti-inflammatory and antiparasitic [4a-e]. Among the bulbs of Amaryllidaceae plants screened previously for AChE inhibitory activity, *Nerine bowdenii* showed strong inhibition [5a,b].

The genus *Nerine*, the second largest within Amaryllidaceae with ca 30 species, is an autumn-flowering perennial bulbous plant group, whose species inhabit areas with summer rainfall and cool, dry winters [6]. Previous investigations led to the isolation of more than 30 Amaryllidaceae alkaloids. Some of these have been tested for their acetylcholinesterase inhibitory activity, but only ungeremine and undulatine showed interesting inhibition

activity with IC_{50} values of 0.35 μ M [7a] and 37 μ M [7b], respectively. In both studies acetylcholinesterase from electric eel was used.

The ethanolic extract of *N. bowdenii* bulbs exhibited significant HuAChE and HuBuChE inhibition activity ($61.5\% \pm 4.0\%$ for HuAChE and $89.8\% \pm 5.2\%$ for HuBuChE, respectively) at a concentration of 500 μ g/mL, thus indicating the presence of cholinesterase inhibitors in this plant taxon. The crude extract from the bulbs was concentrated into an alkaloid fraction and, as expected, this possessed a promising activity with an IC_{50} value of 87.9 ± 3.5 μ g/mL for HuAChE and 14.8 ± 1.1 μ g/mL for HuBuChE (values for reference compounds: belladine 781 ± 12 , galanthamine 6.9 ± 0.3 , huperzine A 0.25 ± 0.01 μ g/mL for HuAChE, and 285 ± 5 , 156 ± 2.9 , > 500 μ g/mL, respectively, for HuBuChE).

In order to identify the compounds in the complex alkaloid fraction of *N. bowdenii* bulbs, capillary GC/MS was employed. In the bulb extract, twenty-two alkaloids of the crinine, lycorine, belladine and haemanthamine types were detected by GC (Fig. 1); nineteen of them were identified based on their retention times (RT) and mass spectra as buphanisine (**1,2**), crinine (**3**), belladine (**4**), caranine (**5**), *N*-demethylbelladine (**7**), acetylscatellatine (**8**), buphanidine (**9**), buphanamine (**10**), crinamine (**11**), powelline (**12**), undulatine (**15**), 11-*O*-acetylbelladine (**16**), ambelline (**17**), 3-acetylnerbowdine (**18**), crinamidine (**19**), bowdensine (**20**), and 11-*O*-acetyl-1,2-epoxyambelline (**21**) (Table 1). Compound **2** showed a longer retention time than buphanisine (**1**) by GC-MS, but its MS pattern was the same, indicating that it is an isomer of **1**. Compounds **13**, **14** and **22** remained unidentified. Considering the low concentration of **13** and **14** ($< 1\%$ of TIC) and the complexity of the alkaloid mixture, their isolation and structural elucidation could be problematic. The mass spectrum of compound **22** was not found either in the databases or in the literature. For the identification of this compound, one of the major components of the alkaloid extract (6.8% of TIC), isolation in larger amount and structure elucidation is needed. The haemanthamine and crinine series of Amaryllidaceae alkaloids only differ in the position of the 5,10*b*-ethano bridge [8a]. The alkaloids of belladine type, which are biosynthetic precursors of Amaryllidaceae alkaloids [8b], have been previously reported only in the genera *Nerine* (*N. filifolia*) and *Crinum* [8c,d]. For *N. bowdenii*, belladine (**4**) and *N*-demethylbelladine (**7**) are reported here for the first time, which is quite surprising when considering their domination of the alkaloid mixture and the number of previous phytochemical investigations of the species. Alkaloids of the lycorine type (**5**, **6** and **8**) were identified only as minor components in the mixture. The relative proportion of each alkaloid was determined as a percentage of the total ion current. The alkaloid pattern of bulbs of *N. bowdenii* was dominated by belladine (**4**; 49.9 % of TIC), ambelline (**17**; 9.1 % of TIC) and *N*-demethylbelladine (**7**; 8.5 % of TIC).

A wide range of Amaryllidaceae alkaloids belonging to various ring types has been evaluated in recent studies for their *in vitro* AChE inhibitory activity with either galanthamine or physostigmine as a positive control [7b, 9a-c]. Results of these studies showed that the AChE inhibitory activity is associated mainly with galanthamine- and lycorine-type alkaloids. The lycorine-type compounds are less active inhibitors than the galanthamine type and their activity is associated with a substitution at positions C-1 and C-2 [9d]. Crinine-type alkaloids have only a weak activity against AChE and it appears that the stereochemistry of the 5,10*b*-ethanobridge has no effect on the inhibition of AChE activity [9c]. In comparison with our previous studies on other Amaryllidaceae plants, such as *Zephyranthes robusta* [9e], *Z. grandiflora* [9f] and *Chlidanthus fragrans* [9g], the alkaloid extract from *N. bowdenii* showed the weakest AChE inhibition activity, which is in agreement with the above mentioned conclusions, because galanthamine-type alkaloids were not detected and the lycorine-type alkaloids represented only a minor fraction (2.9 % of TIC). The alkaloid extracts of *Z. robusta*, *Z. grandiflora* and *Ch. fragrans* were dominated by galanthamine- and lycorine-type alkaloids, and consequently the HuAChE inhibition activity of these extracts was markedly stronger. On the other hand, the extract of *N. bowdenii* showed promising inhibition activity against HuBuChE with an IC_{50} value of 14.8 ± 1.1 μ g/mL. As mentioned, BuChE plays an important role in the late AD stages, but only a limited number of alkaloids have been tested for their BuChE inhibitory activity so far. There are no data on BuChE inhibition activity of pure Amaryllidaceae alkaloids.

Relatively surprising was the weak cholinesterase activity of the isolated alkaloid belladine (**4**), previously isolated only from *N. filifolia* [8c], in comparison with the alkaloid extract due to its dominant amount in the mixture (49.9% of TIC) of *N. bowdenii*, and the very good HuBuChE inhibition activity of this extract. It is clear that the BuChE inhibition activity of *N. bowdenii* must be connected to some compound/compounds present in a smaller amount. The isolation and identification of this/these alkaloids will be the next step in searching for new bioactive natural compounds with BuChE inhibition activity.

Experimental

Plant materials: The fresh bulbs of *Nerine bowdenii* were obtained from Lukon Glads (Sadská, Czech Republic). The botanical identification was performed by Prof. Lubomír Opletal, CSc. A voucher specimen is deposited in the Herbarium of the Faculty of Pharmacy at Hradec Králové.

Extraction of alkaloids: Fresh bulbs (150 g) were extracted 3 times with EtOH (150 mL) at room temperature for 24 h. The solvent was evaporated under reduced pressure and the residue dissolved in 50 mL 2% HCl. After removing the neutral compounds with diethyl ether (3 x 50 mL), the extract was basified with 25% ammonia solution and the alkaloids extracted with EtOAc

Table 1: Alkaloids identified by GC/MS in bulbs of *Nerine bowdenii*.

Compound	Retention time, ^a (min)	[M ⁺] and characteristic ions <i>m/z</i> (% relative intensity)	% ^b	Reference for MS data
1 Buphanisine, isomer I	25.34	285 (100), 270 (23), 253 (25), 230 (28), 215 (80), 201 (22), 187 (20), 157 (15)	1.4	NIST 08, [10c]
2 Buphanisine, isomer II	25.47	285 (100), 270 (23), 253 (25), 230 (28), 215 (80), 201 (22), 187 (20), 157 (15)	0.7	NIST 08, [10c]
3 Crinine	26.45	271 (100), 254 (8), 242 (8), 228 (25), 216 (14), 199 (48), 187 (43)	1.3	NIST 08, [10c]
4 Belladine	26.65	315 (2), 194 (70), 152 (90), 151 (100), 135 (10), 121 (18), 107 (18)	49.9	NIST 08, [10d], ^d
5 Caranine	27.10	271 (86), 270 (46), 252 (52), 250 (10), 240 (12), 227 (48), 226 (100), 212 (5)	0.2	NIST 08
6 Acetylcaranine	27.24	313 (80), 270 (5), 253 (20), 252 (95), 250 (15), 240 (10), 225 (40), 226 (100)	2.6	NIST 08
7 <i>N</i> -Demethylbelladine	27.64	301 (2), 299 (3), 273 (1), 195 (6), 180 (20), 151 (100), 138 (4), 128 (2), 121 (9), 107 (6)	8.5	[8c]
8 Acetylscatelline	28.49	343 (75), 300 (5), 283 (25), 282 (80), 266 (30), 257 (77), 256 (100), 241 (12)	0.1	NIST 08
9 Buphanidine	28.82	315 (100), 300 (26), 287 (30), 272 (10), 260 (45), 245 (60), 231 (30), 228 (22)	2.9	NIST 08, [10c]
10 Buphanamine	28.94	301 (100), 286 (12), 284 (14), 272 (14), 256 (18), 231 (25), 218 (18), 204 (18)	1.1	NIST 08
11 Crinamine	29.34	301 (1), 269 (100), 240 (35), 225 (20), 224 (25), 211 (17), 181 (58)	1.9	NIST 08
12 Powelline	29.67	301 (100), 284 (8), 272 (7), 258 (20), 246 (12), 244 (12), 229 (80), 217 (40)	0.3	NIST 08
13 A13 ^c	30.07	331 (22), 313 (10), 299 (28), 287 (100), 284 (14), 282 (10), 270 (20), 255 (45)	0.4	-
14 A14 ^c	30.19	373 (90), 314 (72), 299 (38), 284 (20), 270 (20), 254 (54), 242 (50), 226 (38), 216 (20), 115 (25)	0.3	-
15 Undulatine	31.14	331 (100), 316 (6), 302 (9), 300 (6), 286 (20), 272 (5), 260 (18), 258 (40)	3.6	[10c]
16 11- <i>O</i> -Acetylbelleline	31.24	373 (100), 358 (15), 342 (15), 330 (10), 314 (70), 313 (69), 298 (30), 282 (55), 270 (25)	4.5	[10c]
17 Ambelline	31.49	331 (100), 316 (5), 299 (30), 287 (55), 270 (25), 260 (50), 257 (35), 255 (23), 241 (27)	9.1	NIST 08
18 Acetylnerbowdine	32.10	361 (75), 318 (30), 302 (60), 273 (20), 254 (50), 244 (25), 43 (100)	0.7	NIST 08
19 Crinamidine	32.62	317 (75), 288 (100), 274 (5), 259 (22), 258 (25), 243 (28), 244 (30), 230 (25), 217 (40), 205 (40)	0.4	NIST 08
20 Bowdensine	32.75	403 (82), 344 (80), 288 (75), 284 (50), 272 (40), 230 (40), 217 (42), 189 (28), 173 (50), 43 (100)	2.9	NIST 08
21 11- <i>O</i> -Acetyl-1,2-epoxybelleline	33.52	389 (60), 330 (75), 316 (100), 274 (30), 256 (55), 231 (40), 228 (12), 205 (25), 203 (27), 190 (18)	0.4	[10c]
22 A22 ^c	33.87	329 (100), 314 (18), 300 (8), 286 (12), 270 (6), 231 (28), 218 (15), 204 (20), 190 (10)	6.8	-

^a For GC conditions see the Experimental section, ^b Values are expressed as a percentage of the total ion current (TIC), ^c not identified, ^d standard.

(3 x 50 mL). The organic solvent was evaporated and 10 mg of alkaloid extract was removed for HuAChE and HuBuChE assay. 10 mg of the dry alkaloid fraction was dissolved in MeOH to a final concentration of 10 mg/mL for further analysis. The rest of the fraction (180 mg) was used for the isolation of pure alkaloids.

GC/MS analysis and identification of alkaloids: The GC/MS analysis of alkaloids from *N. bowdenii* was carried out on a gas chromatograph (Focus Thermo Scientific, USA) with a splitless injector (280°C) and a mass detector (200°C, GC-MS MD 800 Fisons, Manchester, UK). A DB-5MS column (30 m x 0.25 mm x 0.25 µm, Agilent Technologies Santa Clara, CA, USA) and helium gas (constant flow 1 mL/min) were used for separation. The temperature program was: 100 - 180°C at 15°C/min, 1 min held at 180°C, and 180 - 300°C at 5°C/min and 5 min held at 300°C, detection range *m/z* 40-600. The injector temperature was 280°C. The alkaloids were identified by comparison of their MS with those in the NIST library and with those reported in the literature [8c, 9a, 10a-e].

Isolation and identification of belladine: The alkaloid extract of *N. bowdenii* was subjected to preparative TLC (silica gel 60, F₂₅₄, Merck) and eluted with toluene:Et₂NH (95:5, 3 times) to give 4 subfractions A/1-A/4. Subfraction A/1 (35 mg) was further subjected to preparative TLC (cHx: Et₂NH, 95:5, 3 times), which led to the isolation of compound **4** (20 mg). Because of the small amount of other subfractions, they were not used for the isolation of other alkaloids. The structure of belladine (**4**) was determined by comparison of its spectral data with those

reported in the literature [8c]. NMR spectra were recorded on a Varian VNMR500 spectrometer, operating at 500 MHz (¹H) and at 125 MHz (¹³C). ESI-MS were measured on a Thermo Finnigan LCQDuo spectrometer.

Preparation of red blood cell ghosts: Ghosts were prepared from freshly drawn blood (taken from healthy volunteers), to which 1 mL of sodium citrate per 10 mL of blood was added, according to the method of Steck and Kant [11a], with slight modification. Briefly, plasma (HuBuChE) was removed from the whole blood by centrifugation at 4000 rpm in a Boeco U-32R centrifuge with a Hettich 1611 rotor. Red blood cells were transferred to 50 mL tubes and washed 3 times with 5 mM phosphate buffer (pH 7.4) containing 150 mM sodium chloride (12000 rpm, Avanti J-30I, rotor JA-30.50). The washed erythrocytes were stirred with 5 mM phosphate buffer (pH 7.4) for 10 mins to ensure lysis. The lysed cells were centrifuged at 20,000 rpm for 10 mins and then the ghosts (HuAChE) were washed 3 times with phosphate buffer.

Acetylcholinesterase and butyrylcholinesterase assay: HuAChE and HuBuChE activities were determined using a modified method of Ellman *et al.* [11b] at concentrations of 0.5, 2.5, 5, 12.5, 25, 50, 125, 250 and 500 µg/mL using acetylthiocholine iodide and butyrylthiocholine iodide as substrates, respectively. Galanthamine and huperzine A were used as positive standards. The % inhibition was calculated according to the formula: %I = 100 - (ΔA_{BL}/ΔA_{SA})*100, where ΔA_{BL} is increase of absorbance of blank sample and ΔA_{SA} the increase of absorbance of the measured sample.

Statistical analysis: The IC₅₀ values were calculated with the use of GraphPad Prism 5.02 software.

Acknowledgments – This project was supported by grants SVV-2011-263002 and MSM0021620822.

References

- [1] Johnson N, Davis T, Bosanquet N. (2000) The epidemic of Alzheimer's disease; how can we manage the costs? *Pharmacoeconomics*, **18**, 215-223.
- [2] (a) Lahiri DK, Farlow MR, Greig NH, Sambamurti K. (2002) Current drug targets for Alzheimer's disease treatment. *Drug Development Research*, **56**, 267-281; (b) Rahman AU, Choudhary MI. (2001) Bioactive natural products as a potential source of new pharmacophores a theory of memory. *Pure and Applied Chemistry*, **73**, 555-560; (c) Greig NH, Lahiri DK, Sambamurti K. (2002) Butyrylcholinesterase: an important new target in Alzheimer's disease therapy. *International Psychogeriatrics*, **14**, 77-91; (d) Fulton B, Benfield P. (1996) Galanthamine. *Drugs & Aging*, **9**, 60-65; (e) Lleó A. (2007) Current therapeutic options for Alzheimer's disease. *Current Genomics*, **8**, 550-558; (f) Recanatini M, Valenti P. (2004) Acetylcholinesterase inhibitors as a starting point towards improved Alzheimer's disease therapeutics. *Current Pharmaceutical Design*, **10**, 3157-3166.
- [3] Bastida J, Lavilla R, Viladomat F. (2006) Chemical and biological aspects of *Narcissus* alkaloids. In *The Alkaloids*, Vol. **63**, Cordell GA. (Ed.). Elsevier Scientific: Amsterdam, NL, pp. 87-179.
- [4] (a) Evidente A, Kornienko A. (2009) Anticancer evaluation of structurally diverse Amaryllidaceae alkaloids and their synthetic derivatives. *Phytochemistry Reviews*, **8**, 449-459; (b) Toriizuka Y, Kinoshita E, Kogure N, Kitajima M, Ishiyama A, Otaguro K, Yamada H, Omura S, Takayama H. (2008) New lycorine-type alkaloid from *Lycoris traubii* and evaluation of antitrypanosomal and antimalarial activities of lycorine derivatives. *Bioorganic & Medicinal Chemistry*, **16**, 10182-10189; (c) Weniger B, Italiano L, Beck JP, Bastida J, Bergeño S, Codina C, Lobstein A, Anton R. (1995) Cytotoxic activity of Amaryllidaceae alkaloids. *Planta Medica*, **61**, 77-79; (d) Zupkó I, Réthy B, Hohmann J, Molnár J, Ocsóvzki I, Falaky G. (2009) Antitumor activity of alkaloids derived from Amaryllidaceae species. *In Vivo*, **23**, 41-48; (e) Lamoral-Theys D, Andolfi A, van Goietsenoven G, Cimmino A, Le Calvé B, Wauthoz N, Mégalizzi V, Gras T, Bruyère C, Dubois C, Mathieu V, Kornienko A, Kiss R, Evidente A. (2009) Lycorine, the main phenanthridine Amaryllidaceae alkaloid, exhibits significant antitumor activity in cancer cells that display resistance to proapoptotic stimuli: an investigation of structure-activity relationship and mechanistic insight. *Journal of Medicinal Chemistry*, **52**, 6244-6256.
- [5] (a) Rhee IK, van de Meent M, Ingkaninan K, Verpoorte R. (2001) Screening for acetylcholinesterase inhibitors from Amaryllidaceae using silica gel thin-layer chromatography in combination with bioactivity staining. *Journal of Chromatography A*, **915**, 217-223; (b) Marston A, Kissling J, Hostettmann K. (2002) A rapid TLC bioautographic method for the detection of acetylcholinesterase and butyrylcholinesterase inhibitors in plants. *Phytochemical Analysis*, **13**, 51-54.
- [6] Rix M, Phillips R. (1983) *The bulb book: a photographic guide to over 800 hardy bulbs*, Pan Books Ltd., London, GB, pp. 171.
- [7] (a) Rhee IK, Appels N, Hofte B, Karabatak B, Erkelens C, Stark LM, Flippin LA, Verpoorte R. (2004) Isolation of the acetylcholinesterase inhibitor ungeremine from *Nerine bowdenii* by preparative HPLC coupled on-line to a flow assay system. *Biological & Pharmaceutical Bulletin*, **27**, 1804-1809; (b) van Rijn RM, Rhee IK, Verpoorte R. (2010) Isolation of acetylcholinesterase inhibitory alkaloids from *Nerine bowdenii*. *Natural Product Research*, **24**, 222-225.
- [8] (a) De Angelis G, Wildman W. (1969) Identification of Amaryllidaceae alkaloids using ORD and CD spectroscopy. *Tetrahedron Letters*, **9**, 729-732; (b) Eichhorn J, Takada T, Kita Y, Zenk MH. (1998) Biosynthesis of the Amaryllidaceae alkaloid galanthamine. *Phytochemistry*, **49**, 1037-1047; (c) Nair JJ, Campbell WE, Brun R, Viladomat F, Codina C, Bastida J. (2005) Alkaloids from *Nerine filifolia*. *Phytochemistry*, **66**, 373-382; (d) Jin Z. (2009) Amaryllidaceae and *Sceletium* alkaloids. *Natural Product Reports*, **26**, 363-381.
- [9] (a) López S, Bastida J, Viladomat F, Codina C. (2002) Acetylcholinesterase inhibitory activity of some Amaryllidaceae alkaloids and *Narcissus* extracts. *Life Sciences*, **71**, 2521-2529; (b) Elgorashi EE, Stafford GI, van Staden J. (2004) Acetylcholinesterase enzyme inhibitory effects of Amaryllidaceae alkaloids. *Planta Medica*, **70**, 258-260; (c) Houghton PJ, Agbedahunsi JM, Adegbulugbe A. (2004) Choline esterase inhibitory properties of alkaloids from two Nigerian *Crinum* species. *Phytochemistry*, **65**, 2893-2896; (d) Houghton PJ, Ren Y, Howes MJ. (2006) Acetylcholinesterase inhibitors from plants and fungi. *Natural Product Reports*, **23**, 181-189; (e) Cahliková L, Kulhánková A, Urbanová K, Valterová I, Macáková K, Kuneš J. (2010) Analysis of Amaryllidaceae alkaloids from *Zephyranthes robusta* by GC-MS and their cholinesterase activity. *Natural Product Communications*, **5**, 1201-1204; (f) Cahliková L, Valterová I, Macáková K, Opletal L. (2011) Analysis of Amaryllidaceae alkaloids from *Zephyranthes grandiflora* by GC-MS and their cholinesterase activity. *Revista Brasileira de Farmacognosia/Brazilian Journal of Pharmacognosy*, **21**, 575-580; (g) Cahliková L, Macáková K, Zavadil S, Jiroš P, Opletal L, Urbanová K, Jahodář L. (2011) Analysis of Amaryllidaceae alkaloids from *Chlidanthus fragrans* by GC-MS and their cholinesterase activity. *Natural Product Communications*, **6**, 603-606.
- [10] (a) Kreh M, Matusch R, Witte L. (1995) Capillary gas chromatography-mass spectrometry of Amaryllidaceae alkaloids. *Phytochemistry*, **38**, 773-776; (b) Berkov S, Bastida J, Nikolova M, Viladomat F, Codina C. (2008) Rapid TLC/GC-MS identification of acetylcholinesterase inhibitors in alkaloid extract. *Phytochemical Analysis*, **19**, 411-419; (c) Viladomat F, Codina C, Bastida J, Mathee S, Campbell WE. (1995) Further alkaloids from *Brunsvigia josephinae*. *Phytochemistry*, **40**, 961-965; (d) Ghosal S, Shanthi A, Singh SK. (1988) Isocraugsodine, an *N*-arylidene-n-phenethylamine from *Crinum asiaticum* and its *E-Z* isomerism. *Phytochemistry*, **27**, 1849-1852; (e) Machocho A, Chhabra SC, Viladomat F, Codina C, Bastida J. (1999) Alkaloids from *Ammocharis tinneana*. *Phytochemistry*, **51**, 1185-1191.
- [11] (a) Steck TL, Kant JA. (1974) Preparation of impermeable ghosts and inside-out vesicles from human erythrocyte membranes. *Methods in Enzymology*, **31** (Pt A), 172-180; (b) Ellman GL, Courtney KD, Andres V, Featherstone RM. (1961) A new rapid colorimetric determination of acetylcholinesterase activity. *Biochemical Pharmacology*, **7**, 88-95.

HPLC Determination of Majdine in *Vinca herbacea*

Natia Gagua^a, Beatrice Baghdikian^{b,*}, Fathi Mabrouki^b, Riad Elias^b, Valentina Vachnadze^c, Aliosha Bakuridze^a and Evelyne Ollivier^b

^aTbilisi State Medical University, Faculty of Pharmacy, 33 Vaja-Pshavela avenue, 0162, Tbilisi, Georgia

^bLaboratory of Pharmacognosy and Ethnopharmacology, UMR-MD3, Faculty of Pharmacy, University of the Mediterranean Aix-Marseille II, 27 Bd Jean Moulin, F-13385 Marseille cedex 5, France

^cIovel Kutateladze Pharmacochimistry Institute, Laboratory of alkaloids, 36 Saradjishvili street, 0152, Tbilisi, Georgia

beatrice.baghdikian@univmed.fr

Received: May 19th, 2011; Accepted: October 26th, 2011

A reliable HPLC method coupled with DAD detection was developed and validated for determination of majdine in *Vinca herbacea*. The chromatographic separation was carried out on a Symmetry C₁₈ column (250 mm x 4.6 mm, 5 µm, Waters) with an isocratic solvent system of 25 mM potassium phosphate buffer (pH=3.0)-acetonitrile. UV detection was performed at 225 nm. Good linear behavior over the investigated concentration range was observed with the value of $r^2 > 0.9978$. The method was reproducible with intra- and inter-day variations of less than 4.38%. The proposed method was linear, accurate, precise and specific. The validated method was successfully applied to quantify majdine in various parts of *V. herbacea*, which was collected during the flowering months of April and May. The results indicated that the developed HPLC method could be used for the quality control of *V. herbacea* and for the standardization of its extracts in majdine.

Keywords: HPLC-DAD, validation, *Vinca herbacea*, alkaloids, majdine.

Vinca herbacea Waldst. and Kit. (Apocynaceae) is a flowering plant native to eastern and southeastern Europe, from south Austria to Greece and east to the Crimea, and also in southwestern Asia east to the Caucasus, especially in the eastern part of Georgia. *V. herbacea* is an herbaceous perennial plant growing as a trailing vine, growing up to 10-20 cm in height.

The anticancer compounds vinblastine and vincristine are present in *V. rosea* leaves, thus investigations of *Vinca* species became very interesting. Previous phytochemical studies have reported the presence of alkaloids in various parts of *V. herbacea*: akuamicine, akuamine, herbaine, hervine, herbaline, herbaceine, lochnerine, norfluorurarine, 11-methoxytabersonine, tabersonine, reserpine, reserpinine, isoreserpinine, vincamine, vincalherbine, and vincalherbinine. The alkaloids belonging to the oxindole group, such as majdine, isomajdine and carapanaubine, were detected in leaves and roots of *V. herbacea*. Majdine has been shown to be the major alkaloid in the roots and leaves during the period of flower formation. [1-7].

The pharmacological activity of the total alkaloids of *V. herbacea* has been investigated. They markedly reduced blood pressure for an extended period [8]. Also, the blocking action of *V. herbacea* on neuromuscular synapses has been studied. Intravenous injection of total alkaloids in

doses greater than 8 mg/kg was able to induce a curare-like action in rabbits and cats [9]. Similarly, a mixture of alkaloids named Vinherbine, has been shown to produce spasmolytic effects in Guinea pig ileum preparation.

Pharmacological studies have demonstrated that these alkaloids also possess cardio stimulating and antiarrhythmic activity [10-13]. The bacteriostatic effect of the alkaloids has been tested on *Staphylococcus aureus* and *Escherichia coli*. The indole alkaloids were shown to be the most active [14]. In addition, recent pharmacological experiments have revealed that majdine and isomajdine were responsible for apoptotic and antioxidant activity [15].

Majdine is one of the main alkaloids in the plant. Based on pharmacological studies, majdine has been selected as the chemical marker of *V. herbacea*. Therefore, the quantification of this alkaloid is necessary to evaluate the quality of this species. No analytical method has been reported in the literature for alkaloid quantification in *V. herbacea*. Thus, the development of an analytical method is necessary for the quality control of the plant.

HPLC methods have been reported for indole alkaloids such as vincamine. Mostly, mobile phases with buffer are used for determination of these alkaloids [16,17].




Figure 1: Chemical structure of majdine.

The aim of this study was the development and validation of a HPLC-DAD method for the quantification of majdine (Figure 1) in *V. herbacea*.

The separation conditions were optimized to achieve satisfactory resolution. Various reversed phase columns such as Symmetry C18, SunFire C18 and Atlantis C18 were tested. The optimal separation of leaf extract was obtained with a Symmetry C₁₈ column (250 mm x 4.6 mm, 5 μm, Waters). The effect of mobile phase was also examined. In initial experiments, mixtures composed of methanol and water acidified with acetic acid (pH=3.0) in different ratios were tested. This mobile phase led to bad resolution and an unstable baseline. Satisfactory separation was obtained with a mobile phase consisting of 25 mM potassium phosphate buffer-acetonitrile in a ratio of 8 /2, v/v. A pH value of 3.0 was selected for the optimal separation of the compound in the extracts; the pH was adjusted with orthophosphoric acid. The retention time of majdine was observed at 15.6 min. Measurement at 225 nm displayed sufficient sensitivity and a satisfactory chromatographic baseline.

In the present study, quantification with an internal standard was used because majdine was not commercially available. Several alkaloids, such as palmatine, berberine, caffeine, papaverine, quinine and chelidonine, were tested as internal standard. Among the alkaloids, papaverine was chosen because it was stable, commercially available and displayed a suitable chromatographic retention time. The chromatogram showed complete baseline separation of papaverine (2) and majdine (1) in *V. herbacea* (Figure 2).

The method was validated according to the ICH guidelines on the validation of analytical methods. The specificity was investigated. HPLC chromatogram of the sample solution revealed no interference with majdine and other compounds in the extract of *V. herbacea*. The peak purity was verified using Agilent Chemstation software.

The calibration curves were plotted by correlating the area ratio ($y = \text{majdine/internal standard}$) versus the corresponding concentration ratio ($x = \text{majdine/internal standard}$). Assay results of majdine standard (0.006-0.019 mg/mL, $n=15$) were linear within the studied concentration range. The regression equation was $y = 1.5812x - 0.0221$, and the correlation coefficient (r^2) 0.9978.

The intra-day (3 days, $n=6$) and inter-day precisions ($n=18$) were assessed on dried leaves of *V. herbacea*. The




Figure 2: HPLC profile of *Vinca herbacea* leaf extract collected in May 2008. Peaks: 1, majdine; 2, internal standard.

Table 1: Content of majdine in extracts prepared from different parts of *V. herbacea* collected in April/May 2008.

Plant part		Content of majdine %
April	Leaves	3.0
	Stems	0.2
	Roots	1.1
May	Leaves	11.9
	Stems	7.5
	Roots	0.7

data obtained for the repeatability were: day 1 = 12.2% (RSD % = 5.1), day 2 = 11.4% (RSD % = 3.7), day 3 = 12.0% (RSD % = 4.3). The intermediate precision was 11.9 %, RSD % = 5.3%. The mean recoveries of majdine ranged from 94.4 (± 0.1) to 105.9 (± 0.1)%. The limit of quantification and the limit of detection were 13 ng/mL and 65 ng/mL, respectively.

The validated HPLC-DAD method was applied to the determination of majdine in extracts from different parts of *V. herbacea* collected during the flowering period (Table 1). The concentration varied significantly according to the different plant parts. The highest content (11.9%) was found in the leaves, which are considered to be the best raw material.

The content of majdine was also studied during two consecutive months of the flowering period. Important variations were noted between April and May. The content of the leaves was four times higher in May than in April, and forty times higher in the stems. The best harvesting period for the leaves seems to be in May. These results could be completed by analyzing samples from different locations of Georgia and collected at different periods to confirm these first results.

The proposed HPLC-DAD method is simple, linear, accurate, precise and specific. The validation procedure confirmed that the method afforded reliable analyses of majdine. Finally, the developed method has been applied successfully to quantify majdine in different parts of the plant.

Experimental

Plant material: Different parts (leaves, roots, stems) of *Vinca herbacea* were collected in Shiraki, East Georgia in

April and May of 2008. The plant was identified by B. Grigolava (Institute of Pharmacochimistry, Tbilisi, Georgia) and a voucher specimen n° 9 (120) was deposited in the herbarium of the Institute of Pharmacochimistry. The freshly picked parts were air dried at room temperature for 3 weeks and kept in the dark until chemical analysis. The samples were ground to a fine powder before analysis.

Chemicals and reagents: HPLC grade acetonitrile and other solvents of analytical grade were purchased from Carlo Erba (Val de Reuil, France). Potassium dihydrogen phosphate was purchased from Fluka (Saint Quentin Fallavier, France). Ultrapure water (18,2M Ω) for HPLC analysis was obtained from an Elga purelab Classic purification system (Elga, France). Papaverine hydrochloride was purchased from Sigma (ref P3510, purity \geq 98%). Majdine was isolated from *Vinca herbacea* and the purity, determined by HPLC, was 98%.

Chromatographic conditions: The HPLC system consisted of an Agilent Technologies Model 1100 liquid chromatograph, equipped with a vacuum degasser, a quaternary pump, an autosampler and a photodiode array detector (DAD). The system was piloted by Chemstation computer software. The chromatographic separation was achieved using a Symmetry C18 column (250 mm x 4.6 mm, 5 μ m, Waters). The isocratic mobile phase was composed of 25 mM phosphate buffer (pH=3.0, adjusted with orthophosphoric acid) and acetonitrile (8/2, v/v). The mobile phase flow rate was 1mL/min. The injection volume was 20 μ L. UV spectra were recorded in the range 200-400 nm for all peaks. Quantification was carried out at a single wavelength of 225nm.

Standard preparation: A stock solution containing 0.425 mg/mL of internal standard in methanol was used. Standard stock solution of majdine was prepared in methanol to give a final concentration of 0.163 mg/mL. A series of working solutions (n=5) was prepared in order to obtain various concentration levels (0.006-0.019 mg/mL). The appropriate volume of stock solution of majdine was introduced to a 25 mL volumetric flask, 1 mL of internal standard solution was added, and the volume was adjusted to 25.0 mL with the mobile phase. The final concentration of papaverine in each standard solution was 0.017 mg/mL. All prepared standard solutions were filtered through 0.45 μ m membrane filter (Millipore, ref HVPL04700) before HPLC analysis.

Sample preparation: Powdered plant material (100 g) was moistened with water alkalized with ammonia (100/5, v/v). The material was macerated for 4 h and then percolated with methylene chloride at room temperature. Liquid/liquid extraction was performed with 2N HCL. The aqueous layer was alkalized with ammonia (28%) and extracted with methylene chloride. The organic layer was collected and evaporated to dryness to yield the total alkaloids (0.4%). Ten mg of the total alkaloids was introduced to a 100 mL volumetric flask and solubilized in the mobile phase. Afterwards, 1.0 mL of the internal standard solution was added to obtain a final concentration of 0.017 mg/mL of papaverine, and the final volume was adjusted to 100.0 mL with the mobile phase. Then, 2 mL of each solution was filtered through a syringe filter (0.45 mm Millipore) into a HPLC vial.

Validation and assay: A validation procedure was performed to check the performance of the chromatographic technique. The method was validated on a leaf extract of *V. herbacea* according to the ICH guidelines [18]. The specificity was performed by comparing the UV spectrum and retention time of majdine in the sample preparation and the reference compound solution. The linearity of the HPLC method was evaluated by analyzing 5 different concentrations of majdine standard solutions. Each concentration was assayed in triplicate. The limits of detection and quantification were considered to be the concentrations that produced signal-to-noise ratios of 3:1 and 10:1, respectively.

The precision of the method was evaluated with respect to both intra- and inter-day precision. Intra-day precision was calculated from the analysis of 6 sample solutions, which were prepared independently on the same day. Inter-day precision was evaluated by replicating the procedure on 2 consecutive days. The standard deviation and RSD (relative standard deviation) values were calculated for each day.

The accuracy of the method was evaluated using the recovery test. This involved the spiking of known quantities of majdine standard solutions into the tested samples. The standard solutions were prepared at 3 concentration levels (50, 100 and 150%). At each level, samples were analyzed in triplicate according to the established chromatographic conditions.

Acknowledgments - We thank Dr M. De Meo for stylistic and grammatical corrections.

References

- [1] Vachnadze VY, Malikov VM, Mudzhiri KS, Yunusov SY. (1971) A study of the alkaloids of *Vinca herbacea*. *Khimiya Prirodnikh Soedinenii*, **5**, 661-662.
- [2] Ognyanov I, Pyuskyulev B, Kompiš I, Sticzay T, Spiteller G, Shamma M, Shine RJ. (1968) Alkaloids from *Vinca herbacea* W.K. The structures and stereochemistry of majdine and isomajdine. *Tetrahedron*, **24**, 4641-4648.
- [3] Chikvadze G. (1985) Dynamics of the accumulation of alkaloids in *Vinca herbacea* cultivated in eastern Georgia. *Chemistry of Natural Compounds*, **21**, 681-682.

- [4] Chikvadze G. (1978) Isoreserpinine from *Vinca herbacea*. *Chemistry of Natural Compounds*, **14**, 420-422.
- [5] Ognyanov J, Dalev P, Dutschevska H, Mollov M. (1964) Neue alkaloides aus *Vinca herbacea* W. et K C.R. *Bulgarian Academy of Sciences*, **17**, 153-156.
- [6] Ognyanov J, Pyuskyulev B, Bonjanov B, Hesse M. (1967) Alkaloides aus *Vinca herbacea* W et K, mittelung isolierung und structur von Hervin. *Helvetica Chimica Acta*, **50**, 754-759.
- [7] Pyuskyulev B, Ognyanov J, Panov P. (1967) Alkaloides aus *Vinca herbacea*, XI, Norfluokurarine. *Tetrahedron Letters*, **4**, 4559-4562.
- [8] Rusinov K, Zhelyazkov D, Georgiev V. (1962) Pharmacological investigation of the alkaloids of *Vinca herbacea*. *Bulgaria Akademii Nauk*, **5**, 271-306.
- [9] Roussinoff K, Jelyazkoff D, Guerguieff V. (1961) The blocking action of the alkaloids of *Vinca herbacea* on the neuromuscular synapses. *Archivio Italiano di Scienze farmaceutiche*, **11**, 83-91.
- [10] Chkhikvadze G, Vachnadze V, Gambashidze N, Gvishiani E, Mudjiri L. (1981) The method of receipt of compound with antiarrhythmic activity, USSR patent SU 834964, Russian.
- [11] Chkhikvadze G, Gambashidze N, Gvishiani E, Vachnadze V, Giorgadze N. (1982) Chemical and pharmacological investigation of alkaloids of herb *Vinca Herbacea* W. and Kit. Presented at conference *Creation, investigation and complex utilization of medicinal drugs for therapy of cardio vascular diseases*, Tbilisi, Georgia.
- [12] Novikova Z, Gotsiridze I, Abuladze G. (1984) Comparative study of pharmacological activity of individual and combined alkaloids of *Vinca herbacea*. *Izvestiia Akademii Nauk SSSR Seriya Biologicheskaya*, **10**, 54-55.
- [13] Gagua N, Chkhikvadze G, Vachnadze N, Vachnadze V, Bakuridze A. (2010) Vingerbin an indolin alkaloid preparation with antiarrhythmic activity produced by *Vinca herbacea* W. and K. *Allergology and Immunology Journal*, **11**, 160-162.
- [14] Hava M. (1973) The pharmacology of *Vinca* species and their alkaloids. In *Vinca alkaloids*. Taylor WI, Farnsworth N (Eds). Marcel Dekker, New York, USA, 305-338.
- [15] Gülçin İ, Gagua N, Bursal E, Bakuridze A, Beydemir Ş, Gepdiremen A, Bayram R. (2011) Apoptotic, antioxidant, antiradical and antiglaucoma effects of majdine and isomajdine from *Vinca herbacea* Waldst. and Kit. *Journal of Enzyme Inhibition and Medicinal Chemistry*, in press.
- [16] Yongjijian W, Duowel L, Minjie P, Wei L. (2009) Determination of vincamine by RP-HPLC. *Xibex Yoxue Zazhi*, **24**, 23-24.
- [17] El-Saharty, Yasser Shaker I. (2008) Simultaneous determination of piracetam and vincamine by spectrophotometric and high-performance liquid chromatographic methods. *Journal of Association of Official Analytical Chemists International*, **91**, 311-321.
- [18] *Validation of analytical procedures: Text and methodology* - International Conference on Harmonisation. ICH-Q2 (R1) (2005) Geneva.

Pyridine Metabolism and Trigonelline Synthesis in Leaves of the Mangrove Legume trees *Derris indica* (*Millettia pinnata*) and *Caesalpinia crista*

Yuling Yin^{a,b}, Hamako Sasamoto^c and Hiroshi Ashihara^{a,*}

^aDepartment of Biological Sciences, Graduate School of Humanities and Sciences, Ochanomizu University, Tokyo, 112-8610, Japan

^bDepartment of Biotechnology, Dalian Medical University, Dalian, 116044, China

^cFaculty of Environment and Information Sciences, Yokohama National University, Yokohama, Kanagawa, 240-8501, Japan

ashihara.hiroshi@ocha.ac.jp

Received: September 29th, 2011; Accepted: October 5th, 2011

The aim of this study was to reveal the pyridine metabolism in leaves of two mangrove legumes, *Derris indica* (= *Millettia pinnata* or *Pongamia pinnata*) and *Caesalpinia crista*. Radioactivity from [*carboxyl*-¹⁴C]nicotinamide supplied exogenously to young leaf disks was recovered in nicotinic acid, nicotinic acid mononucleotide, NAD, NADP, nicotinamide mononucleotide and trigonelline. These mangrove species, especially *D. indica*, have strong ability to convert nicotinamide to trigonelline, but not to nicotinic acid glucoside. The endogenous trigonelline content in leaves of *D. indica* was more than 830 µg/g dry weight. This value is 5-12 times greater than that in leaves of *Glycine max*. There was little short-term effect of 250 and 500 mM NaCl (equivalent to *ca.* 50% and 100% sea water) on nicotinamide metabolism.

Keywords: Nicotinamide adenine dinucleotide, nicotinamide, nicotinic acid, salt stress, trigonelline.

Derris indica and *Caesalpinia crista* are mangrove plants belonging to the Leguminosae family. They are widely distributed on seashores in Southeast Asia and Pacific Islands. Different organs of these plants possess antimycobacterial activity [1] and have been used in folk medicine. Seeds of *D. indica* contain oils and fatty acids suitable for biodiesel production [2]. These plants grow by seashores with their roots in the sea. They therefore apparently have salt-tolerant properties. These mangrove legumes are interesting from the viewpoint of biotechnology as well as basic plant biology. There have, however, been few physiological studies of these plants.

Nicotinamide is formed as a catabolite of NAD and NADP, and is a key metabolite of pyridine metabolism. In plants, nicotinamide is not directly used for re-synthesis of NAD, as it is in animals [3], but it is readily deaminated to nicotinic acid and then used for pyridine nucleotide synthesis [4-6]. As well as the salvage pathway for NAD synthesis, nicotinic acid is used for the synthesis of secondary metabolites. In a few plants, nicotinic acid is a substrate for the synthesis of unique pyridine alkaloids, such as nicotine and ricinine [7,8], but most plants produce simple and common secondary products: trigonelline (1-*N*-methylnicotinic acid) or nicotinic acid *N*-glucoside (Figure 1) [9,10].




Figure 1: Possible metabolic pathways of nicotinamide in *Caesalpinia crista* and *Derris indica*.

The biosynthesis of these two compounds is an example of biochemical evolution in plant chemosystematics [11]. Diverse physiological functions of trigonelline have been proposed by several researchers [9-10,12], and one role of trigonelline is as a compatible solute to adjust the osmotic pressure when the plants are affected by salt stress [13-16].

As part of our studies of pyridine metabolism in plants, we examined the metabolic fate of [*carbonyl*-¹⁴C]nicotinamide in leaves of two mangrove legumes in the presence and

absence of NaCl. The short-term effect of NaCl on endogenous trigonelline content was also examined in leaf segments of *D. indica*.

We examined the changes in ^{14}C -labelled metabolites, along with the incubation time. Figure 2 shows the results from leaf disks of *C. crista*. [^{14}C]Nicotinamide, taken up by the disks, was readily converted into nicotinic acid. Within 30 min of administration, radioactivity was found in NaMN, NAD and NADP. This implies that nicotinic acid derived from nicotinamide was salvaged to NaMN by nicotinate phosphoribosyltransferase (EC 2.4.2.11), and was then utilized for pyridine nucleotide synthesis.




Figure 2: Time-course studies of [carbonyl- ^{14}C]nicotinamide in leaf disks of *Caesalpinia crista*. Rates of incorporation \pm SD are expressed as a percentage of total radioactivity. The total radioactivity (kBq per 100 mg fresh weight) taken up at 0.5, 1, 3, 6 and 24 h after administration was respectively 1.8 ± 0.1 , 2.6 ± 0.1 , 6.1 ± 0.1 , 11.0 ± 0.7 and 20.6 ± 0.3 .

Radioactivity was also found in trigonelline; the proportion gradually increased with incubation time to 60% of the total radioactivity at 24 h after initiation. In these disks, therefore, nicotinic acid appears to be converted to trigonelline by nicotinic acid *N*-methyltransferase (trigonelline synthase, EC 2.1.1.7); this pyridine alkaloid was accumulated. Release of $^{14}\text{CO}_2$ from [carbonyl- ^{14}C]nicotinamide was low; less than 1.2% of total activity was found during the time of incubation. Small amounts of radioactivity were found in NMN at 6 h after administration of [^{14}C]nicotinamide. Although it is difficult to infer the metabolic sequence from these results alone, our data support the functioning of the pyridine nucleotide cycle shown in Figure 1, which has been reported in other plants [5,16-17].

We examined the metabolic fate of [carbonyl- ^{14}C]nicotinamide in leaf disks from *C. crista* and *D. indica* at 3 h after administration in the presence and absence of NaCl (Figure 3A,B). In leaf disks of both species the radioactivity was recovered in nicotinamide, nicotinic acid, NAD/NADP, NaMN and trigonelline. Only a small amount of radioactivity was found in CO_2 . No radioactivity was detected in nicotinic acid glucoside.




Figure 3: Metabolic fate of [carbonyl- ^{14}C]nicotinamide in leaf disks of *Caesalpinia crista* (A) and *Derris indica* (B) in the presence of 0, 250 and 500 mM NaCl. Incubation took place for 3 h. Values \pm SD ($n = 3$) are expressed as % of total uptake by leaf disks. Total radioactivity (kBq per 100 mg fresh weight) taken up by the leaf disks treated with 0, 250 and 500 mM NaCl was respectively 6.1 ± 0.5 , 6.1 ± 0.4 and 7.8 ± 0.2 (A) and 9.0 ± 0.3 , 7.5 ± 0.1 and 7.2 ± 0.4 (B).

The activity of pyridine salvage, estimated from the sum of the incorporation rate of radioactivity into NAD, NADP and NaMN, varied from 18% to 22% of the total uptake in leaf disks of both species, and no large difference was caused by NaCl. In contrast, widely differing trigonelline synthesis activity was found in the different species; the incorporation rate in *D. indica* (42-47% of total uptake) was much greater than in *C. crista* (15-17%). No increase in trigonelline synthesis due to NaCl took place, and there was even a slight reduction in activity (88% of control) in *D. indica*.

The endogenous concentration of trigonelline in leaf disks of *D. indica* at 3 h after incubation with 0, 250 and 500 mM NaCl was respectively 831.5 ± 53.3 , 857.0 ± 22.0 and 893.9 ± 13.7 μg per g dry weight. Cho *et al.* [13] reported that the trigonelline concentration in leaf tissue of cultivated *Glycine max* ranged from 64 to 162 μg per g dry weight. Trigonelline concentration in *D. indica* leaves is consequently 5-12 times higher than in *G. max* leaves. High concentrations of trigonelline may act as compatible

solutes in this mangrove legume. Although we found that the concentration of trigonelline increased slightly as a result of 3 h salt stress, these differences were not statistically significant ($P > 0.05$).

Our observations suggest that trigonelline synthesis in mangrove leaves does not respond quickly to salt stress. However, it is clear that *D. indica* has a mechanism for producing trigonelline, so that its content in leaves is much greater than in *G. max*, even if the plant is grown in a salt-free environment. It is, therefore, plausible that the strong trigonelline formation in *D. indica* is an inherent property of this mangrove legume.

Experimental

Plant materials and chemicals: Seeds of *Derris indica* were collected from the basin of the Khanom River, and seeds of *Caesalpinia crista* from the Pang Yi River basin, both in Thailand. Young leaves of 6-month-old seedlings grown in a greenhouse were used as plant materials. [Carbonyl- ^{14}C]nicotinamide (2.04 GBq/mmol) was purchased from Moravak Biochemicals Inc., Brea, CA, USA, and other chemicals were from either Sigma-Aldrich, St. Louis, MO, USA, or from Wako Pure Chemical Industries Ltd., Osaka, Japan.

Radioactivity incorporation: Administration of [carbonyl- ^{14}C]nicotinamide to the leaf disks and analysis of labeled metabolites were essentially the same as in a previous paper [18]. Leaf disks (ca. 100 mg fresh weight), together with 2.0 mL of 30 mM potassium phosphate buffer (pH 5.6) containing 10 mM sucrose, 50 mM sodium ascorbate, and 0, 250 or 500 mM NaCl, were placed in the main compartment of a 30 mL Erlenmeyer flask. The flask was fitted with a glass tube containing a piece of folded filter paper that had been impregnated with 0.1 mL of 20% KOH in the centre well. Each reaction was started by adding 10 μL (37 kBq) of a solution of [carbonyl- ^{14}C]nicotinamide to the main compartment of the flask. The flasks were incubated in an oscillating water bath at 27°C. After incubation, the glass tube was removed from the centre well and placed in a 50 mL Erlenmeyer flask containing 10 mL distilled water. At the same time, the leaf disks were harvested and washed with distilled water, frozen with liq. N_2 , and then stored at -80°C prior to extraction. The filter paper, containing CO_2 trapped as potassium bicarbonate, was soaked overnight in distilled water (10 mL), and aliquots of the resulting solution (usually 0.5 mL) were used for determination of radioactivity. Radioactivity was measured with a liquid scintillation counter (Beckman, type LS 6500, Fullerton, CA, USA).

Analysis of metabolites: For analysis of ^{14}C -labelled metabolites, the frozen samples were homogenized using a mortar and pestle with ice-cold 6% perchloric acid (PCA)

containing 0.1 mM NAD. NAD was supplemented to prevent the degradation of labeled nucleotides during extraction and analysis, and also for action as endogenous markers in the TLC analysis. The homogenate was centrifuged at 3,000 $\times g$ for 7 min, the supernatant was collected, and the precipitate was re-suspended in the same extraction reagent and washed 3 times. The ^{14}C -labelled metabolites extracted with 6% PCA were combined, neutralized with 20% KOH, and centrifuged briefly (3,000 $\times g$, 7 min) to remove potassium perchlorate as precipitate. The radioactivity of aliquots of supernatant was measured with a liquid scintillation counter. The neutralized fractions were freeze-dried and then redissolved in a small amount of 50% v/v, ethanol.

The concentrated PCA-soluble metabolites (free nicotinic acid related small molecular weight compounds) were separated by using microcrystalline cellulose TLC sheets (200 \times 200 mm, Merck, Darmstadt, Germany). As solvent systems, system I (*n*-butanol: acetic acid: water, 4:1:2, v/v) and system IV (isobutyric acid: ammonia: water, 660: 17: 330, v/v) as specified by Zheng and Ashihara [19] were used. The distribution of ^{14}C to individual spots on the TLC sheet was estimated using a bio-imaging analyzer (Type, FLA-2000, Fuji Photo Film Co., Ltd. Tokyo, Japan). The total radioactivity of each fraction was determined by a liquid scintillation counter (Beckman, type LS 6500). Incorporation into individual metabolites was calculated using the total radioactivity values in the PCA-soluble fraction, as measured by a liquid scintillation counter, and the % distribution of radioactivity on TLC plates was obtained from the bio-imaging analyzer.

Effects of NaCl: To determine the short-term effect of NaCl on the accumulation of trigonelline, leaf disks (ca. 300 mg fresh weight) were incubated with 2 mL of the medium which had been used for the ^{14}C -tracer experiments, supplemented with 0, 250 or 500 mM NaCl, in a 30 mL Erlenmeyer flask at 27°C for 3 h. After incubation, the leaf disks were washed with distilled water, boiled in 10 mL of hot water (> 95°C) for 10 min, and then homogenized using a mortar and pestle. After brief centrifugation (20,000 $\times g$, for 10 min), the water-soluble supernatant fraction was collected and freeze dried. The samples were dissolved in small amounts of distilled water and analyzed by HPLC, as in our previous paper [20], except that the absorbance was monitored using a Shimadzu Diode Array Detector, type SPD-M10A (Shimadzu Corp., Kyoto, Japan).

Acknowledgements - We thank Dr Reiko Minagawa, Tokyo University of Agriculture, for her generous supply of plant materials. This research was supported by a JSPS Grant-in-Aid for Scientific Research (No. 22510226) to H.A.

References

- [1] Satnami DK, Yadava RN. (2011) Potential phytochemical from *Caesalpinia crista* Linn. *Research Journal of Phytochemistry*, **5**, 22-31
- [2] Scott P, Pregelj L, Chen N, Hadler J, Djordjevic M, Gresshoff P. (2008) *Pongamia pinnata*: An untapped resource for the biofuels industry of the future. *BioEnergy Research*, **1**, 2-11.
- [3] Revollo JR, Grimm AA, Imai S. (2004) The NAD biosynthesis pathway mediated by nicotinamide phosphoribosyltransferase regulates Sir2 activity in mammalian cells. *Journal of Biological Chemistry*, **279**, 50754-50763.
- [4] Wang G, Pichersky E. (2007) Nicotinamidase participates in the salvage pathway of NAD biosynthesis in Arabidopsis. *Plant Journal*, **49**, 1020-1029.
- [5] Zheng XQ, Hayashibe E, Ashihara H. (2005) Changes in trigonelline (*N*-methylnicotinic acid) content and nicotinic acid metabolism during germination of mungbean (*Phaseolus aureus*) seeds. *Journal of Experimental Botany*, **56**, 1615-1623.
- [6] Zrenner R, Ashihara H. (2011) Nucleotide metabolism. In *Plant Metabolism and Biotechnology*, Ashihara H, Crozier A, Komanine A. (Eds), John Wiley & Sons, Ltd, Chichester, England, 135-162.
- [7] Katoh A, Hashimoto T. (2004) Molecular biology of pyridine nucleotide and nicotine biosynthesis. *Frontier in Bioscience*, **9**, 1577-1586.
- [8] Waller GR, Yang KS, Gholson RK, Hadwiger LA, Chaykin S. (1966) The pyridine nucleotide cycle and its role in the biosynthesis of ricinine by *Ricinus communis* L. *Journal of Biological Chemistry*, **241**, 4411-4418.
- [9] Ashihara H. (2008) Trigonelline (*N*-methylnicotinic acid) biosynthesis and its biological role in plants. *Natural Product Communications*, **3**, 1423-1428.
- [10] Barz W. (1985) Metabolism and degradation of nicotinic acid in plant cell cultures, In *Primary and Secondary Metabolism of Plant Cell Cultures*, Neumann KH, Barz W, Reinhard E. (Eds) Springer-Verlag, Berlin, 186-195.
- [11] Harborne JB, Turner BL. (1984) *Plant Chemosystematics*. Academic Press, London.
- [12] Minorsky PV. (2002) Trigonelline: A diverse regulator in plants. *Plant Physiology*, **128**, 7-8.
- [13] Cho Y, Lightfoot DA, Wood AJ. (1999) Trigonelline concentrations in salt stressed leaves of cultivated *Glycine max*. *Phytochemistry*, **52**, 1235-1238.
- [14] Rajasekaran LR, Aspinall D, Jones GP, Paleg LG. (2001) Stress metabolism. IX. Effect of salt stress on trigonelline accumulation in tomato. *Canadian Journal of Plant Science*, **81**, 487-498.
- [15] Tramontano WA, Jouve D. (1997) Trigonelline accumulation in salt-stressed legumes and the role of other osmoregulators as cell cycle control agents. *Phytochemistry*, **44**, 1037-1040.
- [16] Wood AJ. (1999) Comparison of salt-induced osmotic adjustment and trigonelline accumulation in two soybean cultivars. *Biologia Plantarum*, **42**, 389-394.
- [17] Ashihara H, Stasolla C, Yin Y, Loukanina N, Thorpe TA. (2005) *De novo* and salvage biosynthetic pathways of pyridine nucleotides and nicotinic acid conjugates in cultured plant cells. *Plant Science*, **169**, 107-114.
- [18] Katahira R, Ashihara H. (2009) Profiles of the biosynthesis and metabolism of pyridine nucleotides in potatoes (*Solanum tuberosum* L.). *Planta*, **231**, 35-42.
- [19] Zheng XQ, Ashihara H. (2004) Distribution, biosynthesis and function of purine and pyridine alkaloids in *Coffea arabica* seedlings. *Plant Science*, **166**, 807-813.
- [20] Ashihara H, Yin Y, Deng WW, Watanabe S. (2010) Pyridine salvage and nicotinic acid conjugate synthesis in leaves of mangrove species. *Phytochemistry*, **71**, 47-53.

Anti-adipogenic Activity of *Cordyceps militaris* in 3T3-L1 Cells

Qing Liu^a, In Pyo Hong^b, Mi-Jeong Ahn^c, Hwan-Soo Yoo^a, Sang-Bae Han^a, Bang Yeon Hwang^a and Mi Kyeong Lee^{a,*}

^aCollege of Pharmacy, Chungbuk National University, Cheongju 361-763, Korea

^bNational Academy of Agricultural Science, Suwon, Gyunggi 441-707, Korea

^cCollege of Pharmacy, Gyeongsang National University, Jinju 660-751, Korea

mklee@chungbuk.ac.kr

Received: July 14th, 2011; Accepted: October 7th, 2011

Inhibition of adipocytes differentiation is suggested to be an important strategy for prevention and/or treatment of obesity. In our present study, *Cordyceps militaris* showed significant inhibitory activity on adipocyte differentiation in 3T3-L1 preadipocytes as assessed by measuring fat accumulation using Oil Red O staining. Activity-guided fractionation led to the isolation of cordycepin (1), guanosine (2) and tryptophan (3) as active compounds. All the three compounds were more effective in the prevention of early stage of adipogenesis than in lipolysis. In addition, combinational treatment of three compounds significantly increased anti-adipogenic activity.

Keywords: *Cordyceps militaris*, cordycepin, 3T3-L1 preadipocytes, Oil Red O staining, obesity.

Obesity has become a widespread issue in modern society, due to a global rise in obesity-associated problem. It is no longer considered only a cosmetic problem but associated with several pathological disorders, including diabetes, hypertension, atherosclerosis and cancer [1]. Adipose tissue is an important metabolic organ that is crucial for insulin sensitivity and energy homeostasis. Adipocyte differentiation is an important process for its function in normal condition [2]. In obesity, adipocytes undergo abnormal growth and differentiation, which result in increased numbers of fat cells storing their lipids and consequent pathological disorders [3]. Therefore, inhibition of adipocytes differentiation is suggested to be an important strategy for prevention and/or treatment of obesity [4].

Cordyceps, also known as ‘winter worm summer grass’, is a well-known traditional medicine in Asian countries. It has been known as a rich source of biologically active components and used for treatment of several diseases such as cancer, fatigue, hyperglycemia and hyperlipidemia [5,6]. Recently, favorable role of *Cordyceps* in the regulation of obesity has been reported [7]. However, little is known about its active components related to adipocyte differentiation.

In this study, we evaluated the anti-adipogenic activity of *Cordyceps militaris* (Hypocreaceae) employing 3T3-L1, a mouse preadipocyte cell line as an *in vitro* assay system. The *n*-BuOH fraction of *C. militaris* showed potent inhibitory activity on adipocyte differentiation at a




Figure 1: Structure of compounds isolated from *C. militaris* (1–3) and cordycepin derivatives (4, 5).

concentration of 10 µg/mL (6.5% of differentiated control, $p < 0.001$). Activity-guided fractionation of *n*-BuOH fraction resulted in the isolation of three constituents which were identified as cordycepin (1), guanosine (2) and tryptophan (3) by direct comparison of their physicochemical and spectroscopic data with those previously reported (Figure 1) [8].

The inhibitory activity of isolated compounds (1–3) on adipocyte differentiation was also evaluated in our assay system. Among the compounds isolated, guanosine (2) and tryptophan (3) showed the strong inhibitory activity, followed by cordycepin (1) (Figure 2B). At a concentration of 10 µg/mL, guanosine (2) and tryptophan (3) completely inhibited adipocyte differentiation as measured by fat accumulation, which was supported by microscopic observation (Figure 2A).




Figure 2: Effect of cordycepin, guanosine and tryptophan on fat accumulation in 3T3-L1 cells. Cultures were stained with Oil Red O, photographed (A) and quantitated (B). Results are expressed as mean \pm SD of three independent experiments, each performed using triplicate wells. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$ compared with differentiated control.

Adipogenic differentiation is a well-organized process, occurs in several stages [9]. Therefore, we further examined which stage of adipogenic differentiation was effectively inhibited by these compounds. During differentiation, 3T3-L1 cells were incubated with cordycepin, guanosine and tryptophan at different periods of differentiation, namely, during days 0–2, 2–4, 4–8, 0–4, 2–8 and 0–8. As shown in Figure 3A, cordycepin was most effect when treated during days 2–4, whereas guanosine and tryptophan significantly inhibited adipocyte differentiation when treated during days 0–2. However, all the compounds showed weak activity when treated during days 4–8. In addition, these compounds did not show significant effects when treated to differentiated adipocytes (data not shown). These results suggest that cordycepin, guanosine and tryptophan could effectively inhibit the early stage of adipocyte differentiation and might be effective in the prevention of adipogenesis but not in lipolysis.

Cordycepin is a characteristic constituent of *C. militaris* and diverse biological activity such as anticancer and neuroprotective activity [10,11]. Structurally, cordycepin has same structure with adenosine except for the absence of hydroxyl group at C-3' (Figure 1). Adipocytes express adenosine A1 receptor and activation of adenosine A1 receptor by agonists inhibits lipolysis and enhances the lipogenesis, whereas antagonists exerts opposite action [12,13]. Therefore, we compared the effect of cordycepin and other adenosine derivatives such as adenosine and 2'-deoxyadenosine. Consistent with previous report adenosine slightly increased fat accumulation at the concentration of 10 $\mu\text{g/mL}$. In addition, 2'-deoxyadenosine also failed to reduced fat accumulation. Therefore, we suppose that the absence of hydroxyl group at C-3' may play an important role in inhibitory activity of cordycepin on adipocyte differentiation, and cordycepin might act as a




Figure 3: Effect of cordycepin, guanosine and tryptophan on adipocyte differentiation during differentiation process (A) and combinational treatment (B). Compounds were added at different periods of differentiation, namely, during days 0–2, 2–4, 4–8, 0–4, 2–8 and 0–8 (A). In other cultures, Cordycepin, guanosine and tryptophan were treated with combination as indicated (B). On day 8, cells were subjected to Oil Red O staining. Values are expressed as means \pm SD of triplicate experiments. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$ compared with differentiated control.

partial antagonist of adenosine A1 receptor, which needs to be clarified by further investigation.

Natural products are known to exert strong biological activity by additive effects of its constituents. Therefore, combinational effect of cordycepin, guanosine and tryptophan was investigated. Interestingly, combination of each compounds strongly increased anti-adipogenic activity (Figure 3B). Although each compound showed little effect at the concentration of 1 $\mu\text{g/mL}$, combination of two or three compounds showed potent inhibition on fat accumulation. These results suggest that combinational action of each compound contributes to the anti-adipogenic activity of *C. militaris*.

Experimental

Extraction and isolation: Dried *Cordyceps* samples were provided from Rural Development Administration in November 2009. A voucher specimen (CBNU-2009-CM) has been deposited in the Herbarium of College of Pharmacy, Chungbuk National University. The powdered *C. Militaris* were extracted 3 times with EtOH, which yielded the total ethanolic extract. The ethanolic extract was then suspended in H_2O and partitioned successively with *n*-hexane, CHCl_3 and *n*-BuOH. The *n*-BuOH fraction, which showed most potent activity, was subjected to XAD

column chromatography with the mixture of MeOH–H₂O (0%, 20%, 40%, 60%, 80%, 100% MeOH in H₂O) to give 6 fractions (B1–B6). B2 was subjected to column chromatography over Sephadex LH-20 with MeOH to give 9 subfractions (B21–B29). Compound **1** was obtained from B24 by recrystallization using MeOH. Compounds **2** and **3** were isolated from B24 by semipreparative HPLC eluting with MeOH–water (23:77, F=2 min/mL, Rt=7.67 and 8.65 min, respectively).

Cell culture and differentiation induction: 3T3-L1 mouse embryo fibroblasts were obtained from the American Type Culture Collection. Cells were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS) until confluence. Two days after confluence (day 0), cells were stimulated to differentiate with differentiation medium containing DMEM with 10% FBS, 0.5 mM 3-isobutyl-1-methyl-xanthine, 1 µg/mL insulin and 1 µM dexamethasone for 2 days (day 2). Cells were then maintained in DMEM supplemented with 10% FBS and 1 µg/mL for another 2 days (day 4), followed by culturing with DMEM with 10% FBS for an additional 4 days (day 8). All media contained 100 IU/mL penicillin and 100 µg/mL streptomycin. Cells were maintained at 37°C in a humidified atmosphere of 95% air–5% CO₂ [14].

Oil Red O staining: Lipid droplets in cells were stained with Oil Red O. Eight days after differentiation induction, cells were washed three times with PBS and fixed with 10% formalin at room temperature for 1 hr. After fixation, cells were washed twice with PBS and stained with freshly diluted Oil Red O solution (3 parts of 0.6% Oil Red O in isopropyl alcohol and 2 parts of water) for 15 min. Cells were then washed twice with water and visualized. For quantitative analysis, Oil Red O staining was dissolved with isopropyl alcohol and optical density was measured at 550 nm by ELISA plate reader [14].

Statistical analysis: The evaluation of statistical significance was determined by one-way ANOVA test with a value of $p < 0.05$ or less considered to be statistically significant.

Acknowledgments –This work was supported by the grant from the Korean Rural Development Administration (Agenda Program, PJ0067792010) and the grant of the Korean Ministry of Education, Science and Technology (The Regional Core Research Program/Chungbuk BIT research-Oriented University Consortium).

References

- [1] Kopelman PG. (2000) Obesity as a medical problem. *Nature*, **404**, 635–643.
- [2] Rosen ED, Spiegelman BM. (2006) Adipocytes as regulators of energy balance and glucose homeostasis. *Nature*, **444**, 847–853.
- [3] Spiegelman BM, Enerback S. (2006) The adipocyte: a multifunctional cell. *Cell Metabolism*, **4**, 425–427.
- [4] Rayalam S, Cella-Fera MA, Baile CA. (2008) Phytochemicals and regulation of the adipocyte life cycle. *Journal of Nutrition and Biochemistry*, **19**, 717–726.
- [5] Koh JH, Kim KM, Kim JM, Song JC, Suh HJ. (2003) Antifatigue and antistress effect of the hot-water fraction from mycelia of *Cordyceps sinensis*. *Biological & Pharmaceutical Bulletin*, **26**, 691–694.
- [6] Lee H, Kim YJ, Kim HW, Lee DH, Sung MK, Park T. (2006) Induction of apoptosis by *Cordyceps militaris* through activation of caspase-3 in leukemia HL-60 cells. *Biological & Pharmaceutical Bulletin*, **29**, 670–674.
- [7] Shimada T, Hiramatsu N, Kasai A, Mukai M, Okamura M, Yao J, Huang T, Tamai M, Takahashi S, Nakamura T, Kitamura M. (2008) Suppression of adipocyte differentiation by *Cordyceps militaris* through activation of the aryl hydrocarbon receptor. *American Journal of Physiology*, **295**, E859–E867.
- [8] Sawai H, Imai J, Lesiak K, Johnson MI, Torrence PF. (1983) Cordycepin analogues of 2-5A and its derivatives. *Journal of Biological Chemistry*, **258**, 1671–1677.
- [9] Lefterova MI, Lazar MA. (2009) New developments in adipogenesis. *Trends in Endocrinology and Metabolism*, **20**, 107–114.
- [10] Cheng Z, He W, Zhou X, Lv Q, Xu X, Yang S, Zhao C, Guo L. (2011) Cordycepin protects against cerebral ischemia/reperfusion injury *in vivo* and *in vitro*. *European Journal of Pharmacology*, **664**, 20–28.
- [11] Lee EJ, Kim WJ, Moon SK. (2010) Cordycepin suppresses TNF- α -induced invasion, migration and matrix metalloproteinase-9 expression in human bladder cancer cells. *Phytotherapy Research*, **24**, 1755–1761.
- [12] Liang HX, Belardinelli L, Ozeck MJ, Shryock JC. (2002) Tonic activity of the rat adipocyte A1-adenosine receptor. *British Journal of Pharmacology*, **135**, 1457–1466.
- [13] Szkudelski T, Szkudelska K, Nogowski L. (2009) Effect of adenosine A1 receptor antagonism on lipogenesis and lipolysis in isolated rat adipocytes. *Physiological Research*, **58**, 863–871.
- [14] Shin EJ, Choi KM, Yoo HS, Lee CK, Hwang BY, Lee MK. (2010) Inhibitory effects of coumarins from the stem barks of *Fraxinus rhynchophylla* on adipocyte differentiation in 3T3-L1 cells. *Biological & Pharmaceutical Bulletin*, **33**, 1610–1614.

Two New Cyclopeptides and One New Nonenolide from *Xylaria* sp. 101Yao-Yao Li^a, Zhi-Yu Hu^b, and Yue-Mao Shen^{a*}^aSchool of Pharmaceutical Sciences, Shandong University, Jinan, Shandong 250000, P. R. China^bSchool of Life Sciences, Xiamen University, Xiamen, Fujian 361005, P. R. China

yshen@xmu.edu.cn

Received: August 1st, 2010; Accepted: October 26th, 2011


Two novel cyclopeptides, xylarotides A (**1**), and B (**2**), and one novel nonenolide, xylarolide (**3**), along with two known compounds, coriloxin (**4**), and 2-hydroxy-3-methoxy-5-methyl-*p*-benzoquinone (**5**) were isolated from the fungal strain *Xylaria* sp. 101. This strain was isolated from the fruiting body of *Xylaria* sp. collected in Gaoligong Mountain, Yunnan Province. The chemical structures were elucidated by spectroscopic analyses, including 1D- and 2D-NMR spectroscopic experiments, and on the basis of HR-Q-TOF mass spectrometry. Antibacterial assays of **1** - **3** were carried out; no effects on the growth of the tested bacteria and yeast were observed.

Keywords: *Xylaria* sp., cyclopeptides, nonenolide.

Fungi of the genus *Xylaria* are a rich source of bioactive metabolites, including terpenoids [1-5], cyclopeptides [6,7], polyketides [8,9], cytochalasins [10], xanthenes [11,12], and unique unclassified xyloketal [13]. We isolated a fungal strain named 101 from Gaoligong Mountain of southwestern China, and identified it as *Xylaria* sp. (family Xylariaceae) based on its complete ITS1-5.8S-ITS2 sequences. Previously, four novel terpenoids were isolated from the still suspension cultures of *Xylaria* sp. 101 in potato-dextrose (PD) medium [5]. Here we report the isolation and structure elucidation of two novel cyclopeptides, namely xylarotide A (**1**) and xylarotide B (**2**), and one novel nonenolide, xylarolide (**3**), plus the previously known compounds, coriloxin (**4**) [14], and 2-hydroxy-3-methoxy-5-methyl-*p*-benzoquinone (**5**). Antibacterial assays of **1** - **3** were also performed.

The morphological properties of the isolate 101 were examined after incubation for 2 months at 28°C in potato-dextrose agar (PDA) medium. The strain 101 was identified as *Xylaria* sp. according to its ITS rDNA sequence (ITS1-5.8S-ITS2). The culture of 101 was concentrated and extracted successively with AcOEt. This extract was purified by repeated column chromatography (RP-18, Sephadex LH-20, and silica gel) to obtain compounds **1** - **5**.



Compound **1** was obtained as a white amorphous solid. The molecular formula C₂₉H₅₁N₅O₅ was determined by HR-Q-TOF MS and NMR spectroscopic data (Table 1). The IR spectrum indicated the presence of NH groups (3287cm⁻¹) and C=O groups (1640 cm⁻¹), respectively. The ¹³C NMR spectrum of **1** (Table 1) displayed five C=O signals. By extensive analysis of the HSQC, HMBC,





¹H-¹H COSY and NOE spectra, five amino acid residues were established, and shown to be one proline, one leucine, one *N*-methyl-leucine, one isoleucine, and one valine. The seven degrees of unsaturation were deduced from the molecular formula, and attributed to five C=O groups, a cyclic moiety in proline, and the last one suggesting that **1** was a cyclopeptide. The sequence of these amino acid residues was elucidated on the basis of HMBC and NOE correlations (Figures 1 and 2). The HMBC correlations from α -H of proline to C=O of leucine, from α -H of leucine to C=O of isoleucine, from α -H of isoleucine to C=O of valine, from α -H of valine to C=O of *N*-methyl-leucine, and from α -H and *N*-methyl of *N*-methyl-leucine to C=O of proline, together with the NOE correlations between NH of leucine and α -H of isoleucine, and NH of isoleucine and α -H of valine, indicated the sequential connections of these amino acid residues. Thus, compound **1** was established to be cyclo-(Pro-Me-Leu-Val-Ile-Leu), and named as xylaropeptide A.

Table 1: ^1H - and ^{13}C -NMR spectroscopic data of **1** and **2** (600/150 MHz, resp., in CDCl_3 ; δ in ppm, J in Hz).

1				2			
		δ_{H} (multiplicity, J in Hz)	δ_{C}		δ_{H} (multiplicity, J in Hz)	δ_{C}	
Pro ¹	CO		172.3 (s)	Pro ¹	CO		172.5 (s)
	CH (α)	5.15 (d, $J = 6.5$)	59.5 (d)		CH (α)	5.05 (d, $J = 6.5$)	58.9 (d)
	CH ₂ (β)	2.11-2.18 (m), 1.88-1.91 (m)	31.1 (t)		CH ₂ (β)	1.94-2.01 (m), 1.30-1.33 (m)	30.5 (t)
	CH ₂ (γ)	1.86-1.90 (m), 1.65-1.68 (m)	21.0 (t)		CH ₂ (γ)	1.78-1.82 (m), 1.61-1.64 (m)	20.9 (t)
	CH ₂ (δ)	3.88 (t, $J = 8.5$)	45.7 (t)		CH ₂ (δ)	3.77 (br d, $J = 10.5$)	45.9 (t)
Me-Leu ²		3.43 (br d, $J = 9.0$)		Me-Phe ²		3.45 (br d, $J = 7.5$)	
	CO		171.7 (s)		CO		170.4 (s)
	N-CH ₃	3.18 (s)	30.4 (q)		N-CH ₃	3.15 (s)	30.6 (q)
	CH (α)	4.64 (t, $J = 6.5$)	55.3 (d)		CH (α)		57.8 (d)
	CH ₂ (β)	1.62-1.64 (m)	37.4 (t)		CH ₂ (β)	3.18-3.20 (m), 3.04-3.11 (m)	34.5 (t)
	CH (γ)	1.55-1.57 (m)	24.8 (d)		C (γ)		136.8 (s)
	CH ₃ (δ)	0.91 (overlap)	22.1 (q)		CH (δ) $\times 2$	7.21 (d, $J = 6.0$)	128.8 (d)
Val ³	CH (δ)	0.98 (d, $J = 5.5$)	22.8 (q)		CH (δ) $\times 2$	7.25 (d, $J = 6.6$)	128.5 (d)
					CH (ϵ)	7.20 (d, $J = 6.0$)	126.8 (d)
	CO		171.4 (s)	Val ³	CO		171.7 (s)
	NH	7.67 (br s)			NH		
	CH (α)	4.11 (t, $J = 8.5$)	58.1 (d)		CH (α)	4.06 (t, $J = 8.0$)	58.3 (d)
Ile ⁴	CH (β)	2.37-2.41 (m)	26.8 (d)		CH (β)	2.17-2.20 (m)	27.1 (d)
	CH ₃ (γ)	0.94 (d, $J = 5.0$)	18.4 (q)		CH ₃ (γ)	0.75 (d, $J = 4.5$)	18.2 (q)
	CH ₃ (γ)	0.95 (d, $J = 5.0$)	19.7 (q)		CH ₃ (γ)	0.88 (overlap)	19.7 (q)
	CO		171.2 (s)	Ile ⁴	CO		171.0 (s)
	NH	6.78 (d, $J = 7.8$)			NH		
Leu ⁵	CH (α)	4.55 (t, $J = 7.0$)	56.8 (d)		CH (α)	4.33 (br s)	57.0 (d)
	CH (β)	1.62-1.65 (m)	38.0 (d)		CH (β)	1.67-1.70 (m)	37.5 (d)
	CH ₂ (γ)	1.50-1.54 (m), 1.09-1.14 (m)	25.8 (t)		CH ₂ (γ)	1.40-1.47 (m), 1.09-1.12 (m)	25.9 (t)
	CH ₃ (δ)	0.85 (d, $J = 6.0$)	14.4 (q)		CH ₃ (δ)	0.87 (overlap)	14.5 (q)
	CH ₃ (δ)	0.87 (overlap)	11.3 (q)		CH ₃ (δ)	0.91 (overlap)	11.3 (q)
Leu ⁵	CO		169.5 (s)	Leu ⁵	CO		169.5 (s)
	NH	7.67 (br s)			NH		
	CH (α)	4.91 (q, $J = 6.5$)	47.4 (d)		CH (α)	4.85 (q, $J = 6.0$)	47.7 (d)
	CH ₂ (β)	1.62-1.65 (m)	42.1 (t)		CH ₂ (β)	1.64-1.68 (m)	42.2 (t)
		1.56-1.59 (m)				1.50-1.54 (m)	
	CH (γ)	1.53-1.56 (m)	25.1 (d)		CH (γ)	1.45-1.52 (m)	24.8 (d)
	CH ₃ (γ)	0.89 (overlap)	22.6 (q)		CH ₃ (γ)	0.85 (overlap)	22.6 (q)
	CH ₃ (γ)	0.88 (overlap)	22.7 (q)		CH ₃ (γ)	0.89 (overlap)	22.7 (q)

**1****3****Figure 1:** The structures of compounds **1** and **3**, and selected HMBC correlations ($\text{H} \rightarrow \text{C}$) and ^1H , ^1H -COSY (bold line).

Compound **2** was obtained as a white amorphous powder, and established as having a molecular formula of $\text{C}_{32}\text{H}_{49}\text{N}_5\text{O}_5$ by its HR-Q-TOF MS and NMR spectroscopic data (Table 1). The IR spectrum indicated the presence of NH and C=O groups at 3264 and 1627 cm^{-1} , respectively. The ^1H - and ^{13}C -NMR spectral data of **2** (Table 1) were similar to those of **1**, except for the presence of a mono-substituted phenyl group instead of an isopropylidene, indicating that the *N*-methyl-leucine residue in **1** was replaced by *N*-methyl-phenylalanine in **2**. The sequence of these amino acid residues was elucidated on the basis of the same HMBC and NOE correlations as those of **1**, except

**1****3****Figure 2:** The selected NOE correlations for compounds **1** and **3** ($\text{H} \leftrightarrow \text{H}$).

for the HMBC correlations from α -H of *N*-methyl-phenylalanine to C=O of proline, and from α -H of valine to C=O of *N*-methyl-phenylalanine. Thus, the structure of compound **2** was established to be cyclo-(Pro-Me-Phe-Val-Ile-Leu), and named as xylaropeptide B.

Compound **3** was obtained as colorless oil. Its molecular formula was determined to be $\text{C}_{12}\text{H}_{18}\text{O}_4$ according to HR-Q-TOF MS and NMR spectroscopic data. The IR absorption at 3427 cm^{-1} indicated the presence of OH groups. The ^{13}C NMR and DEPT spectra of **3** (Table 2) exhibited 12 signals corresponding to one Me, three CH_2 groups, seven CH (three oxygenated and four olefinic)

groups, as well as one quaternary C-atom (C=O functional group). Inspection of the ^1H - ^1H COSY, HMQC and HMBC data readily revealed a nonenolide core for **3** [15,16]. The HMBC correlations from H-C(2) to C(1) and C(4), from H-C(9) to C(1) and C(11), and from Me(12) to C(10) and C(11), along with ^1H - ^1H COSY correlations H-C(2) \leftrightarrow H-C(3), and H-C(4) \leftrightarrow H-C(5) \leftrightarrow H-C(6) \leftrightarrow H-C(7) \leftrightarrow H₂-C(8) \leftrightarrow H-C(9) \leftrightarrow H₂-C(10), and H₂-C(11) \leftrightarrow H-C(12) established the structure of **3** (Figure 1). The configurations of C(2)/C(3) and C(4)/C(5) double bonds were identified to be *cis* and *trans*, respectively, according to the coupling constants of those corresponding protons.

Table 2: ^1H - and ^{13}C -NMR spectroscopic data of **3** (600/150 MHz, resp., in CDCl_3).

Position	δ_{H} (multiplicity, J in Hz)	δ_{C}
1		168.2 (s)
2	5.94 (d, $J = 10.6$)	126.1 (d)
3	6.66 (d, $J = 10.6$)	139.2 (d)
4	6.26 (d, $J = 15.2$)	130.2 (d)
5	5.48 (dd, $J = 10.1, 15.2$)	133.4 (d)
6	3.90 (t, $J = 9.1$)	78.5 (d)
7	3.53 (t, $J = 7.7$)	76.7 (d)
8	1.87-1.89 (m)	40.0 (t)
9	4.91-4.95 (m)	75.5 (d)
10	1.42-1.54 (m) 1.55-1.62 (m)	38.7 (t)
11	1.35 (m)	18.5 (t)
12	0.93 (t, $J = 7.3$)	13.9 (q)

The relative stereochemistry of **3** was determined by the analysis of the ROESY spectrum. The presence of NOE correlations H-C(7) \leftrightarrow H-C(9) indicated that H-C(7) and H-C(9) were β -oriented, while H-C(6) was in an α -orientation (Figure 2). Thus, the structure of **3** was established as (3*Z*,5*E*)-7,8-dihydroxy-10-propyl-7,8,9,10-tetrahydro-2*H*-oxecin-2-one, and named as xylarolide.

The antibacterial activities of **1** – **3** were tested against bacteria [*Escherichia coli* (CMCC (B) 44103], *Bacillus subtilis* [CMCC (B) 63501], *B. pumilus* [CMCC (B) 63202], and *Staphylococcus aureus* (CMCC (B) 26003)], and a yeast [*Candida albicans* (AS 2.538)] using an Oxford plate assay system. Two replicates were performed for each compound at a concentration of 0.3 mg/mL with a loading volume of 100 μL . Compounds **1**–**3** had no effects on the growth of the tested bacteria and yeast at 30 $\mu\text{g}/\text{plate}$.

Previously, two antibacterial nonenolides, phomolides A and B, were isolated from *Phomopsis* sp.[16]. Phomolides A and B are epoxide analogues of **3**, indicating that the epoxy group between C(6) and C(7) is critical for the antimicrobial activities. The biological activities of cyclopeptides **1** and **2** need to be further explored.

Experimental

General experimental procedures: For column chromatography (CC): silica gel (SiO_2 , 200 - 300 and 80 - 100 mesh; Qingdao Marine Chemical Factory, Qingdao, P. R. China), SiO_2 GF₂₅₄ (Merck), RP-18 (Merck), and

Sephadex LH-20 (Amersham Biosciences) were used. TLC: precoated SiO_2 GF₂₅₄ plates (0.20 - 0.25 mm, Qingdao Marine Chemical Factory). Optical rotations: Perkin-Elmer 341 polarimeter with either CHCl_3 or MeOH as solvent. UV spectra: Amersham Biosciences Ultrospec 1100 pro spectrophotometer; λ_{max} (log ϵ) in nm. IR spectra: Nicolet FT-IR 380 in KBr. HR-Q-TOF-MS: Bruker Daltonics BioTOF-Q mass spectrometers; in m/z (rel. %). ^1H - and ^{13}C -NMR spectra: Bruker DRX-600 spectrometer, at 600 (^1H), and 150 (^{13}C) MHz; in CDCl_3 or $(\text{CD}_3)_2\text{CO}$; δ in ppm rel. to Me_4Si , J in Hz.

Isolation and fermentation of the fungal strain: The fungus was isolated from the fruiting body of *Xylaria* sp., collected at "Gaoligong Mountain National Natural conservation Area", Yunnan Province, P. R. China. Both a traditional morphological assessment and internal transcribed spaces (ITS) sequence analysis were performed to characterize it as *Xylaria* sp. and named 101. Fermentation was performed, and the mycelia of 101 grown on PDA plates were used to inoculate 200 mL PD medium (potato 200 g/L, glucose 20 g/L, pH-neutral) in a 1 L Erlenmeyer flask. The flask was incubated on a rotary shaker for 5 d at 28°C with shaking at 160 rpm. The culture was transferred into 4 flasks (20 L) each containing PD medium (5 L), and cultivated for 2 months at 28°C without agitation.

Extraction and isolation: The culture filtrate (20 L) was concentrated under vacuum at 45°C to a volume of 2 L, and then extracted with an equal volume of AcOEt (3 \times). The combined org. layer, upon evaporation, yielded a brown syrupy extract (4.8 g). This was subjected to MPLC [RP-18 (145g); gradient aq. acetone (0%, 30%, 50%, 70%, and 100% respectively, 2 L each)] to yield 14 fractions (*Fr.a* – *Fr.m*). *Fr.a* (516 mg) was subjected to CC [Sephadex LH-20 (140 g); MeOH] to obtain 2 subfractions *Fr.a.1* and *Fr.a.2*. *Fr.a.1* (286 mg) was purified by CC [Sephadex LH-20 (140 g); MeOH] again, and purified by CC (SiO_2 ; light petroleum (PE)/ Me_2CO 15 : 1) to obtain *Fr.a.1.a* and coriloxin (**4**, 20 mg). *Fr.a.1.a* (6 mg) was further purified by MPLC [RP-18 (30 g); $\text{Me}_2\text{CO}/\text{H}_2\text{O}$ 3:7] to afford **3** (25 mg). *Fr.a.2* (26 mg) was purified by CC (SiO_2 ; PE/ Me_2CO 50: 1) to obtain 2-hydroxy-3-methoxy-5-methyl-*p*-benzoquinone (**5**, 4 mg). *Fr.j* (56 mg) was subjected to MPLC [RP-18 (30g); $\text{Me}_2\text{CO}/\text{H}_2\text{O}$ 1: 1], and further purified by CC (SiO_2 ; chloroform/ Me_2CO 15: 1) to obtain **1** (30 mg). *Fr.k* (160 mg) was subjected to CC [Sephadex LH-20 (140 g); MeOH], and further purified by CC (SiO_2 ; chloroform/ Me_2CO 12: 1), followed by MPLC [RP-18 (30 g); $\text{Me}_2\text{CO}/\text{H}_2\text{O}$ 1:1] to obtain **2** (6 mg).

Xylarotide A (**1**)

Colorless powder.

$[\alpha]_{\text{D}}$: -24.2 (c 0.8, CHCl_3).

IR (KBr): 3287, 2959, 2872, 1640, 1528, 1451, 755 cm^{-1} .

UV/Vis λ_{max} (CHCl_3) nm (log ϵ): 216 (3.09), 225 (2.98), 242 (3.22), 279 (2.85).

^1H NMR and ^{13}C NMR: Table 1.

HR-Q-TOF MS: 550.3921 (calcd. 550.3968 for $\text{C}_{29}\text{H}_{52}\text{N}_5\text{O}_5^+$, $[\text{M} + \text{H}]^+$).

Xylarotide B (2)

Colorless powder.

$[\alpha]_{\text{D}}^{25}$: -3.6 (c 0.18, CHCl_3).

IR (KBr): 3263, 2958, 2925, 2871, 1642, 1529, 698cm^{-1} .

UV/Vis λ_{max} (CHCl_3) nm (log ϵ): 208 (3.44), 212 (3.30), 229 (3.35), 233 (3.34), 239 (3.52), 278 (3.33).

^1H NMR and ^{13}C NMR: Table 1.

HR-Q-TOF MS: 584.3772 (calcd. 584.3812 for $\text{C}_{32}\text{H}_{50}\text{N}_5\text{O}_5^+$, $[\text{M} + \text{H}]^+$).

Xylarolide (3)

Colorless oil.

$[\alpha]_{\text{D}}^{25}$: -18.0 (c 0.54, CHCl_3).

IR (KBr): 3427, 2959, 2934, 2874, 1713, 1387, 1254, 1163, 1091, 1039, 952, 797cm^{-1} .

UV/Vis λ_{max} (CHCl_3) nm (log ϵ): 247 (3.05).

^1H NMR and ^{13}C NMR: Table 2.

HR-Q-TOF MS: 227.1256 (calcd. 227.1283 for $\text{C}_{12}\text{H}_{19}\text{O}_4^+$, $[\text{M} + \text{H}]^+$).

Acknowledgments - This work was financially supported by the National Science Fund for Distinguished Young Scholars to Y.-M. Shen (No. 30325044) and the Key Grant of Chinese Ministry of Education (No. 306010).

References

- [1] Coval SJ, Puar MS, Phife DW, Terracciano JS, Patel M. (1995) SCH57404, an antifungal agent possessing the rare sodaricin skeleton and a tricyclic sugar moiety. *The Journal of Antibiotics*, **48**, 1171-1172.
- [2] Smith CJ, Morin NR, Bills GF, Dombrowski AW, Salituro GM, Smith SK, Zhao A, MacNeil DJ. (2002) Novel sesquiterpenoids from the fermentation of *Xylaria persicaria* are selective ligands for the NPY Y5 receptor. *Journal of Organic Chemistry*, **67**, 5001-5004.
- [3] Deyrup ST, Gloer JB, O'Donnell K, Wicklow DT. (2007) Kolokosides A-D: triterpenoid glycosides from a Hawaiian isolate of *Xylaria* sp. *Journal of Natural Products*, **70**, 378-382.
- [4] Hu ZY, Li YY, Huang YJ, Su WJ, Shen YM. (2008) Three new sesquiterpenoids from *Xylaria* sp. NCY2. *Helvetica Chimica Acta*, **91**, 46-52.
- [5] Li YY, Hu ZY, Lu CH, Shen YM. (2010) Four new terpenoids from *Xylaria* sp. 101. *Helvetica Chimica Acta*, **93**, 796-802.
- [6] Lin YC, Wu XY, Feng S, Jiang GG, Zhou SN, Vrijmoed LLP, Jones EBG. (2001) A novel *N*-cinnamoylcyclopeptide containing an allenic ether from the fungus *Xylaria* sp. (strain # 2508) from the South China Sea. *Tetrahedron Letters*, **42**, 449-451.
- [7] Huang H, She Z, Lin Y, Vrijmoed LL, Lin W. (2007) Cyclic peptides from an endophytic fungus obtained from a mangrove leaf (*Kandelia candel*). *Journal of Natural Products*, **70**, 1696-1699.
- [8] O'Hagana D, Rogers SV, Duffin GR, Edwards RL. (1992) Biosynthesis of the fungal polyketide, cubensis acid from *Xylaria cubensis*. *Tetrahedron Letters*, **33**, 5585-5588.
- [9] Boonphong S, Kittakoop P, Isaka M, Pittayakhajonwut D, Tanticharoen M, Thebtaranonth Y. (2001) Multiplolides A and B, new antifungal 10-membered lactones from *Xylaria multiplex*. *Journal of Natural Products*, **64**, 965-967.
- [10] Jayasuriya H, Herath KB, Ondeyka JG, Polishook JD, Bills GF, Dombrowski AW, Springer MS, Siciliano S, Malkowitz L, Sanchez M, Guan Z, Tiwari S, Stevenson DW, Borris RP, Singh SB. (2004) Isolation and structure of antagonists of chemokine receptor (CCR5). *Journal of Natural Products*, **67**, 1036-1038.
- [11] Healy PC, Hocking A, Tran-Dinh N, Pitt JI, Shivas RG, Mitchell JK, Kotiw M, Davis RA. (2004) Xanthonones from a microfungus of the genus *Xylaria*. *Phytochemistry*, **65**, 2373-2378.
- [12] Davis RA, Pierens GK. (2006) ^1H and ^{13}C NMR assignments for two new xanthonones from the endophytic fungus *Xylaria* sp. FRR 5657. *Magnetic Resonance in Chemistry*, **44**, 966-968.
- [13] Lin Y, Wu X, Feng S, Jiang G, Luo J, Zhou S, Vrijmoed LL, Jones EB, Krohn K, Steingrover K, Zsila F. (2001) Five unique compounds: xyloketal from mangrove fungus *Xylaria* sp. from the South China Sea coast. *Journal of Organic Chemistry*, **66**, 6252-6256.
- [14] Shiono Y, Murayama T, Takahashi K, Okada K, Katohda S, Ikeda M. (2005) Three oxygenated cyclohexenone derivatives produced by an endophytic fungus. *Bioscience, Biotechnology, and Biochemistry*, **69**, 287-292.
- [15] Rivero-Cruz JF, Macias M, Cerda-Garcia-Rojas CM, Mata R. (2003) A new phytotoxic nonenolide from *Phoma herbarum*. *Journal of Natural Products*, **66**, 511-514.
- [16] Du XP, Lu CH; Li YY, Zheng ZH, Su WJ, Shen YM. (2008) Three new antimicrobial metabolites of *Phomopsis* sp. *Journal of Antibiotics*, **61**, 250-253.

A Novel Flavonoid and Furoquinoline Alkaloids from *Vepris glomerata* and their Antioxidant Activity

Joyce J. Kiplimo^a, Md. Shahidul Islam^b and Neil A. Koorbanally^{a*}

^aSchool of Chemistry, University of KwaZulu –Natal, Westville Campus, Private Bag X54001, Durban 4000, South Africa

^bSchool of Biochemistry, Genetics and Microbiology, University of KwaZulu–Natal, Westville Campus, Durban 4000, South Africa

Koorbanally@ukzn.ac.za

Received: June 23rd, 2011; Accepted: October 4th, 2011


The dichloromethane extract of the aerial part of the plant *Vepris glomerata* (Rutaceae) yielded a new flavonoid, which was accorded the trivial name veprisinol (**1**), together with four known furoquinoline alkaloids: haplopine-3,3'-dimethylallyl ether (**2**), anhydroevoxine (**3**), evoxine (**4**) and skimmianine (**5**). The structures of the compounds were established by 1D and 2D NMR spectroscopy, as well as HREIMS. Compounds **1** and **2** have strong antioxidant potential, similar to and in some instances better than ascorbic acid and can be used as beneficial additives to antioxidant supplements.

Keywords: *Vepris glomerata*, veprisinol, furoquinoline alkaloids, antioxidant activity.

The African *Vepris* species have proved to be a good source of furoquinoline and acridone alkaloids that typify the genus as a whole. *V. bilocularis* has been found to have both furoquinoline as well as acridone alkaloids [1,2], while furoquinoline alkaloids alone have been found in *V. ampody* [3], *V. heterophylla* [4], *V. punctata* [5] and *V. stolzii* [6], and acridone alkaloids alone in *V. fitoravina* and *V. macrophylla* [7]. The alkaloids are reported to possess broad spectrum antimicrobial [8], antiradical [9], antioxidant [10], antiplasmodial [11], anticancer [12] and antimutagenic [13] activities. *V. glomerata* is used in African traditional medicine, where its aqueous root extract is used to treat malaria, epilepsy, psychosis and stroke, when mixed with tea [14]. Earlier pharmacological studies on this plant reported antiplasmodial activities of the ethanol extract [15].

Since the species of Rutaceae are often cited as antimalarials or febrifuges in African traditional medicine [14], and the antioxidant activity of alkaloids [10] and flavonoids [16] has previously been demonstrated, all the five compounds isolated were assessed for antioxidant activity using three methods.

Here we report on the isolation and structure elucidation of a new flavonoid, in addition to four known furoquinoline alkaloids: haplopine-3,3'-dimethylallyl ether (**2**), anhydroevoxine (**3**), evoxine (**4**) and skimmianine (**5**) from the dichloromethane extract of *V. glomerata*, together with their antioxidant activities *in vitro*. The structures of the known compounds **2-5** were determined by comparison



of their physical and spectroscopic data with those reported in literature; **2** and **3** [17], **4** [18] and **5** [19]. Only skimmianine was previously reported from the leaves of *V. glomerata* endemic to Ethiopia, in addition to kokusaginine [20]. It is not apparent if the different compounds found in this study are as a result of either geographical or seasonal differences.

Compound **1** was obtained as a yellow solid. Its mass was established to be 388.1573 amu, based on HREIMS data, corresponding to a molecular formula of $C_{21}H_{24}O_7$, which indicates a double bond equivalence of 10, eight being due to the aromatic rings, one being due to the carbonyl group and one to ring C of the flavanone skeleton. The IR spectrum showed a carbonyl stretching band at 1705 cm^{-1} and a hydroxyl absorption band at 3364 cm^{-1} . This compound was identified as a flavanone based on its characteristic ^1H NMR spectral pattern. The characteristic ABX coupling system of H-2 β , H-3 α and H-3 β appeared at δ_{H} 5.29 (1H, dd, $J = 12.84, 2.84\text{ Hz}$, H-2 β), δ_{H} 3.04 (1H, dd, $J = 17.12, 2.84\text{ Hz}$, H-3 α) and δ_{H} 2.75 (1H, dd, $J = 17.12, 12.84\text{ Hz}$, H-3 β). These signals also showed COSY and NOESY correlations with each other.

Another characteristic pattern was that of the trisubstituted aromatic B ring. The proton resonances of this ring occurred as a singlet at δ_H 7.00 (s, H-2') and doublets at δ_H 6.89 and 6.84 (1H each, d, $J = 8.48$ Hz, H-5' and H-6'). The small coupling constant of about 2 Hz for $J_{H2',H6'}$ could not be detected for the H-2' resonance. The 1H NMR spectrum also showed the presence of a methoxy group at δ_H 3.88 (s), its position at C-4' being confirmed by both a 1D NOE and a NOESY correlation with the resonances at δ_H 6.89 and 6.84 (H-5' and H-6'). Five aromatic C-O resonances were seen at δ_C 164.0, 167.2, 162.8, 145.0 and 147.0 attributed to oxygenation at C-5, 7, 9, 3' and 4'.

A pair of doublets at δ_H 6.02 (1H, d, $J = 1.76$ Hz, H-6) and δ_H 6.00 (1H, d, $J = 1.76$ Hz, H-8) were attributed to the *meta* coupled, H-6 and H-8 protons on ring A. These two proton resonances showed NOESY correlations to 2H-1'' at δ_H 4.02, confirming the position of the side chain at C-7. Its corresponding carbon resonance showed HMBC correlations to two multiplets at δ_H 1.87 (overlapping resonances of H-2''a and H-3'') and δ_H 1.61 (H-2''b). The H-2'' resonances were diastereotopic and appeared as two separate resonances. COSY correlations were also observed between H-1'' and H-2''a and H-2''b and between H-2''b and H-3''. The H-3'' methine proton was coupled to the methyl proton resonance at δ_H 0.95 (d, $J = 6.52$ Hz) attributed to 3H-5'' and the methylene proton at δ_H 3.50 (2H-4'') in the COSY spectrum. These correlations formed a side chain which was attached to ring A by an ether linkage at C-7. Compound **1** was thus identified as 4H-1-benzopyran-4-one, 2, 3-dihydro-5-hydroxy-2-(4'-methoxy-3'-hydroxybenzyl)-7-O-(2-methyl butanol) ether, and given the trivial name veprisinol.

The results of the reducing potential (transformation of Fe^{3+} - Fe^{2+}) of the standard (ascorbic acid) and compounds **1-5** are shown in Figure 1. The activity of haplopine-3,3'-dimethylallyl ether, **2** and veprisinol (**1**) was significantly higher than the activity of the other three alkaloids at all concentrations. However, the reducing power of compound **1** was significantly lower than that of compound **2**. The reducing power of the compounds and standard followed the order: ascorbic acid > **2** > **1** > **3** > **4** > **5**.

The DPPH radical scavenging assay results are shown in Fig. 2. The results revealed that the scavenging activity of the standard ascorbic acid was significantly higher than all other compounds tested. At concentrations of $62.5 \mu g mL^{-1}$ and above, the activity decreased in the order ascorbic acid > **1** > **2** > **4** > **3**, whereas at the lower concentrations, 31.25 and $15.625 \mu g mL^{-1}$, evoxine (**4**) had the highest percentage antioxidant activity of 41%. The activity of compounds **1** and **2** was increased with their concentration and significantly higher than other compounds, particularly at higher concentrations (Figure 2).




Figure 1: Free radical reducing potential of compounds **1-5** and standard ascorbic acid as evaluated by the spectrophotometric detection of the Fe^{3+} - Fe^{2+} transformation (FRAP method).




Figure 2: Antioxidant activity of compounds **1-4** and ascorbic acid standard, as measured by the DPPH method.

The hydroxyl radical scavenging activities in the deoxyribose assay are shown in Figure 3. The results revealed that compound **1** possessed significantly higher activity than all the other compounds tested, including the standard, ascorbic acid, at most concentrations. Compounds **1**, **2** and **4** had hydroxyl radical scavenging activity comparable with and in the case of **1** and **2**, better than that of ascorbic acid. Skimmianine (**5**) was not tested in either the DPPH or deoxyribose assays due to insufficient amount.

The three assays revealed that compounds **1** and **2** are good antioxidant compounds, while compound **4** shows high activity at a lower concentration in the DPPH assay. Flavonoids are known to be potent antioxidants and their activity is dependent on their molecular structure. The activity of **1** could be attributed to the hydroxyl (OH) groups in the molecule, which donate hydrogen to reduce the DPPH radical to DPPH-H. The alkaloids **2-5** have the same basic skeleton, the only difference being in their side chain. The reductive ability of **2** may be attributed to the double bond of the isoprenyl unit, rich in delocalized pi-electrons, which are easily donated during reduction of Fe^{3+} to Fe^{2+} . Sang *et al.* also reported that the double bond of the isoprenyl group was responsible for the antioxidant activity of garcinol [21]. The antioxidant activity of **2** in the DPPH assay, like **1**, could also be attributed to the hydroxyl groups in the molecule.




Figure 3: Hydroxyl radical scavenging activity of compounds 1-4 and standard ascorbic acid as measured by the deoxyribose method.

In conclusion, five compounds were isolated (a flavonoid and four alkaloids) from the aerial parts of *V. glomerata*. Verification of their antioxidant activities, as well as comparison with known antioxidants, will provide herbalists and traditional healers with scientific evidence for the use of the aerial parts of this plant as natural antioxidants.

Experimental

General experiment procedures: The melting points were recorded on an Ernst Leitz Wetzlar micro-hot stage melting point apparatus and are uncorrected. UV spectra were obtained on a Varian Cary UV-VIS Spectrophotometer in chloroform. IR spectra were recorded on a Perkin-Elmer Universal ATR Spectrometer. The 1D and 2D NMR spectra were recorded using a Bruker Avance^{III} 400 MHz NMR spectrometer. All the spectra were recorded at room temperature using deuterated chloroform (CDCl₃) as solvent. The HREIMS was measured on a Bruker Micro TOF-QII instrument. Specific rotations were measured at room temperature in chloroform on a PerkinElmerTM, Model 341 Polarimeter with a 10 mm flow tube. The separation, isolation and purification of compounds were carried out by gravity CC and monitored by TLC. Merck silica gel 60 (0.040-0.063 mm) was used for CC. Merck 20 × 20 cm silica gel 60 F₂₅₄ aluminum sheets were used for TLC. TLC plates were analyzed under UV light (254 and 366 nm) before being sprayed with anisaldehyde: concentrated sulfuric acid: methanol [1:2:97] spray reagent and then heated.

Plant material: *Vepris glomerata* was collected from the Rift Valley province of Kenya and identified by Dr S. T. Kariuki from the Department of Botany, Egerton University, Kenya. A voucher specimen (Kiplimo 01) was deposited at the University of KwaZulu-Natal Ward Herbarium, Westville Campus, Durban, South Africa.

Extraction and isolation: The air-dried aerial parts (980 g) of *V. glomerata* were sequentially extracted with *n*-hexane, followed by dichloromethane in a Soxhlet apparatus for 48 h, yielding crude extracts of 46 and 32 g, respectively. The oily residue of the dichloromethane extract obtained after evaporation under vacuum, was separated by CC on silica

gel with *n*-hexane and then increasing the concentration of ethyl acetate from 10 to 80% in *n*-hexane, to give 10 fractions (fr.); fr. 8-16 (1.27 g), fr. 17-19 (0.5 g), fr. 20-26 (2.36 g), fr. 27-32 (2.35 g), fr. 33-39 (1 g), fr. 40-43 (2.1 g), fr. 44-49 (0.5 g), fr. 52-56 (3.9 g), fr. 57-62 (1.75 g) and fr. 63-67 (5.1 g).

Fraction 52-56 was separated by CC with *n*-hexane/EtOAc (7:3) as the solvent to afford sub-fractions A-C. Sub-fraction A was further purified using 100% dichloromethane to afford compound 2, a green solid (51 mg). Sub-fraction B yielded compound 3, a brownish solid (43 mg), which needed no further purification. Sub-fraction C was crystallized in methanol to afford 4 (62 mg). Fraction 44-49 was purified using 100% dichloromethane to afford 5 (60 mg). Fraction 63-67 was separated with *n*-hexane/EtOAc (4:1) to yield 4 sub-fractions A-D. Sub-fraction B was crystallized in methanol to afford yellow crystals of compound 1 (18 mg).

Veprisinol (1)

4H-1-Benzopyran-4-one, 2, 3-dihydro-5-hydroxy-2-(4'-methoxy-3'-hydroxybenzyl)-7-O-(2-methyl butanol) ether

Yellow solid.

M.p: 78-80°C.

[α]_D²⁰: +55.30 (c 0.056, CHCl₃).

IR: 3364 (O-H), 2928, 1705 (C=O), 1636, 1512, 1162 cm⁻¹. UV λ_{max} (CHCl₃) nm (log ε): 337 (4.45), 285 (5.13), 239 (5.44).

¹H NMR (400 MHz, CDCl₃): 11.97 (H, s, OH), 7.00 (H, s, H-2'), 6.89 (H, d, *J* = 8.28 Hz, H-5'), 6.84 (H, d, *J* = 8.28 Hz, H-6'), 6.02 (H, d, *J* = 1.76 Hz, H-6), 6.00 (H, d, *J* = 1.76 Hz, H-8), 5.29 (H, dd, *J* = 12.84, 2.84 Hz, H-2β), 4.02 (2H, dd, *J* = 12.88, 6.24 Hz, 2H-1'), 3.88 (3H, s, OCH₃), 3.50 (2H, d, *J* = 5.68 Hz, 2H-4'), 3.04 (H, dd, *J* = 17.12, 12.84 Hz, H-3α), 2.75 (H, dd, *J* = 17.12, 2.84 Hz, H-3β), 1.87 (2H, m, H-2''a and H-3''), 1.61 (H, m, H-2''b), 0.95 (3H, d, *J* = 6.52 Hz, H-5'').

¹³C NMR: 195.97 (C, C-4), 167.29 (C, C-7), 164.05 (C, C-5), 162.85 (C, C-9), 147.02 (C, C-4'), 145.93 (C, C-3'), 131.52 (C, C-1'), 118.15 (CH, C-5'), 112.71 (CH, C-2'), 110.71 (CH, C-6'), 103.11 (C, C-10), 95.54 (CH, C-6), 94.60 (CH, C-8), 78.92 (CH, C-2), 67.86 (CH₂, C-4''), 66.70 (CH₂, C-1''), 56.06 (OCH₃), 43.15 (CH₂, C-3), 32.94 (CH, C-3''), 32.37 (CH₂, C-2''), 16.60 (CH₃, C-5'').

HREIMS *m/z* 388.1573 [M]⁺ (calcd. for C₂₁H₂₄O₇, 388.1522)

Antioxidant activity: The total reducing power was determined according to the method described previously [22]. The free radical scavenging activity (antioxidant capacity) of the plant phytochemicals on the stable radical 2, 2-diphenyl-β-picrylhydrazyl (DPPH) was evaluated by the method established by Shirwaikar *et al.* [23], and the deoxyribose assay for hydroxyl radical scavenging activity was performed as described previously by Chung *et al.* [24].

Statistical analysis: The data in Figures 1-3 are presented as mean \pm SD of triplicates. ^{a-d}Values with different superscript letters for a given concentration are significantly different from each of the other compounds. The data were statistically analyzed using a statistical software program SPSS (SPSS for Windows, version 18, SPSS Science, Chicago, IL, USA). One-way analysis of

variance (ANOVA) followed by Tukey's multiple range post-hoc test was employed to find the differences. The data were considered significantly different at $p < 0.05$.

Acknowledgments - The authors wish to acknowledge the financial support received from the Organization for Women in Science for the Developing World (OWSDW).

References

- [1] Govindachari TR, Sundararajan VN. (1961) Alkaloids of *Vepris bilocularis*. *Journal of Scientific and Industrial Research*, **20**, 298-299.
- [2] Ganguly AK, Govindachari TR, Manmade A, Mohamed PA. (1966) Alkaloids of *Vepris bilocularis*. *Indian Journal of Chemistry*, **4**, 334-335.
- [3] Kan-Fan B, Das C, Boiteau P, Potier P. (1970) Alcaloïdes de *Vepris ampody* (Rutacées). *Phytochemistry*, **9**, 1283-1291.
- [4] Gomes E, Dellamonica G, Gleye J, Moulis C, Chopin MJ, Stanislas E. (1983) Phenolic compounds from *Vepris heterophylla*. *Phytochemistry*, **22**, 2628-2629.
- [5] Chaturvedula VSP, Schilling JK, Miller JS, Andransiferana R, Rasamison VE, Kingston DGI. (2003) New cytotoxic alkaloids from the wood of *Vepris punctata* from the Madagascar rainforest. *Journal of Natural Products*, **66**, 532-534.
- [6] Khalid SA, Waterman PG. (1982) Chemosystematics in the Rutaceae 15. Furoquinoline and pyrano-2-quinolone alkaloids of *Vepris stolzii*. *Journal of Natural Products*, **45**, 343-346.
- [7] Koffi Y, Gleye J, Moulis C, Stanislas E. (1987) Acridones from *Vepris fitoravina* and *Vepris macrophylla*. *Planta Medica*, **53**, 570-571.
- [8] Karou D, Savadogo A, Canini A, Yameogo S, Montesano C, Simporé J, Colizzi V, Traore AS. (2006) Antibacterial activity of alkaloids from *Sida acuta*. *African Journal of Biotechnology*, **5**, 195-200.
- [9] Račková L, Májeková M, Košťálová D, Štefek M. (2004) Antiradical and antioxidant activities of alkaloids isolated from *Mahonia aquifolium*. Structural aspects. *Bioorganic & Medicinal Chemistry*, **12**, 4709-4715.
- [10] Chung HS, Woo SO. (2001) A quinolone alkaloid with antioxidant activity from the aleurone layer of anthocyanin-pigmented rice. *Journal of Natural Products*, **64**, 1579-1580.
- [11] Waffo KFA, Coombes PH, Crouch NR, Mulholland DA, Amin SMM, Smith PJ. (2007) Acridone and furoquinoline alkaloids from *Teclea gerrardii* (Rutaceae: Toddalioidae) of Southern Africa. *Phytochemistry*, **68**, 663-667.
- [12] Iwasa K, Moriyasu M, Yamori T, Turuo T, Lee DU, Eigrebe W. (2001) *In vitro* cytotoxicity of the protoberberine-type alkaloids. *Journal of Natural Products*, **64**, 896-898.
- [13] Fukuda K, Hibiya Y, Mutoh M, Koshiji M, Akao S, Fujiwara H. (1999) Inhibition by berberine of cyclooxygenase-2 transcriptional activity in human colon cancer cells. *Journal of Ethnopharmacology*, **66**, 227-233.
- [14] Moshi JM, Mbwambo ZH, Nondo RSO, Masimba PJ, Kamuhabwa A, Kapingu MC, Thomas P, Richard M. (2006) Evaluation of ethnomedical claims and brine shrimp toxicity of some plants used in Tanzania as Traditional Medicines. *African Journal of Traditional Complementary and Alternative Medicine*, **3**, 48-58.
- [15] Innocent E, Moshi M, Mbwambo P, Mbwambo Z, Kapingu M, Kamuhabwa A. (2009) Screening of traditionally used plants for *in vivo* antimalarial activity in mice. *African Journal of Traditional Complementary and Alternative Medicine*, **6**, 163-167.
- [16] Dimitrios B. (2006) Sources of natural phenolic antioxidants. *Trends in Food Science Technology*, **17**, 505-512.
- [17] Al-Rehaily AJ, Ahamad MS, Muhammad IM, Al-Thukair AA, Perzanowski HP. (2003) Furoquinoline alkaloids from *Teclea nobilis*. *Phytochemistry*, **64**, 1405-1411.
- [18] Dreyer DL. (1970) Chemotaxonomy of the Rutaceae.VII. Alkaloids in *Evodia zanthoxyloides* F. Muell. *Journal of Organic Chemistry*, **35**, 2420-2422.
- [19] Latip J, Ali MNH, Yamin BM. (2005) 4, 6, 7-Trimethoxyfuro[2, 3-b]quinoline-water (2/3). *Acta Crystallographica*. **E61**, 02230-02232.
- [20] Dagne E, Yenesew A, Waterman PG, Gray AI. (1988) The chemical systematics of the Rutaceae subfamily Toddalioidae, in Africa. *Biochemical Systematics and Ecology*, **16**, 179-188.
- [21] Sang S, Liao CH, Pan MH, Rosen RT, Lin-Shiau SY, Lin JK, Ho CT. (2002) Chemical studies on antioxidant mechanism of garcinol: analysis of radical reaction products of garcinol with peroxy radicals and their antitumor activities. *Tetrahedron*, **58**, 10095-10102.
- [22] Ferreira ICFR, Baptista M, Vilas-Baos, Barros L. (2007) Free radical scavenging capacity and reducing capacity of wild edible mushrooms from northeast Portugal: individual cap and stipe activity. *Food Chemistry*, **100**, 1511-1516.
- [23] Shirwakar A, Shirwakar AR, Rajendran K, Punitha IRS. (2006) *In vitro* antioxidant studies on the benzyltetraisoquinoline alkaloid berberine. *Biological and Pharmaceutical Bulletin*, **29**, 1906-1910.
- [24] Chung SK, Osawa T, Kawakishi S. (1997) Hydroxyl radical-scavenging effects of species and scavengers from brown mustard (*Brassica nigra*). *Bioscience, Biotechnology and Biochemistry*, **61**, 118-123.

Flavonoid Constituents and Free Radical Scavenging Activity of *Alchemilla mollis*

Antoaneta Trendafilova^{a,*}, Milka Todorova^a, Milena Nikolova^b, Anna Gavrilova^b and Antonina Vitkova^b

^aInstitute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, Acad. G. Bonchev Str., bl. 9, 1113 Sofia, Bulgaria

^bInstitute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, 2 Gagarin Str., 1113 Sofia, Bulgaria

trendaf@orgchm.bas.bg

Received: June 29th, 2011; Accepted: October 24th, 2011

Antioxidant capacity of the methanolic extract of *Alchemilla mollis* was measured by its ability to scavenge the DPPH radical. The EtOAc fraction obtained after partition of the total extract was found to be the most active radical scavenger (IC₅₀ 9.8 ± 1.8 µg/mL) and was subjected to fractionation by Sephadex LH-20 CC. Further purification by RP-18 CC led to the isolation of eight flavonoid glycosides: *cis*- and *trans*-tiliroside (**1** and **2**), rhodiogin (**3**), hyperoside (**4**), isoquercitrin (**5**), miquelianin (**6**), sinocrassoside D₂ (**7**), and gossypetin-3-*O*-β-D-galactopyranosyl-7-*O*-α-L-rhamnopyranoside (**8**). It was found that **8** is a new compound and its antioxidant activity is also reported. Identification of the isolated compounds was carried out by spectroscopic and spectrometric analysis (1D and 2D NMR, UV and MS).

Keywords: *Alchemilla mollis*, Rosaceae, flavonoid glycosides, antioxidant activity.

Alchemilla mollis (Buser) Rothm. (Rosaceae) is a herbaceous perennial plant that grows in mountainous regions of the Balkan Peninsula (Romania, Greece and Bulgaria) [1]. The species, with its only known locality in the National Park "Central Balkan", is critically endangered in Bulgaria according to the IUCN criteria and was included in the Red Book of Bulgaria, the Red List of Bulgarian vascular plants, the Biological Diversity Act, and the Medicinal Plants Act [2a-2c]. Because of its very limited occurrence, its cultivation has been started recently [3,4].

A. mollis forms part of the commercial drug "Herba Alchemillae", an aggregate of species collectively referred to as Lady's mantle. The drug possesses astringent, diuretic and antispasmodic properties, and is commonly used in traditional medicine as a cure for excessive menstruation and wounds [5,6]. Different studies showed that the phenolic compounds (for example, tannins, and flavonoids) present in the plant are responsible for the pharmacological activity of Lady's mantle [7a-7d]. Surprisingly, the literature data concerning the chemical composition of *A. mollis* have proved to be quite limited, with only a few reports on total flavonoid and tannin contents of the plant [8a-8c]. Consequently, the aim of the present work was to study the flavonoid constituents of this species and their antioxidant capacity.

The DPPH scavenging assay is widely used for preliminary evaluation of the antioxidant potential of extracts and

Table 1: DPPH radical scavenging activity of *A. mollis* total methanolic extract, fractions and individual compounds.

	IC ₅₀ (µg/mL)
total MeOH extract	31.7 ± 4.9
PE fraction	>200
CHCl ₃ fraction	>200
EtOAc fraction	9.8 ± 1.8
H ₂ O residue	42.5 ± 4.3
Quercetin (Reference)	3.2 ± 0.4
Hyperoside (4)	5.1 ± 0.4
Compound 8	4.9 ± 0.5

individual compounds [10,11]. In this work, the MeOH extract of *A. mollis* and its fractions obtained after re-extraction with light petroleum (PE), chloroform (CHCl₃), and ethyl acetate (EtOAc), as well as the remaining H₂O residue were studied for their general antioxidant effect, as indicated by their potential to scavenge the stable DPPH radical.

The total MeOH extract exhibited a significant dose dependent inhibition of DPPH activity with 50% inhibition (IC₅₀) at a concentration of 31.7 ± 4.9 µg/mL. As shown in Table 1, the scavenging activities of the fractions on the DPPH radical increased in the order of PE < CHCl₃ < H₂O < EtOAc. Although the DPPH free radical scavenging ability of the EtOAc fraction (IC₅₀ 9.8 ± 1.8 µg/mL) was less than that of quercetin (IC₅₀ 3.2 ± 0.4 µg/mL) it was evident that the EtOAc-soluble fraction had hydrogen donating ability and could serve as either a free radical inhibitor or scavenger, possibly acting as a primary antioxidant.





Figure 1: Structures of flavonoid glycosides 1-8.

The most active EtOAc-soluble fraction was subjected to Sephadex LH-20 column chromatography. Further purification by RP-18 silica gel column chromatography afforded eight flavonoid glycosides. The structures (Figure 1) of the known *cis*- and *trans*-tiliroside (**1** and **2**) [11], rhodiolgin (**3**) [12], hyperoside (**4**) [13], isoquercitrin (**5**) [13], miquelianin (**6**) [13] and sinocrassoside D₂ (**7**) [14] were determined by comparison of their spectral data (UV, ¹H NMR, ¹³C NMR and MS) with those published previously.

Figure 2: Significant HMBC and NOE correlations in **8**.Table 2: ¹H and ¹³C NMR spectroscopic data of compound **8** (CD₃OD).

Position	δ _H J (Hz)	δ _C	Position	δ _H J (Hz)	δ _C
2		159.4	1''	5.20 d (7.2)	105.4
3		135.3	2''	3.85 m*	73.2
4		180.1	3''	3.65 m*	75.1
5		154.3	4''	3.85 m*	70.1
6	6.67 s	99.9	5''	3.50 m*	77.2
7		152.2	6''	3.50 m*	62.0
8		128.5			
9		146.2			
10		107.1			
1'		123.0	1'''	5.54 brs	100.9
2'	7.94 d (2.2)	116.1	2'''	4.14 brs	71.7
3'		145.8	3'''	3.99 dd (2.8, 9.4)	72.1
4'		150.2	4'''	3.65 m*	73.8
5'	6.87 d (8.5)	118.1	5'''	3.55 m*	71.3
6'	7.72 dd (2.2, 8.5)	123.5	6'''	1.26 d (6.2)	18.1

*overlapped signals

Compound **8**, isolated as yellowish amorphous powder, displayed spectral properties very similar to those of sinocrassoside D₂ (**7**) [14]. The negative HR-ESI-MS of **8** showed a [M-H]⁺ peak at *m/z* 625.14035, which corresponded to a molecular formula C₂₇H₂₉O₇. The ¹H NMR spectrum (Table 2) contained signals assignable to a gossypetin moiety [δ_H 6.67 (1H, s, H-6), 6.87 (1H, d, *J* = 8.5 Hz, H-5'), 7.72 (1H, dd, *J* = 2.2, 8.5 Hz, H-6'), 7.94 (1H, d, *J* = 2.2 Hz, H-2')], an α-L-rhamnopyranosyl unit [δ_H 1.26 (3H, d, *J* = 6.2 Hz) and δ_H 5.54 (1H, brs)] and the doublet at δ_H 5.20 (*J* = 7.2 Hz) characteristic of an anomeric proton of a sugar moiety with β-configuration. This was identified as D-galactose after acid hydrolysis of **8** and confirmed by co-TLC with an authentic sample. The full NMR assignments and connectivities of **8** were determined by HMBC and NOESY data analysis (Figure 2). Thus, the observed HMBC long-range correlations between anomeric proton H-1'' (δ_H 5.20) and C-3 δ_C (135.3), as well as between the second anomeric proton H-1''' (δ_H 5.54) and C-7 (δ_C 152.2) supported the location of sugar moieties at C-3 and C-7. Furthermore, NOE interactions of the aromatic singlet at δ_H 6.67 (H-6) with H-1''' and H-6''' confirmed the position of the

α-L-rhamnopyranosyl unit at C-7. Thus, the new compound (**8**) was identified as gossypetin-3-β-D-galactopyranosyl-7-α-L-rhamnopyranoside.

Finally, the antioxidant activity of the new compound **8** was evaluated by the DPPH free radical scavenging assay (Table 1) and compared with that of hyperoside (**4**) and quercetin. As can be seen, **8** exhibited antioxidant activity against the DPPH radical with an IC₅₀ value (4.9 ± 0.5 μg/mL) similar to that of hyperoside (IC₅₀ 5.1 ± 0.4 μg/mL) and lower than that of quercetin (IC₅₀ 3.2 ± 0.4 μg/mL). This significant radical scavenging activity of **8** could be explained by the presence of *ortho*-dihydroxyl groups at C-3' and C-4' in ring B of the flavonoid structure [15].

The results described above showed that the EtOAc fraction obtained from the total methanolic extract of *A. mollis* was characterized by the presence of three different types of flavonoid glycosides – kaempferol (**1** and **2**), quercetin (**4**, **5** and **6**) and gossypetin (**3**, **7** and **8**). It is noteworthy, that kaempferol and quercetin glycosides have been previously found in *A. speciosa* [16], *A. xanthochlora* [17,18] and *A. vulgaris* [19], while gossypetin glycosides were detected for the first time in the genus *Alchemilla*. These differences in the chemical composition of the investigated species could be of chemotaxonomic importance.

Experimental

General: UV spectra were taken on a Helios Gamma spectrophotometer (Thermo Electron Corporation). ¹H and ¹³C NMR spectra were recorded on a Bruker Avance II+ 600 NMR spectrometer with an operating frequency of 600.13 MHz (¹H) and 150.903 MHz (¹³C), using TMS as internal standard and CD₃OD as solvent. Chemical shifts (δ) are expressed in ppm and coupling constants (*J*) in Hertz. 1D and 2D NMR spectra were recorded using the standard Bruker pulse sequence. The HR-ESI-TOF-MS

was measured using a 6210 Time-of-flight LC/MS system (Agilent technologies) coupled with a HPLC instrument 1200 Series (Agilent Technologies). TLC analyses were carried out on silica gel 60 F₂₅₄ precoated plates (Merck). The compounds were visualized by spraying with either H₂SO₄ or NP-reagent (Fluka). For open column chromatography, Sephadex LH-20 (Fluka) was used as stationary phase. A LiChroprep RP-18 column (Merck) was used for medium pressure liquid chromatography (MPLC).

Plant material: *A. mollis* plants from a native Bulgarian population (Stara planina mt. -1150 m a.s.l.), cultivated in the experimental field of the Institute of Biodiversity and Ecosystem Research (Sofia – 570 m a.s.l.), were used in the experiments. The aerial flowering parts of the plants were collected within phenophase – full blossoming, air-dried and kept in the dark. The voucher specimen (SOM 159980) was deposited in the Herbarium of the Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, Sofia.

Extraction and isolation: Powdered plant material (70 g) was extracted with MeOH (3 x 1 L) at room temperature in an ultrasonic bath for 30 min each time. After filtration, the solvent from the combined extracts was evaporated under vacuum to give the total methanolic extract (MeOH, 9.10 g). This was further dissolved in distilled water (200 mL) and partitioned with light petroleum (PE, 3 x 80 mL), chloroform (CHCl₃, 3 x 80 mL) and ethyl acetate (EtOAc, 3 x 80 mL) to yield PE (0.57 g), CHCl₃ (0.28 g) and EtOAc (0.44 g) fractions. The remaining aqueous phase was evaporated to dryness (7.51 g).

The EtOAc extract was dissolved in MeOH (15 mL) and filtered through celite in order to remove insoluble parts. The clear methanolic solution was concentrated to 5 mL and applied to a Sephadex LH-20 column (equilibrated with MeOH) to give 4 main fractions [TLC: Silica gel, EtOAc/MeOH/H₂O, 5:0.8:0.6, and EtOAc/HCOOH/CH₃COOH/H₂O, 100:11:11:26]. The flavonoid containing fraction (65 mg) was further subjected to MPLC (Li Chroprep RP-18) and eluted with increasing concentrations of MeOH in H₂O (20 to 70%). Repeated MPLC (Li Chroprep RP-18, MeOH/H₂O, 50:50) of selected fractions yielded mixture of *cis*- and *trans*-tiliroside (**1** and **2**, 4.8 mg), rhodiogin (**3**, 0.8 mg), hyperoside (**4**, 10.8 mg), isoquercitrin (**5**, 2.2 mg), miquelianin (**6**, 4.9 mg), sinocrassoside D₂ (**7**, 1.2 mg) and gossypetin-3-*O*-β-*D*-galactopyranosyl-7-*O*-α-*L*-rhamnopyranoside (**8**, 5.9 mg). Known compounds were identified by comparison of their

spectral data with those reported in the literature and by comparison with authentic samples. It should be noted that the isolated amounts do not represent the real concentration of the compounds in the plant.

Gossypetin 3-*O*-β-*D*-galactopyranoside-7-*O*-α-*L*-rhamnopyranoside (8**)**

Yellowish amorphous powder.

UV λ_{max} nm (MeOH): 260, 275, 303sh, 352; + NaOMe 267, 339sh, 398; + AlCl₃ 277, 315sh, 452; + AlCl₃ / HCl 249sh, 282, 312sh, 365, 429; + NaOAc 270, 336, 390.

¹H NMR (CD₃OD): Table 2.

¹³C NMR (CD₃OD): Table 2.

Negative HR-ESI-TOF-MS: *m/z* 625.14035 (calcd. for C₂₇H₂₉O₇ 625.14102) [M-H]⁺

Acid hydrolysis: Flavonoid **8** (2 mg) was dissolved in MeOH (5 mL) and refluxed with 8% HCl (2 mL) for 2 h. The reaction mixture was evaporated to dryness, dissolved in H₂O (2 mL) and neutralized with NaOH. The neutralized product was analyzed by TLC (silica gel, *n*-PrOH-EtOAc-H₂O, 7:2:1) in the presence of authentic samples.

Free radical scavenging activity of the DPPH radical:

The DPPH radical scavenging method was used for determination of the antioxidant capacity of the extracts and individual compounds [9,10]. Different concentrations of the extracts (10, 20, 50, 100 and 200 µg/mL in MeOH) and individual compounds (5, 10, 20 and 50 µg/mL in MeOH) were added in equal volume (2.5 mL) to the MeOH solution of DPPH[•] (0.3 mM, 1 mL). After 30 min at room temperature, the absorption values were measured at 517 nm on a spectrophotometer (Jenway 6320D) and converted into the percentage antioxidant activity using the following equation: DPPH[•] anti-radical scavenging capacity (%) = [1-(A_{sample} - A_{blank})/A_{control}]*x*100. MeOH (1.0 mL). The solution plus plant extract solution (2.5 mL) was used as a blank, while DPPH[•] solution plus MeOH was used as a control. The extracts were measured in triplicate on 2 different days. The results are presented as a mean ± SD. The IC₅₀ values were calculated by Software Prizm 3.00.

Acknowledgements - The authors are thankful to Dr Milka Jadranin (University of Belgrade, Faculty of Chemistry, Belgrade) for preparing the HRMS. This work was supported by the National Science Fund, Ministry of Education, Youth and Science, Bulgaria, project DTK 02/38 (2009).

References

- [1] Kurtto A, Fröhner SE, Lampinen R. (Eds) (2007) *Atlas Florae Europaeae. Distribution of Vascular Plants in Europe*. Vol. 14. *Rosaceae (Alchemilla and Aphanes)*. The Committee for Mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki. pp.200.

- [2] (a) Biological Diversity Act. State Gazette, №77/02.08. 2002. (in Bulgarian); (b) Vitkova A. (2009) *Alchemilla mollis* (Buser) Rothm. In *Red list of Bulgarian vascular plants*. Petrova A, Vladimirov V. (Eds). *Phytologia Balcanica*, **15**, 63-94; (c) Vitkova A. (2011) *Alchemilla mollis* (Buser) Rothm. In *Red Book of R Bulgaria*. Vol. 1. Plants, Peev D. (Ed.) In press.
- [3] Vitkova A. (1997) Contribution to biological investigations of *Alchemilla mollis* (Buser) Rothm. *Phytologia Balcanica*, **3**, 57-61.
- [4] Vitkova A, Gavrilova A, Tashev A. (2011). *Alchemilla mollis* (Rosaceae) – critically endangered species in Bulgaria. *Phytologia Balcanica*, in press.
- [5] Ivanov I, Landgev I, Neshev G. (1977) *Bilkite v Bulgaria i izpolzvaneto im* (Medicinal plants in Bulgaria and their applications), Sofia. 256-257.
- [6] Bisset NG. (1994) Lady's Mantle In *Herbal Drugs and Phytopharmaceuticals*. Wichtl M and NG Bisset (Eds), Medpharm Scientific Publishers, Stuttgart. 52-54.
- [7] (a) Lamaison JL, Carnat A, Petitjean-Freytet C. (1990) Tannin content and inhibiting activity of elastase in Rosaceae. *Annales Pharmaceutiques Francaises*, **48**, 335-340; (b) Filipek J. (1992) Effect of *Alchemilla xanthochlora* water extracts on lipid peroxidation and superoxide anion scavenging activity. *Pharmazie*, **47**, 717-718; (c) Jonadet M, Meunier MT, Villie F, Bastide JP, Lamaison JL. (1986) Flavonoids extracted from *Ribes nigrum* L. and *Alchemilla vulgaris* L.: 1. *In vitro* inhibitory activities on elastase, trypsin and chymotrypsin. 2. Angioprotective activities compared *in vivo*. *Journal of Pharmacology*, **17**, 21-27; (d) Schimmer O, Lindenbaum M. (1995) Tannins with antimutagenic properties in the herb of *Alchemilla* species and *Potentilla anserine*. *Planta Medica*, **61**, 141-145.
- [8] (a) Tuka L, Tamas M. (1977) Determination of flavonoids in *Alchemilla mollis* (Buser) Rothm and *Alchemilla* L. (Lady's mantle), *Farmacia*, **25**, 247-252; (b) Vitkova A. (1996) A comparative assessment and dynamics of accumulation of flavonoids and tannins in species of *Alchemilla* L. (Rosaceae). *Fitology*, **48**, 11-18; (c) Ayaz F A, Hayirlioglu-Ayaz S. (2001) Total phenols and condensed tannins in the leaves of some *Alchemilla* species. *Biologia*, **6**, 449-453.
- [9] Cimpoiu C. (2006) Analysis of some natural antioxidants by thin-layer chromatography and high performance thin-layer chromatography. *Journal of Liquid Chromatography & Related Technologies*, **29**, 1125-1142.
- [10] Matkowski A, Piotrowska M. (2006) Antioxidant and free radical scavenging activities of some medicinal plants from the Lamiaceae. *Fitoterapia*, **77**, 346-353.
- [11] Matlawska I, Sikorska M, Bylka W. (1999) Flavonoid compounds in *Lavatera thuringiaca* L. (Malvaceae) flowers. *Acta Poloniae Pharmaceutica*, **56**, 453-445.
- [12] Zapesochynaya GG, Kurkin VA, Shchavinskii AN. (1985) Flavonoids of the epigeal part of *Rhodiola rosea*. II. Structures of new glycosides of herbacetin and of gossypetin. *Chemistry of Natural Compounds*, **21**, 464-473.
- [13] Tatsis EC, Boeren S, Exarchou V, Troganis AN, Vervoort J, Gerathanassis IP. (2007) Identification of the major constituents of *Hypericum perforatum* by LC/SPE/NMR and LC/MS. *Phytochemistry*, **68**, 383-393.
- [14] Morikawa T, Xie H, Wang T, Matsuda H, Yoshikawa M. (2008) Bioactive constituents from Chinese natural medicines. XXXII.1) Aminopeptidase N and aldose reductase inhibitors from *Sinocrassula indica*: Structures of sinocrassosides B₄, B₅, C₁, and D₁—D₃. *Chemical & Pharmaceutical Bulletin*, **56**, 1438-1444.
- [15] Zheng CD, Li G, Li HQ, Xu XJ, Gao JM, Zhang AL. (2010) DPPH-scavenging activities and structure-activity relationships of phenolic compounds. *Natural Product Communications*, **5**, 1759-1765.
- [16] Felser C, Schimmer O. (1999) Flavonoid glycosides from *Alchemilla speciosa*, *Planta Medica*, **65**, 668-670.
- [17] Lamaison JL, Carnat A, Petitjean-Freytet C, Carnat AP. (1991) Quercetin-3-glucuronide, the main flavonoid of Lady's Mantle, *Alchemilla xanthochlora* Rothm. (Rosaceae). *Annales Pharmaceutiques Francaises*, **49**, 186-189.
- [18] Fraisse D, Heitz A, Carnat A, Carnat A-P, Lamaison J-L. (2000) Quercetin 3-arabinopyranoside, a major flavonoid compound from *Alchemilla xanthochlora*. *Fitoterapia*, **71**, 463-464.
- [19] D'Agostino M, Dini I, Ramundo E, Senatore F. (1998) Flavonoid glycosides of *Alchemilla vulgaris* L. *Phytotherapy Research*, **12**, (Suppl. 1), S162–S163.

Ultrasound-assisted Extraction of Total Phenols and Flavonoids from Dry Tobacco (*Nicotiana tabacum*) Leaves

Ivana T. Karabegović, Vlada B. Veljković and Miodrag L. Lazić

Faculty of Technology, Bulevar oslobođenja 124, 16000 Leskovac, Serbia

icaneca@gmail.com

Received: October 12th, 2010; Accepted: October 23rd, 2011

Yields of extracted substances, as well as total phenol and flavonoid compounds obtained by classical and ultrasonic extractions from dry leaves of two tobacco types (oriental and Virginia) by two different solvents (acetone and methanol) at two operating temperatures (25 and 40°C) were compared. The yield of extractive, as well as total phenol and flavonoid compounds depended on the type of solvent, operational temperature and the tobacco type. The importance of these factors was assessed using 2⁴ full factorial experiments without replication.

Keywords: Total phenol content, flavonoids, extraction, full factorial experiments.

Tobacco (*Nicotiana tabacum* L.) leaves are the basic raw materials for cigarettes, directly influencing their quality and style, but are also highly appreciated material for perfumes, cosmetics and pharmaceuticals [1a]. Tobacco leaf is rich in polyphenols, which affect its color and quality [1a-1e]. Wang *et al.* extracted dry tobacco leaves with methanol through microwave-assisted technology and reached a total polyphenols yield of 22.4 mg/g (dry weight) [1a]. The yield of polyphenols from tobacco leaves extracted with 80% aqueous ethanol in the presence of ultrasound (40 kHz, 15 minutes, room temperature), reached 23.6 ± 2.3 mg/g dry weight [1e]. The present study deals with the effects of extraction conditions on the yields of total phenolic content and flavonoids from dry tobacco leaves. Both silent and ultrasound-assisted extractions were used to study the recovery of total extracted substances, phenols and flavonoids from oriental and Virginia tobacco leaves by methanol and acetone at 25 and 40°C. The statistical significance of extraction conditions was evaluated by 2⁴ full factorial experiments without replication.

As seen in Table 1, all four factors affected the yields of total phenolic content and flavonoids obtained after a 20 minute extraction from dry tobacco leaves. Independently of the recovery technique, operating temperature and tobacco type, methanol provided much higher extractive yields than acetone. The increase of extraction temperature slightly affected the extractive yields from both tobacco types, while the ultrasound-assisted extraction provided higher extractive substance yields than the classical method, approximately 8 and 24% in the case of oriental and Virginia types of tobacco, respectively. The data given in Table 1 show also that the extracts obtained by classical extraction had higher amounts of phenolics than those obtained by ultrasound extraction. These results confirmed the previously reported benefit of ultrasound action for

extraction from plant materials [2a-2c]. This was explained by oxidation and degradation of some compounds either during sonication or at a higher temperature [2c].

The major results of the analysis of variance (ANOVA), used for screening out the important factors, are given in Table 2. The experiment was not replicated, so the residual variance was determined based on the variance of three- and four-way interactions. That these interactions could be included in the residual variance, it was checked by the Bartlett's test for homogeneity of variances. Bartlett's test statistics of 0.87, 8.53 and 0.55 in the case of yield of extract, and total phenolic and flavonoid content were respectively smaller than the upper critical value of the chi-square distribution with four degrees and with a significance level of 0.05 ($\chi^2_{4;0.05} = 9.49$). Thus, the variances of three- and four-way interactions were judged to be equal. The obtained results confirm that for the total phenolic content, the important factors, with the significance level of 0.05, are the tobacco type, the type of solvent and the operational temperature, while the effects of the extraction technique are ignorable. All four factors, as well as the interaction of factors A and D and factors B and C are important, with the significance level of 0.05 in the case of the total flavonoid content.

Experimental

Plant materials: Dry oriental and Virginia tobacco (*Nicotiana tabacum* L.) leaves harvested in Bulgaria and Brazil, respectively, were used; moisture contents of dry tobacco leaves, determined by drying at 105°C to constant weight, were 6.7 and 7.2%, respectively.

Extraction procedures: Dry leaves were crushed and sieved (0.250 mm sieve). Methanol and acetone were utilized as extraction solvents and the resulting extracts were stored

Table 1: Total phenolic compounds and flavonoids of tobacco leaves extracts*.

Tobacco type	Extraction solvent	Extraction temperature	Extraction technique	Yield of extracted substances, g/100 g dry plant material	Total phenolic content, mg GAE/g dry extract	Total flavonoids, mg RE/g dry extract
Oriental	Acetone	25°C	CE ^a	7.4	122.0±0.97	100.2±1.67
			USE ^b	8.8	121.4±1.80	96.3±2.35
		40°C	CE	7.8	115.3±0.14	94.8±2.34
			USE	9.0	111.7±1.24	87.9±0.64
	MeOH	25°C	CE	22.6	105.9±1.80	51.8±0.12
			USE	23.4	99.7±1.94	50.2±0.30
		40°C	CE	24.7	90.0±1.94	49.7±0.67
			USE	26.6	87.8±3.87	43.9±0.91
Virginia	Acetone	25°C	CE	3.6	164.4±2.77	132.8±0.66
			USE	4.0	145.6±2.49	128.6±1.76
		40°C	CE	4.8	135.9±0.00	122.2±1.64
			USE	5.2	129.7±0.14	108.3±3.59
	MeOH	25°C	CE	18.2	117.3±0.41	59.1±0.83
			USE	21.2	110.6±2.07	54.8±0.51
		40°C	CE	22.4	116.3±1.66	55.7±0.65
			USE	23.4	102.6±3.18	43.6±1.26

*Data were expressed as the mean of three replicates ± standard deviation. ^a Classical (silent) extraction ^b Ultrasound extraction.

Table 2: Results of the variance analysis.

Source of variation	Degrees of freedom	Yield of extract		Total phenolic content		Total flavonoids		F _{1;50.95}
		Mean square	Test statistic	Mean square	Test statistic	Mean square	Test statistic	
A (Tobacco type)	1	62.02	28.66	1612.02	42.87	1061.13	108.01	6.61
B (Extraction technique)	1	2.33	1.08	132.25	3.52	122.66	12.49	
C (Temperature)	1	7.16	3.31	457.96	12.18	286.46	29.16	
D (Type of solvent)	1	1154.30	533.41	2595.90	69.04	13357.6	1359.69	
AB	1	0.77	0.36	125.44	3.34	3.9	0.40	
AC	1	2.98	1.38	28.09	0.75	33.93	3.45	
AD	1	0.031	0.01	180.90	4.81	565.25	57.54	
BC	1	0.53	0.24	13.32	0.35	68.48	6.97	
BD	1	0.031	0.01	4.41	0.12	0.68	0.07	
CD	1	1.38	0.64	64.00	1.70	29.43	3.00	
ABC+ ABD+ ACD+ BCD+ ABCD	5	10.85		187.99		49.12		
Total	15							

at 4°C. The plant material (5 g) and extraction solvent (50 mL) were put in Erlenmayer flasks (100 mL) and placed in an ultrasonic cleaning bath (Sonic, Niš, Serbia; total nominal power: 3 x 50 W; and internal dimensions: 30 x 15 x 20 cm) operating at a frequency of 40 kHz. Sonication was performed in 20 mins. The bath was filled with distilled water up to 1/3 of its volume (about 2.5 L). The extraction was carried out at 25 and 40°C. The temperature was maintained (± 0.1°C) by water circulating from a thermostated bath by means of a pump. The classical extraction was carried out with the ultrasound power switched off. At the end of the extraction procedure, the liquid extract was separated from the solid residue by vacuum filtration. The solid residue was washed twice using fresh solvent (20 mL). The filtrates were collected, the solvent evaporated in a rotary vacuum evaporator at 40°C. Each experiment was carried out in duplicate.

Determination of total phenolic and total flavonoid contents: The total phenolic and flavonoid contents of the extracts was determined by the Folin-Ciocalteu method [3a] and aluminum chloride colorimetric method [3b] respectively. Results were expressed as means of 3 replicate measurements.

Statistical analysis: The comparison of means was analyzed by Student's *t*-test and differences were considered significant when $p < 0.05$. The four factors' effects (tobacco type, extraction technique, solvent and temperature) on the yields of extractive substances, total phenolic content and flavonoids were analyzed by 2⁴ full factorial experiments with no replication.

Acknowledgements - This work was supported by the Ministry of Science of the Republic of Serbia; Projects OI 172047.

References

- [1] (a) Wang HY, Cui C, Zhao MM, Shen GL, Rao GH. (2008) Extraction optimization and component identification of tobacco polyphenols. *Huanan Ligong Daxue Xuebao*, **36**, 64-68; (b) Bazinet L, DeGrandpré Y, Porter A. (2005) Electromigration of tobacco polyphenols. *Separation and Purification Technology*, **4**, 101-107; (c) Bazinet L, DeGrandpré Y, Porter A. (2005) Enhanced tobacco polyphenol electromigration and impact on membrane integrity. *Journal of Membrane Science*, **25**, 111-118; (d) Kierulff JV, Jorgenson OB, Hansen TT, Knox AJ, De Grandpré Y. (2001) Use of a phenol oxidizing enzyme in the treatment of tobacco. *US Patent* 6298859; (e) Wang H, Zhao M, Yang B, Jiang Y, Rao G. (2008) Identification of polyphenols in tobacco leaf and their antioxidant and antimicrobial activities. *Food Chemistry*, **107**, 1399-1406.
- [2] (a) Stojičević S, Stanisavljević I, Veličković D, Veljković V, Lazić M. (2008) Comparative screening of antioxidant and antimicrobial activities of *Sempervivum marmoreum* L. extracts obtained by various extraction techniques. *Journal of the Serbian Chemical Society*, **73**, 597-607; (b) Stanisavljević IT, Lazić ML, Veljković VB. (2007) Ultrasonic extraction of oil from tobacco (*Nicotiana tabacum* L.) seeds. *Ultrasonic Sonochemistry*, **14**, 646-652; (c) Vinatoru M. (2001) An overview of the ultrasonically assisted extraction of bioactive principles from herbs. *Ultrasonic Sonochemistry*, **8**, 303-313.
- [3] (a) Singleton VL, Rossi JA. (1965) Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *American Journal of Enology and Viticulture*, **16**, 144-158; (b) Chang C, Yang M, Wen H, Chern J. (2002) Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *Journal of Food and Drug Analysis*, **10**, 178-182.

Characterization of Polyphenolic Compounds in Unripe Chinotto (*Citrus myrtifolia*) Fruit by HPLC/PDA/ESI/MS-MS

Monica Scordino^{ab*}, Leonardo Sabatino^a, Adalgisa Belligno^b and Giacomo Gagliano^a

^aMinistero delle Politiche Agricole Alimentari e Forestali, Dipartimento dell'Ispettorato Centrale della Tutela della Qualità e della Repressione Frodi dei Prodotti Agroalimentari (ICQRF) - Laboratorio di Catania, Via A. Volta 19, 95122 Catania, Italy

^bUniversità degli Studi di Catania, Dipartimento di Scienze delle Produzioni Agrarie e Alimentari (DISPA) – Sezione Scienze Agrochimiche, Via S. Sofia 98, 95123 Catania, Italy

m.scordino@mpaaf.gov.it

Received: July 1st, 2011; Accepted: September 25th, 2011

The flavonoid and furocoumarin composition was investigated of peel and pulp tissues of unripe fruits of *Citrus myrtifolia* Rafinesque, an ingredient of the popular soft drink "chinotto". Compound separation and identification was made using an HPLC-PDA detector coupled to ESI/MS/MS in positive and negative mode. Eighteen compounds (3-hydroxy-3-methylglutaryl-, C- and O-glycosyl flavonoids, furocoumarins and polymethoxylated flavones) were identified and quantified. Data indicated that the overall amount of flavonoids and furocoumarins in peel was higher than in the pulp, even though their relative distribution did not significantly change, apart from a different distribution of flavones and a lower content of naringin in the peel.

Keywords: chinotto, citrus, flavonoids, furocoumarins, peel, pulp.

Over the past few years, increasing attention has been paid by consumers to the nutritional aspects of products containing significant amounts of biologically active components. Nowadays, many studies are carried out on the thousands of natural phytochemicals that may have important physiological effects [1]. Scientific data demonstrated that an increased consumption of fruit and vegetables may protect against degenerative pathologies such as cancer and atherosclerosis [2]. Epidemiological studies have shown an inverse relationship between dietary flavonoids intake and cardiovascular diseases [3]. Many citrus flavonoids have been shown to have antioxidative activity to inhibit angiogenesis and to slow down cancer cell migration and proliferation [4,5]. Among the phytochemicals, flavonoids are widely contained in Citrus fruits [3].

Recently our research group has studied the unripe fruits of *Citrus myrtifolia* Rafinesque in terms of bioactive compounds and antioxidant activity; results indicated that it is a good source of phytochemicals, mainly vitamin C and flavonoids [6]. *Citrus myrtifolia* is a taxon of Citrus with foliage similar to that of the common myrtle. It is a compact tree with small leaves and no thorns which grows to a height of three meters and can be found in Malta and in the Liguria, Tuscany, Sicily, and Calabria regions of Italy. The fruit of the tree resemble small oranges. They are either sour or bitter and are commonly called by their

Italian name, chinotto. They are an essential flavoring agent of most Italian amari, of the popular Campari aperitif, and of several brands of carbonated soft drinks that are generically called "chinotto". Native of southern China, its origin has not been exactly ascertained; probably it is a mutation of sour orange that eventually evolved into the species known today [7]. The scientific literature about *C. myrtifolia* [8,9] is related prevalently to the juice and no information is available about bioactive compounds quantitation and distribution in the whole fruit. Interest in the Italian citrus industry for chinotto production could be increased by studies on its composition.

The aim of the present work was to elucidate by HPLC/PDA/ESI/MS-MS the identity and relative distribution of flavonoids and furocoumarins in pulp and peel tissues of unripe *C. myrtifolia* used for industrial beverage purposes.

Figure 1 shows the PDA chromatograms of unripe *C. myrtifolia* pulp and peel at different λ allowing for the discrimination of flavanone, furocoumarin and flavone derivatives. All peaks identified were numbered from 1 to 18. The UV spectra, recorded in correspondence with peaks 1, 2, 3, 7 and 9 showed absorptions at 270 and 350 nm (peaks 1, 3 and 9) and 270 and 335 nm (peaks 2 and 7), which can be attributed to bands II (due to the A ring benzoyl system) and I (associated with absorption due to




Figure 1: Typical PDA chromatograms of *C. myrtifolia* pulp and peel from immature fruits recorded at 280 (A, B) and 330 nm (C, D). Compounds 1–18 were identified as follows: 1, lucenin-2; 2, vicianin-2; 3, lucenin-2 4'-methyl ether; 4, eriocitrin; 5, neoeriocitrin; 6, naringin; 7 rhoifolin; 8, neohesperidin; 9, neodiosmin; 10 melitidin; 11, brutieridin; 12, poncirin; 13, bergapten; 14, epoxybergamottin; 15, sinensetin; 16, nobiletin; 17, heptamethoxyflavone; 18, tangeretin.

the B ring cinnamoyl system) of a flavone structure, respectively. The UV spectra of compounds 4, 5, 6, 8, 10, 11 and 12 indicated the flavanone nature of the aglycone, showing absorptions centered at 285 and 330 nm. Compounds 13 and 14 showed UV absorption maxima at 260 and 310, indicating a possible coumarin moiety. The lag-times and UV maxima (335 nm) of peaks from 15 to 18 suggested the presence of less polar polymethoxylated flavones.


The MS/MS spectra recorded in positive and negative mode in correspondence with the studied peaks and comparison with analytical standards and literature data, when available, permitted the unambiguous structural assignment for all compounds (Figures 2, 3 and 4), which were quantified separately in pulp and peel (Table 1).

Peaks 1–3 were identified as the flavone-*C*-glucosides lucenin-2, vicianin-2 and lucenin-2 4'-methyl ether. The precursor ions (m/z 611 ($[M+H]^+$) and 609 ($[M-H]^-$); m/z 595 ($[M+H]^+$) and 593 ($[M-H]^-$); 625 ($[M+H]^+$) and 623 ($[M-H]^-$) for 1, 2 and 3, respectively) were in agreement with their nominal masses. The product ions showed the characteristic loss of 120 amu, ascribable to the fragment mass of *C*-glucosyl flavonoids. These flavones were recently detected in chinotto juice [9, 10] and in taxonomically close Citrus juices, such as bergamot and sour orange [11–13]. Peak 7 was assigned to the flavone-*O*-glycoside rhoifolin, as confirmed by the molecular fragmentation pattern, which showed the precursor ion at 579 m/z and the loss of an *O*-diglycoside fragment mass at 273 m/z . Compound 9 was identified as a diosmetin aglycone *O*-linked with saccharide substituents; the MS/MS fragmentation pattern indicated the loss of a rhamnose unit ($[M+H-146]^+$) and a further loss of a glucose unit ($[M+H-308]^+$), revealing that the substituent is

a rhamnose–glucose disaccharide. Coelution with a standard allowed us to assign to this peak the structure of neodiosmin (7-*O*-neohesperidosil diosmetin). Neodiosmin presence in *Citrus* was earlier reported by Gattuso *et al.* in *Citrus bergamia* Risso juice [12].

All flavanone compounds showed the characteristic loss of 308 amu, corresponding to the *O*-diglycoside breakage. Moreover, two 3-hydroxy-3-methylglutaric acid conjugates of neohesperidin and naringin, namely, brutieridin and melitidin were identified as peaks 10 and 11, respectively. The positive and negative ESI/MS showed a $[M+H]^+$ ion at m/z 725 and 755 and a $[M-H]^-$ ion at m/z 723 and 753, which corresponded to the elemental composition of melitidin and brutieridin, respectively. A detailed analysis by tandem mass spectrometry showed highly diagnostic fragment ions: the loss of 3-hydroxy-3-methylglutaryl moiety was confirmed by the positive daughter ion at m/z 273 for melitidin, and at m/z 303 for brutieridin. These data are in agreement with those reported by Barreca *et al.* [9] and Scordino *et al.* [10] for chinotto juice.


Finally, the present paper reported the occurrence of polymethoxylated flavones (PMFs) (Figure 2) in *C. myrtifolia* (peaks 15–18). PMFs are a class of minor components usually found in the essential oil fraction of citrus peels [14]. Mass spectra generated in positive mode showed $[M+H]^+$ ions at m/z 403, 373 and 433, well-matched in fragmentation, in UV-vis spectra and retention times with data reported by Scordino *et al.* [15]. The parent ions $[M+H]^+$ were further fragmented and the breakage of methyl and methoxy groups were in agreement with the loss of 15 and 30 amu. Compounds 15 to 18 were identified as sinensetin, nobiletin, heptamethoxyflavone and tangeretin, respectively.



Peak	Structure assignment	R1	R2	R3	R4	R5	R6	R7
1	Luteolin 6,8-di-C-glucoside (Lucenin-2)	H	OH	Glu	OH	Glu	OH	OH
2	Apigenin 6,8-di-C-glucoside (Vicenin 2)	H	OH	Glu	OH	Glu	H	OH
3	Diosmetin 6,8-di-C-glucoside (Lucenin-2 4'-Me)	H	OH	Glu	OH	Glu	OH	OCH ₃
7	Apigenin 7-O-neohesperidoside (Rhoifolin)	H	OH	H	O-Nh*	H	H	OH
9	Diosmetin 7-O-neohesperidoside (Neodiosmin)	H	OH	H	O-Nh*	H	OH	OCH ₃
15	Sinensetin	H	OCH ₃	OCH ₃	OCH ₃	H	OCH ₃	OCH ₃
16	Nobiletin	H	OCH ₃	OCH ₃	OCH ₃	OCH ₃	OCH ₃	OCH ₃
17	3,5,6,7,8,3',4'-heptamethoxyflavone	OCH ₃	OCH ₃	OCH ₃	OCH ₃	OCH ₃	OCH ₃	OCH ₃
18	Tangeretin	H	OCH ₃	OCH ₃	OCH ₃	OCH ₃	H	OCH ₃

*Neohesperidoside


Figure 2: Structure assignments of identified flavones.



Peak	Structure assignment	R
13	5-Methoxypsoralen (Bergapten)	OCH ₃
14	5-(6',7'-Epoxy)geranyloxypsoralen (Epoxybergamottin)	

Figure 3: Structure assignments of identified furocoumarins.

Quantitative evaluation: Naringin and neohesperidin are the flavanone-*O*-glycosides found in the highest amounts in chinotto fruit (Table 1), ranging from about 200 to 175 mg/100 g in the pulp and from about 347 to 472 mg/100 g in the peel, respectively. Eriocitrin (6.9 mg/100 g) and poncirin (5.6 mg/100 g) were detected only in pulp, while neoerictin was found in both analyzed matrixes in significant quantity (58.9 and 131.5 mg/100 g in pulp and peel, respectively). The two 3-hydroxy-3-methylglutaryl flavanone glycosides melitidin and brutieridin were found to be present in large amounts (14.2 and 28.6 mg/100 g in the pulp and 30.3 and 95.4 mg/100 g in the peel, respectively). The flavone *O*-glycosides rhoifolin and neodiosmin were the most abundant flavones of chinotto tissues, ranging from about 7 mg/100 g in the pulp to about 29.5 and 57.4 mg/100 g in the peel, respectively. The peel was also characterized by about 30 mg/100 g of the flavone-*C*-glucosides lucenin-2, vicenin-2 and lucenin-2 4'-methyl ether; on the contrary, only traces of these compounds were detected in the pulp. The total amount of



Peak	Structure assignment	R1	R2	R3
4	Eriodictyol 7-O-rutinoside (Eriocitrin)	O-Rutinoside	OH	OH
5	Eriodictyol 7-O-neohesperidoside (Neoerictin)	O-Neohesperidoside	OH	OH
6	Naringenin 7-O-neohesperidoside (Naringin)	O-Neohesperidoside	H	OH
8	Hesperetin 7-O-neohesperidoside (Neoesperidin)	O-Neohesperidoside	OH	OCH ₃
10	Naringenin 7-(2''-α-rhamnosyl-6'''-(3'''-hydroxy-3'''-methylglutaryl)-β-glucoside (Melitidin)	3-Hydroxy-3-Methylglutaryl	OH	OCH ₃
11	Hesperetin 7-(2''-α-rhamnosyl-6'''-(3'''-hydroxy-3'''-methylglutaryl)-β-glucoside (Brutieridin)	3-Hydroxy-3-Methylglutaryl	H	OH
12	Isosakuranetin 7-O-neohesperidoside (Poncirin)	O-Neohesperidoside	H	OCH ₃


Figure 4: Structure assignments of identified flavanones.

the two furocoumarins bergapten and epoxybergamottin in the peel (25.1 mg/100 g) was two times that of the pulp (12.0 mg/100 g). A small but distinctive amount of polymethoxylated flavones (16.4 mg/100 g), among which nobiletin predominated, was detected in the peel; the same components were detectable in traces in the peel. It is well known that polyphenolic contents are usually higher in peels [3], therefore it is not surprising that the total amount of flavonoids and furocoumarins in the peel (1235 mg/100 g) was more than two times higher than that of the pulp (519 mg/100 g). The flavonoid contents were also determined in all samples by the method indicated by the Association of the Industry of Juices and Nectars (AIJN) using the colorimetric Davis method; data obtained were about 423.7 mg/100 g and 736.4 mg/100 g for the pulp and the peel, respectively. Taking into account the relative distribution of the different classes discussed above in the two tissues (Figure 5), no major changes could be observed between the pulp and the peel contribution to the total polyphenolics apart from some differences in the relative distribution of the flavones, and in the naringin amounts, which predominated in the pulp.

In conclusion, our results elucidated the distribution of flavonoids and furocoumarins in *C. myrtifolia* peel and pulp. The HPLC/PDA/ESI/MS-MS technique allowed the identification and relative quantification of 18 compounds, among which many compounds have been detected for the first time in *C. myrtifolia*. Nowadays these aspects are considered to be highly valuable for the commercial valorization of chinotto as a citrus with high potential as a nutraceutical source.

Table 1: Polyphenolic contents of *Citrus myrtifolia* pulp and peel.

		Pulp Mean	Peel Mean
1	Lucenin-2 ^b	traces	4.0 (0.2)
2	Vicenin-2 ^c	0.03 (0.01)	21.3 (1.3)
3	Lucenin-2 4'-OMe ^b	1.9 (0.2)	4.6 (0.2)
4	Eriocitrin	6.9 (0.3)	traces
5	Neoeriocitrin	58.9 (3.3)	131.5 (6.7)
6	Naringin	200.3 (10.7)	347.0 (17.6)
7	Rhoifolin ^c	6.6 (0.4)	29.5 (1.6)
8	Neohesperidin	175.7 (7.3)	472.4 (21.9)
9	Neodiosmin	7.4 (0.3)	57.4 (1.8)
10	Melitidin ^d	14.2 (0.8)	30.3 (1.5)
11	Brutieridin ^e	28.6 (0.9)	95.4 (5.2)
12	Poncirin	5.6 (0.3)	traces
13	Bergapten ^f	7.0 (0.6)	12.0 (0.7)
14	Epoxybergamottin ^f	5.0 (0.1)	13.1 (0.6)
15	Sinensetin	0.24 (0.02)	1.8 (0.2)
16	Nobiletin ^g	0.91 (0.12)	11.2 (0.5)
17	Heptamethoxyflavone ^g	traces	0.39 (0.03)
18	Tangeretin ^g	traces	3.0 (0.1)
	Total flavonoids (HPLC)	519.1 (13.5)	1235.0 (29.5)
	Total flavonoids (according to Davis)	423.7 (22.7)	736.4 (29.7)

^a mean value of five determinations; standard deviation in parentheses.^b expressed as neodiosmin; ^c expressed as apigenin; ^d expressed as naringin; ^e expressed as neohesperidin; ^f expressed as bergamottin; ^g expressed as sinensetin.**Figure 5:** Relative distribution (%) of flavonoids and furocoumarins in *C. myrtifolia* pulp and peel.**Table 2:** Polyphenolic contents in *Citrus myrtifolia* pulp and peel.

Parameter	Mean Value ^a	
Hunter color parameters	Peel	Pulp
a	36.3 (6.2)	5.4 (1.5)
b	-34.4 (4.2)	-14.3 (2.7)
L	66.3 (2.2)	78.8 (2.4)
Mean diameter (cm)	2.5 (0.3)	
Mean weight (g)	27 (2)	
Total soluble solids (Brix) ^b	8.4 (0.1)	
Titrate acidity (% citric acid) ^b	0.83 (0.05)	
pH ^b	3.87 (0.21)	
Ratio (Brix/acidity) ^b	10.1	

^a mean value of five determinations; standard deviation in parentheses.^b referred to the juice.

Experimental

Reagents and standards: HPLC-grade acetonitrile, methanol and formic acid were supplied by Romil (Milan, Italy). Distilled water was purified at 18.2 MΩ cm with a MilliQ ULTRA [Millipore, Vimodrone (MI), Italy] purification system. Apigenin, bergamottin, neodiosmin, neoeriocitrin, naringin, neohesperidin, poncirin and sinensetin were obtained from Extrasynthèse (Genay, France).

Plant materials: Fruits used in this study were grown in Castiglione di Sicilia (Catania, Italy). Immature *Citrus myrtifolia* (chinotto) fruits were picked in October 2010 with the degree of fruit maturity determined from the Hunter colour parameters, mean diameter, total soluble solids, titratable acidity and pH (Table 2). Samples for analysis were prepared from 25 fruits; pulp and peel tissues were separated and analyzed individually. Peels and pulps represented 55% and 38% of the studied sample, while the remaining part was constituted by seeds. The samples were stored at -20°C until needed for the study.

Extraction and analysis of flavonoids and furocoumarins: Five g of sample was extracted for 2 h with 10 mL of 50% aqueous methanol containing 0.3% formic acid at room temperature on an orbital shaker. The mixture was centrifuged at 1000 rpm for 15 min and the supernatant decanted. The pellets were re-extracted under identical conditions. Supernatants were combined and used for the LC analyses, after filtration through 0.45 μm PTFE filters (LabService Analytica, Bologna, Italy). The juice was analyzed before centrifugation and filtration through 0.45 μm PTFE filters. The analyses were performed with a liquid chromatograph consisting of a Finnigan Surveyor MS-pump, autosampler and photodiode-array detector (PDA), coupled with a Finnigan LCQ DECA XP MAX detector (Thermo Fisher Scientific). The analytical column was a Luna C18 250 x 4.6 mm, 5 μm i.d. (Phenomenex), the flow rate was 1 mL/min, the column temperature 30°C and the injection volume 20 μL. Flow rate was split 1/10 before MS interface. A binary gradient of 0.3% formic acid in water (A) and 0.3% formic acid in acetonitrile (B) was employed. The mobile phase gradient was programmed as follows: 0 min, 5% B; 50 min, 28% B; 60 min, 43% B; 60-65 min, 43% B; 70-80 min, 5% B. The range of wavelengths examined by the photodiode-array detector was 190–700 nm, and for quantitative determinations the chromatograms were recorded at 285 nm (flavanones), 310 nm (furocoumarins), 335 nm (polymethoxylated flavones) and 340 nm (flavones). Mass spectral analyses were performed using a LCQ ion-trap mass operating in negative and positive ion mode using an ion spray LC/MS interface. The electrospray ionization (ESI) needle voltage was 4.0 kV. The capillary voltage was 18V and the heated capillary was 250°C. A sheath gas flow rate of 36 (arbitrary units) was used and the auxiliary gas was set to 12 (arbitrary units). The MS-MS spectra were obtained using collision energy of 25% of instrument maximum, operating in selected reaction monitoring (SRM). Preliminary positive and negative tunings were carried out with continuous introduction of dilute solutions of sinensetin and naringin, respectively, at a flow rate of 5 μL/min and the voltages on the lenses were optimized in TunePlus (Excalibur software).

The chinotto polyphenolics were characterized in terms of retention times, lambda max, MS and MS/MS data operating in positive and negative mode as follows:

Compound 1: Rt, 16.9 min; UV, 270, 350 nm. MS, 611 [M + H]⁺ (100); MS, 609 [M - H]⁻ (100). MS-MS focused on [M + H]⁺, 611 [M + H]⁺ (20), 593 [M + H-18]⁺ (100), 575 [M + H - 36]⁺ (10), 545 [M + H - 66]⁺ (5), 491 [M + H - 120]⁺ (15), 473 [M + H - 138]⁺ (15).

Compound 2: Rt, 23.5 min; UV, 270, 335 nm. MS, 595 [M + H]⁺ (100); MS, 593 [M - H]⁻ (100). MS-MS focused on [M + H]⁺, 595 [M + H]⁺ (20), 577 [M + H-18]⁺ (100), 559 [M + H - 36]⁺ (10), 529 [M + H - 66]⁺ (5), 475 [M + H - 120]⁺ (15), 457 [M + H - 138]⁺ (10).

Compound 3: Rt, 26.3 min; UV, 270, 350 nm. MS, 625 [M + H]⁺ (100); MS, 623 [M - H]⁻ (100). MS-MS focused on [M + H]⁺, 625 [M + H]⁺ (70), 607 [M + H -18]⁺ (100), 559 [M + H - 66]⁺ (5), 505 [M + H - 120]⁺ (8), 463 [M + H - 162]⁺ (70), 445 [M + H -180]⁺ (8).

Compound 4: Rt, 29.9 min; UV, 285, 330 nm. MS, 597 [M + H]⁺ (100); MS, 595 [M - H]⁻ (100). MS-MS focused on [M + H]⁺, 597 [M + H]⁺ (100), 451 [M + H - 146]⁺ (40), 289 [M + H - 308]⁺ (80).

Compound 5: Rt, 31.8 min; UV, 285, 330 nm. MS, 597 [M + H]⁺ (100); MS, 595 [M - H]⁻ (100). MS-MS focused on [M + H]⁺, 597 [M + H]⁺ (100), 451 [M + H - 146]⁺ (65), 289 [M + H - 308]⁺ (90).

Compound 6: Rt, 37.4 min; UV, 285, 330 nm. MS, 581 [M + H]⁺ (100); MS, 579 [M - H]⁻ (100). MS-MS focused on [M + H]⁺, 581 [M + H]⁺ (10), 563 [M + H-18]⁺ (30), 545 [M + H - 36]⁺ (25), 435 [M + H - 146]⁺ (65), 419 [M + H - 162]⁺ (100), 315 [M + H - 266]⁺ (25), 271 [M + H - 308]⁺ (10).

Compound 7: Rt, 40.0 min; UV, 265, 335 nm. MS, 579 [M + H]⁺ (100); MS, 577 [M - H]⁻ (100). MS-MS focused on [M + H]⁺, 579 [M + H]⁺ (50), 432 [M + H-146]⁺ (50), 273 [M + H - 308]⁺ (100).

Compound 8: Rt, 40.4 min; UV, 285, 330 nm. MS, 611 [M + H]⁺ (100); MS, 609 [M - H]⁻ (100). MS-MS focused on [M + H]⁺, 611 [M + H]⁺ (50), 593 [M + H-18]⁺ (30), 575 [M + H - 36]⁺ (30), 490 [M + H - 120]⁺ (15), 449 [M + H - 162]⁺ (80), 345 [M + H - 266]⁺ (20), 303 [M + H - 308]⁺ (70).

Compound 9: Rt, 42.2 min; UV, 255, 340 nm. MS, 609 [M + H]⁺ (100); MS, 607 [M - H]⁻ (100). MS-MS focused on [M + H]⁺, 609 [M + H]⁺ (50), 463 [M + H -146]⁺ (40), 302 [M + H - 308]⁺ (100). MS-MS focused on [M - H]⁻, 607 [M - H]⁻ (40), 299 [M - H - 308]⁻ (100).

Compound 10: Rt, 46.2 min; UV, 285, 330 nm. MS, 725 [M + H]⁺ (100); MS, 723 [M - H]⁻ (100). MS-MS focused on [M + H]⁺, 725 [M + H]⁺ (5), 707 [M + H -18]⁺ (100), 461 [M + H - 264]⁺ (90), 273 [M + H - 452]⁺ (10). MS-MS focused on [M - H]⁻, 723 [M - H]⁻ (100), 661 [M - H - 62]⁻ (10), 621 [M - H - 102]⁻ (20), 579 [M - H - 144]⁻ (80).

Compound 11: Rt, 48.2 min; UV, 285, 325 nm. MS, 755 [M + H]⁺ (100); MS-MS focused on [M + H]⁺, 755 [M +

H]⁺ (100), 609 [M + H - 146]⁺ (30), 301 [M + H - 452]⁺ (50); MS-MS focused on [M - H]⁻, 753 [M - H]⁻ (100); 691 [M + H - 62]⁻ (10); 651 [M + H - 102]⁻ (15); 609 [M + H - 144]⁻ (40).

Compound 12: Rt, 50.7 min; UV, 285, 330 nm. MS, 595 [M + H]⁺ (100); MS, 593 [M - H]⁻ (100). MS-MS focused on [M + H]⁺, 595 [M + H]⁺ (20), 577 [M + H-18]⁺ (30), 559 [M + H-36]⁺ (40), 449 [M + H-146]⁺ (70), 433 [M + H-162]⁺ (100), 287 [M + H-308]⁺ (25). MS-MS focused on [M - H]⁻, 593 [M - H]⁻ (10), 473 [M - H-120]⁻ (30), 431 [M - H-162]⁻ (15), 327 [M - H-266]⁻ (30), 285 [M - H-308]⁻ (100).

Compound 13: Rt, 57.4 min; UV, 260, 315 nm. MS, 217 [M + H]⁺ (100). MS-MS focused on [M + H]⁺, 217 [M + H]⁺ (10), 202 [M + H - 30]⁺ (100).

Compound 14: Rt, 59.3 min; UV, 260 (sh), 310 nm. MS, 355 [M + H]⁺ (100). MS-MS focused on [M + H]⁺, 355 [M + H]⁺ (30), 337 [M + H - 18]⁺ (100).

Compound 15: Rt, 61.3 min; UV, 335 nm. MS, 373 [M + H]⁺ (100). MS-MS focused on [M + H]⁺, 373 [M + H]⁺ (100), 358 [M + H - 15]⁺ (50), 312 [M + H - 61]⁺ (30).

Compound 16: Rt, 62.0 min; UV, 335 nm. MS, 403 [M + H]⁺ (100). MS-MS focused on [M + H]⁺, 403 [M + H]⁺ (60), 388 [M + H - 15]⁺ (100), 373 [M + H - 30]⁺ (50).

Compound 17: Rt, 62.8 min; UV, 335 nm. MS, 433 [M + H]⁺ (100). MS-MS focused on [M + H]⁺, 434 [M + H]⁺ (60), 419 [M + H - 15]⁺ (100), 403 [M + H - 30]⁺ (50).

Compound 18: Rt, 63.6 min; UV, 335 nm. MS, 373 [M + H]⁺ (100). MS-MS focused on [M + H]⁺, 373 [M + H]⁺ (40), 358 [M + H - 15]⁺ (100), 343 [M + H - 30]⁺ (40).

The external calibrations were obtained using methanolic standard solutions of known concentration (1-100 mg/L). Lucenin-2 and lucenin-2 4'-methyl ether were expressed as neodiosmin; vicenin-2 and rhoifolin as apigenin; meltidin and brutieridin as naringin and neohesperidin, respectively; bergapten and epoxybergamottin in terms of bergamottin relative amount; and nobiletin, heptamethoxyflavone and tangeretin in terms of sinensetin relative amount.

The total flavonoid content was also determined in all of the samples according to the colorimetric method of Davis [16] and the values were expressed as mg naringin/100 g FW (fresh weight).

Statistical analysis: To verify the statistical significance of all parameters the values of means and standard deviation (SD) were calculated. Where appropriate, the data were tested by two-way ANOVA. P<0.05 was adopted as statistically significant. All data are means of 5 measurements.

References

- [1] Patil BS, Jayaprakasha GK, Murthy KNC, Vikram A. (2009) Bioactive compounds: historical perspectives, opportunities, and challenges. *Journal of Agricultural and Food Chemistry*, 57, 8142-8160.
- [2] Shahidi F, Marian N. (2003) *Phenolics in Food and Nutraceuticals*, CRC Press, Boca Raton.

- [3] Tripoli E, La Guardia M, Giammanco S, Di Majo D, Giammanco M. (2007) Citrus flavonoids: Molecular structure, biological activity and nutritional properties: A review. *Food Chemistry*, **104**, 466–479.
- [4] Cao G, Sofic E, Prior R. (1997) Antioxidant and prooxidant behavior of flavonoids: Structure–activity relationships. *Free Radical Biology and Medicine*, **22**, 749–760.
- [5] Rice-Evans C, Miller NJ, Paganda G. (1996) Structure–antioxidant activity relationship of flavonoids and phenolic acids. *Free Radical and Medicine*, **20**, 933–956.
- [6] Scordino M, Sabatino L, Belligno A, Gagliano G. (2011) Preliminary study on bioactive compounds of *Citrus x myrtifolia* Rafinesque (Chinotto) to its potential application in food industry. *Food and Nutritional Sciences*, **2**, 685–691.
- [7] Hanelt P, Büttner R, Mansfeld, R. (2001) *Mansfeld's encyclopedia of agricultural and horticultural crops (except ornamentals)*. Springer, Germany.
- [8] Cautela D, Pirrello AG, Espositom C, Minasi P. (2004) Caratteristiche compositive del chinotto (*Citrus myrtifolia*). *Essenze dei Derivati Agrumari*, **74**, 51–57.
- [9] Barreca D, Bellocco E, Caristi C, Leuzzi U, Gattuso G. (2010) Flavonoid composition and antioxidant activity of juices from Chinotto (*Citrus x myrtifolia* Raf.) fruits at different ripening stages. *Journal of Agricultural and Food Chemistry*, **58**, 3031–3036.
- [10] Scordino M, Sabatino L, Belligno A, Gagliano G. (2011) Flavonoids and furocoumarins distribution of unripe chinotto (*Citrus x myrtifolia* Rafinesque) fruit: beverage processing homogenate and juice characterization. *European Food Research and Technology*, **233**, 759–767.
- [11] Barreca D, Bellocco E, Caristi C, Leuzzi U, Gattuso G. (2011) Distribution of C- and O-glycosyl flavonoids, (3-hydroxy-3-methylglutaryl)glycosyl flavanones and furocoumarins in *Citrus aurantium* L. juice. *Food Chemistry*, **124**, 576–582.
- [12] Gattuso G, Barreca D, Caristi C, Gargiulli C, Leuzzi U. (2007) Distribution of flavonoids and furocoumarins in juices from cultivars of *Citrus bergamia* Risso. *Journal of Agricultural and Food Chemistry*, **55**, 9921–9927.
- [13] Caristi C, Bellocco E, Gargiulli E, Toscano G, Lezzi E. (2006) Flavone-di-C-glycosides in citrus juices from Southern Italy. *Food Chemistry*, **95**, 431–437.
- [14] Miyake Y, Yamamoto K, Morimitsu Y, Osawa T. (1997) Isolation of C-glucosylflavone from lemon peel and antioxidative activity of flavonoid compounds in lemon fruit. *Journal of Agricultural and Food Chemistry*, **45**, 4619–4623.
- [15] Scordino M, Sabatino L, Traulo P, Gargano M, Panto' V, Gagliano G. (2011) HPLC–PDA/ESI–MS/MS detection of polymethoxylated flavones in highly degraded citrus juice: a quality control case study. *European Food Research and Technologies*, **232**, 275–280.
- [16] Davis WB. (1947) Determination of flavanones in citrus fruits. *Analytical Chemistry*, **19**, 476–477.

Bioactive Compounds, RP-HPLC Analysis of Phenolics, and Antioxidant Activity of Some Portuguese Shrub Species Extracts

Ângelo Luís, Fernanda Domingues and Ana Paula Duarte*

CICS-UBI Health Sciences Research Centre, University of Beira Interior, Av. Infante D. Henrique
6200-506 Covilhã, Portugal

apcd@ubi.pt

Received: August 12th, 2011; Accepted: October 24th, 2011

In the ecosystem of Serra Da Estrela, some plant species have the potential to be used as raw material for extraction of bioactive products. The goal of this work was to determine the phenolic, flavonoid, tannin and alkaloid contents of the methanolic extracts of some shrubs (*Echinopartum ibericum*, *Pterospartum tridentatum*, *Juniperus communis*, *Ruscus aculeatus*, *Rubus ulmifolius*, *Hakea sericea*, *Cytisus multiflorus*, *Crataegus monogyna*, *Erica arborea* and *Ipomoea acuminata*), and then to correlate the phenolic compounds and flavonoids with the antioxidant activity of each extract. The Folin-Ciocalteu's method was used for the determination of total phenols, and tannins were then precipitated with polyvinylpyrrolidone (PVPP); a colorimetric method with aluminum chloride was used for the determination of flavonoids, and a Dragendorff's reagent method was used for total alkaloid estimation. The 2,2-diphenyl-1-picrylhydrazyl (DPPH) and β -carotene bleaching tests were used to assess the antioxidant activity of extracts. The identification of phenolic compounds present in extracts was performed using RP-HPLC. A positive linear correlation between antioxidant activity index and total phenolic content of methanolic extracts was observed. The RP-HPLC procedure showed that the most common compounds were ferulic and ellagic acids and quercetin. Most of the studied shrubs have significant antioxidant properties that are probably due to the existence of phenolic compounds in the extracts. It is noteworthy to emphasize that for *Echinopartum ibericum*, *Hakea sericea* and *Ipomoea acuminata*, to the best of our knowledge, no phytochemical studies have been undertaken nor their use in traditional medicine been described.

Keywords: Phenolic compounds, flavonoids, antioxidant activity, vegetal extracts, RP-HPLC.

The Portuguese forest area occupies around 3.3 million hectares, which represents about 38% of the territory. The Serra Da Estrela Mountain supports a rich plant cover constituting the crown of Portugal's interior ecological network. In this ecosystem grow numerous species of trees and shrubs. Examples of shrubs that grow spontaneously in Portuguese forests in general and in this region in particular are *Echinopartum ibericum* Rivas Mart., Sánchez-Mata & Sancho, *Pterospartum tridentatum* (L.) Willk subsp. *cantabricum* Spach, *Juniperus communis* L. subsp. *alpina* (Suter) Celak., *Ruscus aculeatus* L., *Rubus ulmifolius* Schott, *Hakea sericea* Schrader, *Cytisus multiflorus* (L'Hér) Sweet, *Crataegus monogyna* Jacq., *Erica arborea* L., and *Ipomoea acuminata* (Vahl) Roemer & Schultes (*I. indica* Burm.).

The genus *Echinopartum* (Spach) Fourr. (family Fabaceae), mostly endemic to the Iberian Peninsula, is composed of both silicolous and calcicolous taxa, growing in mountain areas usually above 1000 meters above sea level [1]. The authors are not aware of any known application of these species in folk medicine.

Pterospartum tridentatum (Fabaceae; subfamily Papilionoideae) grows spontaneously in Portugal, where it

is known as Carqueja or Carqueija. The flowers are used in traditional medicine for the treatment of throat irritation conditions and in herbal mixtures for diabetes [2]. No phytochemical, pharmacological and toxicological studies have been reported for this species, but alkaloids and isoflavonoids are characteristic secondary metabolites of Fabaceae species [2].

Juniperus communis or common juniper is a coniferous shrub distributed throughout the Arctic and temperate zone of the northern hemisphere. Its dried bluish-black cones, known as "juniper berries", are said to stimulate the appetite and are used as a flavoring agent for culinary purposes and in the preparation of gin spirits [3]. They have also been used for various medicinal purposes, including as an abortifacient, contraceptive, diuretic, and as a remedy for urinary tract infections, chest complaints, diabetes, rheumatism and backache. More interestingly, juniper has been reported as a traditional cure for chest troubles such as bronchitis and for tuberculosis [3].

An alcoholic infusion of the rhizomes of *Ruscus aculeatus* has been used for the treatment of some venous ailments for decades [4].

Leaves and young shoots of *Rubus ulmifolius* are used in folk medicine for their anti-inflammatory, anti-odontalgic and gastrointestinal spasmolytic properties; crushed young shoots are applied to wounds, infected insect bites and pimples [5].

A key characteristic of *Hakea sericea* is its extreme serotinous habit: all of its seeds are retained in pairs in tough woody follicles, which accumulate along the branches throughout the life of the plant [6]. As far as we know, neither medicinal uses nor phytochemical studies are described to this shrub.

Cytisus spp. is used as a diuretic, hypnotic, sedative, anti-diabetic and also as a hepatoprotective [7].

Crataegus monogyna has been used as a folk medicine due its sedative actions, protective effects against arrhythmias and increase of coronary vessel flow [8]. Previous chemical studies on *C. monogyna* resulted in isolation of known flavonoids and phenolic acids [8].

Erica multiflora is used to treat hyperlipidaemia, and in folk medicine as a diuretic and antiseptic agent [9]. The phytochemical study of this plant showed that tannins, proanthocyanidols and flavonoids represent major compounds of its flowers [9].

Ipomoea species accumulate high levels of anthocyanin pigments in the storage root [10]. To the best of our knowledge, no medical applications are described for this plant.

Several constituents of plant extracts have been shown to have antioxidant activity, such as ascorbic acid, tocopherol, β -carotene, flavonoids, tannins, phenolics, and anthocyanins [11]. Oxidation is essential to many organisms for the production of energy to fuel biological processes. However, reactive oxygen species (ROS) are often over-produced under pathological conditions, resulting in oxidative stress. An over-production of various forms of activated oxygen species, such as free-radical and non-free-radical species is involved in the onset of many diseases such as cancer, cardiovascular diseases, rheumatoid arthritis, and atherosclerosis, as well as in degenerative processes associated with aging [11]. In order to reduce damage to the human body, synthetic antioxidants are used in industrial processing. However, the presence of unwanted side effects is almost unavoidable, and some have been suspected of being responsible for liver damage and carcinogenesis [11]. Thus, it is essential to develop antioxidants that can protect the human body from free radicals and retard the progress of many chronic diseases.

No single chemical component is responsible for the medicinal properties of plant-based drugs, and their synergic action or bioenhancement is due to the presence of several chemical substances in the plant material.

Therefore, the determination of the total amount of different classes of components is essential for the standardization of the plants [12].

The purpose of this work was to determine the phenolic, flavonoid, tannin and total alkaloid contents of the methanolic extracts of the species above mentioned, and then to correlate phenolic compounds and flavonoids with antioxidant activity of corresponding extracts. The Folin-Ciocalteu's method was used for the determination of total phenols, and tannins were then precipitated with polyvinylpyrrolidone (PVPP); a colorimetric method with aluminum chloride was used for the determination of total flavonoids, and a Dragendorff's reagent method was used for alkaloid estimation. A 2,2-diphenyl-1-picrylhydrazyl (DPPH) method and a β -carotene bleaching test were used to assess the antioxidant activity of extracts. The identification of phenolic compounds present in extracts was performed using a RP-HPLC technique.

From the results presented in Table 1 it can be verified that fruits of *Juniperus communis* had a higher extraction yield than the other shrubs. By contrast, the extract of fruits of *Hakea sericea* produced the lowest extraction yield. The extraction yields obtained in the present study are very similar to those reported for some medicinal plants reported in the literature [13].

Medicinal plants are commonly rich in phenolic compounds, such as flavonoids, phenolic acids, stilbenes, tannins, coumarins, lignans and lignins. They have multiple biological properties including antioxidant activity. The antioxidant activity of phenolic acids and flavonoids is due to their redox properties, ability to chelate metals and quenching of singlet oxygen [14]. In this sense, it is reasonable to determine the amount of phenolic compounds in the extracts of these plants.

The quantitative determination of phenolic compounds using Folin-Ciocalteu's method is a widespread assay. [15]. As can be observed in Table 2, in general, the extracts from stems and leaves have the highest amount of phenolic compounds. The extract of stems of *Crataegus monogyna* is the one that had the greatest concentration of total phenols. This result is similar to those obtained by other researchers [16]. Barros *et al.* studied the composition and bioactivity of *Crataegus monogyna* and concluded that phenolics were the major antioxidant components present in this shrub [17]. *Ipomoea acuminata* and *Ruscus aculeatus* are species that are poor in these compounds. Many studies indicate that the roots of sweet potato (*Ipomoea batatas*) are a source of bioactive compounds [18]. In previous studies by our group, we concluded that the ethanolic extract of *Cytisus* spp. had more phenolics than that of the aqueous extract [19]. Methanolic extracts of the different parts of *Cytisus multiflorus* have significant amounts of total phenols. *Pterospartum tridentatum* and *Erica arborea* are also species rich in these compounds. Fruits of *Rubus*

Table 1: Extraction yields of methanolic extracts.

Species	Plant part	Extraction yield (%) *
<i>Echinopartum ibericum</i>	Stems and Leaves	21.1 ± 0.1
	Flowers	19.7 ± 0.4
<i>Pterospartum tridentatum</i>	Stems and Leaves	26.1 ± 1.7
	Flowers	25.1 ± 1.4
<i>Juniperus communis</i>	Stems	15.6 ± 2.1
	Leaves	31.4 ± 2.0
<i>Ruscus aculeatus</i>	Fruits	45.7 ± 4.8
	Stems and Leaves	18.3 ± 3.5
<i>Rubus ulmifolius</i>	Stems	9.9 ± 1.0
	Leaves	23.8 ± 5.6
	Flowers	22.6 ± 0.7
	Fruits	27.5 ± 2.1
<i>Hakea sericea</i>	Stems	11.0 ± 0.1
	Leaves	16.4 ± 2.4
	Fruits	3.2 ± 0.1
	Stems	10.1 ± 0.6
<i>Cytisus multiflorus</i>	Leaves	15.5 ± 1.5
	Flowers	27.6 ± 2.7
	Fruits	25.0 ± 1.1
<i>Crataegus monogyna</i>	Stems	13.1 ± 1.7
	Leaves	23.1 ± 3.1
	Flowers	23.5 ± 9.9
	Fruits	10.2 ± 1.5
<i>Erica arborea</i>	Stems	12.9 ± 0.1
	Leaves	29.5 ± 1.8
	Flowers	38.6 ± 0.7
<i>Ipomoea acuminata</i>	Stems	9.3 ± 1.7
	Leaves	12.0 ± 0.3
	Flowers	14.4 ± 5.4

* Results in terms of mean ± standard deviation

ulmifolius are the part of this shrub that has less phenolic compounds, in contrast with stems, leaves and flowers. Other studies [20] demonstrated that this shrub had a lower concentration of phenols than the one now estimated. There are no literature data for *Echinopartum ibericum* and *Hakea sericea* phenol contents, but in this work they were shown to be rich in these compounds. Öztürk *et al.* conducted an exhaustive study of several species of Turkish juniper, and concluded that *Juniperus communis* is rich in bioactive compounds, and that methanol is one of the best solvents for their extraction [21]. Regarding the determination of tannins (Table 2), it can be concluded that the composition varies in the different extracts (it was not possible to observe a trend). Significant differences were also found in the tannin content among different parts of the same shrub, which is in agreement with the behavior described for tannins in plants.

Flavonoids are polyphenolic compounds that occur ubiquitously in plants and structurally have variations in one of the aromatic rings that characterize the different types, namely, flavonols, flavones, isoflavones, flavonones, flavanol and anthocyanins [14,22]. Flavonoids could act as antioxidants because they can readily donate an electron or a hydrogen atom to a peroxy or alkoxy radical to terminate a lipid peroxidation chain reaction or to regenerate a phenolic/flavonoid compound, which can effectively chelate a pro-oxidant transition metal [22]. The process used in quantitative methods for flavonoid determination is performed by precipitating them with aluminum chloride in an alkalized medium. This precipitate acquires a color and, therefore, lends itself to the colorimetric assay. The flavonoids in the presence of

aluminum chloride have an intense yellow fluorescence when observed under filtered UV light [23]. As can be seen in Figure 1, extracts from flowers and fruits generally have more flavonoids than the extracts of stems and leaves of the same shrub species. However, *Ipomoea acuminata* had the highest flavonoids content of the tested species, especially the extract of leaves. The results obtained for the studied species are very similar to those of our previous studies for other Portuguese shrubs [19]. Andrade *et al.*, have studied the bioactivity of *Cistus ladanifer* and *Arbutus unedo* extracts, and the flavonoids contents were very similar to those presented here [24]. In the present study, *Erica arborea*, *Cytisus multiflorus*, *Rubus ulmifolius* and *Pterospartum tridentatum* are noted for being species with large concentrations of flavonoids. There are no previous studies in the literature about *Hakea sericea* and *Echinopartum ibericum* on the characterization of flavonoids; however, large amounts of these compounds were detected in all these species. Previous research on flavonoids of various species of juniper showed that this shrub could be a source of these biologically active compounds [25]. The results now obtained showed that the leaves of this species are the richest part of the plant in these compounds. Methanol was shown to be a good solvent for flavonoids, as its high polarity favors the extraction of compounds with aromatic rings possessing hydroxyl groups.

Alkaloids are responsible for the therapeutic effect of many plant materials, but several alkaloids are toxic. Over 200 alkaloids have been identified in 300 plant species of up to 13 families. It has been estimated that up to 3% of the world's flowering plants contain toxic alkaloids [26]. The alkaloids were precipitated as BiI₃(Alk·HI) by Dragendorff's reagent (KBiI₄). Bismuth forms a yellow bismuth complex {Bi[CS(NH₂)₃]}(NO₃)₃ in nitric acid medium with thiourea. The bismuth from the alkaloidal complex is completely released by disodium sulfide [12].

Looking at Figure 1 it could be noted that all studied extracts have significant amounts of alkaloids. *Cytisus multiflorus* is the richest species in these compounds, specially the extracts of the stems. This species is known to possess alkaloids, namely spartein, with anti-arrhythmic properties [7]. The extract of leaves of *Ipomoea acuminata* also has a high concentration of alkaloids. By contrast, *Juniperus communis* is the poorest species in alkaloids. The extract of the leaves of that shrub is the one that has the smallest concentration of alkaloids.

One problem in assessing antioxidant activity is that this activity is variable and depends on the method used. It is known that an antioxidant mechanism in various biological matrices is very complex and several factors may intervene [27]. Given this complexity, only one method to determine the antioxidant activity of the extracts is not sufficient to draw relevant conclusions, and thus, we have applied two different methods and determined diverse antioxidant properties.

Table 2: Total phenolic compounds and tannin contents of methanolic extracts.

Species	Plant part	Total phenolic compounds (mg GAE / g dry matter) *	Tannin content (mg GAE / g dry matter)
<i>Echinopartum ibericum</i>	Stems and Leaves	99.1 ± 1.5	26.3
	Flowers	71.9 ± 0.8	N/D
<i>Pterospartum tridentatum</i>	Stems and Leaves	113.6 ± 1.5	43.4
	Flowers	171.4 ± 0.7	55.3
<i>Juniperus communis</i>	Stems	221.3 ± 2.5	79.3
	Leaves	155.6 ± 3.2	60.4
	Fruits	44.7 ± 2.6	N/D
<i>Ruscus aculeatus</i>	Stems and Leaves	32.9 ± 0.3	N/D
	Stems	186.8 ± 3.8	0.5
<i>Rubus ulmifolius</i>	Leaves	167.6 ± 6.1	21.9
	Flowers	291.4 ± 0.4	132.7
	Fruits	54.5 ± 0.6	6.8
<i>Hakea sericea</i>	Stems	267.6 ± 5.9	74.8
	Leaves	217.0 ± 2.7	103.5
	Fruits	110.1 ± 2.7	N/D
<i>Cytisus multiflorus</i>	Stems	142.4 ± 2.0	11.1
	Leaves	176.1 ± 3.6	61.5
	Flowers	120.4 ± 0.9	21.1
	Fruits	155.1 ± 5.6	20.6
<i>Crataegus monogyna</i>	Stems	377.4 ± 4.5	134.0
	Leaves	225.5 ± 2.5	81.1
	Flowers	186.2 ± 2.7	86.5
	Fruits	55.3 ± 2.7	N/D
<i>Erica arborea</i>	Stems	270.2 ± 4.3	133.8
	Leaves	260.2 ± 1.9	92.5
	Flowers	178.1 ± 0.2	83.0
<i>Ipomoea acuminata</i>	Stems	23.5 ± 5.1	N/D
	Leaves	32.2 ± 1.3	N/D
	Flowers	84.3 ± 1.7	N/D

* Results in terms of mean ± standard deviation; N/D – Not detected.

Results for the DPPH radical scavenging assay are reported as IC₅₀ values, which are defined as the amount of antioxidant required to inhibit 50% of DPPH free radicals under the experimental conditions [28]. Table 3 shows that *Echinopartum ibericum*, *Ruscus aculeatus* and *Ipomoea acuminata* had the poorest antioxidant activity in this assay, whereas that of stems of *Crataegus monogyna* had both the greatest concentration of total phenols and, simultaneously, the greatest antioxidant activity. In contrast, *C. monogyna* fruits had only moderate antioxidant activity. Other researchers demonstrated that the ethanolic extract of berries of *C. monogyna* possessed DPPH free-radical-scavenging activity [29]. All *Erica arborea* and *Hakea sericea* extracts had very strong antioxidant activity. As far as we know, no medicinal uses are described for this latter shrub and no previous phytochemical studies have been carried out on it, but it is rich in phenols with antioxidant activity. The results obtained for the parameters of antioxidant activity for the standard compounds are very similar to those determined by the same method by Faustino *et al.*, and Scherer and Godoy [30,31].

The β-carotene/linoleic acid bleaching assay is one of the antioxidant assays suitable for plant samples [28,32]. All the shrub extracts tested presented some degree of antioxidant properties when compared with the synthetic antioxidant BHT. The extracts of *Echinopartum ibericum* (Figure 2) showed a high degree of inhibition of lipid oxidation (linoleic acid) and sequestration of radicals

resulting from its possible oxidation, when compared with the extracts of *Pterospartum tridentatum* and *Hakea sericea* (Figure 2).

In contrast with the results of the DPPH scavenging assay, the extracts of *Echinopartum ibericum* had no significant ability to sequester free radicals such as DPPH, but displayed a high degree of inhibition of lipid oxidation, which may predict a potential use as a food antioxidant. The extract of *Pterospartum tridentatum* flowers showed the weakest inhibition of lipid oxidation. *Rubus ulmifolius* (Figure 3) showed the highest antioxidant potential measured by this method. By contrast, *Ipomoea acuminata* (Figure 3) had no capacity to promote inhibition of lipid oxidation. Stems of *Juniperus communis* (Figure 4) is the part of this shrub that provided higher percentages of inhibition, and the extracts of leaves and fruits presented antioxidant activity similar to that of the extract of *Ruscus aculeatus* (Figure 4).

Other studies [21] undertaken with juniper species allow us to conclude that in the β-carotene-linoleic acid assay, oxidation of linoleic acid was effectively inhibited by methanol extracts of various species of *Juniperus*, which is in agreement with the results now obtained. *Erica arborea*, *Cytisus multiflorus* and *Crataegus monogyna* were grouped in the same graph (Figure 5) because they have very similar antioxidant activity. It should be noted the extract of flowers of *Cytisus multiflorus* (Figure 5) presented percentages of inhibition slightly higher than those of the other extracts.

According to Figure 6, there was a positive linear correlation between antioxidant activity index and total phenolic content of methanolic extracts ($R^2=0.8442$). These results indicate that the phenolic compounds could be the main contributor to the antioxidant properties of these shrubs. This result is in agreement with several previous studies [19,33]. For example Fu *et al.* studied the antioxidant activity and phenolic compounds of 56 wild fruits from South China and found that the results showed a positive linear correlation between the antioxidant capacities and total phenolic content ($R^2>0.86$) [33]. The same analysis was conducted for the flavonoids (Figure 6) and it was verified that there was no correlation ($R^2=0.0083$) between these compounds and the antioxidant activity of the extracts. It is known that only flavonoids of a certain structure and particular hydroxyl position in the molecule determine the antioxidant properties; in general these properties depend on the ability to donate either hydrogen or an electron to a free radical [34].

RP-HPLC analysis was employed to identify major phenolic compounds in the methanolic shrub extracts. Satisfactory separation with good resolution could be achieved with the method now used. Such data will be helpful to compare the antioxidant activities with the




Figure 1: Concentration of flavonoids and alkaloids in studied extracts (* Results in terms of mean \pm standard deviation).




Figure 2: Antioxidant activity of the extracts of *Echinopartum ibericum*, *Pterospartum tridentatum* and *Hakea sericea* measured by β -carotene bleaching test.

phenolic compounds of the different studied shrubs and will be also useful to understand their chemical constituents and functionality. RP-HPLC analysis is the most used method to identify plant phenolics. Because of the diversity and complexity of natural phenolic compounds in hundreds of medicinal plant extracts, it is rather difficult to characterize every compound and elucidate its structure. It is not difficult, however, to identify major categories of phenolic compounds and representative phenolics. In the present study, we identified representative phenolic compounds from selected plants. A total of nine phenolic compounds were identified and quantified in the methanolic extracts of the shrubs (Table 4), including hydroxybenzoic acids, hydroxycinnamic acids and flavonoids. The total amount




Figure 3: Antioxidant activity of the extracts of *Rubus ulmifolius* and *Ipomoea acuminata* measured by β -carotene bleaching test.

of phenolic compounds in all the extracts was 1380.8 mg/g of dry matter. Gallic acid is one of the most important hydroxybenzoic acids, but, surprisingly, in the extracts now analyzed, it was present in low amounts (1.26%). However, the hydroxycinnamic acids identified (caffeic acid, chlorogenic acid, *p*-coumaric acid, ferulic acid) were abundantly present. Ferulic acid was the most dominant hydroxycinnamic acid, taking a percentage of occurrences of 29.0 %. These results concerning phenolic acids are in agreement with previously published data for other biological samples [35]. Ellagic acid and quercetin are also present in large amounts, with percentages of occurrence of 17.9% and 13.7%, respectively. Vanillic acid is the phenolic compound that is present in greater quantity in the extract of the stems and leaves of *Pterospartum tridentatum*, having a concentration of 32.2 mg/g of dry matter, corresponding to a percentage of 27.6 %. *Juniperus*




Figure 4: Antioxidant activity of the extracts of *Ruscus aculeatus* and *Juniperus communis* measured by β -carotene bleaching test.




Figure 5: Antioxidant activity of the extracts of *Erica arborea*, *Cytisus multiflorus* and *Crataegus monogyna* measured by β -carotene bleaching test.




Figure 6: Correlation between antioxidant activity and total phenolics and flavonoids of extracts.

communis, *Ruscus aculeatus*, *Hakea sericea* and *Ipomoea acuminata* are species relatively poor in phenolic compounds. A total of 34.5% of the analyzed phenols of the extract of *Rubus ulmifolius* leaves correspond to ferulic acid, with a concentration of 18.5 mg/g dry matter. In the extract of fruits of the same species, chlorogenic acid is the predominant compound. *Cytisus multiflorus* is rich in ferulic acid and ellagic acid, especially its leaves, flowers and fruits. The extract of stems of *Erica arborea* presents a large amount of ferulic acid, while the extracts of flowers

Table 3: Antioxidant properties of methanolic extracts studied and standard compounds.

Species / Standard	Plant part	IC ₅₀ (mg/L) *	AAI *	Antioxidant Activity
<i>Echinopartum ibericum</i>	Stems and Leaves	162.7 ± 13.8	0.3 ± 0.02	Poor
	Flowers	125.3 ± 12.4	0.3 ± 0.02	Poor
<i>Pterospartum tridentatum</i>	Stems and Leaves	69.7 ± 11.9	0.7 ± 0.06	Moderate
	Flowers	26.1 ± 1.3	1.7 ± 0.06	Strong
<i>Juniperus communis</i>	Stems	12.4 ± 0.3	3.4 ± 0.07	Very Strong
	Leaves	16.8 ± 0.6	2.4 ± 0.07	Very Strong
	Fruits	81.1 ± 6.9	0.6 ± 0.05	Moderate
<i>Ruscus aculeatus</i>	Stems and Leaves	171.9 ± 21.1	0.3 ± 0.03	Poor
	Stems	14.7 ± 1.2	2.7 ± 0.2	Very Strong
<i>Rubus ulmifolius</i>	Leaves	17.3 ± 1.2	2.5 ± 0.4	Very Strong
	Flowers	11.0 ± 0.5	3.2 ± 0.2	Very Strong
	Fruits	78.8 ± 5.9	0.7 ± 0.04	Moderate
<i>Hakea sericea</i>	Stems	9.5 ± 0.1	3.7 ± 0.03	Very Strong
	Leaves	13.4 ± 0.4	2.6 ± 0.06	Very Strong
	Fruits	28.3 ± 1.8	1.2 ± 0.07	Strong
	Stems	64.9 ± 3.2	0.8 ± 0.05	Moderate
<i>Cytisus multiflorus</i>	Leaves	35.1 ± 1.4	1.4 ± 0.05	Strong
	Flowers	71.5 ± 8.8	0.7 ± 0.09	Moderate
	Fruits	77.0 ± 8.6	0.7 ± 0.07	Moderate
<i>Crataegus monogyna</i>	Stems	8.5 ± 0.2	5.3 ± 0.1	Very Strong
	Leaves	16.5 ± 0.3	2.7 ± 0.05	Very Strong
	Flowers	13.6 ± 0.2	3.3 ± 0.05	Very Strong
	Fruits	86.5 ± 12.5	0.6 ± 0.06	Moderate
<i>Erica arborea</i>	Stems	14.6 ± 1.5	3.1 ± 0.3	Very Strong
	Leaves	10.9 ± 0.1	3.1 ± 0.04	Very Strong
	Flowers	16.1 ± 0.7	2.1 ± 0.1	Very Strong
<i>Ipomoea acuminata</i>	Stems	77.3 ± 6.0	0.7 ± 0.06	Moderate
	Leaves	204.1 ± 12.8	0.3 ± 0.09	Poor
	Flowers	57.4 ± 1.4	0.9 ± 0.03	Moderate
Rutin	---	10.7 ± 0.4	4.9 ± 0.2	Very Strong
Quercetin	---	4.3 ± 0.4	12.2 ± 1.7	Very Strong
Trolox	---	8.4 ± 0.1	6.6 ± 0.1	Very Strong
Gallic Acid	---	2.2 ± 0.02	22.8 ± 0.3	Very Strong

* Results in terms of mean ± standard deviation

are rich in ellagic acid. The extract of *Crataegus monogyna* flowers has the greatest concentration of ferulic acid (116.8 mg/g, dry matter), which corresponds to a percentage of 75.7%, relatively to the total phenolics quantified in this extract.

Experimental

Plant material: Aerial parts (stems, leaves, flowers and fruits) of *Echinopartum ibericum*, *Pterospartum tridentatum*, *Juniperus communis*, *Ruscus aculeatus*, *Rubus ulmifolius*, *Hakea sericea*, *Cytisus multiflorus*, *Crataegus monogyna*, *Erica arborea* and *Ipomoea acuminata* were collected in Serra Da Estrela. Table 5 shows the GPS coordinates and altitude at which the different shrubs were collected. Plant materials were dried at 35°C in a ventilated oven during 48 h and reduced to coarse powder (< 2 mm) using a laboratory cutting mill. Harvesting, transport and storage of plant species were authorized by *Instituto da Conservação da Natureza e da Biodiversidade*. All vegetal species were identified by a botanist and a voucher specimen of all species used has been deposited in the Herbarium of the *Instituto Superior de Agronomia (Jardim Botânico d'Ajuda, Lisboa)*. Table 5 lists the full scientific name of the vegetal species and the number of the voucher specimens.

Extraction process: Methanolic extracts were obtained using a Soxhlet apparatus until the solvent became colorless, using approximately 100 g of raw material and 1 L of solvent. The extracts were filtered under vacuum

using a crucible of porosity #2 then distilled under vacuum to remove the solvents to a final volume of 100 mL. Then, 5 mL of each extract was diluted in 45 mL of methanol. Aliquots (5 mL) of the extracts were removed for subsequent evaporation to dryness for the calculation of extraction yield and extract concentration.

Determination of total phenolic compounds: The phenols were determined by Folin-Ciocalteu's colorimetric method. The methanolic solutions of each extract (50 μ L) or gallic acid (standard phenolic compound) were mixed with 450 μ L of distilled water, and then 2.5 mL of Folin-Ciocalteu's reagent 0.2 N (diluted with distilled water) was added. The mixtures were allowed to stand for 5 mins, and then 2 mL of aqueous Na_2CO_3 (75 g/L) was added. After incubation of these reaction mixtures (90 mins / 30°C), the total phenols were determined by colorimetry at 765 nm. The standard curve was prepared using 500, 400, 350, 325, 300, 250, 225, 200, 150, 125, 100 and 50 mg/L solutions of gallic acid in methanol ($y=0.0009x$; $R^2=0.9875$). Total phenol values were expressed as gallic acid equivalents (mg GAE / g of dry mass) [36,37]. The assays were conducted in triplicate.

Tannin content: Tannins were measured as the difference in total phenolics (measured by Folin-Ciocalteu's reagent) before and after treatment with insoluble polyvinylpyrrolidone (PVPP), as this polymer binds strongly to tannins [38]. About 1 mL of each methanolic extract was added to 1 mL of an aqueous solution of PVPP (70 mg/mL). After vigorous shaking, the samples remained for 15 mins at 4°C, to develop the tannin-PVPP complex [39,40]. Then the samples were centrifuged for 10 mins at 3000 rpm; the tannins were found in the residue and the free phenols/non-adsorbed phenolics in the supernatant. Total phenols in the supernatant were determined by the Folin-Ciocalteu colorimetric method, as described above, and the concentration of tannins was calculated as the difference between the total phenols and free phenols, and is expressed in their respective units [39,40]. These determinations were made in triplicate.

Flavonoids determination: The aluminum chloride colorimetric method was used for flavonoids determination according to Pourmorad *et al.* [37]. Each methanol extract (500 μ L) was separately mixed with 1.5 mL of methanol, 0.1 mL 10 % aluminum chloride, 0.1 mL of 1 M potassium acetate, and 2.8 mL of distilled water. This solution remained at room temperature for 30 mins; the absorbance of the reaction mixture was measured at 415 nm using a spectrophotometer. The calibration curve was constructed by preparing 8 quercetin solutions at concentrations ranging from 12.5 to 200 μ g/mL in methanol ($y=0.0074x$; $R^2=0.9980$). Total flavonoid values were expressed as quercetin equivalents (mg QE / g of dry mass) [36]. These determinations were made in duplicate.

Alkaloid estimation: The alkaloid estimation was performed by a spectrophotometric method, as described

by Sreevidya and Mehotra [12]. Briefly, 10 mL of each crude extract was centrifuged for 10 mins (3000 rpm) to remove residual suspended particles and then 5 mL of the supernatant was mixed with 1 mL of HCl, 0.1 N. Then, 2.5 mL of Dragendorff's reagent was added to the previous mixture, for precipitation and the precipitate was centrifuged for 5 mins (3000 rpm). This precipitate was further washed with 2.5 mL of ethanol. The filtrate was discarded and the residue was then treated with 2.5 mL of disodium sulfide solution (1%, w/v). The brownish black precipitate formed was then centrifuged (5 mins, 3000 rpm). This residue was dissolved in 2 mL of concentrated nitric acid, with warming, if necessary; this solution was diluted to 10 mL in a standard flask with distilled water and 1 mL was then pipetted out and mixed with 5 mL of thiourea solution (3%, w/v). The absorbance of this solution was measured at 435 nm against a blank containing 1 mL of concentrated nitric acid and 2.5 mL of thiourea solution (3%, w/v). The standard curve was prepared using 750, 500, 400, 250, 200, 150 and 100 mg/L solutions of pilocarpine nitrate in HCl, 0.1N ($y=0.0013x-0.2750$; $R^2=0.9957$). Alkaloid contents were expressed as pilocarpine nitrate equivalents (mg PNE / g of dry mass) [12]. These tests were performed in duplicate.

Evaluation of antioxidant activity

DPPH scavenging assay: The antioxidant activity of the extracts and standards was determined by the radical scavenging activity method using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical [31]. Briefly, 0.1 mL aliquots of methanolic solutions of the extracts or standards at different concentrations were added to 3.9 mL of a DPPH methanolic solution. Three DPPH solutions were tested, 0.2000, 0.1242 and 0.0800 mM that were prepared by dissolving 39.4, 24.5 and 15.8 mg in 500 mL of methanol, respectively. These concentrations were selected due to the linearity range of DPPH solutions: above 0.2 mM the concentration is very high, and below 0.5 mM the color is very weak having a limited range of absorbance reading. The control sample consisted of a solution of 0.1 mL of methanol mixed with 3.9 mL of DPPH. After a 90 mins incubation period at room temperature in the dark, the absorbance was measured at 517 nm. The radical scavenging activity was calculated as follows: $I \% = [(Abs_0 - Abs_1) / Abs_0] \times 100$, where Abs_0 was the absorbance of the control and Abs_1 was the absorbance in the presence of the test sample at different concentrations. The IC_{50} was calculated graphically using a calibration curve in the linear range by plotting the extract concentration vs. the corresponding scavenging effect. The antioxidant activity was expressed as the antioxidant activity index (AAI), calculated as follows: $AAI = (\text{final concentration of DPPH in the control sample} - \mu\text{g.mL}^{-1}) / (IC_{50} - \mu\text{g.mL}^{-1})$ [31]. Thus, the AAI was calculated considering the mass of DPPH and the mass of the tested sample in the reaction, resulting in a constant for each sample, independent of the concentration of DPPH and sample used. In this work, it was considered that shrub

Table 4: Phenolic content of extracts determined by RP-HPLC.

Species	Plant part	Standard Phenolic Compounds (mg/g dry matter)*								
		Gallic Acid	Vanillic Acid	Caffeic Acid	Chlorogenic Acid	Syringic Acid	p-Coumaric Acid	Ferulic Acid	Ellagic Acid	Quercetin
<i>Echinopartum ibericum</i>	Stems and Leaves	1.9 ± 0.07	N/D	N/D	8.1 ± 1.6	15.9 ± 0.2	5.4 ± 0.2	23.8 ± 3.2	6.2 ± 1.8	25.3 ± 0.1
	Flowers	N/D	5.0 ± 1.0	3.8 ± 0.6	8.5 ± 0.4	16.3 ± 3.1	5.8 ± 0.4	13.5 ± 3.9	40.1 ± 2.6	2.2 ± 0.7
<i>Pteropartum tridentatum</i>	Stems and Leaves	N/D	32.2 ± 2.6	18.2 ± 3.0	17.8 ± 0.6	8.3 ± 0.8	16.8 ± 3.2	7.4 ± 0.8	9.7 ± 1.8	6.6 ± 0.3
	Flowers	N/D	4.2 ± 0.9	3.1 ± 0.3	N/D	1.9 ± 0.4	N/D	22.2 ± 0.6	10.4 ± 0.4	9.3 ± 0.7
<i>Juniperus communis</i>	Stems	N/D	N/D	N/D	2.2 ± 0.8	N/D	2.1 ± 0.4	2.0 ± 0.07	2.9 ± 0.2	6.7 ± 0.1
	Leaves	N/D	N/D	1.1 ± 0.0	N/D	N/D	2.8 ± 0.6	3.7 ± 0.3	12.7 ± 0.3	9.8 ± 0.7
	Fruits	N/D	N/D	N/D	N/D	N/D	1.0 ± 0.04	0.9 ± 0.2	1.8 ± 0.1	N/D
<i>Ruscus aculeatus</i>	Stems and Leaves	N/D	N/D	N/D	N/D	N/D	3.0 ± 0.08	N/D	N/D	4.1 ± 0.4
	Stems	1.8 ± 0.3	2.5 ± 0.1	N/D	8.2 ± 0.9	3.6 ± 0.2	3.5 ± 0.1	13.1 ± 0.8	1.3 ± 0.08	N/D
<i>Rubus ulmifolius</i>	Leaves	7.0 ± 0.4	1.1 ± 0.02	2.1 ± 0.2	4.2 ± 0.8	3.6 ± 0.5	4.1 ± 0.9	18.5 ± 0.6	4.9 ± 0.5	8.2 ± 0.5
	Flowers	3.3 ± 0.3	3.3 ± 0.5	4.6 ± 0.2	6.7 ± 0.3	4.7 ± 0.2	7.6 ± 0.5	7.0 ± 0.2	15.6 ± 0.9	33.3 ± 2.2
	Fruits	N/D	1.4 ± 0.2	0.8 ± 0.09	6.1 ± 0.2	0.9 ± 0.3	1.2 ± 0.08	3.6 ± 0.2	0.7 ± 0.09	1.3 ± 0.1
<i>Hakea sericea</i>	Stems	N/D	N/D	N/D	N/D	N/D	2.1 ± 0.6	11.3 ± 1.1	3.5 ± 0.6	N/D
	Leaves	N/D	1.2 ± 0.08	1.2 ± 0.1	1.9 ± 0.2	2.8 ± 0.4	3.8 ± 0.4	21.5 ± 0.9	2.6 ± 0.06	6.0 ± 0.5
	Fruits	1.2 ± 0.2	0.7 ± 0.07	1.1 ± 0.2	N/D	0.7 ± 0.01	1.7 ± 0.07	1.9 ± 0.2	3.7 ± 0.6	1.6 ± 0.03
<i>Cytisus multiflorus</i>	Stems	N/D	1.2 ± 0.6	1.0 ± 0.2	3.4 ± 0.1	2.7 ± 0.2	6.9 ± 0.8	1.8 ± 0.2	9.2 ± 0.3	37.1 ± 2.6
	Leaves	1.6 ± 0.02	2.3 ± 0.6	2.0 ± 0.1	3.2 ± 0.02	1.0 ± 0.08	3.8 ± 0.8	5.7 ± 0.8	16.1 ± 2.0	7.9 ± 1.9
	Flowers	N/D	N/D	N/D	N/D	0.7 ± 0.04	1.3 ± 0.3	13.0 ± 1.6	5.4 ± 0.4	3.5 ± 0.5
	Fruits	N/D	4.3 ± 0.6	2.0 ± 0.2	2.2 ± 0.4	2.9 ± 0.2	4.2 ± 0.3	25.4 ± 4.2	16.0 ± 1.0	3.0 ± 0.08
<i>Crataegus monogyna</i>	Stems	N/D	8.2 ± 0.5	7.7 ± 1.0	20.5 ± 1.4	4.4 ± 0.6	5.1 ± 0.7	17.2 ± 1.5	9.4 ± 1.0	N/D
	Leaves	N/D	18.9 ± 2.4	4.9 ± 0.9	10.3 ± 1.2	2.0 ± 0.3	5.3 ± 0.6	11.0 ± 1.6	7.4 ± 1.1	1.7 ± 0.2
	Flowers	0.6 ± 0.1	9.7 ± 1.5	4.4 ± 0.3	1.5 ± 0.3	1.8 ± 0.2	1.3 ± 0.06	116.8 ± 8.4	8.4 ± 1.2	9.8 ± 0.5
	Fruits	N/D	0.5 ± 0.05	N/D	3.0 ± 0.3	N/D	0.6 ± 0.1	3.3 ± 0.4	1.2 ± 0.1	0.6 ± 0.03
<i>Erica arborea</i>	Stems	N/D	1.3 ± 0.05	3.6 ± 0.6	0.8 ± 0.1	4.4 ± 0.4	3.1 ± 0.2	7.7 ± 0.9	5.1 ± 0.8	1.2 ± 0.3
	Leaves	N/D	15.1 ± 0.7	3.9 ± 0.3	17.5 ± 1.8	16.3 ± 1.0	7.5 ± 0.6	25.6 ± 1.5	9.3 ± 1.9	N/D
	Flowers	N/D	6.4 ± 0.1	2.3 ± 0.05	1.7 ± 0.3	2.5 ± 0.4	3.5 ± 0.3	6.0 ± 0.3	23.4 ± 0.9	N/D
<i>Ipomoea acuminata</i>	Stems	N/D	2.2 ± 0.1	N/D	N/D	N/D	N/D	2.4 ± 0.2	N/D	N/D
	Leaves	N/D	1.4 ± 0.2	N/D	N/D	N/D	1.0 ± 0.1	1.0 ± 0.1	N/D	9.8 ± 0.4
	Flowers	N/D	4.8 ± 0.2	N/D	N/D	N/D	2.6 ± 0.4	12.8 ± 1.7	20.5 ± 2.3	N/D

* Results in terms of mean ± standard deviation; N/D – Not detected

Table 5: GPS coordinates and altitude at which the different shrubs were collected and number of the vouchers specimens.

Full Scientific Name	GPS coordinates	Altitude (m)	Number of Voucher Specimen
<i>Echinopartum ibericum</i> Rivas Mart., Sánchez-Mata & Sancho	N 40°19.046' W 07°34.365'	1606	LISI 8/2011
<i>Pteropartum tridentatum</i> (L.) Willk subsp. <i>cantabricum</i> Spach	N 40°18.825' W 07°33.630'	1541	LISI 9/2011
<i>Juniperus communis</i> L. subsp. <i>alpina</i> (Suter) Celak.	N 40°19.046' W 07°34.365'	1606	LISI 10/2011
<i>Ruscus aculeatus</i> L.	N 40°19.105' W 07°26.791'	567	LISI 11/2011
<i>Rubus ulmifolius</i> Schott	N 40°19.167' W 07°27.247'	519	LISI 12/2011
<i>Hakea sericea</i> Schrader	N 40°20.296' W 07°27.491'	730	LISI 13/2011
<i>Cytisus multiflorus</i> (L'Hér) Swett	N 40°18.668' W 07°27.076'	504	LISI 14/2011
<i>Crataegus monogyna</i> Jacq.	N 40°18.668' W 07°27.076'	504	LISI 15/2011
<i>Erica arborea</i> L.	N 40°19.046' W 07°34.365'	1606	LISI 16/2011
<i>Ipomoea acuminata</i> (Vahl) Roemer & Schultes	N 40°16.719' W 07°30.744'	649	LISI 18/2011

extracts showed poor antioxidant activity when AAI < 0.5, moderate antioxidant activity between 0.5 and 1.0, strong antioxidant activity between 1.0 and 2.0, and very strong when AAI > 2.0 [31]. Assays were carried out in duplicate and all DPPH solutions were prepared daily.

β-Carotene bleaching test: After preparation of β-carotene solution (20 mg/mL in chloroform), 20 µL was added to 40 µL of linoleic acid, 400 mg of Tween 40 and 1 mL of chloroform. This mixture was then evaporated at 45°C for

5 mins in a rotary vacuum evaporator to remove chloroform and immediately diluted with 100 mL of oxygenated distilled water. The water was added slowly to the mixture and vigorously agitated to form an emulsion. Five mL of the emulsion was transferred into test tubes containing 300 µL of extracts in methanol at different concentrations. About 5 mL of the emulsion and 300 µL of samples in methanol were used as control. Standard butylated hydroxytoluene (BHT) in methanol, at the same concentration as samples, was used as reference. The tubes were then gently shaken and placed at 50°C in a water bath for 2 h. The absorbances of the extracts, standard and control were measured at 470 nm, using a spectrophotometer, against a blank consisting of an emulsion without β-carotene. The measurements were carried out at initial time (t=0 h) and at final time (t=2 h). The antioxidant activity was measured in terms of percentage of inhibition of β-carotene's oxidation by:

$$\% \text{ Inhibition} = (\text{Abs}^{t=2}_{\text{sample}} - \text{Abs}^{t=2}_{\text{control}}) / (\text{Abs}^{t=0}_{\text{control}} - \text{Abs}^{t=2}_{\text{control}})$$

Where Abs^{t=2} was the absorbance of the sample or control at final time of incubation and Abs^{t=0} was the absorbance in the control at initial time of incubation [19].

RP-HPLC analysis of phenolics: Chromatographic analysis was performed with the use of a liquid chromatographic system, which consisted of a Perkin Elmer Binary LC Pump 250, Perkin Elmer UV-visible Spectrophotometric Detector LC 290 and Perkin Elmer LC Oven 101 with a loop of 50 µL. The chromatographic system was connected through the Data Apex U-PAD2

USB Acquisition Device to a PC computer. Software used for data acquisition and evaluation was Clarity Lite Data Apex. The separation was carried out on a 150×4.60 mm, Phenomenex Kinetex Luna 2.6µ PFP 100A reversed phase column equipped with a Phenomenex KrudKatcher Ultra HPLC In-Line Filter with 0.5 µm porosity. Standard solutions were filtered through a 0.22 µm pore membrane filter before injection and crude methanolic extracts were centrifuged and then filtered through a 0.22 µm pore membrane filter before injection. The injection volume for all samples was 50 µL. The mobile phase consisted of 2 solvents: Solvent A, water with acetic acid (pH=3) and Solvent B, acetonitrile/solvent A (6:4; v/v). The mobile phase was filtered through a 0.22 µm pore membrane filter and degasified with nitrogen. Phenolic compounds were eluted under the following conditions: 1 mL/min flow rate and the column temperature was set at 35°C, isocratic conditions from 0 to 10 mins with 0% B, linear gradient conditions from 0% to 5% B in 30 mins, from 5% to 15% B in 18 mins, from 15% to 25% B in 14 mins, from 25% to 50% B in 31 mins, from 50% to 100% B in 3 mins, followed by washing and reconditioning the column. For detection, chromatograms were monitored by ultra-violet-visible spectra (280 nm), which were recorded for all

peaks. Duplicate analyses were performed for each sample. The identification of phenolic compounds was obtained by using authentic standards, while quantification was performed using the external standard method. Stock solutions of standard compounds at a concentration of 1 mg/mL each were prepared in methanol, and several dilutions with methanol were made. The solutions of standards at various concentrations (1.00, 0.75, 0.50, 0.25, 0.15 mg/mL) were injected into the HPLC system and the calibration curves were established for each standard compound. The concentration of the compound was calculated from peak area according to calibration curves. The amount of each phenolic compound was expressed as mg/g of dry mass [35,41,42].

Acknowledgments - Ângelo Luís acknowledges a fellowship from *Fundação Para a Ciência e a Tecnologia* (Reference: SFRH / BD / 65238 / 2009). Authors wish to thank Prof. Dr Dalila Espírito Santo and Dr Paula Paes, from the Herbarium of the *Instituto Superior de Agronomia (Jardim Botânico d'Ajuda, Lisboa)*, for the identification and deposit of the vouchers specimens of the plant species used in this work.

References

- [1] Aparicio A, Guisande R. (1997) Replenishment of the endangered *Echinopartum albigicum* (Genisteae, Fabaceae) from the soil seed bank. *Biological Conservation*, **81**, 267-273.
- [2] Vitor R, Mota-Filipe H, Teixeira G, Borges C, Rodrigues A, Teixeira A, Paulo A. (2004) Flavonoids of an extract of *Pterospartum tridentatum* showing endothelial protection against oxidative injury. *Journal of Ethnopharmacology*, **93**, 363-370.
- [3] Gordien A, Gray A, Franzblau S, Seidel V. (2009) Antimycobacterial terpenoids from *Juniperus communis* L. (Cupressaceae). *Journal of Ethnopharmacology*, **126**, 500-505.
- [4] Mimaki Y, Kuroda M, Yokosuka A, Sashida Y. (1999) A spirostanol saponin from the underground parts of *Ruscus aculeatus*. *Phytochemistry*, **51**, 689-692.
- [5] Sisti M, Santi M, Fraternali D, Ninfali P, Scoccianti V, Brandi G. (2008) Antifungal activity of *Rubus ulmifolius* Schott standardized in *in vitro* culture. *LWT-Food Science and Technology*, **41**, 946-950.
- [6] Le Maitre D, Krug R, Hoffmann J, Gordon A, Mgidi T. (2008) *Hakea sericea*: Development of a model of the impacts of biological control on population dynamics and rates of spread of an invasive species. *Ecological Modelling*, **212**, 342-358.
- [7] Raja S, Ahamed K, Kumar V, Mukherjee K, Bandyopadhyay A, Murkherjee P. (2007) Antioxidant effect of *Cytisus scoparius* against tetrachloride treated liver injury in rats. *Journal of Ethnopharmacology*, **109**, 41-47.
- [8] García M, Sáenz M, Ahumada M, Cert A. (1997) Isolation of three triterpenes and several aliphatic alcohols from *Crataegus monogyna* Jacq.. *Journal of Chromatography A*, **767**, 340-342.
- [9] Harfani H, Bouanani N, Aziz M, Caid H, Ghalim N, Amrani S. (2007) The hypolipidaemic activity of aqueous *Erica multiflora* flowers extract in Triton WR-1339 induced hyperlipidaemic rats: A comparison with fenofibrate. *Journal of Ethnopharmacology*, **109**, 156-160.
- [10] Terahara N, Konczak I, Ono H, Yoshimoto M, Yamakawa O. (2004) Characterization of acylated anthocyanins in callus induced from storage root of purple-fleshed sweet potato, *Ipomoea batatas* L. *Journal of Biomedical Biotechnology*, **5**, 279-286.
- [11] Xu W, Zhang F, Luo Y, Ma L, Kou X, Huang K. (2009) Antioxidant activity of a water-soluble polysaccharide purified from *Pteridium aquilinum*. *Carbohydrate Research*, **344**, 217-222.
- [12] Sreedvidya N, Mehotra S. (2003) Spectrophotometric method for estimation of alkaloids precipitable with Dragendorff's reagent in plant materials. *Journal of AOAC International*, **86**, 1124-1127.
- [13] Silva E, Souza J, Rogez H, Rees J, Larondelle Y. (2007) Antioxidant activities and polyphenolic contents of fifteen selected plant species from the Amazonian regions. *Food Chemistry*, **101**, 1012-1018.
- [14] Surveswaran S, Cai, Y, Corke H, Sun M. (2007) Systematic evaluation of natural phenolic antioxidant from 133 Indian medicinal plants. *Food Chemistry*, **102**, 938-953.
- [15] Cicco N, Lanorte M, Paraggio M, Viggiano M. (2009) A reproducible, rapid and inexpensive Folin-Ciocalteu micro-method in determining phenolics of plant methanol extracts. *Microchemical Journal*, **91**, 107-110.
- [16] Baborun T, Aumjaud E, Ramphul H, Rycha M, Luximon-Ramma A, Trotin F, Aruoma O. (2003) Phenolic constituents and antioxidant capacities of *Crataegus monogyna* (hawthorn) callus extracts. *Nahrung/Food*, **43**, 191-198.
- [17] Barros L, Carvalho A, Ferreira C. (2010) Comparing the composition and bioactivity of *Crataegus monogyna* flowers and fruits used in folk medicine. *Phytochemical Analysis DOI:10.1002/pca.1267*.

- [18] Konczak-Islam I, Yoshimoto M, Hou D, Terahara N, Yamakawa O. (2003) Potential chemopreventive properties of anthocyanin-rich aqueous extracts from *in vitro* tissue of sweetpotato (*Ipomoea batatas* L.). *Journal of Agricultural and Food Chemistry*, **51**, 5916-5922.
- [19] Luís A, Domingues F, Gil C, Duarte AP. (2009) Antioxidant activity of extracts of Portuguese shrubs: *Pterospartum tridentatum*, *Cytisus scoparius* and *Erica* spp. *Journal of Medicinal Plants Research*, **3**, 886-893.
- [20] Dall'Acqua S, Cervellati R, Loi M, Innocenti G. (2008) Evaluation of *in vitro* antioxidant properties of some traditional Sardinian medicinal plants: Investigation of the high antioxidant capacity of *Rubus ulmifolius*. *Food Chemistry*, **106**, 745-749.
- [21] Öztürk M, Tümen İ, Ugür A, Aydoğmuş-Öztürk F, Topçu G. (2011) Evaluation of fruit extracts of six Turkish *Juniperus* species for their antioxidant, anticholinesterase and antimicrobial activities. *Journal of the Science of Food and Agriculture*, **91**, 867-876.
- [22] Arouma O. (2003) Methodological considerations for characterizing potential antioxidant actions of bioactive components in plant foods. *Fundamental and Molecular Mechanisms of Mutagenesis*, **523-524**, 9-20.
- [23] Alim A, Goze I, Goze H, Tepe B. (2009) *In vitro* antimicrobial and antiviral activities of the essential oil and various extracts of *Salvia cedronella* Boiss. *Journal of Medicinal Plants Research*, **3**, 413-419.
- [24] Andrade D, Gil C, Breitenfeld L, Domingues F, Duarte A. (2009) Bioactive extracts from *Cistus ladanifer* L. and *Arbutus unedo* L. *Industrial Crops and Products*, **30**, 165-167.
- [25] Orhan N, Orhan I, Ergun F. (2011) Insights into cholinesterase inhibitory and antioxidant activities of five *Juniperus* species. *Food and Chemical Toxicology* DOI:10.1016/j.fct.2011.06.031.
- [26] Prakash A, Pereira T, Reilly P, Seawright A. (1999) Pyrrolizidine alkaloids in human diet. *Genetic Toxicology and Environmental Mutagenesis*, **443**, 53-67.
- [27] Çam M, Hışıl Y, Durmaz G. (2009) Classification of eight pomegranate juices based on antioxidant capacity measured by four methods. *Food Chemistry*, **112**, 721 – 726.
- [28] Krishnaiah D, Sarbatly R, Nithyanandam R. (2010) A review of the antioxidant potential of medicinal plant species. *Food Bioprocess Processing*, DOI:10.1016/j.fbp.2010.04.008.
- [29] Tadić V, Dobrić S, Marković G, Đorđević S, Arsić I, Menković N, Stević T. (2008) Anti-inflammatory, gastroprotective, free-radical-scavenging, and antimicrobial activities of hawthorn berries ethanol extract. *Journal of Agricultural and Food Chemistry*, **56**, 7700-7709.
- [30] Faustino H, Gil N, Baptista C, Duarte A. (2010) Antioxidant activity of lignin phenolic compounds extracted from kraft and sulphite black liquors. *Molecules* DOI:10.3390/molecules15129308.
- [31] Sherer R, Godoy H. (2009) Antioxidant activity index (AAI) by the 2,2-diphenyl-1-picrylhydrazyl method. *Food Chemistry*, **112**, 654-658.
- [32] Wang B, Zhang W, Duan X, Li X. (2009) *In vitro* antioxidative activities of extract and semi-purified fractions of the marine red alga, *Rhodomela confervoides* (Rhodomelaceae). *Food Chemistry*, **113**, 1101-1105.
- [33] Fu L, Xu B, Xu X, Qin X, Gan R, Li H. (2010) Antioxidant capacities and total phenolic contents of 56 wild fruits from South China. *Molecules*. DOI:10.3390/molecules15128602.
- [34] Miliuskas G, Venskutonis P, van Beek T. (2004) Screening of radical scavenging activity of some medicinal and aromatic plant extracts. *Food Chemistry*, **85**, 231-237.
- [35] Kelebek H, Selli S, Canbas A, Cabaroğlu T. (2009) HPLC determination of organic acids, sugars, phenolic compositions and antioxidant capacity of orange juice and orange wine made from a Turkish cv. Kozan. *Microchemical Journal*, **91**, 187-192.
- [36] Pourmorad F, Hosseini-mehr S, Shahabimajd N. (2006) Antioxidant activity, phenol and flavonoids contents of some selected Iranian medicinal plants. *African Journal of Biotechnology*, **5**, 1142-1145.
- [37] Tawaha K, Alali F, Gharaibeh M, Mohammad M, El-Elmat T. (2007) Antioxidant activity and total phenolic content of selected Jordanian species. *Food Chemistry*, **104**, 1372-1378.
- [38] Vitti D, Abdalla A, Bueno I, Filho J, Costa C, Bueno M, Nozella E, Longo C, Vieira E, Filho S, Godoy P, Mueller-Harvey I. (2005) Do all tannins have similar nutritional effects? A comparison of three Brazilian fodder legumes. *Animal Feed Science and Technology*, **119**, 345-361.
- [39] Wannes W, Mhamdi B, Sriti J, Jemia M, Ouchick O, Hamdaoui G, Kchouk M, Marzouk B. (2010) Antioxidant activities of the essential oils and methanol extracts from myrtle (*Myrtus communis* var. *italica* L.) leaf, stem and flower. *Food and Chemical Toxicology*, **48**, 1362-1370.
- [40] Rubanza C, Shen M, Otsyina R, Bakengesa S, Ichinohe T, Fujihara T. (2005) Polyphenolics and tannins effect on *in vitro* digestibility of selected *Acacia* species leaves. *Animal Feed Science and Technology*, **119**, 129-142.
- [41] Fernández-Arroyo S, Barrajón-Catalán E, Micol V, Segura-Carretero A, Fernández-Gutiérrez A. (2010) High-performance liquid chromatography with diode array detection coupled to electrospray time-of-flight and ion-trap tandem mass spectrometry to identify phenolic compounds from a *Cistus ladanifer* aqueous extract. *Phytochemical Analysis*, **21**, 307-313.
- [42] Tarnawski M, Depta K, Grejciun D, Szelepin B. (2006) HPLC determination of phenolic acids and antioxidant activity in concentrated peat extract - a natural immunomodulator. *Journal of Pharmaceutical and Biomedical Analysis*, **41**, 182-188.

HPLC/PDA/ESI-MS Evaluation of Saffron (*Crocus sativus* L.) Adulteration

Leonardo Sabatino^{a*}, Monica Scordino^{ab}, Maria Gargano^a, Adalgisa Belligno^b, Pasqualino Traulo^a and Giacomo Gagliano^a

^aMinistero delle Politiche Agricole Alimentari e Forestali, Dipartimento dell'Ispettorato Centrale della Tutela della Qualità e della Repressione Frodi dei Prodotti Agroalimentari (ICQRF) - Laboratorio di Catania, Via A. Volta 19, 95122 Catania, Italy

^bUniversità degli Studi di Catania, Dipartimento di Scienze delle Produzioni Agrarie e Alimentari (DISPA) – Sezione Scienze Agrochimiche, Via S. Sofia 98, 95123 Catania, Italy

l.sabatino@mpaaf.gov.it

Received: July 20th, 2011; Accepted: September 28th, 2011

The present study evaluated the reliability of the ISO/TS 3632-2 UV-Vis spectrometric method for saffron classification, making experiments on saffron samples to which were added increasing concentrations of common saffron spice adulterants (safflower, marigold and turmeric). The results showed that the ISO/TS 3632-2 method is not able to detect addition of up to 10-20%, w/w, of saffron adulterants. For additions from 20 to 50%, w/w, of the three adulterants, saffron was classified in a wrong category; addition of higher than 50%, w/w, determined variations in the investigated parameters that did not allow identification of the product as "saffron". In all cases, the method did not permit the recognition of the nature of the adulterant. On the contrary, the specificity of the HPLC/PDA/MS technique allowed the unequivocal identification of adulterant characteristic marker molecules that could be recognized by the values of absorbance and mass. The selection of characteristic ions of each marker molecule has revealed concentrations of up to 5%, w/w, for safflower and marigold and up to 2% for turmeric. In addition, the high dyeing power of turmeric allowed the determination of 2%, w/w, addition using exclusively the HPLC/PDA technique.

Keywords: spice, saffron, safflower, marigold, turmeric, fraud.

Saffron (*Crocus sativus* L., family Iridaceae) is the most valuable spice in the world. The species is a perennial meadow plant that reaches 10 to 25 cm and grows from bulbs. The cultivation of saffron requires a Mediterranean continental climate with cold winters, warm and dry summers, and a dry Mediterranean humidity regime. The plant is resistant to extreme temperatures in the summer and winter [1a]. Saffron is a spice that adds color, taste and aroma to various foods. The odor is related to its essential oil, a component of which is cineole. The bitter taste of saffron is due to picrocrocin and picrocrozioide, and its color from crocin esters that produce glucose and crocetin after hydrolysis [1a].

The world production of saffron is about 180 tons per year of which 90% is produced in Iran and the remainder in India, Greece, Morocco, Italy, Spain and other countries. The biggest worldwide exporters are Iran, followed by Spain. In Italy the annual production is around 400 kg [1a]. Saffron is cultivated mainly in Sardinia and Abruzzo, with about 35 ha and 7 ha, respectively, and to a lesser extent in Umbria, Tuscany, Liguria and Sicily. Some productions (i.e. Abruzzese di Navelli and San Gimignano) have been awarded the Protected Designation of Origin (PDO). The commercialization costs, including the different process stages, could exceed 1,000 €/kg.

Nowadays saffron production faces a crisis, but all nations involved are traditionally committed to saffron cultivation and preserve it actively. In the Mediterranean basin production of saffron has been decreased due to rising standards of living and, inevitably, due to the rise in labor costs. However, Mediterranean saffron bears the best quality features worldwide, which is attributed to deep cultivation knowledge and careful treatment by all European producers.

Due to its high value, saffron spice has been subjected to many adulterants throughout history, such as mixing of extraneous materials, immersing with vegetable oil or glycerin, and addition of various mineral substances, artificial colorants and less valuable colored spices [1a]. Saffron quality is determined after a series of characteristic parameters for the spice itself (moisture content, flower residue, foreign material, ash content, soluble condensate, coloring power, etc) combined with necessary external conditions (absence of parts from other plants, microbiological flora and pesticide residues). Methods applied for quality assurance are widely known and enterprises are able to use the necessary technology in order to guarantee the product quality to consumers. Since 1980 a standard procedure (ISO/TS 3632) allows the quality for saffron classification. ISO/TS 3632 was updated in 2003 with the

Table 1: Specifications of ISO/TS 3632 "Saffron" [1b].

Characteristics	Specifications Categories			Test method
	I	II	III	
Moisture and volatile matter, % (w/w), max.	12 ^a 10 ^b	12 ^a 10 ^b	12 ^a 10 ^b	ISO/TS 3632-2:2003, Clause 7
Total ash, % (w/w), on dry basis, max.	8	8	8	ISO 928:1997, Clause 8, and ISO/TS 3632-2:2003, Clause 12
Acid-insoluble ash, % (w/w), on dry basis, max.	1	1	1.5	ISO 930:1997, Clause 7, and ISO/TS 3632-2:2003, Clause 13
Bitterness, expressed as direct reading of the absorbance of picrocrocin at about 257 nm, on dry basis, min.	70	55	40	ISO/TS 3632-2:2003, Clause 14
Safranal, expressed as direct reading of the absorbance at about 330 nm, on dry wt basis min.	20	20	20	ISO/TS 3632-2:2003, Clause 14
max.	50	50	50	
Coloring strength, expressed as direct reading of the absorbance of crocine at about 440 nm, on dry wt basis, min.	190	150	100	ISO/TS 3632-2:2003, Clause 14
Artificial water soluble acid colorants	Absent	Absent	Absent	ISO/TS 3632-2:2003, Clause 16 and/or Clause 17

^afilaments; ^b powder.

text that currently governs the product quality [1b]. This regulation is applicable to saffron strands, ground saffron and dust. The rule divides saffron into different categories based primarily on physical-chemical criteria (Table 1).

European saffron is considered the best in the world due to its chemical, physical and organoleptic features, as measured by certain parameters. New high quality verification standards and new evaluation methods should be introduced in order to determine accurately color and to prevent fraud, as reported by many scientific works [1c,2a,2b,3].

The present study evaluated the reliability of the ISO/TS 3632 UV-Vis spectrometric method for saffron classification, making analyses on samples of saffron blended with different concentrations of safflower, marigold and turmeric, widely used as saffron spice adulterants [1a,2a]. The results of the spectrometric method were compared and integrated with the HPLC/PDA/ESI-MS technique for the unequivocal identification of adulterants through the identification of specific marker compounds.

In Table 2 are reported the data related to the different parameters measured according to the ISO/TS 3632 (2003) spectrometric method for saffron category assignment performed on unadulterated and on spice-spiked San Gavino ISO Category II saffron. Five independent additions at different concentrations (10-67%) of adulterants (safflower, marigold and turmeric) were made.

Results showed that the ISO/TS 3632-2 spectrometric method is not able to detect the addition of up to 10-20%, w/w, of saffron adulterants, resulting in a correct

Table 2: Results of ISO/TS 3632 UV-vis spectrophotometry on the analyzed mixes.

Mix composition w/w	E ^{1%} 257 nm	E ^{1%} 330 nm	E ^{1%} 440 nm	ISO Category
Saffron 100%	66	32	170	II
90% Saffron - 10% Turmeric	59	27	150	II
80% Saffron - 20% Turmeric	53	25	132	III
67% Saffron - 33% Turmeric	45	21	111	III
50% Saffron - 50% Turmeric	34	15	81	-
33% Saffron - 67% Turmeric	22	9	49	-
90% Saffron - 10% Safflower	63	31	152	II
80% Saffron - 20% Safflower	59	30	136	III
67% Saffron - 33% Safflower	53	29	113	III
50% Saffron - 50% Safflower	48	27	86	-
33% Saffron - 67% Safflower	42	26	60	-
90% Saffron - 10% Marigold	61	30	151	II
80% Saffron - 20% Marigold	55	29	135	III
67% Saffron - 33% Marigold	49	25	112	III
50% Saffron - 50% Marigold	39	22	83	-
33% Saffron - 67% Marigold	31	18	54	-

E^{1%} absorbency at λ_{max} for a 1% solution of the test sample for a 1cm cell

classification of the mixes as saffron ISO category II. For additions from 20 to 50%, w/w, of the three adulterants studied, the mixes were classified as worse than ISO category III saffron. Spikes higher than 50%, w/w, determined variations of the investigated parameters that did not allow the identification of the mixes as "saffron". In all cases the method did not permit the recognition of the kind of adulterant.

Therefore, the use of only the spectrometric technique may underestimate the saffron fraud occurrence due to the addition of less valuable spices. The hyphenated techniques like high performance liquid chromatography coupled with UV-Vis spectrophotometers and mass spectrometry may allow a better assessment of the quality of the saffron products.

Preliminarily, this study has characterized separately the fingerprint HPLC/PDA/ESI-MS of acidified water-methanol extracts of saffron, marigold, safflower and turmeric. Figure 1 showed the UV-Vis chromatograms of the studied spices. In Table 3 are reported the assignment of the characteristic molecules of each botanical species as a function of retention times, UV-Vis and mass properties.

Crocetin glycosides are responsible for the saffron yellow color; their UV-Vis spectra are characterized by an absorption maximum at about 440-460 nm depending on the molecule. *Trans*- and *cis*-crocetin glycoside showed a different spectroscopic behavior because *cis*-crocetins presented an additional absorption band around 325 nm in their UV-Vis spectrum in comparison with their *trans*-isomers. Six crocetin glycosides (**2-7**), together with colorless picrocrocin [4-(α -D-glucopyranosyl)-2,6,6-trimethyl-1-cyclohexene-1-carboxaldehyde] (**1**), were identified in the analyzed San Gavino saffron. As confirmed by ESI-MS analysis, in agreement with literature data [2b,4], *trans*-crocetin di-(β -D-gentibiosyl) ester (**3**), *cis*-crocetin (β -D-glucosyl)-(β -D-gentibiosyl)




Figure 1: UV-Vis chromatograms of analyzed extracts. **A:** Saffron (λ 250 nm + λ 440 nm); **B:** Marigold (λ 350 nm); **C:** Safflower (λ 410 nm + λ 520 nm); **D:** Turmeric (λ 425 nm). For peak identification see Table 3.

Table 3: HPLC/PDA/MS chemical characterization of studied extracts.

Peak	Name	Rt (min)	UV-vis (nm)	[M-H] ⁺ (m/z)
<i>Crocus sativus</i> L. (saffron)				
1	4-(α -D-glucopyranosyl)-2,6,6-trimethyl-1-cyclohexene-1-carboxaldehyde (picrocrocin)	20.2	250	375
2	<i>trans</i> -Crocetin (β -D-neapolitanosyl)-(β -D-gentibiosyl) ester	41.2	260, 440	1137
3	<i>trans</i> -Crocetin di-(β -D-gentibiosyl) ester	46.5	260, 420, 460	975
4	<i>trans</i> -Crocetin (β -D-glucosyl)-(β -D-neapolitanosyl) ester	50.1	260, 440	975
5	<i>cis</i> -Crocetin (β -D-glucosyl)-(β -D-gentibiosyl) ester	51.6	260, 330, 435, 460	813
6	<i>cis</i> -Crocetin di-(β -D-gentibiosyl) ester	57.5	260, 320, 435, 460	976
7	<i>cis</i> -Crocetin di-(β -D-glucosyl) ester	58.8	260, 325, 440, 465	813
<i>Calendula officinalis</i> (marigold)				
a	Quercetin 3- <i>O</i> -rutinosylrhannoside	28.4	255, 355	755
b	Quercetin 3- <i>O</i> -rutinose	30.6	255, 355	609
c	Isorhamnetin-3- <i>O</i> -rutinosylrhannoside	32.2	255, 350	769
d	Narcissin	34.7	255, 355	623
e	Isorhamnetin 3- <i>O</i> -neohesperidoside	38.7	255, 345	623
f	Isorhamnetin-3- <i>O</i> -glucoside	39.9	255, 355	477
<i>Carthamus tinctorius</i> (safflower)				
I	Hydroxysafflor yellow A (safflorin A)	30.1	225, 410	611
II	6-Hydroxykaempferol 3- <i>O</i> - β -D-glucoside	30.6	275, 340	464
III	Kaempferol 3- <i>O</i> - β -rutinose	37.7	265, 350	593
IV	Safflor yellow B	40.2	225, 410	1060
V	Anhydrosafflor yellow B	42.1	225, 410	1044
VI	Prechartamin	58.7	240, 405	955
VII	Chartamin	68.6	370, 520	909
<i>Curcuma longa</i> (turmeric)				
α	Demethoxycurcumin	63.6	250, 425	337
β	Bisdemethoxycurcumin	63.9	250, 420	307
γ	Curcumin	64.3	260, 430	367

ester (5) and *cis*-crocetin di-(β -D-glucosyl) ester (7) were the most abundant crocetin derivatives, followed by *cis*-crocetin di-(β -D-gentibiosyl) ester (6), *trans*-crocetin (β -D-neapolitanosyl)-(β -D-gentibiosyl) ester (2) and *trans*-crocetin (β -D-glucosyl)-(β -D-neapolitanosyl) ester (4).

Marigold extract was characterized by six main peaks (a-f), which displayed identical UV absorptions with maxima at about 255 and 350 nm, typical of flavonols. The ESI-MS [M-H]⁺ molecular ions, together with comparison with scientific references [5], permitted their unequivocal assignment to quercetin 3-*O*-rutinosylrhannoside (a), quercetin 3-*O*-rutinose (b), isorhamnetin-3-*O*-rutinosylrhannoside (c), narcissin (d), isorhamnetin 3-*O*-neohesperidoside (e), and isorhamnetin-3-*O*-glucoside (f). Among them, the isorhamnetin derivatives predominated in the analyzed sample.

As reported by many scientific papers [6,7], the main component of safflower red pigments is carthamin, composed of two chalconoids with conjugated bonds; it is derived from the yellow colored precarthamin by decarboxylation. Both compounds were detected in the analyzed safflower sample as peaks VI and VII. Safflorin A (I), safflor yellow B (IV) and anhydrosafflor yellow B (V) were identified as the quinochalcone C-glycosides responsible for the yellow color of the sample. Moreover, two kaempferol derivatives were identified as 6-hydroxy-kaempferol 3-*O*- β -D-glucoside (II) and kaempferol 3-*O*- β -rutinose (III). Identification was confirmed by spectroscopic and mass spectral data. Three molecules (α - γ) were identified as responsible for the yellow distinctive color of turmeric: all of them clearly possess a maximum absorption wavelength near 420 nm. According to literature data [8], these compounds were identified as the dicinnamoylmethane derivatives demethoxycurcumin (α), bisdemethoxycurcumin (β) and curcumin (γ) on the basis of [M-H]⁺ molecular ions generated by the ESI-MS negative soft ionization. Subsequently, the above-mentioned analytical technique was applied to saffron samples mixed with different concentrations of turmeric, marigold and safflower in the range 2-20%, w/w. For each adulterant, marker molecules have been chosen for their unambiguous identification in the mixture: it was established that their detection was not influenced by the saffron matrix effect.

Extraction of the ion with m/z 623, corresponding to isorhamnetin 3-*O*-neohesperidoside (e), is able to detect the presence of marigold. In the case of safflower, the marker molecules for its identification in the mixture were anhydrosafflor yellow B (V) and chartamin (VII) with ions at m/z 1044 and 909, respectively. The marker molecules could be revealed at concentrations of up to 5%, w/w, of both marigold and safflower. The characteristic turmeric curcuminoids triplet due to the presence of demethoxycurcumin (α), bisdemethoxy-curcumin (β) and curcumin (γ) could easily identify its presence in the mix with saffron, and also at concentrations of 2%, even using only the UV-Vis detector. The above discussed method is currently applied by the Catania Laboratory of Central Inspectorate for Quality Control of Agricultural and Food Productions (ICQRF) on samples of saffron collected in the Italian market in the framework of the Ministry of Agriculture and Forestry institutional quality control activity.

In conclusion, the results of the present study should force the legislative authorities to release new standards for the saffron sector for the maintenance of product purity in order to avoid adulteration and fraud. The methods applied nowadays are outdated, while newer ones are not positively accepted and are rarely used, even though some of them have proven to be efficient in the field. All national and international quality control standards should be reinforced in order to limit the spread of adulterated saffron in the European market deriving from countries that affect considerably the competitiveness of European saffron.

Experimental

Plant material, reagents and standards: Strands of saffron (*Crocus sativus* L.) from San Gavino (Sardinia, Italy), turmeric (*Curcuma longa*) powder, calendula (*Calendula officinalis*) flowers, and safflower (*Carthamus tinctorius*) strands were taken in Italy by ICQRF officials. The plant material was treated in accordance with the specifications of ISO/TS 3632-2:2003 sample preparation for the official analysis [1b]. Different mixes were prepared with saffron with percentages varying from 2 to 70% by weight of each adulterant studied. HPLC-grade acetonitrile, methanol and formic acid were supplied by Romil (Milan, Italy). Distilled water was purified at 18.2 MΩ cm with a MilliQ ULTRA (Millipore, Vimodrone (MI), Italy) purification system.

UV-Vis spectrophotometry: The method used was based on the technical specification ISO/TS 3632-2:2003 [1b] and allowed determination of the main characteristics of saffron. The method is based on the spectrophotometric assessment absorption at 3 wavelengths (λ_{max}): 257 nm (maximum

absorption of picrocrocin), 330 nm (maximum absorption of safranal) and 440 nm (maximum absorption of crocin).

HPLC/PDA/ESI-MS fingerprint: 20±2 mg of sample was extracted with 2 mL of 50% aqueous methanol containing 0.1% formic acid at room temperature in an ultrasonic bath for 30 min. The mixture was centrifuged at 3000 rpm for 5 min and the supernatant decanted and used for the LC analyses, after filtration through 0.45 µm PTFE filters (LabService Analytica, Bologna, Italy). The analyses were performed with a liquid chromatograph consisting of a Finnigan Surveyor MS-pump, autosampler and photodiode-array detector (PDA), coupled with a Finnigan LCQ DECA XP MAX detector (Thermo Fisher Scientific). The analytical column was a Gemini C18 150 x 2.1 mm i.d. 3µm (Phenomenex); the flow rate was 200 µL/min, the column temperature 30°C and the injection volume 10 µL. A binary gradient of 0.3% formic acid in water (A) and 0.3% formic acid in acetonitrile (B) was employed. The mobile phase gradient was programmed as follows: 0 min, 5% B; 50 min, 28% B; 60 min, 43% B; 60-65 min, 43 % B; 70-80 min, 5% B. The range of wavelengths examined by the photodiode-array detector was 200–700 nm and the mass scan range was 100-1600 m/z. Mass spectral analyses were performed using a LCQ ion-trap mass operating in negative ion mode using an ion spray LC/MS interface. The electrospray ionization (ESI) needle voltage was 3.5 kV. The capillary voltage was 18V and the heated capillary was 250°C. A sheath gas flow rate of 36 (arbitrary units) was used and the auxiliary gas was set to 14 (arbitrary units). The main compounds of analyzed plant materials were characterized in terms of retention times, lambda max and MS data.

References

- [1] (a) Kafi M, Koocheki A, Rashid MH, Nassiri N. (2006) *Saffron (Crocus sativus) production and processing*. Science Publishers, Enfield, NH, USA; (b) ISO/TS 3632. (2003) *Saffron (Crocus sativus L.)*. Part 1 (specification) and Part 2 (test methods). International Organization for Standardization, Genève, Switzerland; (c) Alonzo GL, Salinas MR, Garijo J. (1998) Method to determine the authenticity of aroma of saffron. *Journal of Food Protection*, **61**, 1525-1528.
- [2] (a) Haghighi B, Feizy J, Kakhki AH. (2007) LC determination of adulterated saffron prepared by adding styles colored with some natural colorants. *Chromatographia*, **66**, 325–332; (b) Zalacain A, Ordoudi SA, Díaz-Plaza EM, Carmona M, Blázquez I, Tsimidou MZ, Alonso GL. (2005) Near-infrared spectroscopy in saffron quality control: determination of chemical composition and geographical origin. *Journal of Agricultural and Food Chemistry*, **53**, 9337-9341.
- [3] Sujata V, Ravishankar GA, Venkataraman LV. (1992) Methods for the analysis of the saffron metabolites crocin, crocetins, picrocrocin and safranal for the determination of the quality of the spice using thin-layer chromatography, high-performance liquid chromatography and gas chromatography. *Journal of Chromatography A*, **624**, 497-502.
- [4] Carmona M, Zalacain A, Sánchez AM, Novella JL, Alonso GL. (2006) Crocetin esters, picrocrocin and its related compounds present in *Crocus sativus* stigmas and *Gardenia jasminoides* fruits. Tentative identification of seven new compounds by LC-ESI-MS. *Journal of Agricultural and Food Chemistry*, **54**, 973-979.
- [5] Bilia AR, Bergonzi MC, Gallori S, Mazzi G, Vincieri FF. (2002) Stability of the constituents of Calendula, milk-thistle and passion flower tinctures by LC-DAD and LC-MS. *Journal of Pharmaceutical and Biomedical Analysis*, **30**, 613-624.
- [6] Kazuma K, Takahashi T, Sato K, Takeuchi H, Matsumoto T, Okuno T. (2000) Quinochalcones and flavonoids from fresh florets in different cultivars of *Carthamus tinctorius* L. *Bioscience, Biochemistry, and Agrochemistry*, **64**, 1588-1599.
- [7] Jin Y, Zhang XL, Shi H, Xiao YS, Ke YX, Xue XY, Zhang FF, Liang XM. (2008) Characterization of C-glycosyl quinochalcones in *Carthamus tinctorius* L. by ultraperformance liquid chromatography coupled with quadrupole-time-of-flight mass spectrometry. *Rapid Communications in Mass Spectrometry*, **22**, 1275-1287.
- [8] He XG, Lin LZ, Lian LZ, Lindenmaier M. (1998) Liquid chromatography-electrospray mass spectrometric analysis of curcuminoids and sesquiterpenoids in turmeric (*Curcuma longa*). *Journal of Chromatography A*, **818**, 127-132.

Ferric Reducing, Antiradical and β -Carotene Bleaching Activities of Nicotinic Acid and Picolinic Acid Bioconjugates of Curcumin

Archana Pandey^a, Kanti Bhooshan Pandey^b, Ravindra Kumar Gupta^a and Syed Ibrahim Rizvi^{b*}

^aDepartment of Chemistry, C.M.P College, Allahabad, Allahabad-211002, India

^bDepartment of Biochemistry, University of Allahabad, Allahabad-211002, India

sirizvi@gmail.com

Received: June 4th, 2011; Accepted: October 6th, 2011

Although curcumin displays several beneficial properties, its medicinal use is limited by its low bioavailability. In the present study we report the antioxidant potentials of two bioconjugates of curcumin with nicotinic acid and picolinic acid: di-*O*-nicotinoyl curcumin [1,7-bis (4-*O*-nicotinoyl-3-methoxyphenyl)-1,6-heptadiene-3, 5-dione] and di-*O*-picolinoyl curcumin [1,7-bis (4-*O*-picolinoyl-3-methoxyphenyl)-1,6-heptadiene-3, 5-dione], in terms of ferric reducing, radical scavenging and β -carotene bleaching abilities, and comparing the observed activity with that of curcumin. Results demonstrate that both the bioconjugates possess higher antioxidant potentials as evidenced by enhanced ferric reducing, radical scavenging and β -carotene bleaching abilities, in comparison with curcumin. On the basis of our results we conclude that these bioconjugates of curcumin may be better than curcumin for medicinal and pharmacological applications.

Keywords: Curcumin, nicotinic acid, picolinic acid, bioconjugates, antioxidant.

Curcumin, 1,7-bis-(4-hydroxy-3-methoxyphenyl)-1,6-heptadiene-3,5-dione is a natural yellow pigment isolated from turmeric (*Curcuma longa*). Curcumin is known to exhibit anticancer, anti-inflammatory, anti-diabetic, anti-viral and neuroprotective properties [1,2]. The compound is also reported to reduce blood cholesterol, enhance wound healing, protect from liver injuries and cataract formation [3,4]. Although multiple mechanism(s) of action have been reported for curcumin, the antioxidant property associated with this molecule has been a subject of great attention and many of the associated biological and pharmacological effects attributed to it [4,5].

Pleiotropic health beneficial properties make curcumin an important pharmacologically active molecule, but low bioavailability, poor absorption and rapid metabolism in the liver and intestinal wall limits its use as a therapeutic agent [6,7]. Its highly hydrophobic nature also reduces systemic administration, and it has been shown that on intravenous administration, curcumin disappears rapidly from the blood and quickly appears as metabolites in the bile [7,8]. In order to overcome these limitations, curcumin may attach to ligands that can enhance its solubility, bioavailability, slowdown the metabolism and increase the rate of cellular uptake [9]. Keeping this rationale in mind, two bioconjugates of curcumin have been synthesized comprised of nicotinic acid (vitamin B3) and its isomer picolinic acid: di-*O*-nicotinoyl curcumin [1,7-bis (4-*O*-nicotinoyl-3-methoxyphenyl)-1,6-heptadiene-3,5-dione] (DNC) and di-*O*-picolinoyl curcumin [1,7-bis (4-*O*-picolinoyl-3-methoxyphenyl)-1,6-heptadiene-3,5-dione]




Figure 1: Chemical structures of curcumin [1, 7-bis-(4-hydroxy-3-methoxyphenyl)-1,6-heptadiene-3,5-dione], di-*O*-nicotinoyl curcumin [1,7-bis (4-*O*-nicotinoyl-3-methoxyphenyl)-1,6-heptadiene-3,5-dione], di-*O*-picolinoyl curcumin [1,7-bis (4-*O*-picolinoyl-3-methoxyphenyl)-1,6-heptadiene-3,5-dione].

(DPC, Figure 1)]. In the present study these bioconjugates have been screened for their antioxidant potential in terms of ferric reducing, radical scavenging and β -carotene bleaching activities in an effort to assess their efficacy as antioxidant moieties compared with the parent molecule i.e. curcumin.

Both the synthesized bioconjugates, DNC and DPC, showed enhanced ferric reducing ability in comparison with curcumin itself ($p < 0.05$), when assayed with FRAP reagent. The ability of DNC and DPC to reduce the ferric ion was 43% and 45% higher, respectively compared with curcumin (Figure 2). Similar to the ferric reducing ability, both the bioconjugates of curcumin inhibited DPPH \cdot at a higher rate than curcumin ($p < 0.001$). Interestingly, DPC

showed a slightly higher radical scavenging ability/antiradical power than DNC. The ability of DNC and DPC to inhibit DPPH[•] was 42% and 51%, respectively higher than curcumin (Figure 3). Both the bioconjugates showed strong β -carotene bleaching capacity when tested, in comparison with curcumin. DNC elicited 20% and DPC 23% more bleaching capacity than curcumin (Figure 4).

The FRAP assay, based on ferric reducing ability of biological samples, is one of the most frequently used methods to evaluate antioxidant activity since it is independent of the enzymatic/non-enzymatic method to generate free radicals prior to evaluating the anti-radical activity of antioxidants. The ability to reduce the ferric ion at higher rates by both the bioconjugates of curcumin, confirms their higher antioxidant potentials compared with curcumin.




Figure 2: Comparative FRAP values of assayed bioconjugates of curcumin at a concentration of 10 μ M. Values are mean \pm SD of 10–12 independent experiments. PC- Curcumin; DNC-di-O-nicotinoyl curcumin; DPC- di-O-picolinoyl curcumin. * $p < 0.05$ and ** $p < 0.001$, compared with curcumin.

The estimation of radical scavenging activity of curcumin along with its bioconjugates using the DPPH[•] method presented the advantage of using a stable and commercially available free radical. This method has been extensively applied to the study of antiradical power of biological compounds [10,11]. Easy to perform and high reproducibility adds extra advantage over other methods. Inhibition of DPPH[•] by DNC and DPC at higher rates than curcumin supports the results obtained from the FRAP assay.

The β -carotene bleaching assay is a reproducible and authentic method to measure the antioxidant capacity of different biological samples. The enhanced capacity of the synthesized bioconjugates to bleach β -carotene in comparison with curcumin confirms their greater antioxidant potential.

The presence of the carboxyl side chain at the *ortho* position and intramolecular coordination of the carboxylic group and nitrogen of the pyridine nuclei facilitates the proton donating ability of DPC. This may be the reason for the slightly higher DPPH[•] scavenging and β carotene bleaching activity of DPC in comparison with DNC.

Chemically, curcumin is a bis- α,β -unsaturated β -diketone (commonly called diferuloylmethane, Figure1), which




Figure 3: Comparative % DPPH[•] reducing activity of curcumin bioconjugates at the concentration of 10 μ M. Values are mean \pm SD of 10–12 independent experiments. PC- Curcumin; DNC-di-O-nicotinoyl curcumin; DPC- di-O-picolinoyl curcumin. * $p < 0.001$ and ** $p < 0.0005$, compared with curcumin.




Figure 4: Comparative % β -carotene bleaching capacity of curcumin bioconjugates at the concentration of 10 μ M. Values are mean \pm SD of 10–12 independent experiments. PC- Pure curcumin; DNC-di-O-nicotinoyl curcumin; DPC- di-O-picolinoyl curcumin. * $p < 0.05$ and ** $p < 0.001$, compared with pure curcumin.

exhibits keto–enol tautomerism, having a predominant keto form in acidic and neutral solutions and a stable enol form in alkaline media. The double bonds in curcumin provide definite conformational flexibility to the molecule, which accounts for its various properties [12,13].

The inherent disadvantage with respect to curcumin is its poor solubility in aqueous systems, which makes its use in water-based food products difficult. The alkyl and aryl portions of this molecule render it lipophilic and thus restrict its uptake into cells [14,15]. Antioxidant properties of a number of ring-substituted analogues of curcumin show that the highest antioxidant activity is obtained when the phenolic group is sterically hindered [15]. Therefore, one of the most practical approaches is to make biodegradable conjugates of curcumin with suitable ligands to enhance its cellular uptake [16]. The site of attachment of ligands to the curcumin molecule can be at phenolic hydroxyls, ketogroups (enol functions) or the active methylene group.

The masking of phenolic groups with ligands such as amino acids, nucleosides, and peptides has already been reported [7,17,18]. We are the first to select vitamin B3-complex as a ligand. The bioconjugates: DNC and DPC prepared with nicotinic acid and its isomer picolinic acid

have a biodegradable ester linkage. Since both ligands used are natural and are essential components which play a vital role in cellular physiology, the bioconjugates synthesized are predicted to enhance cellular uptake and be more efficacious than pure curcumin.

The vitamin derivatives of curcumin showed higher ferric reducing, free radical scavenging and β -carotene bleaching activities than pure curcumin, thus demonstrating that derivatization at the phenolic position of curcumin improved *in vitro* antioxidant activity. Results of our study on screening of antioxidant potential of curcumin bioconjugates may be useful in developing more efficacious conjugates of curcumin with enhanced bioavailability incorporating better pharmacological and biological effects.

Experimental

Synthesis of di-O-nicotinoyl curcumin [1,7-bis (4-O-nicotinoyl-3-methoxyphenyl)-1, 6-heptadiene-3, 5-dione]: Redistilled thionyl chloride (1.09 mL, 0.015 mol) was added to 1.23 g of nicotinic acid (0.01 mol) and refluxed for 30 min. The reaction mixture was evaporated to dryness on a rotary evaporator under reduced pressure. [yield = 72% (0.890 g, 6 mmol)]. One hundred mg curcumin (0.27 mmol) in 5 mL dimethyl formamide (DMF) was mixed with nicotinic chloride (90 mg \approx 0.6 mmol). A few drops of triethylamine were added as catalyst and the reaction mixture was stirred overnight at room temperature. The completion of reaction was checked by TLC. The reaction mixture was then poured into chilled water (10 mL), shaken well, and then extracted with DCM (dichloromethane) thrice (5 mL each time). Na_2SO_4 (1g) was added to the extracted DCM, the solution kept overnight, then filtered, the filtrate evaporated to dryness, and then purified by silica gel CC using DCM/MeOH gradient.

Yield: 42%.

Rf: 0.81 (DCM/MeOH; 9.8: 0.2).

UV λ_{max} (MeOH): 260 (s), 290,320 (s) nm.

^1H NMR (DMSO- d_6 and D_2O): δ 3.75 (s, 6H, two- OCH_3), 4.52 (s, 2H, C_4 of cur), 6.53 (d, 2H, C_2 and C_6 of cur), 6.80 - 6.95 (m, 6H, Ar-cur), 6.99 (m, 2H, C_5 of pyridine ring), 7.00 (d, 2H, C_4 of pyridine ring), 7.52 (d, 2H, C_1 and C_7 of cur), 8.20 (d, 2H, C_6 of pyridine ring), 8.50 (s, 2H, C_2 of pyridine ring).

Elemental analysis: observed C, 68.49; H, 4.51; N, 4.89% calculated for $\text{C}_{33}\text{H}_{26}\text{O}_8\text{N}_2$: C, 68.51; H, 4.49; N, 4.84%.

Synthesis of di-O-picolinoyl curcumin [1,7-bis (4-O-picolinoyl-3-methoxyphenyl)-1, 6-heptadiene-3, 5-dione]: Di-O-picolinoyl curcumin was prepared similarly to di-O-nicotinoyl curcumin, but by replacing nicotinic chloride by picolinic chloride.

Rf: 0.80 (DCM/MeOH; 9.8: 0.2).

UV λ_{max} (MeOH): 260 (s), 292,326 nm.

^1H NMR (DMSO- d_6 and D_2O): δ 3.72 (s, 6H, two OCH_3), 4.58 (s, 2H, C_4 of cur), 6.67 (d, 2H, C_2 and C_6 of cur), 6.78

- 6.91 (m, 6H, Ar-cur), 6.80 (m, 2H, C_4 of pyridine ring), 6.99 (m, 2H, C_5 of pyridine ring), 7.00 (d, 2H, C_3 of pyridine ring), 7.54 (d, 2H, C_1 and C_7 of cur), 8.20 (d, 2H, C_6 of pyridine ring).

Elemental analysis: observed C, 68.50; H, 4.48; N, 4.86% calculated for $\text{C}_{33}\text{H}_{26}\text{O}_8\text{N}_2$: C, 68.51; H, 4.49; N, 4.84%.

Ferric reducing antioxidant potential (FRAP) assay: The antioxidant capacity of the bioconjugates and curcumin was determined by the ferric reducing antioxidant potential (FRAP) assay, following the method of Benzie and Strain [19]. Working FRAP reagent was prepared by mixing acetate buffer (300 mM, pH 3.6), 2, 4, 6-tri [2-pyridyl]-s-triazine (10 mM in 40 mM HCl) solution and $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ (20 mmol/L) solution in 10:1:1 ratio, respectively. Three mL of FRAP reagent was mixed with 10 μM of each sample solution and the content mixed vigorously. The absorbance was read at 593 nm at intervals of 30 sec for 4 min. An aqueous solution of Fe^{2+} in the range of 100 -1000 $\mu\text{mol/L}$ was used for calibration. Using the regression equation, the FRAP values ($\mu\text{mol Fe (II)/L}$) of the compounds were calculated.

1,1-Diphenyl-2-picrylhydrazyl free radical (DPPH $^\bullet$) assay: Anti-radical activity of curcumin and both the bioconjugates was estimated by the procedure described by Miliauskas *et al.*[20]. Briefly, 0.1 mL of each compound was incubated in a methanolic solution of DPPH $^\bullet$ (0.1 mM). Absorbance at 517 nm was measured after 30 min of incubation with vigorous shaking. Methanol was used as blank reference. All the measurements were performed in triplicate. The free radical DPPH $^\bullet$ scavenging (i.e. reduction) activity was calculated from the equation: Activity [% of DPPH $^\bullet$ reduction] = $[(A - A_x)/A] \times 100\%$, where A – absorbance of DPPH $^\bullet$ solution with methanol, A_x - absorbance of a DPPH $^\bullet$ solution with assayed compounds.

β -Carotene bleaching assay: β -Carotene bleaching capacity of the synthesized bioconjugates was determined according to the method of Burda and Oleszek [21]. Briefly, 1 mL of β -carotene (0.2 mg/mL) was dissolved in chloroform containing 0.02 mL linoleic acid and 0.02 mL Tween 20. The mixture was then dosed with 0.2 mL of compound solution (final concentration 10 μM). Fifty mL of distilled water, saturated for 15 min with oxygen, was added to the flask. The resulting mixture was shaken and kept for 2 h at 50°C. The absorbance of the samples was measured at 470 nm, immediately after their preparation ($t = 0$ min) and at the end of the experiment ($t = 120$ min). Antioxidant activity was calculated as percent inhibition of oxidation versus control, using the equation: % β -carotene bleaching = $100 [1 - (A_s^0 - A_s^t) / (A_c^0 - A_c^t)]$ where A_s^0 : absorbance of sample at 0 min, A_s^t : absorbance of the sample at 120 min, A_c^0 : absorbance of control at 0 min and A_c^t : absorbance of control at 120 min.

Statistical analyses were performed using GraphPad Prism version 4.00 for Windows, GraphPad Software, San Diego California USA. The results are reported as means \pm SD.

Acknowledgments - The authors AP and SIR wish to acknowledge University Grants Commission, New Delhi for financial support in the form of MRP: F-38-38/2009 SR and MRP: F-37-392/2009 SR.

References

- [1] Pandey A, Gupta RK, Srivastava R. (2011) Curcumin- The yellow magic. *Asian Journal of Applied Science*, **4**, 343-354.
- [2] Zandi K, Ramedani E, Mohammadi K, Tajbakhsh S, Deilami I, Rastian Z, Fouladvand M, Yousefi F, Farshadpour F. (2011) Evaluation of antiviral activities of curcumin derivatives against HSV-1 in Vero cell line. *Natural Product Communications*, **12**, 1935-1938.
- [3] Goel A, Kunnumakkara AB, Aggarwal BB. (2008) Curcumin as "Curecumin": from kitchen to clinic. *Biochemical Pharmacology*, **75**, 787-809.
- [4] Ali BH, Marif H, Noureldayem SA, Bakheit AO, Blunden G. (2006) Some biological properties of curcumin: A review. *Natural Product Communications*, **1**, 509-521.
- [5] Zhou H, Beevers CS, Huang S. (2011) The targets of curcumin. *Current Drug Targets*, **12**, 332-347.
- [6] Anand P, Kunnumakkara AB, Newman RA, Aggarwal BB. (2007) Bioavailability of curcumin: problems and promises. *Molecular Pharmacology*, **4**, 807-818.
- [7] Singh RK, Rai D, Yadav D, Bhargava A, Balzarini J, De Clercq E. (2010) Synthesis, antibacterial and antiviral properties of curcumin bioconjugates bearing dipeptide, fatty acids and folic acid. *European Journal of Medicinal Chemistry*, **45**, 1078-1086.
- [8] Priyadarsini KI, Maity DK, Naik GH, Kumar MS, Unnikrishnan MK, Satav JG, Mohan H. (2003) Role of phenolic O-H and methylene hydrogen on the free radical reactions and antioxidant activity of curcumin. *Free Radical Biology & Medicine*, **35**, 475-484.
- [9] Pandey A, Dwivedi V, Misra K. (2010) Drug sensitivity of curcumin analogues and bioconjugates. In *Structural modification and drug development*. Vol 7, Gupta VK, Taneja SC, Gupta BD. (Eds). Studium Press, LLC US. 267-298.
- [10] Sánchez-Moreno C, Larrauri JA, Saura-Calixto F. (1998) A procedure to measure the antiradical efficiency of polyphenols. *Journal of the Science of Food and Agriculture*, **76**, 270-276.
- [11] Villano D, Fernández-Pachón MS, Moyá ML, Troncoso AM, García-Parrilla MC. (2007) Radical scavenging ability of polyphenolic compounds towards DPPH free radical. *Talanta*, **71**, 230-235.
- [12] Weber WM, Hunsaker LA, Abcouwer SF, Deck LM, Vander Jagt DL. (2005) Anti-oxidant activities of curcumin and related enones. *Bioorganic & Medicinal Chemistry*, **13**, 3811-3820.
- [13] Barclay LR, Vinqvist MR, Mukai K, Goto H, Hashimoto Y, Tokunaga A, Uno H. (2000) On the antioxidant mechanism of curcumin: classical methods are needed to determine antioxidant mechanism and activity. *Organic Letters*, **2**, 2841-2843.
- [14] Kumar S, Dubey KK, Tripathi S, Fujii M, Misra K. (2000) Design and synthesis of curcumin-bioconjugates to improve systemic delivery. *Nucleic Acids Symposium Series*, **44**, 75-76.
- [15] Parvathy KS, Negi PS, Srinivas P. (2010) Curcumin–amino acid conjugates: Synthesis, antioxidant and antimutagenic attributes. *Food Chemistry*, **120**, 523–530.
- [16] Kumar S, Misra A, Tripathi S, Misra K. (2001) Study on curcumin-oligonucleotide conjugate as a probable anticancer agent: its hybridisation with telomere target sequence 5'-GGGATTGGGATT-3'. *Nucleic Acids Research Supplement*, **1**, 137-138.
- [17] Elias EJ, Anil S, Ahmad S, Daud A. (2010) Colon targeted curcumin delivery using guar gum. *Natural Product Communications*, **6**, 915-918.
- [18] Dubey SK, Sharma AK, Narain U, Misra K, Pati U. (2008) Design, synthesis and characterization of some bioactive conjugates of curcumin with glycine, glutamic acid, valine and demethylenated piperic acid and study of their antimicrobial and antiproliferative properties. *European Journal of Medicinal Chemistry*, **43**, 1837-1846.
- [19] Benzie IF, Strain JJ. (1996) The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. *Analytical Biochemistry*, **239**, 70-76.
- [20] Miliauskas G, Venskutonis PR, van Beek TA. (2004) Screening of radical scavenging activity of some medicinal and aromatic plant extracts. *Food Chemistry*, **85**, 231–237.
- [21] Burda S, Oleszek W. (2001) Antioxidant and antiradical activities of flavonoids. *Journal of Agricultural and Food Chemistry*, **49**, 2774-2779.

Antiviral Activities of Diarylheptanoids Isolated from *Alpinia officinarum* against Respiratory Syncytial Virus, Poliovirus, Measles Virus, and Herpes Simplex Virus Type 1 *in vitro*

Katsuhiko Konno^a, Rie Sawamura^a, Yi Sun^b, Ken Yasukawa^b, Tomomi Shimizu^a,
Wataru Watanabe^a, Masahiko Kato^a, Ryuichi Yamamoto^a and Masahiko Kurokawa^{a,*}

^aSchool of Pharmaceutical Sciences, Kyushu University of Health and Welfare, 1714-1 Yoshino,
Nobeoka, Miyazaki 882-8508, Japan

^bSchool of Pharmacy, Nihon University, 7-7-1 Narashinodai, Funabashi-shi, Chiba 274-8555, Japan

b2mk@phoenix.ac.jp

Received: August 16th, 2011; Accepted: October 15th, 2011

Alpinia officinarum has been used as a folk medicine and contains diarylheptanoids that have various biological activities. However, their antiviral activities are less elucidated. We examined the antiviral activities of nine diarylheptanoids isolated from *A. officinarum* against respiratory syncytial virus (RSV), poliovirus, measles virus, and herpes simplex virus type 1 (HSV-1) using a plaque reduction assay. The 50% inhibitory concentrations of seven of the nine diarylheptanoids for RSV were moderately but significantly lower than their 50% cytotoxic concentrations, as determined by a trypan blue exclusion assay. Four diarylheptanoids with anti-RSV activity also showed anti-poliovirus and anti-measles virus activities and three of the four exhibited anti-HSV-1 activity. Thus, seven of the nine diarylheptanoids examined exhibited potential antiviral activity against RSV, and most of the diarylheptanoids with anti-RSV activity, including two diarylheptanoids without anti-RSV activity, were effective against poliovirus, measles virus, and/or HSV-1 *in vitro*. Diarylheptanoids were suggested to have a broad spectrum of antiviral activity.

Keywords: diarylheptanoids, antiviral activity, RSV, HSV-1, measles virus, poliovirus.

Alpinia officinarum (*A. officinarum*), family Zingiberaceae, is known as lesser galangal. This rhizome has been used in various Asian cuisines and as a traditional medicine, such as an antiemetic, stomachic, and analgesic in Asia from ancient times.

Respiratory syncytial virus (RSV) infection is very common in children less than 2 years old and sometimes causes serious bronchitis and pneumonia [1]. In elderly and high-risk adults, RSV infection is an important illness [2]. Ribavirin, palivizumab, and motavizumab are used for the treatment and prevention of RSV infection [3–6], but there are few clinically specific and effective anti-RSV drugs.

In a series of studies on the development of bioactive components from natural sources, we found that a methanol extract from the rhizome of *A. officinarum* is effective in inhibiting tumor promotion by 12-*O*-tetradecanoylphorbol-13-acetate in mouse skin [7]. An extract of *A. officinarum* was previously shown to exhibit therapeutic efficacy against herpes simplex virus type 1 (HSV-1) infection in mice [8]. Diarylheptanoids isolated from *A. officinarum* have been shown to exhibit cytotoxic

activity [9], suppressive activity of inducible nitric oxide synthase expression [10], inhibitory activity of biosynthesis of prostaglandin and leukotrienes [11,12], and inhibitory activity of proinflammatory mediators [13]. Although a variety of biological activities associated with diarylheptanoids have been demonstrated, antiviral activity of diarylheptanoids has been reported only against influenza virus [14,15]. In the present study, we examined the potential anti-RSV activity of diarylheptanoids *in vitro*. Their anti-RSV activities were compared with antiviral activities against poliovirus, measles virus, and HSV-1 to characterize the anti-RSV activity.

Diarylheptanoids (**AO-1** to **9**, Figure 1 and Table 1) were examined for their anti-RSV activity and cytotoxicity *in vitro*. As shown in Table 1, the EC₅₀ values of seven diarylheptanoids (**AO-1**, **2**, **4**, **5**, and **7-9**) were significantly lower than their CC₅₀ values. DMSO at 1%, the maximum concentration used to dissolve diarylheptanoids in the culture medium, was not cytotoxic. The therapeutic indexes (CC₅₀/EC₅₀) of 7-(4'-hydroxyphenyl)-1-phenyl-4*E*-hepten-3-one (**AO-2**) and (5*S*)-5-methoxy-1,7-diphenyl-3-heptanone (**AO-7**) were 4.6 and more than 6.1, respectively, and RSV was more




Figure 1: Structures of diarylheptanoids (AO-1 to 9) from *A. officinarum*.

susceptible to **AO-2** and **AO-7** than to the other diarylheptanoids examined. In this assay, the EC₅₀ value of ribavirin, used as a control, was similar to the results reported previously [16,17]. Thus, seven diarylheptanoids (**AO-1**, **2**, **4**, **5**, and **7-9**) were demonstrated to show moderate, but potential antiviral activity against RSV *in vitro*. This is the first evidence demonstrating the anti-RSV activity of diarylheptanoids *in vitro*.

To evaluate the antiviral spectrum of diarylheptanoids with anti-RSV activity, four diarylheptanoids (**AO-1**, **4**, **5**, and **7**) with anti-RSV activity were examined for anti-poliovirus, -measles virus, and -HSV-1 activities. However, three diarylheptanoids (**AO-2**, **8**, and **9**) with anti-RSV activity were not used, because there were not sufficient amounts to perform a plaque reduction assay. As shown in Table 2, the EC₅₀ values of four diarylheptanoids (**AO-1**, **4**, **5**, and **7**) for poliovirus and measles virus were significantly lower than their CC₅₀ values. The EC₅₀ values of three (**AO-4**, **5**, and **7**) of the four for HSV-1 were also significantly lower than their CC₅₀ values. Therefore, among the four diarylheptanoids, three (**AO-4**, **5**, and **7**) with anti-RSV activity exhibited anti-poliovirus, -measles virus, and -HSV-1 activities. **AO-1** exhibited anti-poliovirus and -measles virus activities, but not anti-HSV-1 activity. Because the three diarylheptanoids (**AO-4**, **5**, and **7**) showed antiviral activity against all viruses used in this

study, they were suggested to have broad spectrum antiviral activity.

We also examined anti-poliovirus, -measles virus, and -HSV-1 activities of **AO-3** and **6** that did not exhibit anti-RSV activity *in vitro*. As shown in Table 2, **AO-3** was significantly effective for measles virus, but not for poliovirus and HSV-1. However, all three viruses examined were significantly susceptible to **AO-6**. Only measles virus was susceptible to all of the six diarylheptanoids (**AO-1**, **3**, **4**, **5**, **6**, and **7**) without relation to anti-RSV activity. Diarylheptanoids without anti-RSV activity were also effective against poliovirus, measles virus, and/or HSV-1 *in vitro* and the broad spectrum of antiviral activity was confirmed.

RSV has a different virus structure and replication cycle from poliovirus and HSV-1. However, it has a similar virus structure and replication cycle to measles virus as some paramyxoviruses. In Table 2, six diarylheptanoids (**AO-1**, **3**, **4**, **5**, **6**, and **7**) exhibited anti-measles virus activity. However, two (**AO-3** and **6**) of them had no anti-RSV activity (Table 1). It is possible that **AO-3** and **6** interfered with a replication step specific to measles virus but not RSV in paramyxoviruses. Although we focused on diarylheptanoids with anti-RSV activity in this study, **AO-3** and **6** were suggested to be potent candidates as anti-measles virus compounds. In our screening of anti-RSV activity *in vitro*, the CC₅₀/EC₅₀ value (>6.1) of **AO-7** was highest (Table 1). **AO-7** also exhibited anti-poliovirus, -measles virus, and -HSV-1 activities (Table 2) and may be characterized as a candidate for an anti-RSV compound with a broad antiviral spectrum. Studies of the structure-antiviral activity relationships of many diarylheptanoids isolated from *A. officinarum* [18–20] against various kinds of viruses may be worthwhile to analyze the antiviral actions and to obtain more effective antiviral diarylheptanoids.

RSV was significantly susceptible to seven of the nine diarylheptanoids isolated from *A. officinarum*. Of the nine, six (**AO-1**, **3**, **4**, **5**, **6**, and **7**) with or without anti-RSV activity were effective against poliovirus, measles virus, and/or HSV-1. Thus, diarylheptanoids were suggested to possess a broad spectrum of antiviral activity.

Table 1: Anti-RSV activity and cytotoxicity of diarylheptanoids.

Compounds	EC ₅₀ ^a (μg/mL)	CC ₅₀ ^b (μg/mL)	CC ₅₀ / EC ₅₀
1,7-Diphenyl-4E-hepten-3-one(AO-1)	36.3 ± 4.2 ^c	47.3 ± 1.3	1.3
7-(4''-Hydroxyphenyl)-1-phenyl-4E-hepten-3-one(AO-2)	5.0 ± 0.0 ^c	22.8 ± 2.5	4.6
7-(4''-Hydroxy-3''-methoxyphenyl)-1-phenyl-4E-hepten-3-one (AO-3)	42.7 ± 3.5	39.3 ± 6.4	0.9
(5R)-5-Hydroxy-1,7-diphenyl-3-heptanone (AO-4)	21.7 ± 0.6 ^c	38.3 ± 3.4	1.8
(5R)-5-Hydroxy-7-(4''-hydroxy-3''-methoxyphenyl)-1-phenyl-3-heptanone(AO-5)	37.0 ± 7.2 ^c	84.8 ± 3.8	2.3
(5R)-5-Methoxy-7-(4''-hydroxy-3''-methoxyphenyl)-1-phenyl-3-heptanone (AO-6)	13.3 ± 3.8	17.0 ± 0.8	1.3
(5S)-5-Methoxy-1,7-diphenyl-3-heptanone (AO-7)	16.3 ± 3.5 ^c	>100	>6.1
(5S)-5-Methoxy-7-(4''-hydroxyphenyl)-1-phenyl-3-heptanone (AO-8)	21.7 ± 0.6 ^c	31.5 ± 6.6	1.5
(3R,5R)-1,7-Diphenylheptan-3,5-diol (AO-9)	22.3 ± 0.6 ^c	56.3 ± 3.1	2.5
Ribavirin	0.67 ± 0.08	NT ^d	NT ^d

The structures of these diarylheptanoids are shown in Figure 1.

^aMeans ± SD for four independent experiments; ^bMeans ± SD for three independent experiments; ^cP<0.05 vs. CC₅₀; ^dNot tested.

Table 2: Anti- poliovirus, -measles virus, and -HSV-1 activities and cytotoxicities of diarylheptanoids.

Compounds	EC ₅₀ ^a (μg/mL)			CC ₅₀ ^b (μg/mL)	CC ₅₀ / EC ₅₀		
	Poliovirus	Measles virus	HSV-1		Poliovirus	Measles virus	HSV-1
AO-1	8.3±2.3 ^c	17.3±1.2 ^c	53.7±4.7	45.8±1.7	5.5	2.6	0.9
AO-3	64.3±4.9	47.0±4.6 ^c	59.7±0.6	63.0±10.4	1.0	1.3	1.1
AO-4	22.7±1.5 ^c	17.0±2.0 ^c	54.0±5.6 ^c	69.5±5.2	3.1	4.1	1.3
AO-5	44.3±4.0 ^c	18.3±1.2 ^c	58.7±1.5 ^c	>100	2.3	5.5	1.7
AO-6	3.7±0.6 ^c	6.3±0.6 ^c	5.7±0.6 ^c	10.8±1.3	2.9	1.7	1.9
AO-7	16.7±2.1 ^c	18.0±1.0 ^c	18.3±0.6 ^c	40.5±5.4	2.4	2.3	2.2
Acyclovir	NT ^d	NT ^d	0.23±0.04	NT ^d	NT ^d	NT ^d	NT ^d

^aMeans ± SD for four independent experiments; ^bMeans ± SD for three independent experiments; ^c*P*<0.05 vs. CC₅₀; ^dNot tested.

Experimental

Chemicals: Dimethyl sulfoxide (DMSO) and ribavirin were purchased from Wako Pure Chemical Industries, Ltd., Osaka, Japan. Acyclovir was purchased from GlaxoSmithKline, Tokyo, Japan. Diarylheptanoids (**AO-1** to **9**, Figure 1 and Table 1) were isolated from the rhizome of *A. officinarum*, as described previously [9,20].

Cells and viruses: Human epidermoid carcinoma (HEp-2) cells (American Type Culture Collection CCL-23) were purchased from Dainippon Pharmaceutical, Osaka, Japan, and grown and maintained in Eagle's minimum essential medium (EMEM; Nissui Pharmaceutical Co. Ltd., Tokyo, Japan) supplemented with 10% and 2% heat-inactivated fetal calf serum, respectively. Vero E6 cells were provided by Dr K. Shiraki (Toyama University, Japan) and grown and maintained in EMEM supplemented with 8% and 2% heat-inactivated calf serum, respectively. The A2 strain of RSV was obtained from American Type Culture Collection (Rockville, MD) and grown in HEp-2 cell cultures. Poliovirus type 1 (Sabin strain), measles virus (Tanabe strain), and HSV-1 (7401H strain) were provided by Dr K. Shiraki (Toyama University, Japan) and propagated in Vero cells [8].

Antiviral and cytotoxic assays: The anti-RSV activities of 9 diarylheptanoids were examined by a plaque reduction assay using HEp-2 cells [21,22]. Briefly, HEp-2 cells grown in 24-well plates were infected with 100 plaque-forming units (PFU)/0.2 mL of RSV at 37°C for 1 h. The cells were overlaid with 1 mL of maintenance EMEM containing 0.8% methylcellulose and various concentrations of either diarylheptanoids or ribavirin and maintained in a humidified atmosphere containing 5% CO₂ for 4–5 days.

The anti-poliovirus, -measles virus, and -HSV-1 activities were also examined by a plaque reduction assay using Vero cells [8]. Duplicate cultures of Vero cells in 60-mm plastic dishes were infected with 100 PFU/0.2 mL of poliovirus, measles virus, or HSV-1 for 1 h. Then the cells

were overlaid with 5 mL of nutrient 0.8% methylcellulose medium containing various concentrations of either diarylheptanoids or acyclovir. The virus-infected cultures were incubated for 2–5 days at 37°C. The infected cells were fixed and stained, and the plaques were counted [8]. All diarylheptanoids were dissolved in DMSO and diluted with culture medium to make the various final concentrations. The concentration of DMSO in each medium was less than 1%. Ribavirin and acyclovir were dissolved in distilled water and DMSO, respectively, and used as controls. The 50% effective antiviral concentration (EC₅₀) was the concentration that reduced virus-induced cell destruction by 50%, as described previously [8,21].

The cytotoxicity of diarylheptanoids was assessed by trypan blue exclusion assays using mock-infected HEp-2 or Vero cells. The cells were seeded at a concentration of 5 × 10⁴ cells/mL in 24-well plates. After incubation at 37°C for 24 h, the culture medium was replaced with fresh medium containing one of the diarylheptanoids at various concentrations and the cells were further incubated for 48 h. After 48 h, the cells were trypsinized and the number of viable cells was determined by a trypan blue exclusion assay. The 50% cytotoxic concentration (CC₅₀) was determined as the concentration that reduced cell destruction by 50% [21].

Statistical analysis: Statistical significances of differences between the EC₅₀ and CC₅₀ values were evaluated using Student's *t*-test. A *P* value of 0.05 or less was considered to be significant statistically.

Acknowledgments - We thank Ms Akane Hino for her excellent technical assistance and Ms Katherine Ono for her editorial assistance. This study was partly supported by Grant-in-Aid for Scientific Research (No. 20590131) from the Japan Society for the Promotion of Science, The Uehara Memorial Foundation, and Health and Labor Sciences Research Grants (Research on Risk of Chemical Substances) from the Ministry of Health, Labor and Welfare of Japan.

References

- [1] Villarruel GR, Langley GE, Abedi GR, Anderson LJ. (2009) Respiratory syncytial virus activity-United States, July 2008–December 2009, *MMWR Morbidity and Mortality Weekly Report*, **59**, 230-233.

- [2] Falsey AR, Hennessey PA, Formica MA, Cox C, Walsh EE. (2005) Respiratory syncytial virus infection in elderly and high-risk adults. *The New England Journal of Medicine*, **352**, 1749-1759.
- [3] Cingoz O. (2009) Motavizumab. *MAbs*, **1**, 439-442.
- [4] Gill MA, Welliver RC. (2009) Motavizumab for the prevention of respiratory syncytial virus infection in infants. *Expert Opinion on Biological Therapy*, **9**, 1335-1345.
- [5] Pelaez A, Lyon GM, Force SD, Ramirez AM, Neujahr DC, Foster M, Naik PM, Gal AA, Mitchell PO, Lawrence EC. (2009) Efficacy of oral ribavirin in lung transplant patients with respiratory syncytial virus lower respiratory tract infection. *Transplantation infection*, **28**, 67-71.
- [6] Weisman LE. (2009) Respiratory syncytial virus (RSV) prevention and treatment: past, present, and future. *Cardiovascular & Hematological Agents in Medicinal Chemistry*, **7**, 223-233.
- [7] Yasukawa K, Sun Y, Kitanaka S, Tomizawa N, Miura M, Motohashi S. (2008) Inhibitory effect of the rhizomes of *Alpinia officinarum* on TPA-induced inflammation and tumor promotion in two-stage carcinogenesis in mouse skin. *Journal of Natural Medicines*, **62**, 374-378.
- [8] Kurokawa M, Ochiai H, Nagasaka K, Neki M, Xu H, Kadota S, Sutardjo S, Matsumoto T, Namba T, Shiraki K. (1993) Antiviral traditional medicines against herpes simplex virus (HSV-1), poliovirus, and measles virus *in vitro* and their therapeutic efficacies for HSV-1 infection in mice. *Antiviral Research*, **22**, 175-188.
- [9] Sun Y, Tabata K, Matsubara H, Kitanaka S, Suzuki T, Yasukawa K. (2008) New cytotoxic diarylheptanoids from the rhizomes of *Alpinia officinarum*. *Planta Medica*, **74**, 427-431.
- [10] Lee HJ, Kim JS, Ryu J-H. (2006) Suppression of inducible nitric oxide synthase expression by diarylheptanoids from *Alpinia officinarum*. *Planta Medica*, **72**, 68-71.
- [11] Kiuchi F, Shibuya M, Sankawa U. (1982) Inhibitors of prostaglandin biosynthesis from *Alpinia officinarum*. *Chemical & Pharmaceutical Bulletin*, **30**, 2279-2282.
- [12] Kiuchi F, Iwakami S, Shibuya M, Hanaoka F, Sankawa U. (1992) Inhibition of prostaglandin and leukotriene biosynthesis by gingerols and diarylheptanoids. *Chemical & Pharmaceutical Bulletin*, **40**, 387-391.
- [13] Yadav PN, Liu Z, Rafi MM. (2003) A diarylheptanoid from Lesser Galangal (*Alpinia officinarum*) inhibits proinflammatory mediators via inhibition of mitogen-activated protein kinase, p44/42, and transcription factor nuclear factor- κ B. *The Journal of Pharmacology and Experimental Therapeutics*, **305**, 925-931.
- [14] Sawamura R, Sun Y, Yasukawa K, Shimizu T, Watanabe W, Kurokawa M. (2010) Antiviral activities of diarylheptanoids against influenza virus *in vitro*. *Journal of Natural Medicines*, **64**, 117-120.
- [15] Sawamura R, Shimizu T, Sun Y, Yasukawa K, Miura M, Toriyama M, Motohashi S, Watanabe W, Konno K, Kurokawa M. (2010) Anti-influenza virus activity *in vitro* and *in vivo* of diarylheptanoids isolated from *Alpinia officinarum*. *Antiviral Chemistry and Chemotherapy*, **21**, 33-41.
- [16] Kawana F, Shigeta S, Hosoya M, Suzuki H, De Clercq E. (1978) Inhibitory effects of antiviral compounds on respiratory syncytial virus replication *in vitro*. *Antimicrobial Agents and Chemotherapy*, **31**, 1225-1230.
- [17] Xie YY, Zhao XD, Jiang LP, Liu HL, Wang LJ, Fang P, Shen KL, Xie ZD, Wu YP, Yang XQ. (2006) Inhibition of respiratory syncytial virus in cultured cells by nucleocapsid gene targeted deoxyribozyme (DNAzyme). *Antiviral Research*, **71**, 31-41.
- [18] An N, Xu L-Z, Zou Z-M, Yang S-L. (2006) Diarylheptanoids from *Alpinia officinarum*. *Journal of Asian Natural Products Research*, **8**, 637-641.
- [19] Shin D, Kinoshita K, Koyama K, Takahashi K. (2002) Antiemetic principles of *Alpinia officinarum*. *Journal of Natural Products*, **65**, 1315-1318.
- [20] Sun, Y, Matsubara H, Kitanaka S, Yasukawa K. (2008) Diarylheptanoids from the rhizomes of *Alpinia officinarum*. *Helvetica Chimica Acta*, **91**, 118-123.
- [21] Watanabe W, Shimizu T, Hino A, Kurokawa M. (2008) A new assay system for evaluation of developmental immunotoxicity of chemical compounds using respiratory syncytial virus infection to offspring mice. *Environmental Toxicology and Pharmacology*, **25**, 69-67.
- [22] Watanabe W, Konno K, Ijichi K, Inoue H, Yokota T, Shigeta S. (1994) MTT colorimetric assay system for the screening of anti-orthomyxo- and anti-paramyxoviral agents. *Journal of Virological Methods*, **48**, 257-265.

Isolation of C-glycosyl Xanthones from *Coffea pseudozanguebariae* and Their Location

Pascale Talamond^{1,2}, Geneviève Conejero³, Jean-Luc Verdeil³ and Jean-Luc Poëssel⁴

¹UMR Diversity and Adaptability of Crops, IRD, 911, avenue d'Agropolis, BP 64501, 34394 Montpellier cedex 5, France

²Current address: UMR 226 IRD-ISEM, 361, rue Jean-François Breton, BP 5095, 34196 Montpellier cedex 5, France

³UMR BPMP 5004, UMR DAP 1098, Plate-forme d'Histocytologie et Imagerie Cellulaire Végétale, avenue Agropolis, 34398 Montpellier, cedex 5, France

⁴UR 1052, Genetic Improvement of Fruit and Vegetables, Domaine St Maurice, 84143 Montfavet, France

pascale.talamond@ird.fr

Received: October 21st, 2010; Accepted: July 17th, 2011

The biochemical composition of leaves from *Coffea pseudozanguebariae*, a wild caffeine-free coffee species, was determined. Two phenolic compounds were extracted from leaves, separated and characterized. Their structures were elucidated by mass spectrometry, and 1D and 2D NMR spectroscopy and were shown to be mangiferin (**1**) and isomangiferin (**2**), which were the main polyphenol products. Multiphoton fluorescence imaging was performed to visualize polyphenol distribution in leaf cross sections. Consistent biochemical analysis cell imaging techniques on leaves revealed yellow fluorescence in the epidermis and parenchyma cells corresponding to xanthone compounds.

Keywords: *Coffea pseudozanguebariae* Bridson, xanthones, fluorescence, mangiferin, isomangiferin, Rubiaceae.

Abbreviations: hydroxycinnamoyl quinic acid, HQA, caffeoylquinic acids (3,4,5), CQA; di-caffeoyl quinic acids (3,4; 3,5; 4,5), diCQA.

Coffea pseudozanguebariae Bridson, native to East Africa, is a wild species of coffee tree and the first caffeine-free species discovered in tropical East Africa. It exhibits different morphological and physiological characteristics from other species of coffee: one of the shortest fruiting times [1a], low hydroxycinnamoyl quinic acid (HQA) content [1.2% dmb (dry matter basis)] [1b], morphological aspects (small purple fruits on complete ripeness and small sized leaves), and a small sized genome (1.13 pg). Previous phytochemical investigations described the presence of alkaloids, with the major component trigonelline, diterpenes (cafestol, kahweol), and phenolic compounds (HQA, hydroxycinnamoyl quinic acid, or chlorogenic acids) in *C. pseudozanguebariae*. The unsaponifiable lipid fraction of green beans contains a mozambioside, a diterpene glycoside [2], high cafestol, kahweol, and four unknown diterpenes [3]. The diterpene glycoside, which is not present in the other species of commercial coffee trees, is the origin of the strong bitterness of this coffee [2]. In the soluble fraction, two classes of secondary metabolites have been particularly studied in green coffee beans: alkaloids and phenolic compounds. The content of the major alkaloid, caffeine, varies markedly between species, from 0%, dry wt, in *C. pseudozanguebariae* to 3.2%, dry wt, in *C. canephora* [4]. Trigonelline, a major coffee aroma compound, forms

1.02%, dry wt, of *C. pseudozanguebariae*, 0.67%, dry wt, of *C. canephora*, and 0.57%, dry wt, of *C. liberica* var. *dewevrei* [5]. Only two classes of phenolic compounds have been described in coffee plants: a major one, HQA and a minor one, proanthocyanins, from the flavonoid class [6]. Other phenolic compounds are polymeric, such as tannins. Lignans are also present in coffee seeds, although in minor amounts. This main family is formed from esters between hydroxycinnamate and quinic acid. No previous study on *C. pseudozanguebariae* has been reported in the literature. In this paper, we report the first chemical investigation of *C. pseudozanguebariae* leading to the isolation of compounds including the representative phenolic compounds and the visualization of their accumulation in the tissue.

A methanolic extract of *C. pseudozanguebariae* leaves was chromatographed on a RP-18 column and showed a major peak with a retention time at 10.1 min. This peak exhibited a spectrum with four maximum UV absorbances at 240 nm, 257 nm, 316 nm, and 365 nm. The minor peak at 10.9 min had the same UV characteristics. A peak at 7.3 min was identified, with the help of UV spectra, as 5-*O*-caffeoylquinic acid (or chlorogenic acid). Semi-preparative chromatography was used to isolate the unknown compounds. After evaporation, a yellow powder




Figure 1: Structure of C-glucosyl xanthenes. **1:** C2- β -D-glucoside 1,3,6,7 tetrahydroxyxanthen-9-one (or mangiferin), **2:** C4- β -D-glucoside 1,3,6,7 tetrahydroxyxanthen-9-one (or isomangiferin).

was obtained and used to carry out the first analysis for structure elucidation. Another separation system using a combination of cellulose column and Sephadex LH-20 gel filtration gave yellow crystalline materials (compounds **1**, **2**). From their physical, EI mass spectrometric and NMR spectroscopic features compound **1** was unambiguously established as 2-C- β -D-glucopyranosyl- 1,3,5,7 tetrahydroxy-xanthen-9-one (mangiferin) [7a,7b], and **2** as 4-C- β -D-glucopyranosyl- 1,3,5,7 tetrahydroxyxanthen-9-one (isomangiferin) [8]. Compound **2** led to a slight hypsochrome UV spectrum displacement at 318 nm. The ^1H spectrum of **2** is very close to that of **1** except for signals of two aromatic protons at δ_{H} 6.02 and 6.17.

Cross-sections of fresh *C. pseudozanguebariae* leaves at two stages of development were observed by epifluorescence microscopy. The young leaves (maroon color, and less than 4 cm long) showed a yellow and orange fluorescence in all cells (Figure 2A). The mature leaves (green color) did not show such a fluorescence, except for some cells in the mesophyll, which gave an orange fluorescence (Figure 2B). It is known that phenolic compounds naturally have the ability to emit fluorescence under ultraviolet light. The autofluorescence of polyphenols was used to localize them in tissues of the leaves. In order to visualize the accumulation of phenolic compounds, a spectral analysis was developed on the same fresh leaf sections. This technique was achieved with a special detection system, which enabled separation of the signals from different fluorophores. We used this technique with a multiphotonic microscope because the infra-red laser associated with this microscope gave the possibility of exciting molecules like a UV laser. With an excitation wavelength between 700 and 800 nm, it was possible to excite like a UV laser between 350 and 400 nm. In these optical conditions, spectral signatures were acquired for each pixel of the scanned image (from either leaf section or purified compound) and could subsequently be used for digital separation into component dyes. Then, from these spectral signatures, the Linear Unmix method was used to discriminate between various fluorescence signals, even with widely overlapping emission spectra. The emission spectral curve data obtained from the compounds were recorded in a spectral library and applied on cross-sections of leaves to localize them. The

calculation was realized using a high number of iterations and by keeping a residual channel dedicated to pixels not corresponding to the selected spectral signatures. The spectral acquisition obtained on young leaf cross-sections showed a complex pattern of autofluorescence resulting from various fluorescent compounds. The fluorescence observed in young leaves, identified as the spectrum emission of mangiferin, was found after Linear Unmixing calculation in the upper epidermis and in some mesophyll cells (Figure 2C). The same technique was used with the spectrum emission of 5-O-caffeoylquinic acid (chlorogenic acid), the main phenolic compound in coffee, but no signal was detected on the leaves (data not shown). There were other uncharacterized fluorescent compounds in the mesophyll, which did not correspond to the known compounds and which seems to belong to other families of secondary metabolites. These compounds are not accumulated in the leaves of *C. arabica* and *C. canephora*, the commercial species of coffee. The decrease in yellow fluorescence observed between the very young and mature leaves may tally with biochemical analysis [9], in particular, during leaf development; there is a sharp decrease in the main phenolic compound, mangiferin. Consequently, tissue-specific localization could provide valuable information for understanding the actual role in the mechanisms of acclimatization to several environmental agents such as UV radiation or against pathogen and predator attacks. The fluorescent technologies now available allow researchers to study such dynamic processes in living cells.

Experimental

General experimental procedures: Chromatographic separation was performed using a HPLC series (Shimadzu, Prominence LC) equipped with software, a DGU-20A₃ degasser, an LC-20AD binary gradient pump, a SIL-20AC thermoautosampler, and a SPD-M20A diode array detector. The column used was a LiChrospher 5 μm RP18 (250X4 mm i.d.) from Merck (Darmstadt, Germany), and a guard column of the same material. The mobile phase consisted of 2mM phosphoric acid in water (eluent A) and MeOH (eluent B). The gradient program was as follows: a 25-80% MeOH gradient over 40 min at a flow rate of 0.8 mL.min⁻¹. Separation was at room temperature and the injection volume was 500 μL for isolation and 20 μL for identification. Phenolic compounds were identified by comparing their retention times with appropriate standards: 5-caffeoyl quinic acid, 3-caffeoyl quinic acid and 3,5-dicaffeoylquinic acid. A Q-TOF Micromass (Waters, Milford, MA, USA) mass spectrometer was used to obtain the MS data. The sample was solubilized in 50% water-acetonitrile solution, acidified using 0.1% trifluoroacetic acid solution and introduced at a flow rate of 5 $\mu\text{L min}^{-1}$. All the analyses were performed using an electrospray ionisation source (ESI) set to 100°C in positive ion mode with the following settings: capillary voltage -3000 V, cone voltage = 30 and 50 V, nebuliser gas (N_2) 400 L/h, desolvation temperature 120°C, drying gas (N_2) 20 l/h.




Figure 2: Histological imaging of fluorescent compound accumulation in the leaves of *C. pseudozanguebariae*. Cross-sections of young leaves (A) and mature leaves (B) by epifluorescence microscopy under UV light (long-pass filter 425 nm, real colours). Cross-sections of young leaves by spectral analysis and linear unmixing (C) on a multiphotonic microscope (false colors, yellow: mangiferin, red: chlorophyll, white: unknown). ep: epidermis, pp: palisade parenchyma, sp: spongy parenchyma, bs: bundle sheath cell, bar = 20 μ m.

Mass spectra were monitored using MassLynx4.0 software, with an acquisition rate of 2 spectra s^{-1} in the range $m/z = 70$ -600. NMR spectra were measured using a Bruker DRX-400 spectrometer at 400 MHz (1H) and 100 MHz (^{13}C).

Histology: Thick cross-sections (50 μ m) were obtained from young leaves of *Coffea pseudozanguebariae* using a vibratome (MICROM) and then dipped in 10 mM phosphate buffer saline (7 mM Na_2HPO_4 , 3 mM NaH_2PO_4 , 120 mM NaCl, 2.7 mM KCl). Epi-fluorescence microscopy was carried out on a Leica DMRXA equipped with a Q-Imaging camera (long-pass filter 425 nm). A Zeiss 510 META NLO multiphoton microscope equipped with a Coherent Chameleon Ultra II laser was used to obtain emission fluorescence from fresh leaves. Spectral analysis was carried out using the autofluorescence properties of polyphenol compounds without any dyes. Reference spectra on purified powder of polyphenol compounds (HQA and xanthenes) were obtained from spectral acquisition with $\square\square$ excitation at 780 nm and $\square\square$ emission between 400 and 700 nm. The Linear Unmixing Function of the microscope (method of

Emission Finger printing from Zeiss) was used to visualize the fluorescence of polyphenol compounds in cells from reference spectra.

Plant material: In these experiments we used *Coffea* plants from the collection cultivated at the IRD research centre in Montpellier (France). Leaves of *C. canephora* (DB56, DB57) and *C. pseudozanguebariae* (H65, H70) of various genotypes were taken from trees maintained in a tropical greenhouse (natural daylight, 25°C, 28°C day, 78-82% humidity). The young leaves, under 4 cm long, were cut from the tips of branches. Leaves were harvested from 5 different genotypes and 500 g of the collected leaves were frozen in liquid nitrogen immediately after collection then stored at -80°C. Each batch of leaves was freeze-dried for 48 h then stored in a cold room. *C. pseudozanguebariae* fruits were harvested at the CNRA Station in Divo (Ivory Coast) from field-grown trees.

Extraction and isolation: Fine powder of leaves and beans was obtained in an analytical grinder (IKA, yellow, A10) by grinding for 1 min, repeated 3 times. The leaf powders (2 g) were extracted with 70% MeOH solution (30 mL, X3) at 4°C and bean powders (50 mg) in 5 mL of 70%

MeOH solution. The MeOH extracts were combined, filtered and evaporated under reduced pressure to give a residue. This was suspended in MeOH and filtered through a 0.2- μ m disposable filter tip-syringe assembly and directly analysed by HPLC for separation and identification. A second technique was used with 80 g of freeze-dried leaves suspended in 700 mL of MeOH-H₂O (8:2) solution at room temperature for 20 min with sonication (20 min, 24 KHz, R.E.U.S.-GEX 180, Contes, France), repeating 3 times. MeOH was removed in a rotavapor. After freeze-drying, the aqueous phase was subjected to a Medium Pressure (MP) column (400X47 mm) packed with cellulose (microcristallin Avicel,

Darmstadt, Germany) and successively eluted with H₂O and H₂O-MeOH (1:9). Collected fractions were purified on a Sephadex LH-20 (Fluka) column (500 X 25 mm, Fluka, Basel, Switzerland) and eluted with H₂O for compound **1**. For compound **2**, fractions were purified on a MP column (210 X 30 mm, Buchi) eluted with EtOH- H₂O (8:2). After freeze-drying, both compounds (**1** and **2**) were obtained.

Acknowledgements - The authors thank Dr Maryse Bejaud for support in NMR analysis and helpful discussion. We wish to thank Elisabeth Ambert for assistance in checking bibliographic databases and for valuable help.

References

- [1] (a) Akaffou DS, Ky CL, Barre P, Hamon S, Louarn J, Noirot M. (2003) Identification and mapping of a major gene (Ft1) involved in fructification time in the interspecific cross *Coffea pseudozanguebariae* x *C. liberica* var. 'dewevrei': impact on caffeine content and seed weight. *Theoretical Applied Genetics*, **106**, 1486-1490; (b) Ky CL, Louarn J, Guyot B, Charrier A, Hamon S, Noirot M. (1999) Relations between and inheritance of chlorogenic acid contents in an interspecific cross between *Coffea pseudozanguebariae* and *C. liberica* var. 'dewevrei'. *Theoretical Applied Genetics*, **98**, 628-637.
- [2] Prewo R, Guggisberg A, Lorenzi-Riatsch A, Baumann TW, Wettstein-Bättig M. (1990) Crystal structure of mozambioside, a diterpene glycoside of *Coffea pseudozanguebariae*. *Phytochemistry*, **29**, 990-992.
- [3] de Roos B, van der Weg G, Urgert R, van de Bovenkamp P, Charrier A, Katan MB. (1997) Levels of cafestol, kahweol, and related diterpenoids in wild species of the coffee plant *Coffea*. *Journal of Agricultural and Food Chemistry*, **45**, 3065-3069.
- [4] Anthony F, Noirot M, Clifford MN. (1993) Biochemical diversity in the genus *Coffea* L.: Chlorogenic acids, caffeine, and mozambioside contents. *Genetic Resources and Crop Evolution*, **40**, 61-70.
- [5] Ky CL, Guyot B, Louarn J, Hamon S, Noirot M. (2001) Trigonelline inheritance in the interspecific cross between *Coffea pseudozanguebariae* x *C. liberica* var. dewevrei cross. *Theoretical Applied Genetics*, **102**, 630-634.
- [6] Clifford MN, Gibson CL, Rakotomalala JJ, Cros E, Charrier A. (1991) Caffeine from green beans of *Mascarocoffea*. *Phytochemistry*, **30**, 4039-4040.
- [7] (a) Fujita, M., Inoue, T. (1982) Studies on the constituents of *Iris florentina* L.II.- C-Glucosides of xanthenes and flavones from the leaves. *Chemical & Pharmaceutical Bulletin*, **30**, 2342-2348; (b) Catalano S, Lushi S, Flamini G, Cioni PL, Nieri EM, Morelli I. (1996) A xanthone from *Senecio mikanioides* leaves. *Phytochemistry*, **42**, 1605-1607.
- [8] de Beer D, Jerz G, Joubert E, Wray V, Winterhalter P. (2009) Isolation of isomagiferin from honeybush (*Cyclopia subternata*) using high-speed-counter-current chromatography and high-performance liquid chromatography. *Journal of Chromatography A*, **1216**, 4282-4289.
- [9] Bertrand C, Noirot M, Doulebeau S, de Kochko A, Hamon S, Campa C. (2003) Chlorogenic acid content swap during fruit maturation in *Coffea pseudozanguebariae*. Qualitative comparison with leaves. *Plant Science*, **165**, 1355-1361.

Antifungal Activity and Isomerization of Octadecyl *p*-coumarates from *Ipomoea carnea* subsp. *fistulosa*

Eugene Sebastian J. Nidiry*, Girija Ganeshan and Ankanahalli N. Loksha

Indian Institute of Horticultural Research, Hessaraghatta Lake P.O., Bangalore-560 089, India

nidiry@yahoo.co.in

Received: June 6th, 2011; Accepted: October 25th, 2011

Bioassay monitored HPLC assisted isolation and purification of the chief antifungal fraction of the leaves of *Ipomoea carnea* subsp. *fistulosa* (Convolvulaceae) were achieved using *Colletotrichum gloeosporioides* and *Cladosporium cucumerinum* as test organisms. The activity of the purified fraction was further confirmed by the dose dependent inhibition of the spore germination of *Alternaria alternata* and *A. porri*. The active fraction was identified as a mixture of (*E*)-octadecyl *p*-coumarate and (*Z*)-octadecyl *p*-coumarate. The two isomers were detected on an HPLC column with substantially different retention times, but once eluted from the column, one form was partly converted to the other in daylight. Conclusive evidence for the structures and their isomerization were obtained from the HPLC behavior, IR, UV, HRESIMS, CIMS and NMR spectral data. Important ¹H NMR and ¹³C NMR signals could be separately assigned for the isomers using 2D NMR techniques.

Keywords: *Ipomoea carnea*, *Ipomoea fistulosa*, Convolvulaceae, antifungal, (*E*) octadecyl *p*-coumarate, (*Z*) octadecyl *p*-coumarate, isomerization, 2D NMR techniques.

Ipomoea carnea subsp. *fistulosa* (Mart. ex Choisy) D.F. Austin (Convolvulaceae) is a plant native to South America, but sparsely distributed in India and Bangladesh. It is used in hedgerows along cattle crossings, to fight erosion and as an ornamental. Isolation and chemical characterization of resinous glycosides [1], flavonol glycosides [2] and alkaloids [3] from the leaves, and anthocyanin from the flowers [4] of *I. carnea* have been reported. The leaves are toxic to cattle and the toxicity is attributed to polyhydroxy alkaloids such as swainsonine and calystegines [5]. Recently, a chitinase has been identified in the plant [6]. Antibacterial and antifungal activities of the extractives of the plant have been reported [7], but bioassay monitored isolation and characterizations of the antifungal compounds present in the plant have not yet been carried out. We hereby report the bioassay monitored isolation and characterization of the chief antifungal fraction. The fraction was isolated using *Colletotrichum gloeosporioides* and *Cladosporium cucumerinum* as test organisms and the activity was further confirmed against the spore germination of *Alternaria alternata* and *A. porri*. The active fraction was found to be a mixture of (*E*)-octadecyl *p*-coumarate and (*Z*)-octadecyl *p*-coumarate. The two isomers were detected on the HPLC column with substantially different retention times, but once eluted from the column, one form was partly converted to the other. Conclusive evidence for the structures and their isomerization were obtained from the HPLC behavior, IR, UV, HRESIMS, CIMS, ¹H NMR, ¹³C NMR, DEPT and 2D NMR spectral data [8]. Survey of the literature showed several reports on the isolation of octadecyl *p*-coumarates without any mention of its

antifungal activity [9]. In all these studies, characterization was achieved without resorting to detailed analysis of ¹³C NMR, DEPT and 2D NMR spectra and, for the same reason, isomerization of the (*E*) and (*Z*) forms was not reported. The sole report on antifungal activity [10] is about a mixture of stearyl esters and not of any individual compound.

Thus, this is the first report on the antifungal activity and isomerization of octadecyl *p*-coumarates. This is also the first report in which important ¹H and ¹³C NMR signals have been separately assigned for the (*E*) and (*Z*) isomers of octadecyl *p*-coumarates (Table 1). This may also be the first report in which HSQC data have been used to confirm the isomerization of alkyl *p*-coumarates.




Figure 1: Isomerization of (*E*)-octadecyl-*p*-coumarate (1a) and (*Z*)-octadecyl-*p*-coumarate (1b).

Antifungal activity assay of the crude extractives showed that the ethyl acetate extractive possessed highest activity. Column chromatography of this revealed that the most active fraction was that obtained by elution with *n*-hexane-

Table 1: NMR spectroscopic data (400 MHz, CDCl₃) of *E*-octadecyl *p*-coumarate (**1a**) and *Z*-octadecyl *p*-coumarate (**1b**).

Sl. No.	δ_C , mult	Carbon number in the figure	δ_H , mult. (<i>J</i> in Hz)	Assigned to <i>E/Z/Both</i>	
1.	167.56	C	9	-	<i>E</i>
2.	166.73	C	9	-	<i>Z</i>
3.	157.59	C	4	-	<i>E</i>
4.	156.62	C	4	-	<i>Z</i>
5.	144.22	CH	7	7.62, d, <i>J</i> =12.8 Hz	<i>E</i>
6.	143.18	CH	7	6.84, d, <i>J</i> =10.0 Hz	<i>Z</i>
7.	132.30	CH	6,2	7.63, d, <i>J</i> = 6.8 Hz	<i>Z</i>
8.	129.92	CH	6,2	7.43, d, <i>J</i> =6.8 Hz	<i>E</i>
9.	127.52	C	1	-	<i>Z</i>
10.	127.35	C	1	-	<i>E</i>
11.	117.33	CH	8	5.83, d, <i>J</i> =10.0 Hz	<i>Z</i>
12.	115.84	CH	3,5	6.84, d, <i>J</i> =6.8 Hz	<i>E</i>
13.	115.78	CH	8	6.30, d, <i>J</i> =12.8 Hz	<i>E</i>
14.	114.94	CH	3,5	6.80, d, <i>J</i> =6.8 Hz	<i>Z</i>
15.	64.67	CH ₂	1'	4.19, t, <i>J</i> =5.2 Hz	<i>E</i>
16.	64.35	CH ₂	1'	4.12, t, <i>J</i> =5.2 Hz	<i>Z</i>
17.	31.92	CH ₂	16'	1.25-1.39, m	Both
18.	29.69*	CH ₂	6', 13'	1.25-1.39, m	Both
19.	29.65*	CH ₂	7', 12'	1.25-1.39, m	Both
20.	29.59*	CH ₂	8', 11'	1.25-1.39, m	Both
21.	29.54*	CH ₂	9', 10'	1.25-1.39, m	Both
22.	29.36*	CH ₂	4', 15'	1.25-1.39, m	Both
23.	29.29*	CH ₂	5', 14'	1.25-1.39, m	Both
24.	28.76	CH ₂	2'	1.69, quintet, <i>J</i> =5.6 Hz	Both
25.	25.98	CH ₂	3'	1.25-1.39, m	<i>Z</i>
26.	25.97	CH ₂	3'	1.25-1.39, m	<i>E</i>
27.	22.69	CH ₂	17'	1.25-1.39, m	Both
28.	14.11	CH ₃	18'	0.88, t, <i>J</i> =5.2 Hz	Both
29.	OH (phenolic)	-	5.41		<i>E</i>
30.	OH (phenolic)	-	5.34		<i>Z</i>

*Assignments interchangeable

ethyl acetate (7:3) (Tables 2 and 3). This fraction was subjected to HPLC purification using dichloromethane as eluent and two main peaks (first with t_R =34 min and second t_R =48 min) were detected. Bioassay using *Cladosporium cucumerinum* revealed that fractions corresponding to these two peaks possessed antifungal activity. If the fraction were injected immediately after elution, HPLC showed that peak alone, but after exposure to daylight for a few hours, each of the peaks showed the presence of the other one indicating isomerization of the compounds. Since it was clearly known that the two isomers could not be separated under normal conditions, the compound corresponding to the major peak (t_R =48 min) was collected for further characterization. This purified fraction gave an approximate minimum inhibitory dose of 0.3 mg against the spore germination of *Cladosporium cucumerinum* on a TLC plate (Table 3). Antifungal activity of the purified fraction was further confirmed by the spore germination inhibition of *Alternaria alternata* and *A. porri* (Table 4). Attempts to obtain good quality crystals for X-ray diffraction studies failed. Powder diffraction also did not give any useful information.

Spectral characterization of the fraction having a m.p. 79-80°C (containing **1a** as the main constituent) was achieved using UV, IR, HRESIMS, CIMS, ¹H NMR, ¹³C NMR, DEPT, COSY, HSQC and C, H analysis. The CIMS showed a [M+H]⁺ ion at *m/z* 417, the HRESIMS a [M+Na]⁺ ion at

Table 2: Antifungal activity of the extractives of *Ipomoea carnea* subsp. *fistulosa* against the mycelial growth of *Colletotrichum gloeosporioides*.

Extractive/Fraction/Compound	Conc.	% mycelial growth inhibition
<i>n</i> -Hexane	0.5%	0.0
Ethyl acetate	0.5%	24.2 (±0.8)
Methanol	0.5%	20.4 (±1.4)
Active fraction from column*	0.5%	68.5 (±0.7)
Phenol (standard)	0.05%	78.3 (±0.4)

* Active fraction from column was obtained by elution with *n*-hexane-ethyl acetate (7:3) mixture**Table 3:** Antifungal activity of the compounds of *Ipomoea carnea* subsp. *fistulosa* against *Cladosporium cucumerinum* by TLC bioautography.

Extractive/Fraction/Compound	Dose (mg)	Inhibition**	MID
<i>n</i> -Hexane	5.0	-	ND
Ethyl acetate	5.0	+	ND
Methanol	5.0	++	ND
Active fraction from column*	5.0	+++	ND
Octadecyl <i>p</i> -coumarate	0.3	+	0.3 mg
Phenol (standard)	0.3	+	0.3 mg

* Active fraction from column was obtained by elution with *n*-hexane-ethyl acetate (7:3).

** The observations on TLC plate for *n*-hexane, ethyl acetate and methanol extractives and active column fraction* were made after elution with ethyl acetate (*R_f* value of the inhibition spot ca 0.8). Observations on phenol and octadecyl *p*-coumarate were made by direct bioautography without elution after spotting the compounds quantitatively. '+' indicates observable inhibition, '++' indicates clear inhibition, '+++ indicates very clear inhibition and '-' indicates no inhibition. MID=minimum inhibitory dose; ND=Not determined.

m/z 439.3198, corresponding to a molecular weight of 416.3300, and C, H analysis gave C, 76.71%; H, 10.40%. These data gave the molecular formula as C₂₇H₄₄O₃ (required C, 77.8%; H, 10.45% and M⁺ 416.3291). The ¹H NMR spectrum showed two sets of closely related signal patterns indicating the presence of two isomers in the sample. The ratio of intensity of the peaks based on their coupling constants showed that the *E* and *Z* isomers exist in the ratio 2:1. This was also consistent with the 2:1 ratio of the areas of HPLC peaks with t_R =48 min and t_R =34. ¹H-¹H COSY showed two sets of signal correlations for each of the isomers. In the first set, the signal at δ 7.62 (1H, d, *J*=12.8 Hz) correlated with that at δ 6.30, δ 7.43 (2H, d, *J*=6.8 Hz) with δ 6.84 and δ 4.19 (2H, t, *J*=5.2 Hz) with δ 1.69. In the second set, the signal at δ 7.63 (2H, d, *J*=6.8 Hz) correlated with that at δ 6.84, δ 6.83 (1H, d, *J*=10.0 Hz) with δ 5.83, and δ 4.12 (2H, t, *J*=5.2 Hz) with δ 1.63. ¹H-¹³C HSQC also showed two sets of correlations. In the first set, signal at δ_C 144.22 correlated with that at δ_H 7.62, δ_C 129.92 with δ_H 7.43, δ_C 115.84 with δ_H 6.84, δ_C 115.78 with δ_H 6.30 and δ_C 64.67 with δ_H 4.19. In the second set, signal at δ_C 143.18 correlated with that at δ_H 6.83, δ_C 132.30 with δ_H 7.63, δ_C 117.33 with δ_H 5.83, δ_C 114.94 with δ_H 6.80 and δ_C 64.35 with δ_H 4.12. The ¹³C NMR spectrum showed 28 signals, which were assigned for both *E/Z* isomers taking into consideration DEPT, COSY and HSQC data (Table 1). All these results led to the conclusion that the active fraction is a mixture of (*E*)-octadecyl *p*-coumarate (**1a**) and (*Z*)-octadecyl *p*-coumarate (**1b**).

The isomerization of (*E*)-octadecyl *p*-coumarate and (*Z*)-octadecyl *p*-coumarate is shown in Figure 1. Our finding of the isomerization of octadecyl *p*-coumarate corroborates

the reports on the isomerization of structurally related eicosanyl *p*-coumarates isolated from *Psiadia punctulata* [11] and 21'-hydroxyheicosanyl-4-hydroxy-(*cis*- and *trans*) *p*-coumarate isolated from *Tanacetum longifolium* [12].

The fact that four earlier reports [9b-9e] on the isolation of octadecyl *p*-coumarates were from the genus *Ipomoea* may be of chemotaxonomic interest.

Experimental

General: UV spectra were obtained with a Spectronic UV-Visible spectrophotometer. IR spectra were obtained on a Perkin-Elmer spectrophotometer. ¹H NMR and ¹³C NMR spectra were recorded on a Bruker spectrometer operating at 400 MHz and 100 MHz, respectively (Table 1). HRESIMS were obtained on a Micromass Q-TOF apparatus. Carbon-Hydrogen-Nitrogen analysis was obtained using a ThermoFinnigan Flash EA 1112 CHNS analyzer. HPLC purification was achieved using a Waters HPLC system (515 pump, 7725 Rheodyne injector, Waters 2487 Dual λ absorbance detector) under conditions as follows: Column Prep Nova Pak HR Silica 7.8 x 300 mm, flow rate 1 mL/min, UV detection at 254 nm, eluent CH₂Cl₂. TLC bioassay was achieved with silica gel plates (0.5 mm thickness). Observation of spore germination inhibition was made using a Carl Zeiss Axio Imager AI microscope.

Plant material: The leaves of *Ipomoea carnea* subsp. *fistulosa* were collected from the farm of the Indian Institute of Horticultural Research, Hessaraghatta Lake P.O., Bangalore – 560089, India and a voucher specimen is kept at the Section of Medicinal Crops of the Institute.

Extraction and isolation: The dried plant material (2 Kg) was extracted first with *n*-hexane, then with ethyl acetate and finally with methanol. Column chromatography of the ethyl acetate extractive utilizing silica gel with *n*-hexane-ethyl acetate mixtures with increasing percentages of ethyl acetate and TLC bioassay were conducted side by side. The fraction which showed maximum activity was taken for HPLC purification. Two main peaks (*t_R*=34 min and *t_R*=48 min) were detected and the eluents corresponding to these peaks were collected separately for further investigation.

Antifungal activity assays

Poisoned food technique: Pure culture of *Colletotrichum gloeosporioides* ITCC 4573 obtained from Indian Type Culture Collections, Indian Agricultural Research Institute, New Delhi, India was used for this study [13]. Percent mycelial growth inhibition values presented in Table 2 are the averages of 2 replications, standard deviation being presented in parenthesis. The purified HPLC fraction was not used in this technique because of the paucity of material and its poor solubility both in water and solvents miscible with water.

Table 4: Spore germination inhibition of *Alternaria alternata* and *A. porri* by octadecyl *p*-coumarates (*E* and *Z* isomers in ratio 2:1).

Treatment	<i>Alternaria alternata</i>		<i>Alternaria porri</i>	
	% germination	% inhibition w.r.t. control	% germination	% inhibition w.r.t. control
Control (3% <i>n</i> -propanol in water)	92.0 (±0.5)	-	95.0 (±0.0)	-
Octadecyl coumarates (100 mg/L)	67.0 (±1.4)	26.6 (±0.8)	71.5 (±0.7)	24.8 (±0.8)
Octadecyl coumarates (500 mg/L)	51.5 (±2.1)	44.1 (±2.3)	55.0 (±2.8)	42.1 (±3.0)
Phenol (standard) (100 mg/L)	86.0 (±0.0)	6.5 (±0.0)	91.0 (±1.4)	4.3 (±1.5)
Phenol (standard) (500 mg/L)	71.5 (±0.7)	22.3 (±0.8)	79.0 (±2.8)	16.8 (±3.0)

TLC bioautography: A pure culture of *Cladosporium cucumerinum* IMI 249540 obtained from the International Mycological Institute, U.K., maintained on a potato-dextrose-agar (PDA) medium was used for this assay [14].

Spore germination inhibition study: For this study [13], spores of *Alternaria alternata* from infected tomato fruits and *A. porri* from infected onion leaves collected from the IIHR experimental farm in Hessaraghatta, Bangalore, India were used. Spores were added to a solution of the compound in 3% *n*-propanol in water kept in cavity slides by the hanging drop method. Observation on spore germination was recorded after incubation for 3 h. Percent spore germination inhibition values presented in Table 4 are the averages of 2 replications, standard deviations being given in parenthesis.

(*E*)-Octadecyl *p*-coumarate (1a): It was collected at *t_R* of 48 min. as major peak during HPLC separation. The compound got partly converted to (*Z*)-octadecyl *p*-coumarate after a few hours. White solid with a faint yellowish to greenish tinge (20 mg).

MP: 79-80°C.

IR: 3393 (OH stretching), 2921 (C-H stretching), 2880 (C-H stretching), 1713 (α, β unsaturated ester), 1674 (C=C of phenol), 1604 (C=C of α, β unsaturation), 1586 (aromatic C=C), 1516 (aromatic C=C), 1468 (C-H), 1377 (CH₃), 1307 (C-O stretching), 1274 (C-O stretching), 1170 (C-O stretching), 982 (C=C conjugated to C=O), 835 (C=C-H), 722 (CH₂), 517 cm⁻¹.

UV (MeOH) λ_{max}: 225, 308 nm. Second peak showed bathochromic shift on addition of NaOH.

¹H NMR and ¹³C NMR (CDCl₃): Table 1.

CIMS: 417 [M+H]⁺, 164 (HO-C₆H₄ CH=COOH) 147 (HO-C₆H₄CH=CO), 129, 120, 107.

HRESIMS [M+Na]⁺ 439.3198 (required for C₂₇H₄₄O₃Na 439.3189).

Elemental analysis: Found C, 76.71; H, 10.40 (C₂₇H₄₄O₃ requires C, 77.80; H, 10.45).

Direct bioautography on TLC plate was done using *Cladosporium cucumerinum* (Table 3). The activity was further confirmed by spore germination inhibition of *Alternaria alternata* and *A. porri* (Table 4).

(Z)-Octadecyl p-coumarate (1b): It was collected at t_R of 34 min as minor peak during HPLC separation. It got partly

converted to *E*-octadecyl *p*-coumarate after a few hours. White solid with a yellowish to greenish tinge (10 mg).

MP: 79-80°C.

¹H NMR and ¹³C NMR (CDCl₃): Table 1.

Acknowledgements - The authors thank Dr A.S. Sidhu, Director, I. I. H. R. for encouragement. They are grateful to NMR Research Centre, Indian Institute of Science, Bangalore, India for spectral data.

References

- [1] Legler U. (1965) Die bestandteile des giftigen glykosidharzes aus *Ipomoea fistulosa* Mart. ex Choisy. *Phytochemistry*, **4**, 29-41.
- [2] (a) Lamidi M, Rondi ML, Oliver E, Faure R, Nze EL, Balansard G. (2000) Constituents of *Ipomoea fistulosa* leaves. *Fitoterapia*, **71**, 203-204; (b) Dubey P, Khare N, Gupta PC. (1982) A new flavonoid glycoside from the leaves of *Ipomoea fistulosa*. *Current Science*, **51**, 351-352.
- [3] Umar S, Junior P, Wichtl M. (1980) Isolation and identification of agroclavin and α -dihydrolysergol from the leaves of *Ipomoea fistulosa*. *Planta Medica*, **40**, 328-332.
- [4] Gupta OCD, Gupta R, Gupta PC. (1980) Chemical examination of the flowers of *Ipomoea fistulosa*. *Planta Medica*, **38**, 147-150.
- [5] Haraguchi M, Gorniak SL, Ikeda K, Mikami Y, Kato A, Watson AA, Nash RJ, Molyneux, RJ, Asano N. (2003) Alkaloidal components in the poisonous plant *Ipomoea carnea* (Convolvulaceae). *Journal of Agricultural and Food Chemistry*, **51**, 4995-5000.
- [6] Patel AK, Singh VK, Yadav, RP, Moir, AJG, Jagannadham, MV. (2009) ICChI, a glycosylated chitinase from the latex of *Ipomoea carnea*. *Phytochemistry*, **70**, 1210-1216.
- [7] (a) Reza MS, Khan MOF, Islam MA, Chowdhury AKA. (1994) *In vitro* antimicrobial activity of *Ipomoea fistulosa*. *Fitoterapia*, **65**, 465-466; (b) Chowdhury AKA, Ali MS, Khan MOF. (1997) Antimicrobial activity of *Ipomoea fistulosa* extracts. *Fitoterapia*, **68**, 379-380; (c) Mahmood YAG, Ebrahim MKH, Mageda MA. (2004) Influence of some plant extracts and microbioagents on some physiological traits of faba bean infected with *Botrytis fabae*. *Turkish Journal of Botany*, **28**, 519-528.
- [8] Williams DH, Fleming I. (1988) Spectroscopic methods in organic chemistry. Tata McGraw Hill Publishing Co. Ltd., New Delhi, 1-198.
- [9] (a) Gunatilake AAL, Sultanbawa M, Uvais S. (1973) Chemical investigation of Ceylonese plants. III. Extractives of the fruits of *Argyrea populifolia* (Convolvulaceae). *Journal of the Chemical Society*, **11**, 1155-1157; (b) Dai H, Xiong J, Zhou J, Ding Z. (2000) Chemical constituents from root of *Ipomoea digitata*. *Yunnan Zhiwu Yanjiu*, **22**, 166-168; (c) Ishiguro K, Yoshimoto M, Suzuki M, Yahara S. (2008) Anti-oxidative activity in the lipophilic fraction of sweet potato tubers. *Acta Horticulturae*, **768**, 571-577; (d) Snook ME, Data ES, Kays SJ. (1994) Characterization and quantitation of hexadecyl, octadecyl, and eicosyl esters of *p*-coumaric acid in the vine and root latex of sweet potato [*Ipomoea batatas* (L.)]. *Journal of Agricultural and Food Chemistry*, **42**, 2589-2595; (e) Stevenson PC, Muyinza H, Hall DR, Porter EA, Farman DI, Talwana HM, Robert OM. (2009) Chemical basis for resistance in sweet potato *Ipomoea batatas* to the sweet potato weevil *Cylas puncticollis*. *Pure and Applied Chemistry*, **81**, 141-151.
- [10] Norton RA, Dowd PF. (1996) Effect of stearyl cinnamic acid derivatives from corn bran on *Aspergillus flavus*, corn earworm larvae and dried fruit beetle larvae. *Journal of Agricultural and Food Chemistry*, **44**, 2412-2416.
- [11] Keriko JM, Nakajima S, Baba N, Iwasa J. (1997) Eicosanyl *p*-coumarates from Kenyan plant, *Psiadia punctulata*: plant growth inhibitors. *Bioscience, Biotechnology and Biochemistry*, **61**, 2127-2128.
- [12] Mahmood U, Kaul VK, Acharya R, Jirovetz L. (2003) Sesquiterpene and long chain ester from *Tanacetum longifolium*. *Phytochemistry*, **64**, 851-853.
- [13] Nene YL, Thapliyal PN. (2002) Fungicides in plant disease control. Oxford and IBH Publishing Co. Pvt. Ltd., New Delhi, p. 531.
- [14] Homans AL, Fuchs A. (1970) Direct bioautography on thin layer chromatograms as a method for detecting fungitoxic substances. *Journal of Chromatography*, **51**, 327-329.

New Glucose Esters from the Fresh Leaves of *Jacaranda mimosaeifolia*

Christianah A. Elusiyan^a and Tiwalade A. Olugbade^b

^aDrug Research and Production Unit, Faculty of Pharmacy, Obafemi Awolowo University, Ile-Ife, Osun State, 220005, Nigeria

^bDepartment of Pharmaceutical Chemistry, Faculty of Pharmacy, Obafemi Awolowo University, Ile-Ife, Osun State, 220005, Nigeria

tolugba@oauife.edu.ng

Received: November 21st, 2010; Accepted: August 16th, 2011

From the fresh leaves of *Jacaranda mimosaeifolia* were isolated Phytoquinoids **1-4** established as β -D-glucopyranose 2-benzeneacetate1,6-bis(1-hydroxy-4-oxo-2,5-cyclohexadiene-1-acetate), for which the name Jacaranoside is proposed; β -D-glucopyranose 2-(4-hydroxybenzeneacetate) 1,6-bis(1-hydroxy-4-oxo-2,5-cyclohexadiene-1-acetate), for which the name Jacarandol is proposed; β -D-glucopyranose 2-benzeneacetate 1-(1-hydroxy-4-oxo-2,5-cyclohexadiene-1-acetate) and β -D-glucopyranose 1,6-bis (1-hydroxy-4-oxo-2,5-cyclohexadiene-1-acetate) respectively.

Keywords: *Jacaranda mimosaeifolia*, leaves, glucose esters, jacaranoside, jacarandol, phytoquinoids.

The genus *Jacaranda* (Bignoniaceae) is well known in traditional medicine [1]. *J. mimosaeifolia* D. Don, in particular, has been classified as the World's most widely planted ornamental species [2]. A decoction of the plant is used for the treatment of parasites and for dermatological conditions [3], in the treatment of venereal diseases and for blood purification [4], and an infusion of the bark is applied as a lotion for ulcers [5]. Aqueous extracts of the twigs and seeds of *J. mimosaeifolia* have been reported to show antibacterial properties [6], while the methanol extracts of the leaves showed hypothermic and cardiovascular activities [1]. Antimicrobial activity of the hexane, ethanol and aqueous extract of the leaves has also been reported [7]. Compounds previously isolated from *J. caucana* Pittier include the anticancer agents, jacaranone, its ethyl ester, scutellarein-7-glucuronide, and other triterpenoids with antitumor and cytotoxic activity [8,9]. From the leaves of *J. mimosaeifolia* were isolated jacaranone, phenylacetic- β -glucoside, verbascoside, and jacaranose, with hypotensive and analgesic properties [1,6]. A phenylethanoid dimer, jacaraninoside, was more recently reported from the same species [10].

Compound **1** was isolated as a light yellowish resinous material. The positive ion ESI-MS showed a pseudo-molecular ion at m/z 597 [M-H]⁺, consistent with the molecular formula C₃₀H₂₈O₁₃. The presence of a sugar with acylation at positions 1, 2 and 6 was apparent from the downfield shifts of the corresponding proton signals at δ 5.68, 4.93, and 4.44, 4.21, respectively. The presence of two 1-hydroxy-4-oxo-2,5-cyclohexadien-1-acetyl- (CHD)





Figure 1: Phytoquinoids **1-4** isolated from fresh leaves of *J. mimosaeifolia*.

units was established through the observation of carbon NMR absorptions at δ 184.65 and 184.73 and characteristic *p*-quinoid IR absorptions at 1633.5 and 1673.1 cm⁻¹ [9], with successive links to one methylene group, each observed at δ 44.3 and δ 44.9, and one ester group, each observed at δ 167.1 and δ 168.6. The third acyl group was established as a phenylacetyl unit from the ¹³C NMR spectroscopic data with the corresponding methylene group at δ 40.6 and ester carbonyl at δ 170.1. Compound **1** bears remarkable resemblance to jacaranose [5] in all NMR spectral details except for the non-equivalence of one group of CH₂ protons in **1** ascribable to the CHD unit attached to position 1 of glucose. The AB quartet feature at this position in acetone and methanol was consistent for compounds **1** (δ 2.59, 2.69, J = 14.7 Hz); **2** (δ 2.59, 2.69, J = 14.7 Hz) and **3** (δ 2.50, 2.64, J = 14.7 Hz). Jacaranose was reported in 1988 from the leaves of *J. mimosaeifolia* as a trisubstituted glucose unit with one phenylacetyl moiety and two CHD units. The report gave

Table 1: ^{13}C NMR and DEPT spectral data, for compounds 1, 2, 3 and 4 in acetone- d_6 (75.5 MHz) with assignments supported by COSY, DEPT, HSQC and HMBC.

Position	δ_c ppm							
	1		2		3		4	
Glucose								
1	92.1	CH	92.1	CH	92.3	CH	94.5	CH
2	72.7	CH	72.6	CH	72.9	CH	72.6	CH
3	74.4	CH	74.4	CH	74.6	CH	76.6	CH
4	70.3	CH	70.3	CH	70.2	CH	70.0	CH
5	74.9	CH	74.8	CH	77.7	CH	74.7	CH
6	63.4	CH ₂	63.4	CH ₂	61.2	CH ₂	63.6	CH ₂
4-oxocyclohexadiene acetyl-								
1'	66.8	C	66.8	C	66.9	C	67.0	C
1''	67.0	C	67.0	C	-	-	67.0	C
2'	150.1	CH	150.0	CH	150.2	CH	150.3	CH
6'	150.0	CH	150.2	CH	150.4	CH	150.8	CH
2''	150.5	CH	150.6	CH	-	-	150.5	CH
6''	150.5	CH	150.6	CH	-	-	150.8	CH
3'	127.2	CH	127.2	CH	127.4	CH	127.2	CH
5'	127.4	CH	127.2	CH	127.4	CH	127.4	CH
3''	127.2	CH	127.4	CH	-	-	127.2	CH
5''	127.4	CH	127.4	CH	-	-	127.4	CH
4'	184.7	C	184.8	C	185.0	C	185.1	C
4''	184.7	C	184.9	C	-	-	185.1	C
7'	44.3	CH ₂	44.3	CH ₂	44.4	CH ₂	44.7	CH ₂
7''	44.9	CH ₂	44.9	CH ₂	-	-	44.9	CH ₂
8'	167.1	C	167.1	C	167.1	C	167.6	C
8''	168.6	C	168.6	C	-	-	168.7	C
Arylacetyl-								
1'''	-	-	125.0	C	-	-	-	-
1''''	134.4	C	-	-	134.5	C	-	-
2'''	-	-	130.3x2	CHx2	-	-	-	-
2''''	129.3x2	CHx2	-	-	129.3x2	CHx2	-	-
3'''	-	-	115.2x2	CHx2	-	-	-	-
3''''	128.4x2	CHx2	-	-	128.4x2	CHx2	-	-
4'''	-	-	156.4	C	-	-	-	-
4''''	126.9	CH	-	-	126.8	CH	-	-
7'''	-	-	39.8	CH ₂	-	-	-	-
7''''	40.6	CH ₂	-	-	40.6	CH ₂	-	-
8'''	-	-	170.6	C	-	-	-	-
8''''	170.1	C	-	-	170.1	C	-	-

**Figure 2:** HMBC Correlation of compound 1.

all singlets for the three CH₂ groups in acetone. Although the positions of esterification were reported as positions 1, 2, and 6 of β -glucose, the precise location of the different acyl groups was not established, hence the complete structure of jacaranose remained undetermined.

In the present study, the definite sites of esterification on the glucose molecule of compound 1 were, however, made possible by the combination of COSY, ^1H , ^{13}C , HSQC and HMBC experiments. The phenylacetyl group was located at C-2 of glucose on the basis of the downfield signal observed for H-2 (δ 4.92) and its HMBC correlation to the corresponding ester carbonyl (δ 170.1). A prominent HMBC correlation of the anomeric proton to the carbonyl carbon, C-8' (δ 167.1), established an acylation at the anomeric C-1 position by one CHD unit. The HMBC correlation observed between position 6 protons of glucose

δ_{H6a} 4.44 (dd, $J = 2.1, 12.3$) and δ_{H6b} 4.21 (dd, $J = 5.7, 12.3$) with an ester carbonyl carbon C-8'' (δ 168.5), indicated that the second unit of the CHD moiety was attached at C-6 of glucose. Therefore, the phytoquinoid 1 was established as β -D-glucopyranose 2-benzeneacetate 1,6-bis(1-hydroxy-4-oxo-2,5 cyclohexadiene-1-acetate). Although compound 1 has a similar optical rotation $\{[\alpha]_{\text{D}}^{25} + 6.5$ (c 0.05, MeOH)} as that reported for jacaranose $\{[\alpha]_{\text{D}}^{20} + 6.8$ (c 0.9, MeOH)}, a repeat collection of plant material gave compound 1 with no isomer isolated showing all three singlet methylene groups as previously described for jacaranose [5]. The ABq feature at ambient temperature was consistent at 200, 300 and 500 MHz in acetone and methanol. We propose the name jacaranoside for compound 1.

Compound 2 shows significant IR and NMR similarities to compound 1. Thus, two CHD units at positions 1 and 6 of glucose were readily established with the major difference in the acyl group present at position 2. The presence of an extra oxygen ($[\text{M}+\text{H}]^+ = 613$), a phenolic reaction to ferric chloride, a broad down-field signal in acetone at δ 8.4, a p -substituted aromatic ring with an ortho- coupled proton at 6.77 (d, $J = 8.4$) and a quaternary carbon at δ 156 established the presence of a p -hydroxy phenylacetyl group instead of the phenylacetyl group found in compound 1. HMBC correlation of H-2 of glucose with the corresponding ester carbonyl at δ 170.1 and a methylene at δ 39.8 confirmed this location. Thus, compound 2, for which the name jacarandol is proposed, was established as β -D-glucopyranose 2-(4-hydroxy benzeneacetate) 1,6-bis(1-hydroxy-4-oxo-2,5-cyclohexadiene-1-acetate).

Compound 3 showed the presence of only two acyl groups (ester C=O at δ 167.1 and 170.1). In compound 3, the lack of acylation at position 6 of the sugar is apparent from the relative upfield shift of the protons (δ 3.78 and δ 3.52). An ABq (δ 2.50, 2.64, $J = 14.7$ Hz) was observed for the CH₂ of the CHD unit attached to position 1 of the sugar, as for compounds 1 and 2, while evidence for the presence of phenylacetyl substitution at position 2 was apparent from the downfield shift of sugar protons at position 2 and HMBC correlations. EI-MS did not show the molecular ion, but showed ions diagnostic of a CHD residue (m/z 150 and 122) [9] and a phenylacetyl group (m/z 136, PhCH_2COOH).

Compound 4 showed the presence of only two acyl groups (ester C=O at δ 167.6 and 168.7). The evidence for two CHD units is borne out from two ketonic carbon resonances at δ 185.09 and 185.13. The lack of an ABq for the CHD unit on position 1 of the sugar is consistent with the lack of acylation at position 2, which showed an upfield proton shift (δ 3.40). On the other hand, the presence of the second CHD unit at position 6 of the sugar is evident from the downfield shift of the sugar protons at 6 (δ 4.03 and 4.16) and corroborated the HMBC correlations. The EI-MS showed CHD residues at m/z 150 and 122.

Table 2: ^1H NMR spectral data for compounds **1-4** (δ in ppm, acetone- d_6)

H	δ , ppm			
	1	2	3	4
Glucose				
1	5.68, 1H, d, (8.4)	5.66, 1H, d, (8.4)	5.64, 1H, d, (8.4)	5.50, 1H, d, (8.1)
2	4.93, 4.90, 1H, dd, (8.4; 8.4)	4.89, t, (8.4)	4.88, 1H, t, (9.0)	3.40
3	3.75, 1H, t, (9.6)	3.75 (obs)	3.74 (obs)	3.50 (obs)
4	3.52, 1H, dd, (9.3, 9.3)	3.45 (obs)	3.60 (obs)	3.40 (obs)
5	3.70 (obs)	3.70 (obs)	3.50, m	3.60 (obs)
6a	4.44, dd, (2.1, 12.3)	4.43, d, (12.0)	3.78, dd, (12.9)	4.03, dd, (1.8; 12.0)
6b	4.21, dd, (5.7, 12.3)	4.20, dd, (5.7, 12.0)	3.52 (obs)	4.16, dd, (6.0; 12.0)
4-oxocyclohexadiene acetyl-				
2''	7.00, m	7.00, m	7.07, 7.04, dd, (10.3; 3.3)	-
2'''	7.00, m	7.00, m	-	-
6''	7.00, m	7.00, m	6.99, 7.03, dd, (10.2; 3.3)	-
6'''	7.00, m	7.00, m	-	-
3''	6.10, m	6.10, m	6.10, d, (10.2)	-
3'''	6.10, m	6.10, m	-	-
5''	6.10, m	6.10, m	6.10, d, (10.2)	-
5'''	6.10, m	6.10, m	-	-
7''	ABq, 2.59, 2.69, (14.7)	ABq, 2.59, 2.69, (14.7)	ABq, 2.50, 2.64, (14.7)	2.90, 2H, s
7'''	2.81, 2H, s	2.81, 2H, s	-	2.80, 2H, s
Arylacetyl-				
2''', 6'''	-	7.14, d, (8.4)	-	-
3''', 5'''	-	6.77, d, (8.4)	-	-
2''', 6''''	7.30, m	-	7.29, m	-
3''', 5''''	7.30, m	-	7.29, m	-
4''''	7.30, m	-	7.29, m	-
7''''	-	3.56, 2H, s	-	-
7'''''	3.68, 2H, s	-	3.67, 2H, s	-

Experimental

General experimental procedures: The IR spectra (V_{\max}) were determined on a Perkin–Elmer 2000 FT-IR Spectrophotometer. ESI-MS (negative ion mode) were recorded on an Agilent 1100 series LC/MSD Trap mass spectrometer. NMR measurements were recorded on Varian 200, 500 and Bruker 300 spectrometers. Silica gel 60 (230–400 μm , Merck) was used for vacuum liquid (VLC) and accelerated gradient (AGC) chromatographic separations. Final purifications were performed by medium pressure reverse phase liquid chromatography (MPLC) using a pre-packed Lichroprep RP-18 (40–63 μm) column. All fractions were collected in 15 mL test tubes. Pre-coated silica gel 60 F₂₅₄ aluminum plates (0.2 mm, Merck) and pre-coated silica gel 60 RP-18 F₂₅₄S aluminum TLC plates (Merck) were used for TLC monitoring. TLC mobile phases were 50% methanol and toluene: acetone: water (10:30:1) for reverse phase and normal phase, respectively. TLC plates were visualized by UV light, 1% ferric chloride in methanol, and 1% vanillin in H₂SO₄, followed by heating at 105°C for 1–2 min.

Plant material: The leaves of *J. mimosaeifolia* D. Don, were collected in May 2007 from the Obafemi Awolowo University campus and identified at the Department of Botany. A voucher sample is lodged in the herbarium of the University as no. 16412.

Extraction and isolation: Fresh leaves (950 g) were immediately macerated in ethanol (4L) for 24 h at room temperature. After filtration, the extract was evaporated to complete dryness *in vacuo* at 40°C, yielding a crude ethanolic leaf extract (JL, 71 g). This was fractionated by VLC on silica gel using EtOAc: MeOH mixtures in

gradient to give 5 fractions (Fr 1 = 35.5 g; Fr 2 = 6.14 g; Fr 3 = 2.08 g; Fr 4 = 2.10 g and Fr 5 = 1.06 g). Fraction 1 (16.0 g) was subjected to repeated AGC separation on silica using *n*-Hex: EtOAc: MeOH in gradient to yield 6 sub-fractions (1a–f). Sub-fraction 1d (1.1 g), which eluted from EtOAc: MeOH (9.5: 0.5), was subjected to final purification on reverse phase MPLC using a H₂O: MeOH gradient. Elution with 45% MeOH afforded compound **1** (232 mg). Final purification of sub-fraction 1c (2.4 g), which eluted from EtOAc: MeOH (9.0: 1.0), on reverse phase MPLC, afforded compound **2** (508 mg), eluted with 35% MeOH, compound **3** (63 mg) eluted with 30% MeOH and compound **4** (588 mg) eluted with 45% MeOH. Sub-fraction 1b (1.1 g), which eluted from EtOAc: MeOH (8.5:1.5), was similarly purified by reverse phase MPLC using MeOH: H₂O gradient. Elution with 25% and 40% MeOH afforded compound **4** (120mg) and verbascoside (141 mg), respectively.

Compound 1 [β -D-glucopyranose-2-benzeneacetate1,6-bis(1-hydroxy-4-oxo-2,5-cyclohexadiene-1-acetate)]

Light yellow resinous material.

FTIR: 3396.7, 3440.2, 1673.1, 1633.5 cm^{-1}

$[\alpha]_{25}^D$: +6.5 (*c* 0.05, MeOH).

R_f : 0.34 (MeOH: H₂O, 1:1)

^1H NMR and ^{13}C NMR (acetone- d_6): Tables 1 and 2

ESI-MS: m/z 597 $[\text{M}-\text{H}]^-$, 447

Yield: 232 mg, 1.45 %

Compound 2 [β -D-glucopyranose2-(4-hydroxy benzene acetate) 1,6-bis (1-hydroxy-4-oxo-2,5-cyclohexadiene-1-acetate)]

Brownish resinous compound

FTIR: 3395.1, 3445.6, 1672.7, 1633.4 cm^{-1}

$[\alpha]_{25}^D$: +9.8 (*c* 0.06, MeOH).

R_f : 0.54 (MeOH: H₂O, 1:1)

¹H NMR and ¹³C NMR (acetone-*d*₆): Tables 1 and 2.

ESI-MS: *m/z* 613 [M-H]⁺, 463

Yield: 508 mg, 3.18 %

Compound 3 [β-D-glucopyranose 2-benzeneacetate 1-(1-hydroxy-4-oxo-2,5-cyclohexadiene-1-acetate)]

Brown gum.

R_f : 0.36 (MeOH: H₂O, 1:1)

¹H NMR and ¹³C NMR (acetone-*d*₆): Tables 1 and 2.

ESI-MS: *m/z* 150 (62), 136 (100), 122 (65)

Yield: 63 mg, 0.39%

Compound 4 [β-D-glucopyranose 1,6-bis(1-hydroxy-4-oxo-2,5-cyclohexadiene-1-acetate)]

Brown gum

R_f : 0.76 (MeOH: H₂O, 1:1)

¹H NMR and ¹³C NMR (acetone-*d*₆): Tables 1 and 2.

ESI-MS: *m/z* 150 (90), 122 (100)

Yield: 708 mg, 4.43 %

Verbascoside

R_f : 0.56 (MeOH: H₂O, 1:1)

NMR data were identical to those reported in literature [11]

Yield: 141 mg, 0.88%

Acknowledgments - The authors acknowledge the support of the Carnegie Corporation of New York for OAU Central Science Laboratory, Prof. B. Abegaz, NABSA, Gaborone, for HMBC experiments and EI MS and Rolf Blattler and Uta Wollrab of Varian Inc. for 500 NMR spectra.

References

- [1] Nicasio P, Meckes M. (2005) Hypotensive effect of the hydroalcoholic extract from *Jacaranda mimosaeifolia* leaves in rat. *Journal of Ethnopharmacology*, **97**, 301-304.
- [2] Gentry AH. (1992) A synopsis of Bignoniaceae ethnobotany and economic botany. *Annals of the Missouri Botanical Garden*, **79**, 53-64.
- [3] Tene V, Malagon O, Finzi PV, Vidari G, Armijos C, Zaragoza T. (2007) An ethnobotanical survey of medicinal plants used in Loja and Zamara-Chinchipe, Ecuador. *Journal of Ethnopharmacology*, **111**, 63-81.
- [4] Gachet MS, Schuhly W. (2009) Jacaranda – An ethnopharmacology and phytochemical review. *Journal of Ethnopharmacology*, **121**, 14-27.
- [5] Gambaro V, Garbarino JA, Galeffi C, Nicoletti M, Messana I, Marini-Bettolo GB. (1988) Constituents of *Jacaranda mimosaeifolia*. *Revista Latinoamericana de Quimica*, **19**, 17-19.
- [6] Mahran GH, El-Fishwy AM, Hosny AMS, Hilal AM (1991) Phytochemical and antibacterial study of *Jacaranda mimosaeifolia* D. Don grown in Egypt. *Herba Hungarica*, **30**, 98-104.
- [7] Rojas JJ, Ochoa VJ, Ocampo SA, Munoz JF. (2006) Screening for antimicrobial activity of ten medicinal plants used in Colombian folkloric medicine: a possible alternative in the treatment of non-nosocomial infections. *BioMed Central Complementry and Alternative Medicine*, **6**, 1-6.
- [8] Ogura M, Cordell GA, Farnsworth NR. (1976) Potential anticancer agents. III. Jacaranone, a novel phytoquinoid from *Jacaranda caucana*. *Lloydia*, **39**, 255-257.
- [9] Ogura M, Cordell GA, Farnsworth NR. (1977) Potential anticancer agents. IV. Constituents of *Jacaranda caucana* Pittier (Bignoniaceae). *Lloydia*, **40**, 157-168.
- [10] Moharram FA, Marzouk MSA. (2007) A novel phenylethanoid dimer and flavonoids from *Jacaranda mimosaeifolia*. *Zeitschrift fur Naturforschung B*, **62**, 1213-1220.
- [11] Franzyk H, Husum TL, Jensen SR (1998) A caffeoyl phenyl-ethanoid glycoside from *Plantago myosuros*. *Phytochemistry*, **47**, 1161-1162.

Shamiminol: A New Aromatic Glycoside from the Stem Bark of *Bombax ceiba*

Shaheen Faizi^{a,*}, Sadia Zikr-Ur-Rehman^a and Muhammad Ali Versiani^{b,†}

^aInternational Center for Chemical and Biological Sciences, HEJ Research Institute of Chemistry, University of Karachi, Karachi-75270, Pakistan

^bDepartment of Chemistry, Federal Urdu University of Arts, Sciences and Technology, Gulshan-e-Iqbal Campus, Karachi-75300, Pakistan

shaheenfaizi@hotmail.com; [†]mali.versiani@fuuast.edu.pk

Dedicated to the fond memory of Professor Salimuzzaman Siddiqui *FRS*, (1897–1994), the founding director of HEJ Research Institute of Chemistry, University of Karachi, Karachi, Pakistan

Received: July 30th, 2011; Accepted: October 18th, 2011

A new aromatic glycoside, shamiminol was isolated from the stem bark of *Bombax ceiba* along with the known constituents stigmasta-3,5-diene, lupenone, (±)-lyoniresinol 2a-*O*-β-D-glucopyranoside and opuntiol, obtained for the first time from this plant. The structure of shamiminol was elucidated on the basis of extensive 1D- and 2D-NMR spectroscopic and mass spectrometric studies as 3,4,5-trimethoxyphenol 1-*O*-β-D-xylopyranosyl-(1→2)-β-D-glucopyranoside (**1**).

Keywords: *Bombax cieba*, Bombacaceae, aromatic glycoside, shamiminol.

Bombax cieba Linn. (Syn. *B. malabaricum* D.C. and *Salmalia malabarica* D.C.) commonly known as simbal or red silk-cotton tree, belongs to the family Bombacaceae. It grows in the hotter forest regions of India, Sri Lanka, Malayasia, China, Java and Sumatra. In Pakistan, it is found in Hazara and other parts of the North West frontier province, and also in Sindh province [1]. Various parts of the plant are highly reputed in traditional medicine for the treatment of a variety of ailments [1]. This medicinal plant has immense therapeutic importance, and pharmacological studies showed that it possesses hypoglycemic, hypotensive, antiangiogenic, analgesic, antioxidant, vibriocidal, antimicrobial, anti-inflammatory and antidiabetic activities [2]. Phytochemical investigation of *B. ceiba* has resulted in the isolation of a number of constituents belonging to different classes of compounds [3]. In this communication, we report the isolation and structure elucidation of a new natural product, shamiminol (**1**), along with four known compounds, stigmasta-3,5-diene [4], lupenone [5], (±)-lyoniresinol 2a-*O*-β-D-glucopyranoside [6] and opuntiol [7], which were obtained for the first time from *B. ceiba*.

Shamiminol (**1**) was obtained as a colorless gum from the stem bark of *B. ceiba*. Its IR spectrum indicated the presence of a hydroxyl group (3405 cm⁻¹) and an aromatic ring (1595 and 1507 cm⁻¹), while the UV spectrum showed absorption at 215 and 265 nm, indicative of its aromatic





Figure 1: 3,4,5-Trimethoxyphenol 1-*O*-β-D-xylo-pyranoside-(1→2)-β-D-glucopyranoside [shamiminol (**1**)].

nature. Shamiminol (**1**) has the composition C₂₀H₃₀O₁₃, MW = 478, as derived from positive and negative FABMS and ¹H and ¹³C NMR spectroscopic data (Table 1). It has twenty carbons in the ¹³C NMR spectrum (three methyls, two methylenes, nine methines, two sp² CH, and four sp² quaternary carbons).

The ¹H NMR spectrum of **1** in CD₃OD (Table 1) showed the presence of two equivalent methoxy groups (δ 3.80, 6H, s) and two aromatic protons (δ 6.45, s), along with another methoxy group at δ 3.69, as a three protons singlet, suggesting that the molecule contained an aromatic ring with a symmetrical substitution pattern [8] (Figure 1). The presence of the 3,4,5-trimethoxyphenol system was supported by the ¹³C NMR spectrum which showed signals for three aryl methoxy groups at δ 56.62 (3- and 5-OCH₃) and 61.23 (4-OCH₃), four quaternary carbons at δ 155.93


Table 1: NMR spectroscopic characteristics [δ (ppm) and J (Hz)] of shamiminol (**1**)

Assignments	δ_c	δ_H
1	155.93	
2	96.39	6.45 (s)
3	154.78	
4	134.01	
5	154.78	
6	96.39	6.45 (s)
1'	102.09	4.92 (d, 7.0)
2'	83.74	3.64 (m)
3'	77.89	3.36 (t, 8.7)
4'	71.38	3.25 (m)
5'	78.18	3.46 (ddd, 1.6, 4.0, 9.4)
6 _a '	62.63	3.90 (dd, 1.6, 11.1)
6 _b '	62.63	3.68 (dd, 4.0, 11.1)
1''	106.28	4.62 (d, 7.4)
2''	75.76	3.21 (t, 7.4)
3''	77.39	3.62 (t, 7.4)
4''	71.12	3.49 (m)
5 _a ''	67.24	3.85 (dd, 5.9, 10.9)
5 _b ''	67.24	3.25 (m)
3-OCH ₃	56.62	3.80 (s)
4-OCH ₃	61.23	3.69 (s)
5-OCH ₃	56.62	3.80 (s)
OH		

**Figure 2:** Important HMBC correlations of shamiminol (**1**).

(C-1), 154.78 (C-3 and 5) and 134.01 (C-4), and two methine carbons at δ 96.39 (C-2 and 6). The HSQC spectrum showed one bond correlation of H-2,6 at δ 6.45 with C-2,6, while the HMBC spectrum showed 3J correlation of 3- and 5-OCH₃ protons (δ 3.80) with C-3 and C-5, respectively; 4-OCH₃ protons (δ 3.69) showed correlation with C-4; H-2,6 with C-4, which also showed 2J connectivities with C-1, C-3 and C-5 (Figure 2). These spectral data confirmed that a 3,4,5-trimethoxyphenol moiety was present in the molecule [8–10], which was corroborated by the fragment ions at m/z 184 (C₉H₁₂O₄) and 167 (C₉H₁₁O₃) in the EIMS.

The presence of a disaccharide moiety was also revealed by the 1H NMR spectrum, which exhibited two anomeric proton signals at δ 4.92 (d, J = 7.0, H-1') and 4.62 (d, J = 7.4, H-1'') attributable to those of glucose and xylose units, respectively [11]. The spectrum analyzed with the aid of 1H - 1H shift correlation spectroscopy (COSY) further exhibited signals due to four methines at δ 3.64 (m, H-2'), 3.36 (t, J = 8.7, H-3'), 3.25 (m, H-4'), and 3.46 (ddd, J = 1.6, 4.0, 9.4, H-5'), and one oxymethylene at δ 3.90 (dd, 1.6, 11.1, H-6_a') and 3.68 (dd, J = 4.0, 11.1, H-6_b'), which were ascribed to glucose protons. The ^{13}C NMR spectrum also showed signals corresponding to the sugar unit. In the HMQC plot, chemical shift values at δ 102.09 (C-1'),

**Figure 3:** Important NOESY correlations for shamiminol (**1**).

83.74 (C-2'), 77.89 (C-3'), 71.38 (C-4'), 78.18 (C-5') and 62.63 (C-6') showed direct one bond correlations with H-1', H-2', H-3', H-4', H-5' and H-6', respectively (Table 1).

Additionally, the 1H NMR spectrum exhibited two multiplets at δ 3.49 and 3.25, two triplets at δ 3.21 and 3.62 with the same coupling constant of 7.4 Hz and a double doublet at δ 3.85 (J = 5.9, 10.9) which were assigned to H-4'', H-5_b'', H-2'', H-3'' and H-5_a'', respectively, of the xylose moiety. In the COSY-45° plot the anomeric proton of xylose at δ 4.62 (H-1'') correlated with H-2'', H-3'' with H-4'' and H-2'', and H-4'' with H-5_a'', H-5_b'', and H-3'', while in the HMQC spectrum H-1'', H-2'', H-3'', H-4'' and H-5'' showed direct connectivity with δ 106.28 (C-1''), 75.76 (C-2''), 77.39 (C-3''), 71.12 (C-4'') and 67.24 (C-5''), respectively.

The glycosidic linkages in both glucose and xylose was determined to be β , based on the large coupling constants of the respective anomeric proton signals (7.0 and 7.4 Hz) [11]. That the glucose is attached *via* its anomeric carbon with the oxygen of the trimethoxy aromatic ring was indicated by the high frequency chemical shift of the anomeric proton (δ 4.92) and confirmed by the long-range correlation between the anomeric proton (δ 4.92) and C-1 (δ 155.93) of the benzene ring in the HMBC spectrum (Figure 2). It was reaffirmed by the cross peak observed in the NOESY-plot for the spatial connectivity of the anomeric proton with the aromatic H-2,6 (δ 6.45) (Figure 3).

The ^{13}C NMR spectral data suggested the attachment of xylose with the C-2' of the glucose residue, as it demonstrated marked glycosidation shifts, showing an α -effect on C-2' and β -effects on C-1' and C-3'. Thus C-2' resonated at higher frequency (δ 83.74), and C-1' and C-3' appeared at δ 102.09 and 77.89, respectively. A literature search revealed that when a sugar (xylose or apiose) has a linkage with the C-6 of glucose, that carbon showed a marked α glycosidation shift, while C-5 resonated at a low frequency [9,11a]. When the same sugar was located on C-2 of glucose then this carbon resonated at high frequency, showing the α -effect [10,11b]. Consequently, the disaccharide moiety was identified as β -D-xylopyranosyl-(1 \rightarrow 2)- β -D-glucopyranosyl, which was

supported by the mass fragments at m/z 185 and 183 in the positive and negative ion FABMS, respectively, which arose due to the loss of glucose and xylose moieties from the molecular ion. This was confirmed by the 2D NMR spectral data. Thus, in the HMBC spectrum, H-2' showed crucial long-range coupling with the anomeric carbon (C-1'') of xylose and H-1'' in turn has connectivity with the C-2' of glucose (Figure 2). The NOESY spectrum exhibited an important cross peak for the spatial proximity of H-2' of glucose with H-1'' of xylose (Figure 3).

All the evidence discussed above led to the formulation of shamiminol as 3,4,5-trimethoxyphenol 1-*O*- β -D-xylopranosyl-(1 \rightarrow 2)- β -D-glucopyranoside (**1**), which was corroborated by the important fragment ions in the positive and negative ion FABMS at m/z 461 ($M^+ + 1 - H_2O$) and 459 ($M^+ - 1 - H_2O$), and at m/z 364 ($M^+ - \text{xylose}$), 432 ($M^+ - 3CH_3 - H$), 414 and 137 in the EIMS. The complete 1H and ^{13}C NMR spectral assignments for **1** has been made through the help of 2D NMR spectroscopy. Furthermore, the 1H NMR spectrum of **1**, in C_5D_5N , showed no resonances for the aromatic hydroxyls. However, it exhibited a broad hump at δ 5.20 for the sugar hydroxyprotons, which disappeared on shaking with D_2O . Additionally, all the chemical shifts of the compound shifted to high frequency in C_5D_5N as compared with those in CD_3OD , except for those of 3- OCH_3 and 5- OCH_3 , which resonated at low frequency, and 4- OCH_3 , which appeared at high frequency (Table 1). It is important to mention that the regioisomer of **1**, 3,4,5-trimethoxyphenyl 1-*O*- β -D-xylopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside, has been reported earlier from *Ailanthus integrifolia* [11a].

Stigmasta-3,5-diene [4], lupenone [5], (\pm)-lyoniresinol 2a-*O*- β -D-glucopyranoside [6] and opuntiol [7] were also isolated from the stem bark and identified by comparison with reported data. The literature search revealed that these compounds are mostly found in the stem bark of plants [4–7]. Moreover, *Cinnamomum cassia* (family Lauraceae) and *B. ceiba*, belonging to two different families, contained glucosides of lyoniresinol and 3,4,5-trimethoxyphenol, as well as catechin derivatives in their stem bark [9,2c].

Experimental

General: The IR (in KBr disc) spectrum was recorded on a JASCO A 302 spectrophotometer, and the EI and FAB mass spectra were measured on Finnigan MAT 112 and JMS HX-110 spectrometers. The 1H NMR spectra were run in CD_3OD and C_5D_5N on Bruker Aspect AM-500 and AM-400 spectrometers operating at 500 and 400 MHz, respectively, while ^{13}C NMR spectra were recorded at 125 MHz. Purity of the compounds was checked on silica gel 60GF₂₅₄ precoated cards (0.2 mm thickness), while for flash column chromatography (FCC) (Model Aldrich), silica gel 9385 (E. Merk) was used. For vacuum liquid chromatography (VLC), silica gel 60 GF₂₅₄ was used. Recycling HPLC (LC-908w): Semipreparative ODS (C4

column) used a mixed solvent of MeOH/ H_2O (1:1) at a flow rate of 4.0 mL/min.

Plant material: The stem bark of *B. ceiba* was collected in February 2000 from the Karachi University Campus. The plant was authenticated by Prof. Dr Surrayya Khatoon of the Department of Botany, University of Karachi and a voucher specimen (No. 66854 KUH) was deposited in the same department.

Extraction and isolation of chemical constituents: Fresh, uncrushed and undried stem bark (2.5 kg) of *B. ceiba* was extracted twice with light petroleum at room temperature. The extracts were combined and freed of the solvent *in vacuo* to a residue (BCBP, 5 g), which on treatment with methanol, followed by filtration, gave soluble (BCBPM, 3.47 g) and insoluble (BCBPMX) fractions. The filtrate (BCBPM, 30 mg) was subjected to preparative thin layer chromatography (PTLC) over silica gel (light petroleum: EtOAc, 7.5:2.5), which afforded 10 bands (BCBPM1–10). Spectral studies (mass, peak matching and 1H NMR) on bands 2 and 4 revealed their structures as stigmasta-3,5-diene [4] and lupenone [5].

The marc left after light petroleum extraction was extracted with methanol 3 times. The combined methanolic extracts were freed of the solvent *in vacuo* to give a thickish mass (BCBM) in which insoluble matter settled down on keeping in the cold. This was removed by filtration to give soluble (BCBMM, 36.5 g) and insoluble (BCBMI, 4.50 g) fractions. The former (BCBMM, 36.4 g) was subjected to VLC (silica gel 60 GF₂₅₄, light petroleum, EtOAc, MeOH and H_2O in order of increasing polarity by 10%, which gave 38 fractions). Fractions 1–13 (light petroleum 100% – EtOAc 100%), 14–17 (MeOH:EtOAc, 1:9 – MeOH:EtOAc, 4:6), 18–21 (MeOH:EtOAc, 1:1 – MeOH:EtOAc, 8:2), 22–26 (MeOH:EtOAc, 9:1 – MeOH, 100%) and 27–38 (MeOH: H_2O , 9:1 – H_2O , 100%) were combined on the basis of TLC.

Fractions 14–17 (3.0 g) were subjected to FCC (silica gel 9385, light petroleum, EtOAc, MeOH and H_2O in order of increasing polarity) affording 64 fractions. Fractions 28–32 (EtOAc:MeOH, 7.5:2.5 – EtOAc:MeOH, 5.5:4.5) were purified through FCC (silica gel 9385, light petroleum, $CHCl_3$ and MeOH in order of increasing polarity), affording 43 fractions. FCC fractions S-11 ($CHCl_3$: MeOH, 8:2, 252 mg) and S-12 ($CHCl_3$:MeOH, 7.75:2.25, 21.0 mg) were separately subjected to recycling HPLC semi-preparative ODS (C4 column), eluted with (MeOH: H_2O , 1:1) at a flow rate of 4.0 mL/min. S-11 yielded 2 fractions S11-HP1 (15.0 mg, R_t 24 min.) and S11-HP2 (4.0 mg, R_t 18 min.), which were collected and evaporated to dryness. Spectral studies, along with 2D NMR data, disclosed that S11-HP1 was a mixture of 2 known isomers of lyoniresinol [6]. Eluate S11-HP2 was characterized as a known compound, opuntiol [7]. S-12 also gave 3 sub-fractions in which only S-12HP2 (5.0 mg,

R_t 22 min.) was found to be pure and characterized as a new compound, shamiminol (1).

Shamiminol [3,4,5-trimethoxyphenol 1-O-β-D-xylopyranosyl-(1→2)-β-D-glucopyranoside] (1)

Colorless gum.

[α]_D²⁴: -25.20 (c 0.016, CH₃OH)

IR ν_{max} (KBr) cm⁻¹: 3405, 1595, 1507, 1415, 1165.

UV λ_{max} (MeOH) nm: 215, 265.

¹H and ¹³C NMR: Table 1.

HR FAB MS (+ve) *m/z*: 479.4522 (M⁺+1, calcd. for C₂₀H₃₁O₁₃, 479.4583), 461.4370 (M⁺+1-H₂O, calcd. for C₂₀H₂₉O₁₂, 461.4431), 185.1981 (M⁺+1-glucose-xylose, calcd. for C₉H₁₃O₄, 185.1998).

HR FAB MS (-ve) *m/z*: 477.4363 (M⁺-1, calcd. for C₂₀H₂₉O₁₃, 477.4425), 459.4259 (M⁺-1-H₂O, calcd. for

C₂₀H₂₇O₁₂, 459.4272), 183.1829 (M⁺-1-glucose-xylose, calcd. for C₉H₁₁O₄, 183.1839).

HR EIMS *m/z* (rel. intensity, %): 432.0863 (M⁺-3CH₃-H, calcd. for C₁₇H₂₀O₁₃, 432.0903, 57), 346.1260 (M⁺-xylose, C₁₅H₂₂O₉, 15), 184.0703 (M⁺-glucose-xylose, C₉H₁₂O₄, 35).

EIMS *m/z* (rel. intensity, %): 432 (C₁₇H₂₀O₁₃, 2), 414 (C₁₇H₁₈O₁₂, 5), 346 (C₁₅H₂₂O₉, M⁺-xylose, 2), 311 (C₁₁H₁₉O₁₀, 10), 295 (C₁₁H₁₉O₉, 35), 184 (C₉H₁₂O₄, M⁺-glucose-xylose, 51), 182 (72), 167 (C₉H₁₁O₃, 72), 154 (74), 137 (100), 124 (C₇H₈O₂, 45) 107 (C₇H₇O, 55).

Acknowledgments - One of the authors (Sadia Zikr-Ur-Rehman) is grateful to the Higher Education Commission Islamabad, Pakistan for financial support through the Merit Scholarship Scheme for Ph.D. studies in Science and Technology (200 Scholarships).

References

- [1] (a) Nadkarni KM, Nadkarni AK. (1976) *Indian Materia Medica*, Nadkarni KM (ed.), Popular Parkashan, Bombay, 1; (b) Baquar SR. (1989) *Medicinal and Poisonous Plants of Pakistan*, Printas Karachi (Pakistan), 397; (c) Perry LM, Metzgar J. (1980) *Medicinal Plants of East and Southern Asia*, MIT Press, London, 59.
- [2] (a) Saleem R, Ahmad M, Hussain SA, Qazi AM, Ahmad SI, Qazi MH, Ali M, Faizi S, Akthar S, Husnain SN. (1999) Hypotensive, hypoglycaemic and toxicological studies on the flavonol C-glycoside shamimin from *Bombax ceiba*. *Planta Medica*, **65**, 331-334; (b) You YJ, Nam NH, Kim Y, Bae KH, Ahn BZ. (2003) Antiangiogenic activity of lupeol from *Bombax ceiba*. *Phytotherapy Research*, **17**, 341-344; (c) Saleem R, Ahmed SI, Faizi Z, Zikr-Ur-Rehman S, Ali M, Faizi S. (2003) Hypotensive activity and toxicology of constituents from *Bombax cieba* stem bark, *Biological and Pharmaceutical Bulletin*, **26**, 41-46; (d) Dar A, Faizi S, Naqvi S, Roome T, Zikr-Ur-Rehman S, Ali M, Firdous S, Moin ST. (2005) Analgesic and antioxidant activity of mangiferin and its derivatives: the structure activity relationship. *Biological and Pharmaceutical Bulletin*, **28**, 596-600; (e) Shuping L. (2007) New formulation of traditional Chinese medicine for treating burn. *Faming Zhangli Shenqing Gongkai Shuomingshu* CN 101,032,577 (Cl. A61K36/82), (2007), Appl. 10,027,354, 30 Mar 2007: 4 pp. (Ch); (f) Bapat VR. (2007) A novel ayurvedic drug for treatment of different type of vaginitis. (India) *Indian Patent Application* IN 2007 CH 1,511 (Cl. A61K31/573), 17 Aug 2007, Appl. 2007/CH1,511,13 Jul 2007:23 pp. (Eng); (g) Debasis DK, Ali KM, Mandal S, Barik B, Ghosh D. (2010) Antidiabetic and antioxidative effects of hydro-methanolic extract of sepals of *Salmalia malabarica* in streptozotocin induced diabetic rats. *Journal of Applied Biomedicine*, **8**, 23-33.
- [3] (a) Shahat AA, Hassan RA, Nazif NM, Miert SV, Pieters L, Hammuda FM, Vlietinck AJ. (2003) Isolation of mangiferin from *Bombax malabaricum* and structure revision of shamimin. *Planta Medica*, **69**, 1066-1068; (b) Faizi S, Zikr-Ur-Rehman S, Ali M, Naz A. (2006) Temperature and solvent dependent NMR studies on mangiferin and complete NMR spectral assignments of its acyl and methyl derivatives. *Magnetic Resonance in Chemistry*, **44**, 838-844; (c) Puckhaber LS, Stipanovic RD. (2001) Revised structure for a sesquiterpene lactone from *Bombax malabaricum*. *Journal of Natural Products*, **64**, 260-261; (d) Zhang X, Zhu H, Zhang S, Yu Q, Xuan L. (2007) Sesquiterpenoids from *Bombax malabaricum*. *Journal of Natural Products*, **70**, 1526-1528; (e) Singh P, Mewara DK, Sharma MC. (2008) A new ferulic ester and related compounds from *Bombax malabaricum* DC. *Natural Product Communications*, **3**, 223-225; (f) Wu J, Zhang XH, Zhang SW, Xuan LJ. (2008) Three novel compounds from the flowers of *Bombax malabaricum*, *Helvetica Chimica Acta*, **91**, 136-143; (g) Sampath KN, Madhurambal G. (2010) Quercetagenin glucoside from the flowers of *Bombax cieba*. *Asian Journal of Research in Chemistry*, **3**, 78-80.
- [4] Iribarren AM, Pomilio AB. (1989) Steroidal glucosides, flavonoids and other components of *Bauhinia uruguayensis*. *Anales de la Asociacion Quimica Argentina*, **77**, 461-466.
- [5] Shang-jiang D, Gariboldi P, Jommi G. (1986) Constituents of Shashen (*Adenophora axilliflora*). *Planta Medica*, **4**, 317-320.
- [6] Achenbach H, Lowel M, Waibel R, Gupta M, Solis P. (1992) New lignan glucoside from *Stemmadenia minima*. *Planta Medica*, **58**, 270-272.
- [7] Qiu Y, Chen Y, Pei Y, Matsuda H, Yoshikawa M. (2002) Constituents with radical scavenging effect from *Opuntia dillenii*: structure of new α-pyrone and flavonol glycoside. *Chemical and Pharmaceutical Bulletin*, **50**, 1507-1510.
- [8] Saijo R, Nanaka GI, Nishioka I. (1989) Phenol glucoside gallates from *Mallotus japonicus*. *Phytochemistry*, **28**, 2443-2446.
- [9] Miyamura M, Nohara T, Tomimatso T, Nishioka I. (1983) Seven aromatic compounds from bark of *Cinnamomum cassia*. *Phytochemistry*, **22**, 215-218.
- [10] Higuchi H, Fukui K, Kinjo J, Nohara T. (1992) Four new glycosides from *Albiziaae Cortex*. III. *Chemical and Pharmaceutical Bulletin*, **40**, 534-535.
- [11] (a) Kosuge K, Mitsunaga K, Koike K, Ohmoto T. (1994) Studies on the constituents of *Ailanthus integrifolia*, *Chemical and Pharmaceutical Bulletin*, **42**, 1669-1671; (b) Suttisri R, Chung MS, Kinghorn AD, Sticher O, Yamshimoto Y. (1993) Periandrin V, a further sweet triterpene glycoside from *Periandra dulcis*. *Phytochemistry*, **34**, 405-408.

Two New Phenolic Glycosides from *Viburnum plicatum* var. *plicatum* f. *plicatum*

Saki Katagiri, Yoshiki Watanabe, Yasunori Yaoita, Masao Kikuchi and Koichi Machida*

Department of Molecular Structural Analysis, Tohoku Pharmaceutical University,
4-4-1 Komatsushima, Aoba-ku, Sendai, Miyagi 981-8558, Japan

kmachida@tohoku-pharm.ac.jp

Received: August 31st, 2011; Accepted: October 17th, 2011

Two new glycosides, named dideoxyplicatumoside A (**1**) and *erythro*-syringylglycerol- β -O-4'-(+)-isoeucommin A 4'''-O- β -D-glucopyranoside (**2**), together with seven known compounds, were isolated from the leaves of *Viburnum plicatum* Thunb. ex Murray var. *plicatum* f. *plicatum*. Their structures were established on the basis of NMR, MS and chemical data.

Keywords: *Viburnum plicatum*, Caprifoliaceae, phenolic glycoside, sesquilignan glycoside.

The deciduous shrub *Viburnum plicatum* Thunb. ex. Murray var. *plicatum* f. *plicatum* (Caprifoliaceae) is distributed in Japan, Taiwan and China. We previously reported the isolation of two new glycosides including 13 known compounds from the leaves of *V. plicatum* var. *plicatum* f. *plicatum* [1]. In the course of further studies on the constituents of this plant, two new glycosides, named dideoxyplicatumoside A (**1**) and *erythro*-syringylglycerol- β -O-4'-(+)-isoeucommin A 4'''-O- β -D-glucopyranoside (**2**), and 7 known ones (**3–9**) have been isolated. This article deals with the structural elucidation and identification of these compounds.

The MeOH extract of the fresh leaves of *V. plicatum* var. *plicatum* f. *plicatum* was partitioned with CHCl₃, AcOEt, *n*-BuOH, and H₂O. The *n*-BuOH soluble fraction was separated by a combination of chromatographic procedures to afford two new glycosides (**1** and **2**) and seven known ones (**3–9**). The known compounds **3–9** were identified as (7*S*,8*R*)-dihydro-dehydrodiconiferyl alcohol 4-O- β -D-glucopyranoside (**3**) [2], (7*R*,8*S*)-dihydrodehydrodiconiferyl alcohol 4-O- β -D-glucopyranoside (**4**) [2], kankanoside E (**5**) [3], 7-O-tigloylsecologanol (**6**) [4], (+)-pinoresinol 4-O- β -D-glucopyranoside (**7**) [5], (7*S*,8*R*)-dihydrodehydrodiconiferyl alcohol 9'-O- β -D-glucopyranoside (**8**) [6] and 3'-O-[(2*S*)-2-methylbutanoyl] henryoside (**9**) [4], respectively, by comparison of their spectroscopic data with those previously described in the literature. This is the first record of **3–9** in this plant.

Compound **1** was obtained as an optically active amorphous powder. The molecular formula of **1**, C₂₆H₃₂O₁₃, was confirmed by HR-FAB-MS (*m/z* 575.1749 [M+Na]⁺). The ¹H NMR spectrum of **1** showed signals due to nine aromatic protons, one CH₂ as an AB system, and





Figure 1: Structures of new compounds **1** and **2**.

two anomeric protons at δ_H 4.73 (1H, d, *J*=7.6 Hz, H-1''') and 5.34 (1H, d, *J*=7.3 Hz, H-1''). Acid hydrolysis of **1** yielded only D-glucose, which was identified by retention time and optical rotation using chiral detection by HPLC analysis. The coupling constants of the two anomeric protons indicated that the glycosyl linkages are of β -configuration. Its ¹H- and ¹³C NMR spectra were similar to those of plicatumoside A isolated from the same plant [1]. The ¹³C NMR spectrum of **1**, however, lacked signals from two hydroxylated aromatic quaternary carbons at C-6 and C-2' in plicatumoside A, and instead showed signals for two aromatic CH carbons in **1** [δ_C 132.4 (s, C-6), 129.5

**Figure 2:** Main HMBCs of **1**.


Heavy lines indicate partial structures inferred from ^1H - ^1H COSY.

Table 1: ^{13}C NMR spectroscopic data of **1** (100 MHz, CD_3OD).

Position	1	Position	1
1	121.6	1''	99.7
2	157.7	2''	83.2
3	116.3	3''	77.4
4	134.9	4''	70.7
5	122.6	5''	77.7
6	132.4	6''	61.5
7	167.3	1'''	104.5
1'	137.6	2'''	76.1
2', 6'	129.5	3'''	77.9
3', 5'	129.7	4'''	70.8
4'	129.4	5'''	77.6
7'	67.9	6'''	62.4

(s, C-2', 6'). These spectral data, including the molecular formula, indicated that **1** is the 6, 2'-dideoxy derivative of plicatumoside A. Detailed analyses of the ^1H - and ^{13}C NMR spectra of **1** were undertaken with the aid of ^1H - ^1H COSY and HMBC experiments (Figure 2). Consequently, the structure of **1** was elucidated as shown, and named dideoxyplicatumoside A.

Compound **2** was obtained as an optically active amorphous powder. The molecular formula of **2**, $\text{C}_{44}\text{H}_{58}\text{O}_{22}$, was confirmed by HR-FAB-MS (m/z 961.3316 $[\text{M}+\text{Na}]^+$). In the ^1H NMR spectrum of **2**, a signal pattern was similar to that of (+)-isoeucommin A [7], except for the presence of additional signals due to a 1,3,4,5-tetra-substituted benzene group, two methoxy groups [δ_{H} 3.82 (6H, s, OMe-3''', 5''')] and an anomeric group [δ_{H} 4.81 (1H, d, $J=7.3$ Hz, H-1''')]. Furthermore, the ^{13}C NMR spectrum exhibited the additional signals attributed to two oxygenated CH groups [δ_{C} 74.1 (d, C- α), 87.0 (d, C- β)] and an oxygenated CH_2 group [δ_{C} 61.7 (t, C- γ)]. Acid hydrolysis of **2** yielded only D-glucose, which was identified as described above. The coupling constants of the two anomeric protons indicated that the glycosyl linkages have β -configurations. Detailed analysis of the ^1H - and ^{13}C NMR spectra of the additional signals, with the aid of ^1H - ^1H COSY, HMQC and HMBC experiments (Figure 3), suggested that the additional moiety was a syringylglycerol 4- O - β -D-glucopyranoside unit. From the findings presented above, compound **2** was a sesquiquignan formed by (+)-isoeucommin A and syringylglycerol 4- O - β -D-glucopyranoside units. The additional syringylglycerol 4- O - β -D-glucopyranoside unit of **2** was attached to OH-4' of (+)-isoeucommin A via an ether bridge, as shown by the HMBC cross-peak between H- β and C-4' (Figure 3).

**Figure 3:** Main HMBCs of **2**.

Heavy lines indicate partial structures inferred from ^1H - ^1H COSY.

Table 2: ^{13}C NMR spectroscopic data of (+)-isoeucommin A, **2** and **2a**.

Position	(+)-isoeucommin A [*]	2 [*]	2a [#]
1	55.6	55.8	54.1
2	87.7	87.1	86.1
4	72.8	73.0	72.3
5	55.6	55.5	54.5
6	87.1	87.3	85.6
8	72.8	73.1	71.4
1'	133.2	139.0	138.0
2', 6'	104.6	104.3	102.9
3', 5'	149.4	154.6	153.4
4'	137.5	136.2	134.1
3', 5'-OCH ₃	56.9	56.8	56.3
1''	137.5	137.5	132.7
2''	111.7	111.7	108.6
3''	151.0	151.0	146.7
4''	147.6	147.6	145.3
5''	118.1	118.1	114.3
6''	119.8	119.8	118.9
3''-OCH ₃	56.8	57.0	56.0
α	—	74.1	72.6
β	—	87.0	86.9
γ	—	61.7	60.5
1'''	—	139.5	137.0
2''', 6'''	—	106.0	102.9
3''', 5'''	—	153.8	152.7
4'''	—	135.5	134.5
3''', 5'''-OCH ₃	—	56.7	56.4
1'''	102.9	102.9	—
2'''	75.0	75.0	—
3'''	77.9	77.9	—
4'''	71.4	71.4	—
5'''	78.3	78.4	—
6'''	62.6	62.6	—
1'''	—	105.6	106.5
2'''	—	75.8	74.1
3'''	—	78.3	75.9
4'''	—	71.4	70.5
5'''	—	77.8	76.6
6'''	—	62.5	62.7

^{*}100 Mz, CD_3OD , [#]100 MHz, CDCl_3 .

Enzymatic hydrolysis of **2** afforded **2a**. As shown in Figure 4, the relative configuration of C- α and C- β of syringylglycerol 4- O - β -D-glucopyranoside unit in **2** was determined to be *erythro* by comparing the coupling constant between H- α and H- β ($J=3.9$ Hz) in the ^1H NMR spectrum (CDCl_3) of **2a** with those of the *threo* and *erythro* isomers [8–15], which was also supported by the NOESY correlations (H- α /H- β , H- α /OCH₃-3', 5', H-2''', 6'''/H- α - γ , H-2''', 6'''/H- β) of **2a**. Furthermore, NOE correlations were observed between H-1'/H-2', 6', and H-5'/H-2'', 6''. Consequently, the structure of **2** was elucidated as shown, and named *erythro*-syringylglycerol- β -O-4'-(+)-isoeucommin A 4'''-O- β -D-glucopyranoside.




Figure 4: Main NOEs of **2**.

However, the absolute configurations of the chiral centers at C- α and C- β remain to be determined.

To the best of our knowledge, compound **1** is the first example of a naturally occurring benzyl salicylate glycoside, and there is no report of the isolation of a sesquiquinone constituent from the *Viburnum* species.

Experimental

General: Optical rotations were measured with a JASCO DIP-360 digital polarimeter. UV spectra were recorded with a Beckman DU-64 spectrometer. ^1H - and ^{13}C NMR spectra were recorded on a JEOL JNM-LA 400 (400 MHz, 100 MHz, respectively) spectrometer. Chemical shifts are given on a δ (ppm) scale, with tetramethylsilane (TMS) as internal standard. FAB-MS were recorded on a JEOL JMS-DX 303 mass spectrometer. Column chromatography was carried out on Kieselgel 60 (230–400 mesh, Merck, Darmstadt, Germany) and Sephadex LH-20 (Pharmacia Fine Chemicals, Sweden). HPLC was carried out on a Tosoh HPLC system [pump, CCPS; detector, UV-8020; column, TSK gel ODS 120T (7.8 mm i.d. \times 30 cm, Tosoh, Tokyo, Japan), and Cosmosil 5SL (10 mm i.d. \times 25 cm, Nacalai, Tokyo, Japan).

Plant material: Leaves of *V. plicatum* Thunb. ex Murray var. *plicatum* f. *plicatum* were collected in May, 2007 in Sendai, Miyagi prefecture, Japan, and identified by one of the authors (M. K). A voucher specimen (2007-5-KM3) is held in the laboratory of M. Kikuchi.

Extraction and isolation: Fresh leaves of *V. plicatum* var. *plicatum* f. *plicatum* (860 g) were extracted with MeOH at room temp. for 3 weeks. The MeOH extract was concentrated under reduced pressure and the residue (74.5 g) was suspended in water. This suspension was successively extracted with CHCl_3 (28.8 g) AcOEt (14.8 g), *n*-BuOH (14.4 g) and H_2O (12.4 g). The *n*-BuOH-soluble fraction was concentrated under reduced pressure to produce a residue. This was chromatographed on a silica gel column using CHCl_3 –MeOH– H_2O (40:10:1) and the eluate was separated into 20 fractions (frs. 1–20). Fraction 2 was chromatographed on a Sephadex LH-20 column using 50% MeOH and the eluate was separated into 18 fractions (frs. 2-1–2-18). Fraction 2-6 was subjected to prep. HPLC [column, TSK gel ODS 120T; mobile phase, MeOH– H_2O (5:6); UV detector, 205 nm; flow rate, 1.5 mL/min; column temp., 40°C; Cosmosil 5SL; mobile

phase, CH_2Cl_2 –MeOH– H_2O (7:10:1); UV detector, 225 nm; flow rate, 1.5 mL/min; column temp., 28°C] to give **1** (20.5 mg), **3** (2.8 mg), **4** (3.0 mg), **5** (4.5 mg) and **6** (5.0 mg), respectively. Fraction 2-7 was subjected to prep. HPLC [column, TSK gel ODS 120T; mobile phase, MeOH– H_2O (5:9); UV detector, 205 nm; flow rate, 1.5 mL/min; column temp., 40°C; TSK gel ODS 120T; mobile phase, MeOH– H_2O (2:5); UV detector, 205 nm; flow rate, 1.5 mL/min; column temp., 40°C] to give **7** (18.5 mg), **8** (2.0 mg) and **9** (5.0 mg), respectively. Fraction 4 was chromatographed on a Sephadex LH-20 column using 50% MeOH and the eluate was separated into 10 fractions (frs. 4-1–4-10). Fraction 4-1 was subjected to prep. HPLC [column, TSK gel ODS 120T; mobile phase, MeOH– H_2O (2:5); UV detector, 205 nm; flow rate, 1.5 mL/min; column temp., 40°C] to give **2** (3.2 mg).

Dideoxypticatumoside A (1)

An amorphous powder.

$[\alpha]_{\text{D}}^{25}$: –84.3 (*c* 0.25, MeOH).

UV λ_{max} (MeOH) nm (log ϵ): 205 (4.5), 231 (3.9), 289 (3.4).

^1H NMR (400 MHz, CD_3OD) δ : 3.10–3.15 (2H, m, H-5''', H_A-6'''), 3.19 (1H, dd, J =9.3, 7.6 Hz, H-2'''), 3.33–3.39 (2H, m, H-4'', 3'''), 3.43–3.49 (3H, m, H-5'', 4'', H_B-6'''), 3.66 (1H, dd, J =11.7, 6.0 Hz, H_A-6''), 3.67 (1H, t, J =9.0 Hz, H-3''), 3.73 (1H, dd, J =9.0, 7.3 Hz, H-2''), 3.86 (1H, dd, J =11.7, 2.0 Hz, H_B-6''), 4.73 (1H, d, J =7.6 Hz, H-1'''), 5.30 (1H, d, J =12.5 Hz, H_A-7), 5.34 (1H, d, J =7.3 Hz, H-1''), 5.36 (1H, d, J =12.5 Hz, H_B-7), 7.06 (1H, br t, J =7.8 Hz, H-5), 7.25 (1H, br d, J =8.3 Hz, H-3), 7.36 (1H, dt, J =8.2, 1.4 Hz, H-4'), 7.40 (2H, t, J =8.2 Hz, H-3', 5'), 7.44 (2H, dd, J =8.2, 1.4 Hz, H-2', 6'), 7.51 (1H, br dt, J =7.8, 1.7 Hz, H-4), 7.77 (1H, dd, J =7.8, 1.7 Hz, H-6).

^{13}C NMR (CD_3OD): Table 1.

FAB-MS m/z : 575 $[\text{M}+\text{Na}]^+$. HR-FAB-MS m/z : 575.1749 $[\text{M}+\text{Na}]^+$ (Calcd for $\text{C}_{26}\text{H}_{32}\text{O}_{13}\text{Na}$: 575.1740).

Erythro-syringylglycerol- β -O-4'-(+)-isoeucommin A 4''-O- β -D-glucopyranoside (2)

An amorphous powder.

$[\alpha]_{\text{D}}^{25}$: –14.0 (*c* 0.20, MeOH).

UV λ_{max} (MeOH) nm (log ϵ): 211 (4.6), 230sh (4.3), 274 (3.5).

^1H NMR (400 MHz, CD_3OD) δ : 3.12 (2H, m, H-1, 5), 3.20–3.50 (8H, m, H-2''', 3''', 4''', 5''', 2''', 3''', 4''', 5'''), 3.61 (1H, dd, J =12.0, 3.7 Hz, H_A- γ), 3.64–3.86 (4H, m, H₂-6''', H₂-6'''), 3.80 (6H, s, 3', 5'-OCH₃), 3.82 (6H, s, 3'', 5''-OCH₃), 3.87 (3H, s, 3''-OCH₃), 3.90 (3H, m, H_A-4, H_A-8, H_B- γ), 4.27 (3H, m, H_B-4, H_B-8, H- β), 4.74 (1H, d, J =4.2 Hz, H-2), 4.77 (1H, d, J =4.4 Hz, H-6), 4.81 (1H, d, J =7.3 Hz, H-1'''), 4.87 (H-1'''), overlapped with solvent signal), 4.91 (1H, d, J =5.9 Hz, H- α), 6.67 (2H, s, H-2', 6'), 6.73 (2H, s, H-2'', 6''), 6.92 (1H, dd, J =8.3, 1.7 Hz, H-6'''), 7.03 (1H, d, J =1.7 Hz, H-2''), 7.15 (1H, d, J =8.3 Hz, H-5''). ^{13}C NMR (CD_3OD): Table 2.

FAB-MS m/z : 961 $[\text{M}+\text{Na}]^+$. HR-FAB-MS m/z : 961.3316 $[\text{M}+\text{Na}]^+$ (Calcd for $\text{C}_{44}\text{H}_{58}\text{O}_{22}\text{Na}$: 961.3317).

Enzymatic hydrolysis of 2: An aqueous solution (5.0 mL) containing **2** (2.0 mg) and β -glucosidase (10 mg) was incubated at 40°C for 3 days. The reaction mixture was extracted with CHCl_3 , and the CHCl_3 layer was evaporated under reduced pressure. The residue was purified by HPLC [column, Cosmosil 5SL; mobile phase, CH_2Cl_2 -MeOH- H_2O (70:10:1); UV detector, 225 nm; flow rate, 1.5 mL/min; column temp., 28°C] to give **2a** as an amorphous powder (1.0 mg).

$[\alpha]_{\text{D}}^{25}$: +14.3 (*c* 0.15, MeOH).

UV λ_{max} (MeOH) nm (log ϵ): 231 (4.4), 274 (3.0).

^1H NMR (400 MHz, CDCl_3) δ : 3.10 (1H, m, H-1), 3.16 (1H, m, H-5), 3.40–3.95 (6H, m, H-2''', 3''', 4''', 5''', H-6'''), 3.47 (1H, br d, $J=11.0$ Hz, H_A - γ), 3.86 (6H, s, 3'', 5''-OCH₃), 3.87 (3H, m, H_A -4, H_A -8, H_B - γ), 3.90 (6H, s, 3', 5'-OCH₃), 3.91 (3H, s, 3''-OCH₃), 4.11 (1H, m, H- β), 4.26 (1H, dd, $J=9.3$, 6.3 Hz, H_B -8), 4.34 (1H, dd, $J=9.0$, 7.1 Hz, H_B -4), 4.53 (1H, d, $J=7.6$ Hz, H-1'''), 4.74 (1H, d, $J=5.4$ Hz, H-2), 4.79 (1H, d, $J=4.6$ Hz, H-6), 4.99 (1H, d, $J=3.9$ Hz, H- α), 6.62 (2H, s, H-2'', 6''), 6.64 (2H, s, H-2', 6'), 6.82 (1H, dd, $J=8.1$, 2.0 Hz, H-6''), 6.89 (1H, d, $J=2.0$ Hz, H-2''), 6.90 (1H, d, $J=8.1$ Hz, H-5'').

^{13}C NMR (CDCl_3): Table 2.

FAB-MS m/z : 799 $[\text{M}+\text{Na}]^+$. HR-FAB-MS m/z : 799.2787 $[\text{M}+\text{Na}]^+$ (Calcd for $\text{C}_{38}\text{H}_{48}\text{O}_{17}\text{Na}$: 799.2789).

Acid hydrolysis of 1 and 2: Each compound (*ca.* 1.0 mg) was refluxed with 1M HCl (1 mL) for 5 h. The reaction mixture was neutralized with Ag_2CO_3 and filtered. The solution was concentrated *in vacuo* and dried to give a sugar fraction. The sugar fraction was analyzed by HPLC under the following conditions: column, TSK gel Amide-80; column temperature, 45°C; mobile phase, CH_3CN - H_2O (4:1); flow rate, 1.0 mL/min; chiral detection (JASCO OR-2090). Identification of D-glucose present in the sugar fraction was carried out by comparison of the retention time and optical rotation with those of an authentic sample; t_{R} (min) 39.0 (D-glucose, positive optical rotation).

Acknowledgments - The authors are grateful to Mrs S. Sato and T. Matsuki of Tohoku Pharmaceutical University for conducting the NMR and MS measurements.

References

- [1] Kikuchi M, Onoguchi R, Yaoita Y, Machida K. (2011) Two new glycosides from *Viburnum plicatum* Thunb. ex Murray var. *plicatum* f. *plicatum*. *Journal of Natural Medicines*, **65**, 202–205.
- [2] Matsuda N, Sato H, Yaoita Y, Kikuchi M. (1996) Isolation and absolute structures of the neolignan glycosides with the enantiometric aglycones from the leaves of *Viburnum awabuki* K. Koch. *Chemical & Pharmaceutical Bulletin*, **44**, 1122–1123.
- [3] Xie H, Morikawa T, Matsuda H, Nakamura S, Muraoka O, Yoshikawa M. (2006) Monoterpene constituents from *Cistanche tubulosa*. Chemical structures of kankanosides A–E and kankanol. *Chemical & Pharmaceutical Bulletin*, **54**, 669–675.
- [4] Machida K, Sagawa H, Onoguchi R, Kikuchi M. (2010) Three new glycosides from *Viburnum plicatum* Thunb. var. *tomentosum* MiQ. *Helvetica Chimica Acta*, **93**, 290–297.
- [5] Nishibe S, Chiba M, Hisada S. (1977) Studies on the Chinese crude drug “Forsythiae Fructus.” I. On the constituents of Forsythiae Fructus on the market. *Yakugaku Zasshi*, **97**, 1134–1137.
- [6] Machida K, Sakamoto S, Kikuchi M. (2009) Two new neolignan glycosides from leaves of *Osmanthus heterophyllus*. *Journal of Natural Medicines*, **63**, 227–231.
- [7] Sugiyama M, Kikuchi M. (1991) Studies on the constituents of *Osmanthus* species. VII. Structures of lignan glycosides from the leaves of *Osmanthus asiaticus* Nakai. *Chemical & Pharmaceutical Bulletin*, **39**, 483–485.
- [8] Braga ACH, Zacchino S, Badano H, Sierra MG, Ruveda EA. (1984) ^{13}C NMR spectral and conformational analysis of 8-*O*-4' neolignans. *Phytochemistry*, **23**, 2025–2028.
- [9] Matsuda N, Kikuchi M. (1996) Studies on the constituents of *Lonicera* species. X. Neolignan glycosides from the leaves of *Lonicera gracilipes* var. *glandulosa* Maxim. *Chemical & Pharmaceutical Bulletin*, **44**, 1676–1679.
- [10] Sung S-H, Huh M-S, Kim Y-C. (2001) New tetrahydrofuran-type sesquilignans of *Saururus chinensis* root. *Chemical & Pharmaceutical Bulletin*, **49**, 1192–1194.
- [11] Park S-Y, Hong S-S, Han X-H, Hwang J-S, Lee D, Ro J-S, Hwang B-Y. (2007) Lignans from *Arctium lappa* and their inhibition of LPS-induced nitric oxide production. *Chemical & Pharmaceutical Bulletin*, **55**, 150–152.
- [12] Machida K, Sakamoto S, Kikuchi M. (2008) Structure elucidation and NMR spectral assignments of four neolignan glycosides with enantiometric aglycones from *Osmanthus ilicifolius*. *Magnetic Resonance in Chemistry*, **46**, 990–994.
- [13] Kim K-H, Moon E, Kim S-Y, Lee K-R. (2010) Lignans from the tuber-barks of *Colocasis antiquorum* var. *esculenta* and their antimelanogenic activity. *Journal of Agricultural and Food Chemistry*, **58**, 4779–4785.
- [14] Wang W, Ali Z, Li X-C, Khan I-A. (2010) Neolignans from the leaves of *Casearia sylvestris* SWARTZ. *Helvetica Chimica Acta*, **93**, 139–146.
- [15] Li S-F, Di Y-T, Wang Y-H, Tan C-J, Fang X, Zhang Y, Zheng Y-T, Li L, He H-P, Li S-L, Hao X-J. (2010) Anthraquinones and lignans from *Cassia occidentalis*. *Helvetica Chimica Acta*, **93**, 1795–1802.

Antimicrobial Chemical Constituents from the Endophytic Fungus *Phomopsis* sp. from *Notobasis syriaca*Hidayat Hussain^{a,c,*}, Michel Kenne Tchimené^a, Ishtiaq Ahmed^a, Kathrin Meier^b, Michael Steinert^b, Siegfried Draeger^b, Barbara Schulz^b and Karsten Krohn^{a,*}^aDepartment of Chemistry, Universität Paderborn, Warburger Straße 100, 33098 Paderborn, Germany^bInstitute of Microbiology, Technical University of Braunschweig, Spielmannstraße 7, 31806 Braunschweig, Germany^cDepartment of Biological Sciences and Chemistry, College of Arts and Sciences, University of Nizwa, Birkat Al-Mouz, Nizwa 616, Sultanate of Oman

Hidayat110@gmail.com; k.krohn@upb.de

Received: May 14th, 2011; Accepted: October 31st, 2011

Phomosine K (**1**), a new phomosine derivative, has been isolated from *Phomopsis* sp., in addition to six known compounds: phomosine A (**2**), phenylalanine amide (**3**), 2-hydroxymethyl-4 β ,5 α ,6 β -trihydroxycyclohex-2-en (**4**), (-)-phyllostine (**5**), (+)-epiepoxydon (**6**), and (+)-epoxydon monoacetate (**7**). Preliminary studies showed that compound **1** had strong antibacterial activity, while compounds **4-7** showed good antifungal, antibacterial, and algicidal properties, except compounds **4** and **6**, which lacked antifungal activity.

Key words: Fungal metabolites, *Phomopsis* sp., phomosines, epoxydones, antimicrobial activity.

Endophytic fungi are a rich source of novel organic compounds with interesting biological activities [1]. As part of a program on the isolation of biologically active compounds, we have now investigated *Phomopsis* sp. (internal strain no. 8966), isolated from the plant *Notobasis syriaca* and isolated a number of metabolites (Figure 1).

Phomosine K (**1**), a white solid, showed an $[M]^+$ at $m/z = 332.1267$ (HREIMS), corresponding to the molecular formula $C_{18}H_{20}O_6$. The IR spectrum of **1** showed the presence of an ester carbonyl group (1647 cm^{-1}) and a hydroxyl group (3410 cm^{-1}). The ^1H NMR spectrum showed signals for four methyl groups at δ 2.14 (s, 3H, $\text{CH}_3\text{-2'}$), 2.15 (s, 3H, $\text{CH}_3\text{-5}$), 2.16 (s, 3H, $\text{CH}_3\text{-5'}$), and 2.30 (s, 3H, $\text{CH}_3\text{-2}$), three phenolic OH groups at δ 2.14 (s, 3H, $\text{CH}_3\text{-2'}$), 2.15 (s, 3H, $\text{CH}_3\text{-5}$), and 2.16 (s, 3H, $\text{CH}_3\text{-5'}$), 2.30 (s, 3H, $\text{CH}_3\text{-2}$), three phenolic OH groups at δ 11.87, 7.29, and 6.54 (s, 1H, OH), one methoxy signal at δ 3.94 (s, 3H, CO_2CH_3) and two aromatic protons at δ 5.84 (s, 1H, 6'-H), and 6.46 (s, 1H, 4'-H). The ^{13}C NMR and DEPT spectrum indicated the presence of five CH_3 , two CH, and eleven quaternary carbons. The structure of **1** was determined by comparison of its NMR spectroscopic data with those of phomosine A (**2**) [2]. A fourth methyl group appeared at δ 2.14 (s, 3H), instead of an aldehydic proton at δ 10.43, as found in compound **2**. The fourth methyl group at C-2' was further confirmed from HMBC correlations of $\text{CH}_3\text{-2'}$ to C-1', C-2', and C-3'. Thus, the structure of **1**, was established as methyl 2,4-dihydroxy-5-(3-hydroxy-2,5-dimethylphenoxy)-3,6-dimethylbenzoate. The known compounds phomosine A (**2**) [2], phenyl-




Figure 1: Chemical structure of compounds 1-7.

alanine amide [**3**], 2-hydroxymethyl-4 β ,5 α ,6 β -trihydroxycyclohex-2-en (**4**) [**4**], (-)-phyllostine (**5**) [**5a**], (+)-epiepoxydon (**6**) [**5**], and (+)-epoxydon monoacetate (**7**) [**5b**] were identified by comparison with published data.

The isolated compounds **4-7** were tested in an agar diffusion assay for their antifungal, antibacterial, and algicidal properties towards *Microbotryum violaceum*, *Escherichia coli*, *Bacillus megaterium*, and *Chlorella fusca* (Table 1). Metabolite **1** was tested for antibacterial activity in a MIC (minimum inhibitory concentration)-assay in liquid medium against *Legionella pneumophila* Corby, *Escherichia coli* K12 and *Bacillus megaterium* (Table 2).

Experimental

Plant material: College. Experimental procedures were quite similar to those reported earlier [1b]. Microbiological methods and culture conditions were as previously described [6a,b].

Culture, extraction, and isolation: The endophytic fungus *Phomopsis* sp. was isolated from *Notobasis syriaca*. It was

Table 1: Biological activities of pure metabolites **4-7** against microbial test organisms in agar diffusion assay.

Compound	antibacterial	antibacterial	antialgal	antifungal
	Ec ^a	Bm	Chl	Mv
4	7	PI 8	6	0
5	10	PI 7	10	7
6	7	0	5	0
7	9	PI 7	9	5
Penicillin	14	18	0	0
Tetracycline	18	18	PI 10	0
Nystatin	0	0	0	20
Actidione	0	0	35	50
Acetone	0	0	0	0

^a*Chlorella fusca* (Chl), *Microbotryum violaceum* (Mv), *Escherichia coli* (Ec), and *Bacillus megaterium* (Bm). Application of pure substances at a concentration of 0.05 mg (50 µL of 1 mg/mL). The radius of zone of inhibition was measured in mm. PI = partial inhibition, i.e. there was some growth within the zone of inhibition.

Table 2: Antibacterial activity of **1** in a MIC assay in liquid medium against bacterial test organisms.

Compound 1 µg / mL	Test organisms		
	<i>Lp</i> ^a	<i>Ec</i>	<i>Bm</i>
100	+	+	-
50	+	-	-
25	+	-	-
12.5	-	-	-

cultivated at room temperature for 28 days [7,8] on biomalt solid agar medium. The culture medium was then extracted with ethyl acetate to afford 5.0 g of a residue after removal of the solvent under reduced pressure. The extract was separated into 2 fractions by CC on silica gel, using gradients of dichloromethane / ethyl acetate (85:15, 50:50, 0:100). The less polar fraction 1 (1.5 g) contained mainly fatty acids and lipids. The remaining fraction was further purified by silica gel CC and preparative TLC with *n*-hexane / ethyl acetate (10:1 to 5:1) to give pure compounds **1** (11 mg), **2** (40 mg), **3** (4 mg), **4** (6 mg), **5** (5 mg), **6** (5 mg), and **7** (5 mg).

Phomosine K (**1**)

MP: 199°C.

UV (CHCl₃): λ_{max} (log ε) = 277 (3.70), 310 (3.50) nm.

IR (KBr): ν = 3430 cm⁻¹, 2953, 1650, 1620, 1580, 1440, 1372, 1090, 1020, 795.

References

- [1] (a) Krohn K, Ullah Z, Hussain H, Floerke U, Schulz B, Draeger S, Pescitelli G, Salvadori P, Antus S, Kurtan T. (2007) Massarilactones E-G, new metabolites from the endophytic fungus *Coniothyrium* sp., associated with the plant *Artimisia maritima*. *Chirality*, **19**, 464-470; (b) Dai J, Hussain H, Dräger S, Schulz B, Kurtan T, Pescitelli G, Flörke U, Krohn K. (2010) Metabolites from the fungus *Phoma* sp. 7210, associated with *Aizoon canariense*. *Natural Product Communications*, **5**, 1175-1180.
- [2] Krohn K, Michel A, Roemer E, Flörke U, Aust HJ, Draeger S, Schulz B, Wray V. (1995) Biologically active metabolites from fungi; three new biaryl ethers from *Phomopsis* sp. *Natural Product Letters*, **9**, 309-314.
- [3] Ramallal RM, Liz R, Gotor V. (2010) Enantiopure trans-3-arylaziridine-2-carboxamides: Preparation by bacterial hydrolysis and ring-openings toward enantiopure, unnatural amino acids. *Journal of Organic Chemistry*, **75**, 6614-6624.
- [4] Smetanina OF, Kalinovskii AI, Khudyakov YV, Moiseenko OP, Pivkin MV, Menzorova NI, Sibirtsev YT, Kuznetsova TA. (2005) Metabolites of the marine fungus *Asperigillus varians* KMM 4630. *Chemistry of Natural Compounds*, **41**, 243-244.
- [5] (a) Mehta G, Islam K. (2004) Enantioselective total synthesis of epoxyquinone natural products (-)-phylllostine, (+)-epoxydon, (+)-epiepoxydon and (-)-panepophenanthrin: access to versatile chiral building blocks through enzymatic kinetic resolution. *Tetrahedron Letters*, **45**, 7683-7687; (b) Klemke C, Kehraus S, Wright AD, König GM. (2004) New secondary metabolites from the marine endophytic fungus *Apiospora montagnei*. *Journal of Natural Products*, **67**, 1058-1063.
- [6] (a) Schulz B, Sucker J, Aust HJ, Krohn K, Ludewig K, Jones PG, Doering D. (1995) Biologically active secondary metabolites of endophytic *Pezizula* species. *Mycological Research*, **99**, 1007-1015; (b) Höller U, Wright AD, Matthee GF, König GM, Draeger S, Aust HJ, Schulz B. (2000) Fungi from marine sponges: diversity, biological activity and secondary metabolites. *Mycological Research*, **104**, 1354-1365; (c) Stewart AM, Meier K, Schulz B, Steinert M, Snider BB. (2010) Synthesis and biological evaluation of (±)-dinemasone C and analogues. *Journal of Organic Chemistry*, **75**, 6057-6060.

¹H NMR (500 MHz, CDCl₃): δ = 2.14 (s, 3H, CH₃-2'), 2.15 (s, 3H, CH₃-5), 2.16 (s, 3H, CH₃-5'), 2.30 (s, 3H, CH₃-2), 3.94 (s, 3H, CO₂CH₃), 5.84 (s, 1H, 6'-H), 6.46 (s, 1H, 4'-H), 6.54 (s, 1H, OH), 7.29 (s, 1H, OH), 11.87 (s, 1H, 6-OH).

¹³C NMR (125 MHz, CDCl₃): δ = 8.2 (CH₃-5), 15.4 (CH₃-2), 20.9 (CH₃-2'), 21.7 (CH₃-5'), 51.9 (CO₂CH₃), 104.5 (C-1), 105.8 (C-6'), 108.4 (C-3), 110.8 (C-4'), 112.8 (C-2'), 131.1 (C-6), 133.1 (C-5), 142.2 (C-5'), 152.5 (C-4), 156.6 (C-1'), 156.6 (C-3'), 160.5 (C-6), 172.4 (C-1a).

EIMS (70 eV, 200°C): *m/z* (%) = 332.1 (55) [M⁺], 298 (100), 255(25), 180 (69), 108 (96). HREIMS: *m/z* 332.1215 (calcd. 332.1260 for C₁₈H₂₀O₆).

Assays for biological activity: For the agar diffusion assays, the compounds were dissolved in acetone at a concentration of 1 mg/mL. Fifty µL of the solutions (50 µg) was pipetted onto a sterile filter disk (Schleicher & Schuell, 9 mm), which was placed onto an appropriate agar growth medium for the respective test organism and subsequently sprayed with a suspension of the test organism [6a]. The fungal test organisms were *Escherichia coli*, *Bacillus megaterium*, and *Microbotryum violaceum*. Commencing at the middle of the filter disk, the radius of the zone of inhibition was measured in mm. The MIC-assays were conducted in microtiter plates in liquid media. *Legionella pneumophila* Corby was tested in YEB medium [6c]; *Escherichia coli* and *Bacillus megaterium* in NB medium. The metabolites were dissolved in methanol. In the wells, the bacterial test organisms were initially at a concentration of 2 x 10⁷ cells / mL; final concentrations of the metabolites were 12.5 – 100 µg / mL; incubation of *L. pneumophila* was for 3 days with 5% CO₂ at 37°C, and that of *B. megaterium* and *E. coli* for one day at 24°C.

Acknowledgement - We thank BASF AG and the Bundesministerium für Bildung und Forschung (BMBF), project no. 03F0360A for sponsoring our research work.

Phomosines H–J, Novel Highly Substituted Biaryl Ethers, Isolated from the Endophytic Fungus *Phomopsis* sp. from *Ligustrum vulgare*[‡]

Karsten Krohn^{a,*}, Umar Farooq^{a,e}, Hidayat Hussain^a, Ishtiaq Ahmed^a, Joachim Rheinheimer^b, Siegfried Draeger^c, Barbara Schulz^c and Teunis van Ree^d

[‡]Biologically Active Metabolites from Fungi, 52. Part 51: Ref. [1]

^aDepartment of Chemistry, Universität Paderborn, Warburger Straße 100, 33098 Paderborn, Germany

^bBASF AG, GVA/G-A030, 67056 Ludwigshafen, Germany

^cInstitute of Microbiology, Technical University of Braunschweig, Spielmannstraße 7, 31806 Braunschweig, Germany

^dDepartment of Chemistry, University of Venda, Thohoyandou, 0950, South Africa

^ecurrent address: COMSATS Institute of Information Technology, Abbottabad, Pakistan


k.krohn@upb.de

Received: May 7th, 2011; Accepted: October 29th, 2011

From the endophytic fungus *Phomopsis* sp., four known phomosines A-D (**1-4**) and three new phomosines H-J (**5-7**) have been isolated. The structures of the new compounds were determined on the basis of their spectroscopic data analysis (¹H, ¹³C, ¹H-¹H COSY, HMQC, and HMBC NMR, as well as mass spectrometry). The structures of phomosine H (**5**) and J (**7**) were also confirmed by semisynthesis from phomosine A (**1**). The remaining four known compounds [phomosines A-D (**1-4**)] were identified by comparing their spectroscopic data with those reported in the literature. The four known metabolites were biologically active. Of the novel metabolites, only **6** was antifungal and antibacterial.

Key words: Endophytic fungi, fungal metabolites, phomosines, *Phomopsis* sp.

Endophytic fungi have been shown to be an excellent source of biologically active secondary metabolites [2]. We have investigated novel metabolites from an endophytic fungus, *Phomopsis* sp., which was isolated from *Ligustrum vulgare*. *Phomopsis* species have been found to synthesize a large array of structurally diverse secondary metabolites [3]. These include various classes, for example anthraquinones (e.g. altersolanol A-C [4], cytochalasins [5,6], phomopsichalasin [7], phenochalasin) [8], lactones (convolvulol, convolvulopyrone, convolvulanic acid A) [9], xanthenes [10-12], nitropropionic esters [13,14], terpenoides, such as the oblongolides [13-15] and phomopsolides [16], 2,3-epoxycyclohexenes [17], isocoumarins [17], cyclic peptides, such as phomopsin A and B [18], orselliniic esters (phomozin) [19], macrolides [20,21], polyketides [22] and biaryl ethers, such as phomosines A-C [23] and D-G [24]. In this paper, we describe the isolation, physico-chemical properties, structure elucidation, semisynthesis, and biological activity of three new phomosines H-J (**5-7**), along with four known phomosines A-D (**1-4**). The fungus was cultivated in biomalt semi-solid agar culture media for 26 days at room temperature. The crude extract had good



Scheme 1: Chemical structure of compounds 1-7.

fungicidal, antibacterial, and algicidal activities. The ethyl acetate extract of the culture was separated by silica gel

column chromatography, resulting in the isolation of four known (**1-4**) and three new (**5-7**) biaryl ethers.


Four known metabolites, phomosines A-D (**1-4**) (Figure 1), were identified by comparison of their spectral data with those published in the literature [23,24]. The three new metabolites (**5-7**) of intermediate polarity were elucidated by direct comparison of their spectral data with those of compounds **1-4**.

Phomosine H (**5**) was isolated as a colorless solid with m.p. 163-165°C. The IR spectrum, with bands at 1648 and 3435 cm^{-1} , indicated the presence of conjugated and/or chelated ester carbonyl and hydroxyl groups. The NMR spectra with 25 signals in the ^{13}C NMR and 26 proton signals in the ^1H NMR spectra, together with the molecular ion at $m/z = 438$, suggested a molecular composition of $\text{C}_{25}\text{H}_{26}\text{O}_7$. Although both ^1H NMR and ^{13}C NMR spectra of **5** were in close correspondence to those of phomosines **1-4**, some new signals could be observed such as the resonances at δ 4.98 and 4.72 for two methylene groups and additional signals at δ 7.33-7.43 for five aromatic protons. Resonances at δ 2.33, 2.16 and 2.15 in the ^1H NMR spectrum were typical for three aromatic methyl groups, while two signals at δ 6.39 and 5.86 indicate aromatic protons at C-4' and C-6'. One phenolic hydroxyl was chelated with a carbonyl group and appeared at lower field (δ 11.8).

As in the phomosines A-D (**1-4**), one signal at δ 3.96 showed the presence of an ester methoxy group. The location of the benzyl ether at the benzylic methylene at C-1' was confirmed by two factors. First, the ^{13}C NMR signal for C-2'a in the free alcohol **4** is at δ 53.6, whereas in the methyl ether **3** it is at δ 63.9 and in the benzyl ether at δ 63.6. Secondly, this was unambiguously confirmed by the HMBC experiment with correlations of 2'a-H to C-3'a and 3'a-H- to C-2'a.

In order to further confirm the structure of phomosine H (**5**), chemical synthesis of **5** from **1** was performed (Scheme 2). Protection of three phenolic hydroxyl groups in **1** was achieved with *para*-methoxybenzyl chloride to form the *tris*-PMB ether **8** in 94 % yield. Treatment of **8** with NaBH_4 in the presence of EtOH, led to the alcohol **9** in 99 % yield. Benzylation [25] of the primary alcohol in **9** was achieved with benzyl bromide and NaH in DMS solution. Finally, deprotection of the PMB protecting groups in **10** was achieved with reaction with DDQ [26], yielding phomosine H (**5**) in 80 % yield.

The second new metabolite, phomosine I (**6**), was isolated as a colorless solid with m. p. 152-155°C. The IR spectrum showed strong bands for OH groups at 3394 cm^{-1} and for conjugated and/or chelated carbonyl groups at 1652 cm^{-1} . The structure of **6** was established mainly from its ^1H NMR spectrum which showed great similarities with that of methyl ester **5**, except for the missing signal for a methyl ester group. It was deduced that compound **6** was



Scheme 2: Semisynthesis of phomosine H (**5**) from phomosine A (**1**).

the corresponding acid of ester **5**. The MS and ^{13}C NMR spectra, in which 24 carbon peaks could be detected, supported this assumption.

The most polar compound, phomosine J (**7**), was isolated as a colorless solid with m.p. 225-228°C. The IR spectrum showed an absorption band for OH groups at 3510 cm^{-1} and strong carbonyl absorption at 1665 cm^{-1} . The NMR spectra with 21 signals in the ^{13}C NMR and 22 in the ^1H NMR spectra, together with a molecular ion at $m/z = 386$ suggested a molecular composition of $\text{C}_{21}\text{H}_{22}\text{O}_7$. Although both the ^1H NMR and ^{13}C NMR spectra of **7** were in close correspondence with those of **5** and **6**, some new signals could be observed, such as the two olefinic protons at δ 7.27 (d, $J = 16.4$ Hz, 2H, 2'b-H) and 8.13 (d, $J = 16.4$ Hz, 2H, 2'a-H). The *trans* relationship between the proton signals at δ 7.27 and 8.13 was evidenced from the large coupling constant ($J = 16.4$ Hz). A singlet at δ 2.33 was attributed to a conjugated methyl ketone. This connection was evident from the HMBC correlation to a carbonyl carbon appearing at δ 201.7 and long range correlations with two methylene groups at δ 8.13 and 8.13. From this evidence, structure **7** for phomosine H was proposed. It is the aldol condensation product of phomosine A (**1**) with acetone, and its structure was unambiguously proven by base-catalyzed aldol condensation of phomosine A (**1**) with acetone. However, the possibility cannot be excluded that **7** is an artifact generated during the isolation process.

Biological activity: The known phomosines A – D (**1 - 4**) and the novel phomosines H – J (**5 - 7**) were tested in an agar diffusion assay for antimicrobial activity. As displayed in Table 1, the four previously known phomosines A – D (**1 - 4**) were all active against the fungal test organism *Microbotryum violaceum*; phomosines A – C (**1 - 3**) were antibacterial against *Bacillus megaterium* and phomosines A (**1**) and D (**4**) inhibited the alga *Chlorella fusca*. Of the three new metabolites, only phomosine I (**6**) was antifungal and antibacterial.

Table 1: Biological activities of pure metabolites at a concentration of 50 µg against microbial test organisms in an agar diffusion assay [a].

Substance	<i>Microbotryum violaceum</i>	<i>Bacillus megaterium</i>	<i>Chlorella fusca</i>
1	13	10	6PI
2	6	10	0
3	6PI	7	0
4	7PI	0	7
5	0	0	0
6	9	7PI	0
7	0	0	0
Penicillin	0	18	0
Tetracycline	0	18	10PI
Nystatin	20	0	0
Actidione	50	0	35
Acetone	0	0	0

[a] Application of pure substances at a concentration of 50 µg (50 µL of 1 mg/mL). The radius of zone of inhibition was measured in mm. PI = partial inhibition, i.e. there was some growth within the zone of inhibition.

In summary, the highly antifungal and in part antibacterial group of phomosine biarylether natural products has been extensively extended by analysis of the culture extracts of *Phomopsis* sp., internal strain 5686. The four known metabolites were all biologically active, and the new benzyl ether **6** was antifungal as well as antibacterial.

Experimental

General experimental procedures: Column chromatography: commercial silica gel (Merck, 0.040-0.063 mm), and Sephadex LH-20 (Amersham Biosciences). Analytical and preparative thin-layer chromatography (TLC): Precoated silica gel plates (Merck, G60 F-254 or G50 UV-254), respectively. Optical rotation: Perkin–Elmer 241 MC polarimeter at the sodium D line. CD spectra were recorded on a J-810 spectropolarimeter and concentrations are given as mol/dm³. The CD spectra were measured in millidegrees and normalized into $\Delta\epsilon_{\max}$ [l mol⁻¹cm⁻¹]/ λ [nm] units. For solid-state CD protocol, see refs. [21,22]. IR spectra: Nicolet-510P spectrophotometer; ν_{\max} in cm⁻¹. ¹H and ¹³C NMR spectra: Bruker Avance 500 (500 MHz for ¹H and 125 MHz for ¹³C) spectrometer; chemical shifts δ in ppm, coupling constant J in Hz. EI-MS and HR-EI-MS: MAT 8200 and Micromass LCT mass spectrometers, in m/z . Microbiological methods and culture conditions are as described previously [27,28].

Culture, extraction and isolation: *Phomopsis* sp., internal strain no. 5686, was isolated following surface sterilization from *Ligustrum vulgare* that had been growing in Braunschweig, Germany. The endophyte was cultivated at room temperature for 26 days on biomalt semi-solid agar media. The agar was extracted 4 times with ethyl acetate (4 x 0.6 L) to give the crude extract (19.85 g). This was subjected to CC on silica gel, using first gradients of light petroleum-dichloromethane, then dichloromethane, and then gradients of dichloromethane with up to 10% of methanol. On the basis of TLC analysis, similar fractions were combined and were re-chromatographed on silica gel to yield 9.6 g of **1** (*n*-hexane/ethyl acetate = 7:3, R_f = 0.70), 19.1 mg of **2** (*n*-hexane/ethyl acetate = 6.5:3.5, R_f = 0.65),

33.3 mg of **3** (*n*-hexane/ethyl acetate = 6.5:3.5, R_f = 0.63), 16.2 mg of **4** (*n*-hexane/ethyl acetate = 5:5, R_f = 0.68), 12.6 mg of **5** (*n*-hexane/ethyl acetate = 4:6, R_f = 0.64), 14.3 mg of **6** (*n*-hexane/ethyl acetate = 6: 4, R_f = 0.60), and 6.5 mg of **7** (*n*-hexane/ethyl acetate = 5:5, R_f = 0.68), respectively.

Phomosine H [(Methyl 3-(2-(benzyloxymethyl)-3-hydroxy-5-methylphenoxy)-4-hydroxy-2,5-dimethylbenzoate (5)]

MP: 163-165°C.

IR (KBr): ν = 3435, 2953, 1648, 1620, 1580, 1441, 1372, 1296, 1133, 1102, 1090, 1020, 795 cm⁻¹.

UV (CHCl₃): λ_{\max} (log ϵ) = 277 nm (3.70), 310 (3.50).

¹H NMR (500 MHz, CDCl₃): δ = 2.15 (s, 3 H, 5a-H), 2.16 (s, 3 H, 5'a-H), 2.33 (s, 3 H, 2a-H), 3.96 (s, 3 H, CO₂CH₃), 4.72 (s, 2 H, 2'b-H), 4.98 (s, 2 H, 2'a-H), 5.86 (s, 1 H, 6'-H), 6.39 (s, 1 H, 4'-H), 7.33-7.43 (m, 5 H, 2'd-h-H), 11.87 (s, 1 H, 6-OH).

¹³C NMR (125 MHz, CDCl₃): δ = 8.1 (C-5a), 15.4 (C-2a), 21.5 (C-5'a), 51.8 (CO₂CH₃), 63.6 (C-2'a), 72.9 (C-2'b), 104.2 (C-1), 106.2 (C-6'), 108.7 (C-2'), 110.8 (C-5), 111.5 (C-4'), 128.1-128.5 (C-2'd-h), 130.9 (C-2), 133.8 (C-3), 136.9 (C-2'c), 140.3 (C-5'), 152.8 (C-4), 156.1 (C-1'), 156.3 (C-3'), 160.3 (C-6), 172.4 (C-1a).

EIMS (70 eV, 200 °C): m/z (%) = 438 (71) [M⁺], 330 (94), 298 (100), 255(25), 180 (69), 108 (96). HREIMS: m/z 438.1669 (calcd. 438.1679 for C₂₅H₂₆O₇).

Phomosine I [(3-(2-(benzyloxymethyl)-3-hydroxy-5-methylphenoxy)-4-hydroxy-2,5-dimethyl benzoic acid (6)]

MP: 152-155°C.

IR (KBr): ν = 3394, 3236, 2900, 1652, 1610, 1550, 1494, 1400, 1242, 1105, 1005, 784 cm⁻¹.

UV (MeOH): λ_{\max} (log ϵ) = 282 nm (3.50).

¹H NMR (500 MHz, CDCl₃): δ = 2.15 (s, 3 H, 5'a-H), 2.16 (s, 3 H, 5a-H), 2.33 (s, 3 H, 2a-H), 4.73 (s, 2 H, 2'b-H), 4.98 (s, 2 H, 2'a-H), 5.86 (s, 1 H, 6'-H), 6.39 (s, 1 H, 4'-H), 7.34-7.43 (m, 5 H, 2'd-h-H), 11.87 (s, 1H, 6-OH).

¹³C NMR (125 MHz, CDCl₃): δ = 8.1 (C-5a), 15.4 (C-2a), 21.5 (C-5'a), 63.6 (C-2'a), 72.9 (C-2'b), 104.2 (C-1), 106.2 (C-6'), 108.7 (C-2'), 110.8 (C-5), 111.5 (C-4'), 128.1-128.5 (C-2'd-h), 130.9 (C-2), 133.8 (C-3), 136.9 (C-2'c), 140.3 (C-5'), 152.8 (C-4), 156.1 (C-1'), 156.3 (C-3'), 160.3 (C-6), 172.4 (C-1a).

EIMS (70 eV, 200°C): m/z (%) = 406 [M-H₂O]⁺ (100), 391 (26), 378 (32), 377 (38), 360 (21), 348 (27), 327(20).

HREIMS: m/z 424.1509 (calcd. 424.1522 for C₂₄H₂₄O₇).

Phomosine I [(E)-methyl 4-hydroxy-3-(3-hydroxy-5-methyl-2-(3-oxobut-1-enyl)phenoxy)-2,5-dimethyl benzoate (7)]

MP: 225-228°C.

IR (KBr): ν = 3510, 2942, 1656, 1608, 1553, 1414, 1275, 1324, 1209, 1173, 1076, 822 cm⁻¹.

¹H NMR (500 MHz, CDCl₃): δ = 2.08 (s, 3 H, 5'a-H), 2.12 (s, 3 H, 5a-H), 2.21 (s, 3 H, 2a-H), 2.33 (s, 3 H, 2'd-H), 3.86 (s, 3 H, CO₂CH₃), 5.73 (s, 1H, 6'-H), 6.31 (s, 1 H,

4'-H), 7.27 (d, $J = 16.4$ Hz, 2H, 2'b-H), 8.13 (d, $J = 16.4$ Hz, 2H, 2'a-H).

^{13}C NMR (125 MHz, CDCl_3): $\delta = 8.1$ (C-5a), 15.4 (C-2a), 21.7 (C-5'a), 51.8 (CO_2CH_3), 104.6 (C-1), 105.1 (C-6'), 108.3 (C-2'), 110.6 (C-5), 110.6 (C-4'), 128.6 (C-2'b), 130.9 (C-2), 133.8 (C-3), 135.6 (C-2'a), 142.9 (C-5'), 152.8 (C-4), 158.1 (C-1'), 158.5 (C-3'), 159.8 (C-6), 172.3 (C-1a), 201.7 (C-2'c).

EIMS (70 eV, 200 °C): m/z (%) = 386 [M^+] (57), 368 (84), 336 (100), 311 (87), 308 (37), 284 (17), 175 (65).

HREIMS: m/z 386.1358 (calcd. 386.1366 for $\text{C}_{21}\text{H}_{22}\text{O}_7$).

Compound 8: To a stirred suspension of K_2CO_3 (1.29 g, 9.360 mmol) in dry DMF (20 mL) at room temperature was added phomisine A (**1**) (400 mg, 1.156 mmol) and a catalytic amount of TBAI (20 mg). The mixture was stirred for 30 min and then 4-methoxybenzyl chloride (1.086 g, 1.0 mL, 6.936 mmol) was added. The reaction mixture was stirred at room temperature for 8 h. The reaction was quenched with water (100 mL) and extracted 3 times with ethyl acetate (3x100 mL). The combined organic extracts were washed with water (2 x100 mL), brine (100 mL) and dried (Na_2SO_4). After filtration, the solvent was removed to give the residue, which was purified by silica gel CC using light petroleum/EtOAc (7:1) to give **8** as a pale yellow solid in 94% yield (767 mg, 1.086 mmol).

MP: 144°C.

IR (KBr): $\nu = 1648, 1620, 1580, 1295, 1100, 1090, 1027, 795\text{ cm}^{-1}$.

^1H NMR (500 MHz, CDCl_3): $\delta = 2.15$ (s, 3 H, 5a-H), 2.16 (s, 3 H, 5'a-H), 2.33 (s, 3 H, 2a-H), 3.76 (s, 3 H, PMB-OCH₃), 3.82 (s, 3 H, PMB-OCH₃), 3.83 (s, 3 H, PMB-OCH₃), 3.86 (s, 3 H, CO_2CH_3), 4.84 (s, 2 H, CH_2 -PMB), 5.10 (s, 4 H, CH_2 -PMB), 5.90 (s, 1 H, 6'-H), 6.48 (s, 1 H, 4'-H), 6.80 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H), 6.92 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H), 6.95 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H), 7.17 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H), 7.34 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H), 7.40 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H).

^{13}C NMR (125 MHz, CDCl_3): $\delta = 10.2$ (C-5a), 13.1 (C-2a), 22.7 (C-5'a), 52.3 (CO_2CH_3), 55.2 (PMB-OCH₃), 55.3 (PMB-OCH₃), 70.6 (CH_2 -PMB), 75.2 (CH_2 -PMB), 76.1 (CH_2 -PMB), 107.0 (C-6'), 107.2 (C-2'), 112.2 (C-1), 113.8 (C-5), 113.9 (C-4'), 113.8 (PMB-Ar-C), 114.0 (PMB-Ar-C), 128.8 (C-2), 129.1 (PMB-Ar-C), 129.2 (PMB-Ar-C), 129.2 (PMB-Ar-C), 130.2 (C-3), 130.2 (PMB-Ar-C), 141.8 (C-5'), 151.3 (C-4), 156.1 (C-1'), 159.4 (PMB-Ar-C), 159.5 (C-3'), 161.8 (C-6), 168.2 (C-1a).

HREIMS: m/z 706.2768 (calcd. 706.2778 for $\text{C}_{42}\text{H}_{42}\text{O}_{10}$).

Compound 9: A solution of compound **8** (60 mg; 0.086 mmol) in ethanol (20 mL) was stirred at 0° C and sodium borohydride (15 mg, 0.40 mmol) was immediately added. After the slow gas evolution, the solution was stirred overnight and then the solvent was removed. The white residue was dissolved in EtOAc and washed with water (50 mL), brine (50 mL) and dried (Na_2SO_4). After

filtration, the solvent was evaporated to dryness to afford a white residue, which was purified by silica gel column chromatography using PE/EtOAc (5:1) to afford compound **9** as pale yellow solid in 99 % yield (58 mg, 0.083 mmol).

MP: 130°C.

IR (KBr): $\nu = 3220, 1648, 1620, 1580, 1105, 1080, 1030, 780\text{ cm}^{-1}$.

^1H NMR (500 MHz, CDCl_3): $\delta = 2.08$ (s, 3 H, 5a-H), 2.11 (s, 3 H, 5'a-H), 2.20 (s, 3 H, 2a-H), 3.77 (s, 3 H, PMB-OCH₃), 3.82 (s, 3 H, PMB-OCH₃), 3.83 (s, 3 H, PMB-OCH₃), 3.86 (s, 3 H, CO_2CH_3), 4.84 (s, 2 H, CH_2 -PMB), 4.96 (s, 2 H, 2'a-H), 5.04 (s, 4 H, CH_2 -PMB), 5.91 (s, 1 H, 6'-H), 6.49 (s, 1 H, 4'-H), 6.81 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H), 6.92 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H), 6.94 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H), 7.14 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H), 7.33 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H), 7.38 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H).

^{13}C NMR (125 MHz, CDCl_3): $\delta = 10.3$ (C-5a), 13.1 (C-2a), 22.0 (C-5'a), 52.3 (CO_2CH_3), 55.2 (PMB-OCH₃), 55.3 (C-2'a), 55.4 (PMB-OCH₃), 70.0 (CH_2 -PMB), 75.3 (CH_2 -PMB), 76.5 (CH_2 -PMB), 106.7 (C-6'), 107.0 (C-2'), 113.5 (C-1), 113.7 (C-5), 113.8 (C-4'), 113.9 (PMB-Ar-C), 114.2 (PMB-Ar-C), 125.0 (C-2), 128.8 (PMB-Ar-C), 129.2 (PMB-Ar-C), 129.2 (PMB-Ar-C), 130.2 (C-3), 130.2 (PMB-Ar-C), 139.6 (C-5'), 151.7 (C-4), 156.1 (C-1'), 159.4 (PMB-Ar-C), 159.5 (C-3'), 159.6 (C-6), 168.3 (C-1a).

HREIMS: m/z 708.2919 (calcd. 708.2934 for $\text{C}_{42}\text{H}_{44}\text{O}_{10}$).

Compound 10: To a stirred suspension of 60% NaH (20 mg, 0.833 mmol) in dry DMF (20 mL) at room temperature was added protected primary alcohol **9** (100 mg, 0.141 mmol) and a catalytic amount of TBAI (20 mg). The mixture was stirred for 15 min and then benzyl bromide (144 mg, 0.1 mL, 0.846 mmol) was added. The reaction mixture was stirred at room temperature for 3 h. The reaction was carefully quenched with water (5 mL) and extracted 3 times with ethyl acetate (100 mL). The combined organic extracts were washed with water (2 x 100 mL), brine (100 mL) and dried (Na_2SO_4). After filtration, the solvent was removed to give a residue, which was purified by silica gel CC using light petroleum/EtOAc (7:1) to afford compound **10** as pale yellow solid in 92 % yield (103 mg, 1.29 mmol).

MP: 118°C.

IR (KBr): $\nu = 1655, 1620, 1585, 1110, 1080, 1040, 780\text{ cm}^{-1}$.

^1H NMR (500 MHz, CDCl_3): $\delta = 2.07$ (s, 3 H, 5a-H), 2.10 (s, 3 H, 5'a-H), 2.20 (s, 3 H, 2a-H), 3.75 (s, 3 H, PMB-OCH₃), 3.82 (s, 3 H, PMB-OCH₃), 3.83 (s, 3 H, PMB-OCH₃), 3.85 (s, 3 H, CO_2CH_3), 4.60 (s, 2 H, CH_2 -PMB), 4.83 (s, 2 H, 2'a-H), 4.84 (s, 2 H, 2'a-H), 4.85 (s, 2 H, 2'b-H), 5.04 (s, 4 H, CH_2 -PMB), 5.90 (s, 1 H, 6'-H), 6.49 (s, 1 H, 4'-H), 6.81 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H), 6.74 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H), 6.92 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H), 7.10 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H), 7.19-

7.30 [m, 5 H, 2'd-h-H (Bn)], 7.33 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H), 7.38 (d, $J = 8.50$ Hz, 2 H, PMB-Ar-H).

^{13}C NMR (125 MHz, CDCl_3): $\delta = 10.3$ (C-5a), 13.1 (C-2a), 22.1 (C-5'a), 52.3 (CO_2CH_3), 55.2 (PMB-OCH₃), 55.4 (PMB-OCH₃), 60.9 (C-2'a), 70.8 (CH_2 -PMB), 72.5 (C-2'b), 74.9 (CH_2 -PMB), 76.2 (CH_2 -PMB), 106.4 (C-6'), 107.0 (C-2'), 113.6 (C-1), 113.8 (C-5), 113.9 (C-4'), 113.9 (PMB-Ar-C), 114.2 (PMB-Ar-C), 124.8 (C-2), 128.1-128.5 (C-2'd-h), 128.8 (PMB-Ar-C), 129.2 (PMB-Ar-C), 129.2 (PMB-Ar-C), 129.6 (C-3), 130.2 (PMB-Ar-C), 139.0 (C-2'e), 140.1 (C-5'), 151.6 (C-4), 157.2 (C-1'), 159.3 (C-3'), 159.4 (PMB-Ar-C), 159.5 (C-6), 168.4 (C-1a).

HREIMS: m/z 708.2919 (calcd. 708.2934 for $\text{C}_{42}\text{H}_{44}\text{O}_{10}$).

Phomosine H (5): To a stirred solution of **10** (50 mg, 0.062 mmol) in CH_2Cl_2 (15 mL) containing a small amount of water (1/18-1/20 of CH_2Cl_2), DDQ (22 mg, 0.093 mmol) was added at 0°C. The reaction mixture was stirred at room temperature for 5 h. After completion of the reaction, aq. NaHCO_3 (10 mL) was added, and the mixture extracted with CH_2Cl_2 (100 mL). The extract was washed with aq. NaHCO_3 (3x100 mL) and brine (100 mL) and dried over Na_2SO_4 . The solvent was evaporated and the residue was chromatographed on a silica gel column to give phomosine H (**5**) as white solid in 80% yield (21 mg, 0.050 mmol). The synthetic phomosine H (**5**) was identical in all respects (IR, UV, NMR, HREIMS) to natural phomosine H (**5**), including R_f and melting point.

Semisynthesis of phomosine H (7): A mixture of equimolar amounts of **1** (100 mg, 0.289 mmol) and acetone (170 mg, 0.2 mL, 2.890 mmol) in EtOH (10 mL) was warmed at 50°C. A 25% aqueous solution of KOH (2

mL) was added drop-wise to the reaction mixture over a period of 30 min. and further stirred for 5 h at 50°C. The stirring was terminated and the reaction mixture was kept at room temperature overnight. Ice cold water (20 mL) was added until the solid was dissolved. The reaction mixture was acidified with dilute HCl (5 mL), maintaining the temperature at 0-5°C. The precipitates formed were filtered off, dried and crystallized with aqueous ethanol to give the corresponding phomosine H (**7**) in 90 % yield (97 mg, 0.260 mmol).

Agar diffusion test for biological activity: The metabolites were dissolved in acetone at a concentration of 1 mg/mL. Fifty μL of the solution (50 μg) was pipetted onto a sterile filter disk (Schleicher & Schuell, 9 mm), which was placed onto an appropriate agar growth medium for the respective test organism and subsequently sprayed with a suspension of the test organism [19]. The test organisms were the Gram-positive bacterium *Bacillus megaterium* (grown on NB medium), the fungus *Microbotryum violaceum* and the alga *Chlorella fusca* (both grown on MPY medium). (For media recipes, see ref. [29]) Reference substances were penicillin, nystatin, actidione, and tetracycline. Commencing at the middle of the filter disk, the radius of the zone of inhibition was measured in mm. These microorganisms were chosen because (a) they are nonpathogenic and (b) they had in the past proved to be accurate initial test organisms for antibacterial, antifungal, and antialgal/herbicidal activities.

Acknowledgement - We thank the BMBF (Bundesministerium für Bildung und Forschung, Project-No.: SUA 07/003).

References

- [1] Meng L, Sun P, Li L, Draeger S, Schulz B, Krohn K, Yia Y, Zhang W. (2011) Endophytic fungus *Penicillium chrysogenum*, a new source of hypocrellins. *Biochemical Systematics and Ecology*, **39**, 163-165.
- [2] (a) Schulz B, Boyle C, Draeger S, Römmert A, Krohn K. (2002) Endophytic fungi: a source of novel biologically active secondary metabolites. *Mycological Research*, **106**, 996-1004; (b) Tan RX Zou WZ. (2001) Endophytes: a rich source of functional metabolites. *Natural Products Reports*, **18**, 448-459.
- [3] Cole RJ. (1981) *Handbook of Toxic Fungal Metabolites*. Cox RH. (Ed.), Academic Press, New York, London.
- [4] Wheeler MM, Wheeler DMS, Peterson GW. (1975) Anthraquinone pigments from the phytopathogen *Phomopsis juniperovora* Hahn. *Phytochemistry*, **14**, 288-289.
- [5] Izawa Y, Shimizu T, Koyama K, Natori S. (1989) Six new 10-pheynl-[11]cytochalasans, cytochalasins N - S from *Phomopsis* sp. *Tetrahedron*, **45**, 2323-2335.
- [6] Tomioka T, Izawa Y, Koyama K, Natori S. (1987) Three 10-phenyl [11]cytochalasana, cytochalasins N, O, and P from *Phomopsis* sp. *Chemical & Pharmaceutical Bulletin*, **35**, 902-905.
- [7] Horn WS, Simmonds MSJ, Schwartz RE, Blaney WM (1995) Phomopsichalasin, a novel antimicrobial agent from an endophytic *Phomopsis* sp. *Tetrahedron*, **51**, 3969-3978.
- [8] Tomoda H, Namatame I, Si S, Kawaguchi K, Masuma R, Namikoshi M, Omura S. (1999) Phenochalasin, inhibitors of lipid droplet formation in mouse macrophages produced by *Phomopsis* sp. FT-0211. *Journal of Antibiotics*, **52**, 851-856.
- [9] Tsantrizos YS, Ogilvie KK, Watson AK. (1992) Phytotoxic metabolites of *Phomopsis convolvulus*, a host-specific pathogen of field bindweed. *Canadian Journal of Chemistry*, **70**, 2276-2284.
- [10] Isaka M, Jaturapat A, Rukseree K, Danwisetkanjana K, Tantichareon M, Thebtaranonth Y. (2001) Phomoxanthenes A and B, novel xanthone dimers from the endophytic fungus *Phomopsis* species. *Journal of Natural Products*, **64**, 1015-1018.
- [11] Elsässer B, Krohn K, Flörke U, Root N, Aust H-J, Draeger S, Schulz B, Antus S, Kurtán T. (2005) X-Ray structure determination, absolute configuration and biological activity of phomoxanthone A. *European Journal of Organic Chemistry*, 4563-4570.
- [12] Rukachaisirikul V, Sommart U, Phongpaichit S, Sakayaroj J, Kirtikara K. (2008) Metabolites from the endophytic fungus *Phomopsis* sp. PSU-D15. *Phytochemistry*, **69**, 783-787.
- [13] Claydon N, Grove JF, Pople M. (1985) Elm bark beetle boring and feeding deterrents from *Phomopsis oblonga*. *Phytochemistry*, **24**, 937-943.

- [14] Dai J, Krohn K, Gehle D, Kock I, Flörke U, Aust H-J, Draeger S, Schulz B, Rheinheimer J. (2005) New oblongolides isolated from the endophytic fungus *Phomopsis* sp. from *Melilotus dentata* from the shores of the Baltic Sea. *European Journal of Organic Chemistry*, 4009-4016.
- [17] Hussain H, Akhtar N, Draeger S, Schulz B, Pescitelli G, Salvadori P, Antus S, Kurtán T, Krohn K. (2009) New bioactive 2,3-epoxycyclohexenes and isocoumarins from the endophytic fungus *Phomopsis* sp. from *Laurus azorica*. *European Journal of Organic Chemistry*, 749-756.
- [18] Edgar JA, Frahn JL, Cockrum PA, Culvenor CCJ. (1986) Lupinosis. the chemistry and biochemistry of phomopsins. *Bioactive Molecules*, 169-184.
- [19] Nouguier R, Bertrand MP, Picon P, Perfetti P. (1994) Diastereoselective synthesis of a methylated derivative of phomozin, a phytotoxin isolated from *Phomopsis helianthi*, a phytopathogenic fungus of sunflowers. *Tetrahedron Letters*, **35**, 8171-8172.
- [20] Huang Z, Cai X, Shao C, She Z, Xia X, Chen Y, Yang J, Zhou, S Lin Y. (2008) Chemistry and weak antimicrobial activities of phomopsins produced by mangrove endophytic fungus *Phomopsis* sp. ZSU-H76. *Phytochemistry*, **69**, 1604-1608.
- [21] Du X, Lu C, Li Y, Zheng Z, Su W, Shen Y (2008) Three new antimicrobial metabolites of *Phomopsis* sp. *Journal of Antibiotics*, **61**, 250-253.
- [22] Davis RA, Andjic V, Kotiw M, Shivas RG. (2005) Phomoxins B and C: Polyketides from an endophytic fungus of the genus *Eupenicillium*. *Phytochemistry*, **66**, 2771-2775.
- [23] Krohn K, Michel A, Roemer E, Flörke U, Aust H-J, Draeger S, Schulz B, Wray V. (1995) Biologically active metabolites from fungi, 6. Phomosines A-C. Three new biaryl ethers from *Phomopsis* sp. *Natural Product Letters*, **9**, 309-314.
- [24] Dai J, Krohn K, Flörke U, Gehle D, Aust H-J, Draeger S, Schulz B, Rheinheimer K. (2005) Novel highly substituted biaryl ethers, phomosines D-G, isolated from the endophytic fungus *Phomopsis* sp. from *Adenocarpus foliolosus*. *European Journal of Organic Chemistry*, 5100-5105.
- [25] Kanai K, Sakamoto I, Ogawa S, Suami T. (1987) Synthesis of 1,4-diaminocyclitol antibiotics. III. Synthesis of 4-hydroxy-purpurosamine B derivatives. *Bulletin of the Chemical Society of Japan*, **60**, 1529-1531.
- [26] Horita K, Yoshioka T, Tanaka T, Oikawa Y, Yonemitsu O. (1986) On the selectivity of deprotection of benzyl, mpm (4-methoxybenzyl) and dmpm (3,4-dimethoxybenzyl) protecting groups for hydroxy functions. *Tetrahedron*, **42**, 3021-3028.
- [27] Schulz B, Sucker J, Aust H-J, Krohn K, Ludewig K, Jones PG, Doering D. (1995) Biologically active secondary metabolites of endophytic *Pezizula* species. *Mycological Research*, **99**, 1007-1015.
- [28] Höller U, Wright AD, Matthée GF, König GM, Draeger S, Aust H-J, Schulz B. (2000) Fungi from marine sponges: Diversity, biological activity and secondary metabolites. *Mycological Research*, **104**, 1354-1365.

Isolation and Characterization of a new Benzofuran from the Fungus *Alternaria* sp. (HS-3) Associated with a Sea CucumberXuekui Xia¹, Jun Qi³, Fang Wei⁴, Airon Jia², Wenpeng Yuan², Xiumei Meng², Miansong Zhang¹, Changheng Liu^{2*} and Changyun Wang³¹Key Laboratory for Applied Microbiology of Shandong Province, Biotechnology Center of Shandong Academy of Sciences, Jinan 250014, P. R. China²Biology Institute of Shandong Academy of Sciences, Jinan, 250014, P. R. China³Key Laboratory of Marine Drugs, Ministry of Education, School of Medicine and Pharmacy, Ocean University of China, Qingdao, 266003, P. R. China⁴Food Engineering College, Dalian Ocean University, Dalian, 116023, P. R. China

liuchh@keylab.net

Received: May 15th, 2011; Accepted: August 24th, 2011

A new compound, 4-acetyl-5-hydroxy-3, 6, 7-trimethylbenzofuran-2(3H)-one (**1**), together with two known compounds, 2-carboxy-3-(2-hydroxypropyl) phenol (**2**) and 5-methyl- 6-hydroxy-8-methoxy-3-methylisochroman (**3**) were isolated from the fungus *Alternaria* sp. (HS-3) associated with a sea cucumber from the Yellow Sea in China. Their structures were elucidated by spectral methods.

Keywords: sea cucumber, *Alternaria* sp, metabolism, benzofuran.

Marine-derived fungi have been the sources of novel second metabolites, a considerable number of which display promising biological and pharmacological properties [1]. Seven new indole diketopiperazines, along with twelve known ones, were isolated from the holothurian-derived fungus *Aspergillus fumigatus* [2]. Diterpene glycosides from the fungus *Acremonium striatisporum* isolated from a sea cucumber, *Eupentacta fraudatrix*, exhibited cytotoxic action against Ehrlich carcinoma cells [3-5]. In this article, we report the isolation of a new benzofuran derivative (**1**), together with two known compounds, 2-carboxy-3-(2-hydroxypropyl) phenol (**2**) [6] and 5-methyl-6-hydroxy-8-methoxy-3-methylisochroman (**3**) [7], from the fungus *Alternaria* sp. (HS-3) associated with a sea cucumber.

Compound **1** was assigned as C₁₃H₁₄O₄, as determined by HREIMS. The IR absorptions at 3385 and 1699 cm⁻¹ reflected the presence of hydroxyl and carbonyl group(s). In the ¹H NMR spectrum, there was a chelating phenolic hydroxyl signal at δ 10.8, a methyl at δ 1.57 (d, 6.0 Hz), a methine at δ 4.30 (q, 7.2 Hz), and three methyl singlets at δ 2.31, 2.24 and 2.21. The ¹³C NMR spectrum disclosed the presence of two carbonyl (δ 197.4 and 167.7), four methyl (δ 22.3, 17.3, 14.5 and 11.7), six substituted aromatic carbons (δ 159.2, 148.5, 136.6, 125.3, 124.5 and 102.7), and a methine carbon at δ 39.1. ¹H-¹H COSY revealed a contiguous sequence of coupled signals from H-9 to H-13. The HMBC data (Table 1), especially the correlations from the OH-5 (δ 10.8) to C-4, C-5 and C-6 unambiguously

assigned the positions of the hydroxyl groups. Additionally, the correlations from H-9 to C-2, C-3, C-4 and C-8, indicated that the methine was attached to C-3 of the benzene ring. Thus, compound **1** was established as 4-acetyl-5-hydroxy-3, 6, 7-trimethylbenzofuran-2(3H)-one.




Figure 1: Structures of compounds **1**, **2** and **3**.

Table 1: ¹H and ¹³C NMR (600 and 150 MHz, δ value, J in Hz) spectral data and HMBC correlations of **1** (in acetone-*d*₆).

Position	¹³ C NMR	¹ H NMR	HMBC (C to H)
2	148.5 (C)		H-8, 9
3	136.6 (C)		H-9, 13
4	102.7 (C)		5-OH, H-9, 14
5	159.2 (C)		5-OH, H-11
6	124.5 (C)		5-OH, H-8, 11
7	125.3 (C)		H-8, 11
8	14.5 (CH ₃)	2.31	
9	39.1 (CH)	4.30 (q, 7.2)	H-13
10	197.4 (C)		H-14
11	11.7 (CH ₃)	2.24	
12	167.7 (C)		H-9, 13
13	17.3 (CH ₃)	1.57 (d, 6.0)	H-9
14	22.3 (CH ₃)	2.21 (s)	
5-OH		10.80 (s)	

Experimental

General: Melting points were obtained on a Fisher-Johns hot-stage apparatus and were uncorrected. 1D and 2D

NMR, Varian INOVA-600 MHz spectrometer; MS, VG-ZAB mass spectrometer; IR, Nicolet 5DX-FTIR spectrophotometer; CC was carried out on silica gel (200-300 mesh; Qingdao Haiyang Chemicals), octadecylsilyl silica gel (Unicorn; 45-60 μ m) and Sephadex LH-20 (Amersham Biosciences). Precoated silica gel plates (Yan Tai Zi Fu Chemical Group Co.; G60, F-254) were used for thin layer chromatography.

Fungal material: A strain (HS-3) of the fungus *Alternaria* sp. was isolated from the surface muscle of a sea cucumber from Weihai in the Yellow Sea, China. Voucher specimens are stored in the Biology Institute, Shandong Academy of Sciences, Jinan, PR China. The almost complete 18S rRNA gene sequence of strain HS-3 was deposited in GenBank under the no. JF694748.

Identification of the endophytic isolate: The endophytic fungus was grown on PDA for 5 days at 28°C. Genomic DNA was extracted and purified using the Fungal DNA Kit 50 (OMEGA, USA), according to the manufacture's instructions, suitably modified. For identification and differentiation, the Internal Transcript Spacer regions (ITS1F and ITS2) and the intervening 5.8S rRNA region was amplified and sequenced. The ITS regions of the fungus were amplified by PCR with the universal ITS primers, ITS1F (5'-CTT GGT CAT TTA GAG GAA GTAA-3') and ITS4 (5'-TCC TCC GCT TAT TGA TAT GC-3'). The PCR products were then purified and desalted using the EZ Spin column PCR product purification kit (BBI) and sequenced. The sequencing results were aligned with the nucleotide-nucleotide database (BLASTn) of the U.S. National Center for Biotechnology Information (NCBI) for final identification of the endophytic isolate.

Fermentation, extraction, and isolation: The fungal strain was cultivated in 50 L lipid medium at 28°C without shaking for 28 days. Two hundred and fifty 500 mL Erlenmeyer flasks each contained 2.0 g glucose, 0.4 g yeast extract and 0.4 g peptone dissolved in 200 mL of seawater. The cultures (50 L) were filtered through cheesecloth. The filtrate was concentrated to 5 L below 50°C and extracted 3 times by shaking with an equal volume of ethyl acetate. The combined organic extracts

were subjected to silica gel CC, eluting with a gradient of light petroleum to ethyl acetate. Fractions of 40 mL were collected and combined by TLC examination. Fractions containing the desired compounds were further purified by Sephadex LH-20 chromatography, eluting with mixtures of light petroleum-CHCl₃-MeOH (2:1:1), which yielded compounds **1** (5.0 mg), **2** (31.0 mg), and **3** (6.0 mg).

4-Acetyl-5-hydroxy-3,6,7-trimethylbenzo-furan-2(3H)-one (**1**)

Colorless needles, MP: 113-115°C.

IR (KBr) ν_{\max} : 3385, 3208, 3005, 2963, 2935, 1800, 1699, 1601, 1577 cm⁻¹.

¹H and ¹³C NMR (acetone-*d*₆): Table 1.

EIMS: [M]⁺ 234, [M-CO]⁺ 206, [M-CH₃CO]⁺ 191, HREIMS: 234.0887, relative intensity 100%, C₁₃H₁₄O₄.

2-Carboxy-3-(2-hydroxypropanyl) phenol (**2**)

White solid.

¹H NMR (600 MHz, acetone-*d*₆): δ 1.22 (3H, d, *J* = 6.6 Hz), 3.02 (2H, d, *J* = 7.4 Hz), 4.36 (1H, m), 6.78 (1H, d, *J* = 8.0 Hz), 7.07 (1H, dd, *J* = 8.0, 0.9 Hz), 7.32 (1H, dd, *J* = 8.0, 8.0 Hz), 11.74 (-OH, s).

¹³C NMR (150 MHz, acetone-*d*₆): δ 20.9, 34.8, 76.9, 108.4, 116.4, 118.0, 136.3, 139.5, 162.4, 170.1.

ESIMS: *m/z* [M-1]⁻ 195.

5-Methyl-6-hydroxy-8-methoxy-3-methylisochroman (**3**)

White solid.

¹H NMR (600 MHz, CDCl₃): δ 1.38 (3H, d, *J* = 6.3 Hz), 2.04 (3H, s), 2.42 (1H, dd, *J* = 16.5, 10.6 Hz), 2.61 (1H, dd, *J* = 16.5, 2.3 Hz), 3.73 (1H, m), 3.70 (3H, s), 4.57 (1H, d, *J* = 15.0 Hz), 4.89 (1H, d, *J* = 15.0 Hz), 6.23 (1H, s).

¹³C NMR (150 MHz, CDCl₃): δ 10.0, 21.7, 34.0, 55.0, 64.6, 70.6, 96.2, 112.9, 115.2, 134.1, 152.7, 153.9.

FABMS: *m/z* [M+1]⁺ 209.

Acknowledgement - We wish to acknowledge the financial support of the National Natural Science Foundation of China (31000008), Prize Fund for Excellent Middle Aged and Young Scientists of Shandong Province (BS2009HZ004), Doctoral Prize of Shandong Academy of Sciences (2008, No. 25).

References

- [1] Rateb ME, Rainer E. (2011) Secondary metabolites of fungi from marine habitats. *Natural Product Report*, **28**, 290-344.
- [2] Wang FZ, Fang YC, Zhu TJ, Zhang M, Lin AQ, Gu QQ, Zhu WM. (2008) Seven new prenylated indole diketopiperazine alkaloids from holothurian-derived fungus *Aspergillus fumigatus*. *Tetrahedron*, **64**, 7986-7991.
- [3] Afiyatullo SS, Kalinovsky AI, Pivkin MV, Dmitrenok PS, Kuznetsova TA. (2006) New diterpene glycosides of the fungus *Acremonium striatisporum* isolated from a sea cucumber. *Natural Product Research*, **20**, 902-908.
- [4] Afiyatullo SS, Kuznetsova TA, Isakov V.V, Pivkin MV, Prokofeva NG, Elyakov GB. (2000) New diterpenic altrosides of the fungus *Acremonium striatisporum* isolated from a sea cucumber. *Journal of Natural Products*, **63**, 848-850.
- [5] Afiyatullo SS, Kalinovsky AI, Kuznetsova TA, Pivkin MV, Prokofeva NG, Dmitrenok PS, Elyakov GB. (2004) New glycosides of the fungus *Acremonium striatisporum* isolated from a sea cucumber. *Journal of Natural Products*, **67**, 1047-1051.
- [6] Rahman MM, Sarker SD, Byres M, Gray AI. (2004) New salicylic acid and isoflavone derivatives from *Flemingia paniculata*. *Journal of Natural Products*, **67**, 402-406.
- [7] Masuma R, Tabata N, Tomoda H, Haneda K, Iwai Y, Omura S. (1994) Arohynapenes A and B, new anticoccidial agents produced by *Penicillium* sp. Taxonomy, fermentation, and structure elucidation. *Journal of Antibiotics*, **47**, 46-53.

Potent Toxic Macrocyclic Trichothecenes from the Marine-Derived Fungus *Myrothecium verrucaria* Hmp-F73

Li Zhao^{a,b}, Li Liu^b, Nan Wang^a, Shu-Jin Wang^b, Jing-Chun Hu^{b*} and Jin-Ming Gao^{a*}

^aResearch Centre for Natural Medicinal Chemistry, College of Science, Northwest A & F University, Yangling, Shaanxi 712100, China

^bInstitute of Applied Ecology, Chinese Academy of Sciences, Shenyang 110161, China

jinminggaocn@yahoo.com.cn (J.M. Gao); hujc@iae.ac.cn (J.C. Hu)

Received: July 12th, 2011; Accepted: October 24th, 2011

Activity-guided fractionation of a methanol extract from the culture broth of *Myrothecium verrucaria* Hmp-F73, a fungus associated with the sponge *Hymeniacidon perleve*, afforded six macrocyclic trichothecenes, verrucarins J (1), 8-hydroxyverrucarin J (2), verrucarins A (3), 8-acetoxyroridin H (4), isororidin E (5), and roridin E (6), along with trichoverrin B (7). All seven metabolites displayed potent toxicity to the brine shrimp (*Artemia salina*). In addition, compounds 2, 3, and 6 showed weak phytotoxic activities against lettuce seeds. A preliminary structure-activity relationship of the metabolites is also discussed.

Keywords: *Myrothecium verrucaria* Hmp-F73, *Hymeniacidon perleve*, trichothecenes, mycotoxins, brine shrimp lethality.

In connection with our ongoing screening for bioactive secondary metabolites from fungi [1], we screened in a brine shrimp lethality assay the MeOH extracts of 98 fungal strains associated with the marine sponge *Hymeniacidon perleve* collected in Dalian. The extract of a strain of *Myrothecium verrucaria* Hmp-F73 exhibited potent toxic activity to the brine shrimp. Through activity-guided fractionation, we isolated seven trichothecene derivatives (1-7; Figure 1), namely verrucarins J (1), 8-hydroxyverrucarin J (2), verrucarins A (3), 8-acetoxyroridin H (4), isororidin E (5), roridin E (6) and trichoverrin B (7) [2]. Their structures were identified by comparison of their spectroscopic data with those published [2].

Compounds 1-7 were evaluated for their toxicity toward brine shrimp larvae and phytotoxic activity against lettuce seed germination. As shown in Table 1, they exhibited relatively strong lethality activities to the brine shrimp after exposure for 24 h, with LC₅₀ values ranging between 1 ng/mL and 1.1 µg/mL. Notably, verrucarins (1, in particular 2 and 3) were much more toxic than roridins (4 and 6) and trichoverrin B (7), suggesting that the presence of an ester carbonyl group at position C-6' is essential for activity. Besides, it seems that mono-hydroxylation of a double bond at C-2' of the macrocyclic ring of a verrucaroid (I) series is able to highly improve the toxicity (1 and 2 vs 3), indicating that the activity is associated with the presence of the 2'-OH substituent. For example, verrucarins A (3) showed 6- and 7-fold more activity than verrucarins J (1) and 8-hydroxyverrucarin J (2), respectively. Among the compounds tested, verrucarins A (3) exerted the best activity (LC₅₀ = 0.001 µg/mL), while




Figure 1: Structures of compounds 1-7.

trichoverrin B (7) produced the least (LC₅₀ = 1.138 µg/mL) since it was a nonmacrocyclic trichothecene, thereby indicating that an intact macrolide ring is also an important requirement for a higher toxic effect. Interestingly, verrucarins A (3) was reported to display no significant difference in the inhibition of protein synthesis in Vero cells and rat spleen lymphocytes. Thus, the *in vitro* cell response of a given trichothecene is not always an accurate predictor of toxicity in whole animals [3]. Our findings demonstrated for the first time that some modification of the macrolide ring moiety could highly affect the toxicity of trichothecene derivatives.

Table 1: Brine shrimp lethality and phytotoxicity of compounds **1-7**^a.

compound	brine shrimp assay LC ₅₀ (μg/mL)	phytotoxicity assay 100 μg/mL
verrucarin J (1)	0.006	inactive
isororidin E (5)		
8-hydroxyverrucarin J (2)	0.007	20%
verrucarin A (3)	0.001	30%
8-acetoxysororidin H (4)	0.15	inactive
roridin E (6)	0.221	50%
trichoverrin B (7)	1.138	inactive

^aCompounds **1** and **5** were tested as a mixture (3:1).

Moreover, the phytotoxic activity of the compounds **1-7** toward the lettuce seed germination in a Petri dish bioassay was not observed at a concentration of 1 μg/mL. Only when the concentration was up to 100 μg/mL did the compounds **2**, **3** and **6** exerted a weak inhibiting activity.

From the present study, it can be concluded that the toxicity of the trichothecenes to brine shrimp is not related to the phytotoxicity potency. The trichothecenes **1-7**, identified from *M. verrucaria* Hmp-F73, were found to be most toxic against the brine shrimp larvae. This may constitute chemical defense of the sponge itself to protect it against predators like fishes and shrimps.

Experimental

General: ¹H and ¹³C NMR spectra were acquired using a Bruker ARX 300 MHz spectrometer. ESI MS data were recorded using a Waters Micromass[®] ZQ mass spectrophotometer in positive ion mode.

Microorganism and culture conditions: Fungal strain Hmp-F73 was isolated from the marine sponge *Hymeniacidon perleve* collected from the Bohai Sea at the shore off Lingshui Qiao in Dalian, China, and identified as *Myrothecium verrucaria* (Genbank accession number: HQ625520). Seed culture was prepared with a 500 mL Erlenmeyer flask containing 100 mL of culture medium

containing (g/L): potato 200 g, glucose 20 g and cultivated at 28°C for 48 h with shaking at 150 rpm. The seed culture was inoculated to 3 L Erlenmeyer flasks (×16) containing 1000 mL culture medium, the same as with the seed culture, and incubated for 4 days, at 28°C and 150 rpm on rotary shakers.

Extraction and isolation: To the cell-free broth (filtered, 16 L) of Hmp-F73, 60 mL of HP20 resin per L was added. After shaking at 100 rpm for 1 h, the resin was recovered and successively washed with deionized H₂O and MeOH. After evaporation of MeOH, the resulting crude extract (9.66 g) was subjected to vacuum flash chromatography over silica gel with ethyl acetate-*n*-hexane (0:10→10:0) and pure MeOH to provide Fr.1-Fr.6. Compounds **1** and **5** (26.7 mg, inseparable 3:1), **4** (7.7 mg) and **6** (15.4 mg) were purified from Fr. 3 (86.6 mg) by RP-HPLC (Dionex U3000, USA) using a YMC-pack ODS-A column (5 μm, 250 × 10 mm) eluting with 70% MeOH at a flow rate of 2.5 mL/min with UV detection at 220 nm. Similarly, semi-preparative HPLC of Fr. 4 (90.0 mg) eluting with 77% MeOH afforded **3** (6.4 mg), and purification of Fr. 5 (389.9 mg) with 55% MeOH yielded compounds **2** (6.5 mg) and **7** (13.5 mg).

Brine shrimp lethality assay: This assay was performed according to the reported method [1]. LC₅₀ values for the test compounds were analyzed and calculated with SPSS 13.0.

Phytotoxicity assay: The Petri dish bioassay was carried out by the reported method [4].

Acknowledgments - This work was supported by a grant from the National High Technology Research and Development Program of China (863 Program 2007AA09Z417).

References

- [1] Qin JC, Zhang YM, Hu L, Ma YT, Gao JM (2009) Cytotoxic metabolites produced by *Alternaria* No.28, an endophytic fungus isolated from *Ginkgo biloba*. *Natural Product Communications*, **4**, 1473–1476.
- [2] (a) Breitenstein W, Tamm C. (1978) Verrucarine and voridine.35. Partial synthesis of tetrahydroverrucarin-J. *Helvetica Chimica Acta*, **61**, 1975-1983; (b) Jarvis BB, Midiwo JO, Mazzola EP. (1984) Antileukemic compounds derived by chemical modification of macrocyclic trichothecenes.2. Derivatives of roridin-A and roridin-H and verrucarine-A and verrucarine-J. *Journal of Medicinal Chemistry*, **27**, 239-244; (c) Wagenaar MM, Clardy J. (2001) Two new roridins isolated from *Myrothecium* sp. *Journal of Antibiotics*, **54**, 517-520; (d) Jarvis, BB, Desilva T, McAlphine JB, Swanson SJ, Whittern DN. (1992) New trichoverroids from *Myrothecium verrucaria* isolated by high-speed countercurrent chromatography. *Journal of Natural Products*, **55**, 1441-1446.
- [3] Thompson WL, Wannemacher RW. (1986) Structure-function-relationships of 12,13-epoxytrichothecene mycotoxins in cell-culture - comparison to whole animal lethality. *Toxicon*, **24**, 985-994.
- [4] (a) Abbas HK, Johnson BB, Shier WT, Tak H, Jarvis BB, Boyette CD. (2002) Phytotoxicity and mammalian cytotoxicity of macrocyclic trichothecene mycotoxins from *Myrothecium verrucaria*. *Phytochemistry*, **59**, 309-313; (b) Andolfi A, Boari A, Evidente A, Vurro M. (2005) Metabolites inhibiting germination of *Orobancha ramosa* seeds produced by *Myrothecium verrucaria* and *Fusarium compactum*. *Journal of Agricultural and Food Chemistry*, **53**, 1598-1603.

Synthesis and Bioactivity of Novel Coumarin Derivatives

Ai-Ying Guan, Chang-Ling Liu*, Miao Li, Zhi-Nian Li, Ming-Xing Zhang and Hong Zhang

State Key Laboratory of the Discovery and Development of Novel Pesticide,
Shenyang Research Institute of Chemical Industry Co. Ltd., Shenyang, 110021, China

liuchangling@vip.163.com


Received: July 12th, 2011; Accepted: October 26th, 2011

A series of coumarin derivatives (**6-8**) containing (*E*)-methyl 2-(methoxyimino)-2-phenylacetate, (*E*)-2-(methoxyimino)-*N*-methyl-2-phenylacetamide and methyl methoxy(phenyl)carbamate were synthesized from substituted resorcinols (**1**) and substituted β -keto esters (**2**) as starting material via cyclization and condensation reactions. The test results indicated that (*E*)-methyl 2-{2-[(3-hexyl-4-methyl-coumarin-7-yloxy)methyl]phenyl}-2-(methoxyimino)acetate (**6f**) was the optimal structure with good fungicidal activity against cucumber grey mold (CGM) giving 100% control at 100 mg L⁻¹ concentration, much higher than that of coumoxystrobin. Methyl 2-[(3,4-dimethyl-coumarin-7-yloxy)methyl]phenyl(methoxy)carbamate (**8a**) was another optimal structure with good fungicidal activity against wheat powdery mildew (WPM) showing 100% control at 50 mg L⁻¹ concentration, at the same level as that of the commercial kresoxim-methyl, and very significantly higher than that of coumoxystrobin (no control against WPM at 400 mg L⁻¹).

Keywords: Coumarins, strobilurins, intermediate derivatization method, fungicidal activity, insecticidal activity.

Substituted coumarins and methoxyacrylates derived from strobilurin A exhibit a broad spectrum of bioactivities [1-4]. A series of coumarin derivatives containing (*E*)-methyl 3-methoxy-2-phenylacrylate (MA) were synthesized and bioassayed in our previous study [5a-5c], which resulted in the discovery of coumoxystrobin (SYP-3375), (*E*)-methyl 2-{2-[(3-*n*-butyl-4-methylcoumarin-7-yloxy) methyl] phenyl}-3-methoxyacrylate, having a broad spectrum fungicidal activity against cucumber downy mildew (CDM), cucumber grey mold (CGM), wheat powdery mildew (WPM), rice sheath blight (RSB) and apple valsa canker (AVC). Other non-acrylate compounds, such as Kresoxim-methyl [5d], Dimoxystrobin [6], Orystrobin [7], Pyraclostrobin [8] and Pyrametostrobin (SYP-4155) [9] containing related pharmacophores such as (*E*)-methyl 2-(methoxyimino)-2-phenylacetate (OE), (*E*)-2-(methoxyimino)-*N*-methyl-2-phenylacetamide (OA) and methyl methoxy(phenyl) carbamate (MC), were shown to have excellent fungicidal activity. A number of coumarin derivatives containing OE, OA and MC were synthesized using the intermediate derivatization method based on either bioisosteric replacement [5b,5c,10-12] or active substructure combination [13]. The fungicidal activity and the structure-activity relationship are also discussed in this paper.

Synthesis: As shown in Scheme 1, according to known methods [14,15,16a], substituted coumarins (**3**) were prepared from substituted resorcinols (**1**) and substituted β -keto esters (**2**). Then, intermediates (**3**) were first treated with 1 equiv. of (*E*)-methyl 2-(2-(bromomethyl) phenyl)-2-(methoxyimino)acetate (**4**) in butanone at refluxing



Scheme 1: Reagents and conditions: (a) c. H₂SO₄/5°C-room temperature; (b1) K₂CO₃ in butanone, reflux; (b2) NaH in DMF, room temperature; (c) CH₃NH₂ in CH₃OH, room temperature.

temperature for 5 h to afford compounds (**6**) in 80–90% yields, followed by reacting with 2.1 equiv. of methylamine in CH₃OH at room temperature for 3 h to obtain target compounds (**7**) in 85–90% yields; The desired compounds (**8**) were synthesized via the reaction of intermediates (**3**) with 1 equiv. of methyl 2-(bromomethyl)phenyl(methoxy)carbamate (**5**) in DMF at room temperature for 3 h in 65–90% yields.

Biological activities: Evaluations of biological activities of the title compounds were performed as previously reported [16b-16d]. Biological data were reported in the range 0 (indicates no control) to 100% (complete control). The test results of fungicidal activities of the title compounds are listed in Tables 1 and 2.

Table 1: Chemical structures, physical properties and fungicidal activity against CDM of synthesized coumarin analogues.

Compd	Substituents				mp (°C)	Yield (%)	Fungicidal activity (% control at given concentration mg L ⁻¹)					
	R ₁	R ₂	R ₃	Q			CDM					
							400	200	100	50	25	6.25
6a	H	H	H	Q ₁	106-110	80	70	60	/	/	/	/
6b	CH ₃	H	H	Q ₁	150-152	85	70	70	/	/	/	/
6c	CH ₃	CH ₃	H	Q ₁	116-117	88	95	95	90	80	55	30
6d	CH ₃	H	CH ₃	Q ₁	178-180	81	0	0	/	/	/	/
6e	CH ₃	Cl	CH ₃	Q ₁	198-200	83	0	0	/	/	/	/
6f	CH ₃	<i>n</i> -C ₃ H ₇	H	Q ₁	121-122	80	60	/	/	/	/	/
6g	CH ₃	<i>n</i> -C ₄ H ₉	H	Q ₁	100-102	86	100	98	90	95	80	45
6h	CH ₃	<i>n</i> -C ₆ H ₁₃	H	Q ₁	75-78	87	100	100	100	100	100	85
6i	<i>n</i> -C ₃ H ₇	Cl	H	Q ₁	136-138	83	0	/	/	/	/	/
6j	Ph	Cl	H	Q ₁	166-168	81	98	80	60	50	/	/
7a	H	H	H	Q ₂	129-130	88	80	50	/	/	/	/
7b	CH ₃	H	H	Q ₂	213-214	90	80	80	/	/	/	/
8a	CH ₃	CH ₃	H	Q ₃	131-133	68	100	95	90	100	90	70
8b	CH ₃	Cl	H	Q ₃	139-141	79	100	85	70	85	60	20
8c	CH ₃	F	H	Q ₃	114-116	75	100	75	55	/	/	/
8d	CH ₃	<i>n</i> -C ₄ H ₉	H	Q ₃	65-67	70	0	/	/	/	/	/
8e	CH ₂ CH ₂ CH ₂	H	Q ₃	Q ₃	177-179	85	0	/	/	/	/	/
8f	CH ₂ CH ₂ CH ₂	CH ₃	Q ₃	Q ₃	150-151	88	80	/	/	/	/	/
8g	CH ₂ CH ₂ CH ₂ CH ₂	H	Q ₃	Q ₃	160-161	92	0	/	/	/	/	/
8h	CH ₂ CH ₂ CH ₂ CH ₂	CH ₃	Q ₃	Q ₃	167-169	90	80	/	/	/	/	/
coumoxystrobin							100	100	100	100	100	95
kresoxim-methyl							/	/	/	/	30	0

note: “/” stands for no data

Table 2: Fungicidal activity against RSB, CGM and WPM of synthesized coumarin analogues.

Compd	Fungicidal activity (% control at given concentration mg L ⁻¹)									
	RSB		CGM			WPM				
	400	200	100	50	25	400	200	100	50	25
6a	0	0	/	/	/	50	/	/	/	/
6b	90	70	/	/	/	50	/	/	/	/
6c	80	0	/	/	/	30	/	/	/	/
6d	90	0	/	/	/	0	/	/	/	/
6e	40	0	/	/	/	0	/	/	/	/
6f	0	100	100	100	60	100	100	95	85	/
6g	90	20	/	/	/	100	50	/	/	/
6h	70	50	/	/	/	20	/	/	/	/
6i	0	30	/	/	/	95	95	85	70	/
6j	95	70	/	/	/	0	/	/	/	/
7a	0	50	/	/	/	40	/	/	/	/
7b	60	40	/	/	/	0	/	/	/	/
8a	50	0	/	/	/	100	100	100	100	/
8b	40	85	10	/	/	95	95	85	/	/
8c	0	100	55	/	/	100	30	/	/	/
8d	80	90	/	/	/	0	/	/	/	/
8e	0	80	/	/	/	0	/	/	/	/
8f	0	50	/	/	/	50	/	/	/	/
8g	50	98	/	/	/	0	/	/	/	/
8h	30	0	/	/	/	70	/	/	/	/
coumoxystrobin	70	80	80	65	/	0	/	/	/	/
kresoxim-methyl	/	/	/	/	/	100	100	100	100	/

note: “/” stands for no data

Many compounds showed 100% control against CDM at 400 mg L⁻¹, and these compounds were further tested at a lower dose range (200, 100, 50, 25, and 6.25 mg L⁻¹). The fungicidal activities of all synthesized compounds were lower than that of coumoxystrobin, except for 6h, which was comparable in potency (Table 1).

Some synthesized compounds also have fungicidal activity against RSB, CGM and WPM, besides CDM. Especially, compound 6f showed 100% control against CGM at 100 mg L⁻¹, much higher than that of coumoxystrobin (65%). Furthermore, compounds 6f, 6i and 8a all exhibited excellent activity against WPM, even at a lower dose range

(200, 100, and 50 mg L⁻¹). Particularly, 8a displayed equal fungicidal activity (100%) against WPM at 50 mg L⁻¹ to kresoxim-methyl. However, coumoxystrobin had no activity against WPM at 400 mg L⁻¹, which indicated that 8a is comparable with the commercial kresoxim-methyl (Table 2).

Based on data presented in Table 1, the title compounds (R₁ is methyl and R₂ is halogen or C₁-C₆ alkyl) seem to be more active against CDM at lower concentrations, particularly compound 6h (R₁ is methyl and R₂ is *n*-C₆H₁₃), which was 85% active at 6.25 mg L⁻¹, slightly lower than that of coumoxystrobin, and significantly higher than that of kresoxim-methyl. Replacement of the methyl group at R₁ with either *n*-C₃H₇ or Ph resulted in loss of activity, respectively (6i and 6j). There are, however, a few exceptions, such as 6e, 6f and 8d. When R₁ and R₂ are combined into a five- or six-membered ring, the fungicidal activity decreases considerably. In addition, introduction of methyl to R₃ of the coumarin ring has little effect on the overall fungicidal activity. Furthermore, related pharmacophores, such as OE (Q=Q₁), OA (Q=Q₂) and MC (Q=Q₃) were studied. As shown in the comparison of these three groups of compounds (6g versus 8d; 6a versus 7a; 6b versus 7b), it seemed that the fungicidal activity of OE (Q=Q₁) was at the same level as that of MC (Q=Q₂), significantly greater than that of OA (Q=Q₃), but an exception was noted, the pairs 6c and 8a showed the opposite trend.

Based on the results against WPM presented in Table 2, firstly, of the title compounds (R₁ is methyl or *n*-C₃H₇), all showed good fungicidal activity against WPM, but replacement of *n*-C₃H₇ with Ph led to loss of activity; while replacement of methyl with H had little effect on the

activity of the whole molecule. Secondly, introducing H, halogen, methyl, n -C₄H₉ or n -C₆H₁₃ to R₂ to replace n -C₃H₇ resulted in either reduction or loss of fungicidal activity. The cyclization of R₁ and R₂ made no contribution to the activity of the target compounds. Finally, as for Q, seemingly OE (Q=Q₁) has greater fungicidal activity than OA and MC (Q=Q₂, Q₃), but there is still an exception, the pairs **6a** and **7a** showed nearly equal efficacy.

Insecticidal activity: All synthesized compounds were tested against armyworm, diamondback moth, peach aphid, carmine spider mite and common mosquito. The bioassay results showed that the tested compounds lacked activity against diamondback moth and peach aphid, while three compounds exhibited some insecticidal or acaricidal activities, specifically **6h**, with 100% control against carmine spider mite; **7b** with 50% control against common mosquito, and **8h** with 40% control against armyworm. Overall, the insecticidal activity of this series of compounds was considered modest.

Conclusion: Two compounds with improved fungicidal activity compared with coumoxystrobin, **6f** and **8a**, were discovered as the optimal structure against CGM and WPM respectively, through introducing OE (Q=Q₁), OA (Q=Q₂) and MC (Q=Q₃) to replace MA (β -methoxyacrylate pharmacophore) in coumoxystrobin. Compound **6f** (R₁ methyl, R₂ methyl, R₃ H, and Q Q₃) showed good control against CGM and WPM, especially against CGM (100%) at 100 mg L⁻¹, much higher than that of coumoxystrobin (65%); Compound **8a** (R₁ methyl, R₂ n -C₃H₇, R₃ H, and Q Q₁) exhibited excellent activity against WPM at 50 mg L⁻¹ (100%), at the same level as that of the commercial kresoxim-methyl, and very significantly higher than that of coumoxystrobin (no control against WPM at 400 mg L⁻¹). Another compound, **6h** (R₁ methyl and R₂ n -C₆H₁₃), was also discovered, but although this compound exhibited good activity against CDM (85%) at 6.25 mg L⁻¹, it appeared to be slightly less potent than coumoxystrobin, but superior to kresoxim-methyl. Further syntheses and structure optimization studies are in progress.

Experimental

General: All starting materials and reagents were commercially available. Experimental conditions were quite similar to those reported earlier [9].

Synthesis of substituted coumarins (3): To a solution of substituted resorcinol (**1**) (1 mmol) in 10 mL of conc. H₂SO₄, ethyl 3-oxopropanoate (**2**) (1 mmol) was added drop wise at 5°C. After all the solution has been added (about 30 min), the reaction mixture was stirred at room temperature for 4–5 h and then was poured with vigorous stirring into a mixture of ice (50 g) and water (50 mL), and kept in a refrigerator overnight. The solid was filtered and thoroughly washed with cold water till the pH of the filtrate become neutral. The residue was dried over a

sintered glass funnel by vacuum to afford substituted coumarins **3**.

Synthesis of compound 6b: 7-Hydroxy-4-methylcoumarin (0.43 g, 2.45 mmol) was dissolved in 5 mL of butanone, and anhydrous potassium carbonate (0.68 g, 4.90 mmol) was added to the solution. The solution was stirred for 30 mins, and methyl (*E*)-methyl 2-[2-(bromomethyl)phenyl]-2-(methoxyimino)acetate (0.70 g, 2.45 mmol) was then added. The reaction mixture was stirred at refluxing temperature for 5 h and then was cooled, diluted with 50 mL water and extracted with ethyl acetate (3 × 100 mL). The combined extracts were washed with brine, dried (anhydrous magnesium sulfate), and filtered. The filtrate was evaporated and the crude product was purified via silica gel CC, using a 1:2 (v/v) mixture of ethyl acetate and light petroleum (boiling point range: 60–90°C) as the eluting solution to obtain compound **6b**: 0.83 g.

Synthesis of compound 7b: Compound **6b** (0.43 g, 2.45 mmol) was dissolved in 10 mL of methanol, and methylamine (0.16 g, 5.14 mmol) was added to the solution. After 3 h, the mixture was concentrated, diluted with 50 mL water and extracted with ethyl acetate (3 × 100 mL). The combined extracts were washed with brine, dried (anhydrous magnesium sulfate), and filtered. The filtrate was evaporated and the crude product was purified via silica gel CC, using a 1:1 (v/v) mixture of ethyl acetate and light petroleum to obtain compound **7b**: 0.90 g.

Synthesis of compound 8a: 7-Hydroxy-3,4-dimethylcoumarin (0.52 g, 2.73 mmol) was dissolved in 5 mL of DMF, and 60% sodium hydride (0.22 g, 5.46 mmol, washed with light petroleum) was added to the solution. The solution was stirred for 30 mins, and methyl 2-(bromomethyl)phenyl(methoxy)carbamate (0.75 g, 2.74 mmol) was then added and stirred for 3 h. The reaction mixture was added to 50 mL water and extracted with ethyl acetate (3 × 100 mL). The combined extracts were washed with brine, dried (anhydrous magnesium sulfate), and filtered. The filtrate was evaporated and the crude product was purified via silica gel CC, using a 1:2 (v/v) mixture of ethyl acetate and light petroleum to obtain compound **8a**: 0.82 g.

(*E*)-Methyl 2-(methoxyimino)-2-{2-[(4-methylcoumarin-7-yloxy)methyl]phenyl}acetate (6b**)**

Yield 85% of a white solid.

MP: 150–152°C.

IR (KBr) ν : 2980, 2960 (s, C-H), 1735, 1715 (s, C=O), 1610 (m, C=C), 1390 (m, C-H), 1290 (s, C-O), 1060, 1010 (s, =C-H), 870, 830, 750 (s, Ph-H) cm⁻¹.

¹H NMR (300 MHz, CDCl₃) δ : 7.39–7.54 (4H, m, *H*-3', *H*-5', *H*-6' and *H*-5), 7.21–7.24 (1H, m, *H*-4'), 6.85 (1H, dd, *J*=2.7, 9.0 Hz, *H*-6), 6.80 (1H, d, *J*=2.7 Hz, *H*-8), 6.13 (1H, s, *H*-3), 5.02 (2H, s, OCH₂), 4.05 (3H, s, NOCH₃), 3.87 (3H, s, COOCH₃), 2.39 (3H, s, CH₃-4).

Anal. Calcd. (%) for C₂₁H₁₉NO₆: C, 66.13; H, 5.02; N, 3.67. Found: C, 66.22; H, 4.98; N, 3.64.

(E)-2-(Methoxyimino)-N-methyl-2-{2-[(4-methylcoumarin-7-yloxy)methyl]phenyl}acetamide (7b)

Yield 90% of a white solid.

MP: 213-214°C.

IR (KBr) ν : 3200 (s, N-H), 2980, 2960 (s, C-H), 1720 (s, C=O), 1660 (s, C=C), 1620 (s, N=C), 1500 (s, Ph), 1420 (s, C-H), 1260, 1140 (s, C-O), 1000 (s, =C-H), 930, 750 (s, Ph-H) cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ : 7.41-7.50 (4H, m, *H*-5, *H*-3', *H*-5' and *H*-6'), 7.20-7.23 (1H, m, *H*-4'), 6.82-6.84 (2H, m, *H*-6 and *H*-8), 6.79 (1H, bs, NH), 6.13 (1H, s, *H*-3), 5.02 (2H, s, OCH_2), 3.97 (3H, s, NOCH_3), 2.92 (3H, d, $J=5.1$ Hz, NHCH_3), 2.38 (1H, s, CH_3 -4);

Anal. Calcd: C, 66.31; H, 5.30; N, 7.36. Found: C, 66.25; H, 5.36; N, 7.39.

Methyl 2-[(3,4-dimethylcoumarin-7-yloxy)methyl]phenyl(methoxy)carbamate (8a)

Yield 76% of a white solid.

MP: 121-122°C.

IR (KBr) ν : 2940 (s, C-H), 1730, 1700 (s, C=O), 1600 (m, C=C), 1490 (w, Ph), 1430 (m, C-H), 1375, 1310, 1250,1190 (s, C-O), 1170, 1080 (s, =C-H), 980, 805, 760 (s, Ph-H) cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ : 7.39-7.43 (1H, m, *H*-6'), 7.50 (1H, d, $J=8.7$ Hz, *H*-5), 7.38-7.40 (3H, m, *H*-3', *H*-4' and *H*-5'), 6.91 (1H, dd, $J=2.4, 8.7$ Hz, *H*-6), 6.85 (1H, d, $J=2.4$ Hz, *H*-8), 5.17 (2H, s, OCH_2), 3.83 (3H, s, COOCH_3), 3.77 (3H, s, NOCH_3), 2.36 (3H, s, CH_3 -4), 2.18 (3H, s, CH_3 -3);

Anal. Calcd: C, 65.03; H, 5.18; N, 3.79. Found: C, 65.23; H, 5.11; N, 3.74.

Supplementary data: ^1H NMR, IR, MS and elemental analyses for compounds **6a**, **6c-6j**, **7a** and **8b-8h**.**Acknowledgments** - The project was supported by the National Key Technology Support Program during the 12th Five-Year Plan Period (Grant No. 2011BAE06B00 and 2011BAE06B05) and the National Key Basic Research Program (973 Program) (Grant No. 2010CB735601 and 2010CB126105).**References**

- [1] (a) Singh R, Gupta BB, Malik OP. (1987) Studies on pesticides based on coumarin. I. Antifungal activity of 6-alkyl-3-*n*-butyl-7-hydroxy-4-methylcoumarins. *Pesticide Science*, **20**, 125-130; (b) Daoubi M, Duran-Patron R, Hmamouchi M, Hernandez-Galan R, Benharref A, Collado IG. (2004) Screening study for potential lead compounds for natural product-based fungicides: I. Synthesis and *in vitro* evaluation of coumarins against *Botrytis cinerea*. *Pest Management Science*, **60**, 927-932.
- [2] Beriger E. (1978) Insecticidally active 3-*N*-(4-trifluoromethylphenyl)-carbamoyl-4-hydroxycoumarin. US4078075.
- [3] Vyvyan JR. (2002) Allelochemicals as leads for new herbicides and agrochemicals. *Tetrahedron*, **58**, 1631-1646.
- [4] Liu CL, Li ZM. (2003) Discovery of strobil fungicides. *Chinese Journal of Pesticides (Nong Yao)*, **3**, 43-48.
- [5] (a) Liu CL, Li M, Guan AY, Zhang H, Li ZM. (2007) Design, synthesis and fungicidal activity of novel (*E*)-methyl 2-[(coumarin-7-yloxy)methyl]phenyl-3-methoxyacrylates. *Natural Product Communications*, **2**, 845-848; (b) Guan AY, Liu CL, Li M, Zhang H, Li ZN, Li ZM. (2011) Design, synthesis and structure-activity relationship of novel coumarin derivatives. *Pest Management Science*, **67**, 647-655; (c) Guan AY, Liu CL, Li ZN, Zhang MX, Si NG. (2011) The discovery of fungicide coumoxystrobin. *Chinese Journal of Pesticides (Nong Yao)*, **2**, 90-92; (d) Tomlin CDS. (2009) Kresoxim-methyl, In *The Pesticide Manual*. 15th Ed., British Crop Protection Council, Alton, UK. 688-690.
- [6] Tomlin CDS. (2009) Dimoxystrobin, In *The Pesticide Manual*. 15th Ed., British Crop Protection Council, Alton, UK. 383-384.
- [7] Tomlin CDS. (2009) Orsastrobin, In *The Pesticide Manual*. 15th Ed., British Crop Protection Council, Alton, UK. 840-841.
- [8] Tomlin CDS. (2009) Pyraclostrobin, In *The Pesticide Manual*. 15th Ed., British Crop Protection Council, Alton, UK. 971-972.
- [9] Li M, Liu CL, Yang JC, Li L, Li ZN, Zhang H, Li ZM. (2009) Design, synthesis and biological activity of new strobilurin derivatives with a 1*H*-Pyrazol-5-oxy side chain. *Natural Product Communications*, **4**, 1215-1220.
- [10] Liu CL. (2011) New approach for agrochemical discovery and application (I)—Intermediate derivatization method. *Chinese Journal of Pesticides (Nong Yao)*, **1**, 20-22.
- [11] Liu CL. (2009) New agrochemical candidates from intermediate derivatization method based on bioisosteric replacement. In *Frontiers of Modern Chemical Engineering, Metallurgy, and Material Technologies-7th Academic Conference of Chemical, Metallurgy and Material Engineering Department*, CAE, Wang JK. (Ed). Chemical Industry Press Publishers, Beijing. 86-94.
- [12] (a) Liu CL. (2008) Innovation of pesticide discovery methods and candidate compounds. *High-technology & Industrialization (Chinese)*, **9**, 79-81; (b) Chai BS, Liu CL, Li HC, He XM, Luo YM, Huang G, Zhang H, Chang JB. (2010) Design, synthesis and acaricidal activity of novel strobilurin derivatives containing pyrimidine moieties. *Pest Management Science*, **66**, 1028-1214.
- [13] Menn JJ. (1980) Contemporary frontiers in chemical pesticide research. *Journal of Agricultural and Food Chemistry*, **28**, 2-8.
- [14] (a) Iqbal PF, Bhat AR, Azam A. (2001) Antiamoebic coumarins from the root bark of *Adina cordifolia* and their new thiosemi-carbazone derivatives. *Journal of Medicinal Chemistry*, **44**, 664-671; (b) Khan KM, Saify ZS, Begum S, Noor F, Khan MZ, Hayat S, Choudhary MI, Perveen S, Rahman AU, Ullah Z. (2003) Synthesis and biological screening of 7-hydroxy-4-methyl-2*H*-chromen-2-one, 7-hydroxy-4,5-dimethyl-2*H*-chromen-2-one and some derivatives. *Natural Product Research*, **17**, 115-125.
- [15] Garazd YL, Panteleimonova TN, Garazd MM, Khilya VP. (2002) Modified coumarins. 7. Synthesis and biological activity of mannich bases of substituted 7,8,9,10-tetrahydrobenzo[*c*]chromen-6-ones. *Chemistry of Natural Compounds*, **38**, 532-538.
- [16] (a) Bender DR, Kanne D, Frazier JD, Rapoport H. (1983) Synthesis and derivatization of 8-acetylpsoralens. Acetyl migrations during Claisen rearrangement. *The Journal of Organic Chemistry*, **48**, 2709-2719; (b) Ross R, Fujimoto TT, Shaber SH. (1997) Benzoyloxy substituted aromatics and their use as fungicides and insecticides. EP0811614; (c) Jacobson RM, Nguyen LT. (2000) Phosphoryl hydrazine insecticides. US6147062; (d) Li M, Liu CL, Li L, Yang H, Li ZN, Zhang H, Li ZM. (2010) Design, synthesis and biological activities of new strobilurin derivatives containing substituted pyrazoles. *Pest Management Science*, **66**, 107-112.

Kinase Inhibitory, Haemolytic and Cytotoxic Activity of Three Deep-water Sponges from North Western Australia and their Fatty Acid Composition

Ana Zivanovic^a, Natalie J. Pastro^a, Jane Fromont^b, Murray Thomson^c and Danielle Skropeta^{a,d*}

^aSchool of Chemistry, University of Wollongong, NSW 2500, Australia

^bWestern Australian Museum, Locked Bag 49, Welshpool DC, WA 6986, Australia

^cSchool of Biological Sciences, University of Sydney, NSW 2006, Australia

^dCentre for Medicinal Chemistry, University of Wollongong, NSW 2500, Australia

skropeta@uow.edu.au

Received: August 10th, 2011; Accepted: October 10th, 2011

The c-AMP dependent protein kinase (PKA) inhibition, haemolytic activity, and cytotoxicity of 21 extracts obtained from North Western Australian sponges collected from depths of 84-135 m were investigated. Hexane extracts from *Ircinia/Sarcotragus* sp. and *Geodia* sp. displayed PKA inhibitory activities of 100 and 97% respectively (at 100 µg/mL), while aq. methanol extracts from *Haliclona* sp. exhibited potent haemolytic activity (75%) and hexane extracts from *Geodia* sp. were highly toxic (88%) to the brine shrimp *Artemia franciscana*. As the non-polar extracts gave the greatest PKA inhibition, these were further analysed by GC-MS and 29 fatty acids were identified in the highest proportions in *Ircinia/Sarcotragus* sp. > *Haliclona* sp. > *Geodia* sp. In contrast to shallow-water sponges that are dominated by polyunsaturated fatty acids with a high percentage of long chain fatty acids, LCFAs (C₂₄-C₃₀), the deep-sea sponges investigated herein were all found to be rich in saturated fatty acids, in particular C₁₄-C₂₀ fatty acids, including odd and branched chain fatty acids, with only low levels (0-10%) of LCFAs. Screening of the PKA inhibitory activity of a series of commercially available fatty acids identified C₁₄-C₁₈ fatty acids as possessing significant PKA inhibitory activity that may contribute to the activity observed in the sponges studied.

Keywords: deep-water sponges, fatty acids, enzyme inhibitors, protein kinase A, haemolysis, brine shrimp toxicity, cytotoxicity.

The deep sea is one of the most biodiverse and species-rich habitats on the planet, but also one of the least explored. In recent years, deep-sea organisms have emerged as an important new source of unexplored chemical, genetic and biological diversity [1a]. Yet of the 30,000 marine natural products reported to-date, less than 2% derive from deep-sea organisms [1b]. Moreover, metabolites from deep-sea organisms have exhibited a significantly higher 'hit rate' than their shallow-water counterparts in a wide number of bioassays [1c], including cytotoxicity towards both murine and human cancer cell lines [1b,2]. Of particular interest, are the metabolites from deep-sea sponges, which account for over 60% of the deep-sea natural products reported [1b].

c-AMP Dependent protein kinase (protein kinase A, PKA) is an important enzyme involved in the regulation of an increasing number of physiological processes including immune, cardiovascular and reproductive functions; steroid biosynthesis; adipocyte metabolism; and exocytotic processes [3a]. As such, the inhibition of PKA has become an attractive drug target in a number of areas, in particular in immune function [3b] and for memory disorders such as Alzheimer's and Parkinson's disease and schizophrenia [3c]. Interestingly, despite PKA being one of the most well characterized of all protein kinases [3d],

there are as yet no PKA inhibitors reported from any marine sources [4]. In addition to screening for specific enzyme inhibitors, the brine shrimp lethality assay is a simple and inexpensive way to screen natural product extracts for pharmacological activity [5a]. A strong correlation has been found between brine shrimp toxicity and cytotoxicity in human cancer cell lines [5b], and cytotoxicity is the most commonly reported biological activity in marine fauna, in particular marine sponges [5c]. Haemolytic activity is also prevalent in Porifera, and is another general indicator of bioactivity, in particular membrane-directed cytotoxicity, which is desirable in the development of bactericidal molecules and for targeted intracellular drug delivery.

Herein, we describe the biological screening of 21 extracts obtained from three NW Australian deep-sea sponge samples. The sponge extracts described herein were tested for their ability to inhibit PKA; to lyse equine erythrocytes; and for their toxicity towards the brine shrimp *Artemia franciscana*. Furthermore, the fatty acid composition of the bioactive, non-polar extracts of each sample was identified by GC-MS. Using remotely operated vehicles, a range of deep-sea organisms including sponges, cnidarians, echinoderms, crustaceans and fish were

collected in the North West Shelf of Australia from depths of 84-135 m, as part of wider study on deep-sea ecosystems [6a,6b]. From this diverse collection, the three largest sponge samples were chosen for investigation and identified as *Haliclona* sp. (Order Haplosclerida, Family Chalinidae), *Geodia* sp. (Order Astrophorida, Family Geodiidae) and an intertwined sample of *Sarcotragus* sp. and *Ircinia* sp. (both belonging to the Order Dictyoceratida, Family Irciniidae). Crude extracts were prepared from each sample with dichloromethane and methanol and further partitioned using a modified Kupchan method to give seven partially purified extracts in water, butanol, 50% aq. methanol, 10% aq. methanol, dichloromethane and two hexane extracts, one from the original crude methanol extract (referred to as hexane 1) and the other from the original dichloromethane extract (referred to as hexane 2). This gave a total of 21 extracts that were tested at a final concentration of 100 µg/mL.

PKA inhibitory activity of 97-100% was observed for the hexane 1 extracts of all sponges investigated except *Haliclona*. Other non-polar extracts, including the hexane 2 from *Geodia* sp. and the dichloromethane extract from *Ircinia/Sarcotragus* sp. displayed significant activity (41-84%, Fig. 1 A-C). This is consistent with the large number of non-polar kinase inhibitors described from marine sponges, which include sesterterpenoid mitogen activated protein kinase MSK1 and MAPKAPK-2 inhibitors from *Ircinia* sp. [7a] and a series of prenylhydroquinones isolated from *Sarcotragus muscarum* and *Ircinia fasciculata* that inhibited multiple protein kinases [7b]. Although marine-derived PKA inhibitors are yet to be reported, PKA and related enzymes have been identified from marine sources (mollusks, microalgae), where they play similar roles to their mammalian counterparts [7c].

Haemolytic activity was found in over half of the extracts, with the greatest activity (75%) observed for the 50% aq. methanol extract of the *Haliclona* sp. Extracts showing the highest PKA inhibitory activity (Figure 1, hexane 1 panels B-C) displayed low haemolysis, while almost all *Haliclona* extracts exhibited significant haemolytic activity (Figure 1, panel D), consistent with reports of such activity for this genus [8]. Of the sponges under investigation, haemolytic activity has previously been reported for both *Haliclona* sp. and *Geodia corticostylifera* [9a]. The hexane 1 extract from *Geodia* sp. exhibited 88% toxicity towards the brine shrimp *A. franciscana*; the first report of such activity for this genus (see Supplementary data). Although not observed here, brine shrimp toxicity has been reported previously for the other two sponge genera studied here, including a family of cerebrosidyls from *Haliclona* sp. [9b] and three sulphated polyprenylhydroquinones from *Ircinia spinosula* [9c].

A diverse range of bioactive metabolites have been reported from shallow-water species from the four sponge genera under investigation. There is, however, a paucity of reported metabolites from related deep-water species, with only a small handful to-date including: triterpenoids [10a]




Figure 1: % PKA inhibition and haemolysis activity of deep-sea sponge extracts. Measurements were performed in triplicate at 100 µg/mL and data are presented as means ± S.E.M. DCM, dichloromethane.

and sesterterpene sulfates from the deep-water sponges *Sarcotragus spinulosus* and *Ircinia* sp. [10b]; and the nematocidal polyketide geodin [10c] and antibacterial alkaloid baretin [10d] from two deep-water *Geodia* species. The results described herein are the first report on biological activity in deep-water variants of *Haliclona*.

GC-MS analysis on the lipid component of each sponge was performed (see Table 1), and a total of 23 fatty acids identified for the *Haliclona* sp. as 23-67% of each extract. The major fatty acids were the PUFA C20:4 *n*-5 (21%) and 9-octadecenoic acid (7%) in the hexane 1 extract; hexadecanoic acid (8%) and 12-methyltetradecanoic acid (6%) in the hexane 2 extract; and 3-methylbutanoic acid (14%) and eicosanoic acid (7%) in the dichloromethane extract.

A total of 17 fatty acids were identified for the *Geodia* sp. as 4-24% of each extract: significantly lower than in the other sponges. The C18 fatty acids, 9-octadecenoic acid (5%) and octadecanoic acid (5%) in the dichloromethane extract were the most abundant fatty acids in this sponge, with all other fatty acids present as <5% of each extract. In contrast, the *Ircinia/Sarcotragus* sp. were particularly rich in fatty acids, with a total of 22 identified as 42-83% of each extract. The major fatty acids, accounting for 10% of each extract, were 7-methylhexadecanoic acid/PUFA C20:4 *n*-5, 11-octadecenoic acid, and octadecanoic acid in the hexane 1, hexane 2 and dichloromethane extracts.

A diverse range of fatty acids were identified for the deep-sea sponges including long, branched and odd chain fatty acids common to the phylum Porifera [11a]. The fatty acid profile was dominated by saturated C₁₄-C₂₀ fatty acids, but also varied across species either in type or in relative percentages, including extracts exhibiting similar activities. There were no apparent correlations amongst the bioassay results and/or their lipid profiles. For example, although the bioactivity profiles of the *Geodia* and *Ircinia*/

Table 1: Fatty acid composition of the non-polar extracts from the deep water sponges *Haliclona* sp., *Geodia* sp. and *Ircinia/Sarcotragus* sp.

Fatty acids	Type	<i>Haliclona</i> sp.			<i>Geodia</i> sp.			<i>Ircinia/Sarcotragus</i> sp.		
		A	B	C	A	B	C	A	B	C
1	2-Methylbutanoic acid	5:0	-	2.38	-	-	-	-	-	-
2	3-Methylbutanoic acid	5:0	-	13.52	-	-	-	-	-	-
3	Pentanoic acid	5:0	-	5.29	-	-	-	-	-	-
4	Hexanoic acid	6:0	-	0.32	1.54	-	1.24	-	-	-
5	Decanoic acid	10:0	2.18	4.75	-	0.65	-	2.46	3.66	-
6	10-Methylundecanoic acid	12:0	-	-	-	-	-	-	-	0.21
7	Dodecanoic acid	12:0	-	-	3.65	-	-	2.58	5.85	-
8	Tridecanoic acid	13:0	-	-	-	0.39	-	-	-	-
9	Tetradecanoic acid	14:0	-	1.58	2.03	0.49	2.03	5.72	6.39	-
10	12-Methyltetradecanoic acid	15:0	6.42	3.77	2.51	0.63	1.40	5.78	8.27	6.23
11	5,9,13-Trimethyltetradecanoic acid	17:0	-	-	-	-	-	-	4.37	-
12	Pentadecanoic acid	15:0	0.5	1.53	1.57	0.60	-	-	-	6.36
13	Hexadecanoic acid	16:0	7.74	2.21	-	0.25	-	0.32	3.66	4.39
14	7-Methylhexadecanoic acid	17:0	-	1.31	-	-	2.71	-	9.65	-
15	9-Hexadecenoic acid	16:1 <i>n</i> -9	0.84	5.89	-	-	1.77	0.44	-	-
16	Heptadecanoic acid	17:0	-	-	-	0.35	-	-	7.31	-
17	10-Methylheptadecanoic acid	17:0	-	-	-	-	-	-	-	8.51
18	Octadecanoic acid	18:0	-	3.85	-	-	4.96	-	-	9.75
19	17-Methyloctadecanoic acid	19:0	-	-	-	-	-	-	-	3.41
20	9-Octadecenoic acid	18:1 <i>n</i> -9	0.41	6.47	-	0.41	5.21	-	-	7.84
21	11-Octadecenoic acid	18:1 <i>n</i> -7	-	-	1.90	-	-	9.53	-	7.98
22	9,12,15-Octadecatrienoic acid	18:3 <i>n</i> -9	-	-	-	-	-	-	-	8.16
23	Nonadecanoic acid	19:0	2.98	1.83	-	-	-	-	-	-
24	Eicosanoic acid	20:0	1.98	2.18	-	-	0.52	3.69	6.75	8.22
25	11,14-Eicosadienoic acid	20:2 <i>n</i> -6	-	3.47	-	-	3.16	-	-	-
26	5,8,11,14-Eicosatetraenoic acid	20:4 <i>n</i> -6	-	21.15	-	-	-	3.23	9.62	-
27	Docosanoic acid	22:0	-	-	-	-	0.78	2.55	-	3.16
28	Tetracosanoic acid	24:0	0.15	-	-	-	-	2.54	-	4.65
29	Hexacosanoic acid	26:0	-	1.75	-	-	-	2.87	-	3.96
% of total extract		23.20	61.74	67.23	13.20	3.77	23.78	41.71	65.53	82.83

A: dichloromethane-derived hexane extract (hexane 2); B: methanol-derived hexane extract (hexane 1); C: dichloromethane extract.

Sarcotragus species were similar, their fatty acid profiles were entirely different. For the latter, even extracts displaying similar PKA inhibition activities comprised entirely different fatty acids (*cf.* the *Ircinia* hexane 1 and dichloromethane extracts in Figure 1, panel C and Table 1, columns B and C). Likewise, two of the samples obtained from the identical location and depth (*Geodia* and *Haliclona*) displayed both different bioactivities and fatty acid profiles, indicating that these attributes are largely species dependent. Of the sponge genera investigated, only the fatty acid composition for *Ircinia* sp. has been reported in detail previously, which showed a similar predominance of straight and branched C₁₄-C₁₈ fatty acids to that obtained. Our *Ircinia/Sarcotragus* sp. from 84 m depth contained a greater proportion of saturated fatty acids than reported *Ircinia* specimens from ~10 m, and also lacked the C₂₄-C₂₅Δ^{5,9} demospongiac acids characteristic of shallow-water *Ircinia* sp. [11b]. However, whether these differences are depth related will require further study.

To test which of the fatty acids may contribute to the observed bioactivity, a series of commercially available fatty acids (C14:0, C16:0, C18:0, C18:1, C14:22, C18:22) were examined for their ability to inhibit PKA. Several of these exhibited high levels of PKA inhibition at 100 µg/mL in the order C14:0 (64%), C16:0 (34%), C18:1 (33%) and C18:0 (26%) (see Supplementary data). As these are the dominant fatty acids in the sponges investigated, they may contribute to the observed PKA inhibitory activities, along with other non-polar metabolites, the identification of which is currently underway. In conclusion, these results indicate that deep-sea sponges of NW Australia are a rich source of PKA inhibitory, haemolytic and cytotoxic substances.

Experimental

Collection: The deep-sea sponges, were collected in the North West Shelf of Australia in June-August of 2008 and identified as: *Haliclona* sp., *Geodia* sp., and a mixed sample of intertwined *Sarcotragus* sp. and *Ircinia* sp., the former two from the same location (depth 135 m), and the latter from a location 80 km away (depth 84 m). Samples were collected under permit AU-COM2008032 issued by the Australian Government and vouchers (WAMZ45792-WAMZ45795) are at the Western Australian Museum.

Extracts: Sponge extracts were prepared from methanol and dichloromethane, followed by fractionation using a modified Kupchan partitioning method [12] to give seven extracts: water, butanol, 50% aq. methanol, 10% aq. methanol, dichloromethane, hexane (1) and hexane (2), where hexane (1) was derived from the crude methanol extract and hexane (2) from the crude dichloromethane extract (see Supplementary data). All assays were conducted at 100 µg/mL (in 2% DMSO or 2% ethanol).

Protein kinase A inhibition: This was determined using the Kinase-Glo[®] Luminescent Kinase Assay (Promega Corporation, Madison, USA). Each well of an opaque white 96 well plate (Corning) contained 5 µL of sample, 25 µL of ATP (20 µM), and 20 µL of a mixture of kinase (0.1 unit/µL) and Kemptide substrate (140 µM) in reaction buffer (40 mM Tris, 20mM MgCl₂, BSA 0.1mg/mL, pH 7.4). After incubation (1 h, rt), 50 µL of the Kinase-Glo[®] Reagent was added, the mixture incubated for 15 min and luminescence measured using a BMG Labtech FLUOstar Optima[®] luminometer. Data were determined relative to positive controls containing only solvent, while negative controls contained no substrate and staurosporine (IC₅₀ = 7nM against PKA) was used as an internal standard.

Haemolytic activity: Reaction mixtures containing 100 μL of sample and 900 μL of a 0.5% haemocyte suspension (from defibrinated horse blood, Oxoid, Australia) in PBS were incubated for 30 min at 37°C, centrifuged (5 min, 3000 rpm) and 100 μL of supernatant from each sample transferred to a 96 well plate and absorbance measured at λ 550 nm on a Spectramax PLUS. Data were determined relative to positive controls containing distilled water, and negative controls contained PBS in place of the sample.

Cytotoxicity: Brine shrimp eggs (*Artemia franciscana*) were hatched at 26–28°C for 48 h in brine using a light source and constant aeration. Ten animals were transferred to individual wells of a 96 well plate, along with 20 μL of sample and 180 μL of salt solution. Plates were incubated for 48 h under a light source, then surviving animals counted and percent mortality calculated. Control tests accounted for natural death [5a].

GC/MS Analysis: Performed on a Shimadzu QP-5050A GC-MS system equipped with a BP-5 fused silica Rxi-5ms capillary column (5% phenyl/95% dimethyl polysiloxane, 30 m x 0.25 mm, 0.25 μm film thickness, Resrek), using a helium carrier gas (1.0 mL/min) and samples of 1 μL over the temperature range 80–300°C. Fatty acids were detected as their methyl esters after treatment with methanolic BF_3 and their structural identification was based on comparisons to the NIST 08 mass spectral database.

Supplementary data: Further experimental details are available on-line.

Acknowledgments – We thank D Cummings and A Fowler for sample collections; J Korth and V Valenzuela for technical assistance; and the Australian Research Council (Linkage grant, LP0775183) and Woodside Energy Ltd for financial and logistical support.

References

- [1] (a) Gage JD. (1996) Why are there so many species in deep-sea sediment? *Journal of Experimental Marine Biology and Ecology* **200**, 257-286; (b) Skropeta D. (2008) Deep-sea natural products. *Natural Product Reports*, **25**, 1131-1166; (c) Schupp PJ, Kohlert-Schupp C, Whitefield S, Engemann A, Rohde S, Hemscheidt T, Pezzuto JM, Kondratyuk TP, Park EJ, Marler L, Rostama B, Wright AD. (2009) Cancer chemopreventive and anticancer evaluation of extracts and fractions from marine macro- and micro-organisms collected from twilight zone waters around Guam. *Natural Product Communications*, **4**, 1717-1728.
- [2] Dumdei EJ, Blunt JW, Munro MHG, Pannell LK. (1997) Isolation of calyculins, calyculinamides, and swinholide H from the New Zealand deep-water marine sponge *Lamellomorpha strongylata*. *Journal of Organic Chemistry*, **62**, 2636-2639.
- [3] (a) Tasken K, Aandahl EM. (2004) Localized effects of cAMP mediated by distinct routes of protein kinase A. *Physiological Reviews*, **84**, 137-167; (b) Torgersen KM, Vang T, Abrahamsen H, Yaqub S, Tasken K. (2002) Molecular mechanisms for protein kinase A-mediated modulation of immune function. *Cellular Signalling*, **14**, 1-9; (c) Arnsten AFT, Ramos BP, Birnbaum SG, Taylor JR. (2005) Protein kinase A as a therapeutic target for memory disorders: rationale and challenges. *Trends in Molecular Medicine*, **11**, 121-128; (d) Taylor SS, Yang J, Wu J, Haste NM, Radzio-Andzelm E, Anand G. (2004) PKA: a portrait of protein kinase dynamics. *Biochimica Et Biophysica Acta-Proteins and Proteomics*, **1697**, 259-269.
- [4] Skropeta D, Pastro NJ, Zivanovic A. (2011) Kinase inhibitors from marine sponges. *Marine Drugs*, **9**, 2131-2154.
- [5] (a) Solis PN, Wright CW, Anderson MM, Gupta MP, Phillipson JD. (1993) A microwell cytotoxicity assay using *Artemia salina* (brine shrimp). *Planta Medica*, **59**, 250-252; (b) Carballo J, Hernández-Inda Z, Pérez P, García-Grávalos M. (2002) A comparison between two brine shrimp assays to detect in vitro cytotoxicity in marine natural products. *BMC Biotechnology*, **2**, 17-21; (c) Blunt JW, Copp BR, Munro MHG, Northcote PT, Prinsep MR. (2006) Marine natural products. *Natural Product Reports*, **23**, 26-78.
- [6] (a) Cummings DO, Booth DJ, Lee RW, Simpson SJ, Pile AJ. (2010) Ontogenetic diet shifts in the reef fish *Pseudanthias rubrizonatus* from isolated deep water populations. *Marine Ecology Progress Series*, submitted; (b) Cummings DO, Lee RW, Simpson SJ, Booth DJ, Pile AJ, Holmes SP. Resource partitioning amongst co-occurring decapods on wellheads from Australia's North West Shelf. An analysis of carbon and nitrogen stable isotopes. *Journal of Experimental Marine Biology and Ecology*. submitted
- [7] (a) Buchanan MS, Edser A, King G, Whitmore J, Quinn RJ. (2001) Cheilanthane sesterterpenes, protein kinase inhibitors, from a marine sponge of the genus *Ircinia*. *Journal of Natural Products*, **64**, 300-303; (b) Watjen W, Putz A, Chovolou Y, et al. (2009) Hexa-, hepta- and nonaprenylhydroquinones isolated from marine sponges *Sarcotragus muscarum* and *Ircinia fasciculata* inhibit NF-kappa B signalling in H4IIE cells. *Journal of Pharmacy and Pharmacology*, **61**, 919-924; (c) Dawson JF, Wang KH, Holmes CFB. (1996) Identification and characterization of cAMP-dependent protein kinase and its possible direct interactions with protein phosphatase-1 in marine dinoflagellates. *Biochemistry and Cell Biology-Biochimie Et Biologie Cellulaire*, **74**, 559-567.
- [8] Mebs D, Weiler I, Heinke HF. (1985) Bioactive proteins from marine sponges - screening of sponge extracts for hemagglutinating, hemolytic, ichthyotoxic and lethal properties and isolation and characterization of hemagglutinins. *Toxicon*, **23**, 955-962.
- [9] (a) Rangel M, Konno K, Brunaldi K, Procopio J, De Freitas JC. (2005) Neurotoxic activity induced by a haemolytic substance in the extract of the marine sponge *Geodia corticostylifera*. *Comparative Biochemistry and Physiology C-Toxicology & Pharmacology*, **141**, 207-215; (b) Mansoor TA, Shinde PB, Luo X, Hong J, Lee CO, Sim CJ, Son BW, Jung JH. (2007) Renierosides, Cerebrosides from a marine sponge *Haliclona* (Reniera) sp. *Journal of Natural Products*, **70**, 1481-1486; (c) DeRosa S, Crispino A, Degiulio A, Iodice C, Milone A. (1995) Sulfated polyprenylhydroquinones from the sponge *Ircinia spinosula*. *Journal of Natural Products-Lloydia*, **58**, 1450-1454.
- [10] (a) Ponomarenko LP, Makar'eva TN, Stonik VA. (1998) A new nortriterpenoid from the deep-sea sponge *Sarcotragus spinulosus*. *Russian Chemical Bulletin*, **47**, 2017-2019; (b) Wright AE, McCarthy PJ, Schulte GK. (1989) Sulfircin: a new sesterterpene sulfate from a deep-water sponge of the genus *Ircinia*. *Journal of Organic Chemistry*, **54**, 3472-3474; (c) Capon RJ, Skene C, Lacey E, Gill JH, Wadsworth D, Friedel T. (1999) Geodin A magnesium salt: A novel nematocide from a southern Australian marine sponge *Geodia*. *Journal of Natural Products*, **62**, 1256-1259; (d) Lidgren G, Bohlin L. (1986) Studies of swedish marine organisms VII. A novel biologically active indole alkaloid from the sponge *Geodia baretii*. *Tetrahedron Letters*, **27**, 3283-3284.
- [11] (a) Bergquist PR, Lawson MP, Lavis A, Cambie RC. (1984) Fatty-acid composition and the classification of the Porifera. *Biochemical Systematics and Ecology*, **12**, 63-84; (b) Carballeira NM, Shalabi F, Cruz C, Rodriguez J, Rodriguez E. (1991) Comparative study of the fatty acid composition of sponges of the genus *Ircinia* - identification of the new 23-methyl-5,9-tetracosadienoic acid. *Comparative Biochemistry and Physiology B-Biochemistry & Molecular Biology*, **100**, 489-492.
- [12] Thale Z, Johnson T, Tenney K, Wenzel PJ, Lobkovsky E, Clardy J, Media J, Pietraszkiewicz H, Valeriote FA, Crews P. (2002) Structures and cytotoxic properties of sponge-derived bisannulated acridines. *Journal of Organic Chemistry*, **67**, 9384-9391.

Antimicrobial and Cytotoxic Effects of Mexican Medicinal Plants

Maria del Rosario Jacobo-Salcedo^a, Angel Josabad Alonso-Castro^{bc*}, Luis A. Salazar-Olivo^d, Candy Carranza-Alvarez^e, Luis Ángel González-Espíndola^a, Fabiola Domínguez^f, Sandra Patricia Maciel-Torres^a, Concepción García-Lujan^a, Marisela del Rocio González-Martínez^g, Maricela Gómez-Sánchez^h, Eduardo Estrada-Castillónⁱ, Rocio Zapata-Bustos^d, Pedro Medellín-Milán^e and Alejandro García-Carrancá^{cj*}

^aFacultad de Ciencias Químicas, Universidad Juárez del Estado de Durango, Avenida Artículo 123, Fraccionamiento Filadelfia, Gómez Palacio, Durango, 35010, México

^bFacultad de Química, Universidad Nacional Autónoma de México, 04510 México, D.F., México

^cInstituto Nacional de Cancerología, Av. San Fernando No. 22, Col. Sección XVI, Tlalpan, 14080 México, D.F., México

^dInstituto Potosino de Investigación Científica y Tecnológica, Camino San José 2055, San Luis Potosí, SLP, 78216, México

^eAgenda Ambiental de la Universidad Autónoma de San Luis Potosí, Avenida Manuel Nava 201 tercer piso, Zona Universitaria, San Luis Potosí, San Luis Potosí, 78210, México

^fCentro de Investigación Biomédica de Oriente, IMSS, Carretera Federal Atlixco-Metepec, 74360 Metepec, Puebla, México

^gHospital de Especialidades No. 71, IMSS, Boulevard Revolución 2650 Ote, 27200, Torreón, Coahuila, México

^hFacultad de Ciencias Naturales, Universidad Autónoma de Querétaro, Av de las Ciencias, 76230 Querétaro, México

ⁱFacultad de Ciencias Forestales, Universidad Autónoma de Nuevo León, Carretera Km 145, 67700 Nuevo León, México

^jInstituto de Investigaciones Biomédicas, Universidad Nacional Autónoma de México, 04510 México, D.F., México

angeljosabad@hotmail.com; carranca@biomedicas.unam.mx

Received: May 28th, 2011; Accepted: October 3rd, 2011

The antimicrobial effects of the Mexican medicinal plants *Guazuma ulmifolia*, *Justicia spicigera*, *Opuntia joconostle*, *O. leucotricha*, *Parkinsonia aculeata*, *Phoradendron longifolium*, *P. serotinum*, *Psittacanthus calyculatus*, *Tecoma stans* and *Teucrium cubense* were tested against several human multi-drug resistant pathogens, including three Gram (+) and five Gram (-) bacterial species and three fungal species using the disk-diffusion assay. The cytotoxicity of plant extracts on human cancer cell lines and human normal non-cancerous cells was also evaluated using the MTT assay. *Phoradendron longifolium*, *Teucrium cubense*, *Opuntia joconostle*, *Tecoma stans* and *Guazuma ulmifolia* showed potent antimicrobial effects against at least one multidrug-resistant microorganism (inhibition zone > 15 mm). Only *Justicia spicigera* and *Phoradendron serotinum* extracts exerted active cytotoxic effects on human breast cancer cells (IC₅₀ ≤ 30 µg/mL). The results showed that *Guazuma ulmifolia* produced potent antimicrobial effects against *Candida albicans* and *Acinetobacter baumannii*, whereas *Justicia spicigera* and *Phoradendron serotinum* exerted the highest toxic effects on MCF-7 and HeLa, respectively, which are human cancer cell lines. These three plant species may be important sources of antimicrobial and cytotoxic agents.

Keywords: Mexican medicinal plants, antimicrobial, multidrug-resistance, cancer, cytotoxic, peripheral blood mononuclear cells.

Plant extracts have been an important source of anticancer [1] and antimicrobial [2] agents currently used in clinics, and Mexican medicinal plants play an important role in the search for new treatments. *Guazuma ulmifolia* (Sterculiaceae), *Justicia spicigera* (Acanthaceae), *Opuntia joconostle* (Cactaceae), *O. leucotricha* (Cactaceae),

Parkinsonia aculeata L. (Fabaceae), *Phoradendron longifolium*, *P. serotinum* (Viscaceae), *Psittacanthus calyculatus* (Loranthaceae), *Tecoma stans* (Bignoniaceae) and *Teucrium cubense* (Lamiaceae) are plants traditionally used to treat infections (Table 1). The antimicrobial effects of *Guazuma ulmifolia*, *Justicia spicigera*, *Tecoma stans*

Table 1: Collecting data and ethnomedical information on Mexican plants for antimicrobial and cytotoxic studies.

Plant name (Family)	Native	Collection information	Location collection	Part Used	Ethnomedical uses	References
<i>Guazuma ulmifolia</i> Lam. (Sterculiaceae)	From Mexico to South America	M. Gómez 705, QMEX	Corregidora, Qro., México	Bark	Diarrhea, stomachache, fever, urinary infections,	[4]
<i>Justicia spicigera</i> Schltdl. (Acanthaceae)	From Mexico to South America	García-Pérez 46450, SLPM	Ciudad Valles, Slp, México	Leaves	Fever, diarrhea, dysentery,	[7]
<i>Opuntia joconostle</i> F.A.C Weber (Cactaceae)	Mexico	M. Gómez 705, QMEX	San Miguel de Allende, Gto., México	Fruit	Anti-inflammatory, wounds, fever	[8]
<i>Opuntia leucotricha</i> DC. (Cactaceae)	Mexico	M. Gómez 706, QMEX	Cerritos, Slp., México	Fruit	Anti-inflammatory, wounds, fever	[8]
<i>Parkinsonia aculeata</i> L. (Fabaceae)	From south United States to South America and Caribbean	M. Gómez 705, QMEX	Querétaro, Qro., México	Flowers	Urinary tract infections, fever, stomachache	[9]
<i>Phoradendron longifolium</i> Eichler ex. Trel. (Viscaceae)	Mexico	M. Gómez 705, QMEX	Cerro de San Pedro, SLP., México	Leaves	Anti-inflammatory, urinary tract infections	[8]
<i>Phoradendron serotinum</i> (Raf.) M. C. Johnst. (Viscaceae)	From eastern United States to Mexico	E Estrada 20768, CFNL	Santa Catarina, NL, Mexico	Leaves	Body pain, urinary tract infections, dysmenorrhea	[10]
<i>Psittacanthus calyculatus</i> (DC.) G. Don (Loranthaceae)	Mexico	M. Gómez 754, QMEX	Querétaro, Qro., México	Leaves	Wounds, fever	[9]
<i>Tecoma stans</i> (L.) Juss. ex Kunth (Bignoniaceae)	From southeastern United States to South America and Caribbean	M. Gómez 700, QMEX	Querétaro, Qro., México	Leaves	Gastritis, diarrhea, stomachache	[5]
<i>Teucrium cubense</i> Jacq. (Lamiaceae)	From south United States to South America	M. Gómez 701, QMEX	Querétaro, Qro., México	Leaves and branches	Skin infections, anti-inflammatory,	[9]

QMEX: Herbarium of Facultad de Ciencias Naturales, Universidad Autónoma de Querétaro
 CFNL: Herbarium of Facultad de Ciencias Forestales, Universidad Autónoma de Nuevo León
 SLPM: Herbarium Isidro Palacios of Instituto de Investigación de Zonas Desérticas, Universidad Autónoma de San Luis Potosí

Table 2: Antimicrobial activities of Mexican plant extracts against human pathogens.

	Microorganism/ Concentration (µg/disk)	<i>Enterococcus faecalis</i> 7140203#	<i>Staphylococcus aureus</i> 3090014	<i>Staphylococcus haemolyticus</i> 309220	<i>Acinetobacter baumannii</i> 130412	<i>Acinetobacter lwoffii</i> 130415	<i>Burkholderia cepacia</i> 160255	<i>Escherichia coli</i> 412352	<i>Pseudomonas aeruginosa</i> 160239	<i>Candida albicans</i> 3100268	<i>Candida tropicalis</i> 120120	<i>Trichosporon belgii</i> 80145
GU	75	NA	NA	NA	NA	NA	NA	NA	NA	26 ± 2	NA	NA
	375	NA	NA	NA	NA	19 ± 1	NA	NA	NA	26 ± 3	NA	NA
	750	16 ± 2 *	NA	NA	NA	NA	NA	NA	10 ± 3	NA	NA	NA
JS	75	NA	NA	NA	NA	NA	NA	NA	NA	NA	10 ± 1	NA
	375	NA	NA	NA	NA	11 ± 2	NA	NA	12 ± 3	NA	NA	11 ± 2
	750	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
OJ	75	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
	375	12 ± 3	NA	NA	NA	18 ± 3	NA	NA	NA	NA	NA	NA
	750	11 ± 2	NA	NA	NA	NA	NA	13 ± 2	NA	NA	NA	NA
OL	75	12 ± 2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
	375	12 ± 3	NA	NA	NA	NA	NA	NA	10 ± 2	NA	NA	NA
	750	NA	NA	NA	NA	16 ± 1	NA	NA	10 ± 1	NA	NA	NA
PA	75	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
	375	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
	750	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
PL	75	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
	375	NA	NA	NA	NA	22 ± 2	NA	NA	NA	NA	NA	NA
	750	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
PS	75	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
	375	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
	750	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
PC	75	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
	375	NA	NA	NA	NA	NA	NA	NA	8 ± 2	NA	NA	NA
	750	NA	NA	NA	NA	16 ± 3	16 ± 1	NA	8 ± 2	NA	NA	NA
TS	75	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
	375	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
	750	NA	NA	NA	NA	NA	14 ± 2	NA	NA	NA	10 ± 2	NA
TC	75	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
	375	NA	NA	NA	NA	22 ± 3	NA	NA	NA	NA	NA	NA
	750	NA	NA	NA	NA	NA	NA	NA	10 ± 2	12 ± 3	NA	NA
NEO	25	20 ± 3	19 ± 2	20 ± 2	15 ± 2	21 ± 2	17 ± 3	20 ± 3	15 ± 2	23 ± 3	19 ± 2	20 ± 2

*Diameter of clear zone includes the diameter of the 6 mm disk. # Reference number. Values are shown as the mean ± standard deviations of at least three experiments in triplicate. NA=No inhibitory activity. GU, *Guazuma ulmifolia*; JS, *Justicia spicigera*; OJ, *Opuntia joconostle*; OL, *Opuntia leucotricha*; PA, *Parkinsonia aculeata*; PL, *Phoradendron longifolium*; PS, *Phoradendron serotinum*; PC, *Psittacanthus calyculatus*; TS, *Tecoma stans*; TC, *Teucrium cubense*

and *Parkinsonia aculeata* have been recorded against drug-sensitive bacteria and fungi [3-5].

In traditional medicine the term “cancer” is an ill-defined concept, and plant extracts with active cytotoxic effects are employed for the empirical treatment of dermatological diseases and as anti-inflammatory agents [6]. On this basis, and since the cytotoxicity of *Guazuma ulmifolia*, *Justicia spicigera*, *Opuntia joconostle*, *O. leucotricha*, *Parkinsonia aculeata*, *Phoradendron longifolium*, *P. serotinum*, *Psittacanthus calyculatus*, *Tecoma stans* and *Teucrium*

cubense for human cancer cell lines remain unexplored, we also tested the cytotoxic effects of these plant extracts on human cancer cells used empirically for the treatment of dermatological and inflammatory illnesses (Table 1).

This work demonstrates that *Guazuma ulmifolia*, *Justicia spicigera*, *Opuntia joconostle*, *O. leucotricha*, *Parkinsonia aculeata*, *Phoradendron longifolium*, *P. serotinum*, *Psittacanthus calyculatus*, *Tecoma stans* and *Teucrium cubense* exert antimicrobial and cytotoxic effects and may be important sources of cytotoxic and antimicrobial agents.

Table 3: Cytotoxic activity of Mexican medicinal plants on human cancer cell lines with $IC_{50} < 200 \mu\text{g/mL}$.

	Cancer cell lines ($IC_{50} \mu\text{g/mL}$)		DU-145	SKOV-3
	MCF-7	HeLa		
<i>Justicia spicigera</i>	48 ± 5.1	17 ± 2.6	49 ± 3.4	43 ± 4.5
<i>Parkinsonia aculeata</i>	189 ± 25	>200	>200	137 ± 29
<i>Phoradendron serotinum</i>	28 ± 4.1	46 ± 3.2	87 ± 7.7	37 ± 5.2
Cisplatin	7.1 ± 1.1	3.21 ± 0.9	13.68 ± 1.3	10.7 ± 1.3

We examined aqueous extracts of several Mexican native plants normally used to treat infections or illnesses with cancer symptoms to explore their antimicrobial and cytotoxic effects. In the case of *Justicia spicigera*, we used ethanolic extracts because this is the traditional manner of extraction for this plant [7].

Phoradendron longifolium, *Teucrium cubense*, *Opuntia joconostle* (against *A. lwoffii*), *Tecoma stans* (against *A. lwoffii* and *B. cepacia*), and *Guazuma ulmifolia* (against *A. lwoffii*, *E. faecalis* and *C. albicans*) exhibited high antimicrobial potentials, with $IZ \geq 15 \text{ mm}$ [11]. *Guazuma ulmifolia* showed the highest antimicrobial effects, and tested at $75 \mu\text{g/disk}$ ($IZ=26 \text{ mm}$) displayed a higher ($P \leq 0.05$) antimicrobial potency than the positive control neomycin at $25 \mu\text{g/disk}$ ($IZ=23 \text{ mm}$) against *C. albicans* (Table 2).

The most susceptible microorganism was *A. lwoffii*; eight out of ten plant extracts showed antimicrobial effects against this bacterium. In contrast, the most resistant microorganisms were *S. aureus*, *S. haemolyticus*, *A. baumannii* and *T. belgeii*, which were resistant to all plant extracts (Table 2). The results showed that there was a positive correlation between their traditional use against infections and the antimicrobial experimental effects observed. Experiments are currently being carried out in our laboratory to test fractions of *Phoradendron longifolium*, *Teucrium cubense*, *Opuntia joconostle*, *Tecoma stans* and *Guazuma ulmifolia* on multidrug-resistant microorganisms using the broth dilution method.

The National Cancer Institute (NCI) establishes that plant extracts with cytotoxic ED_{50} (Effective Dose 50) values $\leq 30 \mu\text{g/mL}$ are considered to be active [12]. According to these criteria, only *Phoradendron serotinum* and *Justicia spicigera* exerted active toxic effects against MCF-7 ($IC_{50}=28 \mu\text{g/mL}$) and HeLa ($IC_{50}=17 \mu\text{g/mL}$) cancer cells, respectively (Table 3). Table 3 only shows plant extracts that exerted toxic effects ($IC_{50} < 200 \mu\text{g/mL}$); the other extracts lacked cytotoxicity on the tested human cancer cells. Although the IC_{50} values exhibited by *Phoradendron serotinum* and *Justicia spicigera* were higher when compared with CDDP, these plants did not display toxic effects on PBMCs, and in concentrations of $200 \mu\text{g/mL}$ also increased the proliferation of these cells in a similar manner to PHA $1 \mu\text{g/mL}$ (Fig. 1). The results are of great interest because it is very desirable to find new cytotoxic agents with similar or higher potency and less toxic effects than currently used agents.




Figure 1: Effect of Mexican medicinal plant extracts on the viability of human peripheral blood mononuclear cells. PBMCs were seeded and after 1 day, cultures were fed with different concentrations of plant extracts, cisplatin (CDDP) ($5.7 \mu\text{g/mL}$) or phytohemagglutinin (PHA) ($1 \mu\text{g/mL}$) as control. After 2 days, cell viability was determined by MTT assay. GU, *Guazuma ulmifolia*; JS, *Justicia spicigera*; OJ, *Opuntia joconostle*; OL, *Opuntia leucotricha*; PA, *Parkinsonia aculeata*; PL, *Phoradendron longifolium*; PS, *Phoradendron serotinum*; PC, *Psittacanthus calyculatus*; TS, *Tecoma stans*; TC, *Teucrium cubense*

In summary, these results provide a scientific validation of the ethnomedicinal uses of the plant extracts tested. Thus, it is necessary to perform further phytochemical studies on the isolation and identification of active compounds, mainly *Guazuma ulmifolia*, *Justicia spicigera* and *Phoradendron serotinum*. Also, the mechanisms of action by which these plant extracts exert their antimicrobial and cytotoxic effects should be studied.

Experimental

Microorganisms: Eleven human pathogenic microorganism isolates, obtained and identified as previously described [13], were used for the antimicrobial tests (Table 2). The human pathogens were resistant (diameter of inhibition zones of $\leq 7 \text{ mm}$) to chloramphenicol ($30 \mu\text{g/disk}$), tetracycline ($30 \mu\text{g/disk}$), penicillin ($10 \mu\text{g/disk}$), and erythromycin ($15 \mu\text{g/disk}$), but sensitive to neomycin ($25 \mu\text{g/disk}$), as evaluated by the disk diffusion assay (see below).

Plant material and extraction: Plants were collected, taxonomically validated and preserved at the herbaria indicated in Table 3. The aqueous extracts were obtained as described previously [14]. *Justicia spicigera* extracts were obtained as follows: powdered dry leaves (10 g) were subjected to exhaustive Soxhlet extraction with ethanol for 2 h at 180°C . The extract was filtered and concentrated under reduced pressure until dry. All the supernatants were lyophilized (Freezone 4.5, Labconco, Kansas City, MO).

Cell lines and culture conditions: Human cervical carcinoma (HeLa), colorectal adenocarcinoma (SW-480) and breast carcinoma (MCF-7) cells were maintained in DMEM (GIBCO BRL, Grand Island, NY, USA) supplemented with 7% fetal bovine serum (GIBCO) and antibiotics (100 units/mL penicillin and 100 pg/mL streptomycin), whereas ovarian carcinoma (SKOV-3) and prostate carcinoma (DU-145) cells were maintained in

RPMI with 7% FBS. All human cancer cell lines were obtained from ATCC (Manassas, VA, USA) and grown at 37°C in 5% CO₂.

Antimicrobial assay: Antimicrobial tests were carried out with the disk diffusion method. Filter-paper discs (6 mm diameter) were impregnated with the plant extracts in concentrations ranging from 75 to 750 µg/disk. Negative controls were prepared using impregnated discs with vehicle and virgin discs (blank control). Neomycin (25 µg/disk) was employed as a positive control. Antibacterial activity was defined by measuring the diameter of the growth inhibition zone (IZ) around a disk after 24 h of incubation.

Cytotoxicity assay: Human cancer cells (5 x 10³ cells/well) and PBMCs (2.5 x 10⁴ cells/well), obtained as previously described [13], were seeded in 96 well plates. After 24 h of incubation, plant extracts at concentrations ranging from 1 to 200 µg/mL were added to the cells. Phytohemagglutinin (Sigma) was used as a positive control. The assay was performed as described previously [13] and the optical density (O.D.) was measured at 590 nm in an ELISA

reader (Biorad Laboratories, Hercules, CA, USA). The viability of the treated cells was estimated from the relative growth as follows:

$$\text{relative viability} = \frac{\text{control O.D.} - \text{sample O.D.}}{\text{control O.D.}} \times 100$$

The concentration leading to 50% inhibition of viability (IC₅₀) was also calculated by regression analysis (% survival versus log concentration).

Statistical analysis: Experimental values are expressed as the mean ± the standard deviation of at least 3 experiments, in triplicate. Data were analyzed using the one-way ANOVA and Student's-*t* test when necessary. The level of $p \leq 0.05$ was used to determine statistical significance. All calculations were performed using the JMP 5.1 program (SAS Institute Inc).

Acknowledgments - AJAC (174493) and RZB (233280) were endowed with graduate fellowships from CONACYT. We thank Elizabeth Ortiz-Sanchez, Miriam Guido and Rocio Méndez for their technical assistance.

References

- [1] Saklani A, Kutty SK. (2008) Plant-derived compounds in clinical trials. *Drug Discovery Today*, **13**, 161–171.
- [2] Cowan MM. (1999) Plant products as antimicrobial agents. *Clinical Microbiology Reviews*, **12**, 564–582.
- [3] Heinrich M, Kuhnt M, Wright CW, Rimpler H, Phillipson JD, Schandelmaier A, Warhurst DC. (1992) Parasitological and microbiological evaluation of Mixe Indian medicinal plants (Mexico). *Journal of Ethnopharmacology*, **36**, 81–85.
- [4] Navarro V, Villarreal ML, Rojas G, Lozoya X. (1996) Antimicrobial evaluation of some plants used in Mexican traditional medicine for the treatment of infectious diseases. *Journal of Ethnopharmacology*, **53**, 143–147.
- [5] Kumar VP, Chauhan NS, Padh H, Rajani M. (2006) Search for antibacterial and antifungal agents from selected Indian medicinal plants. *Journal of Ethnopharmacology*, **107**, 182–188.
- [6] Alonso-Castro AJ, Villarreal ML, Salazar-Olivo LA, Gómez-Sánchez M, Domínguez F, García-Carranca A. (2011) Mexican medicinal plants used for cancer treatment: pharmacological, phytochemical and ethnobotanical studies. *Journal of Ethnopharmacology*, **133**, 945–972.
- [7] Domínguez XA, Alcorn JB. (1985) Screening of medicinal plants used by Huastec Mayans of northeastern Mexico. *Journal of Ethnopharmacology*, **13**, 139–156.
- [8] González-Elizondo M, López-Enriquez IL, González-Elizondo MS, Tena-Flores JA. (2004) *Plantas Medicinales del Estado de Durango y Zonas Aledañas*. Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional Unidad Durango. Instituto Politécnico Nacional, México, pp. 210.
- [9] López R, Hinojosa A. (1988) *Catálogo de plantas medicinales sonorenses*. Universidad de Sonora, Hermosillo Sonora, México, pp. 143.
- [10] Adame J, Adame H. (2000) *Plantas curativas del Noreste Mexicano: Nuevo León, Coahuila, Tamaulipas y altiplano potosino*. Monterrey, Nuevo León, México. Ediciones Castillo, pp. 386.
- [11] Mothana RA, Lindequist U, Lindequist U, Gruenert R, Bednarski PJ. (2009) Studies of the *in vitro* anticancer, antimicrobial and antioxidant potentials of selected Yemeni medicinal plants from the island Soqatra. *Complementary and Alternative Medicine*, **9**, 7–17.
- [12] Suffness M, Pezzuto JM. (1990) Assays related to cancer drug discovery. In *Methods in Plant Biochemistry: Assays for Bioactivity*, Hostettmann, K. (Ed.), **6**, 71–133.
- [13] Jacobo-Salcedo MdR, Gonzalez-Espindola LA, Alonso-Castro AJ, Gonzalez-Martinez MdR, Dominguez F, Garcia-Carranca A. Antimicrobial and cytotoxic effects of *Magnolia dealbata* Zucc and its active compounds. *Natural Product Communications*, **6**, 1121–1124.
- [14] Alonso-Castro AJ, Salazar-Olivo LA. (2008) The anti-diabetic properties of *Guazuma ulmifolia* Lam. are mediated by the stimulation of glucose uptake in normal and diabetic adipocytes without inducing adipogenesis. *Journal of Ethnopharmacology*, **118**, 252–256.

Chemometrics Evaluation of the Herbal Drug *Andrographis paniculata*

Shiv Narayan Sharma*, Zenu Jha and D. K. Sharma

Department of Plant Molecular Biology & Biotechnology, IGKV, Raipur, India, 492-006

shivsharma.bt@ovi.com

Received: January 14th, 2011; Accepted: July 22nd, 2011

A fingerprint was developed to evaluate ten genotypes of *Andrographis paniculata* (a herbal drug) collected from different geographical locations. The genotypes were assessed by HPLC and the chromatogram areas were used to construct fingerprints. Between the two fingerprints, vector of differences was defined. The scalar mean of the deviated vector was used for statistical analysis. By incorporating appropriate chemometric methods better differentiation of samples was achieved. This provides tools for extracting relevant chemical information from the obtained data. The developed method can be used as a quality control tool for rapid authentication, identity, assessment and selection of good quality natural drug from a wide variety of herbal samples.

Keywords: *Andrographis paniculata*, quality control, andrographolide, HPLC, fingerprinting, chemometrics.

In recent years, there has been a great demand for plant derived products. These are increasingly being sought as medicinal products, nutraceuticals and cosmetics [1]. *Andrographis paniculata* Nees (family Acanthaceae), one of the most promising herbal medicines in South East Asia, is claimed to have anti-inflammatory [2a], anti-platelet aggregation [2b], antihyperglycemic [2c], hepatoprotective [2d], antitumor [2e], and anti-human immunodeficiency virus activities [2f,2g]. Andrographolide is a principal medicinal component of *A. paniculata* [2h]. It is well known that the therapeutic effects of this herbal drug are based on the synergistic effect of its constituents and, therefore, it has become essential to develop a reliable, specific and sensitive detection and quantification technique for obtaining good quality raw and processed materials.

Conventional chromatographic study focuses mainly on the determination of the active components, while fingerprinting can offer integral characterization of a complex system with a quantitative degree of reliability. In this respect, fingerprinting has gained more attention than any of the other quality control systems [3a-3d]. Fingerprinting is a method that provides chemical information about medicines from spectrograms, chromatograms and other information obtained by analytical techniques [4a,4b]. There are always chances of wide variations with respect to the chemical contents of crude drugs/raw materials of plant origin due to various reasons such as climatic conditions, geographical distribution, source and season. For this reason, it is necessary to establish a fingerprint for a medicinal herb for its quality control. The advancement of chemometric methods has become a leading tool towards faster

analytical results and shorter product development times [5a]. Hence chromatographic fingerprinting with chemometric analysis for controlling the quality of herbal medicines and their products has been accepted by many countries and organizations [5b,5c]. The objective of this work was to establish a simple and reliable technique for quantification of andrographolide and develop fingerprinting for ten genotypes of *A. paniculata* collected from different geographical locations of India.

The quantitative estimation of andrographolide was performed with regard to specificity, precision, accuracy and linearity. This was applicable in fingerprinting and variation analysis within genotype [6a,6b]. The composition of the HPLC mobile phase was optimized to achieve good resolution. The best resolution and peak shape was obtained using acetonitrile, methanol and water (65:20:15 v/v). The HPLC profile of the standard compounds was prepared. Andrographolide had a stable and high content and was completely separated from the other peaks. The presence of andrographolide in *A. paniculata* samples was verified (peak 'd') by comparison of its retention time with that of the standard chromatogram (Figure 1). Specificity was ascertained by comparing the area of the standard peak with that of the sample peak. The precision was evaluated by analysis of the sample in three replications to maintain reproducibility and repeatability. There was no significant difference in the quantification of each sample. The correlation coefficient between the standard and sample graphs was 0.9963 and the R.S.D. value of the peak areas was 1.8%. Also, the sample recoveries were determined by adding reference solutions of variable concentrations. The average value obtained was 99.4% ($n = 3$), with an R.S.D. less than




Figure 1: HPLC profile of crude extract of *A. paniculata* and selected peaks for fingerprinting (a-i).

2%. The results indicate that the method developed has acceptable precision, reproducibility and stability.

The fingerprints were constructed by the peak areas of the HPLC chromatograms of ten *A. paniculata* genotypes presented in Table 4. The chromatogram was represented as vector $a = [a_1, a_2, \dots, a_n]$, where a_i ($i = 1, 2, \dots, n$) is the area of the i^{th} chromatographic peak and n is the number of chosen peaks. The nine selected common peaks are shown in Fig 1 (a – i). The differences between the two fingerprints can be defined as $\bar{r} = a_1 - a_2$, where a_1 and a_2 represent two fingerprints of the same sample. If measurement errors are only from random noise, each peak will have a distribution of $N(\mu_i, \sigma_i^2)$, where μ_i and σ_i are the original value and the standard variance of the i^{th} peak area, respectively. The values for μ_i and σ_i were estimated by repeated measurements on the same sample. With the peak area distribution transformation into $\mu'_i = \mu_i/\sigma_i$, each peak area is divided by its standard variance. If $a_1 = a_2$, then $r = 0$, and the two fingerprints are clearly identical. If r is sufficiently small to be statistically compared with a zero vector, then it can be concluded that there is no significant difference between the two fingerprints and they are considered to be the same. On the basis of r , the similarity matrix was constructed. When $\bar{r} = 0$, it means that two fingerprints have no significant difference, if $\bar{r} \neq 0$, it means that there is a difference between them, where $\bar{r} = \frac{1}{n} \sum_{i=1}^n r_i$ and r^i is the i^{th} element of r . The relative standard deviation (RSD) values of the nine common peaks of the HPLC chromatograms among five batches of samples are shown in Table 1.

RSD values were found to be less than 0.98% for retention time and less than 4.28% for the peak area, which meant the common peaks were in good correspondence in five batches of every sample. This indicates that the pick of same batch was similar. Both the correlation coefficient and the vector cosine were calculated between five batches of fingerprints from same producing area and the results were about 0.9637 (table-2).

The fingerprints were assessed by analysis of PCA. The two principal components represent almost 92% of the total variance in the data set. The first principal components (PCs) consisted of 71% of the total variability

Table 1: RSD values of nine common peaks in HPLC chromatograms among five batches of samples.

Genotypes	No. of Samples	Peak a	Peak b	Peak c	Peak d	Peak e	Peak f	Peak g	Peak h	Peak i
AP1	5	RT	0.69	2.21	1.84	0.48	4.78	2.06	3.16	3.86
		Area	31.93	44.92	35.63	35.83	57.57	33.75	34.40	51.71
AP2	5	RT	0.76	2.20	1.82	0.55	5.89	2.00	3.02	3.61
		Area	83.15	79.44	90.46	122.35	51.05	134.38	71.87	26.39
AP3	5	RT	0.93	1.55	2.06	0.59	4.07	1.98	3.06	2.64
		Area	43.16	36.57	41.01	51.51	63.22	39.98	49.11	32.78
AP4	5	RT	0.86	1.44	0.49	0.59	4.74	2.09	3.07	2.51
		Area	36.13	22.67	29.45	47.61	46.82	31.39	36.52	25.13
AP5	5	RT	1.05	1.60	0.47	0.62	4.32	2.19	3.97	1.50
		Area	135.60	21.12	213.34	152.06	123.29	35.98	82.19	28.26
AP6	5	RT	1.02	1.64	0.41	0.43	3.61	1.56	3.85	1.49
		Area	58.10	17.11	23.09	72.93	56.91	32.88	21.62	27.05
AP7	5	RT	1.05	1.73	0.40	0.42	1.88	1.54	3.83	1.80
		Area	88.97	66.07	69.58	92.37	49.66	123.68	77.14	66.99
AP8	5	RT	0.83	1.64	0.31	0.40	3.12	1.13	3.23	1.72
		Area	55.84	103.13	103.74	48.13	168.45	75.65	83.35	156.40
AP9	5	RT	0.83	2.49	1.78	0.43	2.29	1.12	1.73	1.68
		Area	78.03	89.51	70.94	81.64	102.24	120.56	69.42	104.19
AP10	5	RT	0.66	2.46	2.26	0.41	3.15	1.33	2.23	3.12
		Area	25.69	87.97	45.06	27.52	87.33	66.13	70.15	98.36

Table 2: Result of similarity matrix of same batch of collected genotypes.

Genotypes	No of samples	Correlation coefficient	Cosine value
AP1	5	0.973 ± 0.011	0.932 ± 0.017
AP2	5	0.961 ± 0.021	0.952 ± 0.019
AP3	5	0.977 ± 0.027	0.971 ± 0.018
AP4	5	0.965 ± 0.021	0.937 ± 0.011
AP5	5	0.973 ± 0.023	0.961 ± 0.026
AP6	5	0.971 ± 0.023	0.986 ± 0.029
AP7	5	0.941 ± 0.02	0.976 ± 0.022
AP8	5	0.959 ± 0.021	0.978 ± 0.029
AP9	5	0.980 ± 0.024	0.923 ± 0.027
AP10	5	0.937 ± 0.016	0.969 ± 0.016




Figure 2: Principal component analysis of HPLC data.

followed by the second PCs with 21% variance. Overall, each sample was able to form a distinct cluster in the two-dimensional plot (Fig. 2). Examining the space defined by the first and second PCA of the HPLC data proves the variability found among the collected genotypes. The t -test statistics displayed variation which comes from the differences of μ'_i of the two corresponding peaks (Table 3). The results indicated that the region of origin will cause differences between the fingerprints of *A. paniculata*, and this can be used to formulate standard criteria for selecting the crude drug, and be a quality control standard for the planting of *A. paniculata*, as well as to provide the link for the botanical identity and quality from the number of chemical constituents present in plant [7].

SIMCA classification was employed for performing a real situation task such as assigning upcoming unknown samples [8]. SIMCA classification based on the predefined PCA model yielded 100% correct classification for all classes at 5% significance level. The SIMCA showed that the samples AP3, AP9 and AP7 maintained a higher

Table 3: Result of similarity analysis and compression of mean fingerprinting.

Fingerprint pairs	Correlation coefficient	Vector cosine	#	F = $\frac{1}{\sqrt{1-\cos^2}}$	P
AP1,AP2	-0.04247	0.0431	1.01	3.039282771	0.00
AP1,AP3	0.983874	0.9874	1.68	5.046522	0.00
AP1,AP4	0.902356	0.9023	0.64	1.913784326	0.00
AP1,AP5	0.805445	0.8624	1.53	4.615757373	0.00
AP1,AP6	0.614506	0.6256	0.93	2.790479535	0.00
AP1,AP7	0.727364	0.7364	2.20	6.603093	0.00
AP1,AP8	0.959682	0.9682	4.78	14.346605	0.00
AP1,AP9	0.825119	0.8272	2.48	7.452129	0.00
AP1,AP10	0.939898	0.9898	0.03	0.098381*	0.00
AP2,AP3	-0.01141	0.1143	0.70	2.1134	0.00
AP2,AP4	0.122771	0.1229	0.90	2.7056	0.00
AP2,AP5	0.192286	0.1953	0.149	0.44731*	0.00
AP2,AP6	-0.2859	0.2910	0.55	1.66755*	0.00
AP2,AP7	0.181364	0.1863	0.51	1.539489*	0.00
AP2,AP8	-0.03151	0.0377	0.317	0.951809829*	0.00
AP2,AP9	0.134425	0.1373	0.395	1.187798096*	0.00
AP2,AP10	-0.19422	0.1952	0.53	1.59152*	0.00
AP3,AP4	0.916221	0.9192	0.66	2.0005	0.00
AP3,AP5	0.822485	0.8434	1.19	3.570007	0.00
AP3,AP6	0.663067	0.6630	0.46	1.38715*	0.00
AP3,AP7	0.724001	0.7286	1.85	5.535267	0.00
AP3,AP8	0.944385	0.9740	2.07	6.235882	0.00
AP3,AP9	0.842407	0.8442	2.24	6.713606	0.00
AP3,AP10	0.912364	0.9233	0.43	1.12933*	0.00
AP4,AP5	0.877401	0.8812	1.47	4.432317	0.00
AP4,AP6	0.622719	0.6257	0.69	2.084494	0.00
AP4,AP7	0.908865	0.9101	3.24	9.741992	0.00
AP4,AP8	0.934218	0.9352	2.94	8.821639	0.00
AP4,AP9	0.874958	0.8749	2.66	7.994877	0.00
AP4,AP10	0.785968	0.7879	0.18	0.5403*	0.00
AP5,AP6	0.747936	0.7499	0.75	2.26173	0.00
AP5,AP7	0.740466	0.7404	1.49	2.985292	0.00
AP5,AP8	0.807512	0.8175	0.61	1.842163	0.00
AP5,AP9	0.690674	0.7907	0.76	2.303481	0.00
AP5,AP10	0.670422	0.6724	0.61	1.83601	0.00
AP6,AP7	0.540904	0.5409	1.20	3.613517	0.00
AP6,AP8	0.65615	0.6581	0.77	2.33181	0.00
AP6,AP9	0.581609	0.5886	1.09	3.285674	0.00
AP6,AP10	0.580291	0.5892	0.42	1.28251	0.00
AP7,AP8	0.865307	0.8699	0.02	0.07126*	0.00
AP7,AP9	0.902316	0.9203	0.26	0.7739*	0.00
AP7,AP10	0.584318	0.6143	0.91	2.75372	0.00
AP8,AP9	0.925151	0.9251	0.13	0.39441*	0.00
AP8,AP10	0.861045	0.8810	1.30	3.9104	0.00
AP9,AP10	0.666304	0.6734	0.90	2.7236	0.00

*Shows non-significant differences

distance than the rest of the samples. The training set ranging from 0.1305 (AP10) to 0.7784 (AP3) represents the sample to model distance. The sample distance ranges from 0.56495 (AP3) to 0.98776 (AP10). Principal component mapping by SIMCA described samples into a much lower dimensional subspace and reasonably defined the class of this herbal medicine.

Hierarchical cluster analysis was performed to generate clusters of *A. paniculata* genotypes. The data for the chromatographic peak areas of the samples were normalized to obtain an $m \times k$ matrix (where m is the number of samples and k is the number of chromatographic peaks). All the samples formed two broad categories (Fig. 3). The genotype AP10 was more diverse than the other genotypes. The established relationship between ten genotypes of *A. paniculata* by chromatographic fingerprinting and the cluster analysis




Figure 3: The hierarchical clustering of collected genotypes from different geographic locations.

Table 4: Collections of *A. paniculata* and their geographical status.

Genotype	Region	Latitude	Longitude	Mean Sea Level
AP1	Kondagaon (Chhattisgarh)	19°36' 00.00"N	81°40' 12.00"E	1934 ft.
AP2	Kanker (Chhattisgarh)	20°16' 05.78"N	81°29' 52.31"E	1597 ft.
AP3	Narayanpur (Chhattisgarh)	19°42' 53.81"N	81°14' 44.11"E	1806 ft.
AP4	Bastar (Chhattisgarh)	19°35' 03.34"N	81°41' 28.16"E	1948 ft.
AP5	Jagdalpur (Chhattisgarh.)	19°05' 00.73"N	82°01' 39.45"E	1827 ft.
AP6	Raipur (Chhattisgarh)	21°14' 34.09"N	81°38' 19.35"E	981ft.
AP7	Nagpur (Maharashtra)	21°09' 14.00"N	79°04' 59.00"E	1034 ft.
AP8	Neemach (Madhya Pradesh)	24°27' 45.03"N	74°50' 37.17"E	1619 ft.
AP9	Chitrakoot (Madhya Pradesh)	22°34' 47.09"N	80°49' 49.95"E	1061 ft.
AP10	Indore (Madhya Pradesh)	22°43' 31.13"N	75°51' 56.00"E	1810 ft.

gives the preference of selection of good quality material and is applicable for adoption for a breeding programme for crop improvement [9a,9b].

In summary, the enhanced HPLC fingerprinting method can comprehensively and properly reveal the quality characteristics of medicinal herbs, and will become a very useful complementary technique for quality determination. The results give evidence that *A. paniculata* samples from different geographical origin have a varied and complex chemical mixture.

Experimental

Plant materials and crude extraction: The present study was confined to ten genotypes of *A. paniculata*. Dried leaves of samples from 10 different geographical origins were selected based on their performance and evaluated through field experimentation at the Department of Plant Molecular Biology & Biotechnology, Indira Gandhi Agricultural University, Raipur, India [10]. The collection sites and their geographical status are shown in Table 4. Young leaves were collected from plants at flowering time, but avoiding material containing flowers or damaged by insects or fungi. Leaves were then cut into small pieces to permit homogeneous sampling. Drying was carried out at an ambient temperature of 45°C, followed by grinding. Two g of fine power was extracted by reflux with 100 mL of 70% (v/v) ethanol, with 3 cycles. The ethanolic extracts were filtered and concentrated in a rotary evaporator at 60°C. The extracts were stored at 4°C until used.

HPLC condition and standard preparation: HPLC analysis was carried out on an Agilent Series 1100 chromatograph, equipped with a vacuum degasser, a quaternary pump and a diode array detection system, connected to a reversed-phase column (Diamonsil C18, particle size 5 µm, metachem (150 mm X 4.6 mm column, Agilent, USA). Data collection was performed using Chemstation software (Agilent). The temperature was fixed at 25°C. The injection volume was 20 µL and elution was monitored at 223 nm. Andrographolide (Sigma, USA) at different concentrations (22.0 mg, 11.0 mg, and 5.5 mg) was used for method validation and preparation of calibration curves. Crude extract was evaporated under reduced pressure until dryness and used for HPLC quantification.

Fingerprinting of *A. paniculata*: The 5 samples of each *A. paniculata* genotype were collected for HPLC quantification. The areas of 9 common peaks in the chromatograms were used to construct the fingerprints. The standard variance of each peak area was calculated

from the data array and then used to normalize it. The PCA calculation was based on a singular value decomposition of the data array of the fingerprints. The first 2 scores of PCA results were used to make a projection plot that provided a visual determination of the similarity among the fingerprints. Soft Independent Model Class Analogy (SIMCA) was used to classify samples based on their similarities to principal component models. The data of the training set were used to determine principal component models for each class. One of the ways of determining the appropriate number of principal components is to use the

cross-validation technique. The classification of test set samples was obtained by adjusting each of these to each model and deciding at the 95% confidence level. Cluster analysis classification was used as a decision criterion for the distance between the samples. The average fingerprinting profile of 5 batches of every sample was applied to construct a cluster by measuring Euclidean distance. The cluster analysis was operated with the assistance of Unscrambler® 8.0 [11]. Samples were classified into classes that correspond to the classes of their nearest neighbours.

References

- [1] Bhanu PSS, Zafar R, Panwar R. (2005) Herbal drug standardization. *The Indian Pharmacist*, **4**, 19-22.
- [2] (a) Sheeja K, Shihab PK, Kuttan G. (2006) Antioxidant and anti-inflammatory activities of the plant *Andrographis paniculata* Nees. *Immunopharmacology and Immunotoxicology*, **28**, 129-140; (b) Amroyan E, Gabrielian E, Panossian A. (1996) Inhibitory effect of andrographolide from *Andrographis paniculata* on PAF-induced platelet aggregation. *Phytomedicine*, **1**, 27-31; (c) Yu BC, Hung CR, Chen WC. (2003) Antihyperglycemic effect of andrographolide in streptozotocin-induced diabetic rats, *Planta Medica*, **69**, 1075-1079; (d) Shukla B, Visen PK, Patnaik GK, Dhawan BN. (1992) Choleretic effect of andrographolide in rats and guinea pigs. *Planta Medica*, **58**, 146-149; (e) Menon V, Bhat S. (2010) Anticancer activity of andrographolide semisynthetic derivatives. *Natural Product Communications*, **5**, 717-720; (f) Kumar RA, Sridevi K, Kumar NV, Nanduri S, Rajagopal S. (2004) Anticancer and immunostimulatory compounds from *Andrographis paniculata*. *Journal of Ethnopharmacology*, **92**, 291-295; (g) Handa SS, Sharma A. (1990) Hepatoprotective activity of andrographolide against galactosamine and paracetamol intoxication in rats. *Indian Journal of Medical Research*, **98**, 284-292; (h) Samy RP, Thwin MM, Gopalakrishnakone P. (2007) Phytochemistry, pharmacology and clinical use of *Andrographis paniculata*. *Natural Product Communications*, **2**, 607-618.
- [3] (a) Nederkassel AM, Vijverman V, Massart DL, Heyden YV. (2005) Development of a *Ginkgo biloba* fingerprint chromatogram with UV and evaporative light scattering detection and optimization of the evaporative light scattering detector operating conditions. *Journal of Chromatography A*, **1085**, 230-239; (b) Srivastava A, Misra H, Verma RK, Gupta MM. (2004) Chemical fingerprinting of *Andrographis paniculata* using HPLC, HPTLC and densitometry. *Phytochemical Analysis*, **15**, 280-285; (c) Koll K, Reich E, Blatter A, Veit M. (2003) Validation of standardized high-performance thin-layer chromatographic methods for quality control and stability testing of herbals. *Journal of AOAC International*, **89**, 909-915; (d) Arpini S, Fuzzati N, Giori A, Martino E, Mombelli G, Pagni L, Ramaschi G. (2008) HPLC-DAD-MS fingerprint of *Andrographis paniculata* (Burn. f.) Nees (Acanthaceae). *Natural Product Communications*, **3**, 1981-1984.
- [4] (a) Xu YQ, Sun SQ, Feng XF, Hu SL. (2003) Quick identification of scullcaps in different geographical origins using clustering analysis method and infrared fingerprint spectra. *Spectroscopic Analysis*, **23**, 502-505; (b) Xiao LH, Wang HY, Li FM. (2004) Comparative study on HPLC-FPS of *Radix scutellariae* of various sources. *Journal of Shenyang Pharmaceutical University*, **21**, 21-31.
- [5] (a) Seasholtz MB. (1999) Making money with chemometrics. *Chemometrics and Intelligent Laboratory Systems*, **45**, 55-64; (b) Wang S, Ma HQ, Sun YJ, Qiao CD, Shao SJ, Jiang SX. (2007) Fingerprint quality control of *Angelica sinensis* (Oliv.) Diels by high-performance liquid chromatography coupled with discriminant analysis. *Talanta*, **72**, 434-436; (c) Li W, Deng Y, Dai R, Yu Y, Saeed MK, Li L, Meng W, Zhang X. (2007) Chromatographic fingerprint analysis of *Cephalotaxus sinensis* from various sources by high-performance liquid chromatography-diodearray detection-electrospray ionization-tandem mass spectrometry, *Journal of Pharmaceutical and Biomedical Analysis*, **45**, 38-46.
- [6] (a) Semmar N, Jay M, Farman M, Roux MA. (2008) New approach to plant diversity assessment combining HPLC data, simplex mixture design and discriminate analysis. *Environmental Modeling Assessment*, **13**, 17-33; (b) Giri L, Andola CH, Purohit VK, Rawat MSM, Rawal RS, Bhatt ID. (2010) Chromatographic and spectral fingerprinting standardization of traditional medicines: an overview as modern tools. *Research Journal of Phytochemistry*, **4**, 234-241.
- [7] Gan F, Ye R. (2006) New approach on similarity analysis of chromatographic fingerprint of herbal medicine. *Journal of Chromatography A*, **1104**, 100-105.
- [8] Hatanaka RR, Flumignan DL, de Oliveira JE. (2009) GC fingerprints coupled to pattern-recognition multivariate SIMCA chemometric analysis for Brazilian gasoline quality studies. *Chromatographia*, **70**, 1135-1142.
- [9] (a) Qiang W, Xiao R, Jun-hua H, Ning-yi X, Qi-chuan Y. (2006) Intra-specific genetic relationship analyses of *Elaeagnus angustifolia* based on RP-HPLC biochemical markers. *Journal of Zhejiang University Science B*, **7**, 272-278; (b) Lonni AASG, Scarimino IS, Silva LMC, Ferreria DT. (2003) Differentiation of species of the *Baccharis* genus by HPLC and chemometric methods. *Analytical Science*, **19**, 1013-1018.
- [10] Sharma SN, Sinha RK, Sharma DK, Jha Z. (2009) Assessment of intra-specific variability at morphological, molecular and biochemical level of *Andrographis paniculata* (Kalmegh). *Current Science*, **96**, 402-407.
- [11] Unscrambler® 8.0 CAMO Software <http://www.camo.com/>

Garcinia cambogia Leaf and Seawater for Tannase Production by Marine *Aspergillus awamori* BTMFW032 under Slurry State Fermentation

Beena P. S, Soorej M. Basheer^a, Sarita G. Bhat^a and Chandrasekaran M^{a,b}

^aMicrobial Technology Laboratory, Department of Biotechnology, Cochin University of Science and Technology, Cochin, Kerala, India 682022

^bDepartment of Botany and Microbiology, College of Science, King Saud University, PB No.2455, Riyadh-11451, Kingdom of Saudi Arabia

beena.santhosh@gmail.com

Received: September 15th, 2010; Accepted: October 27th, 2011

Garcinia gummi-gutta (syn. *G. cambogia*, *G. quaesita*), known to have medicinal properties, was evaluated as a substrate and inducer for tannase production by a marine *Aspergillus awamori* BTMFW032, under slurry state fermentation using Czapekdox-minimal medium and sea water as the cultivation medium. Among the various natural tannin substrates evaluated, *Garcinia* leaf supported maximal tannase production. The cultivation conditions and components of the cultivation medium were optimized employing response surface methodology. The experimental results were fitted to a second-order polynomial model at a 92.2% level of significance ($p < 0.0001$). The maximal tannase activity was obtained in a slurry state medium containing 26.6%, w/v, *Garcinia* leaf, supplemented with 0.1% tannic acid as inducer. The optimum values of pH, temperature and inoculum concentration obtained were 5.0, 40°C and 3%, respectively. A Box-Behnken model study of the fermentation conditions was carried out, and the best production of tannase was registered at 40°C without agitation. Optimization strategy employing response surface methodology led to nearly 3-fold increase in the enzyme production from 26.2 U/mL obtained in unoptimized medium to 75.2 Units/mL in Box Behnken design, within 18 h of fermentation. It was observed that sea water could support maximal tannase production by *A. awamori* compared with other media suggesting that the sea water salts could have played an inducer role in expression of tannase encoding genes. To the best of our knowledge, this is the first report on production of tannase, an industrially important enzyme, utilizing *Garcinia* leaf as substrate under slurry state fermentation by marine *A. awamori* and sea water as the cultivation medium.

Keywords: Tannase, *Aspergillus awamori*, *Garcinia cambogia*, slurry state, response surface methodology, Box Behnken model.

Tannin Acyl Hydrolase (E.C.3.1.1.20), commonly referred to as tannase, catalyzes the hydrolysis of ester and depside bonds present in hydrolysable tannins to form glucose and gallic acid. Tannases are enzymes produced by microbes that are responsible for the *in vivo* hydrolysis of tannins, polyphenols that are widely present in plants. Tannase is known as a versatile enzyme with several interesting properties suitable for industrial applications. It is used for the production of gallic acid, which is used in the pharmaceutical industry for production of the anti-bacterial drug trimethoprim [1], as a substrate for the synthesis of propyl gallate, an anti-oxidant, and in the food industry [2]. Tannase is also used in the manufacture of instant tea and acorn wine, in the clarification of beer and fruit juices; the manufacture of coffee flavored soft drinks; improvement in the flavor of grape wine; as an analytical probe for determining the structure of naturally occurring gallic acid esters [3], and for cleavage of polyphenolics, such as dehydrodimer crosslinks present in the cell wall of plants, which is necessary for plant cell wall digestibility [4]. Tannase is useful for hydrolysis of tannins known to

discolor the tooth surface. In animal feed, reduction of the anti-nutritional effects of tannins is minimized by tannase. Tannase can be potentially used for the degradation of tannins present in the effluents of tanneries, which represent serious environmental problems [5].

Tannases are enzymes produced by microbes and are responsible for the *in vivo* hydrolysis of tannins, polyphenols that are widely present in plants. Tannase is produced by a large number of fungi, of which terrestrial *Aspergillus awamori* Nakazawa is one [3,6]. Tannase produced by *A. awamori* was purified and characterized [6]. The fungal strain used in the present study was previously identified as *A. awamori* and as a potential lipase producer [7].

Solid-state fermentation (SSF) is generally preferred for enzyme production owing to the fact that it allows economic production of highly concentrated crude enzymes at low cost of extraction of the pure enzymes [8]. However, it was noted that during SSF production of tannase by *A. awamori* BTMFW032, there was a very high

level of spore formation and consequent difficulty in handling of the fermented moldy bran. So, slurry state fermentation (SLSF) was evaluated as an alternative to submerged fermentation (SmF) and SSF. In fact, enzyme production in SLSF has been employed in limited cases such as pectinase and single cell protein production by *A. niger* and *Trichoderma viride* on pulps from lemon juice clarification [9].

Statistical experimental designs provide an efficient approach to optimization. Response surface methodology (RSM) is a powerful technique for testing multiple process variables, because fewer experimental trials are needed as compared with studying one variable at a time. Also, significant interactions between the variables can be identified and quantified by this technique. The Box Behnken design is especially suitable for accounting for the interactions and identifying the more significant components in a medium. A combination of factors generating a certain optimal response can be identified from the use of a factorial design and response surface methodology. Tannic acid concentration, agitation speed, and pH during the fermentation were identified as important process parameters effecting cell growth and enzyme synthesis by *A. awamori*. These parameters were optimized in a laboratory bioreactor by response surface methodology using Box and Behnken factorial design to determine the optimum conditions for enzyme production and gallic acid accumulation [3].

Commercial success of any fermentation process for industrial enzyme production very much relies on a cheap substrate that is easily obtainable, and which results in a high process yield. Although several natural substrates were tried for different enzyme production, to the best of our knowledge, *Garcinia* sp. (family Clusiaceae) was never tried for any fermentation. *G. cambogia* is a moderate-sized, evergreen tree native to South India and Southeast Asia. The fruit of this species has been traditionally used in food preparation and cooking, having a distinctive taste, but the leaves were not recognized for any utility. *Garcinia* is a source of a revolutionary natural diet ingredient that is currently popular in the US, Japan, Europe, and other western countries. In this study, the prospects of utilizing naturally available *Garcinia* leaves were tried as a source of carbon and inducer of tannase production for SLSF. Previous studies of tannase production by *A. awamori* [3,6], did not explore SLSF, utilization of natural substrate, and sea water as an enzyme production medium. Here we report the role of sea water and natural substrate resources, and the prospects of SLSF in tannase production by the marine fungus *A. awamori* BTMFW032, which holds promise as a tannase producer for further industrial application.

Selection of natural substrates as carbon source: Various natural substrates were tried as a carbon source, both independently and in combination with tannic acid, for tannase production by *A. awamori* BTMFW032. It is

evident from the data presented in Table 1 that, irrespective of the substrate used and whether the medium is prepared with sea water or not, tannic acid influences tannase synthesis by the fungus, whereas in the case of a few substrates, tannase production by the fungus was observed even in the absence of tannic acid. In fact, tamarind fruit in combination with tannic acid in sea water medium recorded maximum enzyme activity (374.67 U/mL), followed by grape wastes and tannic acid in Czapekdox medium (369.2 U/mL), and *Garcinia* leaves and tannic acid in sea water medium (256.0 U/mL). Moreover, it was observed that *Garcinia* leaves alone could induce tannase production by the fungus (22.1 U/mL in Czapekdox medium and 26.2 U/mL in sea water) suggesting their potential for use as a substrate for tannase production. All other substrates were not effective and recorded tannase at very low level compared with the control. Furthermore, tamarind fruit and grape wastes are not economical and not easily available, even though they showed enhanced production when used as a carbon source and inducer. Hence, *Garcinia* leaf was selected as a potential substrate for tannase production. It was also noted that *Garcinia* contains 5 mg tannic acid equivalents in 500 mg of leaf (i.e., 1% tannic acid equivalent was present in *Garcinia* leaves, by the Folin-denisins method). Probably the presence of tannic acid in *Garcinia* might be the causative factor for the production of tannase enzyme by *A. awamori*. The data also provide evidence for the positive role played by sea water as a medium for enhanced production of tannase by *A. awamori* BTMFW032. Maybe the various ions and inorganic salts present in the sea water could influence the fungus by satisfying its requirements for enhanced synthesis of tannase. Of course, a detailed study is warranted to establish this fact. The data further indicate that, in spite of the fact that natural substrates could induce tannase production by the fungus, the organism requires tannic acid in the medium for enhanced synthesis, confirming the inducer role of tannic acid for tannase production by *A. awamori*.

Selection of the variables that significantly affect tannase production: Plackett-Burman (PB) design offers an effective screening procedure and computes the significance of a large number of factors in one experiment, which is time saving and maintains convincing information on each component [15]. The tannase production with PB design experiments showed a wide variation of enzyme concentrations from 26.2 to 57.4 U/mL, which indicated the importance of medium optimization to attain higher yields.

The statistical significance of the model equation, evaluated by the F-test analysis of variance (ANOVA), revealed that the obtained regression is statistically significant. The model F value of 8.36 and values of Prob>F less than 0.05 provide evidence that the model terms are significant. Furthermore, the results presented as a pareto chart (Figure 1) for the effect of individual

Table 1: Effect of various natural substrates as carbon source on tannase production by *Aspergillus awamori* BTMFW032 (enzyme activity assayed after 48 h of fermentation).

No	Natural substrate	Media	Inducer	Enzyme activity U/mL
1	Control	Seawater 50%,v/v, Czapekdox 50%,v/v	Tannic acid (1%,w/v)	240.5
2	Mango leaves	Czapekdox Seawater	Tannic acid	13.4
3	Mango leaves	Seawater	Tannic acid	172.6
4	Grape wastes	Seawater	Tannic acid	369.2
5	<i>Garcinia</i> leaves	Seawater	Tannic acid	256.0
6	<i>Garcinia</i> leaves	Czapekdox	Tannic acid	188.7
7	<i>Garcinia</i> leaves	Czapekdox	-	22.1
8	<i>Garcinia</i> leaves	Seawater	-	26.2
9	Cashew apple	Seawater	Tannic acid	154.8
10	Tea powder	Seawater	Tannic acid	98.7
11	Tamarind fruit	Czapekdox	Tannic acid	374.7
12	Tamarind pods,	Czapekdox	Tannic acid	0.0
13	<i>Psidium</i> leaves,	Czapekdox	Tannic acid	0.0
14	Coconut fiber	Czapekdox	Tannic acid	0.0
15	Coconut pith	Czapekdox	Tannic acid	0.0
16	<i>Averrhoa bilimbi</i> fruits	Czapekdox	Tannic acid	0.0
17	<i>A. bilimbi</i> leaves	Seawater	Tannic acid	0.0




Figure 1: Pareto chart showing the positive and negative effects of individual factors.

parameters studied in PB design testify that tannic acid and inoculum concentration had a positive effect in enhancing enzyme production, along with the increase in their concentrations, whereas glucose, temperature and *Garcinia* leaves had a negative effect, along with an increase in variable. Validation of the PB design was carried out in shake flasks under conditions predicted by the model and it was noted that the experimental values were very close those predicted and hence the model was successfully validated. From the results obtained with PB design it is inferred that among the eighteen variables evaluated, five, namely *Garcinia* leaf, tannic acid, glucose, temperature and inoculum were found to be the most significant.

First order model equation: Tannase activity Y (U/mL) = +44.14250 -0.28315**Garcinia* -76550*glucose +140.0100*tannic acid. 39740*temperature +5.02600*inoculum.

RSM using Box-Behnken design was adopted towards selection of the optimal level of significant variables (*Garcinia* leaf, tannic acid, glucose, inoculum concentration and temperature), based on the PB design experiment. The results obtained were analyzed by ANOVA, with the following regression equation for the level of tannase production.

$$\begin{aligned} \text{Tannase activity } Y(\text{U/mL}) = & +1.29328 -0.18255*\textit{Garcinia} \\ & +1.81366*\text{glucose} +541.516*\text{tannic acid} +1.97696*\text{temperature} \\ & +1.24176*\text{inoculum} -0.047544*\textit{Garcinia}^2 -0.16906*\text{glucose}^2 \\ & -3269.474*\text{tannic acid}^2 -0.046906*\text{temperature}^2 -1.98237*\text{inoculum}^2 \\ & -0.037727*\textit{Garcinia}*\text{Glucose} -0.88075*\textit{Garcinia}*\text{tannic acid} \\ & +0.056280*\textit{Garcinia}*\text{temperature} +0.21372*\textit{Garcinia}*\text{inoculum} \\ & -14.45289*\text{glucose}*\text{tannic acid} +0.022286*\text{glucose}*\text{inoculum} +1.18215* \\ & \text{tannic acid}*\text{temperature} +51.60360*\text{tannic acid}*\text{inoculum}. \end{aligned}$$

The ANOVA analysis of tannase production showed that the model F value of 14.84 and values of Prob>F less than 0.05 were significant. Three linear and six quadratic terms were significant model terms for the response.

The model coefficients estimated by multiple linear regressions and ANOVA showed that the model was significant with coefficient of determination R^2 of 0.9223. This ensured a satisfactory adjustment of the quadratic model to the experimental data and indicated that approximately 92.2% of the variability in the dependent variable (response) could be explained by the model. The adjusted R^2 , which is more suited for comparing models with different number of variables, was 0.8601. All selected parameters were significant and varied levels of interactions were recorded for the variables in their cumulative effect on tannase production. The coefficient of variance was 6.53, and the adequate precision that measures the signal to noise ratio was 15.203. A ratio greater than 4 is desirable as it indicates an adequate signal. Thus this model could be used to navigate the design space.

Experimental data on the effect of five selected physicochemical factors on production of tannase by *A. awamori* BTMFW032 in a total of 46 experiments showed strong dependence on the presence and levels of the selected factors as the enzyme production varied between 37.4 U/mL and 76.8 U/mL under the experimental conditions studied.

Analysis of factors influencing tannase production:

Three dimensional response surface curves were plotted to study interaction among various physicochemical factors and to determine the optimum concentration of each individual factor for maximum tannase production. The model predicted 75.8 U/mL of tannase enzyme activity and obtained a maximum of 75.2 U/mL. It was noted that tannic acid, which is used as an inducer, has a very high influence on the enzyme production and its higher concentration supported maximum enzyme production.

Interaction between factors: The pair wise interactions among the factors in terms of tannase production under the optimized condition were assessed by examining the response surfaces. Three dimensional response surfaces were generated holding three factors constant at a time and plotting the response obtained for varying levels of the other two. With increase in tannic acid, enzyme activity increased, but at higher and lower concentrations of *Garcinia*, there was no interactive effect on the production of enzyme (Figure 2).

At higher concentrations of tannic acid, the medium level of inoculum concentration showed more enzyme activity compared with the higher and lower levels, whereas *Garcinia* showed less interaction with inoculum concentration at its higher, middle and lower levels.




Figure 2: Interaction between tannic acid and *Garcinia* concentrations.

The parabolic nature of the graph indicates that the interactive effect of *Garcinia* and glucose was very high at higher concentrations of tannic acid; the higher and lower levels of *Garcinia* and glucose showed lesser enzyme activity compared with the medium level (Figure 3).




Figure 3: Interaction between glucose and *Garcinia* concentration.

At higher and lower concentrations of glucose and temperature enzyme productivity was less compared with the medium level. A total of 18 variables were checked and, out of these, five parameters were optimized by RSM. These were *Garcinia* leaf (26%), glucose (3.2 mM), tannic acid (0.1%), temperature (40°C) and inoculum concentration (3%). Time course studies using *A. awamori* under final optimized conditions revealed that tannase production increased rapidly during the initial stages of fermentation and maximum enzyme activity was recorded at 24 h (Figure 4). However, with further progress of fermentation, the enzyme activity declined.

In an earlier study [3], tannic acid concentration, agitation speed and pH during the fermentation were identified as important process parameters effecting cell growth and enzyme synthesis by *A. awamori*, and these parameters were optimized in a laboratory bioreactor by RSM using

Box-Behnken factorial design to determine the optimum conditions for enzyme production and gallic acid accumulation. In that report, under optimum process conditions for enzyme synthesis, the fermentation run lasted 60 h with an initial tannic acid concentration of 35.0 g/L,




Figure 4: Time course study under optimized conditions for tannase production by *A. awamori* BTMFW 032.

yielding a biomass concentration of 7.13 g/L containing 771 IU of intracellular tannase per g dry cell weight and 19 g/L of gallic acid. However, maximum gallic acid accumulation (40.3 g/L) was obtained in 24 h with an initial substrate concentration of 45 g/L [3]. Jamun leaves and amla leaves were used for tannase production [16] where a maximum enzyme yield of 69 U/g dry substrate was obtained after 96 h of incubation. At the same time, palm kernel cake and tamarind seed powder as substrate, respectively, supported maximal enzyme activity of 13.0 U/g and 6.4 U/g dry substrate [17], whereas wheat bran enriched with 0.8% tannic acid supported higher enzyme activity of 67.5 U/g dry substrate [18].

Conclusion: Synthetic substrate tannic acid, which is expensive and difficult to obtain on a large scale, is an important issue affecting the viability of production of fungal tannases by industrial fermentation. Hence any natural substrate that can support enhanced production of tannase either alone or in combination with tannic acid would have great impact on enzyme production in the industry. In this context the results obtained in the present study with different natural substrates have strongly indicated the prospects of *Garcinia* leaves, which are normally discarded, as a substrate and inducer for tannase production by *A. awamori* BTMFW032. There was clear indication that when *Garcinia* was used in combination with tannic acid, tannase levels were at a higher level than when tannic acid alone was used as substrate. It is speculated that *Garcinia* is a potential material that can be used as a substrate for tannase production owing to its utilizable tannin content. Further, in the present study, it was observed that sea water could support maximal tannase production by *A. awamori* BTMFW032 compared with other media suggesting that the sea water salts could have played an inducer role in expression of tannase encoding genes, which of course needs further investigation. Optimization strategy employing RSM led to a nearly 3-fold increase in the enzyme production from 26.2 U/mL obtained in unoptimized media to 75.2 U/mL in Box Behnken design, within a short period of incubation. To the best of our knowledge, this is the first report on production of tannase, an industrially important enzyme, utilizing *Garcinia* leaf as substrate under slurry state fermentation by marine *A. awamori* BTMFW032 and sea water as cultivation medium.

Experimental

Strain and cultivation conditions: *A. awamori* BTMFW032, isolated from seawater from the Arabian Sea as part of an earlier investigation and available as stock culture at the Department of Biotechnology, Cochin University of Science and Technology, was used in the present study for the production of tannase. The strain was maintained on Czapekdox -minimal agar media with 1% tannic acid (w/v) as the sole carbon source. Since tannase is an inducible enzyme, pre-induced inoculum was used throughout the study. Inoculum was prepared as a spore suspension by the addition of 5 mL sterile physiological saline to a 10 day old agar slope culture, and mixed well for 30 min on a shaker, at room temperature (RT, 28±2°C). The suspension, with spores adjusted to a spore concentration of 10⁸ spores per mL using sterile physiological saline, was used as inoculum.

Slurry state fermentation (SLSF)

Substrate preparation: The fresh leaves were dried at 60°C for 24 h, coarsely ground, and stored at 4°C until further use [10]. In general, substrates for SLSF were prepared with a coarsely ground matrix of the selected natural substrate. Cultivation of the fungus for tannase production was carried out in 250 mL Erlenmeyer flasks. The coarsely ground matrixes of the substrates and commercially available tea powder were moistened with Czapekdox minimal medium and seawater (10% matrix (w/v) in minimal medium) and inoculated with the prepared spore inoculum at a concentration of 1%, v/v, and incubated for 2 days at RT (28±2°C) and 100 rpm in a rotary shaker. After incubation, the fermented slurry was centrifuged at 4°C and 10,000 rpm for 15 min, and the supernatant was collected and assayed for tannase activity.

Selection of natural substrates as carbon source: A univariable strategy of natural carbon source was employed to screen and select the best natural substrate for induction of tannase production by the fungus as a substitute/supplement to synthetic tannic acid which is currently used for the production of the enzyme. The natural substrates tried were basically tannin containing agro residues, which included fruits, different leaves, and those in combination with tannic acid, like commercially available tea powder, freshly dried and ground mango leaves, tamarind fruit, dry tamarind pods, *Garcinia cambogia* leaves, *Psidium* leaves, coconut fiber and pith, grape waste, cashew apple, and *Averrhoa bilimbi* fruits and leaves. Their concentrations were variably fixed and induced at 10% concentration (w/v). Minimal medium and seawater with tannic acid as sole carbon source were maintained as controls.

Tannin estimation: Tannin content in the natural substrates was estimated following the Folin-Denis method [11]. Tannin-like compounds reduce phosphotungstomolybdic acid in alkaline solution to produce a highly colored blue solution, the intensity of which is proportional to the amount of tannins. An aliquot of 0.5 g of powdered

material was diluted with 75 mL distilled water, boiled for 30 min and centrifuged at 2000 rpm for 20 min. The supernatant was collected and made up to 100 mL. From this solution, 1 mL was diluted with 75 mL distilled water and to it was added 5 mL of Folin-Denis reagent, followed by 10 mL of sodium carbonate and made up to 100 mL. The intensity was measured in a spectrophotometer at 700 nm after 30 min. The tannin content of the sample was calculated as tannic acid equivalents from a tannic acid standard graph.

Tannase assay: Tannase activity was estimated based on the formation of the chromogen between gallic acid (released by the action of tannase on methyl gallate) and Rhodanine (2-thio-4-ketothiazolidine) [12]. The enzyme reaction mixture was prepared by the addition of 0.05M citrate buffer, 0.01M methyl gallate and crude enzyme, and incubated at 30°C for 5 min. Later, the reaction mixture was mixed with 0.667% methanolic rhodanine and incubated further under the same conditions. Then, 0.5M potassium hydroxide was added and incubated further. Finally the enzyme reaction mixture was diluted 5 times with distilled water and the pink color that developed was read at 520 nm using a spectrophotometer (UVShimadzu-160A). A set of blanks and controls were also maintained. One unit of tannase activity (U) is defined as the amount of enzyme required to liberate 1 µM of gallic acid per minute under defined conditions.

Selection of factors that significantly affect tannase production: Factors that significantly influence tannase production were selected statistically employing the PB method [13] in the first phase followed by RSM using Box-Behnken design. The variables and their experimental levels studied using the PB method for selection of the variables that had the largest influence on the experimental response is presented in Table 2. The variables selected by the PB method were later applied to Box-Behnken design in order to evaluate the individualized influence of each variable, as well as the effect of their interactions on enzyme production [14].

Table 2: Variables and their experimental levels adopted in Plackett –Burman method for selection of variables that influence tannase production by *Aspergillus awamori* BTMFW032.

S.No	Variable	Upper level (+)	Lower level (-)
1	<i>Garcinia</i> leaves	30%, w/v	10%
2	Seawater	75%	25%
3	Sodium nitrate	50 mM	10 mM
4	Potassium chloride	10 mM	5 mM
5	Magnesium sulfate	10 mM	5 mM
6	Zinc sulfate	0.05 mM	0
7	Copper sulfate	0.025 mM	0
8	Ferrous sulfate	0.5 mM	0.1 mM
9	Dipotassiumhydrogen phosphate	15 mM	5 mM
10	Glucose	10 mM	0
11	Tannic acid	0.1%, w/v	0
12	pH	5	2
13	Temperature	40°C	25°C
14	Inoculum	3%, v/v	0.5%, v/v
15	Agitation	150 rpm	0
16	Illumination	Dark	Light
17	Incubation hours	48 h	12 h
18	Calcium chloride	5 mM	0

Optimization of production by RSM: Five independent variables were used to obtain the combination of values that optimizes the response within the region of 3-dimensional (3D) observation spaces, which allows one to design a minimal number of experiments. The experiments were designed using the software, Design Expert Version 6.0 (State Ease, Minneapolis, MN, USA). The components (independent variables) selected for the optimization were: concentration of the substrate (*Garcinia* leaves), tannic acid, glucose, inoculum concentration and temperature. Box-Behnken design is a second order design for the estimation of quadratic effect and a 2 level factorial design. Each factor was studied at 3 different levels, the levels coded in units with values -1, 0, and 1 representing lower, middle, and higher values, respectively. The quadratic model chosen to represent the relationship fitted for the 5 variables was:–

$$Y = \beta_0 + \sum \beta_i X_i + \sum \beta_{ii} X_i^2 + \sum \sum \beta_{ij} X_i X_j$$

Y-variable enzyme yield, X1, X2, X3, X4, X5 are the independent variables selected. Analysis of variance (ANOVA) was performed and 3 dimensional response surface curves were plotted by Design Expert software to study the interaction among the various physico-chemical factors. The coded and actual values of independent variables are given in Table 3.

References

- Bajpai B, Patil S. (1996) Tannin acyl hydrolase activity of *Aspergillus*, *Penicillium*, *Fusarium* and *Trichoderma*. *World Journal of Microbiology and Biotechnology*, **12**, 217-220.
- Lekha PK, Lonsane BK. (1997) Production and application of tannin acyl hydrolase: state of the art. *Advances in Applied Microbiology*, **44**, 215-260.
- Seth M, Chand S. (2000) Biosynthesis of tannase and hydrolysis of tannins to gallic acid by *Aspergillus awamori*, optimization of process parameters. *Process Biochemistry*, **36**, 39–44.
- Garcia-Conesa MT, Kauppinen S, Williamson G. (2001) Hydrolysis of diethyl diferulates by a tannase from *Aspergillus oryzae*: Breaking cross-links between plant cell wall polymers, *Carbohydrate Polymer*, **44**, 19-24.
- Van de Lagemaat J, Pyle DL. (2001) Solid-state fermentation and bioremediation: development of a continuous process for the production of fungal tannase. *Chemical Engineering Journal*, **84**, 115-123.
- Kalyani M, Ranjan KN, Subhendu SB, Rintu B, Ashok P, George S. (2005) Purification, characterization and some studies on secondary structure of tannase from *Aspergillus awamori* Nakazawa. *Process Biochemistry*, **40**, 3251-3254.
- Basheer SM. (2008) Lipase production by marine fungus *Aspergillus awamori* Nagazawa Ph.D. thesis, Cochin University of Science and Technology, India.236.
- Tao S, Peng L, Beihui L, Deming L, Zuohu L. (1997) Solid state fermentation of rice chaff for fibrinolytic enzyme production by *Fusarium oxysporum*. *Biotechnology Letters*, **19**, 465-467.
- De Gregorio A, Mandalari G, Arena N, Nucita F, Tripodo MM, Lo Curto RB. (2002) SCP and crude pectinase production by slurry-state fermentation of lemon pulps. *Bioresource Technology*, **83**, 89-94.
- Zhong W, Cen P. (2005) Pectinase production by *Aspergillus niger* P-6021 on citrus Changshan-huyou peel in slurry-state fermentation. *Chinese Journal of Chemical Engineering*, **13**, 510-515 .
- Schanderl SH. (1970) In *Methods in food analysis*, Academic Press, NewYork,709.
- Sharma S, Bhat TK, Dawra RK (2000) A spectrophotometric method for assay of tannase using rhodanine. *Analytical Biochemistry*, **279**, 85-89.
- Plackett RL, Burman, JP. (1946) The design of optimum multifactorial experiments. *Biometrika*, **33**, 305-325.
- Box GEP, Hunter WG, Hunter JS. (1978) *Statistics for Experimenters*. John Wiley & Sons, New York, USA, 653.
- Sharma DC, Satyanarayana T. (2006) A marked enhancement in the production of a highly alkaline and thermostable pectinase by *Bacillus pumilus* Dcsr1 in submerged fermentation by using statistical methods, *Bioresource Technology*, **97**, 727–733.
- Rakesh K, Jitender S, Randhir S. (2007) Production of tannase from *Aspergillus ruber* under solid-state fermentation using jamun (*Syzygium cumini*) leaves. *Microbiological Research*, **162**, 384-390.
- Sabu A, Pandey A, Daud MJ, Szakacs G. (2005) Tamarind seed powder and palm kernel cake: two novel agro residues for production of tannase under solid state fermentation by *Aspergillus niger* ATCC16620. *Bioresource Technology*, **96**, 1223-1228.
- Gustavo ASP, Selma GFL, Selma CT, Sonia C (2001) Selection of tannase-producing *Aspergillus niger* strains. *Brazilian Journal of Microbiology*, **32**, 24-26.

Table 3: Variables and their experimental levels adopted in Box-Behnken method for tannase production by *Aspergillus awamori* BTMFW032.

S.No	Variables	Low level (-1)	Medium level (0)	High level (+1)
1	<i>Garcinia</i> leaves	10%, w/v	20%, w/v	30%, w/v
2	Tannic acid	0	0.05%, w/v	0.1%, w/v
3	Glucose	0	5 mM	10 mM
4	Temperature	25 °C	32.5°C	40°C
5	Inoculum	0.5%, v/v	1.75%, v/v	3%,v/v

All experiments were carried out in triplicate. Replicates at the centre of the domain in 3 blocks permit the checking of the absence of bias between several sets of experiments. The effect of variables and their interactions and all the coefficients were calculated by the software package Expert Version 6.0.

Experimental validation of the optimized conditions: The validation of the response surface was carried out through solutions obtained and a time course study conducted with the optimized parameters in shake flasks in triplicate.

Acknowledgment - The authors wish to thank Department of Science and Technology, India for their financial and technical support under SERC-DST-WOS-A Project given to Beena P.S.

Gas Chromatographic Quantitative Analysis of Methanol in Wine: Operative Conditions, Optimization and Calibration Model Choice

Rosario Caruso*, Grazia Laura Gambino, Monica Scordino, Leonardo Sabatino, Pasqualino Traulo and Giacomo Gagliano

Dipartimento dell'Ispettorato Centrale della Tutela della Qualità e Repressione Frodi dei Prodotti Agroalimentari, Laboratorio di Catania, Via A. Volta 19, 95122 CATANIA, Italy

r.caruso@mpaaf.gov.it

Received: July 26th, 2011; Accepted: September 28th, 2011

The influence of the wine distillation process on methanol content has been determined by quantitative analysis using gas chromatographic flame ionization (GC-FID) detection. A comparative study between direct injection of diluted wine and injection of distilled wine was performed. The distillation process does not affect methanol quantification in wines in proportions higher than 10%. While quantification performed on distilled samples gives more reliable results, a screening method for wine injection after a 1:5 water dilution could be employed. The proposed technique was found to be a compromise between the time consuming distillation process and direct wine injection. In the studied calibration range, the stability of the volatile compounds in the reference solution is concentration-dependent. The stability is higher in the less concentrated reference solution. To shorten the operation time, a stronger temperature ramp and carrier flow rate was employed. With these conditions, helium consumption and column thermal stress were increased. However, detection limits, calibration limits, and analytical method performances are not affected substantially by changing from normal to forced GC conditions.

Statistical data evaluation were made using both ordinary (OLS) and bivariate least squares (BLS) calibration models. Further confirmation was obtained that limit of detection (LOD) values, calculated according to the 3σ approach, are lower than the respective Hubaux-Vos (H-V) calculation method. H-V LOD depends upon background noise, calibration parameters and the number of reference standard solutions employed in producing the calibration curve. These remarks are confirmed by both calibration models used.

Keywords: wine, methanol, distillation, gas chromatography, ordinary least squares, bivariate least squares detection limits.

Due to its cheapness and easy accessibility, methanol has been used for adulteration of alcoholic beverages and wines. In recent times, this fraudulent behavior has lead to either the death or blindness of many people in some developing countries almost every year. In 1986, Italy numbered tens of victims caused by accidental methanol intake. Methanol intake causes headache, vertigo, fatigue, nausea, vomiting, partial or irreversible blindness, and even death. The human minimum lethal dose of methanol ranges from 300 to 1000 mg/kg body weight [1]. A number of methods for the determination of methanol have been proposed. Upadhyay and Gupta developed a spectrophotometric determination of methanol [2]. The AOAC Official Methods of Analysis include a chromotropic colorimetric method [3], a titrimetric method [4], and a gas chromatographic method [5]. Enzymatic methods with spectrophotometric and chemical sensor detection were proposed by Mizgunova *et al.* [6], and Sun *et al.* [7], respectively. A method using a biosensor with chemiluminescence detection was reported by Sekine *et al.* [8]. Van der Berg *et al.* have employed near-infrared spectroscopy in the control of alcohol process quality

assurance [9]. Determination of methanol in biological samples with electrochemical detection in high performance liquid chromatography was performed by Tagliaro *et al.* [10]. Mei-Ling *et al.* proposed several gas chromatographic methods for methanol [11], and simultaneous determination of ethanol and methanol [12]. Although, for methanol analysis, several methods based on different principles have been proposed in the past, a study of the effects of chromatographic conditions and sample pretreatments on its quantification has not been carried out. Therefore, a comprehensive statistical study of detection limits, quantification limits, and calibration curve model has been performed.

In analytical chemistry the detection limit is defined as the lowest concentration that provides an instrumental signal distinguishable from background noise. The Environmental Protection Agency (EPA) procedures identify this limit as three times the standard deviation of the blank. The overlap regions between the signals of the blank and the analyte are distance from zero dependent (Figure 1). The standard deviations (σ) are statistically




Figure 1: Overlap regions between blank signal and a response at 1σ , $z = 0.3085$ (a); 2σ $z = 0.1587$ (b); and 3σ $z = 0.0688$ (c).

compatible with concentration levels. When the instrumental response is σ times higher than the blank, the overlap between blank and analyte distributions is 30.9% (Figure 1 a). The overlap decreases to 15.9% if the blank has a 2σ distance from a blank (Figure 1 b). The percentages of overlap reduce to 6.9 and 2.3 for 3σ (Figure 1 c) and 4σ (data not shown) blank-analyte distances, respectively. The overlap region decreases in an asymptotic way with the distances from zero (Figure 2). These considerations are made under Gaussian distribution conditions.

With the limit of detection (LOD), while the 3σ approach takes into account only false positives α , Hubaux-Vos (H-V) also considers false negatives β . H-V LOD can be carried out by either graphical methods or a numerical iterative process [13]. The choice of a linear calibration model, when experimental data have to be fitted, depends on the uncertainties associated with the axes. The ordinary least squares (OLS) method is probably the most widely used regression technique. Often OLS has a limited scope, since it considers the x-axis to be free of error. An alternative to this model is the bivariate least squares (BLS) linear calibration, which takes into consideration errors in both axes [14]. Taking into consideration x-errors and y-errors, the need of a suitable linear calibration model, like BLS, becomes relevant.




Figure 2: Asymptotic trend of % overlap (ordinate) vs distance from blank signal in σ units (abscissa).

The present work focuses on the OLS and BLS models to GC-FID determination of methanol in wine. In order to ensure the LOD estimation, both OLS and BLS models were used. By application of the two calibration models, the detection limits were calculated according to the EPA 3σ and H-V method. The aim of this work was to test the

influence of sample pretreatment and chromatographic conditions on quantification of methanol in wine, referring to OLS and BLS calibration models.

An appropriate calibration curve for methanol determination in wine was set up in the range 5 – 100 mg/L, by the use of six standard reference levels. The ordinary least squares calibration equation $y_{OLS} = 0.7768 \cdot x + 0.0099$ has a good regression coefficient R^2 (0.9994), whereas bivariate least squares provides an equation $y_{BLS} = 0.7854 \cdot x + 0.0063$. Both equations provide limits of detection, with the 3σ approach, of 1.18 mg/L and 1.17 mg/L for OLS and BLS calibration models, respectively. On the other hand, 10.75 mg/L for OLS and 10.87 mg/L for BLS are the detection limits obtained through the H-V iterative calculation, with $\alpha = \beta = 0.025$, $n=6$. Table 1 shows that methanol content of Feb 2011 (33.3 mg/L) and May 2011 (33.7 mg/L) is very close to quantification limits calculated via H-V (SEE Table 2).

The investigated wine samples were injected into the GC-FID system after simple dilution with water. With the direct injection of wine samples, the dilution operation could appear to be time consuming, but it allows improvement of column stability and shelf life enhancement of the standards. With reference to OENO 19/2004 OIV methanol content limits [15], if the wine is injected directly, the internal standard concentration has to be five times higher than the 1:5 diluted sample. Moreover, the calibration range has to be five times wider as well. Standard aqueous solutions containing methanol and acetonitrile (I.S.) in the 0 – 500 mg/L range are rather stable at room temperature. If the concentration exceeds 1000 mg/L the solution stability with time is not ensured. Results from a study performed on two different concentration ranges is shown in Figure 3, where the relative response factor (RRF) is plotted against the methanol concentration. RRF is calculated by the following formula:

$$RRF = \frac{A_{MeOH}}{A_{ACN}} \cdot \frac{C_{MeOH}}{C_{ACN}}$$

where A_{MeOH} is the peak area of methanol, A_{ACN} is the peak area of the acetonitrile internal standard, and C_{MeOH} and C_{ACN} are the concentrations, in mg/L, of methanol and acetonitrile respectively.

The first evidence was that the change in the RRF was much higher in concentrated than dilute solutions. In the 5 mg/L reference solution the RRF after 5 months was 14% higher than the initial value. When a higher concentration is considered, i.e. 100 mg/L standard solution, after 5 months there is an increasing on RRF value from 0.75 to 1.01, this meaning an enhancement of about 35%. A variation in RRF values between 0.66 and 0.81 was shown by the 10000 mg/L reference solution after 5 days; increasing to 0.99 after 5 months. A more pronounced difference was encountered from 0.66 to 1 (up to 52%), matching 5 months time range, when a 15000 mg/L reference solution was taken into account. Probably, these variations were caused by evaporation of volatile compounds present in the reference standard solution and are, therefore, temperature dependent.




Figure 3: Relative response factors versus methanol concentration level. methanol concentration ranges (a) 0 – 100 mg/L; (b) 0 – 20000 mg/L.

In order to evaluate the influence of non volatile compounds in GC quantification of methanol, a comparative study between distilled wine and diluted wine by direct injection (DWDI) was carried out. Analyses were performed on samples provided from UIV (Unione Italiana Vini). Preliminary tests suggested slightly higher values for methanol in distilled samples. Determined methanol contents and UIV proficiency test data are reported in Figure 4. Except for the March 2011 test, the methanol content, after distillation, seemed to be slightly higher than that of the directly injected diluted samples. From a practical point of view, an exhaustive focus on these differences is worthy of investigation.

The influence of the distillation process sets the methanol recovery data slightly higher than those determined through DWDI. This apparent surplus could not be explained on the basis of background noise as the baselines




Figure 4: Methanol levels in mg/L of seven different UIV proficiency tests.

for both the distilled and undistilled wine were virtually the same. In addition, reproducibility calculated according to ISO 5725:5, provided by the UIV interlaboratory proficiency test, represents a 35-50% average range of the mean data (Table 1). Compared with DWDI, data results of methanol determination through distillation do not affect detected concentrations beyond 10%. This difference is compatible with the UIV reproducibility values.

Table 1: Methanol content and reproducibility provided by UIV interlaboratory proficiency test.

UIV proficiency test month	UIV proficiency test data ^a	Reproducibility ^b
Nov 2010	47.2	21.7
Dec 2010	129.1	49.8
Jan 2011	135.8	46.8
Feb 2011	33.3	15.3
Mar 2011	49.8	25.8
Apr 2011	150.7	52.8
May 2011	33.7	13.6

^a methanol content in mg/L coming from UIV interlaboratory proficiency test.

^b calculated according to ISO 5725:5, $R = 2.8 \cdot S_R$ (mg/L).

According the OENO 19/2004 OIV resolution, the maximum methanol contents in wine are 400 and 250 mg/L for red and white/rosé wines, respectively. The DWDI method provides a methanol concentration on the average of 3% less than that determined for distilled wine samples, but not more than 10%. The DWDI method is faster, and less reagents and energy consuming. So it can be used as a screening check method. When doubtful cases result and quality data assurance has to made, sample distillation can be a valid option to confirm the analytical response.

In the acetal-acetonitrile elution region, chromatograms appear very similar to each other. In the first 10 minutes, there are no differences between distilled and undistilled wine sample for normal conditions (Figure 5 a and b). However, when the temperature program and carrier flow rate are increased, the undistilled samples (Figure 5 c) show many peaks. After 6 minutes there are a number of peaks belonging to the glycol fraction of the wine. The glycol-like compounds are characterized by a high boiling point. Therefore, following a distillation process, they are removed from the samples, as shown in Figure 5 d.




Figure 5: UIV Proficiency Test of April 2011, red dry wine gas chromatograms; sample dilution 1:5. 1) 1,1-diethoxyethane, 2) methanol, 3) ethanol, 4) acetonitrile. (a) normal conditions; (b) normal conditions, distilled sample; (c) forced conditions; (d) forced conditions, distilled sample.

Resolution between methanol and the identified acetal (1,1-diethoxyethane) peak was not less than 0.9. The asymmetry factor, calculated at 10% of the maximum height for methanol, was 1.76. The calibration curve parameters by keeping six concentration levels with normal and forced temperature/flow programs are in agreement. These latter show equations $y_{OLS} = 0.7664 \cdot x + 0.0102$ (R^2 0.9999) and $y_{BLS} = 0.7628 \cdot x + 0.0124$. Method suitability quantification is supported by peaks resolution, band width and peak shape. Recovery performances were assessed by the interlaboratory proficiency test. The method was tested with two spiked commercial red wines and with the UIV proficiency test of the April 2011 sample. In the spiked samples, the mean recoveries ($n=3$) were 101.3% and 103.2%. For the UIV proficiency test of April 2011, the mean recovery was 102.9%.

Table 2: Detection and quantification limits expressed in mg/L.

Detection limits calculation approach	calibration model	Normal chromatographic condition ^a		Forced chromatographic condition ^b	
		LOD	LOQ ^c	LOD	LOQ ^c
H-V	OLS	10.8 (4.5) ^d	35.8	9.1	30.2
	BLS	10.9 (4.5) ^d	36.2	9.0	30.1
3 σ	OLS	1.2 (1.1) ^d	3.9	1.2	4.0
	BLS	1.2 (1.1) ^d	3.9	1.2	4.0

^a He 2 mL/min; oven 50°C \times 6 min; ramp 10°C/min, isotherm 210°C \times 15 min.

^b He 4 mL/min; oven 50°C \times 3 min; ramp 20°C/min, isotherm 210°C \times 7.5 min.

^c Limit of quantification is expressed as (10/3) \times LOD.

^d In parenthesis, limits of detection obtained with fifteen data points, reported in [16].

In Table 2 quantification and detection limits for both normal and forced chromatogram condition are reported. When forced GC-FID conditions were employed, the analytical method performance remained almost identical to the normal GC-FID conditions. In both cases, temperature/flow and total running time did not affect the determination of calibration parameters, LOD and LOQ.

Limits of detection calculated according to the Hubaux-Vos recursive formula are higher than those using the 3 σ approach, in both calibration models BLS and OLS. This is more pronounced when limits of quantification are considered. Comparing the calibration curve made up of fifteen data points [16], while the 3 σ LOD is substantially unvaried, if six levels are employed, the H-V detection limits are rather different. The calibration curve built up on the 5 – 100 mg/L concentration range shows a LOD approximately 2.5 times higher than that of the 0.3 – 510.6 mg/L range [16].

Experimental

Compounds and analysis: Acetonitrile (99.8%) (Sigma-Aldrich) was employed as internal standard (IS). Methanol (99%) (Sigma-Aldrich) was used for standard solution preparation. Ethanol (96% v/v) (VWRTM) was added to standard solution at 3% vol. Stock standard solution was stored at -4°C. Methanol standard solution concentration range 0 – 500 mg/L. IS concentration 76 mg/L, in both standard solution and samples. Wine samples were supplied from UIV (Unione Italiana Vini, Verona, Italy). Analysis was performed in a Trace GC Ultra instrument, equipped with a flame ionization detector (ThermoFisher Scientific, Milan, Italy). Column: retention gap 1 m \times 0.32 mm I.D. (deactivated polyethylene glycol, Varian BV, PL Middelburg, Netherlands) connected to 50 m WCOT fused silica CP-WAX 57 CB capillary column (0.32 mm I.D., film thickness 0.2 μ m, stabilized polyethylene glycol, Varian BV, PL Middelburg, Netherlands). GC conditions: carrier He 2 mL/min, FID temperature 250°C (H₂ 35 mL/min, air 350 mL/min, makeup gas N₂ 30 mL/min), injector temperature 240°C. Oven temperature 50°C (6 min isotherm), ramp program 10°C/min rate to 210°C (15 min isotherm). Injection volume 1 μ L. Split ratio 1:20. Column contamination by non volatile compounds was

avoided by using a Split Precision® Liner 5 mm × 8.0 × 105 filled with deactivate wool (Restek). H-V detection limit calculation was carried out as described by Hubaux

and Vos [13]. BLS calibration was accomplished through calculation proposed by Lisý *et al.* [17] and Riu *et al.* [14].

References

- [1] International Programme on Chemical Safety. (1997) Methanol. *Environmental Health Criteria*, World Health Organization, Geneva **196**.
- [2] Upadhyay MS, Gupta VK. (1984) Reagent system for the spectrophotometric determination of methanol in environmental and biological samples. *Analyst*, **109**, 1427–1429.
- [3] AOAC Official Method 958.04 (1998) Methanol in distilled liquors – chromotropic acid colorimetric method. *AOAC Official Method of Analysis*, Code No. 26.1.34.
- [4] AOAC Official Method 920.132 (1998) Methanol in vanilla extract – titrimetric method. *AOAC Official Method of Analysis*, Code No. 36.2.17.
- [5] AOAC Official Method 972.11 (1998) Methanol in distilled liquors – gas chromatographic method. *AOAC Official Method of Analysis*, Code No. 26.1.36.
- [6] Mizgunova UM, Zolotova GA, Dolmanova IF. (1996) Enzymatic method for the determination of ethanol and methanol with spectrophotometric detection of the rate of the process. *Analyst*, **121**, 431–433.
- [7] Sun LX, Okada T. (2000) Simultaneous determination of the concentration of methanol and relative humidity based on a single Nafion(Ag)-coated quartz crystal microbalance. *Analytica Chimica Acta*, **421**, 83–92.
- [8] Sekine Y, Suzuki M, Takeuchi T, Tamiya E, Karube I. (1993) Selective flow-injection determination of methanol in the presence of ethanol based on a multi-enzyme system with chemiluminescence detection. *Analytica Chimica Acta*, **280**, 179–184.
- [9] Van den Berg FWJ, van Osenbruggen WA, Smilde AK. (1997) Process analytical chemistry in the distillation industry using near-infrared spectroscopy. *Process Control Quality*, **9**, 51–57.
- [10] Tagliaro F, Dorizzi R, Ghielmi S, Marigo M. (1991) Direct injection high performance liquid chromatographic method with electrochemical detection for the determination of ethanol and methanol in plasma using an alcohol oxidase reactor by gas chromatography with direct on-column injection. *Journal of Chromatography*, **566**, 333–339.
- [11] Mei-Ling W, Jih-Terng W, Youk-Meng C. (2004) A rapid and accurate method for determination of methanol in alcoholic beverage by direct injection capillary gas chromatography. *Journal of Food Composition Analysis*, **17**, 187–196.
- [12] Mei-Ling W, Jih-Terng W, Youk-Meng C. (2004) Simultaneous quantification of methanol and ethanol in alcoholic beverage using a rapid gas chromatographic method coupling with dual internal standards. *Food Chemistry*, **86**, 609–615.
- [13] Hubaux A, Vos G. (1970) Decision and detection limits for linear calibration curves. *Analytical Chemistry*, **42**, 849–855.
- [14] Riu J, Rius FX. (1996) Assessing the accuracy of analytical methods using linear regression with errors in both axes. *Analytical Chemistry*, **68**, 1851–1857.
- [15] OIV Resolution OENO 19/2004. (2004) Maximum content limits of methanol in wines, http://news.reseau-concept.net/images/oiv_uk/Client/Resolution_OENO_EN_2004_19.pdf
- [16] Caruso R, Scordino M, Traulo P, Gagliano G. (2011) GC-FID determination of volatile compounds in wine: comparison between EPA 3σ approach and Hubaux-Vos calculation of detection limits, using ordinary and bivariate least-squares. *Journal of AOAC International*, in press.
- [17] Lisý JM, Cholvadová A, Kutej J. (1990) Multiple straight-line least-squares analysis with uncertainties in all variables. *Computers & Chemistry*, **14**, 189–192.

Composition and Biological Potential of Essential Oil from *Thelechitonina trilobata* Growing in South Africa

Jamie Peebles^a, Ephraim Gwebu^a, Opeoluwa Oyedeji^b, Sarah Nanyonga^c, Nokuthula Kunene^d, David Jackson^e, William Setzer^e and Adebola Oyedeji^{f*}

^aDepartment of Chemistry, Geology and Physics, Elizabeth City State University, Elizabeth City, N.C., 27909, USA

^bDepartment of Chemistry, University of Fort Hare, Private Bag X1314, Alice, 5700, South Africa

^cDepartment of Chemistry, University of Zululand, Private Bag X1001, KwaDlangezwa, 3886, South Africa

^dDepartment of Agriculture, University of Zululand, Private Bag X1001, KwaDlangezwa, 3886, South Africa

^eDepartment of Chemistry, University of Alabama in Huntsville, Huntsville, AL 35899, USA

^fDepartment of Chemistry and Chemical Technology, Walter Sisulu University, Private Bag X1, Mthatha, 5099, South Africa

aoyedeji@wsu.ac.za

Received: May 20th, 2011; Accepted: October 4th, 2011

Thelechitonina trilobata is regarded as a troublesome weed that grows to form a dense blanket over the soil preventing the growth of other crops in farmland. Although the plant is regarded as a notorious, invasive plant, its chemical composition and biological potential have not been reported. The essential oil was isolated from the fresh leaves of *T. trilobata* using hydrodistillation. α -Pinene (21.6%), α -phellendrene (21.0%), limonene (12.8%) and germacrene D (7.5%) were the major constituents of the oil. The essential oil was screened against agricultural pests. The anti-tick properties were tested on *Rhipicephalus e. evertsi* found on sheep, while repellency, fumigation, and contact toxicity tests were carried out with maize weevils. Except for the contact toxicity test, all other bioassays gave positive results.

Keywords: *Thelechitonina trilobata*, Asteraceae, essential oil, α -pinene, α -phellendrene, limonene, germacrene D, fumigant, contact toxicity, repellency.

Weeds are classified as plants that are considered by the user to be a nuisance, which interfere with the management objectives for a given area of land at a given point in time. The word weed can therefore be said to be unwanted plants in human-made settings such as garden, farmland or uncultivated land space [1,2].

Thelechitonina trilobata (L.) H. Rob. & Cuatrec. {synonyms *Complaya trilobata* (L.) Strother; *Silphium trilobatum* L.; *Wedelia trilobata* (L.) Hitchc} is a member of the Asteraceae family and considered to be a very noxious weed. It is native not only to Southern Africa but Central America as well. This plant resembles a daisy, which is why its common name is the Singapore Daisy [3,4]. *T. trilobata* is a creeping, mat-forming, perennial plant. The leaves are about 4-9 cm long and 2-5 cm wide. Flowers are yellow in color with 8-13 rays per head [3-10]. It is considered one of the World's one-hundred most alien invasive species, and causes huge problems to the surrounding vegetative life, as well as to farmers. It can, therefore, be said to be a threat to the South African

economy and environment. However, this plant is being overlooked as just a noxious weed without investigating its biological potential. Scientific research was therefore imperative to find end usage for this invading weedy plant, and this stimulated our interest in investigating its essential oil composition and biological potential. To the best of our knowledge, there is no previous report on the chemical composition and biological activities of the essential oils of *T. trilobata*. This paper is, therefore, reporting for the first time the fresh leaf volatile oil composition, its anti-tick property on sheep, and fumigative toxicity, contact toxicity and repellency properties against maize weevils.

Hydrodistillation of the leaves of *T. trilobata* gave a colorless oil with an aromatic pungent smell (yield 0.12%, w/w). Thirty-seven compounds were identified in the oil mixture accounting for 91.9% of the total oil composition. Analysis of the oil revealed a dominance of monoterpenes (65.8%), of which α -pinene, α -phellandrene, *p*-cymene, and limonene were the major components (Table 1). No oxygenated monoterpene was detected in the oil mixture. The percentage composition of sesquiterpenoids in the oil

Table 1: Chemical composition of *T. trilobata* essential oil.

Serial No	RI	Compound	Percent composition
1	941	α -Pinene	21.6
2	953	Camphene	0.5
3	975	Sabinene	0.5
4	977	β -Pinene	2.3
5	991	Myrcene	1.0
6	1005	α -Phellandrene	21.0
7	1025	<i>p</i> -Cymene	4.5
8	1029	Limonene	12.8
9	1039	(<i>Z</i>)- β -Ocimene	0.9
10	1049	(<i>E</i>)- β -Ocimene	0.7
11	1392	β -Elemene	0.3
12	1419	(<i>E</i>)-Caryophyllene	3.5
13	1453	α -Humulene	2.8
14	1481	Germacrene D	7.5
15	1497	Bicyclogermacrene	3.4
16	1511	(<i>E,E</i>)- α -Farnesene	0.2
17	1516	γ -Cadinene	0.1
18	1525	δ -Cadinene	0.2
19	1566	(<i>E</i>)-Nerolidol	0.2
20	1576	Germacrene D-4-ol	0.1
21	1578	Spathulenol	1.4
22	1581	Caryophyllene oxide	0.3
23	1609	Humulene epoxide II	0.1
24	1641	τ -Muurolol	0.1
25	1689	Shyobunol	0.2
26	2082	<i>n</i> -Octadecanol	0.1
27	2100	Heneicosane	0.1
28	2200	Docosane	0.1
29	2300	Tricosane	0.2
30	2400	Tetracosane	0.4
31	2500	Pentacosane	0.6
32	2600	Hexacosane	0.8
33	2700	Heptacosane	0.8
34	2800	Octacosane	0.9
35	2900	Nonacosane	0.8
36	3000	Triacotane	0.5
37	3100	Untriacontane	0.5
Total Identified			91.9

was 21.1% of which β -caryophyllene, α -humulene, germacrene D and bicyclogermacrene were the dominant ones. Fatty acids formed 5.7% of the oil.

The essential oil of *T. trilobata* was effective in killing *Rhipicephalus e. ervertsi* (ticks) found at the anus, udder and tail parts of sheep within one hour at concentrations of both 5 mg/mL and 10 mg/mL (Table 2). The ticks at these parts were nymphs, while those found at the lower part of the stomach toward the rear leg were adults. This means that the oil extract is more effective on the nymphs than the grown adult tick. This is noteworthy as Triatix 125, a synthetic insecticide (also used as control), was only active after 48 hours.

Table 2: Mortality of ticks (number) vs. time of exposure at different concentrations of essential oils.

Time (mins)	Concentrations and body parts where ticks found				
	5 mg/mL		10 mg/mL		
30	Stomach	Anus	Udder	Anus	Tail
30	1	7	1	1	5
45	1	7	0	1	5
60	1	0	0	1	0
75	1	0	0	0	0
90	1	0	0	0	0
120	1	0	0	0	0
720	1	0	0	0	0

Table 3: Percentage repellency of maize weevils to the fresh leaf oil.

Exposure time (h)	(0.47 μ L/cm ² = 15 μ L of the oil)	
	% repellence	No. repelled
1	14.3	12
2	61.8	15
4	62.9	16
20	80.9	19
24	66.6	17
48	42.9	15

The repellency was best at 0.47 μ L/cm² (15 μ L) of the oil (Table 3). Neither lower nor higher concentrations were effective. Furthermore, it was observed that after 20 hours the number of insects in the treated zone increased, thereby reducing the percentage repellency. It can, therefore, be concluded that the oil is most effective at 15 μ L/mL within the first 20 hours as a weevil repellent.

The result of the fumigative toxicity test is presented in Table 4. The oil extract was most effective after 4 days at low concentrations and effective on the first day at higher concentrations. This result is very promising as the oil can be used as a fumigant against maize weevils during storage. The contact toxicity was very poor and hence not reported here.

Experimental

Plant material and isolation of essential oil: Fresh leaves of *Thelechitonina trilobata* (L.) H. Rob. & Cuatrec. were collected from around the Agricultural farm of the University of Zululand in the northern part of Kwa-Zulu Natal Province, South Africa and a voucher specimen (JPKN 1) was deposited in the herbarium at the University of Zululand. The leaves (500 g) were subjected to hydrodistillation using a Clevenger-type apparatus [11,12]. The essential oil was collected 4 h after boiling, weighed and kept at 4°C till used for bioassays.

Table 4: Percentage mortality using fumigation vs. exposure time for maize weevils (*Sitophilus zeamais*) at different concentrations of leaf oil.

Conc.	Day 0		Day 2		Day 4		Day 6		Day 8		Day 12	
	%	No of dead insects	%	No of dead insects	%	No of dead insects	%	No of dead insects	%	No of dead insects	%	No of dead insects
4 μ L/L	0	0	0	0	0	0	8.3	2	41.6	10	79.2	19
8 μ L/L	0	0	0	0	0	0	41.6	10	50	12	79.2	19
16 μ L/L	0	0	20.8	5	41.6	10	79.2	19	83.3	20	87.5	21
32 μ L/L	20.8	5	20.8	5	58.3	14	79.2	19	95.8	23	100	24
40 μ L/L	20.8	5	79.2	19	100	24	100	24	100	24	100	24

% = Percentage mortality

GCMS analysis of the essential oil: The essential oil of *T. trilobata* was analyzed by GC-MS using an Agilent 6890 GC with Agilent 5973 mass selective detector [MSD, operated in the EI mode (electron energy = 70 eV), scan range = 45-400 amu, and scan rate = 3.99 scans/sec], and an Agilent ChemStation data system. The GC column was a HP-5ms fused silica capillary with a (5% phenyl)-polymethylsiloxane stationary phase, film thickness of 0.25 µm, a length of 30 m, and an internal diameter of 0.25 mm. The carrier gas was helium with a column head pressure of 48.7 kPa and a flow rate of 1.0 mL/min. Injector temperature was 200°C and detector temperature 280°C. The GC oven temperature program was used as follows: 40°C initial temperature, held for 10 min; increased at 3°C/min to 200°C; and increased at 2°C/min to 220°C. A 1%, w/v, solution of the sample in CH₂Cl₂ was prepared and 1 µL was injected using a splitless injection technique.

Identification of the oil components was based on their retention indices determined by reference to a homologous series of *n*-alkanes (C₈-C₃₀), and by comparison of their mass spectral fragmentation patterns with those reported in the literature [13-15] and stored on the MS library [NIST database (G1036A, revision D.01.00)/ChemStation data system (G1701CA, version C.00.01.080)]. The percentages of each component are reported as raw percentages based on total ion current without standardization.

Parasite identification and bioassay: The anti-tick bioassay was carried out on a pool of sheep at the University of Zululand's Dairy Unit. The ticks were identified as adult and nymph types of *Rhipicephalus e. ervertsi* by Holeise Heyne of the Parasitology Department, Veterinary Institute, Agricultural Research Council, South Africa. Sheep were caught, identified, and searched for ticks, ears and genital/anal areas being the most popular areas of attachment. Attached ticks in these particular areas were counted and recorded. Essential oil (5 mg/mL and 10 mg/mL) in a 1 mL mixture of water, Tween-20 (Polysorbate 20) and *n*-hexane for an even consistency of oil were prepared. Two drops of each concentration were applied to the area of tick attachment. Triatix 125, a synthetic insecticide, was used as control. The ticks were observed each day to note if any fell off the sheep. This experiment was conducted over a period of 3 days using 3 sheep with similar *Rhipicephalus e. ervertsi* (ticks) attachment. Average results were recorded.

Rearing of test insects: Adults of *Sitophilus zeamais* were obtained from a colony maintained by the Plant Protection Research Institute, Pretoria, South Africa. These were mass reared on whole maize grains in 5 L glass jars in a controlled chamber, at 28 ± 2°C and 56-65% RH in the Department of Chemistry, University of Zululand. Newly emerged, one-week-old insects were used in the bioassay [16]. 21-24 maize weevils were used for the study.

Repellency assay of the essential oil: The repellent effect of *T. trilobata* essential oil against *S. zeamais* was studied using a modified area preference method [17]. The test area consisted of a 9 cm Whatman No.1 filter paper cut into 2 halves. Different oil concentrations were prepared by diluting 2, 4, 8, 10 and 40 µL of the oil in 1 mL *n*-hexane and these corresponded to concentrations of 0.314, 0.472, 0.94 and 1.26 µL of oil/cm² of the filter paper, respectively. The other half was treated with 0.5 mL *n*-hexane alone and this served as a control. Both essential oil treated and *n*-hexane treated filter paper halves were air dried for 10 min to evaporate the solvent. With the aid of a clear adhesive tape, both halves were later joined together into full discs and placed in 9 cm glass Petri dishes. Twenty one-week old, unsexed adult insects were released at the centre of the rejoined filter paper disc and the Petri dish was covered. Each treatment was replicated 4 times for both *S. zeamais*. The number of insects present on the control (Nc) and treated (Nt) areas of the filter paper were recorded for 1, 2, 3, 4, 20, 24 and 48 h. 24 insects were used. Percentage repellency (PR) was calculated as follows:

$$PR = ((Nc - Nt)/(Nc + Nt)) \times 100$$

The weevils were exposed to a filter paper with half the filter paper being oil and the other half being acetone (control) and they were observed to see whether the concentrated oil repelled them. Concentration was calculated by dividing the oil quantity (µL) by the surface area of half the filter paper (31.81 cm²) [17].

Fumigant assay of the essential oil: The fumigation chambers consisted of 500 mL glass jars with screw-on lids. For the bioassay, solutions of 0, 4, 8, 16, 32 and 40 µL of the oil were each diluted with 1 mL *n*-hexane to correspond to concentrations of 0 (control), 4, 8, 16, 32 and 40 µL/mL. One mL of each concentration was then separately applied to 7 mm discs of Whatman No.1 filter paper, air-dried for 10 min and placed at the bottom of the jars. Twenty one-week old adult insects were placed on muslin cloths 21 × 29 mm each with 40 g whole maize grains. The cloths were tied closed with rubber bands and hung at the centre of the jars, which were then sealed with air-tight lids. There were 4 replicates for each concentration. Fumigation was carried out for 24 h, after which the insects were transferred from the fumigation chambers onto clean maize, and mortality was checked for a further 12 days [17].

Contact toxicity of the essential oil: The contact effect of the essential oil of *T. trilobata* on the adults of *S. zeamais* was investigated [16, 17]. Maize grains were treated with concentrations of 0, 25, 50, 100, 200 and 300 µL of essential oil in 1 mL *n*-hexane. The different concentrations of the oil were mixed with 40 g of maize grain respectively. These were thoroughly stirred to allow for homogeneity of the oil on the treated grains. Treated

samples were air dried for 1 h in order to remove the solvent. The grains were thereafter infested with 21-week old *S. zeamais* adults per jar and each jar was covered with a cotton mesh held in place by cover rims. There were 4 replicates per treatment. Dead insects in the jar were counted daily for 6 days.

Acknowledgments – The project described was supported in part by Award Number T37MD001810 from the National Institute on Minority Health and Health Disparities. The authors are also grateful to NRF South Africa, the Directorate of Research, Walter Sisulu University, Govan Mbeki Research and Development Centre, University of Fort Hare and Research Committee of the University of Zululand for research funding.

References

- [1] Janick J. (1979) *Horticultural Science* (3rd ed.). San Francisco, W.H. Freeman. p. 308.
- [2] Whitson T. (Ed.) (1992) *Weeds of the West*. Western Society of Weed Science. p. ix.
- [3] Cuatrecasas J, Robinson H. (1992) *Thelechitonia cuatrecasas*, an older name for *Complaya strother* (Ecliptinae-Heliantheae-Asteraceae). *Phytologia*, **72**, 141-143
- [4] PIER (Pacific Island Ecosystem at Risk) (2003) *Wedelia trilobata*, http://www.hear.org/pier/species/wedelia_tribota.htm [Accessed 13 May 2011]
- [5] Santos I. (1998) *Wedelia trilobata* on Pohnpei. *Aliens*, **7**, 3.
- [6] Thaman RR. (1999) *Wedelia trilobata*: Daisy invader of the Pacific Island. IAS Technical Report, 99/2. Institute of Applied Science, University of the South Pacific, Suva, Fiji Island.
- [7] Henderson L. (2001) *Alien Weeds and Invasive Plants*. Plant Protection Institute Handbook No. 12. Plant Protection Research Institute, Agriculture Research Council, South Africa, p 58.
- [8] Pienaar K. (1984) *The South African what flower is that? (A-Z of Garden flowers in South Africa)*. 2nd Impression, Struik, Cape Town, p 347.
- [9] Sheat WG, Schofield G. (1995) *Complete Gardening in South Africa*. Struik, South Africa, p 285.
- [10] www.sanparks.org/parks/Kruger/conservation [Accessed 16 May 2011]
- [11] Oyedele AO, Afolayan AJ, Hutching A. (2009) Compositional variation of the essential oils of *Artemisia afra* Jacq. from three provinces in South Africa – A case study of its safety. *Natural Product Communications*, **4**, 849-852.
- [12] *British Pharmacopoeia* (1980) **11**, P. A. 109, HMSO.
- [13] Adams RP. (2007) *Identification of Essential Oil Components by Gas Chromatography / Mass Spectrometry*, 4th Ed. Allured Publishing, Carol Stream, Illinois.
- [14] Joulain D, Koenig WA. (1998) *The Atlas of Spectra Data of Sesquiterpene Hydrocarbons*. E.B.-Verlag, Hamburg.
- [15] ESO 2000. (1999) *The Complete Database of Essential Oils*. B.A.C.I.S., The Netherlands.
- [16] Odeyemi OO, Masika P, Afolayan AJ. (2008) Insecticidal activities of essential oil from the leaves of *Mentha longifolia* L. subsp. *capensis* against *Sitophilus zeamais* (Motschulsky) (Coleoptera: Curculionidae). *African Entomology*, **16**, 220-225.
- [17] Tapondjou AL., Adle C, Bouda H, Fontem DA. (2002) Efficacy of powder and essential oil from *Chenopodium ambrosioides* leaves as post-harvest grain protectants against six-stored beetles. *Journal of Stored Products Research*, **38**, 395-402.

Chemical Composition and Antibacterial Activity of Essential oil from *Salvia muckerjeei*

Lalit Mohan^a, Anuradha Negi^a, Anand B. Melkani^{a*} and Vasu Dev^b

^aDepartment of Chemistry, Kumaun University, Nainital 263002, Uttarakhand, India

^bDepartment of Chemistry, California State Polytechnic University, Pomona, CA 91768, USA

melkaniab@rediffmail.com

Received: August 22nd, 2011; Accepted: October 4th, 2011

The composition of steam volatile oil from aerial parts of *Salvia muckerjeei* Bennet & Raizada (Lamiaceae) was analyzed by capillary GC and GC-MS. The oil was rich in sesquiterpene hydrocarbons (67.3%). Among 71 identified constituents representing 91.7% of the oil, β -caryophyllene (28.7%), γ -muurolene (15.5%) and dehydro-aromadendrane (9.5%), were the principal constituents. The oil was tested against ten bacterial strains and was active against *Enterococcus faecalis*, *Erwinia chrysanthemi* and *Agrobacterium tumefaciens*.

Keywords: *Salvia muckerjeei*, Lamiaceae, essential oil composition, β -caryophyllene, γ -muurolene, antibacterial activity.

The genus *Salvia* Linn. contains about 700 species widely distributed in the temperate and warm regions of both the hemispheres. Seven species of *Salvia* have been reported to grow in the central Himalayan region of India [1,2]. Two species have been investigated thoroughly for their steam volatile components [3,4]. The roots of *Salvia muckerjeei* are reported to cure cold and cough, the leaves are applied to wounds and the seeds are used for dysentery and colic [5].

The literature contains several reports on the essential oil composition of *Salvia* species growing worldwide [6] but none for *S. muckerjeei*. The present communication describes a detailed analysis of the oil and its antibacterial activity.

The percentage oil yield from the plant was 0.34% by weight. The GC analysis revealed more than eighty components. Seventy one identified constituents represented 91.7% of the oil, several in trace amounts. The compounds were identified by comparison of their Retention Indices and mass spectra with the literature [7]. The oil was rich in sesquiterpene hydrocarbons (67.3%) including β -caryophyllene (28.7%), γ -muurolene (15.5%), dehydro-aromadendrane (9.5%) and α -guainene (5.4%) in addition to monoterpene hydrocarbons (4.5%) and oxygenated sesquiterpenoids (7.8%) [Table 1].

Since *S. muckerjeei* produced a relatively satisfactory yield of sesquiterpene rich oil, in contrast to the generalization [8] that only the oil poor (<0.2%) Lamiaceae taxa are rich in sesquiterpene hydrocarbons. A comparison between the essential oil composition of *S. muckerjeei* with those of other *Salvia* species indicates it is closely related to those

from *S. aethiopis* (β -caryophyllene 23.3%, α -copaene 23.3%) [9], *S. palanestina* (β -caryophyllene 36.4%) [10], *S. canariensis* (β -caryophyllene 30.2%, α -pinene 9.5%) [11], *S. hydrangea* (β -caryophyllene 33.4%, caryophyllene oxide 25.4%) [12] and *S. bracteata* β -caryophyllene 10.7-41.6%, γ -muurolene 1.8 - 9.9%) [13]. In terms of the essential oil composition, *S. muckerjeei* can also be compared with *S. angulata* from Venezuela which is also rich in sesquiterpene hydrocarbons: β -caryophyllene (27.7%), germacrene D (19.5%), bicyclogermacrene D (18.3%) [14]. However, germacrene D is completely absent in the essential oil from *S. muckerjeei*.

Antibacterial testing of the essential oil showed a maximum zone of inhibition (20mm) against the Gram-positive bacteria, *Enterococcus faecalis* followed by the Gram-negative, *E. chrysanthemi* (18 mm), *A. tumefaciens* (17 mm), *E. coli* (15 mm) and the lowest (10 mm) against *X. phaseoli*. The zone of inhibition values are summarized in Table 2. All bacterial strains were further tested at different concentrations to determine the minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) values.

Erwinia chrysanthemi showed the lowest MIC at a concentration of 31.25 μ L/mL, while the MIC values against *E. coli*, *A. tumefaciens* and *B. subtilis* were at 62.5 μ L/mL and the minimum bactericidal concentration of 125.0 μ L/mL was against *A. tumefaciens*. The results are summarized in Table 3. The *S. palaestina* Oil, with β -caryophyllene as a principal constituent, also showed better activity against the Gram-positive bacteria *S. aureus*, than Gram-negative bacteria (*E. coli*, *P. aeruginosa*.) [10]. The oil from *S. canariensis*, with β -caryophyllene

Table 1: Chemical composition of the essential oil from aerial parts of *S. mukerjeei*.

Compound	Percentage in the oil	RI observed
α -Pinene	0.4	933
Camphene	0.4	953
Benzaldehyde	0.1	967
Sabinene	1.0	978
β -Pinene	1.7	979
δ -2-Carene	0.1	1000
Isosylvestrene	0.1	1005
α -Terpinene	0.1	1017
<i>o</i> -Cymene	0.2	1026
Limonene	0.1	1029
β -Phellandrene	5.0	1031
1,8-Cineole	0.1	1034
(<i>Z</i>)- β -Ocimene	0.3	1036
(<i>E</i>)- β -Ocimene	1.0	1054
γ -Terpinene	0.3	1062
<i>p</i> -Mentha-3,8-diene	0.1	1071
<i>p</i> -Mentha-2,4(8)-diene	0.1	1089
Terpinolene	0.5	1100
<i>n</i> -Nonanal	0.5	1105
Dehydrosabinaketon	0.2	1128
Camphor	0.5	1147
Pinocarvone	0.1	1165
Karahanaenone	0.9	1168
Terpinen-4-ol	0.3	1179
α -Terpineol	0.2	1192
γ -Terpineol	0.1	1197
<i>p</i> -Cymen-9-ol	0.3	1203
Linalyl formate	0.1	1207
Carvone	0.1	1243
Carvacrol methyl ether	0.1	1245
Thymol	1.2	1293
δ -Elemene	0.2	1340
Thymol acetate	trace	1358
α -Copaene	0.7	1378
β -Bourbonene	0.5	1380
(<i>E</i>)- β -Damascenone	0.2	1392
β -Elemene	0.3	1394
Longifolene	trace	1408
β -Doprezianene	0.1	1418
β -Caryophyllene	28.7	1422
γ -elemene	0.1	1432
α -Guaiane	5.4	1447
<i>allo</i> -Aromadendrene	0.2	1452
Dehydro-aromadendrane	9.5	1457
γ -Gurjunene	0.1	1480
γ -Murolene	15.5	1484
γ -Himachalene	0.2	1489
α -Amorphene	0.2	1491
δ -Selinene	0.1	1494
<i>cis</i> - β -Guaiane	0.3	1497
<i>trans</i> -Murola-4-(14),5-diene	0.2	1498
α -Murolene	0.2	1500
(<i>Z</i>)- α -Bisabolene	0.2	1507
δ -Amorphene	0.7	1510
(<i>Z</i>)- γ -Bisabolene	0.1	1514
γ -Cadinene	1.1	1517
δ -Cadinene	1.5	1527
Zonarene	0.1	1529
<i>trans</i> -Cadina-1(2),4-diene	0.1	1530
<i>cis</i> -Calamenene	1.1	1539
α -Cadinene	0.3	1542
Europelargone A	0.2	1547
Germacrene B	0.3	1557
Globulol	0.2	1561
Spathulenol	3.9	1587
Guaiol	0.1	1596
<i>trans</i> -Isolongifolone	0.9	1613
Eremoligenol	1.5	1625
Hinesol	0.2	1634
β -Atlantone	0.1	1636
Guaia-3,10(14)-die-11-ol	0.2	1687
Total identified	91.7 %	

trace= less than 0.1%

Table 2: Antibacterial screening of *S. mukerjeei* (1000 μ L/mL).

Microorganism (Bacteria)	Zone of Inhibition (mm)		
	Oil	Streptomycin	Erythromycin
<i>Erwinia chrysanthemi</i>	18	12	12
<i>Escherichia coli</i>	15	18	18
<i>Agrobacterium tumefaciens</i>	17	30	28
<i>Xanthomonas phaseoli</i>	10	25	12
<i>Bacillus subtilis</i>	15	22	13
<i>Staphylococcus aureus</i>	15	18	10
<i>Klebsiella pneumoniae</i>	14	22	25
<i>Enterococcus faecalis</i>	20	25	12
<i>Salmonella enterica enterica</i>	14	16	12
<i>Pasteurella multocida</i>	13	23	18

Table 3: MIC/MBC values of *S. mukerjeei* oil.

Microorganism (Bacteria)	Concentration (μ L/mL)				
	500.0	250.0	125.0	62.5	31.25
<i>E. chrysanthemi</i>	15*	11	10	8	6 \blacktriangle
<i>E. coli</i>	13	9*	7	7 \blacktriangle	-
<i>A. tumefaciens</i>	14	11	9*	7 \blacktriangle	-
<i>X. phaseoli</i>	7	7	-	-	-
<i>B. subtilis</i>	9	8	7	7 \blacktriangle	-
<i>S. aureus</i>	12	10	7	-	-
<i>K. pneumoniae</i>	11	7 \blacktriangle	-	-	-
<i>E. faecalis</i>	17	12*	9	9 \blacktriangle	-
<i>S. enterica enterica</i>	12	9	7	-	-
<i>P. multocida</i>	10	7 \blacktriangle	-	-	-

*MBC, \blacktriangle MIC

and viridiflorol as the major components, had limited activity against the Gram-positive bacteria *S. aureus* and *S. epidermis*, and were inactive against the Gram-negative bacteria (*E. coli*, *P. aeruginosa*, *Salmonella spp.* and *Protes mirabilis*) [11]. The essential oil of *S. mukerjeei* showed significant activity against all Gram-positive as well as Gram-negative bacteria, thus indicating the importance of this plant for the treatment of infectious diseases.

Experimental

Plant material: The plant material was collected from Lamgara, district Almora, India at an elevation of 1,900 m in March, when the plant was flowering. The aerial part of the plant was used for the extraction of oil. The plant was identified by the Botany Department, Kumaun University, Nainital and its identity was further confirmed by the Botanical Survey of India, Dehradun, where a herbarium specimen was deposited (Herbarium Voucher No. [BSI (NC) Acc No.112203 BSD]).

Extraction and isolation of oil: The oil was obtained by the steam distillation of fresh plant material (2.0 kg) using a copper still fitted with spiral glass condensers. The aqueous distillate was saturated with NaCl and the oil was extracted with hexane. The organic layer was separated and dried over anhydrous sodium sulphate. The solvent was removed using rotovap at reduced pressure and at 30°C.

Gas Chromatography: The GC analysis was performed using a Hewlett Packard 6980 gas chromatograph fitted with an HP-5MS capillary column (30m x 0.25mm, film thickness 0.25 μ m), and a flame ionization detector. The oven was programmed at 50°C for 10 min and then raised

to 240°C (3°C/min) where it was maintained for 5 min. The composition data were taken from area percent data without the use of correction factors and rounded off to the first decimal place. Co-injection of the oil and n-alkanes (C₈-C₂₃) allowed calculation of the Retention Indices [7].

Gas Chromatography-Mass Spectrometry: GC-MS analysis was carried out using an AGILENT 5973 Network Mass Selective Detector interfaced with an AGILENT 6856 GC system fitted with an HP-5MS capillary column (30m X 0.25mm, film thickness 0.25 µm) at 50°C for 10 min then raised by 3°C/min to 230°C where it was maintained for final 10 min. The mass spectra were acquired at 70 eV, at 2.41 scan/second and a mass range of 41-350. Helium gas was used as a carrier gas in both the analyses.

Antibacterial assay

Microbial culture: The test organisms were obtained from the Institute of Microbial Technology, Chandigarh and the Botany Department, Kumaun University, Nainital, India as laboratory isolates of the pure culture of Gram positive *Bacillus subtilis* (MTCC 121), *Staphylococcus aureus* (MTCC 737), *Enterococcus faecalis* (MTCC 439), and Gram negative *Agrobacterium tumefaciens* (MTCC 609), *Erwinia chrysanthemi* (KUMSCC 328), *Xanthomonas phaseoli* (KUMSCC 327) *Escherichia coli* (MTCC 443), *Salmonella enterica enterica* (MTCC 3223), *Klebsiella pneumoniae* (MTCC 109) and *Pasteurella multocida* (MTCC1148). The isolates were separately cultured on nutrient agar plates for 24 hours. Twenty mL of the medium was poured into a sterile Petri plate (80 mm diameter) and allowed to solidify. A colony of each test organism was sub-cultured on ten mL nutrient broth and incubated at 37° ±1°C for 8 hours. One mL of the broth culture was then used to flood the agar plates.

Antibacterial testing: The preliminary screening of antibacterial activity was conducted at 1000 µL/mL concentration by using disc-diffusion method. [15] Streptomycin 30 µg/disc (Himedia) and Erythromycin 15 µg/disc (Himedia) were used as positive control while hexane was used as a negative control on all the bacteria to compare the zones of inhibition with that of the oil sample. The MIC values were determined at five different concentration of the oil (500, 250, 125, 62.5, and 31.25 µL/mL) and each set of experiment was done in triplicate.

Nutrient agar with a pH value of 7.3±0.2 was autoclaved at 121°C for 30 min at 15 psi. Whatman filter paper no. 1 was used to prepare discs of 5 mm diameter. Micro-organism were streaked in a radial pattern on the agar plate [16], rotating the plate approximately 60°C each time to ensure an even distribution of inoculum. Once the plates were inoculated with the micro-organism, four filter discs of 5 mm diameter were applied in a clockwise pattern, one containing plant oil (15 µL), another one of solvent (negative control, *n*-hexane) and two discs of standard antibiotics (positive controls). The plates were inverted and placed in an incubator (37°± 1°C). After 24 hours of incubation, each plate was examined for growth and inhibition. The diameter of the complete inhibition zone was measured to the nearest whole millimeter.

Acknowledgements - We are thankful to Dr. Y.P.S. Pangtey, Botany Department, Kumaun University, Nainital and Botanical Survey of India, Dehradun, India for the identification of plant herbaria specimen. Lalit Mohan is thankful to Uttarakhand Council for Science & Technology, Dehradun, India for financial assistance through a project grant(No.UCS&T/ R&D/CHEM-10/07-08/1835/1).

References

- [1] Strachey R. (1974) *Catalogue of the plant of Kumaon and the portions of Garhwal and Tibet*, reprint ed., Periodical Experts Book Agency, New Delhi, India, 140.
- [2] Gupta RK. (1968) *Flora of Nainitalensis*, Navyug Traders. New Delhi, India.
- [3] Negi A, Javed MS, Melkani AB, Dev V, Beauchamp PS. (2007) Steam volatile terpenoids from *Salvia leucantha*. *Journal of Essential Oil Research*, **19**, 401-403.
- [4] Melkani AB, Negi A, Sati SC, Khulbe K, Dev V. (2010) Terpenoid composition and antimicrobial activity of the essential oil from *Salvia nubicola* Wall ex Sweet. *Journal of Essential Oil Research*, **22**, 575-577.
- [5] Raizada MB, Saxena HO. (1978) *Flora of Mussoorie V-I*, p. 593-595.
- [6] For e.g., (a) Formisano C, Senatore F, Arnold NA, Piozzi F, Rosselli S. (2007) GC and GC/MS analysis of the essential oil of *Salvia hierosolymitana* Boiss. growing wild in Lebanon, *Natural Product Communications*, **2**, 181-184; (b) Rustaiyan A, Masoudi S, Tabatabaei-Anaraki M. (2007) Terpenoids from Iranian *Salvia* species, *Natural Product Communications*, **2**, 1031-1042; (c) Guy P, Kamatou GPP, van Zyl RL, Davids H, van Vuuren SF, Viljoen AM (2008) Synergistic and antagonistic interactions of essential oils on the biological activities of the solvent extracts from three *Salvia* species, *Natural Product Communications*, **3**, 1111-1115; (d) Schmiderer C, Grassi P, Novak J, Franz C. (2008) Diversity of essential oil glands of Spanish Sage (*Salvia lavandulifolia* Vahl, Lamiaceae), *Natural Product Communications*, **3**, 1155-1160; (e) Öztürk M, Tel G, Duru ME, Harmandar M, Gülaçtı Topçu G. (2009) The effect of temperature on the essential oil components of *Salvia potentillifolia* obtained by various methods, *Natural Product Communications*, **4**, 1017-1020; (f) Rojas LB, Visual T, Morillo M, de Rojas YC, Arbola JC, Usubillaga A. (2010) The volatile constituents of *Salvia leucantha*. *Natural Product Communications*, **5**, 937-938; (g) Tabanca N, Demirci B, Turner JL, Pounders C, Demirci F, Başer KHC, Wedge DE. (2010) Microdistillation and analysis of volatiles from eight ornamental *Salvia* Taxa. *Natural Product Communications*, **5**, 1421-1426; (h) Canzoneri M, Bruno M, Rosselli S, Russo A, Cardile V, Formisano C. (2011) Chemical composition and biological activity of *Salvia verbenaca* essential oil. *Natural Product Communications*, **6**, 1023-1026.

- [6] Adams RP. (2007) *Identification of essential oil component by Gas Chromatography /Quadrupole Mass Spectrometry*, 4th Ed., Allured Publishing Co., Carol stream, Illinois, USA.
- [7] Lawrence BM. (1992) Chemical composition of Labiatae oils and their exploitation. In: *Advances in Labiatae Science*. Edts. R.M Harley and T. Reynolds, Royal Botanical Gardens, Kew. UK.
- [8] Chalchat JC, Gorunovic MS, Petrovic SD, Maksimovic ZA. (2001) Chemical composition of two wild Species of the genus *Salvia* L. from Yugoslavia: *Salvia aethiopis* and *Salvia Verticillata*. *Journal of Essential Oil Research*, **13**, 416-418.
- [9] Sennatore F, Formisanno C, Arnold NA, Piozzi F. (2005) Essential Oil from *Salvia* sp. (Lamiaceae). III. Composition and antimicrobial activity of the essential oil of *Salvia palanestina* Benth. growing in Lebanon. *Journal of Essential Oil Research*, **17**, 419-421.
- [10] Garcia Vallejo MC, Moujir L, Burillo J, Guerra LL, Gonzalez M, Penate RD, Andres LS, Luis JG, Blanco FL, de Galarreta CMR. (2006) Chemical composition and biological activities of the essential oil of *Salvia canariensis*. *Flavour and Fragrance Journal*, **21**, 72-76.
- [11] Barazandeh MM. (2004) Volatile constituents of the oil of *Salvia hydrangea* DC ex Benth. from Iran. *Journal of Essential Oil Research*, **16**, 20-21.
- [12] Sefidkon F, Hooshidary F, Jamzad Z. (2007) Chemical variation in the essential oil of *Salvia bracteata* Banks & Soland from Iran. *Journal Essential Oil Bearing Plants*, **10**, 265-272.
- [13] Colmenares NG, Eymargarcia M, Alfredo U. (2006) Volatile constituents from the aerial parts of *Salvia angulata* H.B.K of Venezuela. *Journal of Essential Oil Research*, **18**, 486-488.
- [14] Bauer AW, Kirby WMM, Sherris JC, Turek M. (1966) Antibiotic susceptibility testing by standardized single disc method. *American Journal of Clinical Pathology*, **45**, 493-496.
- [15] Afolayan AJ. (2003) Extracts from the shoots of *Aretotis artotoides* inhibit the growth of bacteria and fungi. *Pharmaceutical Biology*, **41**, 22-25.

Revealing Indigenous Indonesian Traditional Medicine: Anti-infective Agents

Ari S. Nugraha^{a,b,*} and Paul A. Keller^a

^a*School of Chemistry, Faculty of Science, University of Wollongong, Wollongong, New South Wales, Australia 2522*

^b*Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Jember, Jember, Indonesia 68121*

asn559@uowmail.edu.au

Received: May 29th, 2011; Accepted: August 25th, 2011

Indonesia is rich in medicinal plants which the population has used traditionally from generation to generation for curing diseases. Our interest in the treatment of infectious diseases has lead to the investigation of traditional Indonesian treatments. In this review, we present a comprehensive review of ethnopharmacologically directed screening in Indonesian medicinal plants to search for new anti-viral, anti-malarial, anti-bacterial and anti-fungal agents. Some potent drug leads have been isolated from Indonesian medicinal plants. Further research is still required for the lead development as well as the search for new bioactive compounds from the enormous medicinal plant resources.

Keywords: Indonesian medicinal plants, anti-viral, anti-malaria, anti-bacteria, anti-fungi.

Indonesia is an archipelago consisting of approximately 17,508 islands and is covered by tropical rain forest, seasonal forest, mountain vegetation, subalpine shrub vegetation, swamp and coastal vegetation. With its reflective mixture of Asian and Australian native species, Indonesia is said to possess the second largest biodiversity in the world, with around 40,000 endemic plant species including 6,000 medicinal plants [1-5].

Medicinal plants have accompanied the development of indigenous Indonesian traditional treatments which is a combination of physical and spiritual aspects to form a holistic medication. This is heavily influenced by the Indian Ayurveda, ever since the early civilization in Indonesia when Hinduism spread from India to Asia [6,7]. Conversely, the incorporation of indigenous medicinal plants not found in India enhanced the development of local Indonesian traditional medication [6] which has been further enriched by influences from Chinese and Arabian traders [8]. The long indigenous history and the variety of geographical conditions have created a variety of unique Indonesian cultures, of which only a few have recorded their traditional medication; these include the indigenous people of Sumatra, Java and Bali [9-13]. Much of this knowledge was not recorded and was verbally passed from generation to generation, which is common in tribes living in remote areas [14]. The knowledge is commonly practiced which leads to most of the population still relying upon medicinal plants [15,16].

The settlement of Europeans in Indonesia in the early 17th century intervened with the local medication [17]. The lack of knowledge of European physicians of the unique tropical diseases and the limited western medication forced the Western scientists to explore the Indonesian medicinal plants [17]. They subsequently used and published indigenous Indonesian traditional herbal medicine treatments showing them to be understandable and legitimate [8,17]. Findings were published in notable books, including “De medicina Indorum” by Bontius in 1642 [18], “the Ambonese herbal” by Rumphius in 1741 [19], “Materia Indica” by van der Burg in 1885 [17], “De nuttige planten van Nederlandsch Indie” by Heyne in 1927 [20] and “select Indonesian medicinal plants” by Steenis-Kruseman in 1953 [21]. The outcomes were also reported in the Medical Journal of the Dutch East-Indies (1894-1925) [17].

Common infectious diseases occurring in traditional circumstances were able to be correlated with the then modern clinical pathology [19]. This made it easier for investigators, when reading ancient texts and listening to local informers (healers), to re-collect the knowledge through expedition which now helps scientists to perform narrowed pharmacochemical screening in finding new anti-infectious drug leads [22-26]. Since the 1970's, with the use of laboratory based experiments and the development of analytical technology, many bioactive compounds from Indonesian medicinal plants have been revealed. These include anti-viral, anti-malarial, anti-bacterial and anti-fungal agents. This review examines

these anti-infective natural products and lists plant extracts that require further analysis.

Anti-Viral Agents: Traditional knowledge was clearly unable to describe a viral pathology, however ethnopharmacological studies show the Indigenous people could diagnose viral infections and could cure them by using specific plant preparations [27]. Common viral infections known in Indonesian traditional circumstances include human herpes and rhino virus [12,28]. Later diseases such as poliovirus, human immunodeficiency virus, and avian myeloblastosis virus have resulted in extensive screening of Indonesian medicinal plants for activities against these viruses. This section will consider developments in targeting herpes simplex virus (HSV), poliovirus, rhinovirus, human immunodeficiency virus (HIV) and avian myeloblastosis virus-reverse transcriptase (AMV-RT).

Table 1: Extracts of selected Indonesian medicinal plants which have potential anti-HSV-1 and poliovirus activity [33, 34].

Plant	Part of plant	Virus type	
		HSV-1 EC ₅₀ , µg/mL	Poliovirus IC ₅₀ , µg/mL
Guttiferae			
<i>Garcinia griffithii</i> T. A.	-	781	600
<i>Garcinia mangostana</i> L.	LF	40*	
Melastomataceae			
<i>Melastoma malabathricum</i> L.	LF	192	111
Loranthaceae			
<i>Elytranthe globosa</i> B.	LF	336	217
<i>Elytranthe maingayi</i> V.T.	LF	233	41
<i>Elytranthe tubaeflora</i> R.	LF	176	56
<i>Scurrula ferruginea</i> D.	LF	i	62
Meliaceae			
<i>Toona sureni</i> (Bl.) Merr	LF	37*	
Piperaceae			
<i>Piper aduncum</i> L.	FL	344	105
<i>Piper aduncum</i> L.	LF	720	105
Punicaceae			
<i>Punica granatum</i> L.	PR	64*	
Sapindanaceae			
<i>Filicium decipiens</i> T.	SB	68*	
<i>Nephelium lappaceum</i> L.	PR	70*	
Simaroubaceae			
<i>Eurycoma longifolia</i> Jack.	ST	62*	
Verbanaceae			
<i>Vitex pubescens</i> V.	BK	i	i

LF: Leaf; SB: Stem bark; FL: Flower; PR: Pericarp; ST: Stem; HSV-1: Herpes simplex virus type-1; i: inactive. All the samples are methanol extracts. EC₅₀: 50% effective concentration *Expressed as IC₅₀: concentration that inhibits 50%.

Anti-HSV and anti-poliovirus: Selected Indonesian medicinal plant extracts have been tested against HSV-1 and poliovirus (Table 1) [29]. Quassionoid compounds might be responsible for the anti-HSV-1 activity although a drawback to further development is their cytotoxicity [30]. Another study proposed the essential oil and tannin constituents were responsible for the anti-HSV-1 activity [31, 32]. This was based on data suggesting plants that do not contain essential oils or tannins, e.g. *Garcinia* sp, have lower activity against HSV-1 than high essential oil containing plants, e.g. Sapindanaceae plants.

Anti-Rhinovirus: Ginger rhizome is commonly prepared into a traditional hot drink for common cold relief which

can be correlated with rhinovirus infection. Many rhino anti-virals from natural products have been found from higher plants including lipophilic flavonoid type molecules (i.e flavan **1** and 4',6-dichloroflavan **2**) [35]. An investigation into the Indonesian ginger rhizome resulted in the isolation of four lipophilic sesquiterpenes, ar-curcumen **3**, β -sesquiphellandrene **4**, α -zingiberene **5**, β -bisabolene **6** (Figure 1). These compounds possessed rhinovirus IB inhibitor activity at an ED₅₀ of 20.4, 0.9, 1.90, 14.3 µg/10 mL, respectively. The most active is the β -sesquiphellandrene **4** with IC₅₀ of 0.44 µM. A structure analysis would suggest that the sesquiterpene and flavan compounds are equal in size and polarity [36]. However, a minor change in structure results in a significant difference in bioactivity and further studies are therefore required to determine the target of the flavan and the sesquiterpenes molecules.




Figure 1: Rhino anti-viral sesquiterpenes (**3-6**) isolated from *Zingiber officinale* [36].

Anti-HIV agents: Although HIV is an imported disease to Indonesia, it is highly prevalent. While there are no specific traditional medicines to treat HIV, traditional anti-viral concoctions may lead to new potential anti-HIV leads. Therefore at least twenty one Indonesian medicinal plants were screened against HIV-1 protease and HIV-1 replication (Table 2) [37-39].

The methanol extracts of some medicinal plants are more active against HIV-1 protease while the more polar water extracts are more active against HIV-1 replication. There are four methanol extracts of *Terminalia belerica* Roxb, *Swietenia mahagoni* L., *Woodfordia floribunda* Salisb. and *Garcinia mangostana* L. which may contain potential anti-HIV-1 protease inhibitors with IC₅₀ values of 50, 40, 40, 50 µg/mL, respectively [37]. However, there was no further exploration of these plants except *Garcinia mangostana* L. This study isolated mangostin **7** and γ -mangostin **8** (Figure 2) which showed non-competitive inhibition against HIV-1 protease with IC₅₀ values of 5.12 and 4.81 mM, respectively [40].




Figure 2: Mangostins isolated from *Garcinia mangostana* [40].


Table 2: Extracts of selected Indonesian medicinal plants which show anti-HIV-1 protease and replicase activity [37,38].

Plant	Part of plant	HIV-1 protease IC ₅₀ , µg/mL		HIV-1 rep. IC ₅₀ , µg/mL
		MeOH ext.	H ₂ O ext.	
Acanthaceae				
<i>Andrographis paniculata</i> Nees.	LF	500	500	≥170*
Apiaceae				
<i>Foeniculum vulgare</i> Mill	SD	100	>500	
Apocynaceae				
<i>Parameria laevigata</i> Moldenke.	BK	100	>500	
Clusiaceae				
<i>Garcinia mangostana</i> L.	PL	50	100	
Combretaceae				
<i>Terminalia belerica</i> Roxb.	FR	50	220	
Compositae				
<i>Elephantopus scaber</i> L.	WP	500	>500	
Elaeocarpaceae				
<i>Elaeocarpus grandiflorus</i> Smith	FL	100	100	
Fabaceae				
<i>Caesalpinia sappan</i> L.	BK	280	320	
<i>Sindora sumatrana</i> Miq.	FR	260	360	≥134*
Hypoxidaceae				
<i>Curculigo orchoides</i> Gaertn.	FR	400	400	
Lauraceae				
<i>Cinnamomum sintok</i> Bl.	BK	220	320	
Loganiaceae				
<i>Strychnos nux-vomica</i> L.	BK	500	>500	
Loranthaceae				
<i>Loranthus parasiticus</i> (L.) Merr	ST	100	260	≥79.4*
Lythraceae				
<i>Woodfordia floribunda</i> Salisb.	FL	50	50	
Malvaceae				
<i>Helicteres isora</i> L.	FL	380	>500	≥65*
Meliaceae				
<i>Swietenia mahagoni</i> L.	BK	40	100	≥28.4**
Parmeliaceae				
<i>Usnea misaminensis</i> Vain.	WP	220	>500	
Poaceae				
<i>Andropogon zizanioides</i> (L.) Urban	RT	500	>500	
Solanaceae				
<i>Physalis angulata</i> L.	AP	340	>500	
Zingiberaceae				
<i>Curcuma aeruginosa</i> Roxb.	RZ	500	>500	≥323*
<i>Curcuma xanthorrhiza</i> Roxb.	RZ	300	>500	

RT:Root; SD:Seed; FL:Flower; LF: Leaf; AP:Aerial part; BK: Bark; WP:Whole plant; ST: Stem; HSV-1: Herpes simplex virus 1. * H₂O extract. ** MeOH extract. IC₅₀: concentration that inhibits 50%

Anti-AMV-RT: The fruit of the Indonesian medicinal plant *Helicteres isora* L. was screened for potential anti-AMV-RT activity. From these four secondary metabolites were reported: rosmarinic acid **9**, 4'-O-D-glucopyranosyl rosmarinic acid **10**, 4,4'-O-di-β-D-glucopyranosyl rosmarinic acid **11** and 4'-O-D-glucopyranosil isorinic acid **12**, which exhibited xanthine oxidase inhibition [41]. Additional work revealed the flavonoid glucuronides, 3',5,7,8-tetrahydroxy-4'-methoxyflavone 8-O-β-D-glucopyranosiduronic acid methyl ester **13**, 4',5,7,8-tetrahydroxyflavone 8-O-β-D-glucopyranosiduronic acid methyl ester **14** and the first neolignans, helicterins A-F (**15-20**), helisterculin A **21**, helisterculin B **22** and helisorin **23**, isolated from *Helicteres isora* L. obtained from Indonesia (Figure 3) [42-43].

The helicterins were tested against avian myeloblastosis virus-reverse transcriptase (AMV-RT) and presented weak

**Figure 3:** Glycosidic and non glycosidic metabolites isolated from *Helicteres isora* [41-43].**Table 3:** Bioactivity of helicterins A-F **15-20**, helisterculin A **21**, helisterculin B **22**, helisorin **23** and adriamycin **24** against AMV-RT [42,43].

Compounds	AMV-RT, IC ₅₀ µM
15	66
16	172
17	417
18	372
19	120
20	226
21	1600
22	1000
23	460
24	66

activity (Table 3). However, compared to the standard drug for AMV infection, adriamycin **24**, and helicterins A **15** possessed equal activity [43]. There is no specific information regarding the mechanism of inhibition of this compound.

**Figure 4:** Map of Indonesia [46]. * The sites of ethnopharmacological field trips in endemic malaria regions in some islands. Each region is discussed separately in the following sections.

Anti-malarial agents: Ethnopharmacological studies revealed particular medicinal plants were used for malarial

fever therapy in some regions (Figure 4). Further experiments produced the extracts responsible for the claimed activity (Table 4) with only a small number investigated further to reveal individual compounds [44,45].

Table 4: Anti-malarial activity of extracts of some Indonesian medicinal plants [44-45].

Plants	Part of plant	Inhibition rates against <i>Plasmodium falciparum</i> , %
Apocynaceae		
<i>Catharanthus roseus</i> L.	AP	52
<i>Rauvolfia serpentina</i> (L.) Benth.	BK	66
Asteraceae		
<i>Achillea millefolium</i> L.	WP	98
<i>Ageratum conyzoides</i> L.	WP	62
Euphorbiaceae		
<i>Phyllanthus niruri</i> L.	WP	50*
Loganiaceae		
<i>Strychnos lucida</i> R.Br.	WD	100
Meliaceae		
<i>Azadirachta indica</i> Juss	LF	60
<i>Lansium domesticum</i> Corr.	BK	66
<i>Swietenia macrophylla</i> King.	SD	98
Myrtaceae		
<i>Baeckea frutescens</i> L.	LF	90
Rubiaceae		
<i>Morinda citrifolia</i> L.	FR	56
Zingiberaceae		
<i>Curcuma xanthorrhiza</i> Roxb.	RZ	100

AP: Aerial part; BK: Bark; WP: Whole plant; WD: Wood; LF: leaf; SD: Seed; FR: Fruit; RZ: Rhizome. All materials were prepared as a water extract. Concentration for assay, 1 mg/mL. *50% of inhibition at 3.5 µg/mL.

Sumatra Island: An expedition in Sumatra Island revealed *Anthocephalus chinensis*, *Beilschmiedia madang* BL. and *Brucea javanica* L. Merr to be used in malarial therapy [3].

Anthocephalus chinensis (Rubiaceae). The people in Indragiri Hulu area, Riau Province have used the bark, root and leaf of *Anthocephalus chinensis* in malaria therapy. Fourteen compounds were successfully isolated including glycosides, 3'-*O*-caffeoylsveroside **25**, sweroside **26**, loganic acid **27**, loganin **28**, loganol **29**, kelampayoside A (derived from "kelampayan", the local species' name) **30**, 8-epikingiside **31**, kelampayoside B **32** (Figure 5), and alkaloids vallesiachotamine **33**, isovallesiachotamine **34**, cadambine **35**, strictosidine lactam **36**, desoxycordifoline **37**, 5 α -carboxystrictosidine **38** (Figure 6) [47]. Compounds **26-29**, **31**, **33-37** were previously identified from various Rubiaceae species of different origins [48-52].




Figure 5: Glycosides isolated from the bark of *Anthocephalus chinensis* [47].




Figure 6: Alkaloids isolated from the bark of *Anthocephalus chinensis* [47].

Among the fourteen compounds (Figures 5 and 6), only eight were subjected to *in vitro* testing against the *Plasmodium falciparum* K1 strain. The results indicated no anti-malarial activity except the indole type compound cadambine **35**, which showed a moderate inhibitory activity at IC₅₀ of 6.77 µM and IC₉₀ 9.85 µM [47].

Beilschmiedia madang BL. (Lauraceae). The decocted wood of *B. madang* (locally called as 'medang kohat') has been used by people in Kepahiang to treat malaria fever [53]. From this was isolated dehatrine **39** (Figure 7) which was found to have anti-malarial activity with an IC₅₀ of 0.17 µM against the *Plasmodium falciparum* K1 strain, a chloroquine resistant strain [53].




Figure 7: Molecular structure of dehatrine [53].

Brucea javanica L. Merr. (Simaroubaceae). This species was found through an expedition to Rejang Lebong where the plant has been traditionally used to treat malaria. The same species is used by the Bugisse, 'people of Makasasar' for the same purpose. Successfully identified were eleven triterpenoids including, bruceajavanin A **40**, dihydrobruceajavanin A **41**, bruceajavanin B **42**, bruceantanol **43**, bruceantanol B **44**, bruceine A **45**, bruceine B **46**, bruceine C **47**, bruceine D **48**, bruceine J **49**, yandazolid A **50** [54] and five alkaloids including 11-hydroxy-canthine-6-one **51**, 11-hydroxy-1-methoxy-canthine-6-one **52**, canthine-6-one **53**, canthine-6-one-3-*N*-oxide **54** and bruceacanthinoside **55** (Figure 8) [55], of which the last four compounds **51-54** were previously reported from non-Indonesian Simaroubaceous medicinal plants [56-58].

Compounds **40** and **42** possessed moderate anti-malarial activity, e.g. IC₅₀ 1.1 µM and 4.4 µM against the *Plasmodium falciparum* K1 strain, respectively [55]. On the other hand, compound **43** presented only modest anti-malarial activity (IC₅₀, 25 µM) [55]. Some of the




Figure 8: Metabolites isolated from *Brucea javanica* [55].

compounds were also tested against *Babesia gibsoni* parasite of dogs which brucein A **45**, bruceinthinol **43** and B **44** possessed potent activities with IC₅₀ values of 4, 12 and 12 ng/mL, respectively [54].

Java and Lombok Islands: The Javanese and Lomboknesse native people have been infected with malarial fever for centuries. However, they survive by consuming young leaves of *Alstonia scholaris* [59]. Investigation on samples taken from Java and Lombok Island isolated 21 alkaloids, including, picaline **56**, *N*₁-methylburnamine **57**, akuammidina **58**, *φ*-akuammigine **59**, akuammiline **60**, akuammiline *N*₄-oxide **61**, deacetylakuammiline **62**, 5 α -methoxy-akuammiline **63**, tubotaiwinine **64**, tubotaiwinine *N*₄-oxide **65**, scholaricine **66**, *N*₄-methylscholaricine **67**, vallesamine **68**, alstonamine **69**, 6,7-*seco*-alstonamine **70**, 6,7-*seco*-19,20 α -epoxy-alstonamine **71**, leuconolam **72**, mataranine A **73**, mataranine B **74**, kotarajine **75** and (15*S*,16*S*)-losbamine **76** (Figure 9) [59, 60]. Tubotaiwine **64** and mataranine A **73** and B were claimed to be responsible for the anti-malarial activity (Table 5) [60,61].

Flores Island: An ethnopharmacological expedition reported that the people of Flores Island have used *Fagara rhetza* (Roxb.) DC, locally named as Haleza, to treat malarial fever. Eleven compounds were isolated in this study. The bark contains rutaecarpine **77**, evodiamine **78**, skimmianine **79**, zanthobungeanine **80**, *O*-geranylsinapyl




Figure 9: Alkaloids isolated from *Alstonia scholaris* [59, 60].

Table 5: Anti-malarial activity of several alkaloids isolated from *Alstonia scholaris* [60,61].

Alkaloids	EC ₅₀ , μ M
60	18
62	36
63	7
64	^a
69	24
70	17
73	7.4 ^b
74	9.7 ^c

^aNo potency. ^bBioactivity against *Plasmodium falciparum* K1 (an antifolate resistant parasite strain). ^cAgainst *Plasmodium falciparum* TM4 (an anti-folate sensitive parasite strain)




Figure 10: Metabolites isolated from *Fagara rhetza* [62-67].

alcohol **81**, *O*-geranylconiferyl alcohol **82**, hazaleamide **83**, hazaleanin A **84**, hazaleanin B **85**, isopropyl apioglucoside **86** and 4-hydroxyguaiacol apioglucoside **87**, of which compounds **77-80** were previously identified from other plants however no bioactivity data was reported (Figure 10) [62-67]. In this study, only hazaleamide **83** showed a moderate activity against *Plasmodium falciparum* with an IC_{50} value of 43 μ M [62-67].

Kalimantan (Borneo) Island: Malarial fever is a long lasting health problem of Dayak tribes that inhabit the isolated rainforests of Kalimantan Island. An expedition found two plants species, *Eurycoma longifolia* Jack. and *Lansium domesticum* Corr. Ser. have been intensively used for malarial fever treatment [68-69].

Eurycoma longifolia Jack. (Simaroubaceae). This plant is locally called “pasak bumi” by people of Borneo Island where it has been used to treat malaria, dysentery, glandular swelling and persisten fever [68]. Eleven molecules, 9-methoxycanthin-6-one **88**, 9-hydroxycanthin-6-one **89**, 9-methoxycanthin-6-one-*N*-oxide **90**, 9-hydroxycanthin-6-one-*N*-oxide **91**, β -carboline-1-propionic acid **92**, 7-methoxy- β -carboline-1-propionic acid **93**, eurycomanone **94**, 13,21-dihydroeurycomanone **95**, 13 β ,21-dihydroeurycomanone **96**, eurycomanol **97** and longilactone **98**, were isolated of which only the compounds **93** and **94** were shown to have weak anti plasmodium activity [68].




Figure 11: Alkaloids and terpenes isolated from *Eurycoma longifolia* [68].

Lansium domesticum Corr. Ser. (Meliaceae). Isolation revealed five compounds, 3-keto-22-hydroxynoceradiene **99**, onoceradienedione **100**, methyl lansiolate **101**, methyl lansiolate A **102** and methyl 15-acetoxylansiolate **103** (Figure 12) [69]. The anti-malarial activity of the isolates were tested against *Plasmodium berghei* (Table 6).

Table 6: Anti malarial activity of isolates from *Lansium domesticum* Corr. Ser [69].

Quassinoids	IC_{50} , μ M
99	2.41
100	1.66
101	0.65
102	0.69
103	0.17




Figure 12: Anti-malarial agents isolated from *Lansium domesticum* [69].

The most active compound isolated was the methyl ester 15-acetoxylansiolate **103**. This *in vitro* test result agreed with the *in vivo* test by using infected *Plasmodium berghei* mice where **103** possessed 44% parasitemia inhibition at 50 mg/kg daily doses [69].

Sulawesi Island: There are two traditionally prepared medicines for malarial therapy; the seed kernel of *Caesalpinia crista* species (called ‘Bakor’) and decocted root bark and stem of *Quassia indica* (Gaertn.) Nooteboom, named ‘Tobello’ by central Sulawesi people.

Caesalpinia crista (Fabaceae). From this species were isolated at least 24 cassane and norcassane type compounds. The quassonoids are caesalpinins C-P **104-117**, norcaesalpinins A-F **118-123**, caesalmin B **124**, E **125** and G **126**, caesalpin F **127**, 14(17)-dehydrocaesalpin F **128**, caesaldekarin E **129**, 2-acetoxy-3-deacetoxycaesaldekarin E **130**, 1-deacetoxy-1-oxocaesalmin C **131**, 3-deacetoxy-6-acetoxycaesaldekarin E **132**, 2-acetoxycaesaldekarin E **133**, bonducellpins A-C **134-136** and 7-acetoxybonducellpin C **137** (Figure 13) [70-73].

The preliminary anti-malarial testing in mice infected with chloroquinene-resistant *Plasmodium berghei*, showed that norcaesalpinin A **118** suppressed the parasitemia by 48.0, 40.9 and 33.0% at doses of 10, 1 and 0.1 mg/kg, respectively [70].

Most of the isolates were also tested against *Plasmodium falciparum* culture (see Table 7) with **122** as the most active quassonoid with an IC_{50} value of 0.09 μ M [74].

A QSAR study on the cassane and norcassane-type diterpenes observed that the presence of an acetoxy group resulted in a higher anti-malarial activity than when a hydroxyl substituent was present. On the other hand, any additional functional group on the C-ring in 17-norcassane-type diterpenes reduced the activity [74].




Figure 13: Quassonoids isolated from *Caesalpinia crista* [70-73].




Figure 13 (continued): Quassinoids isolated from *Caesalpinia crista* [70-73].

Quassia indica (Gaertn.) Nooteboom. (Simaroubaceae). The chemical screening of the species isolated eight quassinoids, samaderine B-C, E, X-Z, **139-144**, indaquassin C, X **145-146**, simarinolide **147** and 2-*O*-glucosylsamaderine C **148** (Figure 14). Compounds **139-141**, **145**, **147** were previously isolated from other simaroubaceous [76]. Four of the isolates were tested for anti-malarial activity against *Plasmodium falciparum* K1 (Table 8) which showed that samaderine **142** was the most active compound [76].

Table 7: Isolated quassinoids activity against *Plasmodium falciparum* [73,74].

Quassinoids	IC ₅₀ μ M	Quassinoids	IC ₅₀ μ M
104	0.76	121	2.0
105	0.80	122	0.09
106	6.50	123	0.14
107	0.65	124	0.80
109	>10	125	>10
110	>10	126	>10
111	1.00	128	0.20
112	0.4	129	4.0
113	0.65	130	0.098
114	>10	131	2.9
115	0.12	133	6.5
116	>10	135	0.24
117	1.7	136	0.12
118	0.80	137	0.60
119	0.26	138*	0.29
120	5.0		

* Chloroquine **138**, positive control, was discovered in 1934 and it has been used as anti-malarial agent which also shows anti-viral effects [75].




Figure 14 : Quassinoids isolated from *Quassia indica* [76].

Table 8: Anti-malarial activity of isolates from *Quassia indica* (Gaertn.) Nooteboom [76].

Quassinoids	IC ₅₀ , nM
139	210
141	56
142	14
144	71

Anti-bacterial agents: Indigenous Indonesian people has treated diarrhea, swelling, redness and fever with medicinal plant preparations [11,77] in which these symptoms have modern clinical correlations with bacteria causing diseases. Most of the anti-bacterial outcomes were preliminary studies of extracts (see Table 9), which only gave indications of activity. Several extracts such as those from *Terminalia catappa*, *Swietenia mahagoni* Jacq., *Phyllanthus acuminatus*, *Ipomoea* spp., *Tylopora asmathica* and *Hyptis brevipes* possessed high activity which might provide a stimulus for further research [78]. Additional work was performed on a few species resulting in the following:

Clerodendron calamisotum L. and *Clerodendron paniculatum* L. (Verbenaceae): At least five alkaloids were present in *Clerodendron calamisotum* L. and *Clerodendron paniculatum* L. however the alkaloids were not responsible for the anti-bacterial activity [61].

Table 9: Anti-bacterial activities of extracts of Indonesian medicinal plants [24,78].

Plant	Part of plant	Anti-microbial activities**					
		BS	SA	EC	SC	FO	XC
Acanthaceae							
<i>Gendarusa vulgaris</i> Nees	LF, RT, ST		+				
<i>Graptophyllum pictum</i> (L.) Griff.*							DG
<i>Staurogyne sp</i>	LF, RT, ST		+		+	+	
Amaranthaceae							
<i>Celosia argentea</i> L. fma	FR, LF, ST		++	+		+	
<i>Celosia plumose</i> (Voss) Back	FR, LF, ST		++	+		+	
Amaryllidaceae							
<i>Crinum asiaticum</i>	LF		++	+	+		
<i>Crinum lineare</i> L.	LF				++		
<i>Curculigo sp.</i>	LF, RT, ST		++	+			
Anarcadiaceae							
<i>Mangifera foetida</i> Lour.	LF		+				
<i>Mangifera indica</i> L.	BK		++				
Apocynaceae							
<i>Alstonia scholaris</i> (L) R. Br.	LF, ST		++				
<i>Cantharanthus roseus</i> (L.) G. Don*			+				
Araceae							
<i>Acorus calamis</i> L.	LF, RT,RZ		+	+			
Araliaceae							
<i>Aralidium pinnatifidum</i> Miq.	LF		++	+			
Balsaminaceae							
<i>Impatiens balsamina</i> L.	LF, ST		+				
Begoniaceae							
<i>Begonia isopera</i> Bl.	LF, ST		+		+	+	
Bombaceae							
<i>Ceiba petandra</i> (L.) Gaertn.	LF, ST		++				
Bromeliaceae							
<i>Ananas comosus</i> (L.) Merr.	LF		+	+		+	
Capparaceae							
<i>Crateva religiosa</i> Forst.	LF, ST		+	+			
Clusiaceae							
<i>Cratoxylon formosum</i> (Jack)	BK		+				
<i>Garcinia atroviridis</i> Griff.	LF, ST		++	+	+		
<i>Garcinia mangostana</i> L.	BK		+				
<i>Garcinia parvifolia</i> Miq.	LF, ST		++	+	+		
Compositae							
<i>Blumea balsamifera</i> DC.	LF, RT, ST		+	+		+	
<i>Plucea indica</i> (L.) Less.	LF, ST		++	+		+	
Connaraceae							
<i>Rourea mimosoides</i> (Vahl)	LF, ST		+	+			
Convolvaceae							
<i>Ipomoea batatas</i> (L.) Lamk.	LF, ST		+				
<i>Merremia umbellata</i> (L.) Hall.	LF, ST		+				
Cyperaceae							
<i>Scleria purpurascens</i> Steud.	LF, ST		+				
Dilleniaceae							
<i>Dillenia meliosmifolia</i> Hook.	LF		+				
<i>Tetracia asiatica</i> (Lour.)	LF		+		+		
<i>Tetracera sp.</i>	LF		+				
Ebenaceae							
<i>Diospyros sp.</i>	LF, ST		++	+			
Euphorbiaceae							
<i>Breynia racemosa</i> M.	LF, ST		+				
<i>Croton cf. caudatus</i> Geisel.	LF, ST, BK		++				
<i>Galearia filiformis</i> Boerl.	LF				+		
	BK		++	+	+		
<i>Galearia sp.</i>	LF, ST		++		+	+	
<i>Jatropha curcas</i> L.	LF				+		
Euphorbiaceae							
<i>Macaranga gigantea</i> M.	LF		++	+			
<i>Macaranga gigantea</i> M.	BK		+	+	+		
<i>Macaranga triloba</i> M.	LF		+	+		+	
<i>Mallotus ricinoides</i> M.	LF		+				
Euphorbiaceae							
<i>Phyllanthus emblica</i> L.	LF, ST		+	+			
<i>Pimeliendron papaveriodes</i>	LF, ST		+	+			

Table 9 (continued): Anti-bacterial activities of extracts of Indonesian medicinal plants [24,78].


Plant	Part of plant	Anti-microbial activities**					
		BS	SA	EC	SC	FO	XC
Gnetaceae							
<i>Gnetum latifolium</i> Bl.	LF, ST		+			+	
Hanguanaceae							
<i>Hanguana malayana</i> (Jack) Merr.	LF, ST		++				
Labiatae							
<i>Coleus scutellarioides</i> (L.) Bth.	LF, ST		++			+	
<i>Hyptis brevipes</i> *			+++	+			+
<i>Ocimum basilicum</i> L.	FL, LF, ST		+	+			
<i>Ocimum tenuiflorum</i> L.	FL, LF, ST		+	+			
<i>Orthosiphon aristatus</i> (Bl.) Miq.	LF, ST		+				
Lauraceae							
<i>Litsea elliptica</i> Bl.	LF		+				
<i>Litsea robusta</i> Bl.	LF		++				
Lecythidaceae							
<i>Barringtonia lanceolata</i> (Bl.)	LF		++				
<i>Barringtonia sp.</i>	BK		++				
Leguminosae							
<i>Bauhinia sp</i>	LF, ST				+		
<i>Crotalaria juncea</i> L.*		DG					
<i>Cassia cf. nodosa</i> Buch. Ham.	BK		++	++		+	
<i>Flemingia strobilifera</i> (R.Br.)	FR, LF, ST		+				
<i>Mimosa pigra</i> L.	LF, ST		+				
<i>Sesbania aculeate</i> Poir.	LF, RT, ST		++				
<i>Tamarindus indica</i> L.	LF, ST		++	++			
Longaniaceae							
<i>Fagraea cfr. auriculata</i> Jack	LF		+	+	+		
Lythraceae							
<i>Lagerstromia speciosa</i> (L.)	LF, ST		++	++			
Malvaceae							
<i>Abelmoschus ficulneus</i> W. et H.	LF, ST		++		++		
<i>Abelmoschus moschatus</i> Medik.	LF, ST		+				
<i>Gossypium barbadense</i> L. var.	LF, ST		++				
<i>Hibiscus x archeri</i> Wats.	LF, ST		++				
<i>Hibiscus sabdariffa</i> L.	LF, ST		++				
<i>Hibiscus tiliaceus</i> L.	BK		++				
	LF, ST		+				
Melastomataceae							
<i>Dissochaeta gracilis</i> (jack) Bl.	LF, ST		+			+	
<i>Melastoma affine</i> D.Don	LF, ST		++	+	+		
<i>Melastoma malabathricum</i> L.	FL,ST,LF		++	+	+	+	
<i>Phyllagathis rotundifolia</i> (Jack) Bl.	LF, ST		+	+	+		
Meliaceae							
<i>Aglaia eximia</i> Miq.	BK		+	+		+	
<i>Swietenia mahagoni</i> Jcq.*			+	++	DG		+
			+				+
<i>Swietenia mahagoni</i> L. *	PC		+	+			
Menispermaceae							
<i>Arcangelisia flava</i> (L.) Merr.	LF, RT, ST		++		++	+	
<i>Tinomisium petiolare</i> Hook. F. et Thoms.	BK		+				
Moraceae							
<i>Artocarpus elasticus</i> Reinw. Ex Bel.	BK		++			+	
						+	
<i>Artocarpus nitidus</i> Trec.	BK			+			
<i>Ficus grossularioides</i> Burm. f.	LF, ST		+	+			
<i>Ficus macrocarpa</i> Kurz	LF, ST		++	+			
<i>Ficus pandana</i> Burm. f.	LF, ST		++	+			
Myristicaceae							
<i>Horsfieldia glabra</i> (Bl.) Warb.	BK		+				
<i>Horsfieldia wallichii</i> (Hook. F. et Thunb.) Warb.	BK		++				
Myrtaceae							
<i>Eugenia polyantha</i> Wight	LF, ST		+			+	

Table 9 (continued): Anti-bacterial activities of extracts of Indonesian medicinal plants [24,78].


Plant	Part of plant	Anti-microbial activities**					
		BS	SA	EC	SC	FO	XC
Myrtaceae							
<i>Psidium guajava</i> L.	LF, ST		++	+			
<i>Rhodmania cinerea</i> Jack	LF, ST		++	+			
Pandanaceae							
<i>Pandanus</i> sp.	LF		+				
Passifloraceae							
<i>Adenia cordifolia</i> ENgl.	LF, ST		++	+			
Rhamnaceae							
<i>Gouania lepostachya</i> DC.	LF, ST		++		+	+	
Rhizophoraceae							
<i>Anisophyllea disticha</i> (Jack) Baill.	LF, ST		+		+		
Rubiaceae							
<i>Hedyotis capitellata</i> Wall.	LF, ST		+				
<i>Hedyotis leucocarpa</i> Elm.	LF, ST		+				
<i>Mussaenda frondosa</i> L.	LF, ST		+				
<i>Uncaria gambir</i> (Hunt.) Roxb.	LF, ST		++			+	
Rutaceae							
<i>Citrus aurantifolia</i> Swingle	FR		++	++			
Simaroubaceae							
<i>Brucea javanica</i> (L.) Merr.	LF, ST		++		+		
Staphylaceae							
<i>Turpinia sphaerocarpa</i> Hassk.	LF, ST		++	++			
Symplocaceae							
<i>Symplocos cochinchinensis</i> (Lour.) Moore	LF, ST		+	+	++	+	
Theaceae							
<i>Eurya acuminata</i> DC.	LF, ST		+				
Thymelaeaceae							
<i>Aqualaria malaccensis</i> Lamk.	LF, ST		+				
Tiliaceae							
<i>Elaeocarpus cf. mastersii</i> King	LF, ST, BK		++	++			
<i>Grewia acuminata</i> Juss.	BK		++			+	
Ulmaceae							
<i>Trema tomentose</i> (Roxb.) Hara	LF, ST		+				
Urticaceae							
<i>Dendrocnide stimulans</i> (L.f.) Chew	RT		++				
Violaceae							
<i>Rinorea anguifera</i> (Lour.) O.K.	BK		+	+			
Zingiberaceae							
<i>Boesenbergia rotunda</i> (L.) Mansf.	LF, RZ, ST		+				
<i>Costus</i> sp.	LF, ST		+				
<i>Curcuma cf. heyneana</i> Val. Et v. Zijp	RZ					+	
<i>Curcuma</i> sp.	RZ		+		+		
<i>Zingiber purpureum</i> Roxb.	RZ		+		+		

LF: Leaf, ST: Stem, BK: Bark, PC: Pericarp, RZ: Rhizome, BS: *Bacillus subtilis*, SA: *Staphylococcus aureus*, EC: *Escherichia coli*, SC: *Saccharomyces cerevisiae*, FO: *Fusarium oxysporum*, XC: *Xanthomonas campestris*. All samples are methanol extracts except indicated by * extract of DCM. ** + is 25% or less than control; ++, equal to the control; +++, 25-50% more than the control; +++++, 50% and more than the control.


Curcuma xanthorrhiza Roxb.: The species sample collected from Yogyakarta contains sesquiterpenes, α -curcumene **149**, germacrone **150**, zederone **151**, xanthorizol **152** and curcumin **153** (Figure 15). The crude extracts possessed moderate antibacterial activity (MRSA) while the isolated curcumene showed less activity [79]. The extract has potential against *Staphylococcus* mutans for dental caries prevention [80].

**Figure 15:** Sesquiterpenes isolated from *Curcuma xanthorrhiza* [79].

Mitrephora celebica (Annonaceae): Two oropheic acid and four diterpenes were isolated from the bark of the species, oropheic acid **154**, 13,14-dihydrooropheic acid **155**, *ent*-trachyloban-19-oic acid **156**, *ent*-kaur-16-en-19-oic acid **157**, 8(14),15-pimaradien-18-oic acid **158** and 7,15-pimaradien-18-oic acid **159** (Figure 16) [81, 82]. These compounds are proposed to be responsible for the antibacterial activities of the stem bark extracts against methicillin-resistant *Staphylococcus aureus* and *Mycobacterium smegmatis*.

**Figure 16:** Oropheic acids and diterpenes isolated from *Mitrephora celebica* [81,82].

Among the compounds, **154**, **155**, **156** and **157** exhibited significant inhibitory activity with an MIC of 25, 12.5, 6.25, 6.25 μ g/mL, respectively while the others showed an MIC of more than 100 μ g/mL [81,82]. This difference might be correlated with the existence of an extra bridged ring in ring C of the structure.

**Figure 17:** Alkaloids isolated from *Psychotria malayana* [61].

Psychotria malayana Jack (Rubiaceae): The aqueous extracts of the leaves or bark of the species has been used traditionally to treat infections on open wounds by the Lomboknese. Chemical investigation on the locally named as "lolon jalun" plant successfully identified six alkaloids, *meso*-chimonanthine **160**, calcycanthine **161**, *iso*-calcycanthine **162** and hodgekinsine **163** and two

probable new alkaloids named LPM-574 **164** and LMP-186 **165** (Figure 17) [61]. Hodgkinsine **163** presents the major alkaloid while *meso*-chimonanthine **160** exists as a minor alkaloid constituent along with two others which were still under investigation [59]. Further chemical investigation on their activities are discontinued [59]. However, initial bacterial testing (Table 10) revealed that isolate **164** can caused bacteriolysis at 1.0 mg/mL but suppressed the growth at 0.5 mg/mL [61].




Figure 18: Alkaloids isolated from *Voacanga foetida* [84].

Voacanga foetida (Bl.) Rolfe (Apocynaceae): A phytochemical study on a Lombok Island sample isolated seven alkaloids, lombine **166** and coronaridine **167** [84], uncarine **168** voacangine **169**, voacristine **170**, mataranine A **73** and B **74** (Figure 18). The latter two compounds also exist in *Alstonia scholaris* collected from the same region.

Table 10: Bacteriostatic activity of alkaloids isolated from *Psychotria malayana* Jack [61].

Sample	C, mg/mL	Bacteriostatic activity	
		SA	EC
163	1.0	-	-
	0.5	-	-
164	1.0	++	++
	0.5	+	+

Note: C : Concentration ; SA: *Staphylococcus aureus*; EC: *Escherichia coli*.

Table 11: Anti-yeast like fungal activity of extracts of medicinal plants [87].

Plants	Part of plant	Yeast like fungal species				
		CN	SC	CA	WD	AA
Zingiberaceae						
<i>Alpinia galanga</i>	ST				11.9± 0.8	
<i>Alpinia galanga</i>	RZ	10.2±0.7			27.3 ±7.5	9.5 ±0.0
<i>Alpinia mutica</i>	ST				8.2 ±0.7	
<i>Curcuma zedoaria</i>	RZ	14.8±2.1	13.3±0.2	13.5± 0.0	11.5 ±1.0	
<i>Zingiber purpureum</i>	RZ	11.4±0.5		9.3± 0.9	11.3± 1.2	11.1± 0.4
<i>Zingiber officinale</i>	RZ			11.5± 0.0		

CN: *Cryptococcus neoformans*; SC: *Sacharomyces cerevisiae*; CA: *Candida albicans*; WD: *Wangiellia dermatitidis*; AA: *Alternaria alternate*; ST: Stalk; RZ: Rhizome. All sample are ethanolic extracts; Antifungal activity showed by mean diameter (mean ± SE, mm) of fungal growth inhibition zones. Sterile paper disc (7.5 mm diameter) was impregnated into 0.2 mg/μL ethanolic extract for disk diffusion assays.

Table 12: Anti-filamentous fungal activity of extracts of medicinal plants [78,87].

Plants	Filamentous like fungal species									
	PU	RS	SR	AF	PP	FO	MG	PB	Rh	TM
Acanthaceae										
<i>Graptophyllum pictum</i> (L.) Griff.*	+	++	+		+					
<i>Graptophyllum pictum</i> (L.) Griff.**			+							
Apocynaceae										
<i>Alstonia scholaris</i> (L.) R.Br.**	+		+							
<i>Cantharanthus roseus</i> (L.) G. Don**	+		+							
Asclepiadaceae										
<i>Tylophora asthmatica</i> *					DG					

Lombine **166** is responsible for the antibacterial activity of the original extract as this alkaloid exhibited bactericidal activity against *Staphylococcus aureus* and *Escherichia coli* resulting in 94% and 95% cell death, respectively at a concentration of 0.5 mg/mL compared to crude extracts which only partially inhibited at the same concentration [83]. Voacristine **170** showed lower activity with 87% of bacterial cell death at 1.0 mg/mL [61].

Antifungal agents: Indonesian medicinal plants have also been prepared traditionally for fungi caused diseases [84, 85]. Investigations resulted in several species extracts showing significant activities (see Tables 11 and 12) [86, 87]. For example, the extracts of *Tylophora asthmatica*, *Phyllanthus acuminatus* Vahl. and *Swietenia mahogany* Jacq. contain potent compounds against *Phytium ultimum*. On the other hand, extracts of *Ipomoea* spp and *Swietenia mahogany* Jacq. gave significant inhibition against *Sclerotium rolfsii*. Other anti-fungal studies showed *Wangiellia dermatitidis* and *Microsporum gypseum* are more sensitive against Zingiberaceae extracts.

In summary, this review has highlighted the investigations which studied 181 of around six thousand Indonesian medicinal plants which revealed 165 isolates. Some potent drug leads were isolated, eg β -sesquiphellandrene as an anti-rhinoviral agent, dehatrine, mataranine A and B as anti-malarial agents and lombine and LPM-574 as anti-bacterial agents. The remaining preliminary studies on these Indonesian medicinal plants indicate that they are prospective sources of potent anti-microbial constituents and require further studies to establish these outcomes.

Acknowledgments: ASN would like to thank the AusAID program and the University of Wollongong, Australia for ADS and HDR scholarships.

Table 12 (Continued): Anti-filamentous fungal activity of extracts of medicinal plants [78,87].

Plants	Filamentous like fungal species									
	PU	RS	SR	AF	PP	FO	MG	PB	Rh	TM
<i>Tylophora asthmatica</i> **	+++									
Convolvulaceae										
<i>Ipomoea</i> spp.*	++	+	+							
<i>Ipomoea</i> spp.**	++	++	++++							
Combretaceae										
<i>Terminalia catappa</i> L.*	+	+	+	+						
<i>Terminalia catappa</i> L.**	++++				++					
Elaeocarpaceae										
<i>Elaeocarpus grandiflorus</i> S.*			+	+						
<i>Elaeocarpus grandiflorus</i> S.**	++									
Euphorbiaceae										
<i>Phyllanthus acuminatus</i> Vahl.*	++		+	++						
<i>Phyllanthus acuminatus</i> Vahl.**	+++	+								
Fabaceae										
<i>Erythrina variegata</i> L.*	+	+		+						
<i>Antidesma bunius</i> (L.) Spreng*		+	+	+						
Fibaceae										
<i>Sesbania grandiflora</i> Pers*	++			+						
Labiatae										
<i>Hyptis brevipes</i> *	+	+	+	+	++					
Leguminosae										
<i>Crotalaria juncea</i> L.*		+			+					
<i>Crotalaria juncea</i> L.**	+									
Meliaceae										
<i>Swietenia mahagoni</i> Jacq.*	+++	++	+++							
<i>Swietenia mahagoni</i> L. (leaf)*				++	+					
<i>Swietenia mahagoni</i> L. (Pericarp)*	++	+	+	+						
<i>Swietenia mahagoni</i> L. (Pericarp)**		+		+						
Moringaceae										
<i>Moringa oleifera</i> Lam.**	+									
Rubiaceae										
<i>Mussaenda pubescens</i> Ait. f.**	+	++		+						
<i>Morinda citrifolia</i> L.*		+								
<i>Morinda citrifolia</i> L.**	++	+	+	+						
Zingiberaceae										
<i>Alpinia galanga</i> (ST)***								8.2		
<i>Alpinia galanga</i> ***				14.4		11.9	20.6	31.1	12.2	16.6
<i>Curcuma globosus</i> ***				9.5			25.3	9.5	14.2	9.5
<i>Curcuma zedoaria</i> ***							27.9	8.2		26.4
<i>Etlingera elatior</i> ***							9.5			
<i>Etlingera littoralis</i> ***				8.8						
<i>Zingiber purpureum</i> ***				14.2		9.5	27.3	17.1	11.5	20.6
<i>Zingiber officinale</i> ***							20.8	12.2	8.8	18.2

PU: *Phytium ultimum*; RS: *Rhizoctonia solani*; SR: *Sclerotium rolfsii*; AF: *Aspergillus fumigatus*; PP: *Phytophthora parasitica*; FO: *Fusarium oxysporum*; MG: *Microsporium gypseum*; PB: *Pseudallescheria*; RH: *Rhizopus* sp; TM: *Trichophyton mentagrophytes*; DG: Decrease growth. All extracts were obtained from the aerial part of plants except, *S. Mahogany*, whose seed was used and the Zingiberaceae, whose rhizome was used unless stated as ST (stalk). * DCM extract; **MeOH extract; ***EtOH extract. Activity shown as symbol + 25% less than control, ++ equal to the control, +++, 25-50% more than control, ++++ 50% than control. In case of Zingiberaceae extracts, result was showed as mean diameter (in mm) of fungal growth inhibition zones. Sterile paper disc (7.5 mm diameter) was impregnated into 0.2 mg/μL ethanolic extract for disk diffusion assays.

References

- Younce WC. (2001) *Indonesia: issues, historical background and bibliography*. Nova Science Publishers, New York.
- Witton P, Elliott M, Greenway P, Jealous, V. (2003) *Indonesia*. Lonely Planet Publication, Singapore.
- Kitagawa I, Shibuya H. (1995) Pharmacochemical investigation of Indonesian medicinal plants. *Proceedings of the Phytochemical Society of Europe*, 37, 335-358.
- Government of the Republic of Indonesia - State Ministry for Environment. (1997) *Agenda 21-Indonesia. A national strategy for sustainable development*. Government of the Republic of Indonesia - State Ministry for Environment, Jakarta.
- Government of the Republic of Indonesia. (1997) *Country paper on the implementation of biodiversity management in Indonesia*. Government of the Republic of Indonesia, Jakarta.
- Beers S. (2001) *Jamu: The ancient art of herbal healing*. Periplus Editions, Singapore.
- Williamson E. (2006) Systems of traditional medicine from South and Sout East Asia. *The Pharmaceutical Journal*, 276, 539-540.
- Pols H. (2010) The triumph of jamu. *Inside Indonesia*, 100, 1-6.
- Susanto YW. (1996) *Sejarah jamu gendong*. Departemen Kesehatan Republik Indonesia, Jakarta.
- Kozok U. (1999) *Warisan leluhur: sastra lama dan aksara Batak*. Kepustakaan Populer Gramedia, Jakarta.
- Masyarakat Pernaskahan Nusantara. (2000) *Naskah sebagai sumber pengetahuan budaya : kumpulan makalah simposium internasional pernaknahan nusantara II, 26 November 1998*. Masyarakat Pernaskahan Nusantara, Jakarta.
- Ranggasutrasna KN, Pakubuwana V, Marsono D. (1991) *Centhini: Tambangraras-among raga*. Balai Pustaka, Jakarta.
- Tinggen IN. (1996) *Sarining usada bali pusaka leluhur*. Indra Jaya, Singaraja.
- Leaman DJ. (1996) *The medicinal ethnobotany of the Kenyah of East Kalimantan (Indonesian Borneo)*. Ph.D thesis, University of Ottawa, Canada, 1-385.

- [15] Riswan S, Sangat-Rumantyo H. (1991) The opportunity, technical and practical constraints of the plant derived drug development in Indonesia, in *Workshop on drug development, biological diversity and economic growth*. Bethesda, MD, USA, 2-8.
- [16] Stevensen C. (1999) Jamu: an Indonesian herbal tradition with a long past, a little known present and an uncertain future. *Complement Therapy Nursing and Midwifery*, **5**, 1-3.
- [17] Pols H. (2009) European physician and botanists, indigenous medicine in the Dutch East Indies, and colonial network of mediation. *East asia sciences, technology and society: an international journal*, **3**, 173-208.
- [18] Bontius J. (1642) *De medicina Indorum*. Franciscus Hackius, Lugduni Batavorum, Leiden.
- [19] Buenz EJ, Johnson HE, Beekman EM, Motley TJ, Bauer BA. (2005) Bioprospecting Rumphius's Ambonese Herbal: Volume I. *Journal of Ethnopharmacology*, **96**, 57-70.
- [20] Heyne K. (1927) *De nuttige planten van der Nederlandsch Indie (revised and enlarged edition)*. Buitenzorg, Batavia.
- [21] van Steenis-Kruseman MJ. (1953) Select Indonesian medicinal plants. *Organisation for Scientific Research in Indonesia Bulletin*, **18**, 1-90.
- [22] WHO. (2000) *General guidelines for methodologies on research and evaluation of traditional medicine*. World Health Organisation.
- [23] Elliott S, Brimacombe J. (1987) The medicinal plants of Gunung Leuser National Park, Indonesia. *Journal of Ethnopharmacology*, **19**, 285-317.
- [24] Grosvenor PW, Supriono A, Gray DO. (1995) Medicinal plants from Riau province, Sumatra, Indonesia. Part 1: Uses. *Journal of Ethnopharmacology*, **45**, 75-95.
- [25] Grosvenor PW, Supriono A, Gray DO. (1995) Medicinal plants from Riau province, Sumatra, Indonesia. Part 2: Antibacterial and antifungal activity. *Journal of Ethnopharmacology*, **45**, 97-111.
- [26] Roosita K, Kusharto CM, Sekiyama M, Fachrurazi Y, Ohtsuka R. (2008) Medicinal plants used by the villagers of a Sundanese community in West Java, Indonesia. *Journal of Ethnopharmacology*, **115**, 72-81.
- [27] Balick MJ. (1994) Ethnobotany, drug development and biodiversity conservation-exploring the linkages. In *Ethnobotany and the search for new drugs*. Chadwick, DJ, Marsh, J (Eds). John Wiley & Sons, England.
- [28] Limananti AI, Triratnawati A. (2003) Jamu Cekok Components for Treating Children Have No Appetite: An Ethnomedicine Approach. *MAKARA of Health Series*, **7**, 11-20.
- [29] Kurokawa M, Ochiai H, Nagasaka K, Neki M, Xu H, Kadota S, Sutardjo S, Matsumoto T, Namba T, Shiraki K. (1993) Antiviral traditional medicines against herpes simplex virus (HSV-1), poliovirus, and measles virus *in vitro* and their therapeutic efficacies for HSV-1 infection in mice. *Antiviral Research*, **22**, 175-88.
- [30] Morita H, Kishi E, Takeya K, Itokawa H, Tanaka O. (1990) New quassinoids from the roots of *Eurycoma longifolia*. *Chemical letter*, **59**, 749-752.
- [31] Hattori M, Nakabayashi T, Lim Y. (1995) Inhibitory effects of various Ayurvedic and Pnamanian medicinal plants on the infection of herpes simplex virus-1 *in vitro* and *in vivo*. *Phytoteraphy Research*, **9**, 270-276.
- [32] Fukuchi K, Sakagami H, Okuda T. (1989) Inhibition of herpes simplex virus infection by tannins and related compounds. *Antiviral research*, **11**, 285-298.
- [33] Lohezic-Le Devehat F, Bakhtiar A, Bezivin C, Amoros M, Shiraki K. (1999) Antiviral and cytotoxic activities of some Indonesian plants. *Fitoterapia*, **73**, 400-405.
- [34] Nawawi A, Nakamura N, Hattori M, Kurokawa M, Shiraki K. (1999) Inhibitory effects of Indonesian medicinal plants on the infection of herpes simplex virus type 1. *Phytotherapy Research*, **13**, 37-41.
- [35] Bauer DJ, Selway JWT, Batchelor JF, Tisdale M, Caldwell IC, Young DAB. (1981) 4',6-Dichloroflavan, a new antirhinovirus compound. *Nature*, **292**, 369-370.
- [36] Denyer CV, Jackson P, Loakes DM. (1994) Isolation of antirhinoviral sesquiterpenes from ginger (*Zingiber officinale*). *Journal of Natural Products*, **57**, 658-662.
- [37] Kusumoto IT, Kakiuchi N, Hattori M, Namba T, Sutardjo S, Shimotohno K. (1992) Screening of some Indonesian medicinal plants for inhibitory effects on HIV-1 protease. *Shoyakugaku Zasshi*, **46**, 190-193.
- [38] Otake T, Mori H, Morimoto M, Ueba N, Sutradjo S, Kusumoto IT, Hattori M, Namba T. (1995) Screening of Indonesian plant extracts for anti-human immunodeficiency virus type 1 (HIV-1) activity. *Phytotherapy Research*, **9**, 6-10.
- [39] Ma C, Nakabayashi T, Miyashiro H, Hattori M, El-Meckkawy S, Namba T, Shimohtohno K. (1994) Screening of traditional medicines for their inhibitory effects on human immunodeficiency virus protease. *Wakan Iyakugaku Zasshi*, **11**, 416-417.
- [40] Chen SX, Wan M, Loh BN. (1996) Active constituents against HIV-1 protease from *Garcinia mangostana*. *Planta Medica*, **62**, 381-382.
- [41] Satake T, Kamiya K, Saiki Y, Hama T, Fujimoto Y, Kitanaka S, Kimura Y, Uzawa J, Endang H, Umar M. (1999) Studies on the constituents of fruits of *Helicteres isora* L. *Chemical & Pharmaceutical Bulletin*, **47**, 1444-1447.
- [42] Tezuka Y, Terazono M, Kusumoto TI, Kawashima W, Hatanaka Y, Kadota S, Hattori M, Namba T, Kikuchi T, Tanaka K, Supriyatna S. (1999) Helisterculins A and B, 2 new (7.5',8.2')-neolignans, and helisorin, the 1st (6.4',7.5',8.2')-neolignan, from the Indonesian medicinal plant *Helicteres isora*. *Helvetica Chimica Acta*, **82**, 408-417.
- [43] Tezuka Y, Terazono M, Kusumoto TI, Hatanaka Y, Kadota S, Hattori M, Namba T, Kikuchi T, Tanaka K, Supriyatna S. (2000) Helicterins A - F, six new dimeric (7.5',8.2')-neolignans from the Indonesian medicinal plant *Helicteres isora*. *Helvetica Chimica Acta*, **83**, 2908-2919.
- [44] Murnigsih T, Subeki Matsuura H, Takahashi K, Yamasaki M, Yamato O, Maede Y, Katakura K, Suzuki M, Kobayashi S, Chairul, Yoshihara T. (2005) Evaluation of the inhibitory activities of the extracts of Indonesian traditional medicinal plants against *Plasmodium falciparum* and *Babesia gibsoni*. *The Journal of Veterinary Medical Science*, **67**, 829-831.
- [45] Mustofa ET, Sholikhah Wahyuono S. (2007) In vitro and invivo antiplasmodial activity and cytotoxicity of extracts of *Phyllanthus niruri* L. herbs traditionally used to treat malaria in Indonesia. *Southeast Asian Journal of Tropical Medicine and Public Health*, **38**, 609-615.
- [46] Fédération Internationale de Géomètres. (2011) *Facts about Indonesia*.

- [47] Kitagawa I, Wei H, Nagao S, Mahmud T, Hori K, Kobayashi M, Uji T, Shibuya H. (1996) Indonesian medicinal plants. XIV. Characterization of 3'-O-caffeoylsweroside, a new secoiridoid glucoside, and kelampayosides A and B, two new phenolic apioglucosides, from the bark of *Anthocephalus chinensis* (Rubiaceae). *Chemical & Pharmaceutical Bulletin*, **44**, 1162-1167.
- [48] Garcia J, Lavaitte S, Gey C. (1989) 8-Epikingiside and its vanillate ester, isolated from *Gentiana pyrenaica*. *Phytochemistry*, **28**, 2199-2201.
- [49] El-Naggar LJ, Beal JL. (1980) Iridoids. A Review. *Journal of Natural Products*, **43**, 649-707.
- [50] Brown RT, Chapple CL, Lashford AG. (1977) Isolation of strictosidine (isovincoside) lactam from *Nauclea latifolia*. *Phytochemistry*, **16**, 1619-1620.
- [51] Brown RT, Warambwa BFM. (1978) Adina alkaloids: Isolation of desoxycordifoline. *Phytochemistry*, **17**, 1686-1687.
- [52] Unknown. (1989) Short Communication. *Phytochemistry*, **28**, 2503-2504.
- [53] Kitagawa I, Minagawa, K Zhang RS, Hori K, Doi M, Inoue M, Ishida T, Kimura M, Uji T, Shibuya H. (1993) Dehatrine, an antimalarial bisbenzylisoquinoline alkaloid from the Indonesian medicinal plant *Beilschmiedia madang*, isolated as a mixture of two rotational isomers. *Chemical & Pharmaceutical Bulletin*, **41**, 997-999.
- [54] Matsuura H, Takahashi K, Nabeta K, Yamasaki M, Maede Y, Katakura K. (2007) Screening of Indonesian Medicinal Plant Extracts for Antibabesial Activity and Isolation of New Quassinoids from *Brucea javanica*. *Journal of Natural Products*, **70**, 1654-1657.
- [55] Kitagawa I, Mahmud T, Simanjuntak P, Hori K, Uji T, Shibuya H. (1994) Indonesian medicinal plants. VIII. Chemical structures of three new triterpenoids, bruceajavanin A, dihydrobruceajavanin A, and bruceajavanin B, and a new alkaloidal glycoside, bruceacanthinoside, from the stems of *Brucea javanica* (Simaroubaceae). *Chemical & Pharmaceutical Bulletin*, **42**, 1416-1421.
- [56] Anderson LA, Harris A, Phillipson JD. (1983) Production of cytotoxic canthin-6-one alkaloids by *Ailanthus altissima* plant cell cultures. *Journal of Natural Products*, **46**, 374-378.
- [57] Lassak EV, Polonsky J, Jacquemin H. (1977) 5-Hydroxycanthin-6-one from *Simarouba amara*. *Phytochemistry*, **16**, 1126-1127.
- [58] Fukamiya N, Okano M, Aratani T, Negoro K, McPhail AT, Ju-Ichi M, Lee KH. (1986) Antitumor Agents, 79. Cytotoxic Antileukemic Alkaloids from *Brucea antidysenterica*. *Journal of Natural Products*, **49**, 428-434.
- [59] Hadi S, Bremner JB. (2001) Initial studies on alkaloids from Lombok medicinal plants. *Molecules*, **6**, 117-129.
- [60] Yamauchi T, Abe F. (1998) Regional differences of indole alkaloids in *Alstonia scholaris* and *Alstonia macrophylla*. In *Towards natural medicine research in the 21st century*. Ageta, H, Aimi, N, Ebizuka, Y, Fujita, T, Honda, G (Eds). Elsevier Science & Technology Books, 51-58
- [61] Hadi S. (2002) *Bioactive Alkaloids from Medicinal Plants of Lombok*. PhD thesis, School of Chemistry, University of Wollongong, Wollongong.
- [62] Abe F, Furukawa M, Nonaka G, Okabe H, Nishioka I. (1973) Studies on *Xanthoxylum* spp. I. Constituents of the root of *Xanthoxylum piperitum* DC. *Yakugaku Zasshi*, **93**, 624-628.
- [63] Ishii H, Chen IS, Akaike M, Ishikawa T, Lu ST. (1982) Studies on the chemical constituents of rutaceous plants. XLIV. the chemical constituents of *Xanthoxylum integrifolium* (Merr.) Merr. (*Fagara integrifoliola* Merr.). The constituents of the root wood. *Yakugaku Zasshi*, **102**, 182-188.
- [64] Shoji N, Umeyama A, Takemoto T, Kajiwar A, Ohizumi Y. (1986) Isolation of evodiamine, a powerful cardiotonic principle, from *Evodia rutaecarpa* benthana (Rutaceae). *Journal of Pharmaceutical Sciences*, **75**, 612-613.
- [65] Ren LJ, Xie FZ. (1981) Alkaloids from the root of *Xanthoxylum bangeaum* Maxim. *Yaoxue Xuebao*, **16**, 672-677.
- [66] Shibuya H, Takeda Y, Zhang RS, Tanitame A, Tsai YL, Kitagawa I. (1992) Indonesian medicinal plants. IV. On the constituents of the bark of *Fagara rhetza* (Rutaceae). (2). Lignan glycosides and two apioglucosides. *Chemical & Pharmaceutical Bulletin*, **40**, 2639-2646.
- [67] Shibuya H, Takeda Y, Zhang RS, Tong RX, Kitagawa I. (1992) Indonesian medicinal plants. III. On the constituents of the bark of *Fagara rhetza* (Rutaceae). (1): Alkaloids, phenylpropanoids, and acid amide. *Chemical & Pharmaceutical Bulletin*, **40**, 2325-2330.
- [68] Kardono LBS, Angerhofer CK, Tsauri S, Padmawinata K, Pezzuto JM, Kinghorn AD. (1991) Studies on Indonesian medicinal plants. IV. Cytotoxic and antimalarial constituents of the roots of *Eurycoma longifolia*. *Journal of Natural Products*, **54**, 1360-1367.
- [69] Omar S, Zhang J, MacKinnon S, Leaman DJ, Durst T, Philogene BJR, Arnason JT, Sanches-Vindas PE, Poveda L, Tamez PA, Pezzuto JM. (2003) Traditionally-used antimalarials from Meliaceae. *Current Topics in Medicinal Chemistry*, **3**, 137-139.
- [70] Banskota AH, Attamimi F, Usia T, Linn TZ, Tezuka Y, Kalauni SK, Kadota S. (2003) Novel norcassane-type diterpene from the seed kernels of *Caesalpinia crista*. *Tetrahedron Letters*, **44**, 6879-6882.
- [71] Awale S, Linn TZ, Tezuka Y, Kalauni SK, Banskota AH, Attamimi F, Ueda J, Kadota S. (2006) Constituents of *Caesalpinia crista* from Indonesia. *Chemical & Pharmaceutical Bulletin*, **54**, 213-218.
- [72] Linn TZ, Awale S, Tezuka Y, Banskota AH, Kalauni SK, Attamimi F, Ueda J, Asih PBS, Syafruddin D, Tanaka K, Kadota S. (2005) Cassane- and Norcassane-type diterpenes from *Caesalpinia crista* of Indonesia and their antimalarial activity against the growth of *Plasmodium falciparum*. *Journal of Natural Products*, **68**, 706-710.
- [73] Kitagawa I, Simanjuntak P, Mahmud T, Kobayashi M, Fujii S, Uji T, Shibuya H. (1996) Indonesian medicinal plants. XIII. Chemical structures of caesaldekarins c, d, and e, three additional cassane-type furanoditerpenes from the roots of *Caesalpinia major* (Fabaceae). Several interesting reaction products of caesaldekarin a provided by N-bromosuccinimide treatment. *Chemical & Pharmaceutical Bulletin*, **44**, 1157-1161.
- [74] Kalauni SK, Awale S, Tezuka Y, Banskota AH, Linn TZ, Asih PBS, Syafruddin D, Kadota S. (2006) Antimalarial activity of cassane- and norcassane type diterpenes from *Caesalpinia crista* and their structure-activity relationships. *Biological & Pharmaceutical Bulletin*, **29**, 1050-1052.
- [75] Savarino A, Boelaert JR, Cassone A, Majori G, Cauda R. (2003) Effects of chloroquine on viral infections: an old drug against today's diseases. *The Lancet Infectious Diseases*, **3**, 722-727.
- [76] Kitagawa I, Mahmud T, Yokota K, Nakagawa S, Mayumi T, Kobayashi M, Shibuya H. (1996) Indonesian medicinal plants. XVII. Characterization of quassinoids from the stems of *Quassia indica*. *Chemical & Pharmaceutical Bulletin*, **44**, 2009-2014.

- [77] Sudirga SK. (2004) Pemanfaatan tumbuhan sebagai obat tradisional di Desa Trunyan Kecamatan Kintamani Kabupaten Bangli. *Journal of Enviroment*, **4**, 1-12.
- [78] Goun E, Cunningham G, Chu D, Nguyen C, Miles D. (2003) Antibacterial and antifungal activity of Indonesian ethnomedical plants. *Fitoterapia*, **74**, 592-596.
- [79] Sukari MA, Rashid MA, Tang NY, Rahmani M, Lajis NH, Kalid K, Yusuf UK. (2008) Chemical constituents and bioactivity of *Curcuma xanthorrhiza* roxb. *Journal of Ultra Scientist of Physical Sciences*, **20**, 605-610.
- [80] Kim JE, Kim HE, Hwang JK, Lee HJ, Kwon HK, Kim BI. (2008) Antibacterial characteristics of *Curcuma xanthorrhiza* extract on *Streptococcus mutans* biofilm. *Journal of Microbiology*, **46**, 228-232.
- [81] Zgoda JR, Freyer AJ, Killmer LB, Porter JR. (2001) Polyacetylene Carboxylic Acids from *Mitrephora celebica*. *Journal of Natural Products*, **64**, 1348-1349.
- [82] Zgoda-Pols JR, Freyer AJ, Killmer LB, Porter JR. (2002) Antimicrobial diterpenes from the stem bark of *Mitrephora celebica*. *Fitoterapia*, **73**, 434-438.
- [83] Hadi S, Bremner JB. (2006) Constituents of medicinal plants of lombok: a new antibacterial alkaloid from *Voacanga foetida* (Bl.) Rolfe. *Natural Product Communications*, **1**, 825-829.
- [84] Widyawati Subakir. (2006) *Comparison the effectiveness of 2% Alpinia galanga with 2% ketokonazole in vitro to the growth of Malassezia furfur in Pitriasis versicolor*. Hns thesis, Department of Microbiology, Diponegoro University, Semarang.
- [85] Filet GJ. (1855) *De planten in den botanischen tuin bij het Groot militair hospitaal te Weltevreden: hunne inl. benamingen, groeiplaats en gebruik*. Naar de beste schrijvers en inlandsche opgaven zamengesteld.
- [86] Omar S, Lemonnier B, Jones N, Ficker C, Smith ML, Neema C, Towers GH, Goel K, Arnason, JT. (2000) Antimicrobial activity of extracts of eastern North American hardwood trees and relation to traditional medicine. *Journal of Ethnopharmacology*, **73**, 161-170.
- [87] Ficker CE, Smith ML, Susiarti S, Leaman DJ, Irawati C, Arnason, JT. (2003) Inhibition of human pathogenic fungi by members of Zingiberaceae used by the Kenyah (Indonesian Borneo). *Journal of Ethnopharmacology*, **85**, 289-293.

Natural Product Communications

2011

Volume 6

Natural Product Communications 6 (1-12) 1-1968 (2011)

ISSN 1934-578X (print)

ISSN 1555-9475 (online)

EDITOR-IN-CHIEF**DR. PAWAN K AGRAWAL**

Natural Product Inc.
7963, Anderson Park Lane,
Westerville, Ohio 43081, USA
agrawal@naturalproduct.us

EDITORS**PROFESSOR ALESSANDRA BRACA**

Dipartimento di Chimica Bioorganica e Biofarmacia,
Universita di Pisa,
via Bonanno 33, 56126 Pisa, Italy
braca@farm.unipi.it

PROFESSOR DEAN GUO

State Key Laboratory of Natural and Biomimetic Drugs,
School of Pharmaceutical Sciences,
Peking University,
Beijing 100083, China
gda5958@163.com

PROFESSOR YOSHIHIRO MIMAKI

School of Pharmacy,
Tokyo University of Pharmacy and Life Sciences,
Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan
mimakiy@ps.toyaku.ac.jp

PROFESSOR STEPHEN G. PYNE

Department of Chemistry
University of Wollongong
Wollongong, New South Wales, 2522, Australia
spyne@uow.edu.au

PROFESSOR MANFRED G. REINECKE

Department of Chemistry,
Texas Christian University,
Forts Worth, TX 76129, USA
m.reinecke@tcu.edu

PROFESSOR WILLIAM N. SETZER

Department of Chemistry
The University of Alabama in Huntsville
Huntsville, AL 35809, USA
wsetzer@chemistry.uah.edu

PROFESSOR YASUHIRO TEZUKA

Institute of Natural Medicine
Institute of Natural Medicine, University of Toyama,
2630-Sugitani, Toyama 930-0194, Japan
tezuka@inn.u-toyama.ac.jp

PROFESSOR DAVID E. THURSTON

Department of Pharmaceutical and Biological Chemistry,
The School of Pharmacy,
University of London, 29-39 Brunswick Square,
London WC1N 1AX, UK
david.thurston@pharmacy.ac.uk

HONORARY EDITOR**PROFESSOR GERALD BLUNDEN**

The School of Pharmacy & Biomedical Sciences,
University of Portsmouth,
Portsmouth, PO1 2DT U.K.
axuf64@dsl.pipex.com

ADVISORY BOARD

Prof. Berhanu M. Abegaz
Gaborone, Botswana

Prof. Viqar Uddin Ahmad
Karachi, Pakistan

Prof. Oyvind M. Andersen
Bergen, Norway

Prof. Giovanni Appendino
Novara, Italy

Prof. Yoshinori Asakawa
Tokushima, Japan

Prof. Lee Banting
Portsmouth, U.K.

Prof. Julie Banerji
Kolkata, India

Prof. Alejandro F. Barrero
Granada, Spain

Prof. Anna R. Bilia
Florence, Italy

Prof. Maurizio Bruno
Palermo, Italy

Prof. César A. N. Catalán
Tucumán, Argentina

Prof. Josep Coll
Barcelona, Spain

Prof. Geoffrey Cordell
Chicago, IL, USA

Prof. Cristina Gracia-Viguera
Murcia, Spain

Prof. Duvvuru Gunasekar
Tirupati, India

Prof. A.A. Leslie Gunatilaka
Tucson, AZ, USA

Prof. Kurt Hostettmann
Lausanne, Switzerland

Prof. Martin A. Iglesias Arteaga
Mexico, D. F, Mexico

Prof. Jerzy Jaroszewski
Copenhagen, Denmark

Prof. Leopold Jirovetz
Vienna, Austria

Prof. Karsten Krohn
Paderborn, Germany

Prof. Hartmut Laatsch
Gottingen, Germany

Prof. Marie Lacaille-Dubois
Dijon, France

Prof. Shoei-Sheng Lee
Taipei, Taiwan

Prof. Francisco Macias
Cadiz, Spain

Prof. Imre Mathe
Szeged, Hungary

Prof. Joseph Michael
Johannesburg, South Africa

Prof. Ermino Murano
Trieste, Italy

Prof. M. Soledade C. Pedras
Saskatoon, Canada

Prof. Luc Pieters
Antwerp, Belgium

Prof. Peter Proksch
Düsseldorf, Germany

Prof. Phila Raharivelomanana
Tahiti, French Polynesia

Prof. Monique Simmonds
Richmond, UK

Prof. Valentin Stonik
Vladivostok, Russia

Prof. Winston F. Tinto
Barbados, West Indies

Prof. Karen Valant-Vetschera
Vienna, Austria

Prof. Peter G. Waterman
Lismore, Australia

INFORMATION FOR AUTHORS

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site <http://www.naturalproduct.us>.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the Natural Product Inc. (NPI). A photocopy license is available from the NPI for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

To Subscribe: Natural Product Communications is a journal published monthly. 2011 subscription price: US\$1,995 (Print, ISSN# 1934-578X); US\$1,995 (Web edition, ISSN# 1555-9475); US\$2,495 (Print + single site online); US\$595 (Personal online). Orders should be addressed to Subscription Department, Natural Product Communications, Natural Product Inc., 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

Natural Product Communications

Volume 6 (1-12)

2011

Contents

Number 1

A New Eudesmane Sesquiterpene from <i>Pluchea arguta</i> Nikhat Saba, Rasheeda Khatoon, Viqar Uddin Ahmad and Saleha Suleman Khan	1
Bioactive Diterpenes from <i>Clerodendrum kaichianum</i> Mingfeng Xu, Lianqing Shen, Kuiwu Wang, Qizhen Du and Nan Wang	3
Multi-stage Mass Spectrometric Analysis of Saponins in <i>Glycyrrhiza radix</i> Ken Tanaka, Kosuke Hayashi, Abrar Fahad and Masanori Arita	7
5-Methoxyaristolactam I, the First Natural 5-Substituted Aristolactam from <i>Asarum ichangense</i> Bai-Bo Xie, Ming-Ying Shang, Kuo-Hsiung Lee, Xuan Wang, Katsuko Komatsu and Shao-Qing Cai	11
Flavonoid Aglycones from the Leaf and Stem Exudates of Some Geraniaceae Species Eckhard Wollenweber, Marion Dörr and Matthias Christ	15
Qualitative and Quantitative Analysis of the Major Bioactive Phenolic Compounds of <i>Glechoma longituba</i> by LC-Coupled with PAD and ESI-MS Detection Shu-mao Ni, Da-wei Qian, Jin-ao Duan, Nian-yun Yang and Jian-ming Guo	17
Phenolic Compounds of Mountain Tea from the Balkans: LC/DAD/ESI/MSⁿ Profile and Content Jasmina Petreska, Gjose Stefkov, Svetlana Kulevanova, Kalina Alipieva, Vassya Bankova and Marina Stefova	21
Facile Synthesis of Chrysin-derivatives with Promising Activities as Aromatase Inhibitors Hamdoon A. Mohammed, Lalla A. Ba, Torsten Burkholz, Elena Schumann, Britta Diesel, Josef Zapp, Alexandra K. Kiemer, Christina Ries, Rolf W. Hartmann, Mohammed Hosny and Claus Jacob	31
Anthocyanins from <i>Fuchsia</i> Flowers Monica Jordheim, Irene Skaar, Helene Lunder and Øyvind M. Andersen	35
Oxyresveratrol Protects Against DNA Damage Induced by Photosensitized Riboflavin Manussanunt Chatsumpun, Taksina Chuanasa, Boonchoo Sritularak and Kittisak Likhitwitayawuid	41
Bioactive Isocoumarins from a Terrestrial <i>Streptomyces</i> sp. ANK302 Dhafer Saber Zinad, Khaled A. Shaaban, Muna Ali Abdalla, Md. Tofazzal Islam, Anja Schöffler and Hartmut Laatsch	45
Aromatic Compounds from the Liverwort <i>Conocephalum japonicum</i> Na Liu, Dong-Xiao Guo, Yan-Yan Wang, Li-Ning Wang, Mei Ji and Hong-Xiang Lou	49
New Stress Metabolite from <i>Bulbophyllum kwangtungense</i> Jianbo Chen, Huifang Zhang, Li Chen and Bin Wu	53
<i>In vitro</i> Antioxidant Activities of Maillard Reaction Products Produced in the Steaming Process of <i>Polygonum multiflorum</i> Root Zhenli Liu, Yuanyan Liu, Zhimao Chao, Zhiqian Song, Chun Wang and Aiping Lu	55
Targets of Red Grapes: Oxidative Damage of DNA and Leukaemia Cells Jaouad Anter, Noriluz de Abreu-Abreu, Zahira Fernández-Bedmar, Myriam Villatoro-Pulido, Ángeles Alonso-Moraga and Andrés Muñoz-Serrano	59
Extraction and Identification of Isothiocyanates from Broccoli Seeds Bochao Zhang, Xiaoqin Wang, Yanjing Yang and Xuewu Zhang	65
Authentication of Chinese Crude Drug Gecko by DNA Barcoding Hai-Feng Gu, Yun Xia, Rui Peng, Bang-Hui Mo, Li Li and Xiao-Mao Zeng	67
Comparative Biochemical Characterization of 5'-Phosphodiesterase and Phosphomonoesterase from Barley Malt Sprouts Suncica Beluhan and Vladimir Maric	73
Traditional Medicine in Syria: Folk Medicine in Aleppo Governorate Amal Alachkar, Ahmad Jaddouh, Muhammad Salem Elsheikh, Anna Rita Bilia and Franco Francesco Vincieri	79

Essential Oil Composition of <i>Vismia macrophylla</i> Leaves (Guttiferae) Janne Rojas, Alexis Buitrago, Luis Rojas and Antonio Morales	85
Chemical Variability of Essential Oils in Natural Populations of <i>Cupressus dupreziana</i> Messaoud Ramdani, Takia Lograda, Pierre Chalard, Jean Claude Chalchat and Gilles Figueredo	87
Composition of a Monoterpenoid-rich Essential Oil from the Rhizome of <i>Zingiber officinale</i> from North Western Himalayas Suphla Gupta, Pankaj Pandotra, Gandhi Ram, Rajneesh Anand, Ajai Prakash Gupta, Mohd. Kashif Husain, Yashbir Singh Bedi and Gopal Rao Mallavarapu	93
Chemical Composition of the Essential Oil of <i>Croton gossypifolius</i> from Venezuela Alirica I. Suárez, Marly Oropeza, Luis Vásquez, Stephen Tillett and Reinaldo S. Compagnone	97
Volatile Constituents of <i>Festuca nigrescens</i>, <i>Phleum alpinum</i> and <i>Poa alpina</i> from N.W. Italian Alpine Pastures Aldo Tava, Roberto Cecotti, Maris Grecchi, Luca Falchero, Mauro Coppa and Giampiero Lombardi	101
Comparison of <i>Eucalyptus cinerea</i> Essential Oils Produced by Hydrodistillation and Supercritical Carbon Dioxide Extraction Tavleen S. Mann, Garikapati D. Kiran Babu, Shailja Guleria and Bikram Singh	107
2-Undecanone Rich Leaf Essential Oil from <i>Zanthoxylum armatum</i> Deepa Bisht and Chandan S. Chanotiya	111
Chemical Composition of the Essential Oil of <i>Cachrys libanotis</i> from Algeria Nabila Bouderdara, Abdelhakim Elomri, Lakhdar Djarri, Kamel Medjroubi, Elisabeth Seguin and Philippe Vérité	115
The Essential Oil of <i>Artemisia scoparia</i> from Tajikistan is Dominated by Phenylodiacetylenes Farukh S. Sharopov and William N. Setzer	119
Chemical Composition of Essential Oil of <i>Senecio coinnyi</i>, an Endemic Species of the Central Iberian Peninsula Carlos Arrabal, Felipe Martínez García, María Paz Arraiza and Silvia Guerrero García	123
Chemical Composition of the Essential Oil from <i>Carramboa littlei</i> (Asteraceae) Yndra Cordero de Rojas, Luis B. Rojas and Alfredo Usubillaga	127
Terpenoid Compositions and Antioxidant Activities of Two Indian Valerian Oils from the Khasi Hills of North-east India Jayashankar Das, Ashiho A. Mao and Pratap J. Handique	129
Composition, Antioxidant and Antimicrobial Activities of the Seed Essential Oil of <i>Calocedrus formosana</i> from Taiwan Chen-Lung Ho, Yen-Hsueh Tseng, Eugene I-Chen Wang, Pei-Chun Liao, Ju-Ching Chou, Chien-Nan Lin and Yu-Chang Su	133
Comparison of Volatile Constituents, and Antioxidant and Antibacterial Activities of the Essential Oils of <i>Thymus caucasicus</i>, <i>T. kotschyanus</i> and <i>T. vulgaris</i> Shiva Asbaghian, Ali Shafaghhat, Khalil Zarea, Fakhraddin Kasimov and Farshid Salimi	137
Chemical Composition and Antimicrobial Activity of the Essential Oil of the Leaves of <i>Feronia elephantum</i> (Rutaceae) from North West Karnataka Rajesh K. Joshi, Vijayalaxmi M. Badakar, Sanjiva D. Kholkute and Nayeem Khatib	141
Leaf Essential Oil of <i>Manekia naranjoana</i> (Piperaceae) from Costa Rica and its Cytotoxic Activity Carlos Chaverri, Cecilia Díaz and José F. Cicció	145
<u>Review/Account</u>	
Anthocyanins as Antimicrobial Agents of Natural Plant Origin Agnieszka Cisowska, Dorota Wojnicz and Andrzej B. Hendrich	149

Number 2

Chemical Constituents and Antioxidant Activity of <i>Ficus callosa</i> Phan Van Kiem, Nguyen Xuan Cuong, Nguyen Xuan Nhiem, Dan Thi Thuy Hang, Nguyen Hoai Nam, Ninh Khac Ban, Chau Van Minh, Zhou Bing, Hae Dong Jang and Young Ho Kim	159
Antibacterial and Cytotoxic Activities of the Sesquiterpene Lactones Cnicin and Onopordopicrin Sandra M. Bach, Mario A. Fortuna, Rodgoun Attarian, Juliana T. de Trimarco, César A. N. Catalán, Yossef Av-Gay and Horacio Bach	163
Facile Total Synthesis of Xanthorrhizol Zhenting Du, Yin Wang, Weili Ma, Dong Lv and Hongrui Yu	167
Phlomeoic acid: A New Diterpene from <i>Phlomis bracteosa</i> Javid Hussain, Riaz Ullah, Afsar Khan, Farman Ullah Khan, Zia Muhammad and Mohammad Raza Shah	171

Two Minor Diterpene Glycosides from the Leaves of <i>Stevia rebaudiana</i> Venkata Sai Prakash Chaturvedula, Joshua Rhea, Dennis Milanowski, Ulla Mocek and Indra Prakash	175
Two New Triterpenoids from <i>Zygophyllum eurypterum</i> Shazia Iqbal, Afsar Khan, Viqar Uddin Ahmad, Murad Ali Khan, Sadia Bader, Umar Farooq, Saleha Suleman Khan, Aqib Zahoor and Rasool Bakhsh Tareen	179
Production of Lovastatin by Wild Strains of <i>Aspergillus terreus</i> Ravindra H. Patil, Prakash Krishnan and Vijay L. Maheshwari	183
Chemical Constituents of <i>Lycoris albiflora</i> and their Cytotoxic Activities Maki Jitsuno, Akihito Yokosuka, Ken Hashimoto, Osamu Amano, Hiroshi Sakagami and Yoshihiro Mimaki	187
Antimicrobial Activities of Indole Alkaloids from <i>Tabernaemontana catharinensis</i> Maria Rita Furquini Medeiros, Luiz Afonso de Melo Prado, Vanessa Colnaghi Fernandes, Sérgio Souza Figueiredo, Juliana Coppede, Juliana Martins, Giovana Maria Lanchoti Fiori, Nilce Maria Martinez-Rossi, Rene Oliveira Beleboni, Silvia Helena Taleb Contini, Paulo Sérgio Pereira and Ana Lúcia Fachin	193
Camptothecin Distribution and Content in <i>Nothapodytes nimmoniana</i> Ao Mingzhang, Wang Jing, Shi Yue, Guo Wentao and Yu Longjiang	197
α-Glucosidase Inhibitory Activity of Kaempferol-3-O-rutinoside Solomon Habtemariam	201
Secondary Metabolites from the Roots of <i>Paronychia chionaea</i> Sibel Avunduk, Özgen Alankuş-Çalışkan, Tomofumi Miyamoto, Chiaki Tanaka and Marie-Aleth Lacaille-Dubois	205
Asimafoetidnol: a New Sesquiterpenoid Coumarin from the Gum Resin of <i>Ferula assa-foetida</i> Debasish Bandyopadhyay, Manas Banerjee, Subrata Laskar and Bidyut Basak	209
Determination of Oxygen Radical Absorbance Capacity of Black Cumin (<i>Nigella sativa</i>) Seed Quinone Compounds Hana Tesarova, Blanka Svobodova, Ladislav Kokoska, Petr Marsik, Marie Pribylova, Premysl Landa and Jaroslav Vadlejch	213
Short and Regiospecific Synthesis of Echinamine A – the Pigment of Sea Urchin <i>Scaphechinus mirabilis</i> Nikita S. Polonik, Viktor Ph. Anufriev and Sergey G. Polonik	217
Generation of two New Macrolactones through Sequential Biotransformation of Dihydroresorcyllide Jia Zeng, Jonathan Valiente and Jixun Zhan	223
Amides from the Stem of <i>Capsicum annuum</i> Chung-Yi Chen, Yu-Ting Yeh and Woei-Ling Yang	227
Leishmanicidal and Cytotoxic Activities of Extracts and Naturally-Occurring Compounds from two Lauraceae Species Jeysson Sánchez-Suárez, Ericsson Coy-Barrera, Luis Enrique Cuca and Gabriela Delgado	231
Analysis of the Essential Oil of <i>Amsonia illustris</i> Annamaria London, Katalin Veres, Krisztina Szabó, Erzsébet Háznagy-Radnai and Imre Máthé	235
Secondary Metabolites from <i>Asperula lutea</i> subsp. <i>rigidula</i> Olga Tzakou, Konstantinos Lempesis and Anargyros Loukis	237
Variation in the Volatile Constituents of <i>Artemisia annua</i> var. <i>CIM-Arogya</i> during Plant Ontogeny Rajendra C. Padalia, Ram S. Verma, Amit Chauhan, Chandan S. Chanotiya and Anju Yadav	239
Intraspecific Variation in the Essential Oils from <i>Drimys brasiliensis</i> Leaves and Stem Barks João Henrique G. Lago, Larissa Anastácio C. Carvalho, Marcelo J. P. Ferreira, Paulete Romoff, João Luiz B. Zanin, Marisi G. Soares and Oriana A. Fávero	243
Essential Oil from Leaves of <i>Lantana canescens</i> and <i>L. lopez-palacii</i> Grown in Colombia Andrés F. Peralta-Bohórquez, Clara Quijano-Célis, Mauricio Gaviria, Consuelo Vanegas-López and Jorge A. Pino	247
GC/MS Analysis and Analgesic Effect of the Essential Oil of <i>Matricaria pubescens</i> from Algeria Naima Boutaghane, Ahmed Kabouche, Rachid Touzani, Yousriya Ahmad Maklad, Aida El-Azzouny, Christian Bruneau and Zahia Kabouche	251
Chemical Composition, Antioxidant Properties and Anti-cholinesterase Activity of <i>Cordia gillettii</i> (Boraginaceae) Leaves Essential Oil Marco Bonesi, Philippe N. Okusa, Rosa Tundis, Monica R. Loizzo, Federica Menichini, Caroline Stévigny, Pierre Duez and Francesco Menichini	253
Micromorphology of Glandular hairs, Biological Activity and Composition of the Essential Oil of <i>Tanacetum fisherae</i> (Asteraceae-Anthemideae) from Iran Peyman Rajaei, Taher Nejdassattari, Ali Asghar Maassoumi, Valiollah Mozaffarian and Ali Sonboli	259

Composition and Antimicrobial Activity of <i>Seseli montanum</i> subsp. <i>tommasinii</i> Essential Oil Jovana Šiljegović, Jasmina Glamočlija, Marina Soković, Ivan Vučković, Vele Tešević, Slobodan Milosavljević and Danijela Stešević	263
Composition and Antimicrobial Activities of the Leaf Essential Oil of <i>Machilus kusanoi</i> from Taiwan Chen-Lung Ho, Kuang-Ping Hsu, Yen-Hsueh Tseng, Eugene I-Chen Wang, Pei-Chun Liao, Ju-Ching Chou, Chien-Nan Lin and Yu-Chang Su	267
Chemical Composition and Antimicrobial Activity of <i>Anthriscus nemorosa</i> Root Essential Oil Milica Pavlović, Silvana Petrović, Marina Milenković, Maria Couladis, Olga Tzakou and Marjan Niketić	271
Antimicrobial Activity and Chemical Constituents of the Essential Oils from Flower, Leaf and Stem of <i>Gypsophila bicolor</i> from Iran Ali Shafaghat and Masoud Shafaghatlonbar	275
Antimicrobial Activity of the Essential Oil of Greek Endemic <i>Stachys sprunerii</i> and its Main Component, Isoabienol Aikaterini Koutsaviti, Marina Milenković and Olga Tzakou	277
Comparative Chemical, Cytotoxicity and Antileishmanial Properties of Essential Oils from <i>Chenopodium ambrosioides</i> Lianet Monzote, Marcelina R. Nance, Marley García, Ramón Scull and William N. Setzer	281
<u>Review/Account</u>	
Pharmacological Developments Obtained from Marine Natural Products and Current Pipeline Perspective Elkin Galeano J., Jhon J. Rojas and Alejandro Martínez	287

Number 3

Cytotoxic Bibenzyls, and Germacrane- and Pinguisane-type Sesquiterpenoids from Indonesian, Tahitian and Japanese Liverworts Ismiarni Komala, Takuya Ito, Fumihiro Nagashima, Yasuyuki Yagi and Yoshinori Asakawa	303
Total Synthesis of the Bicyclic Marine Sesquiterpenoid Drechslerine B Hisahiro Hagiwara, Masakazu Fukushima, Kimihiko Kinugawa, Takuya Matsui, Takashi Hoshi and Toshio Suzuki	311
Chemosystematics of <i>Porella</i> (Marchantiophyta, Porellaceae) Agnieszka Ludwiczuk, S. Rob Gradstein, Fumihiro Nagashima and Yoshinori Asakawa	315
Melidianolic Acid A and B, New Antimalarial Acyclic Diterpenes from <i>Aphanamixis grandifolia</i> Adil Astulla, Yusuke Hirasawa, Abdul Rahman, Idha Kusumawati, Wiwied Ekasari, Aty Widyawaruyanti, Noor Cholies Zaini and Hiroshi Morita	323
Eight New Clerodane Diterpenoids from the Bark of <i>Ptychopetalum olacoides</i> Wanxia Tang, Kenichi Harada, Miwa Kubo, Hideaki Hioki and Yoshiyasu Fukuyama	327
First Total Synthesis of the Neo-Clerodane Diterpenoid Salvinorin F Hisahiro Hagiwara, Takashi Nojima, Yuhki Suka, Takashi Hoshi and Toshio Suzuki	333
Physico-chemical Effects of the Major Quassinoids in a Standardized <i>Eurycoma longifolia</i> Extract (Fr 2) on the Bioavailability and Pharmacokinetic Properties, and their Implications for Oral Antimalarial Activity Bin-Seng Low, Chin-Hoe Teh, Kah-Hay Yuen and Kit-Lam Chan	337
Chemical Constituents of <i>Melicope ptelefolia</i> Khozirah Shaari, Seema Zareen, Muhammad Nadeem Akhtar and Nordin Hj. Lajis	343
The Quaternary Indole Alkaloids from Two Sumatran <i>Lerchea</i> species Dayar Arbain, Nova Syafni, Friardi, Deddi Prima Putra, Ismiarni Komala, Kentaro Yamaguchi and Yoshinori Asakawa	349
New Acrylamide and Oxazolidin Derivatives from a Termite-Associated <i>Streptomyces</i> sp. Shu Feng Bi, Fang Li, Yong Chun Song, Ren Xiang Tan and Hui Ming Ge	353
The Coloring Constituents of <i>Scleroderma sinamariense</i> (Sclerodermaceae) Deddi Prima Putra, Nurmilasari, Ismiarni Komala, Yoshinori Asakawa and Dayar Arbain	357
Thiophene, Furans, and Related Aromatic Compounds from <i>Eupatorium heterophyllum</i> Yoshinori Saito, Koji Takiguchi, Xun Gong, Chiaki Kuroda and Motoo Tori	361
Synthesis of a Natural γ-Butyrolactone from Nerylacetone by <i>Acremonium roseum</i> and <i>Fusarium oxysporum</i> Cultures Anna Gliszczynska, Marta Świtalska, Joanna Wietrzyk and Czesław Wawrzęńczyk	367
Constituents of the Green Tea Seeds of <i>Camellia sinensis</i> Toshio Hasegawa, Kensuke Akutsu, Yasuhiro Kishi and Kouji Nakamura	371

Estimation of Cholinesterase Inhibitory and Antioxidant Effects of the Leaf Extracts of Anatolian <i>Ficus carica</i> var. <i>domestica</i> and their Total Phenol and Flavonoid Contents Ilkay Erdogan Orhan, Osman Üstün and Bilge Şener	375
Identification and Quantitative Determination of Lignans in <i>Cedrus atlantica</i> Resins using ^{13}C NMR Spectroscopy Anne-Marie Nam, Mathieu Paoli, Vincent Castola, Joseph Casanova and Ange Bighelli	379
Structure-Molluscicidal Activity Relationships of Acylphloroglucinols from Ferns Cecilia Socolsky, Susana Borkosky and Alicia Bardón	387
Syntheses of Marchantins C, O and P as Promising Highly Bioactive Compounds Andreas Speicher, Judith Holz and Alexandra Hoffmann	393
Characteristic Volatile Components of Kabosu (<i>Citrus sphaerocarpa</i> Hort. ex Tanaka) Kenichi Tomiyama, Hirokazu Aoki, Takeshi Oikawa, Kazutoshi Sakurai, Yoko Kasahara and Yukihiro Kawakami	403
Essential Oil of <i>Plectranthus tenuicaulis</i> Leaves from Gabon, Source of (R),(E)-6,7-Epoxyocimene. An Unusual Chemical Composition within the Genus <i>Plectranthus</i> Huguette Agnani, Abdelhamid Agrebi, Raphaël Bikanga, Thomas Makani, Jacques Lebibi, Hervé Casabianca, Alain Morère and Chantal Menut	409
Reliable Identification and Quantification of Volatile Components of Sage Essential Oil Using Ultra HRGC Laura Tedone, Ivana Bonaccorsi, Paola Dugo, Antonella Cotroneo, Giovanni Dugo and Luigi Mondello	417
<u>Review/Account</u>	
Molecular Genetic Study on the Anthocyanin Chemotypes of <i>Perilla frutescens</i> var. <i>crispa</i> Mami Yamazaki and Kazuki Saito	423

Number 4

Myrtenal, a Controversial Molecule for the Proper Application of the CIP Sequence Rule for Multiple Bonds L. Gerardo Zepeda, Eleuterio Burgueño-Tapia and Pedro Joseph-Nathan	429
A Facile and Efficient Four-Step Enantioselective Synthesis of (+)-Vernolepin from (+)-Minimolide, the Major Germacranolide of <i>Mikania minima</i> Sandra M. Bach, Fernando R. Díaz, Horacio Bach and César A. N. Catalán	433
Use of the Plant <i>Bellardia trixago</i> for the Enantiospecific Synthesis of Odorant Products Alejandro F. Barrero, M. Mar Herrador, Pilar Arteaga, Alexis Castillo and Alejandro F. Arteaga	439
Botcinolide/Botcinin: Asymmetric Synthesis of the Key Fragments Jacinto Ramírez-Fernández, José Manuel Botubol, Antonio J. Bustillo, Josefina Aleu, Isidro G. Collado and Rosario Hernández-Galán	443
C-6 Regioselective Bromination of Methyl Indolyl-3-acetate Oscar R. Suárez-Castillo, Myriam Meléndez-Rodríguez, Lidia Beiza-Granados, Indira C. Cano-Escudero, Martha S. Morales-Ríos and Pedro Joseph-Nathan	451
Synthesis of Pyrrolidinoindolines from 2-(2-Oxo-3-indolyl)acetates: Scope and Limitations Martha S. Morales-Ríos, Ernesto Rivera-Becerril, Daphne E. González-Juárez, Juan Benjamín García-Vázquez, Joel J. Trujillo-Serrato, Angelina Hernández-Barragán, and Pedro Joseph-Nathan	457
<u>Review/Account</u>	
Biomimetic Cyclization of Geraniol Derivatives, a Useful Tool in the Total Synthesis of Bioactive Monocyclic Terpenoids Valentina Merlini, Marco Luparia, Alessio Porta, Giuseppe Zanoni and Giovanni Vidari	465
Drimenol: A Versatile Synthone for Compounds with <i>trans</i>-Drimane Skeleton Manuel Cortés, Virginia Delgado, Claudio Saitz and Veronica Armstrong	477
Recent Advances in the Synthesis of Sesquiterpenolides from Umbelliferae Francisco M. Guerra, F. Javier Moreno-Dorado, Zacarías D. Jorge and Guillermo M. Massanet	491
Diterpene Lactones with Labdane, Halimane and Clerodane Frameworks Lúcia Silva, Arlindo C. Gomes and Jesus M. L. Rodilla	497
Stereoselective Synthesis of Five Biologically Active, Naturally Occurring Medium and Large Ring Lactones J. Alberto Marco and Miguel Carda	505
Enantioselective Synthesis of Alkaloids from Phenylglycinol-Derived Lactams Mercedes Amat, Núria Llor, Rosa Griera, Maria Pérez and Joan Bosch	515

<i>Aconitum</i> lipo-alkaloids – Semisynthetic Products of the Traditional Medicine Botond Borcsa, Dezső Csupor, Peter Forgo, Ute Widowitz, Rudolf Bauer and Judit Hohmann	527
Synthesis of Pterocarpans Leticia Jiménez-González, Carmen Hernández-Cervantes, Míriam Álvarez-Corral, Manuel Muñoz-Dorado and Ignacio Rodríguez-García	537
Hyaluronan: From Biomimetic to Industrial Business Strategy Erminio Murano, Danilo Perin, Riaz Khan and Massimo Bergamin	555

Number 5

Bioactivity-guided Study of Antiproliferative Activities of <i>Salvia</i> Extracts Gábor Janicsák, István Zupkó, Milena T. Nikolova, Peter Forgo, Andrea Vasas, Imre Máthé, Gerald Blunden and Judit Hohmann	575
Simultaneous Quantification of <i>Panax</i> and <i>Epimedium</i> Species Using Rapid Resolution Liquid Chromatography (RRLC) Jie Ma, Yuan-Chun Ma, Daniel Wang, Fei Fei Hou, Mai Luo, Shane Lu, Dariusz C. Gorecki and Asmita V Patel	581
Radioprotective Properties of Cumaside, a Complex of Triterpene Glycosides from the Sea Cucumber <i>Cucumaria japonica</i> and Cholesterol Dmitry L. Aminin, Tatiana S. Zaporozhets, Pelageya V. Adryashchenko, Sergey A. Avilov, Vladimir I. Kalinin and Valentin A. Stonik	587
Herbicide Activity of Extracts from <i>Ailanthus altissima</i> (Simaroubaceae) Cristiano Pedersini, Massimo Bergamin, Vincent Aroulmoji, Sanzio Baldini, Rodolfo Picchio, Patricia Gutierrez Pesce, Luca Ballarin and Erminio Murano	593
Redox Properties of 8-Quinolinol and Implications for its Mode of Action Vladimir Chobot, Sigrid Drage and Franz Hadacek	597
Analysis of Amaryllidaceae Alkaloids from <i>Chlidanthus fragrans</i> by GC-MS and their Cholinesterase Activity Lucie Cahlíková, Kateřina Macáková, Stanislav Zavadiš, Pavel Jiroš, Lubomír Opletal, Klára Urbanová and Luděk Jahodář	603
Acetylcholinesterase and Butyrylcholinesterase Inhibitory Compounds from <i>Corydalis cava</i> (Fumariaceae) Jakub Chlebek, Kateřina Macáková, Lucie Cahlíková, Milan Kurfürst, Jiří Kuneš and Lubomír Opletal	607
Effect of Ascorbigen and 1'-Methylascorbigen on Disease Resistance of Bean Plants to <i>Uromyces phaseoli</i> György Kátay, Ernő Tyihák and Edith Kátay	611
Maculansins, Cryptic Phytotoxins from Blackleg Fungi M. Soledade C. Pedras and Paulos B. Chumala	617
Unusual Compounds from Exudates of <i>Dionysia diapsifolia</i> and <i>D. gaubae</i> var. <i>megalantha</i> (Primulaceae) Tshering Doma Bhutia, Karin M. Valant-Vetschera, Eberhard Lorbeer and Lothar Brecker	621
A Novel Biflavonyloxymethane from <i>Pongamia pinnata</i> and its Radical Quenching Activity Anindita Ghosh, Suvra Mandal, Avijit Banerji, Manoj Kar, Kalyan Hazra and Julie Banerji	625
The Thai Medicinal Plant <i>Gynura pseudochina</i> var. <i>hispida</i>: Chemical Composition and <i>in vitro</i> NF-κB Inhibitory Activity Nisarath Siriwatanametanon and Michael Heinrich	627
Multiple Beneficial Effects of Resveratrol and their Chemical-Biochemical Basis Ernő Tyihák, Zsuzsa Király-Véghely and Ágnes M. Móricz	631
Simultaneous Quantification of <i>Echinacea</i> Species, <i>Flos Lonicerae</i>, <i>Radix Scutellaria</i> and <i>Fructus Forsythiae</i> Combinations by Rapid Resolution Liquid Chromatography Jie Ma, Yuan-Chun Ma, Carina Cai, Daniel Wang, Fei Fei Hou, Mai Luo, Shane Lu, Dariusz C. Gorecki, Asmita V. Patel, Alice Chen and Peter Jin	639
Rapid Resolution Liquid Chromatography (RRLC) Analysis for Quality Control of <i>Rhodiola rosea</i> Roots and Commercial Standardized Products Yuan-Chun Ma, Xiao-Qiang Wang, FeiFei Hou, Jie Ma, Mai Luo, Shane Lu, Peter Jin, Nelly Terevsky, Alice Chen, Iris Xu, Asmita V. Patel and Dariusz Gorecki	645
Antioxidant Activity of Polyphenols from Green and Toasted Mate Tea Patrícia de Abreu Marques Coentrão, Valéria Laneuville Teixeira and Annibal Duarte Pereira Netto	651
Investigation of the Antibacterial Mechanism of Aflatoxins in the BioArena System Ágnes M. Móricz, Péter G. Ott, Klára H. Otta and Ernő Tyihák	657

Bioactive Aromatic Derivatives from Endophytic Fungus, <i>Cytospora</i> sp. Shan Lu, Siegfried Draeger, Barbara Schulz, Karsten Krohn, Ishtiaq Ahmed, Hidayat Hussain, Yanghua Yi, Ling Li and Wen Zhang	661
Fatty Acids as Chemotaxonomic Markers of Marine Macrophytes from Rio de Janeiro State, Brazil Beatriz Grosso Fleury, Luciana Figueiredo, Márcia Iara Marconi, Valéria Laneuville Teixeira, Aurélio Baird Buarque Ferreira and Angelo C. Pinto	667
Determination of Absolute Stereochemistry of Natural Alicyclic Glycosides by ¹H NMR Spectroscopy without Application of Chiral Reagents – An Indication Tatyana N. Makarieva, Larisa K. Shubina, Alla G. Guzii, Natalya V. Ivanchina, Vladimir A. Denisenko, Shamil S. Afiyatullo, Pavel S. Dmitrenok, Anatoly I. Kalinovskiy and Valentin A. Stonik	673
Lecythomycin, a New Macrolactone Glycoside from the Endophytic Fungus <i>Lecythophora</i> sp. Noor Erma Sugijanto, Arnulf Diesel, Mostafa Rateb, Alexander Pretsch, Selma Gogalic, Noor Cholies Zaini, Rainer Ebel and Gunawan Indrayanto	677
Concept of Biogenic Ferromanganese Crust Formation: Coccoliths as Bio-seeds in Crusts from Central Atlantic Ocean (Senghor Seamount/Cape Verde) Xiaohong Wang, Florian Peine, Alexander Schmidt, Heinz C. Schröder, Matthias Wiens, Ute Schloßmacher and Werner E.G. Müller	679
Antiprotzoal, Antitubercular and Cytotoxic Potential of Cyanobacterial (Blue-Green Algal) Extracts from Ireland Barbara Broniatowska, Andrea Allmendinger, Marcel Kaiser, Damien Montamat-Sicotte, Suzie Hingley-Wilson, Ajit Lalvani, Michael Guiry, Gerald Blunden and Deniz Tasdemir	689
Chemical Diversity of <i>Ziziphora clinopodioides</i>: Composition of the Essential oil of <i>Z. clinopodioides</i> from Tajikistan Farukh S. Sharopov and William N. Setzer	695
Essential Oil Composition and Antibacterial Activity of <i>Vismia baccifera</i> Fruits Collected from Mérida, Venezuela Janne Rojas, Alexis Buitrago, Luis Rojas, Antonio Morales, María Lucena and Shirley Baldovino	699
Essential Oil Composition of Five Collections of <i>Achillea biebersteinii</i> from Central Turkey and their Antifungal and Insecticidal Activity Nurhayat Tabanca, Betül Demirci, İlhan Gürbüz, Fatih Demirci, James J. Becnel, David E. Wedge and Kemal Hüsnü Can Başer	701
<u>Review/Account</u>	
Ecdysterone and its Activity on some Degenerative Diseases Lucie Cahliková, Kateřina Macáková, Jakub Chlebek, Anna Hošťálková, Andrea Kulhánková and Lubomír Opletal	707
Overpressured Layer Chromatography (OPLC) – A Flexible Tool of Analysis and Isolation Emil Mincsovcics and Ernő Tyihák	719
The Quality of Commercially Available Herbal Products Priya Ruparel and Brian Lockwood	733

Number 6

Iridoid Glucosides from <i>Wendlandia tinctoria</i> roots Biswanath Dinda, Sudhan Debnath, Rajarshi Banik, Noriko Sato and Yoshihiro Harigaya	747
Janceolaroside A and Janceoside A, Two New Compounds from the Stems and Roots of <i>Jasminum lanceolarium</i> Lili Lou, Lingfei Han, Dali Meng, Ning Li and Xian Li	749
<i>Chionanthus virginicus</i> L.: Phytochemical Analysis and Quality Control of Herbal Drug and Herbal Preparations Laurent Boyer, Béatrice Baghdikian, Sok-Siya Bun, Khalil Taoubi, Ana Diaz-Lanza, Riad Elias and Evelyne Ollivier	753
Solid State Structure and Absolute Configuration of Filifolinol Acetate Marcelo A. Muñoz, Alejandro Urzúa, Javier Echeverría, Brenda Modak and Pedro Joseph-Nathan	759
A Novel Sesquiterpene Alcohol from <i>Fimetariella rabenhorstii</i>, an Endophytic Fungus of <i>Aquilaria sinensis</i> Mei-Hua Tao, Jian Yan, Xiao-Yi Wei, Dong-Li Li, Wei-Min Zhang and Jian-Wen Tan	763
Lipophilic Components from the Ecuadorian Plant <i>Schistocarpha eupatorioides</i> Gianluca Gilardoni, Solveig Tosi, Giorgio Mellerio, Maria Elena Maldonado, Ximena Chiriboga and Giovanni Vidari	767
Terpenoid Metabolites from <i>Spongia</i> spp. and Their Effects on Nucleic Acid Biosynthesis in Sea Urchin Eggs Ludmila P. Ponomarenko, Natalia A. Terent'eva, Vladimir B. Krasokhin, Anatoliy I. Kalinovskiy and Valery A. Rasskazov	773
Evaluation of <i>ent</i>-Kaurenoic Acid Derivatives for their Anticariogenic Activity Bruna B. de Andrade, Monique R. Moreira, Sérgio R. Ambrosio, Nieve A. J. C. Furtado, Wilson R. Cunha, Vladimir C. G. Heleno, Aline N. Silva, Marília R. Simão, Erilda M. P. da Rocha, Carlos H. G. Martins and Rodrigo C. S. Veneziani	777

Microbial Transformation of Deoxyandrographolide by <i>Alternaria alternata</i> AS 3.4578 Xiu-Lan Xin, Sha Deng, Bao-Jing Zhang, Shan-shan Huang, Yan Tian, Xiao-Chi Ma, Lei An, Xiao-hong Shu, Ji-Hong Yao and Xun Cui	781
Limonoids from Meliaceae with Lipid Peroxidation Inhibitory Activity Robert G. Fowles, Baldwin S. Mootoo, Russel S. Ramsewak, William F. Reynolds and Muraleedharan G. G. Nair	785
Chemical Diversity of Iridal-Type Triterpenes in <i>Iris delavayi</i> Collected in Yunnan Province of China Yu Hasegawa, Xun Gong and Chiaki Kuroda	789
Triterpene Derivatives from <i>Abies spectabilis</i> Leaves of Nepalese Origin Stefano Dall'Acqua, Paola Minesso, Stefano Comai, Bharat Babu Shrestha, Mohan Bikram Gewali, Pramod Kumar Jha and Gabbriella Innocenti	793
Quinone Reductase Inducing Activity of the Dichloromethane/Ethanol Extract of the Roots of <i>Pulsatilla chinensis</i> Dan Wang, Ling Han and Zengjun Guo	799
Cytotoxic Cardenolide and Sterols from <i>Calotropis gigantea</i> Sonia D. Jacinto, Eunice Alexis C. Chun, Anthony Sebastian Montuno, Chien-Chang Shen, Dinah L. Espineli and Consolacion Y. Ragasa	803
Sartoryglabins, Analogs of Ardeemins, from <i>Neosartorya glabra</i> Anake Kijjoa, Sonia Santos, Tida Dethoup, Leka Manoch, Ana Paula Almeida, Maria Helena Vasconcelos, Artur Silva, Luís Gales and Werner Herz	807
Identification by LC-ESI-MS of Flavonoids Responsible for the Antioxidant Properties of <i>Mallotus</i> Species from Vietnam Van N.T. Hong, Céline Rivière, Quang T. Hong, Gabrielle Chataigné, Nam N. Hoai, Bieke Dejaegher, Christophe Tistaert, Thanh N.T. Kim, Kiem P. Van, Yvan Vander Heyden, Minh C. Van and Joëlle Quetin-Leclercq	813
Antiradical and Cytotoxic Activity of Different <i>Helichrysum plicatum</i> Flower Extracts Dubravka Bigović, Katarina Šavikin, Teodora Janković, Nebojša Menković, Gordana Zdunić, Tatjana Stanojković and Zorica Djurić	819
Chemical Composition of Argentinean Propolis Collected in Extreme Regions and its Relation with Antimicrobial and Antioxidant Activities Nancy Vera, Eliana Solorzano, Roxana Ordoñez, Luis Maldonado, Enrique Bedascarrasbure and María I. Isla	823
Tannins of Tamarind Seed Husk: Preparation, Structural Characterization, and Antioxidant Activities Puksiri Sinchaiyakit, Yohji Ezure, Sarin Sriprang, Supakorn Pongbangpho, Nasapon Povichit and Maitree Suttajit	829
Chemical Constituents of the New Endophytic Fungus <i>Mycosphaerella</i> sp. nov. and Their Anti-parasitic Activity Eufemio Moreno, Titto Varughese, Carmenza Spadafora, A. Elizabeth Arnold, Phyllis D. Coley, Thomas A. Kursar, William H. Gerwick and Luis Cubilla-Rios	835
Synthesis of a 4H-anthra[1,2-b]pyran Derivative and its Antimicrobial activity Hidayat Hussain, Shahid Aziz, Barbara Schulz and Karsten Krohn	841
Antioxidant Activity and Total Phenolic Content in Shiitake Mycelial Exudates Weihuan Huang, Joo-Shin Kim and Hau Yin Chung	845
Phenylethyl Cinnamides as Potential α-Glucosidase Inhibitors from the Roots of <i>Solanum melongena</i> Xuchun Liu, Jianguang Luo and Lingyi Kong	851
New Phenylalkanoids from <i>Zingiber officinale</i> Chung-Yi Chen, Yu-Ting Yeh and Woei-Ling Yang	855
Valine and Phenylalanine as Precursors in the Biosynthesis of Alkamides in <i>Acmella radicans</i> Nancy Cortez-Espinosa, Judit A. Aviña-Verduzco, Enrique Ramírez-Chávez, Jorge Molina-Torres and Patricia Ríos-Chávez	857
A Novel Antinociceptive Sulphated Polysaccharide of the Brown Marine Alga <i>Spatoglossum schroederi</i> Wladimir R. L. Farias, Paula Cristina W. C. Lima, Natália Velloso F. C. Rodrigues, Rômulo Celly L. Siqueira, Renata M. F. Amorim, Maria G. Pereira and Ana Maria S. Assreuy	863
Orally Administered <i>Dendrobium officinale</i> and its Polysaccharides Enhance Immune Functions in BALB/c Mice Xiao-Fei Liu, Jun Zhu, Shao-Yang Ge, Lin-Jing Xia, Hai-Ying Yang, Yong-Tao Qian and Fa-Zheng Ren	867
Hypotensive Effects of the <i>Crotalus durissus cascavella</i> Venom: Involvement of NO Janaina S. A. M Evangelista, João J. F. Evangelista, Inez L. Evangelista, Diva M. B. Nojosa, Nilberto R. F. Nascimento, Marcellus H. L. P. Souza, Renata S. Alves, Alice M. C. Martins, Maria Elisabete A. Moraes and Helena S. A. Monteiro	871
Citrus Essential Oil of Nigeria Part V: Volatile Constituents of Sweet Orange Leaf Oil (<i>Citrus sinensis</i>) Adeleke A. Kasali, Oladipupo A. Lawal, Adeolu O. Eshilokun, Abayomi A. Olaniyan, Andy R. Opoku and William N. Setzer	875

Comparison of the Essential Oils of Leaves and Stem Bark from Two Different Populations of <i>Drimys winteri</i> a Chilean Herbal Medicine	
Orlando Muñoz, Philippe Christen, Silvian Cretton, Alejandro F. Barrero, Armando Lara and M. Mar Herrador	879
A Myristicin-rich Essential Oil from <i>Daucus sahariensis</i> Growing in Algeria	
Tahar Smaili, Amar Zellagui, Pier Luigi Cioni and Guido Flamini	883
Activity of <i>Schinus areira</i> (Anacardiaceae) Essential Oils against the Grain Storage Pest <i>Tribolium castaneum</i>	
Lilian R. Descamps, Carolina Sánchez Chopa and Adriana A. Ferrero	887
Acaricidal Activity against <i>Tetranychus urticae</i> and Essential Oil Composition of Four <i>Croton</i> Species from Caatinga Biome in Northeastern Brazil	
Ilzenayde Araújo Neves and Claudio Augusto Gomes da Camara	893
Composition, Antioxidant and Antimicrobial Activities of Leaf and Twig Essential Oils of <i>Litsea akoensis</i> from Taiwan	
Chen-Lung Ho, Chai-Yi Lin, Eugene I-Chen Wang and Yu-Chang Su	901
Composition and Antimicrobial Activity of <i>Achillea distans</i> Essential Oil	
Angel Konakchiev, Milka Todorova, Bozhanka Mikhova, Antonina Vitkova and Hristo Najdenski	905
Chemical Composition and Antioxidant, Antimicrobial, and Larvicidal Activities of the Essential Oils of <i>Annona salzmännii</i> and <i>A. pickelii</i> (Annonaceae)	
Emmanuel Vilaça Costa, Livia Macedo Dutra, Hugo César Ramos de Jesus, Paulo Cesar de Lima Nogueira, Valéria Regina de Souza Moraes, Marcos José Salvador, Sócrates Cabral de Holanda Cavalcanti, Roseli La Corte dos Santos and Ana Paula do Nascimento Prata	907
<u>Review/Account</u>	
Chemical and Pharmacological Aspects of the Genus <i>Cecropia</i>	
Geison M. Costa, Eloir P. Schenkel and Flávio H. Reginatto	913
<u>Letter to Editor</u>	
Does <i>Pseudomonas fluorescens</i> produce <i>N</i>-mercapto-4-formylcarbostyryl?	
Carsten Christophersen and Ole Hammerich	921
<u>Additions/Corrections</u>	923

Number 7

Use of Dimethyldioxirane in the Epoxidation of the Main Constituents of the Essential Oils Obtained from <i>Tagetes lucida</i>, <i>Cymbopogon citratus</i>, <i>Lippia alba</i> and <i>Eucalyptus citriodora</i>	
Luz A. Veloza, Lina M. Orozco and Juan C. Sepúlveda-Arias	925
Validation of the Ethnopharmacological Use of <i>Polygonum persicaria</i> for its Antifungal Properties	
Marcos Derita and Susana Zacchino	931
On the Isomerization of <i>ent</i>-Kaurenic Acid	
Julio Rojas, Rosa Aparicio, Thayded Villasmil, Alexis Peña and Alfredo Usubillaga	935
Aristolactams from Roots of <i>Ottonia anisum</i> (Piperaceae)	
André M. Marques, Leosvaldo S. M. Velozo, Davyson de L. Moreira, Elsie F. Guimarães and Maria Auxiliadora C. Kaplan	939
Anti-angiogenic Activity Evaluation of Secondary Metabolites from <i>Calycolpus moritzianus</i> Leaves	
Laura Lepore, Maria J. Gualtieri, Nicola Malafronte, Roberta Cotugno, Fabrizio Dal Piaz, Letizia Ambrosio, Sandro De Falco and Nunziatina De Tommasi	943
Chemical and Biological Activity of Leaf Extracts of <i>Chromolaena leivensis</i>	
Ruben D. Torrenegra G. and Oscar E. Rodríguez A.	947
<i>Citrus bergamia</i> Juice: Phytochemical and Technological Studies	
Patrizia Picerno, Francesca Sansone, Teresa Mencherini, Lucia Prota, Rita Patrizia Aquino, Luca Rastrelli and Maria Rosaria Lauro	951
Phenolic Derivatives from the Leaves of <i>Martinella obovata</i> (Bignoniaceae)	
Carolina Arevalo, Ines Ruiz, Anna Lisa Piccinelli, Luca Campone and Luca Rastrelli	957
Phenolic Chemical Composition of <i>Petroselinum crispum</i> Extract and Its Effect on Haemostasis	
Douglas S. A. Chaves, Flávia S. Frattani, Mariane Assafim, Ana Paula de Almeida, Russolina B. Zingali and Sônia S. Costa	961
Bioactivities of <i>Chusqueira straminea</i> Sandwith	
María Elena Mendiondo, Berta E. Juárez, Catiana Zampini, María Inés Isla and Roxana Ordoñez	965

Free Radical Scavenging Activity, Determination of Phenolic Compounds and HPLC-DAD/ESI-MS Profile of <i>Campomanesia adamantium</i> Leaves Aislan C.R.F. Pascoal, Carlos Augusto Ehrenfried, Marcos N. Eberlin, Maria Élide Alves Stefanello and Marcos José Salvador	969
Activity of Cuban Propolis Extracts on <i>Leishmania amazonensis</i> and <i>Trichomonas vaginalis</i> Lianet Monzote Fidalgo, Idalia Sariego Ramos, Marley García Parra, Osmany Cuesta-Rubio, Ingrid Márquez Hernández, Mercedes Campo Fernández, Anna Lisa Piccinelli and Luca Rastrelli	973
Antioxidant Capacity and Phenolic Content of four Myrtaceae Plants of the South of Brazil Marcos José Salvador, Caroline C. de Lourenço, Nathalia Luiza Andreazza, Aislan C.R.F. Pascoal and Maria Élide Alves Stefanello	977
Cytotoxicity of Active Ingredients Extracted from Plants of the Brazilian “Cerrado” Veronica CG Soares, Cibele Bonacorsi, Alana LB Andrela, Lígia V Bortoloti, Stepheny C de Campos, Fábio HR Fagundes, Márcio Piovani, Camila A Cotrim, Wagner Vilegas and Marcos H Toyama	983
Propagation and Conservation of Native Forest Genetic Resources of Medicinal Use by Means of <i>in vitro</i> and <i>ex vitro</i> Techniques Sandra Sharry, Marina Adema, María A. Basiglio Cordal, Blanca Villarreal, Noelia Nikoloff, Valentina Briones and Walter Abedini	985
Genotoxic Evaluation of a Methanolic Extract of <i>Verbascum thapsus</i> using Micronucleus Test in Mouse Bone Marrow Franco Matías Escobar, María Carola Sabini, Silvia Matilde Zanon, Laura Noelia Cariddi, Carlos Eugenio Tonn and Liliana Inés Sabini	989
Study of Antiviral and Virucidal Activities of Aqueous Extract of <i>Baccharis articulata</i> against <i>Herpes suis</i> virus Cristina Vanesa Torres, María Julia Dominguez, José Luis Carbonari, María Carola Sabini, Liliana Inés Sabini and Silvia Matilde Zanon	993
Evaluation of Cytogenotoxic Effects of Cold Aqueous Extract from <i>Achyrocline satureioides</i> by <i>Allium cepa</i> L test María C. Sabini, Laura N. Cariddi, Franco M. Escobar, Romina A. Bachetti, Sonia B. Sutil, Marta S. Contigiani, Silvia M. Zanon and Liliana I. Sabini	995
Toxic Plants Used in Ethnoveterinary Medicine in Italy Lucia Viegi and Roberta Vangelisti	999
Diagnosis of Public Programs focused on Herbal Medicines in Brazil Ely Eduardo Saranz Camargo, Mary Anne Medeiros Bandeira and Anselmo Gomes de Oliveira	1001
Identification of Thiosildenafil in a Health Supplement Marcello Nicoletti	1003
Hypolipidemic Effect of Seed Oil of Noni (<i>Morinda citrifolia</i>) Diana C. Pazos, Fabiola E. Jiménez, Leticia Garduño, V. Eric López and M. Carmen Cruz	1005
Composition of Egyptian Neroli Oil Ivana Bonaccorsi, Danilo Sciarbone, Luisa Schipilliti, Alessandra Trozzi, Hussein A. Fakhry and Giovanni Dugo	1009
Essential oil of <i>Nepeta x faassenii</i> Bergmans ex Stearn (<i>N. mussinii</i> Spreng. x <i>N. nepetella</i> L.): A Comparison Study Niko Radulović, Polina D. Blagojević, Kevin Rabbitt and Fabio de Sousa Menezes	1015
Chemical Composition and Biological Activity of <i>Salvia verbenaca</i> Essential Oil Marisa Canzoneri, Maurizio Bruno, Sergio Rosselli, Alessandra Russo, Venera Cardile, Carmen Formisano, Daniela Rigano and Felice Senatore	1023
Chemical Composition and Antimicrobial Activities of the Essential Oils from <i>Ocimum selloi</i> and <i>Hesperozygis myrtoides</i> Márcia G. Martini, Humberto R. Bizzo, Davyson de L. Moreira, Paulo M. Neufeld, Simone N. Miranda, Celuta S. Alviano, Daniela S. Alviano and Suzana G. Leitão	1027
Chemical Composition and Antibacterial Activity of the Essential Oil of <i>Lantana camara</i> var. <i>moritziana</i> Nurby Rios Tesch, Flor Mora, Luis Rojas, Tulia Díaz, Judith Velasco, Carlos Yáñez, Nahile Rios, Juan Carmona and Sara Pasquale	1031
Activity against <i>Streptococcus pneumoniae</i> of the Essential Oil and 5-(3-Buten-1-ynyl)-2, 2'-bithienyl Isolated from <i>Chrysactinia mexicana</i> Roots Bárbara Missiam Mezari Guevara Campos, Anabel Torres Cirio, Verónica Mayela Rivas Galindo, Ricardo Salazar Aranda, Noemí Waksman de Torres and Luis Alejandro Pérez-López	1035
Antimycotic Effect of the Essential Oil of <i>Aloysia triphylla</i> against <i>Candida</i> Species Obtained from Human Pathologies María de las Mercedes Oliva, María Evangelina Carezzano, Mauro Nicolás Gallucci and Mirta Susana Demo	1039

Secretory Cavities and Volatiles of *Myrrhinium atropurpureum* Schott var. *atropurpureum* (Myrtaceae): An Endemic Species Collected in the Restingas of Rio de Janeiro, Brazil

Cristiane Pimentel Victório, Claudio B. Moreira, Marcelo da Costa Souza, Alice Sato and Rosani do Carmo de Oliveira Arruda 1045

Chemical Composition and *in vitro* Antibacterial Activity of the Essential Oil of *Phthirusa adunca* from Venezuelan Andes

Flor D. Mora, Nurby Ríos, Luis B. Rojas, Tulia Díaz, Judith Velasco, Juan Carmona A and Bladimiro Silva 1051

Number 8

Two New Sesquiterpene Lactones from *Ixeris sonchifolia*

Shao-jiang Song, Ling-yan Zhou, Ling-zhi Li, Pin-yi Gao, Wei-wei Jia and Ying Peng 1055

Additional Minor Diterpene Glycosides from *Stevia rebaudiana*

Venkata Sai Prakash Chaturvedula and Indra Prakash 1059

New Virescenosides from the Marine-derived Fungus *Acremonium striatisporum*

Shamil Sh. Afyatullov, Anatoly I. Kalinovsky and Alexandr S. Antonov 1063

New Clerodane Diterpenoid from the Bulbils of *Dioscorea bulbifera*

Kanlaya Kidyu, Haruthai Thaisuchat, Puttinan Meepowpan, Sukee Sukdee, Narong Nuntasaeen, Sittiporn Punyanitya and Wilart Pompimon 1069

Gastroprotective Activity of Epitaondiol and Sargaol

Carlos Areche, Aurelio San-Martín, Juana Roviroso and Beatriz Sepúlveda 1073

Structure of Cucumariosides H₅, H₆, H₇ and H₈, Triterpene Glycosides from the Sea Cucumber *Eupentacta fraudatrix* and Unprecedented Aglycone with 16,22-Epoxy-group

Alexandra S. Silchenko, Anatoly I. Kalinovsky, Sergey A. Avilov, Pelageya V. Andryjaschenko, Pavel S. Dmitrenok, Ekaterina A. Yurchenko and Vladimir I. Kalinin 1075

Pregnane Derivatives from *Potentilla evestita*

Rehan Khan, Farah Siddiq, Itrat Fatima, Shazia Yasmeen, Aman Karim, Abdul Malik, Nighat Afza and Saira Hameed 1083

Insect Growth Regulatory Activity of *Blechnum chilense*

Carlos A. Hincapié L., Zulma Monsalve F., Katherine Parada, Claudio Lamilla, Julio Alarcón, Carlos L. Céspedes A. and David Seigler 1085

The Therapeutic Potential of *Berberis darwinii* Stem-Bark: Quantification of Berberine and *In Vitro* Evidence for Alzheimer's Disease Therapy

Solomon Habtemariam 1089

A Set of Two Diastereomers of Cyanogenic Glycosides from *Passiflora quadrangularis*

Daisuke Saeki, Takeshi Yamada, Tetsuya Kajimoto, Osamu Muraoka and Reiko Tanaka 1091

Inhibition on HIV-1 Integrase Activity and Nitric Oxide Production of Compounds from *Ficus glomerata*

Kingkan Bunluepuech, Teeratad Sudsai, Chatchai Wattanapiromsakul and Supinya Tewtrakul 1095

Two New Prenylflavanones from *Erythrina sigmoidea*

Muhammad Shaiq Ali, Muhammad Imran Ali, Zeeshan Ahmed and Patricia A. Onocha 1099

Prenylated Flavonoids from the Leaves of *Derris malaccensis* and their Cytotoxicity

Darane Chokchaichamnankit, Vorawan Kongjinda, Nisachon Khunnawutmanotham, Nitirat Chimnoi, Somchai Pisutcharoenpong and Supanna Techasakul 1103

Content of Phenolic Compounds in Aerial Parts of *Chamomilla suaveolens* from Estonia

Ain Raal, Tõnu Püssa, Janne Sepp, Birgit Malmiste and Elmar Arak 1107

Biflavonoids from the Roots of *Wikstroemia indica*

Xiaoli Zhang, Guocai Wang, Weihuan Huang, Wencai Ye and Yaolan Li 1111

Drypetdimer A: A New Flavone Dimer from *Drypetes gerrardii*

Margaret Mwihaki Ng'ang'a, Hidayat Hussain, Sumesh Chhabra, Caroline Langat-Thoruwa, Muhammad Riaz and Karsten Krohn 1115

Chemical Constituents of *Cichorium intybus* and their Inhibitory Effects against Urease and α -Chymotrypsin Enzymes

Sumayya Saied, Shazia Shah, Zulfiqar Ali, Ajmal Khan, Bishnu P. Marasini and Muhammad Iqbal Choudhary 1117

Antimicrobial Activity and Cytotoxic Effects of *Magnolia dealbata* and its Active Compounds

Maria del Rosario Jacobo-Salcedo, Luis Angel Gonzalez-Espindola, Angel Josabad Alonso-Castro, Marisela del Rocio Gonzalez-Martinez, Fabiola Domínguez and Alejandro Garcia-Carranca 1121

DPPH Radical Scavenging and Xanthine Oxidase Inhibitory Activity of <i>Terminalia macroptera</i> Leaves Anh Thu Pham, Karl Egil Malterud, Berit Smestad Paulsen, Drissa Diallo and Helle Wangensteen	1125
Bioactive Metabolites from Biotransformation of Paeonol by the White-Rot Basidiomycete <i>Coriolus versicolor</i> Xiao-Jun Li, Xin-Wei Shi, Qi Shuai, Jin-Ming Gao and An-Ling Zhang	1129
Two Compounds from the Endophytic <i>Colletotrichum</i> sp. of <i>Ginkgo biloba</i> Sheng-Liang Zhou, Song-Lin Zhou, Mei-Xia Wang and Shuang-Lin Chen	1131
Two New Alkylanacardic Acids, Ozorcardic A and B, from <i>Ozoroa pulcherrima</i> Tsague Dongmo Christelle, Hidayat Hussain, Etienne Dongo, Jatsa-Megaptche Boukeng Hermine, Ishtiaq Ahmed and Karsten Krohn	1133
Cordioxime: A New Dioxime γ-Lactam from <i>Cordia platythyrsa</i> Tsague Dongmo Christelle, Hidayat Hussain, Etienne Dongo, Oben Enyong Julius and Javid Hussain	1135
Biosynthesis, Characterization and Biological Evaluation of Fe(III) and Cu(II) Complexes of Neoaspergillilic Acid, a Hydroxamate Siderophore Produced by Co-cultures of two Marine-derived Mangrove Epiphytic Fungi Feng Zhu, Jingshu Wu, Guangying Chen, Weihong Lu and Jiahui Pan	1137
Epoxidation of Soybean Oil Catalyzed by $[\pi\text{-C}_5\text{H}_5\text{NC}_{16}\text{H}_{33}]_3[\text{PW}_4\text{O}_{16}]$ with Hydrogen Peroxide and Ethyl Acetate as Solvent Shuang-Fei Cai and Li-Sheng Wang	1141
GC/MS Analysis of the Essential Oil of <i>Senecio belgaumensis</i> Flowers Rajesh K. Joshi	1145
Composition of the Essential Oils from <i>Anthriscus cerefolium</i> var. <i>trichocarpa</i> and <i>A. caucalis</i> Growing Wild in the Urban Area of Vienna (Austria) Remigius Chizzola	1147
Chemical Composition of the Essential Oil of <i>Pituranthos scoparius</i> Nadhir Gourine, Bahia Merrad, Mohamed Yousfi, Pierre Stocker and Emile M. Gaydou	1151
Characterization of Volatile Components of Tea Flowers (<i>Camellia sinensis</i>) Growing in Kangra by GC/MS Robin Joshi, Poonam, Rikki Saini, Shailja Guleria, Garikapati D. Kiran Babu, Manisha Kumari and Ashu Gulati	1155
Susceptibility of the Multi-drug Resistant Strain of <i>Enterobacter aerogenes</i> EA289 to the Terpene Alcohols from <i>Cistus ladaniferus</i> Essential Oil Elodie Guinoiseau, Vannina Lorenzi, Anne Luciani, Félix Tomi, Joseph Casanova and Liliane Berti	1159
Composition and Antimicrobial Activity of <i>Seseli globiferum</i> Essential Oil Peda Janačković, Marina Soković, Ljubodrag Vujisić, Vlatka Vajs, Ivan Vučković, Zoran Krivošej and Petar D. Marin	1163
Chemical Composition and Antimicrobial activity of <i>Satureja kitaibelii</i> Essential Oil against Pathogenic Microbial Strains Tatjana Mihajilov-Krstev, Dušanka Kitić, Dragan Radnović, Mihajlo Ristić, Mira Mihajlović-Ukropina and Bojan Zlatković	1167
Influence of Growth Phase on the Essential Oil Composition and Antimicrobial Activities of <i>Satureja hortensis</i> Mohammad Jamal Saharkhiz, Kamiar Zomorodian, Mohammad Reza Rezaei, Farshid Saadat and Mohammad Javad Rahimi	1173
Chemical Composition and Antioxidant Activities of the Essential Oil from <i>Tornabenea bischoffii</i> (Apiaceae) Risoleta Ortet, Erik L. Regalado, Olivier P. Thomas, Jorge A. Pino and Miguel D. Fernández	1179
Chemical Composition and Insecticidal Activity of Essential oils of two Aromatic plants from Ivory Coast against <i>Bemisia tabaci</i> G. (Hemiptera: Aleyrodidae) Etienne V. Tia, Augustin A. Adima, Sébastien L. Niamké, Gnago A. Jean, Thibaud Martin, Paul Lozano and Chantal Menut	1183
Influence of Viral Infection on Essential Oil Composition of <i>Ocimum basilicum</i> (Lamiaceae) Alice Nagai, Lígia M.L. Duarte and Déborah Y.A.C. Santos	1189
<i>Neolitsea aciculata</i> Essential Oil Inhibits Drug-Resistant Skin Pathogen Growth and <i>Propionibacterium acnes</i>-Induced Inflammatory Effects of Human Monocyte Leukemia Sang Suk Kim, Jung Eun Kim, Chang-Gu Hyun and Nam Ho Lee	1193
Aroma-therapeutic Effects of Massage Blended Essential Oils on Humans Tapanee Hongratanaworakit	1199
<u>Review/Account</u>	
Biological Activity of Bicyclic and Tricyclic Diterpenoids from <i>Salvia</i> Species of Immediate Pharmacological and Pharmaceutical Interest Maria Carmela Bonito, Carla Cicala, Maria Carla Marcotullio, Francesco Maione and Nicola Mascolo	1205

Number 9

Analysis of Car-3-en-5-hydroperoxide Nicole Lehnert, Ulrich Krings and Ralf G. Berger	1217
Antibacterial Potential of Citral Derivatives Soni A. Singh, Yogesh A. Potdar, Rasika S. Pawar and Sujata V. Bhat	1221
A New Bisabolene from <i>Stevia tomentosa</i> Alejandro Valdez-Calderón, J. Martín Torres-Valencia, J. Jesús Manríquez-Torres, René Velázquez-Jiménez, Mario A. Gómez-Hurtado, Luisa U. Román-Marín, Juan D. Hernández-Hernández, Carlos M. Cerda-García-Rojas and Pedro Joseph-Nathan	1225
Free Radical Scavenging Activity-Guided Isolation of a Diterpenoid from <i>Plectranthus punuctatus</i> Wossen Kebede, Daniel Bisrat and Kaleab Asres	1229
Components from the Steamed Leaves of <i>Acanthopanax koreanum</i> and their Effects on PPAR Activity in HepG2 Cells Jeong Ah Kim, Seok Bean Song, Seo Young Yang and Young Ho Kim	1233
Isolation and X-ray Structure of Deoxycholic Acid from the Sponge <i>Ircinia</i> sp. Keisham Sarjit Singh and Werner Kaminsky	1237
Chemical Constituents of the Gorgonian <i>Dichotella fragilis</i> (Ridleg) from the South China Sea Yuan-Ming Zhou, Chang-Lun Shao, Chang-Yun Wang, Hui Huang, Ying Xu and Pei-Yuan Qian	1239
A New Pyrrolidine Derivative and Steroids from an Algicolous <i>Gibberella zeae</i> Strain Xiang-Hong Liu, Xiao-Zhen Tang, Feng-Ping Miao and Nai-Yun Ji	1243
Szentiamide, an <i>N</i>-formylated Cyclic Depsipeptide from <i>Xenorhabdus szentirmaii</i> DSM 16338^T Birgit Ohlendorf, Sven Simon, Jutta Wiese and Johannes F. Imhoff	1247
Bioactive Constituents from <i>Michelia champaca</i> Yu-Ting Yeh, Jin-Cherng Huang, Po-Lin Kuo and Chung-Yi Chen	1251
Inhibition of Gastric H⁺, K⁺-ATPase Activity by Compounds from Medicinal Plants Cristina Setim Freitas, Cristiane Hatsuko Baggio, Bárbara Mayer, Ana Cristina dos Santos, André Twardowschy, Cid Aimbiré de Moraes Santos and Maria Consuelo Andrade Marques	1253
GC/MS Analysis of Three Amaryllidaceae Species and Their Cholinesterase Activity Lucie Cahlíková, Nina Benešová, Kateřina Macáková, Klára Urbanová and Lubomír Opletal	1255
Astrotricoumarin, an antiproliferative 4'-hydroxy-2',3'-dihydroprenylated methylcoumarin from an <i>Astrotrichilia</i> sp. from the Madagascar dry forest Liva Harinantenaina, Peggy J. Brodie, Martin W. Callmander, Richard Randrianaivo, Stephan Rakotonandrasana, Vincent E. Rasamison, Etienne Rakotobe and David G. I. Kingston	1259
A New Chromene Isolated from <i>Ageratum conyzoides</i> Abiodun Humphrey Adebayo, Chang-Jiu Ji, Yu-Mei Zhang, Wen-Jun He, Guang-Zhi Zeng, Hong-Jin Han, Jun-Ju Xu, Afolabi Akintunde Akindahunsi and Ning-Hua Tan	1263
Antifibrotic Constituents from <i>Garcinia mangostana</i> Young-Won Chin, Eunjin Shin, Bang Yeon Hwang and Mi Kyeong Lee	1267
Antioxidant and Antimutagenic Polyisoprenylated Benzophenones and Xanthenes from <i>Rheedia acuminata</i> Giovanna R. Almanza, Raúl Quispe, Patricia Mollinedo, Gloria Rodrigo, Odette Fukushima, Rodrigo Villagomez, Bjorn Akesson and Olov Sterner	1269
Antraquinone Profile, Antioxidant and Antimicrobial Properties of Bark Extracts of <i>Rhamnus catharticus</i> and <i>R. orbiculatus</i> Marcello Locatelli, Francesco Epifano, Salvatore Genovese, Giuseppe Carlucci, Marijana Zovko Končić, Ivan Kosalec and Dario Kremer	1275
Quantitative Analysis of Euglobals in <i>Eucalyptus loxophleba</i> Leaves by qNMR Jasmeen Sidana, William J. Foley and Inder Pal Singh	1281
Evaluation of Antioxidant Activity of Isoferulic Acid <i>in vitro</i> Xiaozhen Wang, Xican Li and Dongfeng Chen	1285
Determination of Caffeoyl Quinic Acids and Flavonoids in <i>Acanthopanax trifoliatum</i> Leaves by HPLC Pongtip Sithisarn, Sarinthip Muensaen and Siripen Jarikasem	1289
Amides and Esters of Phenylpropenoic Acids from the Aerial Parts of <i>Trifolium pallidum</i> Barbara Szajwaj, Jaroslaw Moldoch, Milena Masullo, Sonia Piacente, Wieslaw Oleszek and Anna Stochmal	1293
Amides from the Stems of <i>Cinnamomum burmannii</i> Zi-Ling Hong, Jin-Cherng Huang, Soong-Yu Kuo and Chung-Yi Chen	1297

Phenolic Composition, Antioxidant Capacity and Antibacterial Activity of Selected Irish Brassica Vegetables Amit Kumar Jaiswal, Gaurav Rajauria, Nissreen Abu-Ghannam and Shilpi Gupta	1299
Dietary Burden of Phenolics Per Serving of “Mountain Tea” (<i>Sideritis</i>) from Macedonia and Correlation to Antioxidant Activity Jasmina Petreska, Marina Stefova, Federico Ferreres, Diego. A. Moreno, Francisco. A. Tomás-Barberán, Gjose Stefkov, Svetlana Kulevanova and Angel Gil-Izquierdo	1305
Aqueous Extract from <i>Vitis vinifera</i> Tendrils is Able to Enrich Keratinocyte Antioxidant Defences Daniele Fraternale, Roberta De Bellis, Cinzia Calcabrini, Lucia Potenza, Luigi Cucchiari, Umberto Mancini, Marina Dachà and Donata Ricci	1315
A New Acylated Neohesperidoside from <i>Geranium purpureum</i> Didem Şöhretoğlu, Tibor Liptaj, M. Koray Sakar and Olov Sterner	1321
Synthesis and Field Test of Three Candidates for Soybean Pod Borer’s Sex Pheromone Tao Zhang, Juntao Feng, Chonglin Cai and Xing Zhang	1323
Chemical Investigation of Carrageenan from the Red alga <i>Sarconema filiforme</i> (Gigartinales, Rhodophyta) of Indian Waters Sanjay Kumar, Gaurav K Mehta, Kamalesh Prasad, Ramavtar Meena and Arup K Siddhanta	1327
Monitoring the Emission of Volatile Organic Compounds from Flowers of <i>Jasminum sambac</i> Using Solid-Phase Micro-extraction Fibers and Gas Chromatography with Mass Spectrometry Detection VPPalayam Shanmugam Pragadheesh, Anju Yadav, Chandan Singh Chanotiya, Prasanta Kumar Rout and Girish Chandra Uniyal	1333
Volatile Components from Aerial parts of <i>Centaurea gracilentia</i> and <i>C. ovina</i> ssp. <i>besserana</i> Growing Wild in Bulgaria Carmen Formisano, Daniela Rigano, Felice Senatore, Svetlana Bancheva, Maurizio Bruno, Antonella Maggio and Sergio Rosselli	1339
Variability of Essential Oils of <i>Betonica officinalis</i> (Lamiaceae) from Different Wild Populations in Kosovo Avni Hajdari, Behxhet Mustafa, Chlodwig Franz and Johannes Novak	1343
Analysis of Essential Oils from <i>Scutellaria orientalis</i> ssp. <i>alpina</i> and <i>S. utriculata</i> by GC and GC-MS Carmen Formisano, Daniela Rigano, Felice Senatore, Franco Piozzi and Nelly Apostolides Arnold	1347
Antibacterial Activity and GC/MS Analysis of the Essential Oils from Flower, Leaf and Stem of <i>Origanum vulgare</i> ssp. <i>viride</i> Growing Wild in North-west Iran Ali Shafaghat	1351
Composition of <i>Satureja kitaibelii</i> Essential Oil and its Antimicrobial Activity Tatjana Kundaković, Marina Milenković, Saša Zlatković, Nada Kovačević and Nikolić Goran	1353
Composition and Antifungal Activities of the Leaf Essential Oil of <i>Neolitsea parvigemma</i> from Taiwan Chen-Lung Ho, Pei-Chun Liao, Eugene I-Chen Wang and Yu-Chang Su	1357
Antioxidant Capacity and Larvicidal and Antifungal Activities of Essential Oils and Extracts from <i>Piper krukoffii</i> Joyce Kelly R. da Silva, Eloisa Helena A. Andrade, Massuo J. Kato, Léa Maria M. Carreira, Elsie F Guimarães and José Guilherme S. Maia	1361
Evaluation of <i>Clausena anisata</i> Essential Oil from Cameroon for Controlling Food Spoilage Fungi and its Potential Use as an Antiradical Agent Aoudou Yaouba, Léopold Ngoune Tatsadjieu, Pierre Michel Jazet Dongmo, François Xavier Etoa, Carl Moses Fontum Mbofung, Paul Henri Amvam Zollo and Chantal Menut	1367
Chemical Diversity in <i>Mentha spicata</i>: Antioxidant and Potato Sprout Inhibition Activity of its Essential Oils Shailendra S. Chauhan, Om Prakash, Rajendra C. Padalia, Vivekanand, Anil K. Pant and Chandra S. Mathela	1373
Role of Direct Bioautographic Method for Detection of Antistaphylococcal Activity of Essential Oils Györgyi Horváth, Noémi Jámbor, Erika Kocsis, Andrea Böszörményi, Éva Lemberkovics, Éva Héthelyi, Krisztina Kovács and Béla Kocsis	1379
Antiphytoviral Activity of Essential Oil from Endemic Species <i>Teucrium arduini</i> Valerija Dunkić, Nada Bezić and Elma Vuko	1385
Toxic Effects of <i>Citrus aurantium</i> and <i>C. limon</i> Essential Oils on <i>Spodoptera frugiperda</i> (Lepidoptera: Noctuidae) Emilio Villafañe, Diego Tolosa, Alicia Bardón and Adriana Neske	1389
Neutralizing Effects of <i>Nectandra angustifolia</i> Extracts against <i>Bothrops neuwiedi</i> Snake Venom Ana M. Torres, Francisco J. Camargo, Gabriela A. L. Ricciardi, Armando I. A. Ricciardi and Eduardo Dellacassa	1393

Review/Account

Artocarpus Plants as a Potential Source of Skin Whitening Agents Enos Tangke Arung, Kuniyoshi Shimizu and Ryuichiro Kondo	1397
Mining Invertebrate Natural Products for Future Therapeutic Treasure Youmie Park	1403

Number 10

Vitexin Inhibits Polyubiquitin Synthesis by the Ubiquitin-conjugating Enzyme E2-25K Kimberli M. Helms, Randall C. Wilson, Ifedayo V. Ogungbe, William N. Setzer and Pamela D. Twigg	1411
Variability of Total Flavonoid, Polyphenol and Tannin Contents in Some <i>Lythrum salicaria</i> Populations Timea Bencsik, Györgyi Horváth and Nóra Papp	1417
Peptidyl-tRNA Hydrolase Screening Combined with Molecular Docking Reveals the Antibiotic Potential of <i>Syzygium johnsonii</i> Bark Extract Sarah M. Harris, Hana McFeeters, Ifedayo V. Ogungbe, Luis R. Cruz-Vera, William N. Setzer, Betsy R. Jackes and Robert L. McFeeters	1421
Betalain: A Particular Class of Antioxidant Pigment Hasna El Gharras	1425
Evaluation of <i>in vivo</i> and <i>in vitro</i> Biological Activities of Different Extracts of <i>Cuscuta arvensis</i> Ufuk Koca, Esra Küpeli-Akkol and Nazim Sekeroglu	1433
<i>In Vitro</i> Anti-inflammatory and Xanthine Oxidase Inhibitory Activity of <i>Tephrosia purpurea</i> Shoot Extract Shivraj H. Nile and Chandrasah N. Khobragade	1437
<i>In Vivo</i> Anti-Inflammatory and <i>In Vitro</i> Antioxidant Activities of Moroccan Medicinal Plants Mina Moussaid, Abd Elaziz Elamrani, Nourdinne Bourhim and Mohamed Benaissa	1441
<i>In Vitro</i> Effect of Wheat Bran (<i>Triticum aestivum</i>) Extract on Calcium Oxalate Urolithiasis Crystallization Khaled Sekkoum, Abdelkrim Cheriti and Safia Taleb	1445
Antioxidant and Antimicrobial Activities of <i>Withania frutescens</i> Laila El Bouzidi, Mustapha Larhsini, Mohamed Markouk, Abdelaziz Abbad, Lahcen Hassani and Khalid Bekkouche	1447
Analytical Evaluation of Three Wild Growing Omani Medicinal Plants Muhammad Asif Hanif, Ahmed Yahya Al-Maskri, Zeyana Mohammed Hamed Al-Mahruqi, Jamal Nasser Al-sabahi, Ahlam Al-Azkawi and Masoud Yahya Al-Maskari	1451
Ethnobotanical Study of Some Aromatic and Medicinal Plants in the Middle Atlas Mountains of Morocco Mohammed El Midaoui, Abdelwahed Maataoui, Mohamed Benbella, Abdelhadi Ait Houssa and Nadia Labazi	1455
Traditional Uses of Medicinal Plants for Respiratory Diseases in Transylvania Nóra Papp, Sámuel Bartha, Gyöngyvér Boris and Lajos Balogh	1459
<i>Perilla frutescens</i>: Interesting New Medicinal and Melliferous Plant in Italy Cinzia Barbieri and Paola Ferrazzi	1461
Benefits of Environmental Conditions for Growing Coriander in Banat Region, Serbia Milica Acimovic, Snezana Oljaca, Goran Jacimovic, Slobodan Drazic and Slavoljub Tasic	1465
Selenium Concentrations of Selected Medicinal and Aromatic Plants in Turkey Faruk Ozkutlu, Nazım Sekeroglu, Ufuk Koca and Gizem Yazıcı	1469
Lead Concentrations of Herbs Used in Van Herby Cheese Murat Tuncurk, Ruveyde Tuncurk, Nazım Sekeroglu, Mehmet M. Ertus and Fevzi Ozgokce	1473
Histological Study of Some <i>Echium vulgare</i>, <i>Pulmonaria officinalis</i> and <i>Symphytum officinale</i> Populations Nóra Papp, Timea Bencsik, Kitti Németh, Kinga Gyergyák, Alexandra Sulc and Ágnes Farkas	1475
Chemical Composition of the Essential Oil from Corsican <i>Mentha aquatica</i> - Combined Analysis by GC(RI), GC-MS and ¹³C NMR Spectroscopy Sylvain Sutour, Félix Tomi, Pascale Bradesi and Joseph Casanova	1479
Chemical Polymorphism of Essential Oils from Populations of <i>Laurus nobilis</i> Grown on Tunisia, Algeria and France Hanen Marzouki, Abdelhamid Khaldi, Bruno Marongiu, Alessandra Piras and Fethia Harzallah-Skhiri	1483
Essential Oil from <i>Ocimum basilicum</i> (Omani Basil): A Desert Crop Ahmed Yahya Al-Maskri, Muhammad Asif Hanif, Masoud Yahya Al-Maskari, Alfie Susan Abraham, Jamal Nasser Al-sabahi and Omar Al-Mantheri	1487

Chemical Composition and Anticandidal Properties of the Essential Oil Isolated from Aerial parts of <i>Cotula cinerea</i>: A Rare and Threatened Medicinal Plant in Morocco Leila El Bouzidi, Abdelaziz Abbad, Karine Fattarsi, Lahcen Hassani, David Leach, Mohammed Markouk, Laurent Legendre and Khalid Bekkouch	1491
Characterization and Comparison of Volatile Constituents of Juice and Peel from Clementine, Mandarin and their Hybrids Toussaint Barboni, Julien Paolini, Pierre Tomi, François Luro, Alain Muselli and Jean Costa	1495
Characterization of Essential Oil and Effects on Growth of <i>Verbena gratissima</i> Plants Treated with Homeopathic Phosphorus Fúlvia M. Santos, Lucila E. F. Monfort, Daniel M. Castro, José E. B. P. Pinto, Michele Leonardi and Luisa Pistelli	1499
Antibacterial Activity of the Essential Oils of <i>Pistacia lentiscus</i> Used in Moroccan Folkloric Medicine Fatima Zohra Mharti, Badiaa Lyoussi and Abdelfattah Abdellaoui	1505
Chemical Composition, Antibacterial and Antioxidant Activities of the Essential Oils from <i>Thymus satureioides</i> and <i>Thymus pallidus</i> Ghalbane Ichrak, Belaqqiz Rim, Ait Said Loubna, Oufdou Khalid, Romane Abderrahmane and El Messoussi Said	1507
Essential Oil Analysis and Antibacterial Activity of <i>Rosmarinus tournefortii</i> from Algeria Mansouria Souria Bendeddouche, Hachemi Benhassaini, Zouaoui Hazem and Abderrahmane Romane	1511
Chemical Composition and Antibacterial Activity of the Essential Oil of Moroccan <i>Juniperus phoenicea</i> Bahri Fouad, Romane Abderrahmane, Arjouni Youssef, Harrak Rajae and M. Ahmed El Alaoui El Fels	1515
Chemical Composition and Antimicrobial Activity of Essential Oil of <i>Cupressus atlantica</i> My Youssef Arjouni, Fouad Bahri, Abderrahmane Romane and My Ahmed El Alaoui El Fels	1519
Isolation of the Volatile Oil from <i>Satureja thymbra</i> by Supercritical Carbon Dioxide Extraction: Chemical Composition and Biological Activity Alessandra Piras, Viviana Cocco, Danilo Falconieri, Silvia Porcedda, Bruno Marongiu, Andrea Maxia, Maria Assunta Frau, Maria J. Gonçalves, Carlos Cavaleiro and Ligia Salgueiro	1523
Chemical Composition and Biological Activity of the Volatile Extracts of <i>Achillea millefolium</i> Danilo Falconieri, Alessandra Piras, Silvia Porcedda, Bruno Marongiu, Maria J. Gonçalves, Céla Cabral, Carlos Cavaleiro and Ligia Salgueiro	1527
Chemical Composition and Biological Activities of Essential Oils of <i>Pinus patula</i> Ismail Amri, Hamrouni Lamia, Samia Gargouri, Mohsen Hanana, Mariem Mahfoudhi, Tarek Fezzani, Ferjani Ezzeddine and Bassem Jamoussi	1531
Composition and Antioxidant Activities of Leaf and Root Volatile Oils of <i>Morinda lucida</i> Sunday O. Okoh, Olayinka T. Asekun, Oluwole B. Familoni and Anthony J. Afolayan	1537
Anti-inflammatory Activity of <i>Pistacia lentiscus</i> Essential Oil: Involvement of IL-6 and TNF-α Andrea Maxia, Cinzia Sanna, Maria Assunta Frau, Alessandra Piras, Manvendra Singh Karchuli and Veena Kasture	1543
Essential Oil of <i>Myrtus communis</i> Inhibits Inflammation in Rats by Reducing Serum IL-6 and TNF-α Andrea Maxia, Maria Assunta Frau, Danilo Falconieri, Manvendra Singh Karchuli and Sanjay Kasture	1545
<u>Review/Account</u>	
Drugs from the Cloudforest: The Search for New Medicines from Monteverde, Costa Rica William N. Setzer	1549
Effect of Salt, Drought and Metal Stress on Essential Oil Yield and Quality in Plants Shreyasee Biswas, Monika Koul and Ashok Kumar Bhatnagar	1559

Number 11

Antifungal Activity of Plumericin and Isoplumericin Dharmendra Singh, Umakant Sharma, Parveen Kumar, Yogesh K. Gupta, M. P. Dobhal and Sarman Singh	1567
A New Diacylated Labdane Diterpenoid from <i>Andrographis wightiana</i> Jalli Madhu Sudhana, Rachakunta Munikishore, Mopuru Vijayabhaskar Reddy, Duvvuru Gunasekar, Alain Blond and Bernard Bodo	1569
Triterpenoid Acids and Lactones from the Leaves of <i>Fadogia tetraquetra</i> var. <i>tetraquetra</i> (Rubiaceae) Dulcie A. Mulholland, Abdelhafeez M.A. Mohammed, Philip H. Coombes, Shafiul Haque, Leena L. Pohjala, Päivi S.M. Tammela and Neil R. Crouch	1573
Isolation of Friedelin from Black Condensate of Cork Ricardo A. Pires, Ivo Aroso, Susana P. Silva, João F. Mano and Rui L. Reis	1577

Novel Microbial Transformation of Resibufogenin by <i>Absidia coerulea</i> Jian Zheng, Dong-hai Su, Dong-sheng Zhang, Xiu-Lan Xin, Jun-ying Liu, Yan Tian, Qing Wei and Xun Cui	1581
Antidiabetic Activity of <i>Terminalia sericea</i> Constituents Nolitha Nkobile, Peter James Houghton, Ahmed Hussein and Namrita Lall	1585
X-ray Crystallographic Study of Ranaconitine Yang Li, Jun-hui Zhou, Gui-jun Han, Min-juan Wang, Wen-ji Sun and Ye Zhao	1589
Obscure: a New Cyclostachine Acid Derivative from <i>Beilschmiedia obscura</i> Bruno Ndjakou Lenta, Jean Rodolphe Chouna, Pepin Alango Nkeng-Efouet, Samuel Fon Kimbu, Etienne Tsamo and Norbert Sewald	1591
Alkaloids from <i>Papaver coreanum</i> Dong-Ung Lee, Jong Hee Park, Ludger Wessjohann and Jürgen Schmidt	1593
Isolation, Structure Elucidation, and Biological Activity of a New Alkaloid from <i>Zanthoxylum rhetsa</i> Karsten Krohn, Stephan Cludius-Brandt, Barbara Schulz, Mambatta Sreelekha and Pottachola Mohamed Shafi	1595
Amplexicine, an Antioxidant Flavan-3-ol from <i>Polygonum amplexicaule</i> Mudasir A. Tantry and Aziz A. Rahman	1597
A New Flavonoid Glycoside from <i>Vaccaria hispanica</i> Haijiang Zhang, Kuiwu Wang, Jie Wu, Yao Chen and Peipei He	1599
Flavonoids from Algerian Endemic <i>Centaurea microcarpa</i> and their Chemotaxonomical Significance Souheila Louaar, Amel Achouri, Mostefa Lefahal, Hocine Laouer, Kamel Medjroubi, Helmut Duddeck and Salah Akkal	1603
On-line (HPLC-NMR) and Off-line Phytochemical Profiling of the Australian Plant, <i>Lasiopetalum macrophyllum</i> Michael Timmers and Sylvia Urban	1605
Chemical Fingerprint Analysis of Phenolics of <i>Albizia chinensis</i> Based on Ultra-Performance LC-Electrospray Ionization-Quadrupole Time-of-Flight Mass Spectrometry and Antioxidant Activity Abha Chaudhary, Pushpinder Kaur, Neeraj Kumar, Bikram Singh, Shiv Awasthi and Brij Lal	1617
Application to Classification of Mulberry Leaves using Multivariate Analysis of Proton NMR Metabolomic Data Eriko Fukuda, Motoyuki Yoshida, Masaki Baba, Yoshihiro Uesawa, Ryuichiro Suzuki, Osamu Kamo, Koji Tsubono, Kazunori Arifuku, Kazuhisa Yatsunami and Yoshihito Okada	1621
Phenolic Constituents of <i>Knautia arvensis</i> Aerial Parts Jaroslaw Moldoch, Barbara Szajwaj, Milena Masullo, Lukasz Pecio, Wieslaw Oleszek, Sonia Piacente and Anna Stochmal	1627
New Acylated Anthocyanins and Other Flavonoids from the Red Flowers of <i>Clematis</i> Cultivars Masanori Hashimoto, Toshisada Suzuki and Tsukasa Iwashina	1631
Prenylated Isoflavonoids from <i>Rhynchosia edulis</i> Ifedayo V. Ogungbe, Gabrielle M. Hill, Rebecca A. Crouch, Bernhard Vogler, Meenakshi Nagarkoti, William A. Haber and William N. Setzer	1637
Antiparasitic and Antimicrobial Isoflavanquinones from <i>Abrus schimperi</i> Aziz A. Rahman, Volodymyr Samoylenko, Surendra K. Jain, Babu L. Tekwani, Shabana I. Khan, Melissa R. Jacob, Jacob O. Midiwo, John P. Hester, Larry A. Walker and Ilias Muhammad	1645
Two New Rotenoids from <i>Boerhavia repens</i> Mamona Nazir, Muhammad Saleem, Naheed Riaz, Maria Hafeez, Misbah Sultan, Abdul Jabbar and Muhammad Shaiq Ali	1651
A Comparison of the Diastereoisomers, Silybin A and Silybin B, on the Induction of Apoptosis in K562 cells Jiyong Zhang, Qiuying Luan, Yanze Liu, David Y-W Lee and Zhao Wang	1653
Effects of pH, Sample Size, and Solvent Partitioning on Recovery of Soluble Phenolic Acids and Isoflavonoids in Leaves and Stems of Red Clover (<i>Trifolium pratense</i> cv. Kenland) Isabelle A. Kagan	1657
Arbutin Derivatives from the Seeds of <i>Madhuca latifolia</i> Shazia Khan, M. Nadeem Kardar and Bina S. Siddiqui	1661
Quinic Acids from <i>Aster caucasicus</i> and from Transgenic Callus Expressing a β-Amyrin Synthase Paola Pecchia, Maria Cammareri, Nicola Malafronte, M. Federica Consiglio, Maria Josefina Gualtieri and Clara Conicella	1665
Cytotoxic Activity and Cell Cycle Analysis of Hexahydro-curcumin on SW 480 Human Colorectal Cancer Cells Chung-Yi Chen, Woei-Ling Yang and Soong-Yu Kuo	1671
Does the Combination of Resveratrol with Al (III) and Zn (II) Improve its Antioxidant Activity? Karina Dias and Sofia Nikolaou	1673
Photosensitization Mechanisms of Triplet Excited State β-Lapachone. A Density Functional Theory Study Liang Shen	1677

AFLP Marking and Polymorphism among Progenies of <i>Gymnema sylvestre</i>: an Important Medicinal Plant of India Magda Abbaker Osman, Sunita Singh Dhawan, Janak Raj Bahl, Mahendra P Darokar and Suman P S Khanuja	1679
Antioxidant Activity of Protein Hydrolysates from Aqueous Extract of Velvet Antler (<i>Cervus elaphus</i>) as Influenced by Molecular Weight and Enzymes Lei Zhao, Yang-Chao Luo, Cheng-Tao Wang and Bao-Ping Ji	1683
Effects of <i>Sideritis euboea</i> (Lamiaceae) Aqueous Extract on IL-6, OPG and RANKL Secretion by Osteoblasts Eva Kassi, Anna Paliogianni, Ismene Dontas, Nektarios Aligiannis, Maria Halabalaki, Zoi Papoutsis, Alexios-Leandros Skaltsounis and Paraskevi Moutsatsou	1689
<i>In vitro</i> Antiprotozoal Activity of Extracts of five Turkish Lamiaceae Species Hasan Kirmizibekmez, Irem Atay, Marcel Kaiser, Erdem Yesilada and Deniz Tasdemir	1697
Antimicrobial Investigation of <i>Linum usitatissimum</i> for the Treatment of Acne Pratibha Nand, Sushma Drabu and Rajinder K Gupta	1701
Chemical and Biological Diversity in Fourteen Selections of Four <i>Ocimum</i> Species Bhaskaruni R. Rajeswara Rao, Sushil K. Kothari, Dharmendra K. Rajput, Rajendra P. Patel and Mahendra P. Darokar	1705
Environmental Effect on Essential Oil Composition of <i>Aloysia citriodora</i> from Corrientes (Argentina) Gabriela Ricciardi, Ana Maria Torres, Ana Laura Bubenik, Armando Ricciardi, Daniel Lorenzo and Eduardo Dellacassa	1711
Essential Oil of Three <i>Uvaria</i> species from Ivory Coast Koffi A. Muriel, Tonzibo Z. Félix, Gilles Figueredo, Pierre Chalard and Yao T. N'guessan	1715
Composition of the Essential Oil of <i>Origanum tyttanthum</i> from Tajikistan Farukh S. Sharopov, Muhamadsho A. Kukaniev and William N. Setzer	1719
Volatiles of French Ferns and “fougère” Scent in Perfumery Didier Froissard, Françoise Fons, Jean-Marie Bessière, Bruno Buatois and Sylvie Rapior	1723
Volatile Constituents of Two Species of <i>Protium</i> from the Atlantic Rainforest in the State of Pernambuco, Brazil José Gildo Rufino de Freitas, Claudio Augusto Gomes da Camara, Marcílio Martins de Moraes and Henrique Costa Hermenegildo da Silva	1727
Volatile Constituents of Two <i>Hypericum</i> Species from Tunisia Karim Hosni, Kamel Msaâda, Mouna Ben taârit, Thouraya Chahed and Brahim Marzouk	1731
Chemical Composition and Possible <i>in vitro</i> Antigermination Activity of Three <i>Hypericum</i> Essential Oils Aurelio Marandino, Laura De Martino, Emilia Mancini, Luigi Milella and Vincenzo De Feo	1735
Antioxidant, Antimicrobial Activities and Fatty Acid Components of Flower, Leaf, Stem and Seed of <i>Hypericum scabrum</i> Ali Shafaghat	1739
Composition of Three Essential Oils, and their Mammalian Cell Toxicity and Antimycobacterial Activity against Drug Resistant-Tuberculosis and Nontuberculous Mycobacteria Strains Juan Bueno, Patricia Escobar, Jairo René Martínez, Sandra Milena Leal and Elena E. Stashenko	1743
Antimicrobial and Antioxidant Activities of the Flower Essential Oil of <i>Halimodendron halodendron</i> Jihua Wang, Hao Liu, Haifeng Gao, Jianglin Zhao, Ligang Zhou, Jianguo Han, Zhu Yu and Fuyu Yang	1749
Composition and Antimicrobial Activity of the Leaf and Twig Oils of <i>Litsea acutivena</i> from Taiwan Chen-Lung Ho, Pei-Chun Liao, Eugene I-Chen Wang and Yu-Chang Su	1755
Chemical Composition and Antimicrobial Activity of the Volatile Oil from <i>Fusarium tricinctum</i>, the Endophytic Fungus in <i>Paris polyphylla</i> var. <i>yunnanensis</i> Ying Zhang, Jianglin Zhao, Jihua Wang, Tijiang Shan, Yan Mou, Ligang Zhou and Jingguo Wang	1759
Antifungal Activity of Essential Oil from <i>Asteriscus graveolens</i> against Postharvest Phytopathogenic Fungi in Apples Mohamed Znini, Gregory Cristofari, Lhou Majidi, Hamid Mazouz, Pierre Tomi, Julien Paolini and Jean Costa	1763
Interspecies Comparison of Chemical Composition and Anxiolytic-like Effects of Lavender Oils upon Inhalation Mizuho Takahashi, Tadaaki Satou, Mai Ohashi, Shinichiro Hayashi, Kiyomi Sadamoto and Kazuo Koike	1769
<u>Review/Account</u>	
Essential Oils from the <i>Hyptis</i> genus- A Review (1909-2009) Megil McNeil, Petrea Facey and Roy Porter	1775
<u>Additions/Corrections</u>	1797

Number 12

Sibiralactone: A New Monoterpene from <i>Sibiraea angustata</i> Guangbo Xie, Xianlong Wang, Tibor Kurtán, Attila Mándi and Tianzhi Wang	1799
---	------

Bioconversion of Proposed Precursors into Theobroxide and Related Compounds Peng Li, Kosaku Takahashi, Ahmed Elkhateeb, Hideyuki Matsuura, Teruhiko Yoshihara and Kensuke Nabeta	1801
Microbial Hydroxylation of <i>S</i>-(-)-Perillyl Alcohol by <i>Fusarium heterosporium</i> Ismail Kiran	1805
A Phytochemical Investigation of <i>Zanthoxylum setulosum</i> Tameka M. Walker, Bernhard Vogler, Debra M. Moriarity, William A. Haber and William N. Setzer	1807
Cytotoxic Cembranoids from the Red Sea Soft Coral <i>Sarcophyton glaucum</i> Mohamed-Elamir F. Hegazy, Ahmed A. El-Beih, Alaa Y. Moustafa, Abdelhamed A. Hamdy, Montaser A. Alhammad, Rehab M. Selim, Mohamed Abdel-Rehim and Paul W. Paré	1809
C-Lactam Derivatives of Oleanolic Acid. The synthesis of C-lactam by Beckmann rearrangement of C-oxime Barbara Bednarczyk – Cwynar	1813
Analysis of Native Carotenoid Composition of Sweet Bell Peppers by Serially Coupled C₃₀ Columns Daniele Giuffrida, Paola Dugo, Giacomo Dugo, Germana Torre and Luigi Mondello	1817
New Antifungal Cholestane and Aldehyde Derivatives from the Red Alga <i>Laurencia papillosa</i> Walied M. Alarif, Sultan S. Al-Lihaibi, Ahmed Abdel-Lateff and Seif-Eldin N. Ayyad	1821
Steroidal Saponins from the Fruits of <i>Cestrum ruizteranianum</i> Elier Galaraga M., Anne-Claire Mitaine-Offer, Juan Manuel Amaro-Luis, Tomofumi Miyamoto, Chiaki Tanaka, Laurent Pouységu, Stéphane Quideau, Luis B. Rojas and Marie-Aleth Lacaille-Dubois	1825
Isolation and Cholinesterase Activity of Amaryllidaceae Alkaloids from <i>Nerine bowdenii</i> Lucie Cahliková, Stanislav Zavadil, Kateřina Macáková, Irena Valterová, Andrea Kulhánková, Anna Hošťálková, Jiří Kuneš and Lubomír Opletal	1827
HPLC Determination of Majdine in <i>Vinca herbacea</i> Natia Gagua, Beatrice Baghdikian, Fathi Mabrouki, Riad Elias, Valentina Vachnadze, Aliosha Bakuridze and Evelyne Ollivier	1831
Pyridine Metabolism and Trigonelline Synthesis in Leaves of the Mangrove Legume trees <i>Derris indica</i> (<i>Millettia pinnata</i>) and <i>Caesalpinia crista</i> Yuling Yin, Hamako Sasamoto and Hiroshi Ashihara	1835
Anti-adipogenic Activity of <i>Cordyceps militaris</i> in 3T3-L1 Cells Qing Liu, In Pyo Hong, Mi-Jeong Ahn, Hwan-Soo Yoo, Sang-Bae Han, Bang Yeon Hwang and Mi Kyeong Lee	1839
Two New Cyclopeptides and One New Nonenolide from <i>Xylaria</i> sp. 101 Yao-Yao Li, Zhi-Yu Hu, and Yue-Mao Shen	1843
A Novel Flavonoid and Furoquinoline Alkaloids from <i>Vepris glomerata</i> and their Antioxidant Activity Joyce J. Kiplimo, Md. Shahidul Islam and Neil A. Koorbanally	1847
Flavonoid Constituents and Free Radical Scavenging Activity of <i>Alchemilla mollis</i> Antoaneta Trendafilova, Milka Todorova, Milena Nikolova, Anna Gavrilova and Antonina Vitkova	1851
Ultrasound-assisted Extraction of Total Phenols and Flavonoids from Dry Tobacco (<i>Nicotiana tabacum</i>) Leaves Ivana T. Karabegović, Vlada B. Veljković and Miodrag L. Lazić	1855
Characterization of Polyphenolic Compounds in Unripe Chinotto (<i>Citrus myrtifolia</i>) Fruit by HPLC/PDA/ESI/MS-MS Monica Scordino, Leonardo Sabatino, Adalgisa Belligno and Giacomo Gagliano	1857
Bioactive Compounds, RP-HPLC Analysis of Phenolics, and Antioxidant Activity of Some Portuguese Shrub Species Extracts Ángelo Luís, Fernanda Domingues and Ana Paula Duarte	1863
HPLC/PDA/ESI-MS Evaluation of Saffron (<i>Crocus sativus</i> L.) Adulteration Leonardo Sabatino, Monica Scordino, Maria Gargano, Adalgisa Belligno, Pasqualino Traulo and Giacomo Gagliano	1873
Ferric Reducing, Antiradical and β-Carotene Bleaching Activities of Nicotinic Acid and Picolinic Acid Bioconjugates of Curcumin Archana Pandey, Kanti Bhooshan Pandey, Ravindra Kumar Gupta and Syed Ibrahim Rizvi	1877
Antiviral Activities of Diarylheptanoids Isolated from <i>Alpinia officinarum</i> against Respiratory Syncytial Virus, Poliovirus, Measles Virus, and Herpes Simplex Virus Type 1 <i>in vitro</i> Katsuhiko Konno, Rie Sawamura, Yi Sun, Ken Yasukawa, Tomomi Shimizu, Wataru Watanabe, Masahiko Kato, Ryuichi Yamamoto and Masahiko Kurokawa	1881
Isolation of C-glycosyl Xanthones from <i>Coffea pseudozanguebariae</i> and Their Location Pascale Talamond, Geneviève Conejero, Jean-Luc Verdeil and Jean-Luc Poëssel	1885

Antifungal Activity and Isomerization of Octadecyl <i>p</i>-coumarates from <i>Ipomoea carnea</i> subsp. <i>fistulosa</i> Eugene Sebastian J. Nidiry, Girija Ganeshan and Ankanahalli N. Loksha	1889
New Glucose Esters from the Fresh Leaves of <i>Jacaranda mimosaeifolia</i> Christianah A. Elusiyan and Tiwalade A. Olugbade	1893
Shamiminol: A New Aromatic Glycoside from the Stem Bark of <i>Bombax ceiba</i> Shaheen Faizi, Sadia Zikr-Ur-Rehman and Muhammad Ali Versiani	1897
Two New Phenolic Glycosides from <i>Viburnum plicatum</i> var. <i>plicatum</i> f. <i>plicatum</i> Saki Katagiri, Yoshiki Watanabe, Yasunori Yaoita, Masao Kikuchi and Koichi Machida	1901
Antimicrobial Chemical Constituents from the Endophytic Fungus <i>Phomopsis</i> sp. from <i>Notobasis syriaca</i> Hidayat Hussain, Michel Kenne Tchimine, Ishtiaq Ahmed, Kathrin Meier, Michael Steinert, Siegfried Draeger, Barbara Schulz and Karsten Krohn	1905
Phomosines H–J, Novel Highly Substituted Biaryl Ethers, Isolated from the Endophytic Fungus <i>Phomopsis</i> sp. from <i>Ligustrum vulgare</i>[†] Karsten Krohn, Umar Farooq, Hidayat Hussain, Ishtiaq Ahmed, Joachim Rheinheimer, Siegfried Draeger, Barbara Schulz and Teunis van Ree	1907
Isolation and Characterization of a new Benzofuran from the Fungus <i>Alternaria</i> sp. (HS-3) Associated with a Sea Cucumber Xuekui Xia, Jun Qi, Fang Wei, Airong Jia, Wenpeng Yuan, Xiumei Meng, Miansong Zhang, Changheng Liu and Changyun Wang	1913
Potent Toxic Macrocylic Trichothecenes from the Marine-Derived Fungus <i>Myrothecium verrucaria</i> Hmp-F73 Li Zhao, Li Liu, Nan Wang, Shu-Jin Wang, Jing-Chun Hu and Jin-Ming Gao	1915
Synthesis and Bioactivity of Novel Coumarin Derivatives Ai-Ying Guan, Chang-Ling Liu, Miao Li, Zhi-Nian Li, Ming-Xing Zhang and Hong Zhang	1917
Kinase Inhibitory, Haemolytic and Cytotoxic Activity of Three Deep-water Sponges from North Western Australia and their Fatty Acid Composition Ana Zivanovic, Natalie J. Pastro, Jane Fromont, Murray Thomson and Danielle Skropeta	1921
Antimicrobial and Cytotoxic Effects of Mexican Medicinal Plants Maria del Rosario Jacobo-Salcedo, Angel Josabad Alonso-Castro, Luis A. Salazar-Olivo, Candy Carranza-Alvarez, Luis Ángel González-Espíndola, Fabiola Domínguez, Sandra Patricia Maciel-Torres, Concepción García-Lujan, Marisela del Rocio González-Martínez, Maricela Gómez-Sánchez, Eduardo Estrada-Castillón, Rocio Zapata-Bustos, Pedro Medellín-Milán and Alejandro García-Carrancá	1925
Chemometrics Evaluation of the Herbal Drug <i>Andrographis paniculata</i> Shiv Narayan Sharma, Zenu Jha and D. K. Sharma	1929
<i>Garcinia cambogia</i> Leaf and Seawater for Tannase Production by Marine <i>Aspergillus awamori</i> BTMFW032 under Slurry State Fermentation Beena P. S, Soorej M. Basheer, Sarita G. Bhat and Chandrasekaran M	1933
Gas Chromatographic Quantitative Analysis of Methanol in Wine: Operative Conditions, Optimization and Calibration Model Choice Rosario Caruso, Grazia Laura Gambino, Monica Scordino, Leonardo Sabatino, Pasqualino Traulo and Giacomo Gagliano	1939
Composition and Biological Potential of Essential Oil from <i>Thelechitonina trilobata</i> Growing in South Africa Jamie Peebles, Ephraim Gwebu, Opeoluwa Oyedeji, Sarah Nanyonga, Nokuthula Kunene, David Jackson, William Setzer and Adebola Oyedeji	1945
Chemical Composition and Antibacterial Activity of Essential oil from <i>Salvia mukerjeei</i> Lalit Mohan, Anuradha Negi, Anand B. Melkani and Vasu Dev	1949
<u>Review/Account</u>	
Revealing Indigenous Indonesian Traditional Medicine: Anti-infective Agents Ari S. Nugraha and Paul A. Keller	1953

Natural Product Communications

Volume 6 (1-12)

2011

Author Index

Abbad, A.....	1447,1491	Andersen, ØM.....	35	Balogh, L.....	1459	Bonito, MC.....	1205
Abdalla, MA.....	45	Andrade, EHA.....	1361	Ban, NK.....	159	Borcsa, B.....	527
Abdellaoui, A.....	1505	Andreazza, NL.....	977	Bancheva, S.....	1339	Boris, G.....	1459
Abdel-Lateff, A.....	1821	Andrela, ALB.....	983	Bandeira, MAM.....	1001	Borkosky, S.....	387
Abdel-Rehim, M.....	1809	Andryjaschenko, PV.....	1075	Bandyopadhyay, D.....	209	Bortoloti, LV.....	983
Abderrahmane, R..	1507,1515,1519	Anter, J.....	59	Banerjee, M.....	209	Bosch, J.....	515
Abedini, W.....	985	Antonov, AS.....	1063	Banerji, A.....	625	Böszörményi, A.....	1379
Abraham, AS.....	1487	Anufriev, VP.....	217	Banerji, J.....	625	Botubol, JM.....	443
Abreu-Abreu, ND.....	59	Aoki, H.....	403	Banik, R.....	747	Bouderdara, N.....	115
Abu-Ghannam, N.....	1299	Aparicio, R.....	935	Bankova, V.....	21	Bourhim, N.....	1441
Achouri, A.....	1603	Aquino, RP.....	951	Bárbara Mayer, B.....	1253	Boutaghane, N.....	251
Acimovic, M.....	1465	Arak, E.....	1107	Barbieri, C.....	1461	Bouzidi, LE.....	1447,1491
Adebayo, AH.....	1263	Aranda, RS.....	1035	Barboni, T.....	1495	Boyer, L.....	753
Adema, M.....	985	Arbain, D.....	349,357	Bardón, A.....	1389	Bradesi, P.....	1479
Adima, AA.....	1183	Areche, C.....	1073	Bardón, A.....	387	Breckler, L.....	621
Adryjashchenko, PV.....	587	Arevalo, C.....	957	Barrero, AF.....	439,879	Briones, V.....	985
Afiyatullof, SS.....	673,1063	Arifuku, K.....	1621	Bartha, S.....	1459	Brodie, PJ.....	1259
Afolayan, AJ.....	1537	Arita, M.....	7	Basak, B.....	209	Broniatowska, B.....	689
Afza, N.....	1083	Arjouni, MY.....	1519	Başer, KHC.....	701	Bruneau, C.....	251
Agnaniet, H.....	409	Armstrong, V.....	477	Basheer, SM.....	1933	Bruno, M.....	1023
Agrebi, A.....	409	Arnold, AE.....	835	Bauer, R.....	527	Bruno, M.....	1339
Ahmad, VU.....	1,179	Arnold, NA.....	1347	Becnel, JJ.....	701	Buatois, BG.....	1723
Ahmed, I.....	661,1133,1905	Aroso, I.....	1577	Bedascarrasbure, E.....	823	Bubenik, AL.....	1711
Ahmed, Z.....	1099	Aroulmoji, V.....	593	Bedi, YS.....	93	Bueno, J.....	1743
Ahn, MJ.....	1839	Arrabal, C.....	123	Bednarczyk – Cwynar, B.....	1813	Buitrago, A.....	699
Akesson, B.....	1269	Arraiza, MP.....	123	Beena, PS.....	1933	Buitrago, A.....	85
Akindahunsi, AA.....	1263	Arruda, ACO.....	1045	Beiza-Granados, L.....	451	Bun, SS.....	753
Akkal, S.....	1603	Arteaga, AF.....	439	Bekkouche, K.....	1447,1491	Bunluepuech, K.....	1095
Akutsu, K.....	371	Arteaga, P.....	439	Beleboni, RO.....	193	Burgueño-Tapia, E.....	429
Alachkar, A.....	79	Arung, ET.....	1397	Belligno, A.....	1857,1873	Burkholz, T.....	31
Alankuş-Çalışkan, Ö.....	205,429	Asakawa, Y.....	303,315,349,357	Bellis, RD.....	1315	Bustillo, AJ.....	443
Alarcón, J.....	1085	Asbaghian, S.....	137	Beluhan, S.....	73		
Alarif, WM.....	1821	Asekun, OT.....	1537	Ben taarit, M.....	1731	Cabral, C.....	1527
Al-Azkawi, A.....	1451	Ashihara, H.....	1835	Benaissa, M.....	1441	Cahliková, L.....	1827
Aleu, J.....	443	Asres, K.....	1229	Benbella, M.....	1455	Cahliková, L.....	603,607,707,1255
Alhammady, MA.....	1809	Assafim, M.....	961	Bencsik, T.....	1417,1475	Cai, C.....	639,1323
Ali, MI.....	1099	Assreuy, AMS.....	863	Beneddouch, MS.....	1511	Cai, SF.....	1141
Ali, MS.....	1099,1651	Astulla, A.....	323	Benešová, N.....	1255	Cai, SQ.....	11
Ali, Z.....	1117	Atay, I.....	1697	Benhassaini, H.....	1511	Calcabrini, C.....	1315
Aligiannis, N.....	1689	Attarian, R.....	163	Bergamin, M.....	555,593	Callmander, MW.....	1259
Alipieva, K.....	21	Av-GayY.....	163	Berger, RG.....	1217	Camara, CAG.....	1727
Al-Lihaibi, SS.....	1821	Avilov, SA.....	587,1075	Berti, L.....	1159	Camargo, EES.....	1001
Allmendinger, A.....	689	Aviña-Verduzco, J.....	857	Bessière, JM.....	1723	Camargo, FJ.....	1393
Al-Mahruqi, ZMH.....	1451	Avunduk, S.....	205	Bezić, N.....	1385	Cammareri, M.....	1665
Al-Mantheri, O.....	1487	Awasthi, S.....	1617	Bhat, SG.....	1933	Campone, L.....	957
Almanza, GR.....	1269	Ayyad, SN.....	1821	Bhat, SV.....	1221	Campos, BMMG.....	1035
Al-Maskri, AY.....	1451,1487	Aziz, S.....	841	Bhatnagar, AKJ.....	1555	Cano-Escudero, IC.....	451
Almeida, AP.....	807,961			Bhutia, TD.....	621	Canzoneri, M.....	1023
Alonso-Castro, AJ.....	1121,1925	Ba, LA.....	31	Bi, SF.....	353	Carbonari, JL.....	993
Alonso-Moraga, A.....	59	Baba, M.....	1621	Bighelli, A.....	379	Carda, M.....	505
Al-sabahi, JN.....	1451,1487	Babu, GD.....	1155	Bigović, D.....	819	Cardile, V.....	1023
Álvarez-Corral, M.....	537	Bach, H.....	163	Bikanga, R.....	409	Carezzano, ME.....	1039
Alves, RS.....	871	Bach, H.....	433	Bilia, AR.....	79	Cariddi, LN.....	989,995
Alviano, CS.....	1027	Bach, SM.....	163,433	Bing, Z.....	159	Carlucci, G.....	1275
Alviano, DS.....	1027	Bachetti, RA.....	995	Bisht, D.....	111	Carmona, A, J.....	1051
Amano, O.....	187	Badakar, VM.....	141	Bisrat, D.....	1229	Carmona, J.....	1031
Amaro-Luis, JM.....	1825	Bader, S.....	179	Biswas, S.....	1555	Carranza-Alvarez, C.....	1925
Amat, M.....	515	Baggio, CH.....	1253	Bizzo, HR.....	1027	Carreira, LMM.....	1361
Ambrosio, L.....	943	Baghdikian, B.....	753,1831	Blagojević, PD.....	1015	Caruso, R.....	1939
Ambrosio, SR.....	777	Bahl, JR.....	1679	Blond, A.....	1569	Carvalho, LAC.....	243
Aminin, DL.....	587	Bahri, F.....	1519	Blunden, G.....	575,689	Casabianca, H.....	409
Amorim, RMF.....	863	Bakuridze, A.....	1831	Bodo, B.....	1569	Casanova, J.....	379,1159,1479
Amri, I.....	1531	Baldini, S.....	593	Bonaccorsi, I.....	417,1009	Castillo, A.....	439
An, L.....	781	Baldovino, S.....	699	Bonacorsi, C.....	983	Castola, V.....	379
Anand, R.....	93	Ballarin, L.....	593	Bonesi, M.....	253	Castro, DM.....	1499

- Catalán, CAN.....163,433
 Cavalcanti, SCH.....907
 Cavaleiro, C.....1523,1527
 Cecotti, R.....101
 Cerda-García-Rojas, CM.....1225
 Céspedes, A. CL.....1085
 Chahed, T.....1731
 Chalard, P.....87,1715
 Chalchat, JC.....87
 Chan, KL.....337
 Chandrasekaran M.....1933
 Chanotiya, CS.....111,239,1333
 Chao, Z.....55
 Chataigné, G.....813
 Chatsumpun, M.....41
 Chaturvedula, VSP.....175,1059
 Chaudhary, A.....1617
 Chauhan, A.....239
 Chauhan, SS.....1373
 Chaverri, C.....145
 Chaves, DSA.....961
 Chen, A.....639,645
 Chen, CY.....1251,1297
 Chen, CY.....227,855,1671
 Chen, D.....1285
 Chen, G.....1137
 Chen, J.....53
 Chen, L.....53
 Chen, SL.....1131
 Chen, Y.....1599
 Cheng, HY.....845
 Cheriti, A.....1445
 Chhabra, S.....1115
 Chimnoi, N.....1103
 Chin, YW.....1267
 Chiriboga, X.....767
 Chizzola, R.....1147
 Chlebek, J.....607,707
 Chobot, V.....597
 Chokchaichamnankit, D.....1103
 Chopra, CS.....887
 Chou, JC.....133,267
 Choudhary, MI.....1117
 Chouna, JR.....1591
 Christ, M.....15
 Christelle, TD.....1133,1135
 Christen, P.....879
 Christophersen, C.....921
 Chuanaasa, T.....41
 Chumala, PB.....617
 Chun, EAC.....803
 Cicala, C.....1205
 Ciccio, JF.....145
 Cioni, PL.....883
 Cirio, AT.....1035
 Cisowska, A.....149
 Cludius-Brandt, S.....1595
 Cocco, V.....1523
 Coentrão, PAM.....651
 Coley, PD.....835
 Collado, IG.....443
 Comai, S.....793
 Compagnone, RS.....97
 Conejero, G.....1885
 Conicella, C.....1665
 Consiglio, MF.....1665
 Contigiani, MS.....995
 Contini, SHT.....193
 Coombes, PH.....1573
 Coppa, M.....101
 Coppede, J.....193
 Cordal, MAB.....985
 Cortés, M.....477
 Cortez-Espinosa, N.....857
 Costa, EV.....907
 Costa, GM.....913
 Costa, J.....1495,1763
 Costa, SS.....961
 Cotrim, CA.....983
 Cotroneo, A.....417
 Cotugno, R.....943
 Couladis, M.....271
 Coy-Barrera, E.....231
 Cretton, S.....879
 Cristofari, G.....1763
 Crouch, NR.....1573
 Crouch, RA.....1637
 Cruz, MC.....1005
 Cruz-Vera, LR.....1421
 Csupor, D.....527
 Cubilla-Rios, L.....835
 Cuca, LE.....231
 Cucchiari, L.....1315
 Cuesta-Rubio, O.....973
 Cui, X.....781,1581
 Cunha, WR.....777
 Cuong, NX.....159
 da Camara, CAG.....893
 da Rocha, EMP.....777
 da Silva, JKR.....1361
 Dachà, M.....1315
 Dall'Acqua, S.....793
 Darokar, MP.....1679,1705
 Das, J.....129
 de Andrade, BB.....777
 de Campos, SC.....983
 de Jesus, HCR.....907
 de Lourenço, CC.....977
 de Oliveira, AG.....1001
 de Trimarco, JT.....163
 Debnath, S.....747
 Dejaegher, B.....813
 Delgado, G.....231
 Delgado, V.....477
 Dellacassa, E.....1393,1711
 Demirci, B.....701
 Demirci, F.....701
 Demo, MS.....1039
 Deng, S.....781
 Denisenko, VA.....673
 Derita, M.....931
 Descamps, LR.....887
 Dethoup, T.....807
 Dev, V.....1949
 Dhawan, SS.....1679
 Diallo, D.....1125
 Dias, K.....1673
 Díaz, C.....145
 Diaz, FR.....433
 Diaz, T.....1031,1051
 Diaz-Lanza, A.....753
 Diesel, A.....677
 Diesel, B.....31
 Dinda, B.....747
 Djarri, L.....115
 Djurić, Z.....819
 Dmitrenok, PS.....673,1075
 Dobhal, MP.....1567
 Domingues, F.....1121,1863,1925
 Domínguez, MJ.....993
 Dongmo, PMJ.....1367
 Dongo, E.....1133,1135
 Dontas, I.....1689
 Dörr, M.....15
 dos Santos, AC.....1253
 dos Santos, RLC.....907
 Drabu, S.....1701
 Draeger, S.....661,1907
 Drage, S.....597
 Drazic, S.....1465
 Du, Q.....3
 Du, Z.....167
 Duan, J.....17
 Duarte, AP.....1863
 Duarte, LML.....1189
 Duddeck, H.....1603
 Duez, P.....253
 Dugo, G.....417,1009,1817
 Dugo, P.....417,1817
 Dunkić, V.....1385
 Dutra, LM.....907
 Ebel, R.....677
 Eberlin, MN.....969
 Echeverria, J.....759
 Ehrenfried, CA.....969
 Ekasari, W.....323
 Elamrani, AE.....1441
 El-Azzouny, A.....251
 El-Beih, AA.....1809
 Elias, R.....753,1831
 Elier Galarraaga M., E.....1825
 Elkhateeb, A.....1801
 Elomri, A.....115
 Elsheikh, MS.....79
 Elusiyan, CA.....1893
 Epifano, F.....1275
 Ertus, MM.....1473
 Escobar, FM.....989,995
 Escobar, P.....1743
 Eshilokun, AO.....875
 Espineli, DL.....803
 Estrada-Castillón, E.....1925
 Etoa, FX.....1367
 Evangelista, IL.....871
 Evangelista, JF.....871
 Evangelista, JSAM.....871
 Ezure, Y.....829
 Ezzeddine, F.....1531
 Facey, P.....1775
 Fachin, AL.....193
 Fagundes, FHR.....983
 Fahad, A.....7
 Faizi, S.....1897
 Fakhry, HA.....1009
 Falchero, L.....101
 Falco, SD.....943
 Falconieri, D.....1523,1527,1545
 FAMILONI, OB.....1537
 Farias, WRL.....863
 Farkas, A.....1475
 Farooq, U.....179
 Fatima, I.....1083
 Fattarsi, K.....1491
 Fávero, OA.....243
 Félix, TZ.....1715
 Fels, MAEAE.....1515,1519
 Feng, J.....1323
 Feo, VD.....1735
 Fernandes, VC.....193
 Fernández, MC.....973
 Fernández, MD.....1179
 Fernández-Bedmar, Z.....59
 Ferrazzi, P.....1461
 Ferreira.....667
 Ferreira, MJP.....243
 Ferreres, F.....1305
 Ferrero, AA.....887
 Fezzani, T.....1531
 Fidalgo, LM.....973
 Figueiredo, L.....667
 Figueiredo, SS.....193
 Figueredo, G.....1715
 Figueredo, G.....87
 Fiori, GML.....193
 Flamini, G.....883
 Fleury, BG.....667
 Foley, WJ.....1281
 Fons, F.....1723
 Forgo, P.....527,575
 Formisano, C.....1023,1339,1347
 Fortuna, MA.....163
 Fouad, B.....1515
 Fowles, RG.....785
 Franz, C.....1343
 Fraternale, D.....1315
 Frattani, FS.....961
 Frau, MA.....1523,1543,1545
 Freitas, CS.....1253
 Freitas, JFR.....1727
 Friardi.....349
 Froissard, D.....1723
 Fromont, J.....1921
 Fukuda, E.....1621
 Fukushima, M.....311
 Fukushima, O.....1269
 Fukuyama, Y.....327
 Furtado, NAJC.....777
 Gagliano, G.....1857,1873,1939
 Gagua, N.....1831
 Galeano JE.....287
 Gales, L.....807
 Galindo, VMR.....1035
 Gallucci, MN.....1039
 Gambino, GL.....1939
 Ganeshan, G.....1889
 Gao, H.....1749
 Gao, JM.....1129,1915
 Gao, P.....1055
 García, FM.....123
 García, M.....281
 García, SG.....123
 García-Carranca, A.....1121
 García-Carranca, A.....1925
 García-Lujan, C.....1925
 García-Vázquez, JB.....457
 Garduño, L.....1005
 Gargano, M.....1873
 Gargouri, S.....1531
 Gaviira, M.....247
 Gavrilova, A.....1851
 Gaydou, EM.....1151
 Ge, HM.....353
 Ge, SY.....867
 Genovese, S.....1275
 Gerwick, WH.....835
 Gewali, MB.....793
 Gharras, HE.....1425
 Ghosh, A.....625
 Gilardon, G.....767
 Gil-Izquierdo, A.....1305
 Giuffrida, D.....1817
 Glamočlija, J.....263
 Gliszczynska, A.....367
 Gogalic, S.....677
 Gomes, ac.....497
 Gómez-Hurtado, MA.....1225
 Gómez-Sánchez, M.....1925
 Gonçalves, MJ.....1523,1527
 Gong, X.....361
 Gong, X.....789
 Gonzalez-Espindola, LA 1121,1925
 González-Juárez, DE.....457
 Gonzalez-Martinez, MR 1121,1925
 Goran, N.....1353
 Gorecki, D.....645
 Gorecki, DC.....581,639
 Gourine, N.....1151
 Gradstein, SR.....315
 Grecchi, M.....101
 Griera, R.....515
 Gu, HF.....67
 Gualtieri, M.....943
 Gualtieri, MJ.....1665
 Guan, AY.....1917
 Guerra, FM.....491
 Guimarães, EF.....939,1361
 Guinoiseau, E.....1159
 Guiry, M.....689
 Gulati, A.....1155
 Guleria, S.....107
 Guleria, S.....1155
 Gunasekar, D.....1569

Guo, DX	49
Guo, J	17
Guo, Z	799
Gupta, AP	93
Gupta, RK	1701
Gupta, RK	1877
Gupta, S	93,1299
Gupta, YK	1567
Gürbüz, I	701
Guzii, AG	673
Gwebu, E	1945
Gyergyák, K	1475
Haber, WA	1637,1807
Habtemariam, S	201,1089
Hadacek, F	597
Hafeez, M	1651
Hagiwara, H	311,333
Hajdari, A	1343
Halabalaki, M	1689
Hamdy, AA	1809
Hameed, S	1083
Hammerich, O	921
Han, GJ	1589
Han, HJ	1263
Han, J	1749
Han, L	749,799
Han, SB	1839
Hanana, M	1531
Handique, PJ	129
Hang, DTT	159
Hanif, MA	1451,1487
Haque, S	1573
Harada, K	327
Harigaya, Y	747
Harinantenaina, L	1259
Harris, SM	1421
Hartmann, RW	31
Harzallah-Skhiri, F	1483
Hasegawa, T	371
Hasegawa, Y	789
Hashimoto, K	187
Hashimoto, M	1631
Hassani, L	1447,1491
Hayashi, K	7
Hayashi, S	1769
Hazem, Z	1511
Háznagy-Radnai, E	235
Hazra, K	625
He, P	1599
He, WJ	1263
Hegazy, MF	1809
Heinrich, M	627
Heleno, VCG	777
Helms, KM	1411
Hendrich, AB	149
Hermine, JMB	1133
Hernández, IM	973
Hernández-Barragán, A	457
Hernández-Cervantes, C	537
Hernández-Galán, R	443
Hernández-Hernández, JD	1225
Herrador, MM	439,879
Herz, W	807
Hester, JP	1645
Héthelyi, E	1379
Heyden, YV	813
Hill, GM	1637
Hincapié L.CA	1085
Hingley-Wilson, S	689
Hioki, H	327
Hirasawa, Y	323
Ho, CL	133,267,901,1357,1755
Hoai, NN	813
Hoffmann, A	393
Hohmann, J	527,575
Holz, J	393
Hong, IP	1839
Hong, QT	813

Hong, VNT	813
Hong, ZL	1297
Hongratanaworakit, T	1199
Horváth, G	1379,1417
Hoshi, T	311,333
Hosni, K	1731
Hosny, M	31
Hošťáková, A	707,1827
Hou, FF	581,639,645
Houghton, PJ	1585
Houssa, AA	1455
Hsu, KP	267
Hu, JC	1915
Hu, ZY	1843
Huang, H	1239
Huang, JC	1251,1297
Huang, SS	781
Huang, W	1111
Huang, W	845
Husain, MK	93
Hussain, H	1133,1135,1905
Hussain, H	661,841,1115
Hussain, J	171,1135
Hussein, A	1585
Hwang, BY	1267,1839
Hyun, CG	1193
Ichrak, G	1507
Imhoff, JF	1247
Indrayanto, G	677
Innocenti, G	793
Iqbal, S	179
Isla, MI	823,965
Islam, MS	1847
Islam, MT	45
Ito, T	303
Ivanchina, NV	673
Iwashina, T	1631
Jabbar, A	1651
Jacimovic, G	1465
Jacinto, SD	803
Jackes, BR	1421
Jackson, D	1945
Jacob, C	31
Jacob, MR	1645
Jacobo-Salcedo, MR	1121,1925
Jaddouh, A	79
Jahodář, L	603
Jain, SK	1645
Jaiswal, AK	1299
Jámbor, N	1379
Jamoussi, B	1531
Janačković, P	1163
Jang, HD	159
Janicsák, G	575
Janković, T	819
Jarikasem, S	1289
Jean, GA	1183
Jha, PK	793
Jha, Z	1929
Ji, BP	1683
Ji, CJ	1263
Ji, M	49
Ji, NY	1243
Jia, A	1913
Jia, W	1055
Jiménez, FE	1005
Jiménez-González, L	537
Jin, P	639,645
Jing, W	197
Jiroš, P	603
Jitsuno, M	187
Jordheim, M	35
Jorge, ZD	491
Joseph-Nathan, P	429,451,457
Joseph-Nathan, P	759,1225
Joshi, R	1155
Joshi, RK	141,1145

Juárez, BE	965
Julius, OE	1135
Kabouche, A	251
Kabouche, Z	251
Kagan, IA	1657
Kaiser, M	689,1697
Kajimoto, T	1091
Kalinin, VI	587,1075
Kalinovsky, AI	673,773,1063,1075
Kaminsky, W	1237
Kamo, O	1621
Kaplan, MAC	939
Kar, M	625
Karabegović, IT	1855
Karchuli, MS	1543,1545
Kardar, MN	1661
Karim, A	1083
Kasahara, Y	403
Kasali, AA	875
Kasimov, F	137
Kassi, E	1689
Kasture, S	1545
Kasture, V	1543
Katagiri, S	1901
Katar, MN	343
Kátay, E	611
Kátay, G	611
Kato, M	1881
Kato, MJ	1361
Kaur, P	1617
Kawakami, Y	403
Kebede, W	1229
Keller, PAQ	1953
Khalid, A	1483
Khalid, O	1507
Khan, A	171,179,1117
Khan, FU	171
Khan, MA	179
Khan, R	555,1083
Khan, S	1661
Khan, SI	1645
Khan, SS	1,179
Khanuja, SPS	1679
Khatib, N	141
Khatoon, R	1
Khobragade, CN	1437
Kholkute, SD	141
Khunnawutmanotham, N	1103
Kidyu, K	1069
Kiem, PV	159
Kiemer, AK	31
Kijjoa, A	807
Kikuchi, M	1901
Kim, JA	1233
Kim, JE	1193
Kim, JS	845
Kim, SS	1193
Kim, TNT	813
Kim, YH	159,1233
Kimbu, SF	1591
Kingston, DGI	1259
Kinugawa, K	311
Kiplimo, JJ	1847
Király-Véghely, Z	631
Kiran Babu, GD	107
Kiran, I	1805
Kirmizibekmez, H	1697
Kishi, Y	371
Kitic, D	1167
Koca, U	1433,1469
Kocsis, B	1379
Kocsis, E	1379
Koike, K	1769
Kokoska, L	213
Komala, I	303,349,357
Komatsu, K	11
Konakchiev, A	905
Končić, MZ	1275

Kondo, T	1397
Kong, L	851
Konginda, V	1103
Konno, K	1881
Koorbanally, NA	1847
Kosalec, I	1275
Kothari, SK	1705
Koul, M	1555
Koutsaviti, A	277
Kovačević, N	1353
Kovács, K	1379
Krasokhin, VB	773
Kremer, D	1275
Krings, U	1217
Krishnan, P	183
Krivošej, Z	1163
Krohn, K	1115,1133,1595,1907
Krohn, K	661,841
Kubo, M	327
Kukaniev, MA	1719
Kulevanova, S	21,1305
Kulhánková, A	707,1827
Kumar, N	1617
Kumar, P	1567
Kumar, S	1327
Kumari, M	1155
Kundaković, T	1353
Kunene, N	1945
Kuneš, J	607,1827
Kuo, PL	1251
Kuo, SY	1297
Kuo, SY	1671
Küpeli-Akkol, E	1433
Kurfürst, M	607
Kuroda, C	361,789
Kurokawa, M	1881
Kursar, TA	835
Kurtán, T	1799
Kusumawati, I	323
Laatsch, H	45
Labazi, N	1455
Lacaille-Dubois, MA	205,1825
Lago, JHG	243
Lajis, NH	343
Lal, B	1617
Lall, N	1585
Lalvani, A	689
Lamia, H	1531
Lamilla, C	1085
Landa, P	213
Langat-Thoruwa, C	1115
Laouer, H	1603
Lara, A	879
Larhsini, M	1447
Laskar, S	209
Lauro, MR	951
Lawal, OA	875
Lazić, ML	1855
Leach, D	1491
Leal, SM	1743
Lebib, J	409
Lee, DU	1593
Lee, DYW	1653
Lee, KH	11
Lee, MK	1267
Lee, MK	1839
Lee, NH	1193
Lefahal, M	1603
Legendre, L	1491
Lehnert, N	1217
Leitão, SG	1027
Lemberkovics, E	1379
Lempesis, K	237
Lenta, BN	1591
Leonardi, M	1499
Lepore, L	943
Li, DL	763
Li, F	353

Li, L	67,661,1055	Maia, JGS	1361	Miao, FP	1243	Najdenski, H	905
Li, M	1917	Maione, F	1205	Midaoui, ME	1455	Nakamura, K	371
Li, N	749	Majidi, L	1763	Midiwo, JO	1645	Nam, AM	379
Li, P	1801	Makani, T	409	Mihajilov-Krstev, T	1167	Nam, NH	159
Li, X	749,1285	Makarieva, TN	673	Mihajlović-Ukropina, M	1167	Nance, MR	281
Li, XJ	1129	Maklad, YA	251	Mikhova, B	905	Nand, P	1701
Li, Y	1111,1589	Malafrente, N	943,1665	Milanowski, D	175	Nanyonga, S	1945
Li, YY	1843	Maldonado, L	823	Milella, L	1735	Nascimento, NRF	871
Li, ZN	1917	Maldonado, ME	767	Milenković, M	271,277,1353	Nazir, M	1651
Liao, PC	133,267,1357,1755	Malik, A	1083	Milosavljević, S	263	Negi, A	1949
Likhititayawuid, K	41	Mallavarapu, GR	93	Mimaki, Y	187	Nejadsattari, T	259
Lima, PCWC	863	Malmiste, B	1107	Mincsovcis, E	719	Németh, K	1475
Lin, CN	133,267	Malterud, KE	1125	Minesso, P	793	Neske, A	1389
Lin, CY	901	Mancini, E	1735	Mingzhang, A	197	Netto, ADP	651
Liptaj, T	1321	Mancini, U	1315	Minh, CV	159	Neufeld, PM	1027
Liu, C	1913	Mandal, S	625	Miranda, SN	1027	Neves, IA	893
Liu, CL	1917	Mándi, A	1799	Mitaine-Offier, A	1825	Ng'ang'a, MM	1115
Liu, H	1749	Mann, TS	107	Miyamoto, T	205,1825	Nhiem, NX	159
Liu, JY	1581	Mano, JF	1577	Mo, BH	67	Ni, S	17
Liu, L	1915	Manoch, L	807	Mocek, U	175	Niamké, SL	1183
Liu, N	49	Manríquez-Torres, JJ	1225	Modak, B	759	Nicoletti, M	1003
Liu, Q	1839	Mao, AA	129	Mohammed, AMA	1573	Nidiry, ESJ	1889
Liu, X	851	Marandino, A	1735	Mohammed, HA	31	Niketić, M	271
Liu, XF	867	Marasini, BP	1117	Mohan, L	1949	Nikolaou, S	1673
Liu, XH	1243	Marco, JA	505	Moldoch, J	1293,1627	Nikoloff, N	985
Liu, Y	55,1653	Marconi, MI	667	Molina-Torres, J	857	Nikolova, M	1851
Liu, Z	55	Marcotullio, MC	1205	Mollinedo, P	1269	Nikolova, MT	575
Llor, N	515	Maric, V	73	Mondello, L	417,1817	Nile, SH	1437
Locatelli, M	1275	Marin, PD	1163	Monfort, LEF	1499	Nkeng-Efouet, PA	1591
Lockwood, B	733	Markouk, M	1447,1491	Monsalve F, Z	1085	Nkobole, N	1585
Lograda, T	87	Marongiu, B	1483,1523,1527	Montamat-Sicotte, D	689	Nogueira, PCL	907
Loizzo, MR	253	Marques, AM	939	Monteiro, HSA	871	Nojima, T	333
Loksha, AN	1899	Marques, MCA	1253	Montuno, AS	803	Nojosa, DMB	871
Lombardi, G	101	Marsik, P	213	Monzote, L	281	Novak, J	1343
London, A	235	Martin, T	1183	Mootoo, BS	785	Nugraha, AS	1953
Longjiang, Y	197	Martínez, A	287	Mora, F	1031	Nuntasaen, N	1069
López, VE	1005	Martínez, JR	1743	Mora, FD	1051	Nurmilasari,	357
Lorbeer, E	621	Martínez-Rossi, NM	193	Moraes, MEA	871		
Lorenzi, V	1159	Martini, MG	1027	Moraes, MM	1727	Ogungbe, IV	1411,1421,1637
Lorenzo, D	1711	Martino, LD	1735	Moraes, VRS	907	Ohashi, M	1769
Lou, HX	49	Martins, AMC	871	Morales, A	85,699	Ohlendorf, B	1247
Lou, L	749	Martins, CHG	777	Morales-Rios, MS	451,457	Oikawa, T	403
Louaar, S	1603	Martins, J	193	Moreira, CB	1045	Okada, Y	1621
Loubna, AS	1507	Marzouk, B	1731	Moreira, DL	939,1027	Okoh, SO	1537
Loukis, A	237	Marzouki, H	1483	Moreira, MR	777	Okusa, PN	253
Low, BS	337	Mascolo, N	1205	Moreno, DA	1305	Olaniyan, AA	875
Lozano, P	1183	Massanet, GM	491	Moreno, E	835	Oleszek, W	1293,1627
Lu, A	55	Masullo, M	1293,1627	Moreno-Dorado, FJ	491	Oliva, MM	1039
Lu, S	581,639,645	Máthé, I	235,575	Moriarity, DM	1807	Oljaca, S	1465
Lu, S	661	Mathela, CS	1373	Móricz, AM	631,657	Ollivier, E	753,1831
Lu, W	1137	Matsui, T	311	Morita, H	323	Olugbade, TA	1893
Luan, Q	1653	Matsuura, H	1801	Mou, Y	1759	Onocha, PA	1099
Lucena, M	699	Maxia, A	1523,1543,1545	Moussaid, M	1441	Opletal, L	603,607,707,1255,1827
Luciani, A	1159	Mazouz, H	1763	Moustafa, AY	1809	Opoku, AR	875
Ludwiczuk, A	315	Mbofung, CMF	1367	Moutsatsou, P	1689	Ordoñez, R	823,965
Luis, Á	1863	McFeeters, H	1421	Mozaffarian, V	259	Orhan, IE	375
Lunder, H	35	McFeeters, RL	1421	Msaâda, K	1731	Oropeza, M	97
Luo, J	851	McNeil, M	1775	Muensaen, S	1289	Orozco, LM	925
Luo, M	581,639,645	Medeiros, MRF	193	Muhammad, I	1645	Ortet, R	1179
Luo, YC	1683	Medellin-Milán, P	1925	Muhammad, Z	171	Osman, MA	1679
Luparia, M	465	Medjroubi, K	115,1603	Mulholland, DA	1573	Ott, PG	657
Luro, L	1495	Meena, R	1327	Müller, WEG	679	Otta, KH	657
Lv, D	167	Meepowpan, P	1069	Munikishore, R	1569	Oyedéji, A	1945
Lyoussi, B	1505	Mehta, GK	1327	Muñoz, MA	759	Oyedéji, O	1945
		Meier, K	1907	Muñoz, O	879	Ozgokce, F	1473
Ma, J	581,639,645	Meléndez-Rodríguez, M	451	Muñoz-Dorado, M	537	Ozkutlu, F	1469
Ma, W	167	Melkani, AB	1949	Muñoz-Serrano, A	59		
Ma, XC	781	Mellerio, G	767	Murano, E	555,593	Padalia, RC	239,1373
Ma, Y-C	581,639,645	Mencherini, T	951	Muraoka, O	1091	Paliogianni, A	1689
Maassoumi, AA	259	Mendiondo, ME	965	Muriel, KA	1715	Pan, J	1137
Maataoui, A	1455	Menezes, FS	1015	Muselli, A	1495	Pandey, A	1877
Mabrouki, F	1831	Meng, D	749	Mustafa, B	1343	Pandey, KB	1877
Macáková, K	1255,1827	Meng, X	1913			Pandotra, P	93
Macáková, K	603,607,707	Menichini, F	253	N'guessan, YT	1715	Pant, AK	1373
Machida, K	1901	Menković, N	819	Nabeta, K	1801	Paoli, M	379
Maciel-Torres, SP	1925	Menut, C	1183,1367	Nagai, A	1189	Paolini, J	1495,1763
Maggio, A	1339	Merlini, V	465	Nagarkoti, M	1637	Papoutsis, Z	1689
Maheshwari, VL	183	Merrad, B	1151	Nagashima, F	303,315	Papp, N	1417,1459,1475
Mahfoudhi, M	1531	Mharti, FZ	1505	Nair, MGG	785	Parada, K	1085

Paré, PW	1809	Quetin-Leclercq, J	813	Sabatino, L	1857,1873,1939	Shen, L	3,1677
Park, JH	1593	Quideau, S	1825	Sabini, LI	989,993,995	Shen, YM	1843
Park, y	1403	Quijano-Célis, C	247	Sabini, MC	989,993,995	Shi, XW	1129
Parra, MG	973	Quispe, R	1269	Sadamoto, K	1769	Shimizu, K	1397
Pascoal, ACRF	969,977			Saeki, D	1091	Shimizu, T	1881
Pasquale, S	1031	Raal, A	1107	Saharkhiz, MJ	1173	Shin, E	1267
Pastro, NJ	1921	Rabbitt, K	1015	Said, EM	1507	Shrestha, BB	793
Patel, AV	581,639,645	Radnović, D	1167	Saied, S	1117	Shu, XH	781
Patel, RP	1705	Radulović, N	1015	Saini, R	1155	Shuai, Q	1129
Patil, RH	183	Ragasa, CY	803	Saito, K	423	Shubina, LK	673
Paulsen, BS	1125	Rahimi, MJ	1173	Saito, Y	361	Sidana, J	1281
Pavlović, M	271	Rahman, A	323	Saitz, C	477	Siddhanta, AK	1327
Pawar , RS	1221	Rahman, AA	1597,1645	Sakagami, H	187	Siddiq, F	1083
Pazos, DC	1005	Rajae, H	1515	Sakar, MK	1321	Siddiqui, BS	1661
Pecchia, P	1665	Rajaei, P	259	Sakurai, K	403	Silchenko, AS	1075
Pecio, L	1627	Rajauria, G	1299	Salami, F	137	Šiljegović, J	263
Pedersini, C	593	Rajput, DK	1705	Salazar-Olivo, LA	1925	Silva, A	807
Pedras, MSC	617	Rakotobe, E	1259	Saleem, M	1651	Silva, AN	777
Peebles, J	1945	Rakotonandrasana, S	1259	Salgueiro, L	1523,1527	Silva, B	1051
Peine, F	679	Ram, G	93	Salvador, MJ	907,969,977	Silva, HCH	1727
Peña, A	935	Ramdani, M	87	Samoylenko, V	1645	Silva, L	497
Peng, R	67	Ramírez-Chávez, E	857	Sánchez-Suárez, J	231	Silva, SP	1577
Peng, Y	1055	Ramírez-Fernández, J	443	San-Martin, A	1073	Simão, MR	777
Peralta-Bohórquez, AF	247	Ramos, IS	973	Sanna, C	1543	Simon, S	1247
Pereira, MG	863	Ramsewak, RS	785	Sansone, F	951	Sinchaiyakit, P	829
Pereira, PS	193	Randrianaivo, R	1259	Santos, CA	1253	Singh, B	107
Pérez, M	515	Rao, BRR	1705	Santos, DYAC	1189	Singh, B	1617
Pérez-López, LA	1035	Rapior, S	1723	Santos, FM	1499	Singh, D	1567
Perin, D	555	Rasamison, VE	1259	Santos, S	807	Singh, IP	1281
Pesce, PG	593	Rasskazov, VA	773	Sasamoto, H	1835	Singh, KS	1237
Petreska, J	21,1305	Rastrelli, L	951,957,973	Sato, A	1045	Singh, S	1567
Petrović, S	271	Rateb, M	677	Sato, N	747	Singh, SA	1221
Pham, AT	1125	Reddy, MV	1569	Satou, T	1769	Siqueira, RL	863
Piacente, S	1293,1627	Ree, Tv	1907	Šavikin, K	819	Siriwatanametanon, N	627
Piaz, FD	943	Regalado, EL	1179	Sawamura, R	1881	Sithisarn, P	1289
Picchio, R	593	Reginatto, FH	913	Schenkel, EP	913	Skaar, I	35
Piccinelli, AL	957,973	Reis, RL	1577	Schipilliti, L	1009	Skaltsounis, AL	1689
Picerno, P	951	Ren, FZ	867	Schloßmacher, U	679	Skropeta, D	1921
Pino, JA	247,1179	Reynolds, WF	785	Schmidt, A	679	Smaili, T	883
Pinto, AC	667	Rezaei, MR	1173	Schmidt, J	1593	Soares, MG	243
Pinto, JEBP	1499	Rhea, J	175	Schröder, HC	679	Soares, VCG	983
Piovani, M	983	Riaz, M	1115	Schüffler, A	45	Socolsky, C	387
Piozzi, F	1347	Riaz, N	1651	Schulz, B	661,841,1595,1907	Şöhretoğlu, D	1321
Piras, A	1483,1523,1527,1543	Ricci, D	1315	Schumann, E	31	Soković, M	1163
Pires, RA	1577	Ricciardi, A	1711	Sciarrone, D	1009	Soković, M	263
Pistelli, L	1499	Ricciardi, AIA	1393	Scordino, M	1857,1939,1973	Solorzano, E	823
Pisutcharoenpong, S	1003	Ricciardi, G	1711	Scull, R	281	Sonboli, A	259
Pössel, JL	1885	Ricciardi, GAL	1393	Seguin, E	115	Song, S	1055
Pohjala, LOL	1573	Ries, C	31	Seigler, D	1085	Song, SB	1233
Polonik, NS	217	Rigano, D	1023,1339,1347	Sekeroglu, N	1433,1469,1473	Song, YC	353
Polonik, SG	217	Rim, B	1507	Sekkoum, K	1445	Song, Z	55
Pompimon, W	1069	Rios, N	1031,1051	Selim, RM	1809	Souza, MC	1045
Pongbangpho, S	829	Ríos-Chávez, P	857	Senatore, F	1023	Souza, MHL P	871
Ponomarenko, LP	773	Ristić, M	1167	Senatore, F	1339,1347	Spadafora, C	835
Poonam	1155	Rivera-Becerril, E	457	Şener, B	375	Speicher, A	393
Porcedda, S	1523,1527	Rivière, C	813	Sepp, J	1107	Sreelekha, M	1595
Porta, A	465	Rizvi, SI	1877	Sepúlveda, B	1073	Sriprang, S	829
Porter, R	1775	Rodilla, JML	497	Sepúlveda-Arias, JC	925	Sritularak, B	41
Potdar , YA	1221	Rodrigo, G	1269	Setzer, W	1945	Stanojković, T	819
Potenza, L	1315	Rodrigues, NVFC	863	Setzer, WN	119,281,695,875,1411	Stashenko, EE	1743
Pouységu, L	1825	Rodríguez A, OE	947	Setzer, WN	1421,1549,1637	Stefanello, MEA	969,977
Povichit, N	829	Rodríguez-García, I	537	Setzer, WN	1719,1807	Stefkov, G	21,1305
Prado, LAM	193	Rojas, J	85,699,935,1031	Sewald, N	1591	Stefova, M	21,1305
Pragadheesh, VPPS	1333	Rojas, JJ	287	Shaaban, KA	45	Steinert, M	1907
Prakash, I	175,1059	Rojas, L	85,699	Shaari, K	343	Sternier, O	1269,1321
Prakash, O	1373	Rojas, LB	127,1051,1825	Shafaghath, A	137,275,1351,1739	Stević, D	263
Prasad, K	1327	Rojas, YCD	127	Shafaghathlonbar, M	275	Stévigny, C	253
Prata, APN	907	Romane, A	1511	Shafi, PM	1595	Stochmal, A	1293,1627
Pretsch, A	677	Román-Marín, LU	1225	Shah, MR	171	Stocker, P	1151
Pribylova, M	213	Romoff, P	243	Shah, S	1117	Stonik, VA	587,673
Prota, L	951	Rosselli, S	1023,1339	Shan, T	1759	Su, DH	1581
Punyanitya, S	1069	Rout, PK	1333	Shang, MY	11	Su, YC	133,267,901,1357,1755
Püssa, T	1107	Rovirosa, J	1073	Shao, CL	1239	Suárez, AI	97
Putra, DP	349,357	Ruiz, I	957	Sharma, DK	1929	Suárez-Castillo, OR	451
		Ruparel, P	733	Sharma, SN	1929	Sudhana, JM	1569
Qi, J	1913	Russo, A	1023	Sharma, U	1567	Sudsai, T	1095
Qian, D	17			Sharopov, FS	119,695,1719	Sugijanto, NE	677
Qian, PY	1239	Saadat, F	1173	Sharry, S	985	Suka, Y	333
Qian, YT	867	Saba, N	1	Shen, CC	803	Sukdee, S	1069

- Sulc, A1475
 Sultan, M1651
 Sun, Y1881
 Sun, WJ1589
 Sutil, SB995
 Soutour, S1479
 Suttajit, M829
 Suzuki, R1621
 Suzuki, T311,333,1631
 Svobodova, B213
 Świtalska, M367
 Syafni, N349
 Szabó, K235
 Szajwaj, B1293,1627

 Tabanca, N701
 Takahashi, K1801
 Takahashi, M1769
 Takiguchi, K361
 Talamond, P1885
 Taleb, S1445
 Tammela, PSM1573
 Tan, JW763
 Tan, NH1263
 Tan, RX353
 Tanaka, C205,1825
 Tanaka, K7
 Tanaka, R1091
 Tang, W327
 Tang, XZ1243
 Tantry, MA1597
 Tao, MH763
 Taoubi, K753
 Tareen, RB179
 Tasdemir, D689,1697
 Tasic, S1465
 Tatsadjieu, LN1367
 Tava, A101
 Tchimine, MK1905
 Techasakul, S1103
 Tedone, L417
 Teh, CH337
 Teixeira, VL651,667
 Tekwani, BL1645
 Terent'eva, NA773
 Terevsky, N645
 Tesarova, H213
 Tesch, NR1031
 Tešević, V263
 Tewtrakul, S1095
 Thaisuchat, H1069
 Thomas, OP1179
 Thomson, M1921
 Tia, EV1183
 Tian, Y781,1581
 Tillett, S97
 Timmers, M1605
 Tistaert, C813
 Todorova, M905,1851
 Tolosa, D1389
 Tomás-Barberán, FA1305
 Tomi, F1159,1479
 Tomi, P1495,1763
 Tomiyama, K403
 Tommasi, ND943
 Tonn, CE989
 Tori, M361
 Torre, G1817
 Torrenegra G., RD947
 Torres, AM1393,1711
 Torres, CV993
 Torres, NW1035
 Torres-Valencia, JM1225
 Tosi, S767

 Touzani, R251
 Toyama, MH983
 Traulo, P1873,1939
 Trendafilova, A1851
 Trozzi, A1009
 Trujillo-Serrato, JJ457
 Tsamo, E1591
 Tseng, YH133,267
 Tsubono, K1621
 Tuncturk, M1473
 Tuncturk, R1473
 Tundis, R253
 Twardowschy, A1253
 Twigg, PD1411
 Tyihák, E611,631,657,719
 Tzakou, O237,271,277

 Uesawa, Y1621
 Ullah, R171
 Uniyal, GC1333
 Urban, S1605
 Urbanová, K603,1255
 Urzúa, A759
 Üstün, O375
 Usubillaga, A127,935

 Vachnadze, V1831
 Vadlejch, J213
 Vajs, V1163
 Valant-Vetschera, KM621
 Valdez-Calderón, A1225
 Valiente, J223
 Valterová, I1827
 Van, KP813
 Van, MC813
 Vanegas-López, C247
 Vangelisti, R999
 Varughese, T835
 Vasas, A575
 Vasconcelos, MH807
 Vásquez, L97
 Velasco, J1031,1051
 Velázquez-Jiménez, R1225
 Veljković, VB1855
 Veloza, LA925
 Velozo, LSM939
 Veneziani, RCS777
 Vera, N823
 Verdeil, JL1885
 Veres, K235
 Vérité, P115
 Verma, RS239
 Versiani, MA1897
 Victório, CP1045
 Vidari, G465,767
 Viegi, L999
 Vilegas, W983
 Villafañe, E1389
 Villagomez, R1269
 Villarreal, B985
 Villasmil, T935
 Villatoro-Pulido, M59
 Vincieri, FF79
 Vitkova, A905,1851
 Vivekanand1373
 Vogler, B1637,1807
 Vučković, I263,1163
 Vujisić, L1163
 Vuko, E1385

 Walker, LA1645
 Walker, TM1807
 Wang, CY1239
 Wang, C55,1913

 Wang, CT1683
 Wang, D581,639,799
 Wang, EIC 133,267,901,1357,1755
 Wang, G1111
 Wang, J1749,1759
 Wang, K3,1599
 Wang, LN49
 Wang, LS1141
 Wang, MJ1589
 Wang, MX1131
 Wang, N3,1915
 Wang, SJ1915
 Wang, T1799
 Wang, X11,65,679,1285,1799
 Wang, X-Q645
 Wang, Y167
 Wang, YY49
 Wang, Z1653
 Wangenstein, H1125
 Watanabe, W1881
 Watanabe, Y1901
 Wattanapiromsakul, C1095
 Wawrzęczyk, C367
 Wedge, DE701
 Wei, F1913
 Wei, Q1581
 Wei, XY763
 Wentao, G197
 Wessjohann, L1593
 Widowitz, U527
 Widyawaruyanti, A323
 Wiens, M679
 Wiese, J1247
 Wietrzyk, J367
 Wilson, RC1411
 Wojnicz, D149
 Wollenweber, E15
 Wu, B53
 Wu, J1137,1599

 Xia, LJ867
 Xia, X1913
 Xia, Y67
 Xie, BB11
 Xie, G1799
 Xin, XL781,1581
 Xu, I645
 Xu, JJ1263
 Xu, M3
 Xu, Y1239

 Yadav, A239,1333
 Yagi, Y303
 Yamada, T1091
 Yamaguchi, K349
 Yamamoto, R1881
 Yamazaki, M423
 Yan, J763
 Yáñez, C1031
 Yang, F1749
 Yang, HY867
 Yang, N17
 Yang, SY1233
 Yang, WL227,855,1671
 Yang, Y65
 Yao, JH781
 Yaoita, Y1901
 Yaouba, A1367
 Yasmeen, S1083
 Yasukawa, K1881
 Yatsunami, K1621
 Yazıcı, G1469
 Ye, W1111
 Yeh, YT1251

 Yeh, YT227,855
 Yesilada, E1697
 Yi, Y661
 Yin, Y1835
 Yokosuka, A187
 Yoo, HS1839
 Yoshida, M1621
 Yoshihara, T1801
 Yousfi, M1151
 Youssef, A1515
 Yu, H167
 Yu, Z1749
 Yuan, W1913
 Yue, S197
 Yuen, KH337
 Yurchenko, EA1075

 Zacchino, S931
 Zahoor, A179
 Zaini, NC323,677
 Zampini, C965
 Zanin, JLB243
 Zanon, SM989,993,995
 Zanon, G465
 Zapata-Bustos, R1925
 Zaporozhets, TS587
 Zapp, J31
 Zarea, K137
 Zareen, S343
 Zavadil, S603,1827
 Zdunić, G819
 Zellagui, A883
 Zeng, GZ1263
 Zeng, J223
 Zeng, XM67
 Zepeda, LG429
 Zhan, J223
 Zhang, AL1129
 Zhang, B65
 Zhang, BJ781
 Zhang, DS1581
 Zhang, H53,1599
 Zhang, HJ1917
 Zhang, J1653
 Zhang, M1913
 Zhang, MX1917
 Zhang, T1323
 Zhang, W661
 Zhang, WM763
 Zhang, X65,1111,1323
 Zhang, Y1759
 Zhang, YM1263
 Zhao, J1749,1759
 Zhao, L1683,1915
 Zhao, Y1589
 Zheng, J1581
 Zhou, JH1589
 Zhou, L1055,1749,1759
 Zhou, SL1131
 Zhou, YM1239
 Zhu, F1137
 Zhu, J867
 Zikr-Ur-Rehman, S1897
 Zinad, DS45
 Zingali, RB961
 Zivanovic, A1921
 Zlatković, B1167
 Zlatković, S1353
 Znini, M1763
 Zollo, PHA1367
 Zomorodian, K1173
 Zupkó, I575

Natural Product Communications

Volume 6 (1-12) 2011 Keyword Index

- A2780 1259
AAS 1473
Abies spectabilis 793
Abietane diterpenes 3,575,1229
Abrus schimperi 1645
Absidia coerulea 1581
Absolute configuration 429,759,1225
Absolute stereochemistry 673
Acacia nilotica 1451
Acanthaceae 1569
Acanthopanax koreanum 1233
Acanthopanax trifoliatum 1289
Acaricidal activity 893
Acetoxy-3-phenyl-propiofenone 621
Acetoxypinoresinol glucoside 753
Acetylcholinesterase 603,607,1089,1255,1827
Acetylenes 361
Acharan sulfate 1403
Achillea biebersteinii 701
Achillea distans 905
Achillea millefolium L. 1527
Achyrocline satureioides 995
Acmella radicans 857
Acne 1701
Acne vulgaris 1193
Aconitum 527
Acremonium roseum 367
Acremonium striatisporum 1063
Action mechanism 287
Acuminophenone A 1269
Acyclic diterpenes 323
Acylated anthocyanins 1631
Acylated neohesperosides 1321
Acylglucosylated sterols 767
Acylphloroglucinols 387
Adult activity 701
Adults 887
AEDA 403
Aedes aegypti 701
Affinin 857
Aflatoxin 657
AFLP 1679
Aframomum latifolium 1183
Agarwood 763
Ageratum conyzoides 1263
Agrobacterium-mediated transformation 1665
Agro-homeopathy 1499
Agro-waste 183
Ailanthone 593
Ailanthus altissima 593
Albizia chinensis 1617
Alchemilla mollis 1851
Aldehydes 1821
Aleppo region 79
Algeria 1511
Alkaloids
187,515,857,939,1253,1593,1595,1831
Alkylanthracenic acids 1133
Alkylating agents 457
Allium cepa L. test 995
Allyldimethoxybenzene 1147
Allylsilanes 465
Aloysia citriodora 1711
Aloysia gratissima 1499
Aloysia triphylla 1039
Alpine Grasses 101
Alternaria sp. 1913
Altruronic acid 1063
Alyxia reinwardtii 677
Alzheimer's disease 607,1089,1411
Amaryllidaceae 187,603
Ambergris related compounds 477
Amblypelta nitida 409
Ambrinol 439
Amides 227,1293,1297,1591
Amorphaquinone 1645
Amplexicine 1597
Amsonia illustris 235
Amylase 1585
Anacardiaceae 1133
Analytical techniques 733
Andrographis paniculata 1929
Andrographis wightiana 1569
Andrographolide 1929
Anethole 141
Angiogenesis 943
Annona pickelii 907
Annona salzmannii 907
Annonaceae 1715
Anthocyanins 35,149,423
Anthrapyran 841
Anthraquinone profile 1275
Anthriscus caucalis 1147
Anthriscus cerefolium 1147
Anthriscus nemorosa 271
Anti-bacteria 1953
Antibacterial activities 1705
Antibacterial activity 163,699,823,905,1023,
1031,1051,1163,1299,1351,1379,1505,1507,
1511,1515,1705,1949,
Antibacterials 597,1221,1421,1573
Anticandidal properties 1491
Anticariogenic activity 777
Anticorrosive activity 1425
Antifungal activities 1705
Antifungal activity 1527,1531,1763,1821
Antifungal activity 433,905,931,1163,1357,
1361,1367,1523
Antifungals 1221,1487,1889
Anti-fungi 1953
Antigenotoxicity 59
Anti-inflammatory 627,1441,1433
Anti-inflammatory activity 1437
Anti-malaria 1953
Antimalarial activity 323,337
Antimicrobial activity 1027,1035,1167
Antimicrobial activity 1173,1275,1353,1447,
1519,1701
Antimicrobial activity 133,141,149,247,259,263,
267,271,277, 353,661,841,901,907,965,1739,
1749,1759,1905
Antimicrobial activity
Antimicrobial agents 193
Antimicrobial effect 275
Antimicrobials 1121,1645,1925
Antimutagenicity 1269
Antimycobacterial 1743
Antimycotic activity 1039
Antinociception 863
Antinociceptive 1433
Antioxidant 597,627,829,907
Antioxidant activities 55,1537,813,1229,1275,1285
Antioxidant activity 133,253,375,651,823,845,
901,965,1305,1315,1373,1447,1507,1597,1617,
1739,1749,1847,1851,1863
Antioxidant capacity 1269,1299
Antioxidants 59,785,1179,1361,1433,1441,
1487,1673,1683,1877
Antioxidative activity 213
Antiparasitic 1645
Anti-parasitic activity 835
Antiproliferative 575,1259
Antiprotozoal 973
Antiprotozoal activity 1697
Antiradical activity 1367
Anti-snake venoms 1393
Antitumor drugs 555
Antiviral activity 993,1881
Antivirals 1573,1953
Anti-wood-decay fungal activity 133,267,1357,
1755
Anxiolytic-like effect 1769
Aphanamixis grandifolia 323
Apiaceae 115,263,491,883,1147,1163,1179
Apigenin-8-C- β -D-glucopyranoside 1131
Apiole 1361
Apocynaceae 235
Apoptosis 187,947,1653
Aporphine 1251
Aqueous extract 993
Aquilaria sinensis 763
Arbutin derivatives 1661
Ardeemin analogs 807
Argentinean propolis 823
Aristolactams 939
Aristolochiaceae 11
Aromatase inhibition 31
Aromatic and medicinal plants 1455
Aromatic compounds 361
Aromatic glycoside 1897
Artemisia annua var. CIM-Arogya 239
Artemisia scoparia 119
Artocarpus 1397
Artocarpus lakoocha 41
Asarum ichangense 11
Ascaridole 281
Asclepiadaceae 803
Asimafoetidinol 209
Aspergillides 505
Aspergillus awamori 1933
Aspergillus terreus 183
Asperula lutea subsp. *rigidula* 237
Asteraceae 1059,1117,1339,1603,1665,1945
Asteraceae 119,123,127,175,251,259,627,767
Asteriscus graveolens 1763
Astrortricoumarin 1259
Atherogenic Index 1005
Atlantic rain forest 1045,1727
Autonomic arousal 1199
Autophagy 187
Avicennia marina 1137
Azaderivatives 1813
B16 melanoma cells 1397
Baccharis articulata 993

- Barley malt sprouts 73
 Baylis-Hillman adducts 1221
 Beckmann rearrangement 1813
 Beekeeping 1461
 Behavioral arousal 1199
 Beijing genotype 1743
Beilschmiedia obscura 1591
 Belladine 1827
Bellardia trixago 439
Bemisia tabaci 1183
 Beneficial effect,
 Benzaldehydes 1023
 Benzofurans 1913
 Benzyl acetate 1333
 Berberidaceae 1089
 Berberine 1089
Berberis darwinii 1089
 Bergamot 1199
 Berries 149
 Betalain 1425
Betonica officinalis 1343
 Bibenzyl 303
 Bicyclic alicyclic compound 311
 Bicyclogermacrene 275
 Bicyclononalides 785
 Biflavonoids 1111
 Biflavonyloxymethane 625
 Bignonaceae 957
 Bioactive compounds 823
 Bioactivity 1129,1239
 BioArena 631
 Bioassay 1657
 Bioassay-guided separation 851
 Bioassay-oriented study 943
 Bioautography 701
 Bioavailability 337
 Biochemical characterization 73
 Biochemical immunization 611
 Biocide 1183
 Bioconjugates 1877
 Bioconversion 1217,1801
 Biological activity 707,1137,1205,1595
 Biological detection 719
 Biologically induced mineralization 679
 Biomass and essential oil yields 1705
 Biomimetic cyclization 465
 Biomimetic synthesis 537
 Biomineralization 679
Biomphalaria peregriana 387
 Biosynthesis 555,773
 Biotransformation 367,1581,1805
 Bis(3,4-dimethoxyphenyl)cyclobutane 49
 Bis(3,4-dimethoxyphenyl)cyclobutane 49
 Bis(benzyls) 393
 Bisabolene 123,1537
 Bisabolene derivatives 1225
 Bislactone 497
 Bivariate least squares detection limits 1939
 Black condensate 1577
Blechnum chilense 1085
 Blood pressure 871
 Boerharotenoid A 1651
 Boerharotenoid B 1651
Boerhavia repens 1651
 Bombacaceae 1897
Bombax cieba 1897
 Bone marrow 989
 Botcinin 443
 Botcinolide 443
Bothrops neuwiedi diporus 1393
Botrytis cinerea. 443
 Box Behnken model 1933
 Brassica 65
 Brassica vegetables 1299
 Brazilian Cerrado 983
 Brazilian pepper tree 887
 Brazilian-lavender 1499
 Brine shrimp lethality 1915
 Brine shrimp toxicity 1921
 Broccolini 65
 Bromine/carbon tetrachloride 451
 Bromoindole natural products 451
 Brown algae 863
 Bufadienolides 1581
Bulbophyllum kwangtungense 53
 Burseraceae 1727
 Butenynyl-bithienyl 1035
 Butyrolactones 367
 Butyrylcholinesterase 603,607,1827

Cachrys libanotis 115
 Cactus pear 1425
 Cadin-1,4-dienes 699
 Cadinene 271
 Cadinol 251,1755
 Caffeoyl quinic acids 1289
 Caffeoylmussaenosidic acid 747
 Caffeoylquinic acids 1885
 Calcium oxalate 1445
Calocedrus formosana 133
Calotropis gigantea Linn 803
Calycolpus moritzianus 943
Camellia sinensis 371,1155
 Camphor 239,1705
Campomanesia adamantium 969
 Camptothecin 197
 Cancer 1925
Candida 1173
 Candidiasis 1039,1567
 Capillene 119
 Caprifoliaceae 1901
Capsicum annum 227
 Car-3-en-5-hydroperoxide 1217
 Car-3-en-5-one 1217
 Car-3-ene 1217
 Carbohydrate-degrading enzymes 1403
 Carbon monoxide insertion 311
 Cardiovascular disease 961
 Carene 875
 Carissone 621
 Carotenoids 1817
 Carrageenan 1327,1543
Carramboa littlei 127
 Carvacrol 137,1173,1719
 Carvone 925,1373
 Caryophyllaceae 275
 Caryophyllene 85,111,247,267,699,1145,1351,
 1357,1385,1949
 Caryophyllene-oxide 1385
 Cascade reaction 465
Cecropia 913
 Cedrene oxide 97
Cedrus atlantica 379
 Cell cycle 1671
 Cembrane diterpenes 1809
Centaurea 163
Centaurea gracilentia 1339
Centaurea microcarpa 1603
Centaurea ovina ssp. *besserana* 1339
 Central nervous system activity 913
 Cercosporin 835
Cestrum ruizteranianum 1825
 CF₃COOH 935
 Chalcone 823
Chamomilla suaveolens 1107
 Chemical composition 107,1183,1367,1507,
 1775
 Chemical investigation 53
 Chemical markers 1711
 Chemical polymorphism 1483
 Chemometrics 1015,1929
 Chemoprevention 799
 Chemosystematics 315
 Chemotaxonomic significance 1603
 Chemotaxonomy 667,1393
 Chemotypes 315,423,1511
Chenopodium ambrosioides 281
 Chinamine A 217
 Chinotto 1857
Chionanthus virginicus 753
 Chiral synthon 477
Chlidanthus fragrans 603
 Chlorination 223
 Chlorogenic acids 639,1627
 Cholestane derivatives 1821
 Cholinesterase inhibition 375
 Cholinesterase inhibitory activity 253
 Chromene 537
Chromolaena leivensis 947
Chrysactinia mexicana 1035
 Chrysanthemylacetate 1147
 Chrysin 31
 Chrysoeriol 205
Chuquiraga straminea 965
 Chymotrypsin 1117
Cichorium intybus 1117
 Cineole 93,107,111,137, 239,259,,1351,1483
Cinnamomum burmannii 1297
 CIP rules 429
 Circular Diachroism 1645
Cistus ladaniferus 1159
 Citral 925
 Citronellal 925
 Citrus 1857
Citrus aurantium 1009,1389
Citrus bergamia Risso 951
 Citrus juice 1495
Citrus limon 1389
 Citrus peel 1495
Citrus sinensis 875
 Claisen-Johnson rearrangement 167
Clausena anisata 1367
Clematis 1631
 Clementine 1495
 Clerodane diterpenoid 1069
 Clerodanes 327,333,497
Clerodendrum kaichianum 3
 Clusiaceae 1267
 Cluster analysis 695,1151
 CMV 1385
 Cnicin 163
 Coccolithophores 679
 Coccoliths 679
Codium fragile 1243
 Cold aqueous extract 995
Colletotrichum 1131
Colletotrichum 701
 Colors 35
 Common diseases 79
 Compositae 1,175,1059,1263
 Conjugated acid derivatives 1221
Conocephalum japonicum 49
 Conotoxins 1403
 Contact toxicity 887,1945
 Content-activity relationship 1769
 Convulvulaceae 1433,1889
Cordia gillettii 253
Cordia platythyrsa 1135
 Cordycepin 1839
Cordyceps militaris 1839
Coriandrum sativum 1465
Corioliolus versicolor 1129
 Cork 1577
Corydalis cava 607
 Cosmetics 1397
 Cosmosiin 961
Cotula cinerea 1491
 Coumarins 1107,1723,1917
 Coupling reaction 1323
 CQA 1885
 Crosslinked HA 555
Crotalus durissus cascavella venom 871
Croton gossypifolius 97
Croton jacobinensis 893
Croton micans 893
Croton muscicapa 893
Croton rhamnifolius 893

- Cryptococcosis 1567
 Crystal structure 759,1589
 Crystallography 1237
 Cucumariosides 1075
 Cumaside 587
 Cupressaceae 87,133
Cupressus atlantica 1519
Cupressus dupreziana 87
 Curcumin 1877
Cuscuta arvensis 1433
 Cyanidin glycosides 1631
 Cyanobacteria 689
 Cyanogenic glycoside 1091
 Cyclic voltammetry 597
 Cyclodepsipeptide 1247
 Cyclopeptides 1843
 Cyclostachine acid derivative 1591
 Cymene 275,1353
 Cytogenotoxicity 995
Cytospora sp.661
 Cytotoxic 303,803,1573,1645,1925
 Cytotoxic activity 3,49,59,145,163,187,281,353,505
 Cytotoxicity 813,819,835,947,983,993
 1263,1637,1671,1807,1921

 DAD-APCI-MS 1817
 Damascone 465
Daucus sahariensis 883
 Decarboxylase activity 857
 Deep-water sponges 1921
 Degenerative diseases 707
Dendrobium officinale 867
 Dendrogram 1499
 Density functional theory 1677
 Density functional theory calculations 759
 Deoxyandrophthalide 781
 Deoxycholic acid 1237
 Depsilairdin 617
 Dereplication 1605
Derris malaccensis 1103
 Development stages 1465
 Di[but-3(E)-en-2-onyl]naphthalene 1117
 Diabetes 201
 Diabetes mellitus 1585
 Dialkylloxindoles 457
 Diarylheptanoids 1881
 Diastereoisomers 1091,1653
 Dicafeoyl quinic acids 1885
 Dichloronaphthazarin 217
Dichotella fragilis 1239
 DiCQA 1885
 Dietary supplement 1003
 Dihydrochalcone 621
 Dihydroflavopereirine 349
 Dihydromethoxyldibenz[b, f]oxepin 53
 Dihydroprenylmethylcoumarin 1259
 Dihydroresorcylic acid 223
 Dihydrostreptazolin 353
 Dihydrothymoquinone 213
 Dihydroxydeepoxysarcophine 1809
 Dihydroxy-methoxyacetophenone 1129
 Dihydro- γ -ionone 439
 Dill apiole 1151
 Dimethoxy citral 93
 Dimethoxyvulpinic acid 357
 Dimethylchromene glucopyranoside 1263
 Dimethyldioxirane 925
Dionysia 621
Dioscorea bulbifera 1069
 Dioscoreaceae 1069
 Dioxime γ -lactam 1135
 Dioxopiperazine 617
 Direct bioautography 1379
 Distillation 1939
 Diterpene glycosides 1059
 Diterpene glycosides 175
 Diterpenes 1253
 Diterpenoid alkaloid 1589

 Diterpenoids 327,333,497,1205
 Diversity 361,789
 DNA barcode 67
 DNA markers 1679
 DNA polymorphism 1679
 DNA protective 41
 Dodecadienolactone 275
 Double effect 631
 Double immune response 611
 DPPH 129,213,819,969,977,1125,1447,1673
 Drechslerine B 311
 Drimane terpenoids 477
 Drimanes 315
 Drimenol 477
Drimys brasiliensis 243
Drimys winteri 879
Drosophila 59
 Drought 1559
 Drug candidates 1403
 Drugs 287
Dryopteris wallichiana 387
 Drypetdimer A 1115
Drypetes gerrardii 1115

 Ecdysterone 707
 Echinacea 639
Echium vulgare 1475
 Edema 1543,1545
Elaphoglossum 387
 Electrospray ionization 7
 Elemene 85
 Ellagitannins 1125
 Embryos 773
 Enantiomeric distribution 403
 Endophytic fungi 1131,1907
 Endophytic fungus 661,677,763,835,1759
Enterobacter aerogenes 1159
Ent-kaur-15-en-19-oic acid 935
Ent-Kaur-16-en-19-al 127
Ent-kaur-16-en-19-oic acid 935
Ent-kaurane diterpenes 1233
Ent-Kaurenoic acid derivatives 777
 Ent-labdanes 781
 Environmental and seasonal effects 1711
 Enzyme inhibitors 1921
 Epidermis 1475
 Epimedin 581
Epimedium 581
 Epiphytic fungus 1137
 Epitaondiol 1073
 Epoxidation 925,1141
 Epoxydecenal 403
 Epoxydones 1905
 Epoxyocimene 409
 Epoxyoleananolide 1573
 Erva-mate 651
Erythrina sigmoidea 1099
 ESI-FTICR-MS 1593
 ESI-MS 977
 Essential oil 85,97,111,115,123,129,133,141,145,243,251,253,263,267,271,275,277,409,699,883,901,905,1015,1031,1035,1039,1051,1145,1147,1151,1163,1173,1183,1189,1193,1339,1343,1347,1353,1357,1373,1385,1465,1487,1505,1515,1519,1523,1527,1715,1743,1749,1755,1763,1775,1945
 Essential oil composition 93,101,119,127,137,235,247,281,695,875,879,893,907,1179,1351,1361,1479,1491,1511,1705,1719,1727,1949
 Essential oil plants 1559
 Essential oils 239,887,1027,1167,1379,1483,1507,1735
 Estragol 1147
 Estragole 925,1147
 Ethnobotanical survey 1459
 Ethnopharmacological use 931
 Ethnoveterinary medicine 999
Eucalyptus cinerea 107
Eucalyptus loxophleba 1281

 Eudesmane sesquiterpene 1
 Eudesmol 267,1339,1357
Eugenia 977
 Eugenol 1705
 Euglobals 1281
Eupatorium heterophyllum 361
Eupuntia fraudatrix 1075
 Euphorbiaceae 97,813,1115
Eurycoma longifolia 337
 Excelside B 753
 Extraction 439,1577,1855
 Exudates 621,845

 Fabaceae 625,1099
Fadogia tetraquetra 1573
 Farnesene 1333
 Fatty acids 237,667,1739,1921
 Ferns 1723
Feronia elephantum 141
 Ferromanganese crusts 679
 Ferruginol 133
Ferula assa-foetida L. 209
Festuca nigrescens 101
 Ficalloside 159
Ficus callosa 159
Ficus carica 375
Ficus glomerata 1095
 Fig 375
 Filifolinol acetate 759
Fimetiaria rabenhorstii 763
 Fingerprinting 1929
 Flavone dimer 1115
 Flavone glucosides 1627
 Flavones 1569
 Flavonoid content 1441
 Flavonoid glycosides 1599,1605,1851
 Flavonoid-7-O-glycosides 21
 Flavonoids 15,149,205,237,651,813,819,913,947,965,1107,1289,1603,1855,1857,1863
 Flavonoids Flavonols 651,1631
Flos Lonicerae 639
 Flower colors 1631
 Flowering 1455
 Flowers 1155
 Fluoro-*p*-anidine 1537
 Foodborne disease 1167
 Formaldehyde 657
 Formaldehyde cycle 611,631
 Formaldehyde generator 611
 Formoxanthone C 1269
 Formylpyrrole-4-acrylamide 353
 Fougere 1723
 Fragmentation 7
 FRAP129
 Fraud 1873
 Free radical scavenging 597
 Fresh juice 951
 Friedelin 1577
 Fruiting 1455
 Fruits 883
Frullania 303
 FTIR 1445
Fuchsia 35
 Full factorial experiments 1855
 Fumariaceae 607
 Fumigant 1945
 Fumigant toxicity 887
 Fungal biosynthesis 1801
 Fungal metabolites 1905,1907
 Fungicidal activity 1917
 Furan 361
 Furocoumarins 1857
 Furoquinoline alkaloids 1847
Fusarium graminearum 1243
Fusarium heterosporium 1805
Fusarium oxysporum 367
Fusarium tricinctum 1759

Garcinia cambogia 1933

- Garcinia mangostana* 1267
 Gas chromatography 1939
 Gas chromatography-quadrupole mass spectrometry 1333
 Gastroprotection 1253
 Gastroprotective activity 1073
 GC 141,1009,1145,1343,1519
 GC/MS 101,141,253,603,1145,1155,1255,1505, 1511,1519,1827
 GC/MS-LRI 1009
 GC-C-IRMS 1009
 GC-FID 235
 GC-MS 115,129,235,857,1179,1343,1487,1499, 1701
 Gecko, adulterants 67
Gekko gekko 67
 Genetic variability 87
 Genotoxicity 989
 Geraniaceae 15,1321
 Geranial 93
 Geraniol 93
 Geraniol derivatives 465
Geranium 15,1321
Geranium purpureum 1321
 Geranylacetoxy coumaric acid 343
 Geranylisovalerate 251
 Germacrene 303
 Germacrene D 85,115,127,247,271,275,1519
 Germanicol 1505
Gibberella zeae 1243
 Ginger oil 93
Ginkgo biloba 1131
 Ginsenoside 581
Glechoma longituba 17
 Global markets 555
 Glucose esters 1893
 Glucosidase 1585
 Glucosidase 201
 Glucosidase inhibitors 851
 Glucosides 361
 Glucosinolates 65
 Glucosylation 223
 Glutanol 171
 Glycerides 371
 Glycoside-bound volatiles 107
 Glycosides 21,673,1679
 Glycyrrhiza radix 7
 Gorgonian 1239
 Green tea seeds 371
 GSH 1315
 Gurmar 1679
 Guttiferae 85,699
 Guttiferone K 1269
Gymnema sylvestre 1679
Gynura pseudochina var. *hispida* 627
Gypsophyla bicolor 275

 H⁺, K⁺-ATPase inhibition 1253
 Haemolysis 1921
 Haemostasis 961
 Halimane 497
Halimondendron halodendron 1749
 Hallucinogen 333
 Hardwickic acid 327
 Harvesting 1455
 HCHO 631
 Headspace (HS) 1155
 Heavy metals 1473,1559
 Heck arylation 537
Helichrysum plicatum 819
Heliotropium filifolium 759
 Hengduan Mountains 789
 Herb 1461
 Herbal medicinal products 733
 Herbal medicines 1001
 Herbal preparations 753
 Herbicidal activity 1531
 Herbicide activity 593
 Herbs 1473

Herpes suis virus 993
Hesperozygis 1027
 Hexadecanoic acid 1023
 Hexadecanol 271
 Hexahydrocurcumin 1671
 Hexahydrofarnesylacetone 1347
 Hexenal 1723
 Hexenoic acid 1723
 Hexenyl acetate 1333
 Hierarchical digraphs 429
 Highly sensitive analytical methods 527
 Histochemistry 1045
 Histological variance 1475
 HIV-1 integrase 1095
 HL-60 cells 187
 Holothurians 587
 Homosines 1905
 Honokiol 1121
 Horminone 575
 HPLC 197,753,1289,1299,1657,1929
 HPLC profile 639,645
 HPLC-DAD 1831
 HPLC-DAD/ESI-MS 969
 HPLC-MS 1599
 HPLC-NMR 1605
 HPLC-UV/DAD 977
 HPTLC 1003
 HQA 1885
 HRGC-MS, LRI 417
 HSC-2 cells 187
 HSC-T6 cells 1267
 HSV-1 1881
 Human tumor cell lines 807
 Huntington's Disease 1411
 Hyaluronan 555
 Hyaluronic acid 555
 Hybrids 1495
 Hydrodistillation 107
 Hydrogen peroxide 1141
 Hydrolysis 65
 Hydroxycinnamic acids 21
 Hydroxycinnamoyl quinic acid 1885
 Hydroxycyclonerolidol 1763
 Hydroxylation 781
 Hydroxymethylthioethanol 1131
 Hydroxyquinoline 597
 Hydroxyursenolide 1573
Hymeniacidon perleve 1915
 Hypericaceae 1731,1739
Hypericum 1735
Hypericum ericoides ssp. *roberti* 1731
Hypericum perforatum 1731
Hypericum scabrum 1739
 Hypoglycemia 1679
 Hypoglycemic activity 913
 Hypolipidemic effect 1005
 Hypotensive activity 913
 Hypotensive effects 871
Hyptis 1775

 Icacinaceae 197
 Icariin 581
 ICP 1469
 Iksut 1433
 Ikuskut 1433
 IL-6 1543,1545,1689
Ilex canariensis 661
Ilex paraguariensis 651
 Illecebraceae 205
 Immunostimulating activity 867
In vitro disaggregation tests 951
In vitro dissolution tests 951
in vitro evaluation 231
 Indian valerian 129
 Indole 515,1333
 Indole alkaloids 193
 Indonesian medicinal plants 1953
 Inflammation 1193
 Infusion and transfusion OPLC 719

 Inhalation 1769
 Insecticidal activity 1085,1917
 Insolation 1465
 Intermediate derivatization method 1917
 Interspecies comparison 1769
 Intraspecific variation 243
 Invertebrate source 1403
 Ionone 465
Ipomoea carnea 1889
Ipomoea fistulosa 1889
Ircinia sp. 1237
 Iridals 789
 Iridoid glucosides 747
 Iridoids 237
Iris delavayi 789
 Iron complex 597
 Irones 465
 Isoabienol 277
 Isochrysanthemide ester 251
 Isocoumarins 45
 Isoferulic acid 1285
 Isomerization 1889
 Isoplumericin 1567
 Isothiocyanates 65
 Isoxanthochymol 1269
Ixeris sonchifolia 1055
 ixoside 747

Jacaranda mimosaefolia 1893
 Jacarandol 1893
 Jacaranoside 1893
 Janceolaroside A 749
 Janceoside A 749
Jasminum lanceolarium 749
Jasminum sambac 1333
Juniperus phoenicea L 1515

 Kabosu 403
 Kaempferol 201
 Kaempferol-3-O-rutinoside 201
 Kenland 1657
 Khenifra 1455
Knautia arvensis 1627
 Kolavelool 327
 Kolavenol 327
 Kosovo 1343

 Labdane 497
 Labdane diterpenoids 1569
 Labiatae 137
 Lactams 515,1135,1813
 Lactone 497
 Lamiaceae 171,1015,1023,1027,1205, 1229, 1343,1347,1351,1373, 1487,1511,1689,1697, 1775,1949
Lantana camara 1031
Lantana canescens 247
Lantana lopez-palcii 247
 Lapachone 1677
 Lariciresinol 379
 Larvae 887
 Larvicidal activity 701,907,1361
Lasiodiplodia theobromae 1801
 Lauraceae 267,901,1297,1357,1591,1755
Laurencia papillosa 1821
Laurus nobilis 1483
 Lavender 1199
 Lavender essential oils 1769
 LC/ESI-MS/MS 1593
 LC-ESI-MS 17
 LC-MS/MS 7,21,1305
 Lead 1473
 Leaf anatomy 1045
 Leaves 97,883,1893
 Lectins 1403
 Lecythomycin 677
Lecythophora sp 677
 Leguminosae 1437,1749
Leishmania braziliensis 231

- Leishmania panamensis* 231
Leishmania 689
Leishmania amazonensis 281
Leishmania amazonensis 973
Leptosphaeria biglobosa 617
Leptosphaeria maculans 617
Lerchea cf. *bracteata* 349
Lerchea parviflora 349
Lethaloxin 505
Lichexanthone 1807
Lignans 379,1361
Limonene 901,1945
Linalool 111,875,1333,1347,1705,
Linolenic acid 1005
Linum usitatissimum 1701
Lipid peroxidation 785,1285
Lipo-alkaloid 527
Lippia multiflora 1183
Liquid chromatography 1817
Liriodenine 1251
Litsea acutivena 1755
Litsea akoensis 901
Liver fibrosis 1267
Liverwort constituents 393
Liverworts 49,303
Loranthaceae 1051
Lovastatin 183
Lövéte 1459
Lozenge 1353
Lupeol 1807
Lycoris albiflora 187
Lythrum salicaria L. 1417

Machilus kusanoi 267
Macluraxanthone 1269
Macrocycles 393
Macrolactone glycosides 677
Macrolactones 223
Macrolactonization 505
Macrolides 505
Macropropagation 985
Maculansin A 617
Madhuca latifolia 1661
Madhuglucoside 1661
Magnolia dealbata Zucc 1121
Magnoliaceae 1121
Magnolol 1121
Maillard reaction products 55
Majdine 1831
Malic acid esters 1293
Mallotus 813
Maltodextrin/juice powder 951
Mammalian cell toxicity 1743
Mandarin 1495
Manekia (Sarcorhachis) naranjoana 145
Manganese deposition 679
Mangostin 1267
Mannich reaction 31
Marchantins 393
Marigold 1873
Marine natural products 287
Marine sponges 1237
Marine-derived fungus 1063
Marrubium 1697
Marschalk alkylation 841
Martinella obovata 957
Massage aromatherapy 1199
Mate 651
Matricaria pubescens 251
Matricaria suaveolens 1107
MCF-7 807
Measles virus 1881
Medicinal plants 79,1001,1459,1461,1469
Mediterranean 1483
Melanin 1397
Meliaceae 785,1259
Melicoester 343
Melicope ptelefolia 343
Melicopeprenoate 343
Melidianolic acid A 323
Melidianolic acid B 323
Mentha aquatica L. 1479
Mentha spicata 1373
Menthane lactone 1479
Menthofuran 1479
Menthone 695
Mesenteric bed 871
Metabolism 1913
Metabolomic analysis 1621
Metal complex 1137
Methanol 1939
Methanolic extract 989
Methoxyaristololactam I 11
Methoxydaphnodorin D₁ 1111
Methoxydaphnodorin D₂ 1111
Methyl 4,4'-dimethoxyvulpinate 357
Methyl 6-bromoindolyl-3-acetate 451
Methyl chavicol 1705
Methyl cinnamate 1705
Methyl eugenol 1705
Methylascorbigen 611
Methylated carrageenan 1327
Methylchavicol 1027
Methyleugenol 1189
Mexican medicinal plants 1925
MIC 1159
Michelia champaca 1251
Microbial transformation 781,1129
Microcarpalide 505
Microclimatic factors 243
Microelements 1469
Micromorphology 259
Micronucleus 989
Micropropagation 985
Middle Atlas mountainous area 1455
Mini-barcode sequence 67
Minimolide 433
Molecular docking 1411,1421,1637
Molecular genetic study 423
Molecular identification 67
Molecular modeling 1225
Molluscicidal activity 387
Monoterpene citral 1221
Monoterpenes 879,1151,1799
Monoterpenes esters 49
Monoterpenoids 93
Moraceae 375,1095
Morinda citrifolia seed oil 1005
Morinda lucida 1537
Moringa peregrina 1451
Moroccan ethnopharmacology 1441
Morphofunctional groups 667
Mountain tea 1305
MPO 1545
MRSA 1379
MS 35,175,1059
MTT 803
Mulberry leaves 1621
Multidrug-resistant 1159,1743,1925
Multivariate analysis 1711
Mutagenicity 1269
Murolene 1949
Mycobacterium 689
Mycosphaerella sp. nov 835
Mycotoxins 657,1915
Myrcia 977
Myristicin 883,1151,1361
Myrothecium verrucaria Hmp-F73 1915
Myrtaceae 969,977,1045
Myrtanal 429
Myrtus communis 1545

N(b)-methylantirrhine 349
Nagybacon 1459
Naphthalenones 661
Naphthazarin 217
NaphthoquinoneS 217
Naringenin glycoside 371
Naringin 951
Native forest species 985
Natural compounds 1205
Natural insecticide 1389
Natural limonoids 785
Natural products 393,983,1133,1135,1605
Natural products drug discovery 1549
NCTC 2544 1315
Nectandra angustifolia 1393
Neospergilliacid 1137
Neolitsea aciculata 1193
Neolitsea parvigemma 1357
Neomenthol 695
Neosartorya glabra 807
Neotropics 1549
Nepeta × *faassenii* Bergmans ex Stearn 1015
Nepeta 1697
Nepeta cataria 1015
Nepeta mussinii 1015
Nepeta nepetella L. 1015
Nepeta nuda 1015
Nepetalactone 1015
Neral 93
Nerine bowdenii 1827
Nerine undulata 1255
Neroli oil 1009
Nerylacetone 367
Neurotrophic activity 327
NF-κB inhibitors 627
Nicotinamide 1835
Nicotinamide adenine dinucleotide 1835
Nicotinic acid 1835,1877
Nitric oxide 1095
N-mercapto-4-formylcarbostyryl 921
NMR 35,175,793,9571003,1059,1225,1479,
1621,1645, 1801,1889
NMR quantitative analysis 379
NMR spectroscopy 209,673
Nonane 271
Nonenolide 1843
Northern Syria 79
Nosocomial infection 1505
Nostoc commune 689
Nothapodytes nimmoniana 197
N-thiols 921
Nucleic acids 773
Nyctaginaceae 1651

Obesity 1839
Ocimene 875
Ocimenone 1367
Ocimum 1027
Ocimum basilicum 1189,1487
Ocimum species 1705
Ocotea 231
Octadecenoic acid 1023
Octadecyl *p*-coumarate 1889
Ostenol 247,1723
Odor-active components 403
Odour Products 439
Off-line processes analysis 719
Oil composition 1393,1559
Oil Red O staining 1839
Olacaceae 327
Oleanolic acid 1573,1813
Olefin metathesis 505
Oleuropein 753
Omani basil 1487
Onagraceae 35
On-line processes analysis 719
Onopordopicrin 163
Ontogeny 239
Oomycetes 45
OPG 1689
ORAC 213
Ordinary least squares 1939
Origanum tyttanthum 1719
Origanum vulgare ssp. *viride* 1351
Osteoblasts 1689

- Ottonia anisum* 939
 Overpressured layer chromatography 657
 Oxidation potential 1673
 Oxine 597
 Oxocyclonerolidol 1763
 Oxofuroindolines 457
 Oxygen radical absorbance capacity 159
 Oxygenated components 403
 Oxyresveratrol 41
 Ozone 657
Ozoroa pulcherrima 1133

 Paeonol 1129
 Palisade cells 1475
Panax 581
Papaver coreanum 1593
 Papaveraceae 1593
Paris polyphylla var. *yunnanensis* 1759
Paronychia 205
 Parsley 961
 Partial purification 73
Passiflora quadrangularis 1091
 PCA 21,1621
 Peel 1857
Pelargonium 15
 Pendulone 1645
 Peonidin glucopyranosides 35
 Peptide fractions 1683
 Peptidyl-tRNA hydrolase 1421
Perilla frutescens 1461
Perilla frutescens var. *crispa* 423
 Perillyl alcohol 1805
 peripheral blood mononuclear cells 1925
Petroselinum crispum 961
 pH 1425
 Pharmacokinetics 337
 Pharmacology 287
 Phellandrene 1023,1051, 1531,1945
 Phenolic apiosides 957
 Phenolic compounds 17,1863
 Phenolic contents 845,1441
 Phenolic extraction 1657
 Phenolic glycosides 205,1901
 Phenolics 819,1305
 Phenylalanine 857
 Phenylalkanoid 855
 Phenylethanoid glycosides 21,1305
 Phenylethyl cinnamides 851
 Phenylglycinol 515
 Phenylpentadiyne 119
 Phenylpropanoid glycoside 1599
 Phenylpropenoic acids 1293
 Pheromone 409
 Phillyrin 753
Phleum alpinum 101
 Phlomeic acid 171
Phlomis bracteosa 171
Phoma lingam 617
 Phomalide 617
Phomopsis sp. 1905,1907
 Phomosines 1907
 Phosphodiesterase 73
 Phosphomonoesterase 73
 Photo reaction 41
 Photosensitization mechanisms 1677
Phthirusa adunca 1051
 Physico-chemical factors 337
 Phytoalexins 537
 Phytochemical polymorphism 1417
 Phytochemical screening 1421
 Phytochemistry 1549
 Phytoecdysteroids 1085
 Phytopathogenic fungi 921
 Phytoquinoids 1893
 Phytotoxic activity 1735
 Phytotoxins 593,617
 Phytyl fatty acid esters 767
 Picolinic acid 1877
 Piedmont 1461

 PIFA 537
 Pinaceae 793
 Pineapple weed 1107
 Pinene 119,141,271,1515,1531,1945
 Pinguiculate 303,315
 Pinorensin 379
Pinus patula 1531
 Piperaceae 145,939
 Piperitenone oxide 1373
 Piperidine 515
Pistacia lentiscus 1505,1543
Pituranthos scoparius 1151
 Plant pathogens 701
 Plant virus 1189
Plasmodium 689
Plasmodium falciparum 3D7 323
 Platelet aggregation 961
Plectranthus punctatus 1229
Plectranthus tenuicaulis 409
Pleurothyrium 231
Pleurotus sapidus 1217
 Pluchagen A 3
Pluchea arguta 1
Plumeria bicolor 1567
 Plumericin 1567
 PMOR 1315
Poa alpina 101
 Poliovirus 1881
 Pollinator 1461
 Polygodial 315
 Polygonaceae 55,1597
Polygonum 931
Polygonum amplexicaule 1597
Polygonum multiflorum 55
 Polyisoprenylated benzophenones 1269
 Polyketide 617
 Polyphenols 651,1107,1299
 Polyphenols complexes 1673
 Polyphenols content 951
 Polysaccharides 867
 Pomiferin-4'-O-methyl ether 1103
Pongamia pinnata 625
Porella 315
Porella perrottetiana 303
Potentilla evestita 1083
 PPAR activity 1233
 Preadipocytes 1839
 Precipitation 1465
 Pre-emergence 593
 Pregnane derivatives 1083
 Prenylated chalcone 1103
 Prenylated isoflavone 1103
 Prenylated polyphenols 1397
 Prenylated xanthones 1269
 Prenylflavanones 1099
 Proanthocyanidins 829,913
 Production cost 1461
 Profitability 1461
 Propolis 973
 Prospective application 707
 Protein hydrolysates 1683
 Protein kinase A 1921
 Protein ubiquitination 1411
Protium aracouchine 1727
Protium giganteum 1727
 Provenances 123
Pseudomonas fluorescens 921
 Pteridophyta 1723
 Pterocarpan 537
 Ptycholid 327
Ptychopetalum olacoides 327
 Public health 1001
 Pulegone 695,1027
Pulmonaria officinalis 1475
 Pulp 1857
Pulsatilla chinensis 799
 Pyranoflavone 31
 Pyrrolidin-2-ones 1243
 Pyrrolidinoindolines 457

 Pyruvated α carrageenan 1327

 qNMR 1281
 QR inducing activity 799
 Quadruple immune response 631
 Quality 733
 Quality control 17,753,1929
 Quassinoids 337
 Quercetin 201
 Quinic acids 1665

 Radical quenching activity 625
 Radical scavenger 969,977,1441
 Radical scavenging 1285
 Radioprotection 587
 Ranaconitine 1589
 RANKL 1689
 Rapid Resolution Liquid Chromatography 581, 645
 Reaction-controlled phase-transfer catalyst 1141
 Reactive oxygen species 1677
 Red alga 1821
 Red blood cells 607
 Red flour beetle 887
 Reducing power 1285,1447
 Regioselective bromination 451
 Relaxing effect 1199
 Repellence 887
 Repellency 1945
 Resibufogenin 1581
 Resin 379
 Respiratory diseases 1459
 Response factors 417
 Response surface methodology 1933
 Restinga 1045
 Resveratrol 631,1673
 Reverse prenylated indole alkaloids 807
Rhamnus cathartica 1275
Rhamnus orbiculatus 1275
Rhazya stricta 1451
Rheedia acuminata 1269
 Rhodesain inhibition 1637
Rhodiola rosea L. 645
Rhynchosia edulis 1637
 Rhynedulin 1637
 Rhynedulin 1637
 Riboflavin 41
Rivularia bischoffiana 689
 ROS 597
 Rosaceae 1083,1851
 Rosavin 645
 Rose oxide 403
Rosmarinus tournefortii 1511
 Rotenoids 1651
 Royleanone 575
 RP-HPLC 1863
 RRLC 581,639,645
 RSV 1881
 Rubiaceae 237,747,1573
 Rutaceae 343,875
 Rutin 201

 Sabinene 875
 Sabinyl acetate 275
 Safflower 1873
 Saffron 1873
Sage oil 417
 Sahara 1445
 Sakurai-Hosomi reaction 537
 Salidroside 645
 Salinity 1559
 Salt stress 1835
Salvia 575,1697
Salvia mukerjeei 1949
Salvia officinalis 417
Salvia species 1205
Salvia verbenaca 1023
 Sapotaceae 1661
 SAR 303

- Sarconema filiforme* 1327
 Sarcophine 1809
Sarcophyton glaucum 1809
 Sargaol 1073
 Sartoryglabris 807
Satureja 1167
Satureja hortensis 1173
Satureja kitaibelii 1353
Satureja thymbra L 1523
Scadoxus multiflorus 1255
Scaphechinus mirabilis 217
 Sch642305 505
Schistocarpa eupatorioides 767
Scleroderma sinnamariense 357
Scutellaria orientalis ssp. *alpina* 1347
Scutellaria utriculata 1347
 Sea cucumbers 1075,1913
 Sea urchin 773
 Sea urchin pigments 217
 Seagrass 667
 Seaweeds 667
 Secretory cavity 1045
 Selenium 1469
 Semi-synthesis 433
 Semi-synthetic analogues 785
 Semliki Forest virus 1573
Senecio belgaumensis (Wight) Cl 1145
Senecio coincy 123
 Senghor Seamount 679
 Sequential biotransformation 223
 Sesamin 1807
Seseli globiferum 1163
Seseli montanum subsp. *tommasinii* 263
 Sesquichamaenol 1807
 Sesquillignan glycoside 1901
 Sesquiterpene lactones 163,491,1055
 Sesquiterpenoid 1,311,477,621,763,879
 Sesquiterpenoid coumarin 209
 Sesquiterpenolides 491
 Sex pheromone 1323
 Shamiminol 1897
 Shiitake mushroom 845
Sibiraea angustata 1799
 Sibiralactone 1799
Sideritis 21,1305,1689
 Siderophore 1137
 Sigmone 1099
 Sigmotriol 1099
 Sildenafil 1003
 Silver nitrate chromatography 935
 Silybin A1653
 Silybin B 1653
 SIMCA 1621
 Simultaneous distillation extraction (SDE) 1155
 Sirodesmins 617
 Skin pathogen 1193
 Skin whitening 1397
 Slurry state 1933
 Snails 1403
 Snake venom 1393
 Solanaceae 227,1825,
Solanum melongena 851
 Somatic embryogenesis 985
 Soybean oil 1141
 Soybean pod borer 1323
 Spathulenol 97,251
Spatoglossum schroederi 863
 Spice 1873
 SPME 1333
 SPME-GC/MS 1495
Spodoptera frugiperda 1389
Spongia 773
 Spongian diterpenoids 773
Sprekelia formosissima 1255
 Sprout inhibition activity 1373
 SSF 183
 Stability 1425
 Stability testing 639
 Stabilizers 1425
Stachys spruneri 277
 Standardized extract Fr 2 337
 Steaming process 55
 Steroidal glycosides 1239
 Steroidal saponins 1825
 Sterol 1237
Stevia rebaudiana 175,1059
Stevia tomentosa 1225
Streptococcus mutans 777
Streptococcus pneumoniae 1035
 Stress metabolite 53
 Strobilurins 1917
Strongylocentrotus intermedius 773
Stypopodium flabelliforme 1073
 Submerged cultures 183
 Sugars 673
 Sulfated polysaccharide 863
 Summer savory 1173
 Sun 1475
 Supercritical carbon dioxide extraction 107
 Supercritical CO₂ 1523,1527
 Supercritical fluid extraction (SFE) 1155
 Superoxide 41
 Surface flavonoids 15
 SW480 1671
 Sweet basil 1189
 Sweet bell peppers 1817
Symphytum officinale 1475
 Synergism 777
 Synthesis 167,439,443,491,1323
Syzygium johnsonii 1421
Tabernaemontana catharinensis 193
 Tablets 951
 Tajikistan 1719
 Tamarind seed husk 829
Tanacetum 259
 Tannase 1933
 Tannin 829
 Tassili n'Ajjer 87
 TDDFT ECD calculation 1799
 Teas 651
 TEM 1159
 Temperature 1425,1465
Tephrosia purpurea 1437
Terminalia macroptera 1125
Terminalia sericea 1585
 Terpene alcohols 1159
 Terpenes 1499,1577
 Terpenoids 87,315,1205
 Terpinen-4-ol 875
 Terpinene 1173,1537,1715
 Terpeneol 107
 Terpinolene 275
 Terrestrial *Streptomyces* 45
 Tetrahydroxychalcone 1117
Teucrium arduini 1385
 Thailand 627
Thelechitonia trilobata 1945
 Theobroxide 1801
 Thin layer chromatography 1379
 Thiophene 361
 Thiosildenafil 1003
 Thymidine kinases 773
 Thymol 137,901,1719
 Thymoquinone 213
Thymus caucasicus 137
Thymus kotschyanus 137
Thymus pallidus 1507
Thymus saturoioides 1507
Thymus vulgaris 137
 Time-kill curve 777
 Tissue engineering 555
 TMV 1385
 TNF- α 1543
 TNF- α 1545
Tornabenea bischoffii 1179
 Total flavonoid 375
 Total flavonoid composition 1417
 Total flavonoid content 1617
 Total phenol 375
 Total phenol content 1855
 Total phenolic content 1617
 Total polyphenol composition 1417
 Total synthesis 333,393,515
 Totarol 133
 Toxic plants 999
 Traditional Arabic Medicine (TAM) 79
 Traditional medicine 627
 Traditional processing methods of drugs 527
 Transesterification 527
Trichomonas vaginalis 973
 Trichothecenes 1915
 Tridecanone 111
 Tridecene 123
Trifolium pallidum 1293
Trifolium pratense 1657
 Trigonelline 1835
 Trihydroxy-methyl-norpregna-trienone 1083
 Trihydroxypregna-diene-dione 1083
 Triplet excited state 1677
 Triquinane sesquiterpenes 767
 Triterpene glycosides 587,1075
 Triterpene saponins 799
 Triterpene synthase 1665
 Triterpenes 575,1813
 Triterpenes 789,793
 Triterpenoid saponins 1679
 Triterpenoids 179
Triticum aestivum L 1445
 Trixagol 439
 Tropical plant extracts 1421
Trypanosoma 689
 TTN 537
 Tuberculosis 1743
 Turbidimetry 1445
 Turmeric 1873
 Two-spotted spider mite 893
 Tyloxapol 1005
 Tyrosinase 1397
Ultra HRGC 417
 Ultra performance liquid chromatography 1617
 Ultrafiltration 1683
 Ultra-high dilutions 1499
 Umbelliferae 209,491
 Undecanol 1145
 Undecanone 111
 Urease 1117
 Ursolic acid 171,1573
 Uscharin 803
Uvaria anonoides 1715
Uvaria ovata 1715
Uvaria tortilis 1715
Vaccaria hispanica 1599
 Vaccarin H 1599
Valeriana hardwickii 129
Valeriana jatamansi 129
 Validation 931,1831
 Valine 857
 Van Herby Cheese 1473
 Vapor toxicity 1183
 Vegetal extracts 1863
 VEGF and PIGF 943
 VEGFR1/Flt-1 943
 Velvet antler 1683
Vepris glomerata 1847
 Veprisinol 1847
Verbascum thapsus L. 989
 Verbenaceae 3,247,1031
 Vernolepin 433
 Vibrational circular dichroism 759
Viburnum plicatum 1901
 Vietnam 813
Vinca herbacea 1831
 Virescenosides 1063
 Virucidal 993

Vismia baccifera 699
Vismia macrophylla 85
 Vitexin, E2-25K 1411
Vitis vinifera 59
Vitis vinifera tendrils 1315
 Volatile components 1023
 Volatile compounds 1045
 Volatile oils 107,1537,1759
 Volatile organic compounds 1723

Wendlandia tinctoria 747
Wikstroemia indica 1111
 Wind 1475

Wine 1939
 Wine lactone 403
Winteraceae 879
Withania frutescens 1447
 Withanolides 1447
 Wound and skin infections 1167

Xanthine oxidase 1125,1437
 Xanthone 1267
 Xanthorrhizol 167
Xenorhabdus 1247
 X-ray diffraction analysis 1589
Xylaria sp. 1843

Yield 1559

Zanthoxylum armatum 111
Zanthoxylum rhetsa 1595
Zanthoxylum setulosum 1807
Zingiber officinale 93,855,1671
Zingiberaceae 93,855
Ziziphora clinopodioides 695
 Zoosporicide 45
 Zygophyllaceae 179
Zygophyllum eurypterum 179

Natural Product Communications

Manuscripts in Press Volume 6, Number 12 (2011)

***Spartium junceum* Aromatic Water: Chemical Composition and Antitumor activity**

Teresa Cerchiara, Serafina V. Straface, Giuseppe Chidichimo, Emilia L. Belsito, Angelo Liguori, Barbara Luppi, Federica Bigucci and Vittorio Zecchi

Effect of *Hibiscus sabdariffa* and its Anthocyanins on some Reproductive Aspects in Rats

Badreldin H. Ali, Intisar Al-Lawati, Sumyia Beegam, Amal Ziada, Suhail Al salam, Abderrahim Nemmar and Gerald Blunden

Luteolin Induces Mitochondria-dependent Apoptosis in Human Lung Adenocarcinoma Cell

Qing Chen, Shengming Liu, Jinghong Chen, Qianqian Zhang, Shijie Lin, Zhiming Chen and Jianwei Jiang

Chemical Composition, Antimicrobial, Antiradical and Anticholinesterase activity of the Essential Oil of *Pulicaria stephanocarpa* from Soqatra

Nasser A. Awadh Ali, Rebecca A. Crouch, Mohamed A. Al-Fatimi, Norbert Arnold, Axel Teichert, William N. Setzer and Ludger Wessjohann

A New Megastigmane Glycoside, Phoenixoside A, from *Phoenix dactylifera*

Sumbul Azmat, Aqib Zahoor, Rehana Ifzal, Viqar Uddin Ahmad and Faryal Vali Mohammed

Identification and Quantification of the Antimicrobial Components of a Citrus Essential Oil Vapor

Carol A. Phillips, Konstantinos Gkatzionis, Katie Laird, Jodie Score, Avinash Kant and Mark D. Fielder

Ternary Liquid-Liquid Equilibria Measurement for Epoxidized Soybean Oil + Acetic Acid + Water

Shuang-Fei Cai, Li-Sheng Wang, Guo-Qing Yan, Yi Li, Yun-Xia Feng and Rong-Gang Linghu

Composition, Antioxidant and Antimicrobial Activities of the Leaf Essential Oil of *Machilus japonica* from Taiwan

Chen-Lung Ho and Yu-Chang Su

Composition of the Essential Oil of *Pogostemon travancoricus* var. *travancoricus*

Ramar Murugan and Gopal Rao Mallavarapu

New 3,4-*Seco-ent*-kaurene Dimers from *Croton micans*

Elsa Mateu, Katuska Chavez, Ricarda Riina, Reinaldo S. Compagnone, Franco Delle Monache and Alirica I. Suárez

Synthesis of Novel Anti-acetylcholinesterase Inhibitors from Oleanolic Acid

Ahlem Beyaoui, Fayçal Hichri, Amel Omri and Hichem Ben Jannet

Microbial Conversion of Tomato by a Plant Pathogenic Bacterium *Pectobacterium atrosepticum*: a Plant-Microbial Approach to Control Pathogenic *Candida* Species

Vivek K. Bajpai, Sun Chul Kang, Soon-Gu Lee and Kwang-Hyun Baek

Xanthones with Antiproliferative Effects on Prostate Cancer Cells from the Stem Bark of *Garcinia xanthochymus*

Feng Ji, Zhanlin Li, Gaofeng Liu, Shengli Niu, Nan Zhao, Xiaoqiu Liu and Huiming Hua

Chemical Constituents of the Essential Oil from Aerial Parts and Fruit of *Anisosciadium orientale*

Vahid Rowshan, Ahmad Hatami, Atefeh Bahmanzadegan and Mahnaz Yazdani

Physical Properties and Molecular Conformations of Indole Alkaloids and Model Protein Interactions – Theoretical vs. Experimental Study

Bojidarka Ivanova and Michael Spitteller

Constituents of Kenyan *Gardenia volkensii*

Esther W. Kinuthia, Moses K. Langat, Elizabeth M. Mwangi and Peter K. Cheplogoi

Analysis of Danshen and Twelve Related *Salvia* Species

Luyang Lu, Yuan Liu, Zhifeng Zhang and Hao Zhang

Flowers from *Kalanchoe pinnata* are a Rich Source of T Cell-Suppressive Flavonoids

Marcela A. S. Coutinho, Michelle F. Muzitano, Elaine A. Cruz, Maria C. Bergonzi, Carlos R. Kaiser, Luzineide W. Tinoco, Anna R. Bilia, Franco F. Vincieri, Bartira Rossi-Bergmann and Sônia S. Costa

Fast Quality Assessment of German Chamomile (*Matricaria chamomilla* L.) by Headspace Solid-Phase Microextraction: Influence of Flower Development Stage

Mohammad Rafieiohossaini, An Adams, Hamid Sodaieizadeh, Patrick Van Damme and Norbert De Kimpe

Composition of the Essential Oils and Antibacterial Activities of *Hymenocrater yazdianus*, *Stachys obtusirena* and *Nepeta asterotricha* Three Labiatae Herbs Growing Wild in Iran

Shiva Masoudi, Abdolhossein Rustaiyan, Raziheh Mohebat and Mohammad Hossein Mosslemian

GC-MS Analysis of *Ziziphora clinopodioides* Essential Oil from North Xinjiang, China

Xiaoying Zhou, Qian Yu, Haiyan Gong and Shuge Tian

Analysis of the Essential Oil of *Teucrium polium* ssp. *capitatum* from the Balkan Peninsula

Violeta Mitić, Olga Jovanović, Vesna Stankov-Jovanović, Bojan Zlatkovic and Gordana Stojanovic

Application of Mixture Analysis to Crude Materials from Natural Resources (III)^[1]: NMR Spectral Studies to Analyze Chalcones from *Angelica keiskei*

Eriko Fukuda, Masaki Baba, Yoshihiro Uesawa, Osamu Kamo, Kazunori Arifuku, Koji Tsubono, and Yoshihito Okada

Diacarperoxide S, New Norterpene Cyclic Peroxide from the Sponge *Diacarnus megaspinorhabdosa*

Sabrin R. M. Ibrahim

Complete NMR Assignments of Tubulosine

Venkata Siva Satyanarayana Kantamreddi and Colin W. Wright

Ferric Reducing, Antiradical and β-Carotene Bleaching Activities of Nicotinic Acid and Picolinic Acid Bioconjugates of Curcumin	
Archana Pandey, Kanti Bhooshan Pandey, Ravindra Kumar Gupta and Syed Ibrahim Rizvi	1877
Antiviral Activities of Diarylheptanoids Isolated from <i>Alpinia officinarum</i> against Respiratory Syncytial Virus, Poliovirus, Measles Virus, and Herpes Simplex Virus Type 1 <i>in vitro</i>	
Katsuhiko Konno, Rie Sawamura, Yi Sun, Ken Yasukawa, Tomomi Shimizu, Wataru Watanabe, Masahiko Kato, Ryuichi Yamamoto and Masahiko Kurokawa	1881
Isolation of C-glycosyl Xanthenes from <i>Coffea pseudozanguebariae</i> and Their Location	
Pascale Talamond, Geneviève Conejero, Jean-Luc Verdeil and Jean-Luc Poëssel	1885
Antifungal Activity and Isomerization of Octadecyl <i>p</i>-coumarates from <i>Ipomoea carnea</i> subsp. <i>fistulosa</i>	
Eugene Sebastian J. Nidiry, Girija Ganeshan and Ankanahalli N. Lokesh	1889
New Glucose Esters from the Fresh Leaves of <i>Jacaranda mimosaeifolia</i>	
Christianah A. Elusiyan and Tiwalade A. Olugbade	1893
Shamiminol: A New Aromatic Glycoside from the Stem Bark of <i>Bombax ceiba</i>	
Shaheen Faizi, Sadia Zikr-Ur-Rehman and Muhammad Ali Versiani	1897
Two New Phenolic Glycosides from <i>Viburnum plicatum</i> var. <i>plicatum</i> f. <i>plicatum</i>	
Saki Katagiri, Yoshiki Watanabe, Yasunori Yaoita, Masao Kikuchi and Koichi Machida	1901
Antimicrobial Chemical Constituents from the Endophytic Fungus <i>Phomopsis</i> sp. from <i>Notobasis syriaca</i>	
Hidayat Hussain, Michel Kenne Tchimine, Ishtiaq Ahmed, Kathrin Meier, Michael Steinert, Siegfried Draeger, Barbara Schulz and Karsten Krohn	1905
Phomosines H–J, Novel Highly Substituted Biaryl Ethers, Isolated from the Endophytic Fungus <i>Phomopsis</i> sp. from <i>Ligustrum vulgare</i>	
Karsten Krohn, Umar Farooq, Hidayat Hussain, Ishtiaq Ahmed, Joachim Rheinheimer, Siegfried Draeger, Barbara Schulz and Teunis van Ree	1907
Isolation and Characterization of a new Benzofuran from the Fungus <i>Alternaria</i> sp. (HS-3) Associated with a Sea Cucumber	
Xuekui Xia, Jun Qi, Fang Wei, Airong Jia, Wenpeng Yuan, Xiumei Meng, Miansong Zhang, Changheng Liu and Changyun Wang	1913
Potent Toxic Macrocyclic Trichothecenes from the Marine-Derived Fungus <i>Myrothecium verrucaria</i> Hmp-F73	
Li Zhao, Li Liu, Nan Wang, Shu-Jin Wang, Jing-Chun Hu and Jin-Ming Gao	1915
Synthesis and Bioactivity of Novel Coumarin Derivatives	
Ai-Ying Guan, Chang-Ling Liu, Miao Li, Zhi-Nian Li, Ming-Xing Zhang and Hong Zhang	1917
Kinase Inhibitory, Haemolytic and Cytotoxic Activity of Three Deep-water Sponges from North Western Australia and their Fatty Acid Composition	
Ana Zivanovic, Natalie J. Pastro, Jane Fromont, Murray Thomson and Danielle Skropeta	1921
Antimicrobial and Cytotoxic Effects of Mexican Medicinal Plants	
Maria del Rosario Jacobo-Salcedo, Angel Josabad Alonso-Castro, Luis A. Salazar-Olivo, Candy Carranza-Alvarez, Luis Ángel González-Espíndola, Fabiola Domínguez, Sandra Patricia Maciel-Torres, Concepción García-Lujan, Marisela del Rocio González-Martínez, Maricela Gómez-Sánchez, Eduardo Estrada-Castillón, Rocio Zapata-Bustos, Pedro Medellín-Milán and Alejandro García-Carrancá	1925
Chemometrics Evaluation of the Herbal Drug <i>Andrographis paniculata</i>	
Shiv Narayan Sharma, Zenu Jha and D. K. Sharma	1929
<i>Garcinia cambogia</i> Leaf and Seawater for Tannase Production by Marine <i>Aspergillus awamori</i> BTMFW032 under Slurry State Fermentation	
Beena P. S, Soorej M. Basheer, Sarita G. Bhat and Chandrasekaran M	1933
Gas Chromatographic Quantitative Analysis of Methanol in Wine: Operative Conditions, Optimization and Calibration Model Choice	
Rosario Caruso, Grazia Laura Gambino, Monica Scordino, Leonardo Sabatino, Pasqualino Traulo and Giacomo Gagliano	1939
Composition and Biological Potential of Essential Oil from <i>Thelechitonina trilobata</i> Growing in South Africa	
Jamie Peebles, Ephraim Gwebu, Opeoluwa Oyediji, Sarah Nanyonga, Nokuthula Kunene, David Jackson, William Setzer and Adebola Oyediji	1945
Chemical Composition and Antibacterial Activity of Essential oil from <i>Salvia mukerjeei</i>	
Lalit Mohan, Anuradha Negi, Anand B. Melkani and Vasu Dev	1949
<u>Review/Account</u>	
Revealing Indigenous Indonesian Traditional Medicine: Anti-infective Agents	
Ari S. Nugraha and Paul A. Keller	1953

Natural Product Communications

2011

Volume 6, Number 12

Contents

<u>Original Paper</u>	<u>Page</u>
Sibiralactone: A New Monoterpene from <i>Sibiraea angustata</i> Guangbo Xie, Xianlong Wang, Tibor Kurtán, Attila Mándi and Tianzhi Wang	1799
Bioconversion of Proposed Precursors into Theobroxide and Related Compounds Peng Li, Kosaku Takahashi, Ahmed Elkhateeb, Hideyuki Matsuura, Teruhiko Yoshihara and Kensuke Nabeta	1801
Microbial Hydroxylation of <i>S</i>-(-)-Perillyl Alcohol by <i>Fusarium heterosporium</i> Ismail Kiran	1805
A Phytochemical Investigation of <i>Zanthoxylum setulosum</i> Tameka M. Walker, Bernhard Vogler, Debra M. Moriarity, William A. Haber and William N. Setzer	1807
Cytotoxic Cembranoids from the Red Sea Soft Coral <i>Sarcophyton glaucum</i> Mohamed-Elamir F. Hegazy, Ahmed A. El-Beih, Alaa Y. Moustafa, Abdelhamed A. Hamdy, Montaser A. Alhammad, Rehab M. Selim, Mohamed Abdel-Rehim and Paul W. Paré	1809
C-Lactam Derivatives of Oleonic Acid. The synthesis of C-lactam by Beckmann rearrangement of C-oxime Barbara Bednarczyk – Cwynar	1813
Analysis of Native Carotenoid Composition of Sweet Bell Peppers by Serially Coupled C₃₀ Columns Daniele Giuffrida, Paola Dugo, Giacomo Dugo, Germana Torre and Luigi Mondello	1817
New Antifungal Cholestane and Aldehyde Derivatives from the Red Alga <i>Laurencia papillosa</i> Walied M. Alarif, Sultan S. Al-Lihaibi, Ahmed Abdel-Lateff and Seif-Eldin N. Ayyad	1821
Steroidal Saponins from the Fruits of <i>Cestrum ruizteraniamum</i> Elier Galarraga M., Anne-Claire Mitaine-Offer, Juan Manuel Amaro-Luis, Tomofumi Miyamoto, Chiaki Tanaka, Laurent Pouységu, Stéphane Quideau, Luis B. Rojas and Marie-Aleth Lacaille-Dubois	1825
Isolation and Cholinesterase Activity of Amaryllidaceae Alkaloids from <i>Nerine bowdenii</i> Lucie Cahliková, Stanislav Zavadil, Kateřina Macáková, Irena Valterová, Andrea Kulhánková, Anna Hošťálková, Jiří Kuneš and Lubomír Opletal	1827
HPLC Determination of Majdine in <i>Vinca herbacea</i> Natia Gagua, Beatrice Baghdikian, Fathi Mabrouki, Riad Elias, Valentina Vachnadze, Aliosha Bakuridze and Evelyne Ollivier	1831
Pyridine Metabolism and Trigonelline Synthesis in Leaves of the Mangrove Legume trees <i>Derris indica</i> (<i>Millettia pinnata</i>) and <i>Caesalpinia crista</i> Yuling Yin, Hamako Sasamoto and Hiroshi Ashihara	1835
Anti-adipogenic Activity of <i>Cordyceps militaris</i> in 3T3-L1 Cells Qing Liu, In Pyo Hong, Mi-Jeong Ahn, Hwan-Soo Yoo, Sang-Bae Han, Bang Yeon Hwang and Mi Kyeong Lee	1839
Two New Cyclopeptides and One New Nonenolide from <i>Xylaria</i> sp. 101 Yao-Yao Li, Zhi-Yu Hu, and Yue-Mao Shen	1843
A Novel Flavonoid and Furoquinoline Alkaloids from <i>Vepris glomerata</i> and their Antioxidant Activity Joyce J. Kiplimo, Md. Shahidul Islam and Neil A. Koorbanally	1847
Flavonoid Constituents and Free Radical Scavenging Activity of <i>Alchemilla mollis</i> Antoaneta Trendafilova, Milka Todorova, Milena Nikolova, Anna Gavrilova and Antonina Vitkova	1851
Ultrasound-assisted Extraction of Total Phenols and Flavonoids from Dry Tobacco (<i>Nicotiana tabacum</i>) Leaves Ivana T. Karabegović, Vlada B. Veljković and Miodrag L. Lazić	1855
Characterization of Polyphenolic Compounds in Unripe Chinotto (<i>Citrus myrtifolia</i>) Fruit by HPLC/PDA/ESI/MS-MS Monica Scordino, Leonardo Sabatino, Adalgisa Belligno and Giacomo Gagliano	1857
Bioactive Compounds, RP-HPLC Analysis of Phenolics, and Antioxidant Activity of Some Portuguese Shrub Species Extracts Ángelo Luís, Fernanda Domingues and Ana Paula Duarte	1863
HPLC/PDA/ESI-MS Evaluation of Saffron (<i>Crocus sativus</i> L.) Adulteration Leonardo Sabatino, Monica Scordino, Maria Gargano, Adalgisa Belligno, Pasqualino Traulo and Giacomo Gagliano	1873

Continued inside backcover