


NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering all
Aspects of Natural Products Research



Volume 8. Issue 12. Pages 1665-1830. 2013
ISSN 1934-578X (printed); ISSN 1555-9475 (online)
www.naturalproduct.us

EDITOR-IN-CHIEF

DR. PAWAN K AGRAWAL

Natural Product Inc.
7963, Anderson Park Lane,
Westerville, Ohio 43081, USA
agrawal@naturalproduct.us

EDITORS

PROFESSOR ALEJANDRO F. BARRERO

Department of Organic Chemistry,
University of Granada,
Campus de Fuente Nueva, s/n, 18071, Granada, Spain
afbarre@ugr.es

PROFESSOR ALESSANDRA BRACA

Dipartimento di Chimica Bioorganica e Biofarmacia,
Università di Pisa,
via Bonanno 33, 56126 Pisa, Italy
braca@farm.unipi.it

PROFESSOR DEAN GUO

State Key Laboratory of Natural and Biomimetic Drugs,
School of Pharmaceutical Sciences,
Peking University,
Beijing 100083, China
gda5958@163.com

PROFESSOR YOSHIHIRO MIMAKI

School of Pharmacy,
Tokyo University of Pharmacy and Life Sciences,
Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan
mimaki@ps.toyaku.ac.jp

PROFESSOR STEPHEN G. PYNE

Department of Chemistry
University of Wollongong
Wollongong, New South Wales, 2522, Australia
spyne@uow.edu.au

PROFESSOR MANFRED G. REINECKE

Department of Chemistry,
Texas Christian University,
Forts Worth, TX 76129, USA
m.reinecke@tcu.edu

PROFESSOR WILLIAM N. SETZER

Department of Chemistry
The University of Alabama in Huntsville
Huntsville, AL 35809, USA
wssetzer@chemistry.uah.edu

PROFESSOR YASUHIRO TEZUKA

Institute of Natural Medicine
Institute of Natural Medicine, University of Toyama,
2630-Sugitani, Toyama 930-0194, Japan
tezuka@inm.u-toyama.ac.jp

PROFESSOR DAVID E. THURSTON

Department of Pharmaceutical and Biological Chemistry,
The School of Pharmacy,
University of London, 29-39 Brunswick Square,
London WC1N 1AX, UK
david.thurston@pharmacy.ac.uk

HONORARY EDITOR

PROFESSOR GERALD BLUNDEN

The School of Pharmacy & Biomedical Sciences,
University of Portsmouth,
Portsmouth, PO1 2DT U.K.
axuf64@dsl.pipex.com

ADVISORY BOARD

Prof. Berhanu M. Abegaz
Gaborone, Botswana

Prof. Viqar Uddin Ahmad
Karachi, Pakistan

Prof. Øyvind M. Andersen
Bergen, Norway

Prof. Giovanni Appendino
Novara, Italy

Prof. Yoshinori Asakawa
Tokushima, Japan

Prof. Lee Banting
Portsmouth, U.K.

Prof. Julie Banerji
Kolkata, India

Prof. Anna R. Bilia
Florence, Italy

Prof. Maurizio Bruno
Palermo, Italy

Prof. César A. N. Catalán
Tucumán, Argentina

Prof. Josep Coll
Barcelona, Spain

Prof. Geoffrey Cordell
Chicago, IL, USA

Prof. Ana Cristina Figueiredo
Lisbon, Portugal

Prof. Cristina Gracia-Viguera
Murcia, Spain

Prof. Duvvuru Gunasekar
Tirupati, India

Prof. Kurt Hostettmann
Lausanne, Switzerland

Prof. Martin A. Iglesias Arteaga
Mexico, D. F., Mexico

Prof. Leopold Jirovetz
Vienna, Austria

Prof. Vladimir I Kalinin
Vladivostok, Russia

Prof. Niel A. Koobanally
Durban, South Africa

Prof. Karsten Krohn
Paderborn, Germany

Prof. Chiaki Kuroda
Tokyo, Japan

Prof. Hartmut Laatsch
Göttingen, Germany

Prof. Marie Lacaille-Dubois
Dijon, France

Prof. Shoen-Sheng Lee
Taipei, Taiwan

Prof. Francisco Macias
Cadiz, Spain

Prof. Imre Mathe
Szeged, Hungary

Prof. Ermino Murano
Trieste, Italy

Prof. M. Soledade C. Pedras
Saskatoon, Canada

Prof. Luc Pieters
Antwerp, Belgium

Prof. Peter Proksch
Düsseldorf, Germany

Prof. Phila Raharivelomanana
Tahiti, French Polynesia

Prof. Luca Rastrelli
Fisciano, Italy

Prof. Monique Simmonds
Richmond, UK

Dr. Bikram Singh
Palampur, India

Prof. John L. Sorensen
Manitoba, Canada

Prof. Valentin Stonik
Vladivostok, Russia

Prof. Winston F. Tinto
Barbados, West Indies

Prof. Sylvia Urban
Melbourne, Australia

Prof. Karen Valant-Vetschera
Vienna, Austria

INFORMATION FOR AUTHORS

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site <http://www.naturalproduct.us>.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the Natural Product Inc. (NPI). A photocopy license is available from the NPI for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

To Subscribe: Natural Product Communications is a journal published monthly. 2013 subscription price: US\$2,395 (Print, ISSN# 1934-578X); US\$2,395 (Web edition, ISSN# 1555-9475); US\$2,795 (Print + single site online); US\$595 (Personal online). Orders should be addressed to Subscription Department, Natural Product Communications, Natural Product Inc., 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

Natural Product Communications

2013

Volume 8, Number 12

Contents

<u>Original Paper</u>	<u>Page</u>
New Humulenes from <i>Hyptis incana</i> (Labiatae) Mitsuru Satoh, Yoshio Satoh, Yasuhiro Anzai, Daisuke Ajisawa, Keiichi Matsuzaki, Mitsuko Makino and Yasuo Fujimoto	1665
Inhibitory Effects against Pasture Weeds in Brazilian Amazonia of Natural Products from the Marine Brown Alga <i>Dictyota menstrualis</i> Rainiomar Raimundo Fonseca, Antonio Pedro Silva Souza Filho, Roberto Campos Villaca and Valéria Laneuville Teixeira	1669
Isolation of the Plant Hormone (+)-Absciscic acid as an Antimycobacterial Constituent of the Medicinal Plant Endophyte <i>Nigrospora</i> sp. Trevor N. Clark, Katelyn Ellsworth, Haoxin Li, John A. Johnson and Christopher A. Gray	1673
New Cembranoid Diterpene from the South China Sea Soft Coral <i>Sarcophyton</i> sp. Fei Cao, Jing Zhou, Kai-Xia Xu, Meng-Qi Zhang and Chang-Yun Wang	1675
Crotofolane Diterpenoids from <i>Croton caracasanus</i> Katuska Chávez, Reinaldo S. Compagnone, Ricarda Riina, Alexander Briceño, Teresa González, Emilio Squitieri, Carlos Landaeta, Humberto Soscún and Alírica I. Suárez	1679
Development and Validation of a Modified Ultrasound-Assisted Extraction Method and a HPLC Method for the Quantitative Determination of Two Triterpenic Acids in <i>Hedyotis diffusa</i> Yu-Chiao Yang, Ming-Chi Wei, Hui-Fen Chiu and Ting-Chia Huang	1683
New Triterpenoid Saponins from the Roots of <i>Saponaria officinalis</i> Barbara Moniuszko-Szajwaj, Łukasz Pecio, Mariusz Kowalczyk, Ana M. Simonet, Francisco A. Macias, Małgorzata Szumacher-Strabel, Adam Cieślak, Wiesław Oleszek and Anna Stochmal	1687
Minor Triterpene Saponins from Underground Parts of <i>Lysimachia thyriflora</i>: Structure elucidation, LC-ESI-MS/MS Quantification, and Biological Activity Irma Podolak, Paweł Zmudzki, Paulina Koczurkiewicz, Marta Michalik, Paweł Zajdel and Agnieszka Galanty	1691
Variation of Saponin Contents and Physiological Status in <i>Quillaja saponaria</i> Under Different Environmental Conditions Angélica Grandón S, Miguel Espinosa B, Darcy Ríos L, Manuel Sánchez O, Katia Sáez C, Víctor Hernández S. and José Becerra A	1697
New Access to 7,17-seco C₁₉-Diterpenoid Alkaloids via Vacuum Pyrolysis of <i>N</i>-Deethyl-8-acetyl Derivatives Ling Wang, Qi-Feng Chen and Feng-Peng Wang	1701
Alkaloids from <i>Boophone haemanthoides</i> (Amaryllidaceae) Jerald J. Nair, Lucie Rárová, Miroslav Strnad, Jaume Bastida and Johannes van Staden	1705
Supinidine Viridiflorates from the Roots of <i>Chromolaena pulchella</i> Mario A. Gómez-Hurtado, J. Martín Torres-Valencia, Rosa E. del Río, Gabriela Rodríguez-García, Virginia Motilva, Sofia García-Mauriño, Carlos M. Cerda-García-Rojas and Pedro Joseph-Nathan	1711
<i>N</i>-Containing Metabolites from the Marine Sponge <i>Agelas clathrodes</i> Fan Yang, Rui-Hua Ji, Jiang Li, Jian-Hong Gan and Hou-Wen Lin	1713
Two New Compounds and Anti-complementary Constituents from <i>Amomum tsao-ko</i> Jiahong Jin, Zhihong Cheng and Daofeng Chen	1715
Antiangiogenic Activity of Flavonoids from <i>Melia azedarach</i> Shigenori Kumazawa, Satomi Kubota, Haruna Yamamoto, Naoki Okamura, Yasumasa Sugiyama, Hirokazu Kobayashi, Motoyasu Nakanishi and Toshiro Ohta	1719
Application of Mixture Analysis to Crude Materials from Natural Resources (IV)^[1(a-c)]: Identification of <i>Glycyrrhiza</i> Species by Direct Analysis in Real Time Mass Spectrometry (II) Eriko Fukuda, Yoshihiro Uesawa, Masaki Baba and Yoshihito Okada	1721
Comparison of Total Phenolic Content, Scavenging Activity and HPLC-ESI-MS/MS Profiles of Both Young and Mature Leaves and Stems of <i>Andrographis paniculata</i> Lee Suan Chua, Ken Choy Yap and Indu Bala Jaganath	1725
Xanthones from aerial parts of <i>Hypericum laricifolium</i> Juss. Irama Ramírez-González, Juan Manuel Amaro-Luis and Ali Bahsas	1731
A New Xanthone from the Pericarp of <i>Garcinia mangostana</i> Manqin Fu, Samuel X. Qiu, Yujuan Xu, Jijun Wu, Yulong Chen, Yuanshan Yu and Gengsheng Xiao	1733
Isolation of a Phomoxanthone A Derivative, a New Metabolite of Tetrahydroxanthone, from a <i>Phomopsis</i> sp. Isolated from the Mangrove, <i>Rhizophora mucronata</i> Yoshihito Shiono, Takehiro Sasaki, Fumiaki Shibuya, Yukito Yasuda, Takuya Koseki and Unang Supratman	1735
Anti-allergic Inflammatory Effects of Cyanogenic and Phenolic Glycosides from the Seed of <i>Prunus persica</i> Geum Jin Kim, Hyun Gyu Choi, Ji Hyang Kim, Sang Hyun Kim, Jeong Ah Kim and Seung Ho Lee	1739
Isolation, Synthesis and Biological Evaluation of Phenylpropanoids from the Rhizomes of <i>Alpania galanga</i> Sumit S Chourasiya, Eppakayala Sreedhar, K. Suresh Babu, Nagula Shankaraiah, V. Lakshma Nayak, S. Ramakrishna, S. Sravani and M.V. Basaveswara Rao	1741

Continued Overleaf

Rosmarinic Acid Interaction with Planktonic and Biofilm <i>Staphylococcus aureus</i> Livia Slobodniková, Silvia Fialová, Helena Hupková and Daniel Grančai	1747
New Butenolide and Pentenolide from <i>Dysidea cinerea</i> Phan Van Kiem, Nguyen Xuan Nhiem, Ngo Van Quang, Chau Van Minh, Nguyen Hoai Nam, Nguyen Thi Cuc, Hoang Le Tuan Anh, Bui Huu Tai, Pham Hai Yen, Nguyen Xuan Cuong, Nguyen Phuong Thao, Nguyen Thi Hoai, Nan Young Kim, Seon Ju Park and Kim Seung Hyun	1751
A New Cyclopeptide from Endophytic <i>Streptomyces</i> sp. YIM 64018 Xueqiong Yang, Yabin Yang, Tianfeng Peng, Fangfang Yang, Hao Zhou, Lixing Zhao, Lihua Xu and Zhongtao Ding	1753
Involvement of Trypsin-Digested Silk Peptides in the Induction of RAW264.7 Macrophage Activation Kyoung-Ho Pyo, Min-Ki Kim, Kwang-Soon Shin, Hyang Sook Chun and Eun-Hee Shin	1755
Low-Volatile Lipophilic Compounds in Needles, Defoliated Twigs, and Outer Bark of <i>Pinus thunbergii</i> Alexander V. Shpatov, Sergey A. Popov, Olga I. Salnikova, Ekaterina A. Khokhrina, Emma N. Shmidt and Byung Hun Um	1759
Lipid Constituents of the Edible Mushroom, <i>Pleurotus giganteus</i> Demonstrate Anti-Candida Activity Chia-Wei Phan, Guan-Serm Lee, Ian G. Macreadie, Sri Nurestri Abd Malek, David Pamela and Vikineswary Sabaratnam	1763
Effect of Trehalose Addition on Volatiles Responsible for Strawberry Aroma Mirela Kopjar, Janez Hribar, Marjan Simčič, Emil Zlatič, Tomaž Požrl and Vlasta Piližota	1767
<i>Pogostemon hirsutus</i> Oil, Rich in Abietane Diterpenes Ramar Murugan, Gopal Rao Mallavarapu, Veerappan Sudha and Pemaiah Brindha	1771
Combined Analysis of the Root Bark Oil of <i>Cleistopholis glauca</i> by Chromatographic and Spectroscopic Techniques Zana A. Ouattara, Jean Brice Boti, Coffy Antoine Ahibo, Félix Tomi, Joseph Casanova and Ange Bighelli	1773
Bioactivities and Compositional Analyses of <i>Cinnamomum</i> Essential Oils from Nepal: <i>C. camphora</i>, <i>C. tamala</i>, and <i>C. glaucescens</i> Prabodh Satyal, Prajwal Paudel, Ambika Poudel, Noura S. Dosoky, Kiran Kumar Pokharel, and William N. Setzer	1777
Essential Oil Characterization of Two Azorean <i>Cryptomeria japonica</i> Populations and Their Biological Evaluations Cristina Moiteiro, Teresa Esteves, Luís Ramalho, Rosario Rojas, Sandra Alvarez, Susana Zacchino and Helena Bragança	1785
Antioxidant, Antiproliferative and Antimicrobial Activities of the Volatile Oil from the Wild Pepper <i>Piper capense</i> Used in Cameroon as a Culinary Spice Verlaine Woguem, Filippo Maggi, Hervet P. D. Fogang, Léon A. Tapondjou, Hilaire M. Womeni, Luana Quassinti, Massimo Bramucci, Luca A. Vitali, Dezemona Petrelli, Giulio Lupidi, Fabrizio Papa, Sauro Vittori and Luciano Barboni	1791

Review/Account

Boldine and Related Aporphines: From Antioxidant to Antiproliferative Properties Darina Muthna, Jana Cmielova, Pavel Tomsik and Martina Rezacova	1797
New Therapeutic Potentials of Milk Thistle (<i>Silybum marianum</i>) Nataša Milić, Nataša Milošević, Ljiljana Suvajdžić, Marija Žarkov and Ludovico Abenavoli	1801
Biomedical Properties of Edible Seaweed in Cancer Therapy and Chemoprevention Trials: A Review Farideh Namvar, Paridah Md. Tahir, Rosfarizan Mohamad, Mahnaz Mahdavi, Parvin Abedi, Tahereh Fathi Najafi, Heshu Sulaiman Rahman and Mohammad Jawaid	1811
Methods for Extraction and Determination of Phenolic Acids in Medicinal Plants: A Review Agnieszka Arceusz, Marek Wesolowski and Pawel Konieczynski	1821

<u>Manuscripts in Press</u>	1830
------------------------------------	------

Cummulative Index

Contents	<i>i-xviii</i>
Author Index	<i>i-viii</i>
Keywords Index	<i>i-vii</i>

LIST OF AUTHORS

Abedi, P.....	1811	Grandón S, A.....	1697	Nakanishi, M.....	1719	Sravani, S.....	1741
Abenavoli, L.....	1801	Gray, CA.....	1673	Nam, NH.....	1751	Sreedhar, E.....	1741
Ahibo, CA.....	1773			Namvar, F.....	1811	Stochmal, A.....	1687
Ajisawa, D.....	1665	Hernández S., V.....	1697	Nhiem, NX.....	1751	Strnad, M.....	1705
Alvarez, S.....	1785	Hoai, NT.....	1751			Suárez, AI.....	1679
Amaro-Luis, JM.....	1731	Hribar, J.....	1767	Ohta, T.....	1719	Sudha, V.....	1771
Anh, HLT.....	1751	Huang, TC.....	1683	Okada, Y.....	1721	Sugiyama, Y.....	1719
Anzai, Y.....	1665	Hupková, H.....	1747	Okamura, N.....	1719	Supratman, U.....	1735
Arceusz, A.....	1821	Hyun, KS.....	1751	Oleszek, W.....	1687	Suresh Babu, K.....	1741
				Ouattara, ZA.....	1773	Suvajdžić, L.....	1801
Baba, M.....	1721	Jaganath, IB.....	1725			Szumacher-Strabel, M.....	1687
Bahsas, A.....	1731	Jawaid, M.....	1811	Pamela, D.....	1763		
Barboni, L.....	1791	Ji, RH.....	1713	Papa, F.....	1791	Tahir, PM.....	1811
Basaveswara Rao, MV.....	1741	Jin, J.....	1715	Park, SJ.....	1751	Tai, BH.....	1751
Bastida, J.....	1705	Johnson, JA.....	1673	Paudel, P.....	1777	Tapondjou, LA.....	1791
Becerra A, J.....	1697	Joseph-Nathan, P.....	1711	Pecio, L.....	1687	Teixeira, VL.....	1669
Bighelli, A.....	1773			Peng, T.....	1753	Thao, NP.....	1751
Boti, JB.....	1773	Khokhrina, EA.....	1759	Petrelli, D.....	1791	Tomi, F.....	1773
Bragança, H.....	1785	Kiem, PV.....	1751	Phan, CW.....	1763	Tomsik, P.....	1797
Bramucci, M.....	1791	Kim, GJ.....	1739	Piližota, V.....	1767	Torres-Valencia, JM.....	1711
Briceño, A.....	1679	Kim, JA.....	1739	Podolak, I.....	1691		
Brindha, P.....	1771	Kim, JH.....	1739	Pokharel, KK.....	1777	Uesawa, Y.....	1721
		Kim, MK.....	1755	Popov, SA.....	1759	Um, BH.....	1759
Cao, F.....	1675	Kim, NY.....	1751	Poudel, A.....	1777		
Casanova, J.....	1773	Kim, SH.....	1739	Požrl, T.....	1767	van Staden, J.....	1705
Cerda-García-Rojas, CM.....	1711	Kobayashi, H.....	1719	Pyo, KH.....	1755	Villaça, RC.....	1669
Chávez, K.....	1679	Koczurkiewicz, P.....	1691			Vitali, LA.....	1791
Chen, D.....	1715	Konieczynski, P.....	1821	Qiu, SX.....	1733	Vittori, S.....	1791
Chen, QF.....	1701	Kopjar, M.....	1767	Quang, NV.....	1751		
Chen, Y.....	1733	Koseki, T.....	1735	Quassinti, L.....	1791	Wang, L.....	1701
Cheng, Z.....	1715	Kowalczyk, M.....	1687			Wang, CY.....	1675
Chiu, HF.....	1683	Kubota, S.....	1719	Rahman, HS.....	1811	Wang, FP.....	1701
Choi, HG.....	1739	Kumazawa, S.....	1719	Ramakrishna, S.....	1741	Wei, MC.....	1683
Chourasiya, SS.....	1741			Ramvalho, L.....	1785	Wesolowski, M.....	1821
Chua, LS.....	1725	Lakshma Nayak, V.....	1741	Ramírez-González, I.....	1731	Woguemi, V.....	1791
Chun, HS.....	1755	Landaeta, C.....	1679	Rárová, L.....	1705	Womeni, HM.....	1791
Cieślak, A.....	1687	Lee, GS.....	1763	Rezacova, M.....	1797	Wu, J.....	1733
Clark, TN.....	1673	Lee, SH.....	1739	Riina, R.....	1679		
Cmielova, J.....	1797	Li, H.....	1673	Río, RE.....	1711	Xiao, G.....	1733
Compagnone, R S.....	1679	Li, J.....	1713	Ríos LD.....	1697	Xu, KX.....	1675
Cuc, NT.....	1751	Lin, HW.....	1713	Rodríguez-García, G.....	1711	Xu, L.....	1753
Cuong, NX.....	1751	Lupidi, G.....	1791	Rojas, R.....	1785	Xu, Y.....	1733
Ding, Z.....	1753	Macias, FA.....	1687	Sabaratham, V.....	1763	Yamamoto, H.....	1719
Dosoky, NS.....	1777	Macreadie, IG.....	1763	Sáez C, K.....	1697	Yang, F.....	1713
		Maggi, F.....	1791	Salnikova, OI.....	1759	Yang, F.....	1753
Ellsworth, K.....	1673	Mahdavi, M.....	1811	Sánchez O, M.....	1697	Yang, X.....	1753
Espinosa B, M.....	1697	Makino, M.....	1665	Sasaki, T.....	1735	Yang, Y.....	1753
Esteves, T.....	1785	Malek, SNA.....	1763	Satoh, M.....	1665	Yang, YC.....	1683
		Mallavarapu, GR.....	1771	Satoh, Y.....	1665	Yap, KC.....	1725
Fialová, S.....	1747	Matsuzaki, K.....	1665	Satyral, P.....	1777	Yasuda, Y.....	1735
Filho, APSS.....	1669	Michalik, M.....	1691	Setzer, WN.....	1777	Yen, PH.....	1751
Fogang, HPD.....	1791	Milić, N.....	1801	Shankaraiah, N.....	1741	Yu, Y.....	1733
Fonseca, RR.....	1669	Milošević, N.....	1801	Shibuya, F.....	1735		
Fu, M.....	1733	Minh, CV.....	1751	Shin, EH.....	1755	Zacchino, S.....	1785
Fujimoto, Y.....	1665	Mohamad, R.....	1811	Shin, KS.....	1755	Zajdel, P.....	1691
Fukuda, E.....	1721	Moiteiro, C.....	1785	Shiono, Y.....	1735	Zarkov, M.....	1801
		Moniuszko-Szajwaj, B.....	1687	Shmidt, EN.....	1759	Zhang, MQ.....	1675
Galanty, A.....	1691	Motilva, V.....	1711	Shpatov, AV.....	1759	Zhao, L.....	1753
Gan, JH.....	1713	Murugan, R.....	1771	Simčić, M.....	1767	Zhou, H.....	1753
García-Mauriño, S.....	1711	Muthna, D.....	1797	Simonet, AM.....	1687	Zhou, J.....	1675
Gómez-Hurtado, MA.....	1711			Slobodníková, L.....	1747	Zlatić, E.....	1767
González, T.....	1679	Nair, JJ.....	1705	Soscún, H.....	1679	Žmudzki, P.....	1691
Grančai, D.....	1747	Najafi, TF.....	1811	Squitieri, E.....	1679		

New Humulenes from *Hyptis incana* (Labiatae)Mitsuru Satoh^{a,*}, Yoshio Satoh^b, Yasuhiro Anzai^a, Daisuke Ajisawa^a, Keiichi Matsuzaki^c, Mitsuko Makino^c and Yasuo Fujimoto^d^aLaboratory of Drug Design and Medicinal Chemistry, Showa Pharmaceutical University, 3-3165 Higashitamagawagakuen, Machida, Tokyo 194-8543 Japan^bDepartment of Clinical and Biomedical Sciences, Showa Pharmaceutical University, 3-3165 Higashitamagawagakuen, Machida, Tokyo 194-8543 Japan^cSchool of Pharmacy, Nihon University, 7-7-1 Narashinodai, Funabashi, Chiba 274-8555, Japan^dDepartment of General Studies, College of Humanities and Sciences, Nihon University, 3-25-40 Sakurajosui, Setagaya-ku, Tokyo 156-8550, Japan

ms@ac.shoyaku.ac.jp

Received: July 16th, 2013; Accepted: September 8th, 2013

Two new humulene-type sesquiterpenes, named hyptishumulene I (**1**) and II (**2**), have been isolated, together with eight known compounds, a humulene-type sesquiterpene (**3**), a monoterpene (**4**) and six abietane-type diterpenoids (**5-10**) from the aerial parts of *Hyptis incana* (Labiatae). The cytotoxic activity of the isolated compounds against mouse leukemia cells (L1210) was examined. The abietane-type diterpenoids (**5-10**) showed rather potent growth inhibitory activity ($IC_{50} < 15 \mu M$), while the new humulene-type compounds (**1** and **2**) exhibited moderate activity ($IC_{50} > 50 \mu M$).

Keywords: *Hyptis incana*, Labiatae, Humulene-type sesquiterpene, Cytotoxicity, Leukemia cell.

Hyptis incana Willd. ex Steudel (Labiatae, local name in Brazil: Salva do Marajo) is distributed widely throughout the Amazon area in South America. The plant has been used by native people as a tonic, to control perspiration, and to treat acne [1]. During the course of our studies on the isolation of biologically active substances from South American medicinal plants [2,3], we investigated the cytotoxic activity of alcohol extracts of this plant and found that the extracts exhibited potent growth inhibitory activity against leukemia cells (L1210). In this paper, we describe the isolation from *H. incana*, structure elucidation and growth inhibitory activity of two new humulene-type sesquiterpenes, named hyptishumulene I (**1**) and II (**2**), along with eight known compounds, a humulene-type sesquiterpene (**3**), which was isolated from *Torilis japonica* [4] and the red alga *Laurencia lobata* [5], a monoterpene (**4**) [6], six abietane-type diterpenes {7-methoxyrosmanol (**5**) [7], 7-ethoxyrosmanol (**6**) [7], rosmanol (**7**) [7,8], epirosmanol (**8**) [8], isorosmanol (**9**) [8] and saffcinolide (**10**) [9]}.

The dried aerial parts of *H. incana* were extracted with 80% EtOH. The crude extract was fractionated on a Diaion HP-20 column eluted successively with H₂O, 40% MeOH, 70% MeOH, MeOH, and acetone. The growth inhibitory activity of each fraction against L1210 cells was tested, and the 70% MeOH and MeOH fractions (>90% inhibition at the concentration of 50 $\mu g/mL$) were found to exhibit potent activities. Compounds (**1-10**) were isolated from these fractions by purification using silica gel chromatography and HPLC (Figure 1).

Hyptishumulene I (**1**) was obtained as a colorless amorphous powder, $[\alpha]_D^{20} -24.6$ ($c = 0.1$, CHCl₃). Its high-resolution HR-EI-MS showed the $[M]^+$ ion peak at m/z 236.1791, corresponding to the molecular formula C₁₅H₂₄O₂ (calcd. for 236.1777). The ¹H-, ¹³C-, ¹H-¹H correlation spectroscopy (COSY) and heteronuclear single quantum coherence (HSQC) spectra of **1** showed signals due to two tertiary methyls (δ_H 0.98- δ_C 30.8, δ_H





Figure 1: Structures of Compounds 1-10.

1.16- δ_C 24.3), a vinyl methyl (δ_H 1.73- δ_C 17.0), three methylenes, (δ_H 1.45, 1.85- δ_C 49.7, δ_H 2.03, 2.74- δ_C 35.3, δ_H 2.66, 2.98- δ_C 35.0), an exomethylene (δ_H 4.87, 4.99- δ_C 113.2), two *sec*-alcohols (δ_H 4.41- δ_C 67.3, δ_H 4.06- δ_C 77.2), a *trans*-disubstituted double bond [δ_H 5.34 (d, $J = 16.0$ Hz) - δ_C 141.5, δ_H 5.39 (ddd, $J = 6.0, 6.2, 16.0$ Hz) - δ_C 124.0], a vinyl proton (δ_H 5.11- δ_C 122.8), an sp³ quaternary carbon (δ_C 36.0), and two sp² quaternary carbons (δ_C 138.3, 149.6).

Table 1: ^1H - and ^{13}C -NMR data for hyptishumulene **1** (**1**) and **2** (**2**).

position	^1H -NMR (600 MHz, δ , CDCl_3)		^{13}C -NMR (150 MHz, δ , CDCl_3)	
	1	2	1	2
1			36.0	35.1
2 α	1.85 dd (6.3, 14.1)	1.76 dd (11.6, 12.2)	49.7	46.0
2 β	1.45 dd (2.1, 14.1)	1.40 dd (2.2, 12.2)		
3	4.41 dd (2.1, 6.3)	4.13 dd (2.2, 11.6)	67.3	75.8
4			138.3	139.9
5	5.11 ddd (1.8, 1.8, 12.0)	5.09 ddd (0.8, 7.7, 7.9)	122.8	125.0
6 α	2.03 ddd (1.8, 6.0, 14.0)	2.64 ddd (4.1, 7.9, 12.0)	35.3	37.0
6 β	2.74 dd (7.2, 12.0, 14.0)	2.14 dd (7.7, 10.6, 12.0)		
7	4.06 dd (6.0, 7.2)	3.92 dd (4.1, 10.6)	77.2	72.9
8			149.6	153.3
9 α	2.98 dd (6.0, 14.0)	2.93 dd (5.3, 12.2)	35.0	41.3
9 β	2.66 dd (6.2, 14.0)	2.52 dd (10.3, 12.2)		
10	5.39 ddd (6.0, 6.2, 16.0)	4.85 ddd (5.3, 10.3, 15.5)	124.0	124.3
11	5.34 d (16.0)	5.05 d (15.5)	141.5	138.8
12	1.16 s	1.14 s	24.3	32.5
13	0.98 s	0.98 s	30.8	23.8
14	1.73 t (1.8)	1.59 d (0.8)	17.0	10.1
15	4.87 d (1.8)	5.01 s	113.2	112.0
	4.99 d (1.8)	5.21 s		


The numbers in parentheses are J values in Hz**Figure 2:** HMBC correlations for **1** and **2**.

The presence of the *sec*-alcohol groups was further confirmed by acetylation. Treatment of **1** with acetic anhydride-pyridine gave a diacetate whose oxymethine proton signals showed acetylation shifts (δ_{H} 4.06 \rightarrow δ_{H} 4.77, δ_{H} 4.41 \rightarrow δ_{H} 5.44). Connectivity of these functional groups was investigated by measuring the heteronuclear multiple bond coherence (HMBC) spectrum.

As shown in Figure 2, the proton signals at δ_{H} 1.16 and δ_{H} 0.98 assignable to a *geminal* dimethyl group (C-12 and -13) showed correlations with carbon signals at δ_{C} 141.5 (C-11), δ_{C} 36.0 (C-1) and δ_{C} 49.7 (C-2), suggesting the connectivity from C-10 to C-2 through C-1. The vinyl methyl proton signal at δ_{H} 1.73 (H-14) exhibited long-range correlations with the carbon signals at δ_{C} 138.3 (C-4), 122.8 (C-5) and δ_{C} 67.3 (C-3), which showed the cross peaks with the proton signals at δ_{H} 1.45 and 1.85 (H-2). These results indicated the presence of the C-3-C-5 sequence, including the vinyl methyl group at C-14 and the connectivity of C-3 and C-2. In addition, the HMBC spectrum of **1** showed correlations between the proton signal at δ_{H} 4.06 (H-7) due to the other *sec*-alcohol group and the carbon signals at δ_{C} 122.8 (C-5), 35.3 (C-6), δ_{C} 113.2 (C-15) and δ_{C} 149.6 (C-8), which, in turn, correlated with the proton signals at δ_{H} 2.66 and 2.98 (H-9) and at δ_{H} 5.39 (H-10). Thus, the structure of **1** was deduced to be a humulene-type sesquiterpene (Table 1).


The stereostructure of **1** was investigated by analysis of its nuclear Overhauser effect enhancement (NOE) difference spectra. As shown in Figure 3, the proton signal at δ 1.73 (H-14), due to a vinyl methyl group, showed a cross peak with the vinyl proton signal at δ 5.11 (H-5), which had correlation with the proton signal at δ 4.06 (H-7), thereby suggesting the *Z*-configuration of the C-(4)-C-(5) double bond and α -configuration of the hydroxy group at C-7 [10].

The coupling constants of H-7 (J = 6.0, 7.2 Hz) are also supportive of the β -configuration for H-7 by taking into account the Karplus' rule, because the dihedral angles between H-7 and H-6 α , and H-6 β in hyptishumulene are approximately 170° and 20°, respectively, on inspection using a molecular model. The stereochemistry at C-3 was also deduced by difference NOE experiments on **1**. Irradiation at δ 4.41 (H-3) produced significant enhancement of the proton signal at δ 1.16 (H-12) and δ 2.74 (H-6 β), indicating an α -configuration of the hydroxy group at C-3. Furthermore, NOEs were observed between the following proton signals: H-7 and H-5, H-15; H-9 α and H-11; H-10 and H-9 β , H-12; H-11 and H-9 α , H-13; H-2 α and H-13, H-14; and H-2 β and H-12. These spectral data are explained satisfactorily by the stereostructure shown in Figure 3. The structure of **1** was also supported by it being the most stable (Figure 3', minimized energy: 13.2963 kcal/mol) by molecular mechanics calculation (MM2) [11].

**Figure 3:** NOE correlations for **1**.**Figure 3'**

Hyptishumulene **2** was obtained as a colorless amorphous powder [α]_D²⁰ -30.4° (c = 0.1, CHCl_3). Its HR-EI-MS showed the $[\text{M}]^+$ ion peak at m/z 236.1786, corresponding to the molecular formula $\text{C}_{15}\text{H}_{24}\text{O}_2$ (calcd. for 236.1777), which is the same as that of compound **1**.

The ^1H - and ^{13}C -NMR spectra of **2** were very similar to those of **1**, except for a few differences (Tables 1). The spectral data indicated that **2** should be a stereoisomer of **1**. Comparison of the NOESY spectra of **1** and **2**, showed that the H-5 signal (δ 5.09) of **2** had cross peaks with H-3 (δ 4.13) and H-6 β (δ 2.14), while the corresponding signal (δ 5.11) of **1** exhibited correlations with H-7 (δ 4.06) and H-14 (δ 1.73). These results indicated that the C-(4)-C-(5) double bond of **2** is the *E*-form. In addition, the β -orientation of the hydroxy group at C-7 was confirmed by observation of NOE correlations between H-7 and H6 α , H-9 α , as well as the NOEs shown in Figure 4. Thus, the structure of **2** was represented as shown in Fig. 1.

**Figure 4:** NOE Correlations for **2**.

The structures of compounds **3-10** were identified by comparison of their spectral data with those described in the literature.

Growth inhibitory activity: The effects of the isolated compounds (**1-10**) on the growth of leukemia cells (L1210) were examined. The abietane-type diterpenoids (**5-10**) showed rather potent growth inhibitory activity (IC_{50} < 15 μM), while the new humulene-type

compounds (**1** and **2**) exhibited moderate activity ($IC_{50} > 50 \mu M$) (Table 2). Compounds **5-9** induced apoptosis of neuroblastoma cells [12].

Table 2: Growth inhibitory effects of the compounds isolated from *H. incana* against leukemia cells (L1210) ($n=5$).

Compounds	IC_{50} (μM)
1	> 50
2	> 50
3	> 50
4	> 50
5	13
6	13
7	12
8	12
9	12
10	15

Experimental

General: 1H and ^{13}C NMR spectra were measured on Bruker AV300 and AV600 spectrometers in $CDCl_3$, with tetramethylsilane (TMS) as an internal standard. MS were recorded on a JEOL JMS-600 spectrometer. Optical rotations were recorded using a JASCO DIP-360 digital polarimeter. CC was performed on either Diaion HP-20 resin (Mitsubishi Kasei) or silica gel (Wakogel C-200). HPLC was conducted with JASCO PU-980 or Senshu SSC 3220 pumps equipped with an ERMA ERC-7522 (RI) detector. TLC was performed using silica gel 60 F_{254} (Merck) precoated plates and detection was carried out by spraying with 10% sodium phosphomolybdate in methanol, followed by heating.

Plant material: *Hyptis incana* was purchased in 1999, and identified by Pharmacist Gilbert Rubens Biancalana (Laboratorio Farmaervas Ltda. in Sao Paulo, Brazil). The voucher specimen was deposited in the herbarium of Showa Pharmaceutical University.

Extraction and isolation: The dried aerial parts of *H. incana* (2858 g) were extracted with 80% EtOH (20 l \times 3) with ultrasonication. The extract was concentrated *in vacuo* to give a brown solid (750 g). This (550 g) was chromatographed on a Diaion HP-20 column eluted successively with stepwise gradients of 10 L each of H_2O , 40% MeOH, 70% MeOH, MeOH, and acetone, and then each eluate was concentrated *in vacuo* to afford 5 fractions [H_2O (168.0 g), 40% MeOH (57.7 g), 70% MeOH (74.8 g), MeOH (143.6 g), and acetone (21.1 g)]. The cytotoxicity of each fraction against L1210 cells was tested, and the MeOH and 70% MeOH fractions were found to be the most active.

The 70% MeOH fraction (50.0 g) was chromatographed on a silica gel column eluted successively with solvents of increasing polarity [*n*-hexane: EtOAc = 1: 1 (5 L, Fr. 1-4), 1: 5 (5 L, Fr. 5-9), and EtOAc (5 L, Fr. 10)] to afford 10 fractions. Fraction 3 (3.0 g) was further fractionated by HPLC [Senshu Pak, PEGASIL-Silica 60-5, 10 \times 250 mm (column A), *n*-hexane: EtOAc = 1: 1, flow rate: 3.0 mL/min] to afford 15 fractions (Fr. 3-1 to 3-15). Fraction 3-11 (retention time (Rt): 18.0–19.5 min) was further purified by HPLC [PEGASIL-ODS, 10 \times 250 mm (column B), MeOH: H_2O = 8: 2, flow rate: 3.0 mL/min] to give **1** (13 mg, Rt: 8.0 min).

Hyptishumulene I (1)

Colorless amorphous powder.

$[\alpha]_D^{20}$: -24.6 (c 0.1, $CHCl_3$).

HR-EI-MS m/z : 236.1791 (calcd. for $C_{15}H_{24}O_2$ 236.1777),

1H NMR ($CDCl_3$, 600 MHz) and ^{13}C NMR ($CDCl_3$, 150 MHz): Table 1.

Fraction 3-12 (Rt: 20.5–22.0 min) was purified using HPLC (column B, MeOH: H_2O = 8: 2, flow rate: 3.0 mL/min) to give **2** (6 mg, Rt: 7.0 min).

Hyptishumulene II (2)

Colorless amorphous powder.

$[\alpha]_D^{20}$: -30.4 (c 0.1, $CHCl_3$).

HR-EI-MS m/z : 236.1786 (calcd. for $C_{15}H_{24}O_2$ 236.1777),

1H NMR ($CDCl_3$, 600 MHz) and ^{13}C NMR ($CDCl_3$, 150 MHz): Table 1.

Fraction 3-10 (retention time (Rt): 16.0–17.5 min) was further purified by HPLC [PEGASIL-ODS, 10 \times 250 mm (column B), MeOH: H_2O = 8: 2, flow rate: 3.0 mL/min] to give **3** (10 mg, Rt: 6.0 min).

Fraction 3-2 (retention time (Rt): 8.0–8.8 min) was further purified by HPLC [column A, *n*-hexane: EtOAc = 1: 1, flow rate: 3.0 mL/min] to give **4** (12 mg, Rt: 8.5 min). Purification of a part (100 mg) of fraction 1 by HPLC (column A, *n*-hexane: EtOAc = 3: 1, flow rate: 5.0 mL/min) gave **5** (methoxyrosmanol, 15 mg, Rt: 6.0 min) and **6** (ethoxyrosmanol, 33 mg, Rt: 7.5 min). Separation of a part (100 mg) of fraction 2 by HPLC (column A, *n*-hexane: EtOAc = 5: 2, flow rate: 5.0 mL/min) afforded **7** (rosmanol, 20.5 mg, Rt: 6.5 min), **8** (epirosmanol, 9.8 mg, Rt: 8.0 min) and **9** (isorosmanol, 7.3 mg, Rt: 13.0 min). Purification of part (50 mg) of fraction 2 by HPLC [PEGASIL-ODS, 20 \times 150 mm, MeOH, flow rate: 8.0 mL/min] gave **10** (safficinolide, 8.5 mg, Rt: 4.5 min).

Acetylation of hyptishumulene I (1): A mixture of **1** (1 mg) and five drops (large excess) of acetic anhydride in pyridine (0.5 mL) was allowed to stand for 12 h at room temperature. The mixture was poured into water and then extracted with AcOEt (10 mL). The AcOEt solution was washed with brine (10 mL \times 2), dried over Na_2SO_4 and concentrated *in vacuo* to give an acetate (1 mg).

1H NMR δ : 0.94 (3H, s), 0.99 (3H, s), 1.18 (3H, s), 1.57 (1H, m), 1.64 (1H, m), 1.92 (3H, s, -Ac), 1.99 (3H, s, -Ac), 2.00 (1H, m), 2.02 (1H, m), 2.76 (1H, m), 2.83 (1H, m), 4.77 (1H, m), 4.82 (1H, s), 4.89 (1H, s), 5.11 (1H, m), 5.36 (1H, m), 5.38 (1H, m), 5.44 (1H, m).

MS: m/z 320 $[M]^+$.

Cytotoxicity assay: The effects of isolated compounds on the growth of leukemia cells (L1210) were investigated as follows. Cells were suspended in RPMI 1640 medium (NISSUI) containing 1% kanamycin sulfate, 10% fetal bovine serum, and supplemented with L-glutamine. Cells were plated in a 96-well microplate (100 μL /well) at a density of 3.0×10^4 cells/well. After incubating the plate for 24 h at 37°C in a humidified 5% CO_2 atmosphere, compounds **1-10** in MeOH (2 μL /well) at various concentrations ($n = 5$) were added and the plate was incubated for an additional 48 h under the same conditions. Control wells ($n = 5$) containing the same volume of MeOH (2 μL /well) were incubated in each assay. Ten μL of Cell Counting Kit-8 [13, 14] was added to each well, then, the microplate was incubated for 3 h in a 5% CO_2 atmosphere at 37°C. The absorbance (A) of each well was measured at 450 nm using a Microplate reader (MTP-450, Corona Electric). The percent inhibition was calculated by comparing $A_{control}$ and $A_{treated}$.

$$\text{inhibition (\%)} = (1 - A_{\text{treated-blank}} / A_{\text{control-blank}}) \times 100$$

References and Notes

- [1] Hashimoto G. (1996) *Illustrated Cyclopedia of Brazilian Medicinal Plants*. p574, Aboc-Sha, Kamakura city, Japan.
- [2] Satoh M, Satoh Y, Fujimoto Y. (2000) Cytotoxic constituents from *Erythroxylum catuaba* isolation and cytotoxic activities of cinchonain. *Natural Medicines*, **54**, 97-100.
- [3] Satoh M, Satoh Y, Isobe K, Fujimoto Y. (2003) Studies on the constituents of *Leonurus sibiricus* L. *Chemical and Pharmaceutical Bulletin*, **51**, 341-342.
- [4] Kitajima J, Suzuki N, Satoh M, Watanabe M. (2002) Sesquiterpenoids of *Torilis japonica* fruit. *Phytochemistry*, **59**, 811-815
- [5] Takeda S, Iimura Y, Tanaka K, Kurosawa E, Suzuki T. (1990) A new naturally occurring racemic compound from the marine red alga *Laurencia obtusa* (Hudson) Lamouroux. *Chemistry Letters*, 155-156.
- [6] Marcelo JPF, Gilberto VR, Antonio JCB, Vicente PE. (2001) An auxiliary program for pattern recognition and substructure elucidation of monoterpenes. *Spectroscopy*, **15**, 65-98
- [7] Arisawa M, Hayashi T, Ohmura K, Nagayama K, Shimizu M, Morita N. (1987) Chemical and pharmaceutical studies on medicinal plants in Paraguay: studies on “ Romero “ Part 2. *Journal of Natural Products*, **50**, 1164-1166. 7-Ethoxyrosmanol (**6**) may be an artifact produced by extraction procedure with EtOH.
- [8] Nakatani N, Inatani R. (1984) Two antioxidative diterpenes from rosemary (*Rosmarinus officinalis* L.) and a revised structure for rosmanol. *Agricultural and Biological Chemistry*, **48**, 2081-2085.
- [9] Tada M, Okuno K, Chiba K, Ohnishi E, Yoshii T. (1994) Antiviral diterpenes from *Salvia officinalis*. *Phytochemistry*, **35**, 539-541.
- [10] The diastereomeric isomer at C-7 of **1** has been prepared by acid hydrolysis of humulene diepoxide A (Smith RJ, Mahiou B, Deinzer ML. (1991) Hydrolysis products of humulene diepoxide. *Tetrahedron*, **47**, 933-940). A comparison of the spectral data of **1** with those of the compound from humulene diepoxide A showed that these compounds were not identical.
- [11] CS Chem 3D Pro v. 12.0.2. Cambridge Soft Corp.: Cambridge, MA.
- [12] The experimental results on the apoptosis inducement were already published in *Anticancer Research*. (Tabata K, Kim M, Makino M, Satoh M, Satoh Y, Suzuki T. (2012) Phenolic diterpenes derived from *Hyptis incana* induce apoptosis and G₂/M arrest of neuroblastoma cells. *Anticancer Research*, **32**, 4781-4790.
- [13] Ishiyama N, Miyazono Y, Sasamoto K, Ohkura Y, Ueno K. (1997) A highly water-soluble disulfonated tetrazolium salt as a chromogenic indicator for NADH as well as cell viability. *Talanta*, **44**, 1299-1305.
- [14] Tominaga H, Ishiyama M, Ohseto F, Sasamoto K, Hamamoto T, Suzuki K, Watanabe K. (1999) A water-soluble tetrazolium salt useful for colorimetric cell viability assay. *Analytical Communications*, **36**, 47-50.

Inhibitory Effects against Pasture Weeds in Brazilian Amazonia of Natural Products from the Marine Brown Alga *Dictyota menstrualis*

Rainiomar Raimundo Fonseca^a, Antonio Pedro Silva Souza Filho^b, Roberto Campos Villaça^c and Valéria Laneuville Teixeira^{c,*}

^aPrograma de Pós-Graduação em Química, Instituto de Química, Universidade Federal Fluminense, Niterói, RJ, 24020-150, Brazil

^bLab. Agroindústria, EMBRAPA Amazônia Oriental/ CPATU, Belém, Pará, 66095-100, Brazil

^cDepartamento de Biologia Marinha, Instituto de Biologia, P.O. Box 100.644, Niterói, RJ, 24001-970, Brazil

valerialaneuville@gmail.com

Received: July 3rd, 2013; Accepted: October 7th, 2013

Fractions of the acetone extract and a mixture of two diterpenes from the marine brown alga *Dictyota menstrualis* were prepared with the aim of identifying potential effects on the germination of seeds and on elongation of the radicle and hypocotyl of the weeds *Mimosa pudica* and *Senna obtusifolia*. The bioassay on seed germination was performed in controlled conditions of 25°C temperature and a 12 hour photoperiod, while the one concerning radicle and hypocotyl elongation was performed at the same temperature, though adopting a photoperiod of 24 hours. The results varied according to the receptor species, the fraction utilized, and its concentration. TLC analysis of the fractions and comparison with isolated products indicated that the diterpenes pachydictyol A and isopachydictyol A were the most abundant compounds in fraction HE, whereas the diterpene 6-hydroxy-dichotomano-2, 13-diene-16, 17-dial (**3**) was the most abundant compound in fractions DC and EA. Analysis of less polar fractions (in *n*-hexane, dichloromethane and ethyl acetate) revealed values of less than 30% inhibition. On the other hand, the ethanol/water fraction was the most active in all instances. The biological activity of these fractions must be due to the presence of known diterpenes and/or sulfated polysaccharides isolated in earlier studies.

Keywords: *Dictyota menstrualis*, Marine natural product, *Mimosa pudica*, *Senna obtusifolia*, Inhibitory effects.

In the Amazon region, the occurrence of weeds is regarded as the most serious biological problem faced by the ranchers, and weed control represents one of the highest costs of production. Therefore, the definition of new strategies that enable reduction of infestation in pastures to an acceptable level may guarantee higher productivity in the long term by assuring the longevity of crops. In addition to increasing the economic profit, these strategies may be useful to soften the social and environmental dissatisfaction that this type of activity has awakened nationally and internationally due to the use of aggressive bio defensives [1-3]. Our group has been studying the natural products of marine brown algae for over 25 years. During this period, many biological activities were recognized for the natural products of benthic algae. In the previous study, we have shown the inhibitory effects of the crude extract from the marine red alga *Plocamium brasiliense* (Greville) M. A. Howe & W. R. Taylor [4]. The positive results obtained stimulated new tests with other species of seaweed.

The marine brown alga *Dictyota menstrualis* (Hoyt) Schnetter, Hörning & Weber-Peukert is very common along the Brazilian coast and known as a prolific producer of bioactive diterpenes [5-9]. The diterpenes are the main components of the hexane, dichloromethane and acetone extracts of these algae. On the other hand, the sulfated polysaccharides (heterofucans) are the major components of the more polar extracts from the Brazilian population [10].




Fifteen diterpenes are known for *D. menstrualis*, isolated from five populations in several regions of the world: one from North Carolina (USA) [11,12], one from the area around Sydney (Australia) [13], two from the coast of the State of Rio de Janeiro (Brazil) and two from the São Pedro and São Paulo Archipelago (Brazil) [8,9]. Besides, studies using diterpenes isolated from

D. menstrualis from the Brazilian littoral indicate that the secondary metabolites are used as a defense mechanism against amphipod herbivores [14].

Indeed, investigation of allelopathy has traditionally been conducted utilizing phytochemicals and metabolites from terrestrial plants [e.g. 15]. Allelopathy is considered to be a phenomenon in which a plant (including microorganisms) directly or indirectly causes either positive or negative effects on the development of other plants by liberating chemical substances in the environment. Nevertheless, a study on allelopathy may well be applied to a seaweed species such as *D. menstrualis* since it produces various bioactive diterpenes [8]. Allelopathy may represent an excellent alternative strategy for the management of pastures, allowing the control of infestations of undesired plants as weeds and, as a consequence, avoid great losses, especially regarding livestock. Moreover, this phenomenon has ecological importance, notably when it comes to providing alternative sources of chemical structures that can be used to produce agricultural biodefensives [16, 17].

In the present study, the inhibitory effects of fractions of different polarities, and a mixture of pachydictyol A and isopachydictyol A obtained from the acetone extract of *D. menstrualis* was evaluated in bioassays concerning germination and development of radicle and hypocotyl using as receptor species the weeds *Mimosa pudica* L. and *Senna obtusifolia* (L.) Irwin & Barney.

The inhibitory effects were evaluated of different fractions {*n*-hexane (HE), dichloromethane (DC), ethyl acetate (EA), and ethanol/water (HA)} of an acetone extract of *D. menstrualis*. The aim of the first assay was to analyze the effects of these fractions on the germination of seeds (Figure 1a-c). The results presented in Figure 1 revealed that the ethanol/water fraction(HA) had a






■ *Mimosa pudica* ■ *Senna obtusifolia*

Figure 1: Effects of fractions HE, DC, EA and HA on (a) the germination of seeds, (b) on radicle elongation, and (c) on hypocotyl elongation of two weeds (*Mimosa pudica* and *Senna obtusifolia*). The data are expressed as percentage of inhibition in relation to control treatment (distilled water). HE = *n*-Hexane fraction, DC = Dichloromethane fraction, EA = Ethyl acetate fraction, and HA = Ethanol/water (7:3) fraction. Extracts with the same solvent from *M. pudica* and *S. obtusifolia* do not differ by Tukey (5%).

potent inhibitory effect on the germination of seeds, ranging from 88% to 77% inhibition for *M. pudica* and *S. obtusifolia*, respectively. On the other hand, the apolar (HE) and intermediate polarity fractions (DC and EA) presented values ranging from 8 to 20% inhibition.

These results revealed that the fractions from *D. menstrualis* were more active than the crude extract of the red alga *P. brasiliense* [4], since the dichloromethane extract of the latter contains active components capable of inhibiting the germination of seeds of *M. pudica* and *S. obtusifolia* by merely 25 and 14%, respectively.

The effect of each fraction on radicle elongation is presented in Figure 1b. The results show that the ethanol/water fraction of *D. menstrualis* (HA) contains active components that inhibit by 75 and 68% the radicle elongation of *M. pudica* and *S. obtusifolia*, respectively.




■ *Mimosa pudica* ■ *Senna obtusifolia*

Figure 2: Effects of pachydictyol A+isopachydictyol A (in ppm) (a) on seed germination, (b) on radicle elongation, and (c) on hypocotyl elongation of two weeds. The data are expressed as percentage inhibition in relation to control (distilled water).

Finally, the effect of each fraction on hypocotyl development of the weeds is shown in Figure 1c. Again, the ethanol/water fraction was the most effective (35% for *M. pudica* and 30% for *S. obtusifolia*). In fact, it is important to note that the *D. menstrualis* fractions tested were more efficient against *M. pudica* in all experiments.

The inhibition values observed (Figure 1a, b and c) demonstrate that the ethanol/water fraction has a far more intense inhibitory effect on seed germination (88 and 77%) and radicle elongation (75 and 68%) than on hypocotyl elongation (35 and 30%, respectively).

Considering that the results reflect the inhibitory potential of chemical substances according to their polarities, it was possible to verify that the compounds present in HA are strongly active, whereas those in HE, DC and EA presented low inhibition rates. Thus, the results obtained qualify HA as a potential alternative source of chemical substances that might inspire more advanced studies regarding the production of agricultural bio defenses.



The chemical composition of *D. menstrualis* is characterized by cyclic diterpenes known as prenylated guaianes, α , β -unsaturated aldehydes [5-9, 11-13], and sulfated polysaccharides [10]. TLC analysis of the fractions and comparison with isolated products indicated that the diterpenes pachydictyol A (1) and isopachydictyol A (2) were the most abundant compounds in fraction HE. In the fractions DC and EA, the most abundant compound was the diterpene 6-hydroxy-dichotomano-2, 13-diene-16, 17-dial (3).

The effects of the unseparated diterpenes pachydictyol A and isopachydictyol A are presented in Figure 2. The mixture had a satisfactory inhibitory effect in dose-dependent concentrations (5-20 ppm) on the germination of seeds, ranging from 14-48% inhibition of *M. pudica* and from 6-28% of *S. obtusifolia*.

The results obtained indicate an effective inhibitory potential for the ethanol/water fraction of the alga. This inhibitory effect must be due to polar diterpenes and/or sulfated polysaccharides [18,19]. The other fractions (HE, DC and AC), however, presented low inhibitory potential compared with HA. Therefore, we may conclude that the ethanol/water fraction of *D. menstrualis* can become a promising alternative for the management of pastures.

Experimental

Biological material: *Dictyota menstrualis* (Dictyotaceae, Phaeophyceae) was collected at Atol das Rocas reef, Rio Grande do Norte State (lat. 03°51'03"S, long. 33°40' 29"W), Brazil, in February 2005. Atol das Rocas reef is a marine biological reserve in the northeast of Brazil, and is the only atoll in the South Atlantic. The collection was performed with federal authorization for collection, SISBIO/IBAMA number 17532, by scuba diving at a depth of 2-5 m. The algae were triaged in the place of collection and air-dried. They were then placed in thermic boxes to be transported to ALGAMAR laboratory where the identification was performed by one of us (RCV). Voucher specimens were deposited in the Herbarium of Universidade do Estado do Rio de Janeiro (HRJ). The receptor plants used in bioassays were two of the main weeds that occur in areas of cultivated pastures in the Amazon region, known as sensitive plant (*Mimosa pudica*) and mata-pasto (*Senna obtusifolia*). Seeds of these 2 plants were collected in the Municipality of Castanhal, Pará State, in the Brazilian Amazon region. The seeds were cleaned and purged in seed conservation chambers. Seed dormancy was overcome as established by earlier studies [2,3].

Extraction: The dried algae were powdered in an industrial blender, yielding 50 g of powder, which was extracted with acetone (100%) at room temperature for 10 days. The acetone extracts were

obtained by exhaustive extraction (using a decanting funnel as a container). The solvent was evaporated under reduced pressure, yielding a brownish residue (6 g). This was mixed with 40 g of silica gel. First, the mixture was stirred and filtered through filter paper with 800 mL *n*-hexane (HE, 92 mg). The second extraction was performed with 800 mL of dichloromethane (CH_2Cl_2), yielding 62 mg of extract (DC). The third extraction was made with 800 mL of ethyl acetate (EtOAc), yielding 130 mg after evaporation of the solvent (EA). Finally, the ethanol/water extract (HA) was prepared using 800 mL of ethyl alcohol/water (7:3). The ethanol was evaporated in a rotary evaporator; this was followed by lyophilization of the residue, yielding 240 mg (HA). All fractions were prepared at a 1% concentration for biological tests. The *n*-hexane fraction (HE) was submitted to silica gel chromatography (*n*-hexane/EtOAc), eluting with 30% EtOAc in *n*-hexane to afford 20 mg of unseparated diterpenes pachydictyol A and isopachydictyol A, in relative proportions 3:2. The NMR and IR spectra of the hydroalcoholic extracts presented characteristic signals of sulfated polysaccharides.

Bioassays: The bioassays on germination were developed in controlled conditions: constant temperature (25°C) and photoperiod of 12 h. Germination was monitored for 15 days, involving daily counting and elimination of germinated seeds, considered as those presenting radicle extension equal to or superior to 2 mm [2-3]. Each transparent Petri dish measured 9.0 cm in diameter and received 20 seeds. Regarding radicle and hypocotyl elongation, bioassays were performed under controlled conditions of constant temperature (25°C) and a photoperiod of 24 h in a germination chamber with cool white fluorescent lamps and a luminous flux of $10 \mu\text{mol.m}^{-2}\text{s}^{-1}$. Six pre-germinated seeds were placed in a transparent Petri dish for approximately 2-3 days. At the end of 10 days of growth, extension of the radicle and hypocotyl were measured. Each Petri dish was covered with a qualitative filter paper and received 3.0 mL of fraction, diterpene or an equal volume of distilled water (control). In the evaluation of fractions, a concentration was used of 1%, w/v, of each fraction and the volume of water replaced was 3.0 mL. In the evaluation of diterpenes, 4 concentrations were used, 5, 10, 15 and 20 ppm. Only after the beginning of each bioassay was either the fraction or diterpenes added. From this moment on, only distilled water was added when necessary.

Statistical analysis: The experimental design for all bioassays was entirely randomized and 3 repetitions were performed. The data were submitted to analysis of variance by the F test and the averages were compared by Tukey test ($P < 5\%$). All the analyses were performed in SAS program [20].

Acknowledgments - We are all thankful to CNPq and FAPERJ for financial support and productivity fellowships for VLT. DNC thanks CNPq for offering postdoc. RRF and FAOR thank CAPES for DSc fellowships. We are also grateful to EMBRAPA-Amazonia Oriental, represented by APSSE, and to SEDUC/PA for giving a professional development license to RRF.

References

- [1] Lobo LT, Castro KCF, Arruda MSP, Silva MN, Arruda AC, Muller AH, Arruda GMSP, Santos AS, Souza Filho APS. (2008) Potencial alelopático de catequinas de *Tachigali myrmecophylla* (Leguminosae). *Química Nova*, **31**, 493-497.
- [2] Souza Filho APS, Bayma JC, Guilhon GMSP, Zoghbi MGB. (2009) Atividade potencialmente alelopática do óleo essencial de *Ocimum americanum*. *Planta Daninha*, **27**, 499-505.
- [3] Souza Filho APS, Vasconcelos MAM, Zoghbi MGB, Cunha RL. (2009) Efeitos potencialmente alelopáticos dos óleos essenciais de *Piper hispidinervium* C. DC. e *Pogostemon heyneanus* Benth sobre plantas daninhas *Acta Amazonica*, **39**, 389-395.
- [4] Fonseca RR, Ortiz-Ramirez FA, Cavalcanti DN, Ramos CJB, Teixeira VL, Souza Filho APS. (2012) Allelopathic potential of extracts from marine macroalga *Plocamium brasiliense* and their effects on pasture weed. *Brazilian Journal of Pharmacognosy*, **22**, 850-853.

- [5] Teixeira VL, Cavalcanti DN, Pereira RC (2001) Chemotaxonomy study of the diterpenes from the brown alga *Dictyota menstrualis*. *Biochemical Systematics and Ecology*, **29**, 313-316.
- [6] Pereira HS, Leão-Ferreira LR, Moussatché N, Teixeira VL, Cavalcanti DN, Costa LJ, Diaz R, Frugulhetti ICPP. (2004) Antiviral activity of diterpenes isolated from the Brazilian marine alga *Dictyota menstrualis* against human immunodeficiency virus type 1 (HIV-1). *Antiviral Research*, **64**, 69-74.
- [7] Pereira HS, Leão-Ferreira LR, Moussatché N, Teixeira VL, Cavalcanti DN, Costa LJ, Diaz R, Frugulhetti ICPP. (2005) Effects of diterpenes isolated from the Brazilian marine alga *Dictyota menstrualis* on HIV-1 reverse transcriptase. *Planta Medica*, **71**, 1019-1024.
- [8] Cavalcanti DN, Rezende CM, Pinto AC, Teixeira VL. (2006) Diterpenoid constituents from the brown alga *Dictyota menstrualis* (Dictyotaceae, Phaeophyta). *Natural Product Communications*, **1**, 609-611.
- [9] Ortiz-Ramirez FA, Cavalcanti DN, Villaza RC, De-Paula JC, Yoneshigue-Valentin Y, Teixeira VL. (2008) Chemical variation in the diterpenes from the Brazilian brown alga *Dictyota menstrualis* (Dictyotaceae, Phaeophyceae). *Natural Product Communications*, **3**, 1879-1884.
- [10] Albuquerque IRL, Queiroz KCS, Alves LG, Santos EA, Leite EL, Rocha HAO (2004) Heterofucans from *Dictyota menstrualis* have anticoagulant activity. *Brazilian Journal of Medical and Biological Research*, **37**, 167-171.
- [11] Cronin G, Lindquist N, Hay ME, Fenical W. (1995) Effects of storage and extraction procedure on yields of lipophilic metabolites from the brown seaweed *Dictyota ciliolata* and *D. menstrualis*. *Marine Ecology Progress Series*, **119**, 265-273.
- [12] Taylor RB, Lindquist N, Kubanek J, Hay ME. (2003) Intraspecific variation in palatability and defensive chemistry of brown seaweeds: Effects on herbivore fitness. *Oecologia*, **136**, 412-423.
- [13] Blount JF, Dunlop RW, Erickson KL, Wells RJ. (1982) Two diterpenes with new carbocyclic ring. Systems from an Australian collection of the brown alga *Dictyota dichotoma*. *Australian Journal of Chemistry*, **35**, 145-163.
- [14] Pereira RC, Cavalcanti DN, Teixeira VL. (2000) Effects of secondary metabolites from the tropical Brazilian brown alga *Dictyota menstrualis* on the amphipod *Parhyale hawaiensis*. *Marine Ecology Progress Series*, **205**, 95-100.
- [15] Chou SC, Krishna V, Chou CH. (2009) Hydrophobic metabolites from *Rhododendron formosanum* and their allelopathic activities. *Natural Product Communications*, **4**, 1189-1192.
- [16] Junttila O. (1976) Seed and embryo germination in *Syringa vulgaris* and *S. reflexa* as affected by temperature during seed development. *Physiologia Plantarum*, **29**, 264-268.
- [17] Duran JM, Tortosa ME. (1985) The effect of mechanical and chemical scarification on germination of charlock (*Sinapis arvensis* L.) seeds. *Seed Science and Technology*, **13**, 155-163.
- [18] Shalaby EA. (2011) Algae as promising organisms for environment and health. *Plant Signal Behavior*, **6**, 1338-1350.
- [19] Renjun W, Hui X, Peiyu Z, Liand Q, Hengjang C, Xuexi T. (2006) Allelopathic effects of *Ulva pertusa*, *Corallina pilufera* and *Sargassum thunbergii* on the growth of the dinoflagellates *Heterosigma akashiwo* and *Alexandrium tamarense*. *Journal of Applied Phycology*, **19**, 109-121.
- [20] SAS Statistical Analysis System (1989) *User's guide. Version 6.12*, SAS Inst Inc Cary, North Caroline, 846 p.

Isolation of the Plant Hormone (+)-Absciscic acid as an Antimycobacterial Constituent of the Medicinal Plant Endophyte *Nigrospora* sp.

Trevor N. Clark^a, Katelyn Ellsworth^a, Haoxin Li^a, John A. Johnson^a and Christopher A. Gray^{a,b,*}.

^aDepartment of Biology, University of New Brunswick, 100 Tucker Park Rd, Saint John, NB, Canada, E2L 4L5.

^bDepartment of Chemistry, University of New Brunswick, 100 Tucker Park Rd, Saint John, NB, Canada, E2L 4L5.

cgray@unb.ca

Received: September 12th, 2013; Accepted: October 8th, 2013

An extract of the endophytic fungus *Nigrospora* sp. (isolate TC2-054) from the Canadian medicinal plant *Fragaria virginiana* exhibited significant antimycobacterial activity against *Mycobacterium tuberculosis* H37Ra. Bioassay guided fractionation revealed that linoleic acid derivatives and the plant hormone (+)-absciscic acid (ABA) were responsible for the observed antimycobacterial activity. This activity of ABA has not been previously reported.

Keywords: Absciscic acid, Antimycobacterial activity, Endophyte, *Fragaria virginiana*, Canadian medicinal plant, *Nigrospora* sp.

Endophytic fungi are increasingly being recognized as an important source of bioactive natural products [1] and the biosynthetic potential of endophytes isolated from medicinal plants has recently been highlighted [1f]. As part of our research on Canadian medicinal plants, we have so far isolated 81 endophytic fungi from a selection of twelve plants that have been used therapeutically by the First Nations peoples of the Canadian Maritimes [2]. Antimicrobial screening of this endophyte library indicated that an extract of the spent fermentation broth of *Nigrospora* sp. isolate TC2-054 obtained from the medicinal plant *Fragaria virginiana* exhibited significant antimycobacterial activity and, therefore, warranted further investigation.

The EtOAc extract of a bench-scale (2 L), two-week fermentation of TC2-054 was subjected to a modified Kupchan partition protocol with the *n*-hexane and CH₂Cl₂ fractions showing significant activity towards *M. tuberculosis* H37Ra. NMR analysis of the *n*-hexane fraction showed it to contain significant amounts of linoleic acid derivatives that are known to exhibit antimycobacterial activity [3] and was not further fractionated. However, normal phase HPLC of the CH₂Cl₂ fraction led to the isolation of (+)-absciscic acid [4] (ABA; 6.5 mg/L; Figure 1) as the only antimycobacterial constituent [*M. tuberculosis* H37Ra; MIC: 200 µg/mL; IC₅₀ (± SD): 40.8 ± 3.7 µg/mL]. LC-MS screening did not detect ABA in a *F. virginiana* extract prepared from leaves collected at the same time as those used for the isolation of the *Nigrospora* sp.




Figure 1: (+)-Absciscic acid (ABA).

The role of ABA as a plant stress hormone has been well documented since its isolation in 1961 [4a]. More recently, it has been shown to be involved in a wide range of developmental physiological processes in higher plants [5a] and found also to be produced by lower plants, lichens and fungi [5b], including a

species of *Nigrospora* isolated from a Thai sea-fan [5c]. Many of these fungi are recognized endophytes and phytopathogens [5b, d] and the production of ABA by these fungi may be a factor in the development and maintenance of plant-fungus relationships [5b]. ABA has recently been shown to be an important factor in fruit ripening in the strawberry, *Fragaria ananassa* [5e], posing intriguing questions relating to the role of endophyte derived ABA in *F. virginiana*.

The antimycobacterial activity of ABA has not been previously described, and although it exhibited only moderate activity against *M. tuberculosis* H37Ra, it did not show any observable toxicity to mammalian cells; no difference in viability was observed between HEK293 cells treated with absciscic acid (at concentrations up to 250 µg/mL) and the corresponding vehicle controls [analysis of 250 µg/mL data: t-test, $F = 0.052$, $P = 0.83$, 4 d.f. and Mann-Whitney, $U = 6.0$, $P = 0.70$; mean cell viability (± SD) $93 \pm 10\%$ compared with negative control]. Given its low toxicity to human cells, further work should focus on defining the mode of action of ABA against mycobacteria as it may reveal a selective cellular target that could be exploited in future tuberculosis drug development.

Experimental

Isolation and identification of *Nigrospora* sp. (TC2-054):

Endophytes were isolated from the leaves of *Fragaria virginiana* Duchesne collected from the wood-lot on the UNB campus in Saint John, New Brunswick, Canada (N 45° 18.375' W 66° 05.616') in August 2010. Plants were identified by Dr Stephen Clayden of the New Brunswick Museum and a voucher specimen has been deposited in the New Brunswick Museum Herbarium (Number: NBM VP-37478). Leaf surfaces were sterilized by immersion in 5.25% aqueous sodium hypochlorite for 5 sec, sterile distilled water for 10 sec and 70% EtOH for 15 sec. The sterile tissue was rinsed with sterile distilled water, blotted dry on an autoclaved paper towel, and aseptically cut into pieces that were placed onto 2% malt extract agar and incubated at ambient room temperature. Endophytic fungi were subcultured on 2% malt extract agar until pure cultures were obtained. Isolate TC2-054 was identified taxonomically as *Nigrospora* sp. through examination of colony and

spore morphology, with the taxonomic classification being confirmed by comparison of the internal transcribed spacer and 5.8S rRNA gene (ITS) DNA regions with corresponding sequences available in the GenBank database (NCBI, US Government), as previously described. The sequence data derived from isolate TC2-054 has been submitted to GenBank and assigned accession number KC916673.

Biological assays: Anti-mycobacterial activity against *M. tuberculosis* strain H37Ra (ATCC 25177) and cytotoxicity against HEK 293 was evaluated as previously described [6].

Fermentation and extraction: TC2-054 was fermented in 2% malt extract broth at room temperature for 2 weeks (2 L; 20 × 100 mL batches in 250 mL Erlenmeyer flasks). The fungal material was separated from the broth using vacuum filtration, and the broth was extracted with EtOAc (3 × 300 mL). The organic fraction was concentrated *in vacuo* to give a crude extract (79 mg).

Bioassay guided fractionation: The crude extract (79 mg) was dissolved in 9:1 MeOH/H₂O (50 mL) and extracted with hexanes (3 × 25 mL) before being diluted with H₂O (25 mL) and extracted with CH₂Cl₂ (3 × 25 mL). The aqueous fraction was concentrated *in vacuo*, dissolved in H₂O (50 mL) and extracted with EtOAc (3 × 25 mL) and *n*-BuOH (3 × 25 mL). The 5 partition fractions were concentrated *in vacuo* with the CH₂Cl₂ fraction (46 mg) showing strong bioactivity. The CH₂Cl₂ fraction (46 mg) was subjected to normal phase HPLC (1:1, hexanes/EtOAc) to give **1** (13 mg)

Extraction and LC-MS screening of *Fragaria virginiana*: Freeze-dried ground plant material (20.5 g) was extracted using a Soxhlet apparatus for 11.5 h with MeOH (200 mL). The crude extract was

then concentrated *in vacuo* to give a green oil (6.54 g). The crude extract (0.5 mg/mL in MeOH, 20 µL injections) was subjected to LC/MS using a Phenomenex Kinetex C18 column (50 × 4.60 mm, 2.6 µm) eluted with a 0.1% formic acid H₂O/MeOH gradient (1 min at 95:5 H₂O/MeOH, 7 min linear transition to 1:99 H₂O:MeOH for a further 7 min) at 500 µL/min on an Agilent 1100 LC system coupled to an AB Sciex API 2000 triple quadrupole MS (TurboIonSpray source run in positive mode). The presence of ABA was analyzed by selected reaction monitoring of the pseudo-molecular ion (*m/z* 265) to dehydrated product ion (*m/z* 247) transition. Under these conditions, an ABA standard eluted with a retention time of 6.6 min and was detectable in MeOH solutions at a concentration of ≥ 25 ng/mL. No peak corresponding to ABA was observed in the *F. virginiana* crude extract. Subsequent co-injections of the crude extract and ABA standard confirmed that ABA would be detectable at ≥ 100 ng/g in the crude extract.

Acknowledgments - The authors would like to thank Stephen Clayden (New Brunswick Museum), Christopher Martyniuk (University of New Brunswick), Gilles Robichaud (Université de Moncton), Larry Calhoun (University of New Brunswick), and Fabrice Berrue and Patricia Boland (University of Prince Edward Island) for their assistance with plant identification, isolation and amplification of DNA, HEK293 cytotoxicity assays, recording NMR spectra, and obtaining ESIMS data, respectively. Financial support for this research was provided by NSERC (Discovery Grant to CAG and CGS-M to TNC), the NBHRF (Seed Operating Grant to CAG and JAJ), the NBIF (Research Assistantship Initiative grants to CAG), and UNB (University Research Fund grants to CAG and JAJ) and is gratefully acknowledged.

References

- [1] (a) Strobel G, Daisy B, Castillo U, Harper J. (2004) Natural products from endophytic microorganisms. *Journal of Natural Products*, **67**, 257-268; (b) Guo B, Wang Y, Sun X, Tang K. (2008) Bioactive natural products from endophytes: A review. *Applied Biochemistry and Microbiology*, **44**, 136-142; (c) Verma VC, Kharwar RN, Strobel GA. (2009) Chemical and functional diversity of natural products from plant associated endophytic fungi. *Natural Product Communications*, **4**, 1511-1532; (d) Aly AH, Debbab A, Kjer J, Proksch P. (2010) Fungal endophytes from higher plants: a prolific source of phytochemicals and other bioactive natural products. *Fungal Diversity*, **41**, 1-16; (e) Aly AH, Debbab A, Proksch P. (2011) Fungal endophytes: unique plant inhabitants with great promises. *Applied Microbiology and Biotechnology*, **90**, 1829-1845; (f) Kaul S, Gupta S, Ahmed M, Dhar M. (2012) Endophytic fungi from medicinal plants: a treasure hunt for bioactive metabolites. *Phytochemistry Reviews*, **11**, 487-505.
- [2] Ellsworth K, O'Neill T, Webster D, Johnson JA, Gray CA. (2012) Bioactivity of endophytes from traditionally used medicinal plants in New Brunswick, Canada. *Planta Medica*, **78**, 1157.
- [3] Kanetsuna F. (1985) Bactericidal effect of fatty acids on mycobacteria, with particular reference to the suggested mechanism of intracellular killing. *Microbiology and Immunology*, **29**, 127-141.
- [4] (a) Addicott FT, Lyon JL, Ohkuma K, Thiessen WE, Carns HR, Smith OE, Cornfort JW, Milborro BV, Ryback G, Wareing PF. (1968) Absciscic acid - a new name for abscisin 2 (dormin). *Science*, **159**, 1493; (b) Constantino MG, Donate PM, Petagnani N. (1986) An efficient synthesis of (±)-abscisic acid. *Journal of Organic Chemistry*, **51**, 253-254; (c) Assante G, Merlini L, Nasini G. (1977) (+)-Absciscic acid, a metabolite of fungus *Cercospora rosicola*. *Experientia*, **33**, 1556-1557.
- [5] (a) Wasilewska A, Vlad F, Sirichandra C, Redko Y, Jammes F, Valon C, Frey NFD, Leung J. (2008) An update on abscisic acid signaling in plants and more. *Molecular Plant*, **1**, 198-217; (b) Hartung W. (2010) The evolution of abscisic acid (ABA) and ABA function in lower plants, fungi and lichen. *Functional Plant Biology*, **37**, 806-812; (c) Trisuwan K, Rukachaisirikul V, Sukpondma Y, Preedanon S, Phongpaichit S, Rungjindamai N, Sakayaroj J. (2008) Epoxydons and a pyrone from the marine-derived fungus *Nigrospora* sp PSU-F5. *Journal of Natural Products*, **71**, 1323-1326; (d) Tudzynski B, Sharon A. (2002) *Biosynthesis, biological role and application of fungal phytohormones*. In *The Mycota: A comprehensive treatise on fungi as experimental systems for basic and applied research. Industrial applications*, Springer-Verlag, New York, 183-211; (e) Jia HF, Chai YM, Li CL, Lu D, Luo JJ, Qin L, Shen YY. (2011) Absciscic acid plays an important role in the regulation of strawberry fruit ripening. *Plant Physiology*, **157**, 188-199.
- [6] (a) Carpenter CD, O'Neill T, Picot N, Johnson JA, Robichaud GA, Webster D, Gray CA. (2012) Anti-mycobacterial natural products from the Canadian medicinal plant *Juniperus communis*. *Journal of Ethnopharmacology*, **143**, 695-700; (b) O'Neill TE, Webster D, Johnson JA, Gray CA. (2013) The Canadian medicinal plant *Heracleum maximum* contains antimycobacterial diynes and furanocoumarins. *Journal of Ethnopharmacology*, **147**, 232-237.

New Cembranoid Diterpene from the South China Sea Soft Coral
Sarcophyton sp.

Fei Cao, Jing Zhou, Kai-Xia Xu, Meng-Qi Zhang and Chang-Yun Wang*

Key Laboratory of Marine Drugs, Ministry of Education, School of Medicine and Pharmacy,
Ocean University of China, Qingdao 266003, China

changyun@ouc.edu.cn

Received: May 8th, 2013; Accepted: September 3rd, 2013

One new oxygenated cembranoid diterpene, named sarcophytol W (**1**), along with six known analogues (**2–7**) were isolated from a soft coral *Sarcophyton* sp. collected from the South China Sea. Their structures were elucidated by spectroscopic analyses and by comparison with previously reported data. The absolute configuration of **1** was assigned on the basis of the absolute configuration of the related congener **2**, which was determined by application of the modified Mosher's method. All the compounds were evaluated for their antibacterial and antifouling activities.

Keywords: Soft coral, *Sarcophyton* sp., Cembrane diterpenoid, Absolute configuration, Bioactivity.

Diterpenoids from marine organisms possess a wide array of unusual structures. Among them, cembrane diterpenoids with the 14-carbon cyclic nucleus have been frequently found in soft corals [1]. These cembrane diterpenoids possess various biological activities such as ichthyotoxic [2], antifouling [3], antitumor [4], neuroprotective [5], antiinflammatory [6] and cytotoxic activities [7].

Recently, in the course of our investigation of new bioactive substances from soft corals collected from the South China Sea [8–13], a soft coral *Sarcophyton* sp. attracted our attention because the EtOAc portion showed lethal activity toward brine shrimps, *Artemia salina*. Chemical investigation of the active extract led to the isolation of seven cembrane diterpenoids, named sarcophytol W (**1**), (2*E*,7*E*)-4,11-dihydroxy-1,12-oxidocembra-2,7-dien (**2**) [14], (–)-marasol (**3**) [15], (+)-11,12-epoxy-11,12-dihydrocembra-C (**4**) [16], (+)-11,12-epoxysarcophytol A (**5**) [17], sarcophytol B (**6**) [18], and sarcolactone A (**7**) [3]. Their structures were elucidated by NMR spectroscopic methods and comparison with previously reported data in the literature. Among these isolated compounds, **1** is a new cembrane diterpenoid possessing 3,14-epoxy and 11,12-epoxy groups (Figure 1).

Sarcophytol W (**1**) was obtained as a colorless oil. Its molecular formula was determined as C₂₀H₃₂O₃ on the basis of positive HRESIMS (*m/z* 321.2422 [M + H]⁺; calc. 321.2430), with five degrees of unsaturation, demonstrating the presence of an additional oxygen with respect to (–)-marasol (**3**) [15]. Careful comparison of the ¹H and ¹³C NMR data of **1** with those of **3** showed a close structural relationship between them. The main differences were the presence of the typical signals for a trisubstituted epoxy functionality (δ_H 2.57, 1H, dd, *J* = 2.4, 6.6 Hz; δ_C 66.6, CH and 60.8, C) in **1**, instead of a double bond (δ_H 5.11, 1H, br d, *J* = 9.2 Hz, H-11; δ_C 131.2, CH, C-11 and 129.8, C, C-12) in **3**, suggesting **1** is an 11,12-epoxy derivative of **3**. Analysis of 2D NMR experiments and in particular of HMBC correlations from H₃-20 to C-11, C-12, and C-13, and ¹H-¹H COSY correlations (Figure 1) confirmed the proposed structure and allowed us to assign all carbon and proton resonances (Table 1).






Figure 1: ¹H-¹H COSY and key HMBC correlations for compound **1** and structures of compounds **1–7**.

The relative configuration of **1** was assigned on the basis of 1D NOE measurements and 2D NOESY experiments. In the NOESY spectrum, correlations between H-3/H-14, H-3/H-11, H-14/H₃-20 and H-11/H₃-18 revealed the β orientation of H-3, H-11, H-14, H₃-18, and H₃-20. Also, in the selective NOE experiments, the result of irradiation of H-3, H-11, H-14, H₃-18 and H₃-20 was compatible with the 2D NOESY experiments (Figure 2).

The absolute configuration of **1** was assigned on the basis of a shared biogenesis with **2**. The relative and absolute configurations of **2** were determined by 1D NOE experiments and the modified Mosher's method. In the selective 1D NOE experiments with **2**, irradiation of H₃-17 resulted in the enhancement of H₃-18 and H₃-20, and irradiation of H-11 resulted in the enhancement of H₃-20, indicating that they should be on the same face of the

Table 1: ^1H and ^{13}C NMR spectroscopic data (CDCl_3) of **1**.

position	δ_{C}	δ_{H} J in Hz
1	151.8	—
2	118.1	5.43 (1H, d, 1.8)
3	87.5	4.54 (1H, m)
4	74.6	—
5	40.6	1.93 (Ha, ddd, 14.4, 10.8, 1.8), 1.60 (Hb, ddd, 14.4, 7.8, 1.8)
6	21.7	2.18 (Ha, m), 2.13 (Hb, m)
7	126.0	5.35 (1H, t, 6.0)
8	132.6	—
9	36.5	2.20 (2H, m)
10	25.0	1.76 (2H, m)
11	66.4	2.58 (1H, dd, 6.6, 2.4)
12	60.6	—
13	45.5	2.18 (Ha, dd, 13.6, 4.2), 1.01 (Hb, d, 13.6)
14	82.8	4.80 (1H, m)
15	26.2	2.17 (1H, m)
16	22.8	1.03 (3H, d, 6.6)
17	21.2	1.12 (3H, d, 6.6)
18	22.8	1.00 (3H, s)
19	16.6	1.59 (3H, s)
20	16.5	1.34 (3H, s)

**Figure 2:** Selected NOE correlations for compounds **1** and **2**.**Figure 3:** Values of $\Delta\delta_{\text{H}(S,R)}$ (measured in CDCl_3) of the MTPA esters of compound **2**.

molecule (Figure 2). Thus, the relative configurations of all asymmetric carbons of **2** were assigned as β -faces of H_3 -18, H-11, H_3 -20 and the isopropyl group. The absolute configuration of **2** was established by the modified Mosher's method [19]. Treatment of **2** with (*R*)-(+)- α - and (*S*)-(–)- α -methoxy- α -(trifluoromethyl)phenylacetyl chloride (MTPA-Cl) gave the corresponding (*S*)- and (*R*)-MTPA esters, **2s** and **2r**, respectively. The ^1H NMR signals of the two MTPA esters were assigned on the basis of their ^1H - ^1H COSY spectra. The $\Delta\delta_{\text{H}(S-R)}$ values were then calculated (Figure 3). Following the literature [19], the results indicated that the absolute configuration of C-11 was *S*. Therefore, the absolute configuration of **2** was assigned as 1*S*,4*S*,11*S*,12*S*. Correspondingly, the absolute configuration of **1** could also be inferred according to a shared biogenesis as 3*S*,4*S*,11*S*,12*S*,14*R*. Compound **1** was thus identified as (1*Z*,3*S*,4*S*,7*E*,11*S*,12*S*,14*R*)-3,14-epoxy-11,12-epoxy-4-hydroxycembra-1,7-diene.

In this study, six known cembranoid diterpenoids (**2**–**7**) were isolated from the *Sarcophyton* sp.; these had been obtained from soft corals of the family Alcyoniidae in previous reports [16,18,20]. Interestingly, (–)-marasol (**3**) was isolated from *S. infundibuliforme* and *S. glaucum* [13, 20], whereas the enantiomer of **3**, (+)-marasol, was found in the gorgonian *Plexaura flexuosa* of the Plexauridae family [21]. More interestingly, their biosynthetic precursor, sarcophytol A, was found explicitly in both families. These results could partly support the conclusion that the families Alcyoniidae and Plexauridae are closely related taxonomically, which was inferred from our previous chemical investigations [13].

The antibacterial activity of **1**–**7** was evaluated against a panel of pathogenic bacteria *in vitro*. Compound **6** showed antibacterial activity against *Bacillus cereus*, *Staphylococcus albus* and *Vibrio parahaemolyticus* with MIC values of 3.13, 1.56, and 0.50 μM , respectively. The antifouling activity of all the isolated compounds was also evaluated, but only **3** exhibited an antifouling effect on larval settlement of the barnacle *Balanus amphitrite* at a concentration of 10.0 $\mu\text{g/mL}$.

In conclusion, seven oxygenated cembranoid diterpenes (**1**–**7**) were obtained from the soft coral *Sarcophyton* sp., of which **1** is a new compound possessing 3,14-epoxy and 11,12-epoxy groups. The absolute configuration of **1** was assigned on the basis of the absolute configuration of the related congener **2**, which was determined by application of the modified Mosher's method. In addition, the results reflected part of the chemotaxonomic significance among Alcyoniidae and Plexauridae.

Experimental

General: Optical rotations were measured on a JASCO P-1020 digital polarimeter at room temperature. UV spectra were recorded on a UV-2501PC spectrophotometer, and IR spectra on a Nicolet Nexus 470 spectrometer. ^1H and ^{13}C NMR spectra were obtained using a JEOL Eclips-600 (600 and 150 MHz) spectrometer. ESI-MS and HR-ESI-MS were recorded on a Q-TOF Ultima Global GAA076 LC mass spectrometer. HPLC separation was performed with a Waters 1525 semi-preparative HPLC system coupled with a Waters 2996 photodiode array detector. A Kromasil C_{18} preparative HPLC column (250 \times 10 mm, 5 μm) was used. All solvents were of analytical grade (Shanghai Chemical Plant, Shanghai, People's Republic of China). Silica gel (200–300 mesh; Qingdao Marine Chemical Group Co., Qingdao, People's Republic of China), octadecylsilyl (ODS) silica gel (45–60 μm ; Merck KGaA, Darmstadt, Germany), and Sephadex LH-20 (Amersham Biosciences Inc., Piscataway, NJ, USA) were used for column chromatography. Precoated silica gel GF254 plates (Yantai Zifu Chemical Group Co., Yantai, People's Republic of China) were used for analytical TLC analyses.

Biological material: The sample of *Sarcophyton* sp. was collected by scuba diving in Xuwen Coral Reef Area, Guangdong Province, China, and was subsequently identified by Prof. Hui Huang of South China Sea Institute of Oceanology, Chinese Academy of Sciences. A voucher specimen (XW-18) was deposited at the Key Laboratory of Marine Drugs, Ministry of Education, Ocean University of China, Qingdao, China.

Extraction and isolation: *Sarcophyton* sp. (950.0 g, wet weight) was homogenized, followed by extraction with ethanol (95%, v/v) and $\text{MeOH}/\text{CHCl}_3$ (1:1), successively. The organic layer was filtered and concentrated under reduced pressure to give a residue (50.6 g), which was partitioned between EtOAc and H_2O (EtOAc: H_2O = 2:1), 6 times. The EtOAc extract (8.4 g) was first applied to silica gel CC eluting with light petroleum containing increasing amounts of EtOAc to yield 6 fractions (Fr.1 – Fr.6). Fr.2 was separated by silica gel CC (light petroleum: EtOAc = 98:2), followed by Sephadex LH-20 CC (light petroleum: CHCl_3 : MeOH = 2:1:1) to obtain **3** (38.6 mg) and **4** (1.20 g). Fr.3 was separated by silica gel CC (light petroleum: EtOAc = 15:1) to offer Fr.3-1 and Fr.3-2. Fr.3-1 was further separated by Sephadex LH-20 CC (light petroleum: CHCl_3 : MeOH = 2:1:1) to yield **1** (40.0 mg). Fr.3-2 was purified by semi-preparative HPLC (MeOH : H_2O = 85:15) to obtain **5** (38.1 mg). Fr.4 was subjected to repeated chromatography over Sephadex LH-20 to give **6** (25.2 mg). Fr.5 was separated by silica gel CC (light petroleum: EtOAc = 4:1), followed by Sephadex

LH-20 CC (light petroleum: CHCl_3 : MeOH = 2:1:1), and further purified by semi-preparative HPLC (MeOH : H_2O = 80:20) to yield **7** (93.5 mg). Fr.6 was first subjected to silica gel CC, then purified by semi-preparative HPLC to give **2** (16.4 mg).

Sarcophytol W (1)

Colorless oil.

$[\alpha]_D^{20}$: +69 (c 0.41, CHCl_3).

UV λ_{max} (MeOH) nm: 206 (3.01).

IR (KBr): 3046, 2933, 1453, 1366, 1029, 971, 943 cm^{-1} .

^1H NMR and ^{13}C NMR: Table 1.

HRESIMS: m/z $[\text{M} + \text{H}]^+$, calcd for $\text{C}_{20}\text{H}_{33}\text{O}_3$: 321.2430; found: 321.2422.

Preparation of the (S)- and (R)-MTPA ester derivatives of 2: To a stirred solution of **2** (1.5 mg) in pyridine (500 μL) was added 4-(dimethylamino)pyridine (2 mg) and (R)-(+)- α -methoxy- α -(trifluoromethyl)phenylacetyl chloride (MTPA-Cl, 10 μL). The mixture was stirred at room temperature for 12 h. The reaction mixture was then passed through a disposable pipet packed with silica gel and eluted with light petroleum and EtOAc (3:1) to give the (R)-Mosher ester **2s**. Treatment of **2** (1.5 mg) with (S)-MTPA-Cl (10 μL) as described above yielded the corresponding (R)-Mosher ester **2r**.

2-(S)-MTPA ester (2s)

^1H NMR (600 MHz CDCl_3): 5.89 (1H, d, J = 15.6 Hz, H-3), 5.58 (1H, d, J = 15.6 Hz, H-2), 5.47 (1H, d, J = 2.4, 10.2 Hz, H-11), 5.22 (1H, d, J = 9.0 Hz, H-7), 3.51 (3H, s, OCH_3 -MTPA), 2.17 (1H, m, H-9a), 2.14 (1H, m, H-13a), 2.05 (1H, m, H-14a), 1.83 (1H, m, H-14b), 1.80 (1H, m, H-13b), 1.76 (1H, m, H-9b), 1.67 (3H, s, H-19), 1.66 (2H, m, H-10), 1.65 (1H, m, H-15), 1.31 (3H, s, H-18), 1.06 (3H, s, H-20), 0.84 (3H, d, J = 6.6 Hz, H-17), 0.81 (3H, d, J = 6.6 Hz, H-16).

ESIMS: m/z 561 $[\text{M} + \text{Na}]^+$.

2-(R)-MTPA ester (2r)

^1H NMR (600 MHz, CDCl_3): 5.89 (1H, d, J = 15.6 Hz, H-3), 5.58 (1H, d, J = 15.6 Hz, H-2), 5.48 (1H, d, J = 2.4, 10.2 Hz, H-11), 5.23 (1H, d, J = 9.0 Hz, H-7), 3.53 (3H, s, OCH_3 -MTPA), 2.18 (1H, m, H-9a), 2.10 (1H, m, H-13a), 2.02 (1H, m, H-14a), 1.83 (1H, m, H-9b), 1.81 (1H, m, H-14b), 1.78 (1H, m, H-13b), 1.74 (2H, m, H-10), 1.68 (3H, s, H-19), 1.63 (1H, m, H-15), 1.30 (3H, s, H-18), 1.02 (3H, s, H-20), 0.83 (3H, d, J = 6.6 Hz, H-17), 0.80 (3H, d, J = 6.6 Hz, H-16).

ESIMS: m/z 561 $[\text{M} + \text{Na}]^+$.

Antibacterial activity assay: Antibacterial activity was evaluated by the conventional broth dilution assay [22]. Eight bacterial strains, *Bacillus cereus*, *Tetragenococcus halophilus*, *Staphylococcus albus*, *S. aureus*, *Escherichia coli*, *Pseudomonas putida*, *Nocardia brasiliensis*, and *Vibrio parahaemolyticus* were used, and ciprofloxacin was employed as a positive control.

Antifouling bioassay: Antifouling activity of the compounds on the larval settlement of the barnacle *Balanus amphitrite* was tested according to literature procedures [23].

Supplementary data: ^1H , ^{13}C , ^1H - ^1H COSY, HMQC, HMBC, 1D NOE, 2D NOESY and mass spectra of **1**; 1D NOE spectra of **2**; ^1H and mass spectra of the (R)- and (S)-MTPA esters of **2**; ^1H and ^{13}C data of **3** are also available.

Acknowledgments - We thank Dr Chang-Lun Shao, School of Medicine and Pharmacy, Ocean University of China, for sample collection, and Prof. Pei-Yuan Qian and Dr Ying Xu, Division of Life Science, the Hong Kong University of Science and Technology, China, for antifouling bioassay. This work was supported by the National Natural Science Foundation of China (Nos. 41130858; 40976077), the National High Technology Research and Development Program of China (863 Program) (No. 2013AA093001), and the Natural Science Foundation of Shandong Province (No. ZR2011HM085).

References

- [1] Blunt JW, Copp BR, Keyzers RA, Munro MHG, Prinsep MR. (2012) Marine natural products. *Natural Product Reports*, **29**, 144–222.
- [2] Iwagawa T, Nakamura S, Masuda T, Okamura H, Nakatani M, Siro M. (1995) Irregular cembranoids containing a 13-membered carbocyclic skeleton isolated from a soft coral, *Sarcophyton* species. *Tetrahedron*, **51**, 5291–5298.
- [3] Sun XP, Wang CY, Shao CL, Li L, Li XB, Chen M, Qian PY. (2010) Chemical constituents of the soft coral *Sarcophyton infundibuliforme* from the South China Sea. *Natural Product Communications*, **5**, 1171–1174.
- [4] Yokomatsu H, Satake K, Hiura A, Tsutsumi M, Suganuma M. (1994) Sarcophytol A: a new chemotherapeutic and chemopreventive agent for pancreatic cancer. *Pancreas*, **9**, 526–530.
- [5] Badria FA, Guirguis AN, Perovic S, Steffen R, Müller WEG, Schööder HC. (1998) Sarcophytolide: a new neuroprotective compound from the soft coral *Sarcophyton glaucum*. *Toxicology*, **131**, 133–143.
- [6] Cheng SY, Wen ZH, Wang SK, Chiou SF, Hsu CH, Dai CF, Chiang MY, Duh CY. (2009) Unprecedented hemiketal cembranolides with anti-inflammatory activity from the soft coral *Lobophytum durum*. *Journal of Natural Products*, **72**, 152–155.
- [7] Shaker KH, Müller M, Ghani MA, Dahse HM, Seifert K. (2010) Terpenes from the soft corals *Litophyton arboretum* and *Sarcophyton ehrenbergi*. *Chemistry & Biodiversity*, **7**, 2007–2015.
- [8] Han L, Wang CY, Huang H, Shao CL, Liu QA, Qi J, Sun XP, Zhai P, Gu YC. (2010) A new pregnane analogue from Hainan soft coral *Scleronephthya gracillimum* Küekenthal. *Biochemical Systematics and Ecology*, **38**, 243–246.
- [9] Li R, Shao CL, Qi X, Li XB, Li J, Sun LL, Wang CY. (2012) Polyoxygenated sterols from the South China Sea soft coral *Sinularia* sp. *Marine Drugs*, **10**, 1422–1432.
- [10] Li L, Sheng L, Wang CY, Zhou YB, Huang H, Li XB, Li J, Molloy E, Gavagnin M, Guo YW. (2011) Diterpenes from the Hainan soft coral *Lobophytum cristatum* Tixier-Durivault. *Journal of Natural Products*, **74**, 2089–2094.
- [11] Li L, Wang CY, Shao CL, Guo YW, Li GQ, Sun XP, Han L, Huang H, Guan HS. (2009) Sarcoglycosides A–C, new O-glycosylglycerol derivatives from the South China Sea soft coral *Sarcophyton infundibuliforme*. *Helvetica Chimica Acta*, **92**, 1495–1502.
- [12] Li L, Wang CY, Shao CL, Han L, Sun XP, Zhao J, Guo YW, Huang H, Guan HS. (2009) Two new metabolites from the Hainan soft coral *Sarcophyton crassocaule*. *Journal of Asian Natural Products Research*, **11**, 851–855.
- [13] Wang CY, Chen AN, Shao CL, Li L, Xu Y, Qian PY. (2011) Chemical constituents of soft coral *Sarcophyton infundibuliforme* from the South China Sea. *Biochemical Systematics and Ecology*, **39**, 853–856.
- [14] Rao CB, Satyanarayana C, Rao DS, Rao DV, Fahy E, Faulkner DJ. (1993) Metabolites of the soft coral *Sinularia ovispiculata* from the Indian Ocean. *Journal of Natural Products*, **56**, 2003–2007.
- [15] Kobayashi M, Nakano E. (1990) Stereochemical course of the transannular cyclization, in chloroform, of epoxycebranoids derived from the geometrical isomers of (14S)-14-hydroxy-1,3,7,11-cebranetraens. *The Journal of Organic Chemistry*, **55**, 1947–1951.

- [16] Bowden BF, Coll JC, Hicks W, Kazlauskas R, Mitchell SJ. (1978) Studies of Australian soft corals. X. The isolation of epoxyisoneocembrene-A from *Sinularia grayi* and isoneocembrene-A from *Sarcophyton ehrenbergi*. *Australian Journal of Chemistry*, **31**, 2707–2712.
- [17] Bowden BF, Coll JC, Tapiolas DM. (1983) Studies of Australian soft corals. XXXIII. New cembranoid diterpenes from a *Lobophytum* species. *Australian Journal of Chemistry*, **36**, 2289–2295.
- [18] Kobayashi M, Nakagawa T, Mitsuhashi H. (1979) Marine terpenes and terpenoids. I. Structures of four cembrane-type diterpenes; sarcophytol-A, sarcophytol-A acetate sarcophytol-B, and sarcophytonin-A, from the soft coral *Sarcophyton glaucum*. *Chemical & Pharmaceutical Bulletin*, **27**, 2382–2387.
- [19] Ohtani I, Kusumi T, Kashman Y, Kakisawa H. (1991) High-field FT NMR application of Mosher's method. The absolute configurations of marine terpenoids. *Journal of the American Chemical Society*, **113**, 4092–4096.
- [20] Kobayashi M, Iesaka T, Nakano E. (1989) Marine terpenes and terpenoids. IX. Structures of six new cembranoids, sarcophytols F, K, P, Q, R and S, from the soft coral *Sarcophyton glaucum*. *Chemical & Pharmaceutical Bulletin*, **37**, 2053–2057.
- [21] Peniston M, Rodriguez AD. (1991) The isolation of (–)-sarcophytola and (+)-marasol from the Caribbean gorgonian *Plexaura flexuosa*. *Journal of Natural Products*, **54**, 1009–1016.
- [22] Appendino G, Gibbons S, Giana A, Pagani A, Grassi G, Stavri M, Smith E, Rahman MM. (2008) Antibacterial cannabinoids from *Cannabis sativa*: a structure-activity study. *Journal of Natural Products*, **71**, 1427–1430.
- [23] Shao CL, Wu HX, Wang CY, Liu QA, Xu Y, Wei MY, Qian PY, Gu YC, Zheng CJ, She ZG, Lin YC. (2011) Potent antifouling resorcylic acid lactones from the gorgonian-derived fungus *Cochliobolus lunatus*. *Journal of Natural Products*, **74**, 629–633.

Crotofolane Diterpenoids from *Croton caracasanus*

Katiuska Chávez^a, Reinaldo S. Compagnone^b, Ricarda Riina^c, Alexander Briceño^d, Teresa González^d, Emilio Squitieri^b, Carlos Landaeta^b, Humberto Soscún^d and Alirica I. Suárez^{a,*}

^aFacultad de Farmacia, Universidad Central de Venezuela, Caracas, Venezuela

^bEscuela de Química, Facultad de Ciencias, Universidad Central de Venezuela, Caracas, Venezuela

^cReal Jardín Botánico, RJB-CSIC, Plaza Murillo 2, Madrid 28014, Spain

^dCentro de Química, Instituto Venezolano de Investigaciones Científicas (IVIC), Caracas, Venezuela

alirica.suarez@ciens.ucv.ve

Received: August 16th, 2013; Accepted: September 9th, 2013

Four new crotofolane-type diterpenoids, crotoctarasin (A-D) (1-4), together with the known crotofolin E, were isolated from a dichloromethane extract of the stems of *Croton caracasanus* Pittier. The structures of the new compounds were determined by spectroscopic methods, and the structure of **3** was further confirmed by single-crystal X-ray data analyses.

Keywords: Diterpenoids, Crotofolanes, *Croton caracasanus* Pittier, Euphorbiaceae.

The genus *Croton* (Euphorbiaceae), one of the largest genera of flowering plants [1], is a rich source of terpenoids. A great variety of structurally diverse diterpenoids such as abietanes [2], neoclerodanes [3] labdanes [4], and *ent*-kauranes [5] are frequently found in *Croton* species. The presence of cembranoids [6] and nor-crotofolane-type diterpenoids [7] has also been reported. The unusual structure of one crotofolane diterpenoid was reported for the first time from *C. corylifolius* [8]; the compound was named crotofolin A (**6**). Furthermore, a series of compounds, crotofolins B, C, D and E, were identified from the same species [9] and from *C. dichogamus* [10]. More recently, new crotofolanes were reported from *C. cascarilloides* [11]. As part of our continuing interest in the chemistry of *Croton* species occurring in Venezuela [12-15], we now report the isolation and structural elucidation of four new diterpenoids with a crotofolane skeleton, isolated from stems of *C. caracasanus* Pittier, a species closely related to *C. corylifolius*. The *in vitro* cytotoxicity of the isolated metabolites against PC-3, HeLa, and MCF-7 tumor cell lines was also assessed. From a chemotaxonomic point of view, it is of interest to note that these crotofolane diterpenoids have been isolated only from related species of *Croton*.

Crotoctarasin A (**1**) was isolated as a colorless sticky solid, $[\alpha]_D^{25}$ -33.3 (*c* 1.00, CHCl₃). Its molecular formula, C₂₀H₂₂O₄, was established by HREIMS as 326.1500, and its ¹³C NMR spectrum indicated a structure with 10 indices of hydrogen deficiency. The IR spectrum of **1** revealed absorption bands for an α,β -unsaturated γ -lactone (1753 cm⁻¹), α,β -unsaturated carbonyl (1663 and 1630 cm⁻¹) and terminal olefin (901 cm⁻¹) functionalities. The ¹H and ¹³C NMR spectra (Tables 1 and 2) showed signals due to three different methyl groups, at δ_H 1.18 (d, *J* = 7.7 Hz, H-19), at δ_H 1.20 (s, H-20) and, a vinylic one at δ_H 1.93 (s, H-17). A terminal olefinic methylene group at δ_H 4.46 and 4.94 (s, H-18), and one allylic proton at δ_H 3.14 (s, H-5) were also evident. ¹³C NMR and DEPT data (Table 2) indicated the presence of three methyl, four methylene and five methine carbons, one of them oxygenated, at δ_C 82.3 ppm (C-9). The crotofolane skeleton was proposed on the basis of the mentioned features and the COSY, HMQC, and HMBC (Figure 2) correlations, and also by comparison of spectroscopic data





Figure 1: Structures of compounds 1-6.

of compound **1** with those of crotofolin E (**5**), isolated many years ago from *C. corylifolius* and also isolated in this work (Figure 1).

The differences between the NMR spectra of **1** and **5** could be explained by the absence of the hydroxyl group. Instead, the presence of a broad singlet integrating for one proton at δ_H 5.00 (brs, H-9) identified another oxo-allylic proton on C-9, which was corroborated by the difference in the chemical shifts of the carbon in this position (δ_C 82.3 vs 108.7) (Figure 1). NOESY NMR experiments were performed for crotoctarasin A (**1**); NOE correlations between the methyl group (H-20) and the oxo-allylic proton (H-5) were observed. The relationship between H-13 and

Table 1: ^{13}C NMR spectral data at 67.5 MHz [CDCl_3 (**1-4**) and CD_3OD (**5**)].

Position	1	2	3	4	5
			δ_{C}		
1	207.6	207.3	208.0	208.8	209.6
2	40.9	41.1	40.8	40.6	40.9
3	36.4	36.2	36.5	37.4	36.0
4	165.8	166.2	165.7	166.2	167.9
5	54.6	55.0	55.0	56.4	54.5
6	59.4	59.8	60.0	61.2	59.8
7	52.7	53.3	52.8	41.3	53.0
8	161.9	141.6	159.5	162.1	160.2
9	82.3	148.8	110.0	108.7	108.7
10	38.3	113.4	36.9	36.3	41.5
11	35.9	39.0	34.7	28.5	34.5
12	145.8	146.3	146.2	137.7	147.0
13	40.8	40.6	40.2	44.6	39.9
14	144.9	147.1	145.2	145.7	144.4
15	128.5	130.2	129.7	123.4	129.4
16	173.1	169.4	170.9	171.4	171.6
17	10.2	11.3	10.0	11.0	8.6
18	112.8	113.0	112.9	110.1	111.6
19	17.3	17.4	17.4	17.5	16.2
20	19.7	19.4	20.1	22.0	18.9
21	----	----	60.0	59.0	----
22	----	----	15.5	15.3	----

**Figure 2:** HMBC correlations for crotofolin A (**1**).

H-9 was evident, indicating that both of them are on the same face of the structure. Due to this, the stereochemistry of the previous isolated crotofolanes was established on the basis of X-ray studies and similarities of the NMR data; here we propose that crotofolin A (**1**) has a similar relative configuration to crotofolin E (**5**).

Crotofolin B (**2**) was isolated as a yellow sticky solid with $[\alpha]_{\text{D}}^{25} -56.6$ (c 1.00, CHCl_3). The ^{13}C NMR and mass spectrometric data of **2** indicated that it has the molecular formula $\text{C}_{20}\text{H}_{20}\text{O}_4$, thus accounting for eleven elements of unsaturation. The IR spectrum showed bands at 1735, 1710 and 1458 cm^{-1} . In the ^{13}C NMR spectrum of **4** (Table 2), a trisubstituted olefin was deduced from the signals at δ_{C} 148.8 (C-9) and δ_{C} 113.4 (C-10), additional to the terminal olefin also found in the previously described compounds. The relationship of each proton and carbon was established from HMQC and HMBC spectra. Close examination of the NMR data (Tables 1 and 2) shows that **2** has the same framework as **1** and **5**, and differs only at C-9 and C-10 (Figure 1). An olefin between these two carbons, confirmed by the strong correlations observed in the HMBC experiment [δ_{H} 5.97 (H-10) with δ_{C} 148.8 (C-9), δ_{C} 141.6 (C-8) and δ_{C} 39.0 (C-11)] agrees with the proposed structure of **2**, which could be obtained biogenetically by dehydration of crotofolin E (**5**), and is named by us as crotofolin B.

Crotofolin C (**3**) was isolated as white crystals, mp $236\text{--}238^\circ\text{C}$, $[\alpha]_{\text{D}}^{25} -78.4$ (c 1.00, CHCl_3), and gave a molecular formula of $\text{C}_{22}\text{H}_{26}\text{O}_5$, as determined by HREIMS. Comparison of the ^1H and ^{13}C NMR (Tables 1 and 2) with those of **5**, suggested a similar structure, but with additional signals. The presence of a multiplet at δ_{H} 3.80 (m, H-21), integrating for two protons, and a new methyl signal at δ_{H} 1.19 (t, $J = 7.2\text{ Hz}$, H-22), showed the presence of an ethoxy group in this compound. ^{13}C NMR and DEPT spectra showed 22 carbon atoms with 26 attached protons. The signals that identify the principal subunits found in **1** and **5** were also visualized. Close examination of 2D NMR, HMQC, and HMBC data indicated that the crotofolane skeleton was the same as in **5**, and that the ethoxy group found in **3** was attached to C-9. In the NOESY experiment on crotofolin C (**3**), the NOE correlations more evident were between methyl H-20 and the oxo-allylic H-5. None was observed for protons H-7 and H-13, and with the same criteria used above for crotofolin A, the relative stereochemistry of the chiral centers was proposed for crotofolin C (**3**). In order to verify the stereochemical assignment made by NMR experiments, the structure of **3** was determined by single crystal X-ray analysis (Figure 3). The molecular structure of **3** is shown in Figure 1 with the respective labels.

Table 2: ^1H NMR spectral data at 270 MHz [CDCl_3 (**1-4**) and CD_3OD (**5**)].

Position	1	2	3	4	5
		δ_{H} (mult, J in Hz)			
1	---	---	---	---	---
2	2.47 (m)	2.48 (m)	2.48 (m)	2.48 (m)	2.48 (m)
3	2.44 (m), 2.88 (m)	2.43 (m)	2.43 (m), 2.93 (2dd, 6.4; 3.7)	2.54 (m), 2.99 (m)	2.41 (m), 3.04 (m)
4	---	---	---	---	---
5	3.14 (s)	3.16 (s)	3.14 (s)	3.40 (s)	3.30 (s)
6	---	---	---	---	---
7	2.83 (brs)	2.80 (d, 11.6)	2.77 (brs)	3.18 (brs)	2.74 (d, 11.6)
8	---	---	---	---	---
9	5.00 (brs)	---	---	---	---
10	1.23 (m), 2.61 (d, 3.7)	5.97 (dd, 6.4; 3.5)	1.34 (dt, 14.0; 4.9), 2.81 (brt, 2.7)	2.04 (m), 2.34 (m)	1.48 (dt, 12.9; 5.7), 2.58 (m)
11	2.23 (dt, 12.6; 3.7), 2.64 (m)	3.18 (m)	2.20 (dt, 13.4; 4.2), 2.42 (m)	2.39 (m), 2.72 (m)	2.53 (m)
12	---	---	---	---	---
13	3.01 (m)	3.21 (m)	3.48 (td, 11.6; 3.5)	3.15 (s)	3.59 (td, 11.6; 3.5)
14	---	---	---	---	---
15	---	---	---	---	---
16	---	---	---	---	---
17	1.93 (s)	2.08 (s)	1.89 (s)	1.83 (s)	1.85 (s)
18	4.46 (s), 4.94 (s)	4.53 (s), 5.04 (s)	4.44 (s), 4.92 (s)	4.30 (s), 4.76 (s)	4.44 (s), 4.90 (s)
19	1.18 (d, 7.7)	1.20 (d, 7.4)	1.17 (d, 7.4)	1.22 (d, 7.3)	1.17 (d, 7.7)
20	1.20 (s)	1.23 (s)	1.25 (s)	1.46 (s)	1.29 (s)
21	---	---	3.80 (m)	3.55 (m)	---
22	---	---	1.19 (t, 7.2)	1.22 (t, 6.9)	---




Figure 3: ORTEP drawing of compound **3**. No absolute configuration was determined (Hydrogen atoms were omitted for clarity).

Crotoctarasin D (**4**) was isolated as a colorless oil, $[\alpha]_D^{25} -48.8$ (c 1.00, CHCl_3), and gave the same molecular formula $\text{C}_{22}\text{H}_{26}\text{O}_5$ as **3**, deduced from HREIMS and ^{13}C NMR. The ^1H NMR spectrum resembled compound **3**, but with the relevant difference of one methyl signal integrating for 6 protons, which was identified at δ_{H} 1.22 (d, $J = 7.3$ Hz, H-19 and t, $J = 6.9$ Hz, H-22); the correlations between carbons and protons at this chemical shift indicated that two methyl groups are involved with this signal; the carbon at δ_{C} 17.5 (C-19) correlates with the doublet for H-19, and the signal at δ_{C} 15.3 (C-22) is correlated with the triplet for H-22, respectively. The other two methyl groups present in **3** were found in **4**, also as two singlets at δ_{H} 1.83 (s, H-17) and δ_{H} 1.46 (s, H-20). The differences in ^1H and ^{13}C NMR spectra between **3** and **4** (see Tables 1 and 2) involved mainly the seven membered ring and the gamma lactone. These differences could be explained by possible conformational changes in these rings. The NOESY NMR experiment of crotoctarasin D (**4**) did not show any correlation between H-7/H-13. GIAO/CAM-B3LYP/6-31+G(d,p) calculations were performed on the three possible diastereomers to compare theoretical and experimental ^1H and ^{13}C NMR spectra, and thus predict the configuration of this new crotofolane, but were not conclusive (results not shown). Different experiments are necessary to propose the relative stereochemistry in the diastereomer of **3** named as crotoctarasin D (**4**). An attempt to obtain a crystal for this compound was unsuccessful.

Crotoctarasins A-D (**1-4**) were tested for cytotoxic activity against human tumor cell lines PC-3, HeLa, and MCF-7, using the MTT method [16]. Adriamycin was used as a positive control. IC_{50} values > 50 μM were obtained, which indicate that these crotofolanes were not cytotoxic.

Experimental

General experimental procedures: Melting points were measured on a Kofler hot-stage melting point instrument and are uncorrected. Specific rotations were acquired with an ATAGO Polax-2L polarimeter. IR spectra were recorded on a Shidmazu 470 spectrophotometer, EIMS on a Varian Saturn 2000, and HREIMS with a Finnigan Trace mass spectrometer. NMR spectra were measured on a JEOL 270 MHz, and Bruker AMX-500. Chemical shifts are given in ppm referenced to the residual non-deuterated solvent signal (CHCl_3 7.24 and 77.0 ppm). CC was performed using Si gel (70-230 mesh) from Scharlau. TLC analysis was carried out using plastic precoated plates (Merck, Si gel plates GF₂₅₄, 0.2 mm) and the compounds were visualized using either a UV lamp $\lambda = 254$ nm or by spraying with *p*-anisaldehyde. All solvents used were of analytical grade.

Plant material: Aerial parts of *C. caracasanus* Pittier were collected around Caracas, Venezuela, in August 2009 and identified by Dr Ricarda Riina. A voucher specimen (VEN 382364) has been deposited at the Herbario Nacional de Venezuela.

Extraction and isolation: The air-dried and powdered stems (213 g) of *C. caracasanus* were extracted by maceration with MeOH (2 x 2L) at room temperature for 15 days, and the combined extracts were concentrated *in vacuo*. The residue was dissolved in a mixture of MeOH: H₂O (1:1), and then partitioned in turn with *n*-hexane, dichloromethane, and EtOAc, to afford *n*-hexane (1.38 g), CH_2Cl_2 (0.99 g), EtOAc (0.70 g) and a residual hydro-methanolic fraction (2.75 g). CC of the dichloromethane fraction over Si gel, using a CH_2Cl_2 : EtOAc (90:10) mixture with increasing polarity afforded fractions A-M. Fraction D furnished crotoctarasin B (**2**) (55.0 mg); fraction E was subjected to chromatography with Si-gel eluting with *n*-hexane:EtOAc:MeOH (65:30:5) to yield crotoctarasin C (**3**) (29.3 mg). Crotoctarasin D (**4**) (23.1 mg) was obtained pure after purification of fraction J by CC on Si gel using a step gradient of CH_2Cl_2 :*n*-hexane (0-5%) as eluent. In the same manner, fraction G gave crotofolin E (**5**) (51.5 mg). Preparative TLC (CHCl_3 : EtOAc, 88:12) of fraction I afforded crotoctarasin A (**1**) (19.2 mg).

Crotoctarasin A (1)

Colorless sticky solid.

$[\alpha]_D^{25}$: -33.3 (c 1.00, CHCl_3).

IR ν_{max} film: 3473, 1753, 1704, 1663, 1630, 1447, 901 cm^{-1}

^1H NMR and ^{13}C NMR: Table 2 and 1.

HREIMS: m/z 326.1510 $[\text{M}]^+$; calcd for $\text{C}_{20}\text{H}_{22}\text{O}_4$: 326.1512.

Crotoctarasin B (2)

Yellow sticky solid.

$[\alpha]_D^{25}$: -56.6 (c 1.00, CHCl_3).

IR ν_{max} film: 3233, 1735, 1710, 1651, 1632, 1458, 902 cm^{-1} .

^1H NMR and ^{13}C NMR: Table 2 and 1.

HREIMS: m/z 324.1360 $[\text{M}]^+$; calcd for $\text{C}_{20}\text{H}_{20}\text{O}_4$: 324.1356.

Crotoctarasin C (3)

White crystals.

MP: 236-238°C.

$[\alpha]_D^{25}$: -78.4 (c 1.00, CHCl_3).

IR ν_{max} film: 3491, 1758, 1707, 1670, 1650, 1630, 1441, 940 cm^{-1} .

^1H NMR and ^{13}C NMR: Table 2 and 1.

HREIMS: m/z 370.1775 $[\text{M}]^+$; calcd for $\text{C}_{22}\text{H}_{26}\text{O}_5$: 370.1773.

Crotoctarasin D (4)

Colorless oil.

$[\alpha]_D^{25}$: -48.8 (c 1.00, CHCl_3).

IR ν_{max} film: 3464, 1763, 1705, 1597, 1433, 1120, 938, 910 cm^{-1} .

^1H NMR and ^{13}C NMR: Table 2 and 1.

HREIMS m/z : 370.1771 $[\text{M}]^+$; calcd for $\text{C}_{22}\text{H}_{26}\text{O}_5$: 370.1773.

Crotofolin E (5)

White crystals.

MP: 207-209°C.

$[\alpha]_D^{25}$: -76.4 (c 1.00, CHCl_3).

IR ν_{max} film: 3465, 1737, 1695, 1651, 1628, 1456, 952, 902 cm^{-1} .

^1H NMR and ^{13}C NMR: Table 2 and 1.

HREIMS: m/z 342.1459 $[\text{M}]^+$; calcd for $\text{C}_{20}\text{H}_{22}\text{O}_5$: 342.1461.

Note: several NMR assignments were different from those reported in the reference and here are reassigned according to 2D NMR data, which was not available in the original publication [9].

Crystal data of crotoctarasin C (3): Compound **3** was crystallized from CHCl_3 solution by slow solvent evaporation. A colorless block crystal was selected for crystallographic measurements. Intensity data were recorded at room temperature on a Rigaku AFC-7S diffractometer equipped with a Mercury CCD detector using monochromatic $\text{Mo}(\text{K}\alpha)$ radiation ($\lambda = 0.71070$ Å). Experimental details on unit cell and intensity measurements can be found in the

CIF file deposited at the Cambridge Crystallographic Data Centre with CCDC number 867457. The structure was solved by Direct Methods and refined by full-matrix least-squares on F^2 . The H-atoms on C were placed in calculated positions using a riding atom model with fixed C-H distances [0.93 Å for C(sp²), 0.96 Å for C(sp³, CH₃), and 0.97 Å for C(sp³, CH₂)]. All the H atoms were refined with isotropic displacement parameters set to $1.2 \times U_{eq}$ for C(sp²) and 1.5 for C(sp³) of the attached atom. All the refinement calculations were made using SHELXTL-NT [16]. Molecular formula = C₂₂H₂₆O₅, molecular mass = 370.4393, crystal system = orthorhombic, space group = $P2_12_12_1$, unit cell dimensions: $a = 10.478$ (3) Å, $b = 12.250$ (3) Å, $c = 15.590$ (3) Å; $V = 2001.0$ (9) Å³, 4; $\rho = 1.230$ Mg/m³; $\mu = 0.086$ mm⁻¹, $F(000) = 792$, $\lambda(\text{Mo K}\alpha)$. Theta range for data collection = 4.20° to 55.40° ; limiting

indices = $-12 \leq h \leq 12$, $-11 \leq k \leq 14$, $-18 \leq l \leq 18$; collected reflections 23828; independent reflections 3272 [$R(\text{int}) = 0.0360$]; Data/Restraints/Parameters = 3948/0/245; Goodness-of-fit on $F^2 = 1.096$; Final R indices [$I > 2\sigma(I)$]; R indices (all data) $R_1 = 0.0546$, $wR_2 = 0.1006$; Largest diff. Peak and hole 0.137 and -0.139 eÅ⁻³.

Supplementary data: Spectral data, theoretical calculations and cytotoxicity values are available.

Acknowledgments - Financial support for this work was provided by Consejo de Desarrollo Científico de la Universidad Central de Venezuela. Grant PG O6.7342.2008 and by FONACIT Grant PEII-705.

References

- [1] Riina R, van Ee B, Wiedenhoeft AC, Cardozo A, Berry PB. (2010) Sectional rearrangement of arborescent clades of *Croton* (Euphorbiaceae) in South America: evolution of arillate seeds and a new species *Croton domatifer*. *Taxon*, **59**, 1147-1160.
- [2] Santos HS, Barros WA, Albuquerque MRJR, Bandeira PN, Pessoa C, Braz-Filho R, Monte FJQ, Leal-Cardoso JH, Lemos TLG. (2009) Cytotoxic diterpenoids from *Croton argyrophyllodes*. *Journal of Natural Products*, **58**, 1884-1887.
- [3] Vigor C, Fabre N, Fourasté I, Moulis C. (2002) Neoclerodane diterpenoids from *Croton eluteria*. *Journal of Natural Products*, **65**, 1180-1182.
- [4] Roengsumran S, Petsom A, Kuptiyanuwat M, Vilaivan T, Ngamrojanavanich N, Chaichanipyuth C, Phuthong S. (2001) Cytotoxic labdane diterpenoids from *Croton oblongifolius*. *Phytochemistry*, **56**, 103-107.
- [5] Yang X-D, Chen W, Zhao J-F, Yang L-J, Zhang H-B, Li L. (2009) Ent-kaurane diterpenes and phenolic compounds from *Croton kongensis* (Euphorbiaceae). *Biochemical Systematic and Ecology*, **37**, 23-240.
- [6] Mulholland DA, Langat MK, Crouch NR, Coley HM, Mutambi EM, Nuzillard JM. (2010) Cembranolides from the stem bark of the southern African medicinal plant, *Croton gratissimus* (Euphorbiaceae). *Phytochemistry*, **71**, 1381-1386.
- [7] Kawakami S, Matsunami K, Otsuka H, Shinzato T, Takeda Y, Kawahata M, Yamaguchi K. (2010) A crotofolane-type diterpenoid and a rearranged nor-crotofolane-type diterpenoid with a new skeleton from the stems of *Croton cascarilloides*. *Tetrahedron Letters*, **51**, 4320-4322.
- [8] Chan WR, Prince EC, Manchand PS, Springer JP, Clardy J. (1975) The structure of Crotofolin A, a diterpene with a new skeleton. *Journal of the American Chemical Society*, **97**, 4437-4439.
- [9] Burke BA, Chan WR, Pascoe KO, Blount JF, Manchand PS. (1979) The structure of Crotofolin E, a novel tricyclic diterpene from *Croton corylifolius*. *Tetrahedron Letters*, **36**, 3345-3348.
- [10] Jogia MK, Andersen RJ, Párkani L, Clardy J, Dublin HT, Sinclair ARE. (1989) Crotofolane diterpenoids from the African shrub *Croton dichogamus* Pax. *Journal of Organic Chemistry*, **54**, 1654-1657.
- [11] Kawakami S, Toyoda H, Harinantenaina L, Matsunami K, Otsuka H, Shinzato T, Takeda Y, Kawahata M, Yamaguchi K. (2013) Eight new diterpenoids and two new nor-diterpenoids from the stems of *Croton cascarilloides*. *Chemical & Pharmaceutical Bulletin*, **61**, 411-418.
- [12] Suárez AI, Blanco Z, Delle Monache F, Compagnone RS. (2004) Three new glutarimide alkaloids from *Croton cuneatus*. *Natural Products Research*, **18**, 421-426.
- [13] Suárez AI, Chávez K, Delle Monache F, Vásquez L, Delannoy DM, Orsini G, Compagnone RS. (2008) New 3,4-Seco-ent-kaurenes from *Croton caracasana* flowers. *Natural Product Communications*, **3**, 319-322.
- [14] Suárez AI, Blanco Z, Compagnone RS, Salazar-Bookaman MM, Zapata V, Alvarado C. (2006) Anti-inflammatory activity of *Croton cuneatus* aqueous extract. *Journal of Ethnopharmacology*, **105**, 99-101.
- [15] Morales A, Alvarez A, Arvelo F, Suárez AI, Compagnone RS, Galindo-Castro I. (2011) The natural diterpene ent-16 β -17 α -dihydroxykaurane down-regulates Bcl-2 by disruption of the Ap-2 α /Rb transcription activating complex and induces E2F1 up-regulation in MCF-7 cells. *Apoptosis*, **16**, 1245-1252.
- [16] Mosmann, T. (1983) Rapid colorimetric assay for cellular growth and survival. Application to proliferation and cytotoxicity assays. *Journal of Immunology Methods*, **65**, 55-63.

Development and Validation of a Modified Ultrasound-Assisted Extraction Method and a HPLC Method for the Quantitative Determination of Two Triterpenic Acids in *Hedyotis diffusa*

Yu-Chiao Yang^a, Ming-Chi Wei^{b,*}, Hui-Fen Chiu^a and Ting-Chia Huang^c

^aDepartment and Graduate Institute of Pharmacology, Kaohsiung Medical University, Kaohsiung 80708, Taiwan

^bDepartment of Environmental Engineering & Science, Chia Nan University of Pharmacy and Science, Tainan 71710, Taiwan

^cDepartment of Chemical Engineering, National Cheng Kung University, Tainan 70101, Taiwan

s120702@mail.chna.edu.tw; ycyang@kmu.edu.tw

Received: October 26th, 2012; Accepted: September 20th, 2013

In the present study, the oleanolic acid (OA) and ursolic acid (UA) contents of *Hedyotis diffusa* and *H. corymbosa* were determined by a rapid, selective and accurate method combining modified ultrasound-assisted extraction (MUAE) and HPLC. Compared with traditional extraction methods, MUAE reduced the extraction time, the extraction temperature and the solvent consumption and maximized the extraction yields of OA and UA. Furthermore, the combined MUAE–HPLC method was applied to quantitate OA and UA in plant samples and exhibited good repeatability, reproducibility and stability. The mean recovery studies (one extraction cycle) for OA and UA were between 91.3 and 91.7% with RSD values less than 4.5%. The pioneer method was further applied to quantitate OA and UA in six samples of *H. diffusa* and five samples of *H. corymbosa*. The results showed that the OA and UA content in the samples from different sources were significantly different. This report is valuable for the application of *H. diffusa* and *H. corymbosa* obtained from different regions in clinical research and pharmacology.

Keywords: *Hedyotis diffusa*, *Hedyotis corymbosa*, Oleanolic acid, Ursolic acid, Modified ultrasound-assisted extraction.

The dried whole plants of *Hedyotis diffusa* Willd. (synonym *Oldenlandia diffusa* Willd., family Rubiaceae) are called “Peh-Hue-Juwa-Chi-Cao” in Chinese. The extracts of *H. diffusa* possess various pharmacological properties, such as anti-oxidative, anti-angiogenic, antimutagenic, neuroprotective, anti-inflammatory, hepatoprotective and anticancer activities [1,2]. Up to now, four major classes of compounds, triterpenoids, flavonoids, anthraquinones and steroids, have been reported as bioactive compounds from this herb [1,3]. Among these components, triterpenic acids such as oleanolic acid (OA) and ursolic acid (UA) have been extensively studied for their pharmacological properties due to their high abundance in medicinal herbs. Both have minimal toxicity and have been shown to display numerous biological properties with therapeutic potential [4]. In recent years, studies have shown that UA and OA induce apoptosis in a wide variety of cancer cells, including hepatocellular carcinoma [5], prostate carcinoma [6], colorectal cancer [7], acute myelogenous leukemia [8], skin tumorigenesis [9], cervical carcinoma [10] and lung carcinoma [11].

Several publications have reported the analysis of OA and UA in other raw materials using HPLC methods [12–14]. However, the simultaneous determination of OA and UA in plant extracts is a difficult task due to their structural similarity [15]. In this paper, a simple, rapid and accurate HPLC method coupled with a photo diode array (PDA) detector is described for the analysis of OA and UA from *H. diffusa*. The optimal conditions for the analytical method were also investigated for the best resolution and highest sensitivity of detection. In addition, *H. corymbosa* (L.) Lam has also been used in Taiwan for treating the same conditions as *H. diffusa*. To compare the efficacy of the two herbs, the OA and UA contents of *H. diffusa* and *H. corymbosa* were also analyzed.

Recently, traditional solvent [16], ultrasonic-assisted [12,13,17,18] and supercritical carbon dioxide [13] extraction techniques have been employed in the extraction of OA and UA from different materials. In the case of ultrasonic-assisted extraction (UAE), the principal advantage is that it is the simplest and most economical technique and is easy to scale up to industrial level. Several probable mechanisms for the ultrasonic enhancement of extraction, such as cell disruption, improved penetration and enhanced swelling and hydration processes, have been cited as leading to enhanced extraction [19–20].

In this study, we focussed primarily on the development of an efficient technique for the extraction and determination of OA and UA contents in *H. diffusa* and *H. corymbosa*. The novel method consists of heat-reflux extraction (HRE) (5–10 min), followed by UAE, which is the first time that this novel method has been investigated. The effects of processing parameters were further evaluated. The optimized conditions giving the highest OA and UA yields were recorded and used to compare the results obtained with those found using conventional UAE and HRE. Moreover, a simple, rapid and accurate HPLC method was developed and optimized for the analysis of OA and UA in extracts of *H. diffusa* and *H. corymbosa*.

Because various parameters potentially affect the UAE process, the optimization of the experimental conditions represents a critical step in the development of a UAE method. In this study, UAE parameters such as the type of solvent, solvent consumption, extraction time, extraction temperature, duty cycle, liquid/solid ratio, material particle size and ethanol-water mixture, were optimized. However, the extraction yields of OA and UA (shown in Table 1) still did not match those of the extracts prepared by HRE

(one extraction cycle). Therefore, MUAE in the presence of two stages of extraction was employed in order to enhance the extraction efficiency. Moreover, the conventional HRE of OA and UA from *H. diffusa* was also optimized and compared with MUAE and UAE. The optimized conditions and extraction yields of OA and UA obtained by MUAE, UAE and HRE are listed in Table 1 (one extraction cycle). The application of MUAE significantly reduced the extraction time, extraction temperature and solvent consumption, and achieved superior OA and UA yields. This was attributed to the cavitation effects of ultrasound disruption of plant cell walls and facilitating the release of the target constituents. This release enhanced the mass transport of the solvent from the continuous phase into the plant cells. These results clearly demonstrated that the developed MUAE method provided a good alternative for the extraction of OA and UA from *H. diffusa*.

Table 1: Comparison of optimum extraction parameters for HRE, UAE and MUAE methods.

Extraction parameters	Extraction method		
	HRE	UAE	MUAE
Herbal sample	HD1	HD1	HD1
Particle size (mm)	0.2–0.5	0.2–0.5	0.2–0.5
Plant weight (g)	5	5	5
Frequency (kHz)	–	40	40
Duty cycle (%)	–	79	79
Stirring rate (rpm)	200	–	200
Time (min)	60	40	40
Temperature (°C)	Boiling point	55	55
Liquid/solid ratio (mL/g)	14	12	12
Extraction cycle	1	1	1
OA			
Yield (mg/g) ^a	0.59 ± 0.03	0.56 ± 0.03	0.60 ± 0.02
Ethanol (%) ^b	70	60	60
RSD (%) ^c	4.42	5.35	3.82
UA			
Yield (mg/g) ^a	3.28 ± 0.14	3.06 ± 0.14	3.37 ± 0.13
Ethanol (%) ^b	90	80	80
RSD (%) ^c	4.20	4.57	3.94

^aValues are written as the mean (mg/g) ± standard deviation (SD) of six replications. The amounts are expressed as mg of the target compound per g of the plant on a dry weight basis. ^bEthanol concentration in water (v/v). ^cRSD (%) = (SD/mean) × 100.

The optimization of the HPLC conditions for the analysis of triterpenic acids in *H. diffusa* was achieved by varying the mobile phase composition, flow rate and column temperature. The chromatograms of standard and ultrasonically extracted *H. diffusa* are compared. OA and UA were satisfactorily separated under the optimum conditions described above, with retention times of 50.44 ± 0.06 and 56.62 ± 0.08 min, respectively. No interference peaks from the endogenous constituents of the extract of *H. diffusa* were found in the region of the investigated compounds. Furthermore, the two compounds of interest were unequivocally confirmed by the retention times and UV spectra of the authentic standards. Therefore, this HPLC system was simple, easy to use, and effective for the identification and quantification of OA and UA in *H. diffusa*.

The calibration curve was constructed using a linear regression of the theoretical concentration of an analyte versus the corrected peak area. The mean regression equations and their correlation coefficients were calculated to be $Y = 1.8083 \times 10^6 X + 7.8839 \times 10^4$ and $R^2 = 0.997$ for OA, and $Y = 1.5469 \times 10^6 X - 1.6539 \times 10^4$ and $R^2 = 0.998$ for UA. The analytical procedure was also sensitive with respect to the limits of detection (LOD) and limits of quantification (LOQ) for OA and UA (0.056 and 0.094 µg/mL, respectively, for the LOD and 0.18 and 0.31 µg/mL, respectively, for the LOQ).

The precision and accuracy of the developed method was evaluated by measuring intra- and inter-day variability in terms of the relative

standard deviation (RSD) and the relative error (RE). The standard solutions, at four different concentration levels, were analyzed at least six times within the same day, and the RSD and RE values obtained were less than 5.0 and 4.0% for OA and less than 4.5 and 4.0% for UA. Similarly, to measure the inter-day variability, the same concentration of the two standards was run over at least six consecutive days, and the values were 6.5 and 5.7% for OA and 6.3 and 5.9% for UA. Therefore, the criteria for the precision and accuracy for analyzing the OA and UA samples were fulfilled in the developed analytical method.

Table 2: Repeatability and reproducibility of the developed MUAE–HPLC method and stability of the sample.

Analytes	Repeatability ^a		Reproducibility ^b		Stability (RE (%)) ^c	
	Mean (mg/g)	RSD (%) ^d	Mean (mg/g)	RSD (%) ^d	24 h	48 h
OA	0.60	3.82	0.60	5.98	–1.07	1.46
UA	3.37	3.94	3.21	6.63	–0.98	–1.12

^aSix samples of HD1 from the same source were extracted and analyzed under the optimum conditions during one day. ^bThe extraction yields obtained from six independent extractions (HD1) performed on five consecutive days. ^cRE (%) = [(mean observed concentration after 24 h (or 48 h) – initial mean concentration)/initial mean concentration] × 100. ^dRSD (%) = (SD/mean) × 100.

To study the repeatability of the developed MUAE–HPLC method, six herbal samples (HD1) from the same source were extracted under the optimized conditions and analyzed using the established method. Table 2 (one extraction cycle) reveals that the RSD of the OA and UA content in six replicate herbal samples was 3.82 and 3.94%, respectively, indicating that the developed method had good repeatability. Furthermore, the reproducibility was evaluated by calculating the extraction yields obtained from six independent extractions performed on five consecutive days. The obtained RSD for OA and UA were less than 6.7%, which were within the acceptance criteria and indicated that the combined MUAE–HPLC method was accurate, reliable, and reproducible.

The same sample solution was stored at 25°C and analyzed every 12 h over 48 h to evaluate the stability of the samples. The results showed that the relative error (RE) of OA and UA were 1.07 and 0.98%, respectively, after 24 h and 1.46 and 1.12%, respectively, after two days. The results listed in Table 2 indicate that OA and UA were relatively stable in 60% ethanol solution for at least two days.

Table 3: Extraction recoveries of spiked analytes.

Analytes ^a	Content (mg)	Spiked (mg)	Found ^b (mg)	Recovery ^c (%)	Mean ^d (%)	RSD ^d (%)	Mean ^e (%)	RSD ^e (%)
OA	2.67	1.5	4.06 ± 0.19	92.67 ± 4.34	91.72	4.5	100.84	5.67
		3.0	5.47 ± 0.26	93.33 ± 4.45				
		4.5	6.79 ± 0.28	90.16 ± 3.81				
UA	14.95	8.0	22.20 ± 0.89	90.63 ± 3.71	91.29	4.3	100.39	5.41
		15.0	28.78 ± 1.21	92.20 ± 3.87				
		25.0	37.71 ± 1.75	91.04 ± 4.26				

^aThree different quantities of the authentic standards were added to the sample solution (HD1). The mixtures were extracted (1 extraction cycle) and analyzed using the optimized MUAE–HPLC method. ^bThe experiments were repeated six times at each level. Values are expressed as the mean ± standard deviation (SD). ^cRecovery (%) = [(found value – content)/spiked value] × 100. ^dOne extraction cycle. ^eThree extraction cycles.

Table 3 shows the recoveries of OA and UA after applying the combined MUAE–HPLC method. As listed in Table 3, the mean recoveries of OA and UA were 91.72%, with an average RSD of 4.56% (n = 6), and 91.29%, with an average RSD of 4.32% (n = 6), respectively (1 extraction cycle). The mean extraction recoveries of OA and UA reach 100% after three extraction cycles using MUAE in the present work. The recovery results revealed that the MUAE method was adequate and appropriate for the investigated analysis. The recovery tests also confirmed that the HPLC method was reliable and accurate for OA and UA.

Using optimized experimental conditions, the developed MUAE–HPLC method was applied for the simultaneous quantitation of OA and UA in the extracts of six samples of *H. diffusa* (HD1 to HD6) and five of *H. corymbosa* (HC7 to HC11) collected from different areas. The calculated content of the two components are summarized in Table 4 (one extraction cycle). There were significant variations in the OA and UA contents in the different samples of *H. diffusa* and *H. corymbosa*. The OA and UA contents in the six batches of *H. diffusa* ranged from 0.32 to 0.80 mg/g and from 1.75 to 3.42 mg/g, respectively. The RSD values calculated for the average content (HD1 to HD6) were 21.7% for OA and 28.2% for UA. For *H. corymbosa*, the OA and UA contents in the five batches ranged from 1.53 to 2.15 mg/g and from 6.36 to 8.54 mg/g, respectively. The RSD values calculated for the average content (HC7 to HC11) were 12.4% for OA and 11.4% for UA. These results were not surprising, as previous studies have revealed that the content of bioactive components in plants could vary greatly between cultivars [21]. Notwithstanding, further studies on the influence of genetic and environmental factors on bioactive content should be performed. Table 4 also reveals that the amount of OA in all the plant samples was markedly lower than the amount of UA. These results are valuable for the application of *H. diffusa* and *H. corymbosa* to clinical research and pharmacology.

Table 4: The mean contents of OA and UA present in 11 samples of *H. diffusa* and *H. corymbosa*.

Sample ^a	OA		UA	
	Content ^b (mg/g)	RSD ^c (%)	Content ^b (mg/g)	RSD ^c (%)
HD1	0.60 ± 0.02	3.17	3.37 ± 0.13	3.94
HD2	0.32 ± 0.01	3.84	1.75 ± 0.07	4.05
HD3	0.80 ± 0.02	2.97	3.42 ± 0.13	3.76
HD4	0.59 ± 0.02	4.25	2.16 ± 0.07	3.27
HD5	0.76 ± 0.03	3.87	2.94 ± 1.01	3.09
HD6	0.50 ± 0.01	3.06	1.97 ± 0.08	4.23
HC7	1.79 ± 0.08	4.22	7.04 ± 0.25	3.51
HC8	1.53 ± 0.05	3.43	6.36 ± 0.19	2.96
HC9	2.15 ± 0.08	3.58	8.54 ± 0.33	3.85
HC10	1.94 ± 0.06	2.94	7.85 ± 0.31	4.01
HC11	1.80 ± 0.07	4.03	7.99 ± 0.30	3.74

^aSix different *H. diffusa* and five different *H. corymbosa* samples were extracted using the optimized MUAE process (1 extraction cycle), and the extracts were analyzed by the developed HPLC method. ^bValues are expressed as the mean ± standard deviation (SD) of six replications and are calculated based on the dry weight of the plant. ^cRSD (%) = (SD/mean) × 100.

Experimental

Plant material: Six batches of dried, whole *H. diffusa* plants (Sample HD1 to HD6) and 5 samples of *H. corymbosa* (Sample HC7 to HC11) were purchased from different local Chinese medicinal shops (Taiwan). The air-dried whole plants were pulverized in a knife mill, and parts were sieved to produce samples with sizes of 0.6–0.85, 0.5–0.6, 0.21–0.5 and <0.21 mm. The moisture content (% dry weight basis) was determined by drying at 105°C to a constant mass and was 11.32 to 12.82% for samples HD1 to HC11. All the yields and compositions were calculated based on a moisture-free basis and represent the mean values of at least 6 experiments.

Chemicals: Oleanolic acid (97%) and ursolic acid (95%) were purchased as HPLC reference standards from the Sigma Chemical Co. (St. Louis, MO, USA). Methanol (99.9%), ethanol (99.9%), acetone (99.7%), acetonitrile (99.9%), ethyl acetate (99.9%), *n*-hexane (95%) and 85% phosphoric acid were bought from Merck Co. (Darmstadt, Germany), while cyclohexane and sulfuric acid were supplied by Tedia (Fairfield, OH, USA) and Acros Organics (Morris Plains, NJ, USA), respectively. Deionized water was prepared using a Milli-Q reverse osmosis unit from Millipore (Bedford, MA, USA).

Preparation of standard solutions: Standard stock solutions of OA and UA were prepared daily. The standard mixture solutions containing OA and UA at low (40 µg/mL), medium (200 µg/mL) and high (500 µg/mL) concentrations were prepared in a similar manner to yield the quality-control (QC) samples.

Modified ultrasound-assisted extraction (MUAE): MUAE was carried out with a working frequency of 40 kHz and 185W (Branson B-33510E-DTH, USA) of power; this extraction was operated in a pulse mode that provided 95 s of pulse on, followed by 25 s of pulse off for every 120 s (79% on/off time, work with the hands). There were 2 stages of extraction: HRE and UAE. An HRE (200 rpm) was used to obtain good solvent-to-plant material contact (10 min). For all experiments, 10 min HRE was followed by UAE. The MUAE conditions were then utilized at various ratios of solvent to raw material (6–22 mL/g), ultrasound amplitude (0–100%), ethanol concentrations (0–100 %), particle sizes (0.85–1.0, 0.6–0.85, 0.5–0.6, 0.21–0.5 and <0.21 mm), extraction times (5–60 min), extraction temperatures (30–80°C) and extraction cycles (1–4). At least 6 replicates were performed for each extraction condition tested, and all analyses were repeated 6 times.

Heat-reflux extraction (HRE): HRE was performed by mixing 5 g of dried, powdered plant with an appropriate amount of extraction solvent (100 mL) in a 250 mL conical flask that was immersed in a temperature-controlled water-bath, with stirring (200 rpm). The follow-up extractions were performed using the method detailed above for MUAE. The post-treatment of the extracts was also the same as that mentioned for MUAE.

High-performance liquid chromatography (HPLC): The HPLC analysis of OA and UA was performed on a Jasco HPLC system (Jasco, Tokyo, Japan) with a LiChrospher® C-18 analytical column (250 mm × 4 mm i.d., 5-µm particle size) (Merck, Darmstadt, Germany). The mobile phase was composed of acetonitrile (A) and 0.1% aqueous H₃PO₄ (B). The gradient was as follows: 0–25 min, 22–23% (solvent A), flow rate of 1 mL/min; 25–40 min, 23–23% (solvent A), flow rate of 1.0–1.5 mL/min; and 40–60 min, 23–90% (solvent A), and flow rate of 1.5–1.0 mL/min. The temperature of the column was maintained at 40°C, and the effluent was monitored at 210 nm. Detailed descriptions are available elsewhere [12].

Calibration curves: Eleven-point calibration curves for OA and UA were constructed. Working solutions at concentrations of 3, 6, 8, 10, 40, 90, 200, 400, 600, 800 and 1000 µg/mL were prepared for OA, and 10, 20, 40, 70, 100, 120, 200, 400, 600, 800 and 1000 µg/mL for UA. This operation was repeated on 6 consecutive days with freshly prepared calibration standards to select the most appropriate regression model.

Limits of detection and quantification: The LOD and LOQ of the analysis method were determined as the analyte concentrations giving rise to signal-to-noise ratios of 3 and 10, respectively. For this purpose, the baseline noise was also evaluated by the injection of 20 µL of the mobile phase B (blank) in 6 replications.

Precision and accuracy, analysis repeatability and stability: The precision and accuracy of the HPLC analysis method were established across the analytical range for OA and UA at 3 concentrations of OA and UA (40, 200 and 500 µg/mL). The intra-day precision was established as the repeatability of the assay and determined by calculating the RSD (%) for the repeated measurements, while the inter-day precision was evaluated as the intermediate precision of the assay. To determine the intra-day precision and accuracy for each standard, the same mixed standard

solutions were examined 6 times within one day. The inter-day precision and accuracy were established by analyzing each sample on 6 consecutive days; each sample was injected 5 times on each day. The assay precision was expressed by the RSD (%) between the replicate measurements, while accuracy was defined as the RE (%), which was calculated using the formula $RE (\%) = (\text{mean of observed concentration} - \text{spiked concentration}) / \text{spiked concentration} \times 100$. The injection repeatability was analyzed by injecting the same standard mixture solution 6 times sequentially, while analysis repeatability was examined by injecting 6 different samples obtained through the same sample preparation procedure. For the stability test, the same real sample used for analysis repeatability was analyzed after 48 h at room temperature.

Extraction recovery: To assess the extraction recoveries of OA and UA from *H. diffusa*, 3 different quantities (low, medium and high) of the authentic standards were added to the sample solution before extraction. The follow-up extractions and HPLC analyses were performed in the same manner as described above.

Acknowledgments - The authors thank the National Science Council of Taiwan and Chuang Song Zong Pharmaceutical Co. Ltd (Kaohsiung, Taiwan) for financial support. Finally, the authors would like to acknowledge the editors and referees for their constructive comments.

References

- [1] Xu GH, Kim YH, Chi SW, Choo SJ, Ryoo IJ, Ahn JS, Yoo ID. (2010) Evaluation of human neutrophil elastase inhibitory effect of iridoid glycosides from *Hedyotis diffusa*. *Bioorganic & Medicinal Chemistry Letters*, **20**, 513-515.
- [2] Wu PK, Tai WCS, Liang ZT, Zhao ZZ, Hsiao WLW. (2009) Oleanolic acid isolated from *Oldenlandia diffusa* exhibits a unique growth inhibitory effect against ras-transformed fibroblasts. *Life Sciences*, **85**, 113-121.
- [3] Kim Y, Park EJ, Kim J, Kim Y, Kim SR, Kim YY. (2001) Neuroprotective constituents from *Hedyotis diffusa*. *Journal of Natural Products*, **64**, 75-78.
- [4] Attard E, Attard H. (2006) The potential angiotensin-converting enzyme inhibitory activity of oleanolic acid in the hydroethanolic extract of *Crataegus monogyna* Jacq. *Natural Product Communications*, **1**, 381-385.
- [5] Yan SL, Huang CY, Wu ST, Yin MC. (2010) Oleanolic acid and ursolic acid induce apoptosis in four human liver cancer cell lines. *Toxicology in Vitro*, **24**, 842-848.
- [6] Zhang YX, Kong CZ, Wang HQ, Wang LH, Xu CL, Sun YH. (2009) Phosphorylation of Bcl-2 and activation of caspase-3 via the c-Jun N-terminal kinase pathway in ursolic acid-induced DU145 cells apoptosis. *Biochimie*, **91**, 1173-1179.
- [7] Ramos AA, Pereira-Wilson C, Collins AR. (2010) Protective effects of ursolic acid and luteolin against oxidative DNA damage include enhancement of DNA repair in Caco-2 cells. *Mutation Research*, **692**, 6-11.
- [8] Ovesná Z, Kozics K, Slameňová D. (2006) Protective effects of ursolic acid and oleanolic acid in leukemic cells. *Mutation Research*, **600**, 131-137.
- [9] Huang MT, Ho CT, Wang ZY, Ferraro T, Lou YR, Stauber K, Ma W, Georgiadis C, Laskin JD, Conne AH. (1994) Inhibition of skin tumorigenesis by rosemary and its constituents carnosol and ursolic acid. *Cancer Research*, **54**, 701-708.
- [10] Yim EK, Lee KH, Namkoong SE, Um SJ, Park JS. (2006) Proteomic analysis of ursolic acid-induced apoptosis in cervical carcinoma cells. *Cancer Letters*, **235**, 209-220.
- [11] Hsu YL, Kuo PL, Lin CC. (2004) Proliferative inhibition, cell-cycle dysregulation, and induction of apoptosis by ursolic acid in human non-small cell lung cancer A549 cells. *Life Sciences*, **75**, 2303-2316.
- [12] Yang YC, Wei MC, Huang TC. (2012) Optimisation of an ultrasound-assisted extraction followed by RP-HPLC separation for the simultaneous determination of oleanolic acid, ursolic acid and oridonin content in *Rabdosia rubescens*. *Phytochemical Analysis*, **23**, 627-636.
- [13] Yang YC, Wei MC, Hong SJ, Huang TC, Lee SZ. (2013) Development/optimization of a green procedure with ultrasound-assisted improved supercritical carbon dioxide to produce extracts enriched in oleanolic acid and ursolic acid from *Scutellaria barbata* D. Don *Industrial Crops and Products*, **49**, 542-553.
- [14] Fang X, Wang J, Yu X, Zhang G, Zhao J. (2010) Optimization of microwave-assisted extraction followed by RP-HPLC for the simultaneous determination of oleanolic acid and ursolic acid in the fruits of *Chaenomeles sinensis*. *Journal of Separation Science*, **33**, 1147-1155.
- [15] Claude B, Morin P, Lafosse M, Andre P. (2004) Evaluation of apparent formation constants of pentacyclic triterpene acids complexes with derivatized beta- and gamma-cyclodextrins by reversed phase liquid chromatography. *Journal of Chromatography A*, **1049**, 37-42.
- [16] Domingues RMA, Sousa GDA, Freire CSR, Silvestre AJD, Pascoal Neto C. (2010) *Eucalyptus globulus* biomass residues from pulping industry as a source of high value triterpenic compounds. *Industrial Crops and Products*, **31**, 65-70.
- [17] Xia EQ, Yu YY, Xu XR, Deng GF, Guo YJ, Li HB. (2012) Ultrasound-assisted extraction of oleanolic acid and ursolic acid from *Ligustrum lucidum* Ait. *Ultrasonics Sonochemistry*, **19**, 772-776.
- [18] Yang YC, Wei MC, Lian FY, Huang TC. (2013). Simultaneous extraction and quantitation of oleanolic acid and ursolic acid from *Scutellaria barbata* D. Don by ultrasound-assisted extraction and high-performance liquid chromatography. *Chemical Engineering Communications*, 201DOI: 10.1080/00986445.2013.777901.
- [19] Wei MC, Yang YC, Chiu HF, Hong SJ. (2013) Development of a hyphenated procedure of heat-reflux and ultrasound-assisted extraction followed by RP-HPLC separation for the determination of three flavonoids content in *Scutellaria barbata* D. Don, *Journal of Chromatography. B Analytical Technologies in the Biomedical and Life Sciences*, **940**, 126-134.
- [20] Yang YC, Wei MC, Huang TC, Lee SZ, Lin SS. (2013) Comparison of modified ultrasound-assisted and traditional extraction methods for the extraction of baicalin and baicalein from *Radix Scutellariae*. *Industrial Crops and Products*, **45**, 182-190.
- [21] Zhou C, Sheng Y, Zhao D, Wang Z, Tao J. (2010) Variation of oleanolic and ursolic acid in the flesh of persimmon fruit among different cultivars. *Molecules*, **15**, 6580-6587.

New Triterpenoid Saponins from the Roots of *Saponaria officinalis*Barbara Moniuszko-Szajwaj^{a,*}, Łukasz Pecio^a, Mariusz Kowalczyk^a, Ana M. Simonet^b, Francisco A. Macias^b, Małgorzata Szumacher-Strabel^c, Adam Cieślak^c, Wiesław Oleszek^a and Anna Stochmal^a^aDepartment of Biochemistry, Institute of Soil Science and Plant Cultivation, State Research Institute, Czartoryskich 8, 24-100 Puławy, Poland^bGrupo de Alelopatia, Departamento de Química Organica, Facultad de Ciencias, Universidad de Cadiz, C/Republica Saharaui, s/n, 11510 Puerto Real, Cadiz, Spain^cDepartment of Animal Nutrition and Feed Management, RUMEN PULS, Poznan University of Life Sciences, Wołyńska 33, 60-637 Poznań, Poland

bszajwaj@iung.pulawy.pl


Received: May 29th, 2013; Accepted: October 2nd, 2013

Three new triterpenoid saponins (**1-3**), along with nine known saponins, were isolated from the roots of *Saponaria officinalis* L. Two of them: vaccaroside D (**4**) and dianchinenoside B (**5**) are known, but not previously reported for *S. officinalis*, and seven others: saponarioside C (**6**), D (**7**), F (**8**), G (**9**), I (**10**), K (**11**), and L (**12**) have been previously isolated from this plant. The structures of the new saponins were established as 3-*O*- β -D-xylopyranosyl-16 α -hydroxygypsogenic acid-28-*O*-[β -D-glucopyranosyl-(1 \rightarrow 6)]- β -D-glucopyranoside (**1**), 3-*O*- β -D-xylopyranosyl-16 α -hydroxygypsogenic acid-28-*O*-[β -D-glucopyranosyl-(1 \rightarrow 3)]-[α -D-galactopyranosyl-(1 \rightarrow 6)]- β -D-glucopyranosyl-(1 \rightarrow 6)]- β -D-glucopyranoside (**2**) and 3-*O*- β -D-xylopyranosyl-gypsogenic acid-28-*O*-[β -D-glucopyranosyl-(1 \rightarrow 3)]-[6-*O*-(3-hydroxy-3-methylglutaryl)- β -D-glucopyranosyl-(1 \rightarrow 6)]- β -D-glucopyranoside (**3**). Their structures were elucidated by extensive spectroscopic methods, including 1D- (¹H, ¹³C) and 2D-NMR (D QF-COSY, TOCSY, ROESY, HSQC and HMBC) experiments, as well as high-resolution electrospray ionization mass spectrometry (HR-ESI-MS), ESI-MS/MS and acid hydrolysis.

Keywords: *Saponaria officinalis* L., Caryophyllaceae, Triterpenoid saponins, NMR.

Saponaria officinalis L., commonly named soapwort or bouncing-bet, is a perennial plant belonging to the family Caryophyllaceae. Its native range extends throughout Europe, northern Africa and west to central Asia, but currently is also cultivated in many countries across the world. It contains a high level of saponins and, like other such plants, it is used as an expectorant to remove catarrh from the upper respiratory tract. In ancient times, when soap was unknown, *S. officinalis* was used for cleaning and washing because of its detergent properties [1-3]. The saponin fraction of *S. officinalis* has shown an anti-inflammatory activity *in vitro* against carrageenan-induced rat-paw edema and inhibited prostaglandin synthetase. Purified saponins have indicated hypocholesterolemic effects *in vitro*, which is believed to be due to the ability of saponin to form an insoluble complex with cholesterol, preventing its absorption from the small intestine. Saponins also indicate spermicidal activity, which may result from their hemolytic property [4]. Unfortunately, apart from its beneficial effects, this plant is toxic for ruminants. It leads to photosensitization followed by liver and kidney degeneration and gut problems [5a]. Beside the pharmacological properties of saponins from *S. officinalis*, like anticancer, immunomodulatory and cytotoxicity activities [5b], Czaban and co-workers reported antifungal activities of the saponin fraction against *Gaeumannomyces graminis* var. *tritici* and *Fusarium culmorum*, which are pathogens of cereals [5c].

S. officinalis is a rich source of triterpenoid glycosides [6] with gypsogenic acid, hydroxygypsogenic acid and quillaic acid aglycones, mostly occurring as bisdesmosides. The highest yield of saponins in Caryophyllaceae species is usually located in roots or seeds [5b], therefore the study of root material led to the isolation of three novel saponins and two not described previously in this plant. The structural elucidation of these triterpenoid saponins was based on chemical methods and spectroscopic techniques.

**Figure 1:** Structure of compounds **1-3**.

The 70% MeOH extract of the roots of *S. officinalis* was subjected to successive chromatographic steps to provide three new triterpene glycosides (**1-3**), along with nine known saponins.

Compound **1** exhibited in the HR-ESI-MS, the [M-H]⁻ peak at *m/z* 957.4684 consistent with the molecular formula C₄₇H₇₃O₂₀. Its positive-ion ESI-MS displayed a quasimolecular ion at *m/z* 981 [M+Na]⁺ and fragmentation ion peaks were observed at *m/z* 849 [M+Na-132]⁺, 831 [M+Na-H₂O-132]⁺, 819 [M+Na-162]⁺, 657 [M+Na-162-162]⁺, 525 [M+Na-162-162-132]⁺, 365 [Na+162+162+H₂O]⁺, and 347 [Na+162+162]⁺. Its spectral features and physiochemical properties suggested **1** to be a triterpenoid saponin. From the 47 carbons, 30 were assigned to the aglycone part and 17 to the oligosaccharide moiety (Tables 1 and 2). The six *sp*³ hybrid carbon signals at δ_c 13.2, 16.6, 17.9, 25.0, 27.6, and 33.5, and the two *sp*² hybrid carbon signals at δ_c 122.9 and 144.9 together with the information from ¹H NMR analysis [six methyl

Table 1: ^{13}C and ^1H -NMR data for the aglycone moieties of compounds **1-3** in pyridine- d_5 (δ in ppm, J in Hz).^{a,b)}

	1		2		3	
	δ_{C}	δ_{H}	δ_{C}	δ_{H}	δ_{C}	δ_{H}
1	39.4	1.64 ^b , 1.20 ^b	39.4	1.61 ^b , 1.16 ^b	39.1	1.57 ^b , 1.11 ^b
2	26.8	2.20 ^b , 1.97 ^b	26.8	2.20 ^b , 1.95 ^b	26.6	2.20 ^b , 1.94 ^b
3	85.8	4.65 dd (11.9, 3.9)	85.2	4.62 ^b	85.4	4.62 dd (12.0, 3.1)
4	53.8	-	53.8	-	53.7	-
5	52.5	2.04 ^b	52.6	1.96 ^b	52.4	1.90 ^b
6	21.8	1.55 ^b , 1.48 ^b	21.8	1.60 ^b , 1.47 ^b	21.6	1.58 ^b , 1.46 ^b
7	33.5	1.76 ^b , 1.35 ^b	33.5	1.68 ^b , 1.23 ^b	32.7	1.82 ^b , 1.68 ^b
8	40.9	-	40.8	-	40.5	-
9	47.8	1.96 ^b	47.9	1.91 ^b	48.6	1.76 ^b
10	37.1	-	37.2	-	37.0	-
11	24.2	2.02 ^b	24.3	1.98 ^b , 1.92 ^b	24.1	1.95 ^b , 1.87 ^b
12	122.9	5.61 t (3.1)	122.9	5.58 t (3.4)	123.0	5.40 t (3.3)
13	144.9	-	144.8	-	144.4	-
14	42.4	-	42.5	-	42.4	-
15	36.5	2.51 d (14.0), 1.69 ^b	36.5	2.36 ^b , 1.62 ^b	28.5	2.17 ^b , 1.04 ^b
16	74.7	5.26 br s	74.6	5.20 br s	23.6	1.98 ^b , 1.85 ^b
17	49.5	-	49.5	-	47.3	-
18	41.6	3.53 dd (14.2, 4.3)	41.6	3.44 dd (14.2, 3.8)	41.9	3.12 ^b
19	47.5	2.77 t (13.6) 1.36 dd (12.5, 4.1)	47.6	2.70 t (13.6) 1.29 ^b	46.5	1.68 ^b , 1.19 ^b
20	31.2	-	31.2	-	31.0	-
21	36.3	2.40 ^b , 1.27 ^b	36.3	2.35 ^b , 1.25 ^b	34.2	1.55 ^b , 1.23 ^b
22	32.6	2.40 ^b , 2.17 ^b	32.5	2.31 ^b , 2.06 ^b	33.4	1.27 ^b , 1.05 ^b
23	181.7	-	181.6	-	180.8	-
24	13.2	1.58 s	13.2	1.58 s	13.0	1.55 s
25	16.6	1.01 s	16.6	1.00 s	16.4	0.92 s
26	17.9	1.15 s	17.9	1.10 s	17.7	1.08 s
27	27.6	1.79 s	27.6	1.79 s	26.3	1.19 s
28	176.4	-	176.1	-	176.5	-
29	33.5	0.97 s	33.5	0.95 s	33.1	0.86 s
30	25.0	1.05 s	25.1	1.01 s	24.0	0.83 s

^{a)}The assignments were based upon 2D-COSY, 2D-TOCSY, HSQC and HMBC spectra. ^{b)}Overlapped signals.

proton singlets at δ_{H} 0.97, 1.01, 1.05, 1.15, 1.58, and 1.79 and a broad triplet vinyl proton at δ_{H} 5.61 (t, $J = 3.1$ Hz) indicated that the aglycone possesses an olean-12-ene skeleton. The appearance of H-16 as a broad singlet at δ_{H} 5.26 indicated that 16-OH was α -oriented. This was supported by the high-frequency shifts of C-15 (δ_{C} 36.5) and C-16 (δ_{C} 74.7) in the ^{13}C NMR spectrum of **1**. Two high-frequency shifts at δ_{C} 181.7 (indicating an unsubstituted carboxylic group) and 176.4 were assigned to C-23 and C-28 on the basis of 1D ^{13}C and HMBC experiment, respectively. Detailed NMR analysis identified the aglycone as 16 α -hydroxygypsogenic acid [7a]. The high-frequency shift of the carbinol function at δ_{C} 85.8 (C-3) and low-frequency shift of the carboxylic function at δ_{C} 176.4 (C-28) showed the bisdesmosidic structure of **1**. The observation of two overlapping anomeric signals at δ_{H} 5.03 (d, $J = 7.8$ Hz) and a signal at δ_{H} 6.26 (d, $J = 8.2$ Hz) correlating with signals at δ_{C} 105.7, 106.6 and at δ_{C} 96.2 in the g-HSQC spectrum, respectively, suggested that this compound possesses three sugar moieties. The individual sugar units were identified on the basis of proton resonance analysis of each sugar unit. These determinations were carried out by a combination of DQF-COSY and 1D-TOCSY NMR experiments. 1D TOCSY subspectra were obtained from selective excitation of the anomeric protons. The selective TOCSY experiment for the signal at δ 6.26 revealed spin system characteristics of a β -glucose unit. The anomeric signals at δ_{H} 5.03 were too close to obtain independent subspectra in the 1D TOCSY experiment, but a different ratio of the pair of signals could be observed, indicating the presence of β -glucose and β -xylose units. These observations were supported by the 1D ROESY correlation between H-1 and H-3/H-5 protons of the sugar units. Analysis of the g-HSQC experiments allowed the assignments of signals of the ^{13}C NMR spectrum corresponding to the three sugar units. The sequence of the three sugar units was deduced from the 1D ROESY and HMBC spectra, in which long range correlations were observed from H-1_{Glc} (δ 5.03) to H-6b_{Glc} (δ 4.35) and C-6_{Glc} (δ 69.8), H-1_{Glc} (δ 6.26) to C-28 of the aglycone (δ 176.4), H-1_{Xyl} (δ 5.03) to H-3 of the aglycone (δ 4.65) and C-3 of the aglycone (δ 85.8). Thus the structure of **1** was established as 3-*O*- β -D-xylopyranosyl-16 α -

Table 2: ^{13}C and ^1H -NMR data for the sugar moieties of compounds **1-3** in pyridine- d_5 (δ in ppm, J in Hz).^{a,b,c)}

	1		2		3	
	δ_{C}	δ_{H}	δ_{C}	δ_{H}	δ_{C}	δ_{H}
Xyl-1	106.6	5.03 d (7.8)	106.6	5.02 d (7.3)	106.5	5.01 d (7.4)
2	75.7	3.96 t (8.1)	75.8	3.98 t (8.0)	75.6	3.98 ^b
3	78.4	4.11 ^b	78.3	4.10 ^b	78.0	4.09 ^b
4	71.5	4.19 ^b	71.5	4.20 ^b	71.4	4.22 ^b
Sax.	67.5	3.70 t (10.8)	67.5	3.71 t (11.0)	67.4	3.70 t (10.7)
Seq.		4.32 dd (10.5, 5.0)		4.36 dd (11.0, 5.0)		4.36 dd (11.2, 5.0)
Glc-1	96.2	6.26 d (8.2)	95.5	6.21 d (8.3)	95.1	6.22 d (8.1)
2	74.3	4.06 t (8.6)	72.3	4.06 t (8.0)	73.3	4.18 t (8.3)
3	79.1	4.20 t (8.9)	88.7	4.22 ^b	88.8	4.27 ^b
4	71.3	4.29 t (9.3)	68.6	4.22 ^b	69.3	4.24 ^b
5	78.4		77.9	4.05 ^b	78.0	4.14 ^b
6a	69.8	4.71 dd (11.1, 1.5)	69.5	4.61 dd (11.3, 3.7)	69.9	4.80 d (9.6)
6b		4.35 dd (11.2, 3.5)		4.20		4.31 dd (11.3, 4.0)
Glc'-1			106.0	5.24 d (8.0)	106.1	5.30 d (7.8)
2			75.0	4.00 t (8.2)	75.8	4.04 t (8.6)
3			78.1	4.16 t (8.7)	78.5	4.16 t (9.0)
4			71.3	4.12 t (8.7)	71.9	4.09 t (9.0)
5			78.9	4.00 ^b	78.7	3.98 ^b
6a			62.9	4.51 d (10.0)	62.5	4.53 dd (11.9, 1.6)
6b				4.24 dd (11.1, 3.8)		4.23 dd (12.0, 4.0)
Glc''-1	105.7	5.03 d (7.8)	105.8	4.88 d (7.8)	105.5	4.98 d (7.8)
2	75.5	4.01 t (8.0)	75.2	3.93 t (8.3)	75.3	3.98 t (8.3)
3	78.9	4.22 ^b	78.4	4.07 t (9.5)	78.4	4.18 t (8.7)
4	71.9	4.22 ^b	72.1	4.11 t (8.5)	72.1	3.99 ^b
5	78.7	3.90 ddd (7.8, 4.9, 2.4)	76.5	3.87 ddd (9.0, 5.0, 2.4)	78.4	4.03 ^b
6a		4.49 dd (11.8, 2.3)		4.51 dd (11.5, 3.5)		5.11 ^b
6b	63.0	4.37 dd (11.7, 5.0)	68.1	4.14 dd (10.5, 4.7)	65.2	4.64 ^b
Gal-1			100.8	5.42 d (3.7)		
2			70.5	4.62 dd (10.0, 3.4)		
3			71.8	4.49 dd (11.0, 3.5)		
4			71.5	4.57 ^b		
5			70.5	4.73 t (6.2)		
6a			68.6	4.47 dd (10.1, 6.5)		
6b				4.24 dd (10.3, 5.8)		
Gal'-1			101.2	5.55 d (3.7)		
2			70.6	4.64 dd (10.0, 2.3)		
3			72.1	4.55 dd (9.7, 3.3)		
4			72.0	4.63 d (3.8)		
5			73.1	4.55 dd (5.9, 5.9)		
6			63.0	4.40 2H, d (6.2)		
HMG-1					172.2	-
2					47.3	3.10 ^b
3						2.96 d (14.0)
4					70.4	-
5					47.3	3.10 ^b
6						3.05 d (13.8)
					172.2	-
					28.8	1.69 s

^{a)}The assignments were based upon 1D-TOCSY, 2D-COSY, 2D-TOCSY, HSQC, HSQC-TOCSY and HMBC spectra. ^{b)}Overlapped signals. ^{c)} ^{13}C chemical shifts of substituted residues are underlined.

hydroxygypsogenic acid-28-*O*-[β -D-glucopyranosyl-(1 \rightarrow 6)]- β -D-glucopyranoside, a new triterpenoid saponin. A similar compound was reported in *Dianthus superbus* (another member of Caryophyllaceae family), but containing an α -L-arabinopyranosyl unit attached to C-3 of the aglycone [7a].

Compound **2**, $\text{C}_{65}\text{H}_{103}\text{O}_{35}$ (HR-ESI-MS [$\text{M}-\text{H}$] $^-$ at m/z 1443.6276), exhibited a fragment-ion peak at m/z 1263 [$\text{M}-\text{H}-\text{H}_2\text{O}-162$] $^-$, corresponding with the loss of a hexosyl moiety. Other fragmentation ion peaks were observed at m/z 1119 [$\text{M}-\text{H}-162-162$] $^-$, 939 [$\text{M}-\text{H}-162-162-\text{H}_2\text{O}-162$] $^-$, 809 [$\text{M}-\text{H}-162-162-162-\text{H}_2\text{O}-132$] $^-$, 633 [$\text{M}-\text{H}-162-162-162-162-162$] $^-$, and 485 [$\text{M}-\text{H}-162-162-162-162-162-\text{H}_2\text{O}-132$] $^-$. Comparison of ^1H and ^{13}C NMR spectra (Tables 1 and 2) of **2** with those of **1** indicated that **2** had the same aglycone as **1**. The difference was observable in the saccharide moieties. Thus the aglycone of **2** was also determined to be 16 α -hydroxygypsogenic acid. Structure elucidation of the sugar portions was achieved by 1D TOCSY, DQF-COSY, g-HSQC, g-HSQC-TOCSY and g-HMBC experiments. The 1D TOCSY subspectra obtained by irradiating the six anomeric proton signals, allowed the determination of the nature of the sugar units. The signals at δ 6.21 (d, $J = 8.3$ Hz), 5.24 (d, $J = 8.0$ Hz) and 4.88 (d,

$J = 7.8$ Hz) showed the typical spin system of β -glucose units, whereas the subspectra obtained by irradiating the signal at δ 5.02 (d, $J = 7.3$ Hz) were established as belonging to a β -xylose unit. The characteristic spin system for the two α -galactoses was obtained by irradiation of signals at δ 5.55 (d, $J = 3.7$ Hz) and 5.42 (d, $J = 3.7$ Hz). The doublet at δ 5.55 generated a 1D TOCSY subspectrum with only two signals at δ 4.64 (dd, $J = 10.0, 2.3$ Hz) and 4.55 (dd, $J = 9.7, 3.3$ Hz). Similarly, the doublet at δ 5.42 generated a 1D TOCSY subspectrum with two signals at δ 4.62 (dd, $J = 10.0, 3.4$ Hz) and 4.49 (dd, $J = 11.0, 3.5$ Hz). Additional 1D TOCSY experiments conducted for isolated signals at δ 4.73 (t, $J = 6.2$ Hz) and 4.40 (d, 6.2) led to the identification of the H-6/H-4 sequences. These observations were supported by the 1D ROESY correlation between H-3/H-5 protons of the sugar units. Analysis of the g -HSQC and g -HSQC-TOCSY experiments allowed the assignments of signals of the ^{13}C NMR spectrum corresponding to the six sugar units. The sequence of the six sugar units was deduced from the 1D ROESY and HMBC spectra, in which long range correlations were observed from H-1_{Glc}' (δ 5.24) to H-3_{Glc} (δ 4.22) and C-3_{Glc} (δ 88.7), H-1_{Gal}' (δ 5.55) to H-6b_{Gal} (δ 4.24) and C-6_{Gal} (δ 68.6), H-1_{Gal} (δ 5.42) to H-6b_{Glc}' (δ 4.14) and C-6_{Glc}' (δ 68.1), H-1_{Glc}' (δ 4.88) to H-6b_{Glc} (δ 4.20) and C-6_{Glc} (δ 69.5), H-1_{Glc} (δ 6.21) to C-28 of the aglycone (δ 176.1), H-1_{Xyl} (δ 5.02) to H-3 of the aglycone (δ 4.62) and C-3 of the aglycone (δ 85.2). Thus the structure of **2** was established as 3-*O*- β -D-xylopyranosyl-16 α -hydroxygypsogenic acid-28-*O*-[β -D-glucopyranosyl-(1 \rightarrow 3)]-[α -D-galactopyranosyl-(1 \rightarrow 6)- α -D-galactopyranosyl-(1 \rightarrow 6)- β -D-glucopyranosyl-(1 \rightarrow 6)]- β -D-glucopyranoside. A triterpenoid saponin with α -D-galactopyranose as a component sugar is rare in nature, but has been reported from *Saponaria officinalis* and *Silene rubicunda*. Compound **2** is the first triterpenoid saponin containing two α -D-galactose units [2a,7b].

Compound **3** exhibited in its HR-ESI-MS, the $[\text{M}-\text{H}]^-$ peak at m/z 1247.5732 consistent with the molecular formula $\text{C}_{59}\text{H}_{91}\text{O}_{28}$. Fragment-ion peaks were observed at m/z 1103 $[\text{M}-\text{H}-144]^-$ and 923 $[\text{M}-\text{H}-144-\text{H}_2\text{O}-162]^-$, corresponding with the successive elimination of the 3-hydroxy-3-methylglutaryl (HMG) moiety and hexosyl moiety, respectively. Other fragmentation ion peaks were observed at m/z 1066 $[\text{M}-\text{H}-\text{H}_2\text{O}-162]^-$, 617 $[\text{M}-\text{H}-144-162-162-162]^-$, and 485 $[\text{M}-\text{H}-144-162-162-162-132]^-$. The aglycone of **3** was similar to **1** and **2** except for the lack of a hydroxyl group at C-16. This was supported by the low-frequency shifts of C-15 (δ_{C} 28.5) and C-16 (δ_{C} 23.6) in the ^{13}C NMR spectrum of **3** (Tables 1 and 2). Thus the aglycone of **3** was determined to be gypsogenic acid. Structure elucidation of the sugar portions was achieved by 1D TOCSY, DQF-COSY, g -HSQC, g -HSQC-TOCSY and g -HMBC experiments. The 1D TOCSY subspectra obtained by irradiating the four anomeric proton signals, allowed the determination of the nature of the sugar units. Signals at δ 6.22 (d, $J = 8.1$ Hz), 5.30 (d, $J = 7.8$ Hz), and 4.98 (d, $J = 7.8$ Hz) showed the typical spin system of β -glucose units, whereas the subspectra obtained by irradiating the signal at δ 5.01 (d, $J = 7.4$ Hz) were established as belonging to a β -xylose unit. These observations were supported by the 1D ROESY correlation between H-1 and H-3/H-5 protons of the sugar units. Analysis of the g -HSQC and g -HSQC-TOCSY experiments allowed the assignment of signals of the ^{13}C NMR spectrum corresponding to the four sugar units. The sequence of the four sugar units was deduced from the 1D ROESY and HMBC spectra, in which long range correlations were observed from H-1_{Glc}' (δ 5.30) to H-3_{Glc} (δ 4.27) and C-3_{Glc} (δ 88.8), H-1_{Glc}' (δ 4.98) to H-6b_{Glc} (δ 4.31) and C-6_{Glc} (δ 69.9), H-1_{Glc} (δ 6.22) to C-28 of the aglycone (δ 176.5), H-1_{Xyl} (δ 5.01) to H-3 of the aglycone (δ 4.62) and C-3 of the aglycone (δ 85.4). Additional signals, not belonging to the sugar and aglycone parts, were observed in the ^1H and ^{13}C

spectra of **3**, suggesting the presence of a 3-hydroxyl-3-methylglutaryl (HMG) moiety: $-\text{CH}_3$ group at δ_{H} 1.69 (s) and δ_{C} 28.8, two methylenes [δ_{H} 3.10 and 2.96 (d, $J = 14.0$ Hz)], δ_{C} 47.3 and [δ_{H} 3.10 and 3.05 (d, $J = 13.8$ Hz)], δ_{C} 47.3 and two overlapping carboxylic groups at δ_{C} 172.2, and one quaternary carbinol carbon at δ_{C} 70.4. The ester bond between HMG-1 and C-6_{Glc}' was suggested by high-frequency shifts for the H-6_{Glc}' (δ_{H} 5.11 and 4.64) and for the C-6_{Glc}' (δ_{C} 65.2), as the HMBC connectivity was not observed. Thus the structure of **3** was established as 3-*O*- β -D-xylopyranosyl-gypsogenic acid-28-*O*-[β -D-glucopyranosyl-(1 \rightarrow 3)]-[6-*O*-(3-hydroxy-3-methylglutaryl)- β -D-glucopyranosyl-(1 \rightarrow 6)]- β -D-glucopyranoside. A similar compound was reported in *Dianthus superbus*, but containing a β -D-glucopyranosyl unit attached to C-3 of the aglycone [7a].

Moreover, nine known compounds were identified as vaccaroside D (**4**) [7c], dianchinenoside B (**5**) [8a], saponarioside C (**6**), D (**7**), F (**8**) and G (**9**) [2a] and saponarioside I (**10**), K (**11**) and L (**12**) [2b], by comparing their NMR data with those in the literature. Compounds **4** and **5** were isolated from this genus for the first time.

Usually, several oleane-type saponins are accompanied in one species but *Saponaria officinalis* like *Vaccaria segetalis* contains saponins that are based on more than three different sapogenins [5b]. Both new and known saponins have gypsogenic acid, hydrogypsogenic acids and their derivatives [2b,c] as an aglycone component. Existing research suggests that *S. officinalis* is lacking gypsogenin, in contrast to the rest of the Caryophyllaceae family, in which the occurrence of gypsogenin is more frequent than that of gypsogenic acid [5b].

Almost all isolated saponins had a bisdesmosidic linkage, with the exception of saponarioside K and vaccaroside B, which are monodesmosides. The linking points for sugar moieties to the triterpenoid skeletons in *S. officinalis* are at C-3 and C-28. For the first time in this plant we proved the presence of 3-hydroxyl-3-methylglutaryl in a sugar chain – compound **3**.

In spite of the fact that *Saponaria officinalis* is widespread over different climes we still know little about its phytochemical composition, especially about the presence of saponins and their structures. Since this plant is used in folk medicine there is a need for confirmation of its composition by modern methods and proper standardization of plant material used in experiments.

Experimental

General experimental procedure: 1D and 2D NMR, Varian INOVA-600 spectrometer; ESI-MS/MS, Thermo LCQ Advantage Max mass spectrometer; HR-ESI-MS, Maldi SYNAPT G2-S HDMS Qq-TOF spectrometer; CC, Sephadex LH-20; VLC, LiChroprep C18 (Merck). HPLC, was performed on a Kromasil C18 column (10 x 250 mm, 5 μm) using a Gilson semi-preparative HPLC equipped with a PrepELS II (Evaporative Light Scattering) detector (the drift tube temperature was maintained at 65°C and the pressure of the nebulizer gas – nitrogen – was 47 psi).

Plant material: Dry, ground roots of *Saponaria officinalis* were purchased from a commercial source (Herbapol in Cracow, Poland). A voucher sample has been deposited at the Department of Biochemistry and Crop Quality, Institute of Soil Science and Plant Cultivation, State Research Institute, Pulawy.

Extraction and isolation: The isolation of the saponin fraction was performed according to previously developed procedures [8b]. Powdered (500 g) and defatted (497 g) roots of *S. officinalis* were

extracted 3 times with 80% MeOH at room temperature for 24 h. After each extraction the crude extract was filtered. The combined extracts were concentrated under reduced pressure and lyophilized (233 g). The extract was suspended in H₂O and applied to a vacuum liquid chromatograph (100 x 60 mm, LiChroprep C18, 40-63 µm) preconditioned with water. Sugars and phenolics were removed with water and 30% MeOH. Saponins were then eluted from the column with 90% MeOH. The solvent was evaporated *in vacuo* and the residue lyophilized, obtaining 49.9 g of crude saponin mixture, which was suspended in H₂O and extracted with EtOAc saturated with water. This extraction separated nonpolar compounds, which went into the EtOAc layer, such as catechins and yielded 45.41 g of saponins. The purified saponin fraction was suspended in H₂O and then partitioned successively between H₂O and *n*-BuOH saturated with water. The *n*-BuOH soluble fraction was fractionated by CC (Sephadex LH-20, 44 x 450 mm, 25-100 µm, Sigma-Aldrich) preconditioned with 40% MeOH containing 0.1% HCOOH, and eluted with 40% MeOH containing 0.1% HCOOH. This separation gave 5 subfractions. Individual saponins were separated from the subfractions using a semi-preparative high-pressure chromatograph (Gilson) equipped with a PrepELS II (Evaporative Light Scattering) detector and a Kromasil C18 column (10 x 250 mm, 5µm; Eka Chemicals AB). The flow rate was 5 mL min⁻¹. 0.1% HCOOH was used as mobile phase A and acetonitrile containing 0.1% HCOOH as mobile phase B. The drift tube temperature was maintained at 65°C and the pressure of the nebulizer gas – nitrogen – was 47 psi. In this way all compounds were isolated.

3-*O*-β-D-Xylopyranosyl-16α-hydroxygypsogenic acid-28-*O*-[β-D-glucopyranosyl (1→6)]-β-D-glucopyranoside (1)

¹H and ¹³C NMR: Tables 1 and 2.

ESI-MS/MS: *m/z* 957 [M-H]⁻ and 981 [M+Na]⁺, 849, 831, 819, 809, 657, 525, 365, 347.

HR-ESI-MS: *m/z* 957.4684 [M-H]⁻, calcd. for C₄₇H₇₃O₂₀: 957.4695.

3-*O*-β-D-Xylopyranosyl-16α-hydroxygypsogenic acid-28-*O*-[β-D-glucopyranosyl-(1→3)]-[α-D-galactopyranosyl-(1→6)]-α-D-galactopyranosyl-(1→6)-β-D-glucopyranosyl-(1→6)]-β-D-glucopyranoside (2)

¹H and ¹³C NMR: Tables 1 and 2.

ESI-MS/MS: *m/z* 1443 [M-H]⁻, 1263, 1119, 939, 633, 485.

HR-ESI-MS: *m/z* 1443.6276 [M-H]⁻, calcd. for C₆₅H₁₀₃O₃₅: 1443.6280.

3-*O*-β-D-Xylopyranosyl-gypsogenic acid-28-*O*-[β-D-glucopyranosyl-(1→3)]-[6-*O*-(3-hydroxy-3-methylglutaryl)-β-D-glucopyranosyl-(1→6)]-β-D-glucopyranoside (3)

¹H and ¹³C NMR: Tables 1 and 2.

ESI-MS/MS: *m/z* 1247.5 [M-H]⁻, 1103, 1066, 923, 617, 485.

HR-ESI-MS: *m/z* 1247.5707 [M-H]⁻, calcd. for C₅₉H₉₁O₂₈: 1247.5732.

Acid hydrolysis of saponins and determination of absolute configuration of monosaccharides: Each compound (2.5 mg) was hydrolysed in 1.5 mL solution of 3 M HCl (dioxane-H₂O, 1:2) at 90°C for 5 h. After dioxane was removed from the hydrolysis reaction mixture under a stream of nitrogen, the aglycones were extracted with EtOAc (3×1 mL) and dried. The aqueous layer containing sugars was neutralized using an Amberlite IRA-400 ion exchange resin in its hydroxide form and concentrated under a stream of nitrogen at 60°C and then used for sugar determination. To determine the absolute configuration of the monosaccharide ingredients of isolated compounds, the method of Tanaka *et al.* [8c] was used, with slight modification. Sugars of each sample were dissolved in pyridine (0.5 mL) to which 1 mg of L-cysteine methyl ester hydrochloride was added. The mixture was kept at 60°C for 1 h, and then 0.2 µL of *o*-tolyl isothiocyanate was added to the mixture, which was heated at 60°C for 1 h. Each mixture was directly analysed by reversed-phase HPLC using a 616-Pump, 966-Photodiode Array detector 717-plus Autosampler (Waters) and an Eurospher-100 C18 (250 mm x 4 mm; 5 µm Knauer) column. MeCN-H₂O (25:75, v/v) containing 50 mM H₃PO₄ was used as mobile phase at a flow rate 0.8 mL min⁻¹ and a column temperature of 35°C. Detection was performed by UV at 250 nm. The absolute configuration was determined by comparing the retention times of sugars with derivatives prepared in a similar way from standard monosaccharides (Sigma-Aldrich). The following sugars were detected for **1** and **3**: D-glucose and D-xylose, and for **2**: D-glucose, D-xylose and D-galactose.

Supplementary data: NMR spectra (¹H and ¹³C NMR, HSQC, HMBC, TOCSY, ROESY and DQF-COSY), HR-ESI-MS and ESI-MS/MS spectra for the new compounds (**1-3**).

Acknowledgments - This work has been supported by the Polish Ministry of Science and Higher Education (Project: NN 311 4763 39) and in part by European Community 7th Framework Program project Proficiency (FP7-REGPOT-2009-1-245751).

References

- [1] Jia Z, Koike K, Nikaido T. (1998) Major triterpenoid saponins from *Saponaria officinalis*. *Journal of Natural Products*, **61**, 1368-1373.
- [2] (a) Jia Z, Koike K, Nikaido T. (1999) Saponarioside C, the first α-D-galactose containing triterpenoid saponin, and five related compounds from *Saponaria officinalis*. *Journal of Natural Products*, **62**, 449-453; (b) Koike K, Jia Z, Nikaido T. (1999) New triterpenoid saponins and saponinins from *Saponaria officinalis*. *Journal of Natural Products*, **62**, 1655-1659; (c) Wood M. (2008) *Materia medica*. In *The earthwise herbal: a complete guide to old world medicinal plants*. North Atlantic Books, Berkeley, California, 463.
- [3] Oleszek W. (2000) Saponins. In *Natural Food Antimicrobial System*, Naidu AS. (Ed.), CRC Press, Inc., USA, 295-324.
- [4] Jia Z, Koike K, Sahu NP, Nikaido T. (2002) Triterpenoid saponins from Caryophyllaceae family. In *Studies in Natural Products Chemistry, Bioactive Natural Products*, Atta-ur-Rahman (Ed.), Elsevier, Amsterdam, The Netherlands, **26**, 3-61.
- [5] (a) Wina E, Muetzel S, Becker K. (2005) The impact of saponins or saponin-containing plant materials on ruminant production – a review. *Journal of Agriculture and Food Chemistry*, **53**, 8093-8105; (b) Bötger S, Melzig MF. (2011) Triterpenoid saponins of the Caryophyllaceae and Illecebraceae family. *Phytochemistry Letters*, **4**, 59-68; (c) Czaban J, Molodoch J, Wróblewska B, Szumacher-Strabel M, Cieślak A, Oleszek W, Stochmal A. (2013) Effect of triterpenoid saponins of field scabious, alfalfa, red clover and common soapwort on growth of *Gaeumannomyces graminis* var. *tritici* and *Fusarium culmorum*. *Allelopathy Journal*, **32**, 79-90.
- [6] Vincken J-P, Heng L, de Grot A, Gruppen H. (2007) Saponins, classification and occurrence in the plant kingdom. *Phytochemistry*, **68**, 275-297.
- [7] (a) Luo J-G, Chen X, Kong L-Y. (2011) Three new triterpenoid saponins from *Dianthus superbus*. *Chemical and Pharmaceutical Bulletin*, **59**, 518-521; (b) Fu H, Koike K, Li W, Nikaido T, Lin W, Guo D. (2005) Silenorubicosides A – D, triterpenoid saponins from *Silene rubicunda*. *Journal of Natural Products*, **68**, 754-758; (c) Koike A, Jia Z, Nikaido T. (1998) Triterpenoid saponins from *Vaccaria segetalis*. *Phytochemistry*, **47**, 1343-1349.
- [8] (a) Li H-Y, Koike K, Ohmoto T. (1993) Dianchinenosides A and B, two new saponins from *Dianthus chinensis*. *Journal of Natural Products*, **56**, 1065-1070; (b) Szajwaj B, Molodoch J, Masullo M, Piacente S, Oleszek W, Stochmal A. (2011) Amides and esters of phenylpropenoic acids from the aerial parts of *Trifolium pallidum*. *Natural Product Communications*, **6**, 1293-1296; (c) Tanaka T, Nakashima T, Ueda T, Tomii K, Kouno I. (2007) Facile discrimination of aldose enantiomers by reversed-phase HPLC. *Chemical and Pharmaceutical Bulletin*, **55**, 899-901.

Minor Triterpene Saponins from Underground Parts of *Lysimachia thyrsoflora*: Structure elucidation, LC-ESI-MS/MS Quantification, and Biological Activity

Irma Podolak^{a,*}, Paweł Żmudziński^c, Paulina Koczurkiewicz^{a,b}, Marta Michalik^b, Paweł Zajdel^c and Agnieszka Galanty^a

^aDepartment of Pharmacognosy, Pharmaceutical Faculty, Medical College, Jagiellonian University, Medyczna 9, 30-688 Cracow, Poland

^bDepartment of Cell Biology, Faculty of Biochemistry, Biophysics and Biotechnology, Jagiellonian University, Gronostajowa 7, 30-387 Cracow, Poland

^cDepartment of Medicinal Chemistry, Pharmaceutical Faculty, Medical College, Jagiellonian University, Medyczna 9, 30-688 Cracow, Poland

mfpodola@cyf-kr.edu.pl

Received: August 8th, 2013; Accepted: September 8th, 2013

Two minor triterpene saponins, one previously not reported, were isolated from the underground parts of *Lysimachia thyrsoflora* L. The structures were determined based on a combination of one- and two-dimensional NMR techniques, and mass spectrometry, as 30-*O*- β -D-glucopyranosyl-3 β ,16 α ,30-trihydroxy-olean-12-en-28-yl acetate 3-*O*- β -D-glucopyranosyl-(1 \rightarrow 4)-*O*- β -D-glucopyranosyl-(1 \rightarrow 2)-*O*- α -L-arabinopyranoside (denoted as thyrsoilide A), and davuricoside L. Quantitative determination of these two compounds, along with other saponins present in *L. thyrsoflora* underground parts, was performed by a developed UPLC-ESI-MS/MS method, validated according to the ICH guidelines. The relative total content of triterpene saponins amounted to 1.19 μ g/mg dw. Both **4** and **5** were present in minor quantities, 0.031 and 0.077 μ g/mg dw, respectively. Assays performed on a panel of human cancer cell lines showed antiproliferative activity against prostate carcinoma PC-3 (GI₅₀ after 48 h = 20 and 24 μ g/mL, respectively).

Keywords: *Lysimachia thyrsoflora*, Olean-12-ene saponins, UPLC-ESI-MS/MS quantification, Cytotoxicity, Antifungal activity.

Lysimachia L. is a genus of over hundred species, both wild and cultivated in Europe and Asia. Many of them have been used in traditional folk medicine for their anticholelithic, anthelmintic, and antihypertensive properties. Pentacyclic triterpene glycosides are the most characteristic constituents within this genus, and generally have oleanane-derived sapogenols of two structural types: **I**. 13 β ,28-epoxy and **II**. Δ^{12} -17-CH₂OH. Compounds of type **I**, with a completely saturated pentacyclic skeleton, are fairly uncommon and are found almost exclusively in the families Myrsinaceae and Primulaceae [1]. Although the genus *Lysimachia* is traditionally classified in Primulaceae, some recent phylogenetic data suggest its relocation to the family Myrsinaceae [2]. *L. thyrsoflora* L. (tufted loosestrife) is among five representatives of this genus reported in Poland. In addition to flavonoids, ecdysteroids and benzoquinones, the plant contains triterpene saponins. While tri- and tetrasaccharides of priverogenin A were reported in the aerial parts [3,4], in our own previous investigations of *L. thyrsoflora* we isolated cyclaminorin (**1**), ardisicrispin A (**2**), and ardisicrenoside B (**3**) from the underground parts [5,6]. These compounds exhibited interesting biological activities, such as the uterocontractile activity of **1** and **2** [7,8], cytotoxicity of **2** towards human cancer cell lines [9,10], and antifungal activity of **1** against *Candida* yeasts [7].

As part of our continuing phytochemical investigations of the *Lysimachia* genus, in this paper we report the isolation and structure elucidation of two additional triterpene saponins, 30-*O*- β -D-glucopyranosyl-3 β ,16 α ,30-trihydroxy-olean-12-en-28-yl acetate 3-*O*- β -D-glucopyranosyl-(1 \rightarrow 4)-*O*- β -D-glucopyranosyl-(1 \rightarrow 2)-*O*- α -L-arabinopyranoside (**4**) and davuricoside L (**5**) from the underground parts of *L. thyrsoflora*. Of these, **4** is a new compound and both were isolated for the first time from this species.




Figure 1: Structures of compounds **4** and **5**.

Furthermore, an UPLC-ESI-MS/MS method was developed and validated for the quantitative determination of **4** and **5**, along with other triterpene glycosides (**1-3**) in this plant species. In the context of biological activity of *Lysimachia* saponins, our present work has also focused on the screening of isolated compounds against human tumor and normal cell lines, as well as fungal strains, to establish potential structure-activity correlations.

A combination of repetitive open column and preparative silica gel chromatography of the MeOH extract of the underground parts of *L. thyrsoflora* yielded compounds **4** and **5** (Figure 1).

Compound **4** showed a [M + Na]⁺ quasimolecular ion peak at *m/z* = 1157. 57 in the positive HR-ESI-TOF-MS (C₅₅H₉₀O₂₄Na)

indicating a molecular mass of 1134 Da. The negative-ion FAB-MS showed the major ion peak at m/z 1133.4, which was assigned to $[M - H]^-$, a peak corresponding to the loss of one hexose unit at m/z 971.4 $[M - H - 162]^-$, and two hexose units at m/z 809.3 $[M - H - 162 - 162]^-$. The IR spectrum, apart from absorption bands due to the presence of hydroxyl groups (3333 cm^{-1}), showed a significant absorption peak at 1737 cm^{-1} (ester). The ^{13}C NMR spectrum displayed 55 carbons, of which 30 were assigned to the aglycone part, 23 to the oligosaccharide moiety, and the remaining 2 to an acetyl group (Tables 1 and 2). The ^1H NMR spectrum showed the presence of six tertiary methyl groups [δ_{H} 0.86, 0.95, 0.96, 0.98, 1.06, 1.37] and one olefinic proton [δ_{H} 5.36 (t, $J = 3.6\text{ Hz}$)]. The corresponding carbons were identified by a HSQC experiment [δ_{C} 16.9, 17.7, 28.1, 16.2, 28.5, 27.6, respectively]. The chemical shifts of the two olefinic carbons [δ_{C} 124.1 and 144.2], and comparison of the remaining chemical shifts (analyzed with DEPT and HSQC) with those established for pentacyclic triterpenes [11], revealed the characteristic carbon pattern of an olean-12-ene skeleton. Moreover, characteristic signals due to an acetyl group were observed [δ_{H} 2.03, δ_{C} 20.9, 173.2]. The structural assignment was initiated from the long-range coupling networks observed between the methyl protons and the adjacent carbons in the HMBC experiment (Figure 2). Two methyl groups (C-23 and C-24) and a methyl and CH_2OR group (C-29 and C-30, respectively) were shown to be geminal pairs, and the six methyls exhibited the following key connectivities: H-23 (δ_{H} 1.06) and H-24 (δ_{H} 0.86) to C-3 (δ_{C} 91.4), C-4 (δ_{C} 40.4), C-5 (δ_{C} 57.0), and carbon resonances of each other (δ_{C} 28.5; 16.9); H-25 (δ_{H} 0.98) to C-5 (δ_{C} 57.0), C-9 (δ_{C} 48.2), C-10 (δ_{C} 37.8); H-26 (δ_{H} 0.95) to C-7 (δ_{C} 34.0), C-9 (δ_{C} 48.2), and C-14 (δ_{C} 42.7); H-27 (δ_{H} 1.37) to C-8 (δ_{C} 41.0), C-14 (δ_{C} 42.7), and C-15 (δ_{C} 35.2); H-29 (δ_{H} 0.96) to C-19 (δ_{C} 43.4), C-20 (δ_{C} 35.6), and C-30 (δ_{C} 76.0).




Figure 2: Detail (methyl region) of the HMBC spectrum of compound 4.

Two oxygenated methine protons at δ_{H} 3.16 (dd, $J = 11.4, 4.4\text{ Hz}$) and 3.92 (br d, $J = 5.8\text{ Hz}$), which showed correlations in the HSQC spectrum with carbon signals at δ_{C} 91.4 and 73.7, respectively, were assigned to H-3 and H-16. Moreover, the ^1H NMR spectrum displayed two primary alcoholic functions [δ_{H} 3.68, 3.91], and [δ_{H} 3.35, 3.88] which were located at C-28 and C-30, respectively, on the basis of HMBC correlations from H-28 to C-17 (δ_{C} 40.7) and C-22 (δ_{C} 28.3) and from H-30 to C-19 (δ_{C} 43.4), C-20 (δ_{C} 35.6),

Table 1: ^{13}C and ^1H NMR data (J in Hz) of the aglycone moieties of compounds 4 and 5 (600 MHz, δ ppm, in CD_3OD).

No.	4 δ_{C}	δ_{H}^a (J in Hz)	5 δ_{C}	δ_{H}^a (J in Hz)
1	40.0	1.66, 0.99, m	40.0	1.65, 0.99, m
2	26.9	1.84, 1.75, m	27.0	1.85, 1.75, m
3	91.4	3.16, dd (11.4, 4.4)	91.4	3.16, dd (11.7, 4.3)
4	40.4	-	40.4	-
5	57.0	0.79 d (11.4)	57.0	0.79, d (11.2)
6	19.3	1.58, 1.43, m	19.3	1.57, 1.43, m
7	34.0	1.60, 1.40, m	34.0	1.59, 1.40, m
8	41.0	-	41.0	-
9	48.2	1.63, m	48.3	1.63, m
10	37.8	-	37.8	-
11	24.4	1.89 (2H), m	24.5	1.89 (2H), m
12	124.1	5.36, t (3.6)	123.7	5.35, t (3.4)
13	144.2	-	144.8	-
14	42.7	-	42.6	-
15	35.2	1.88, 1.37, m	35.0	1.90, 1.34, m
16	73.7	3.92, br d (5.8)	74.8	3.92, br d (5.7)
17	40.7	-	41.5	-
18	43.2	2.15, m	42.5	2.10, m
19	43.4	2.03, 1.36, m	43.6	2.06, 1.35, m
20	35.6	-	35.7	-
21	31.8	1.65, 1.47, m	33.9	1.59, 1.40, m
22	28.3	1.77, 1.49, m	32.2	1.62, 1.46, m
23	28.5	1.06, s	28.5	1.07, s
24	16.9	0.86, s	16.9	0.86, s
25	16.2	0.98, s	16.2	0.98, s
26	17.7	0.95, s	17.7	0.95, s
27	27.6	1.37, s	27.6	1.37, s
28	72.3	3.91, 3.68	70.2	3.20 (2H), d (9.2)
29	28.1	0.96 s	28.3	0.95, s
30	76.0	3.88, 3.35	75.9	3.89, 3.33
CH_3CO	20.9	2.03, s	-	-
	173.2	-	-	-

^a Multiplicity is not clear for some resonances due to overlapping.

Table 2: ^{13}C and ^1H NMR spectroscopic data (J in Hz) of the sugar portions of compounds 4 and 5 (600 MHz, δ ppm, in CD_3OD).

No.	4 δ_{C}	δ_{H}^a (J , Hz)	5 δ_{C}	δ_{H}^a (J , Hz)
α-L-Ara (at C-3)				
1	105.1	4.52, d (5.5)	105.2	4.52, d (5.8)
2	78.7	3.92	78.6	3.91
3	73.4	3.89	73.4	3.89
4	78.3	3.98	78.3	3.98
5	64.5	4.12, dd (12.3, 4.3)	64.5	4.12, dd (12.4, 4.4)
		3.56 dd (12.3, 1.8)		3.56, dd (12.4, 1.8)
β-D-GlcI (at C-2 Ara)				
1	104.4	4.62, d (7.7)	104.4	4.62, d (7.7)
2	75.7	3.20	75.8	3.19
3	78.2	3.36	78.2	3.36
4	71.9	3.22	71.9	3.21
5	77.8	3.28	77.9	3.28
6	63.1	3.84, dd (11.9, 2.0)	63.1	3.84, dd (11.3, 2.0)
		3.63, dd (11.9, 4.9)		3.63 dd (11.3, 4.9)
β-D-GlcII (at C-4 Ara)				
1	105.6	4.48, d (7.7)	105.6	4.48, d (7.8)
2	75.5	3.26	75.4	3.23
3	78.0	3.28	78.0	3.28
4	71.8	3.30	71.8	3.29
5	77.8	3.37	77.8	3.37
6	62.7	3.86, 3.67	62.7	3.86, 3.67
β-D-GlcIII (at C-30)				
1	105.2	4.25, d (7.7)	105.0	4.24, d (7.8)
2	75.3	3.21	75.3	3.21
3	78.1	3.28	78.1	3.27
4	71.5	3.29	71.5	3.29
5	78.0	3.36	78.0	3.36
6	62.9	3.87, dd (11.9, 2.4)	62.9	3.87, dd (11.4, 3.0)
		3.68		3.68

^a Multiplicity is not clear for some resonances due to overlapping.

C-29 (δ_{C} 28.1). Spatial proximities observed in the ROESY experiment, between H-3 and H-23, H-3 and H-5, H-16 and H-18, H-16 and H-28, confirmed the β -orientation of the hydroxyl group at C-3 and α -orientation at C-16, respectively.

The location of the acetyl group was established at C-28 from the long-range HMBC coupling between H-28 (δ_{H} 3.68, 3.91) and C-1 of the acetyl (δ_{C} 173.2). From the above evidence, the aglycone of compound 4 was identified as $3\beta,16\alpha,30$ -trihydroxy-olean-12-en-28-yl acetate.

The downfield shifts of C-3 (δ_C 91.4) and C-30 (δ_C 76.0) suggested that compound **4** was a bidesmosidic saponin. The sugar residues were determined to be L-arabinose and D-glucose after acid hydrolysis followed by derivatization [12]. In the 1H NMR spectrum, signals corresponding to four anomeric protons were found at [δ_H 4.25 (d, J = 7.8 Hz), 4.48 (d, J = 7.7 Hz), 4.52 (d, J = 5.5 Hz), and 4.62 (d, J = 7.7 Hz)]. These were correlated by HSQC experiment to the corresponding carbon resonances at δ_C 105.2, 105.6, 105.1, and 104.4, respectively (Table 2). The remaining proton resonances within each sugar unit were established by TOCSY, DQF-COSY, and ROESY experiments. HSQC was used to assign the respective carbons. All monosaccharides were determined to be in the pyranose forms by ^{13}C NMR evidence [13]. From the relatively large H-1 coupling constants (~ 7.7 Hz) it was clear that the anomeric hydroxyls of all three glucose moieties should have a β -configuration. The value of the $J_{1,2}$ coupling constant for the anomeric proton of arabinose (5.5 Hz), which was reported not to be diagnostic on its own, together with NOE connectivities between H-1_{ara}, H-3_{ara} and H-5_{ara} [14] indicated an α -orientation of this sugar unit. Sequencing and points of attachment in the oligosaccharidic chains were deduced from HMBC experiment and ^{13}C shift differences between individual sugar residues and model compounds [13]. For the trisaccharide side chain at C-3 the following key-correlation peaks were observed in the HMBC spectrum (Figure 3): H-1_{ara} (δ_H 4.52) and C-3 (δ_C 91.4), H-1_{glcI} (δ_H 4.62) and C-2_{ara} (δ_C 78.7), H-1_{glcII} (δ_H 4.48) and C-4_{ara} (δ_C 78.3). The side chain at C-30 of the aglycone was identified as β -D-glucose, with its anomeric proton at δ_H 4.25, and its anomeric carbon at δ_C 105.2. Its location was confirmed by the HMBC correlation between H-1_{glcIII} and C-30 (δ_C 76.0).




Figure 3: Detail (sugar region) of the HMBC spectrum of compound **4**.

The same conclusion with regard to the sugar sequence was also drawn from the ROESY experiment. Correlations were observed between H-1_{ara} (δ_H 4.52) and H-3 (δ_H 3.16); H-1_{glcI} (δ_H 4.62) and H-2_{ara} (δ_H 3.92); H-1_{glcII} (δ_H 4.48) and H-4_{ara} (δ_H 3.98); and H-1_{glcIII} and H-30 (δ_H 3.35, 3.88). Based on the above findings the structure of compound **4** was elucidated to be 30- O - β -D-glucopyranosyl-3 β ,16 α ,30-trihydroxy-olean-12-en-28-yl acetate 3- O - β -D-glucopyranosyl-(1 \rightarrow 4)- O -[β -D-glucopyranosyl-(1 \rightarrow 2)]- O - α -L-arabinopyranoside. This is the first known report of compound

4, which was named thyrsoilside A. MS, IR and NMR data comparison of compound **4** with **5**, indicated that **5** lacks an acetyl group. Thus, it was identified as davuricoside L, which has been previously reported in *L. davurica* [15].

The developed UPLC-ESI-MS/MS method was optimized for negative-ion mode, which is less dependent on the specific conditions of analysis. The chromatographic profile obtained by MRM analysis exhibited all the peaks corresponding to the compounds under investigation (Table 3). Two transitions per compound were used. For saponins **1**, **4** and **5**, one transition was the loss of formate from the formate adduct. For **5**, the second transition was the additional loss of CH_3OH from one of the terminal sugar moieties. In the case of compound **4**, the second transition was deacetylation leading to **5**. In the case of saponins **2** and **3**, the first transition was the loss of the xylopyranose moiety, and the second was the loss of the β -D-xylopyranosyl-(1 \rightarrow 2)- β -D-glucopyranose fragment.

Table 3: Optimized settings for quantitative analysis of saponosides from *Lysimachia thyrsoflora*.

Compound	Rt [min]	Transition	Cone potential [V]	Collision energy [eV]
chloramphenicol	3.95	320.85 \rightarrow 151.83	-32	22
		320.85 \rightarrow 256.93	-32	10
		795.55 \rightarrow 471.39	-32	64
α -hederin	6.39	795.55 \rightarrow 749.69	-32	30
		1137.64 [*] \rightarrow 1059.81	-32	42
5	3.17	1137.64 [*] \rightarrow 1091.80	-32	40
		1133.62 \rightarrow 1091.82	-148	50
4	3.80	1179.69 [*] \rightarrow 1133.72	-30	18
		1061.59 \rightarrow 767.53	-150	44
3	4.25	1061.59 \rightarrow 929.64	-150	42
		1059.59 \rightarrow 765.53	-150	44
2	4.97	1059.59 \rightarrow 927.64	-150	42
		927.51 \rightarrow 765.53	-114	32
1	5.08	973.51 [*] \rightarrow 927.65	-30	28

* Formate adduct

The calibration curve was constructed by plotting the peak area (y) versus concentration (x) by analyzing a set of 15 external standard (α -hederin) solutions. Linearity of response was confirmed in the range 0.1164 – 1.50 μ g/mL (y-intercept of the linear equation was statistically significant $p < 0.005$; for details see Table 4).

Table 4: Calibration data for α -hederin.

Calibration curve	Correlation coefficient (R^2)	Determination coefficient (R^2)	Linear range (μ g/mL)
$y = 0.4932x - 0.03339$	0.9925	0.9850	0.1164-1.50
	LOD (μ g/mL)	LOQ (μ g/mL)	Shapiro-Wilk test
	0.0384	0.1164	$p > 0.11$

After preliminary estimation of saponin concentrations in the analyzed material, appropriate dilution levels were selected and used. In order to quantify the saponin content, six biological samples, obtained from six individual plants were analyzed in triplicate. Table 5 reports the quantitative analysis results. The major constituents were triterpene saponins of the 13,28-epoxy-oleanane type (98.7%), compound **1** being the most abundant. The plant samples were collected from the same stand and so were fairly homogenous in terms of similarity due to the effect of growth conditions. To reflect plant-to-plant physiological differences, which are attributed to variations in their microenvironments or developmental stage, biological variability was quantified. The average coefficient of variation in *L. thyrsoflora* roots was 23%, and this value is similar to those obtained for other saponin accumulating plant species [16]. The RSD is also indicative of the effect of the several steps involved in the extraction/purification process, and is comparable with other studies [17].

Validation data showed good precision and intermediate precision, with a % RSD less than 8%. ANOVA test showed no statistically significant differences between different days ($p > 0.23$). In all the deliberately varied chromatographic conditions (flow rate, column temperature, mobile phase composition), the examined saponosides were adequately resolved, and the order of elution remained unchanged.

Table 5: Concentrations of individual and total saponins in the underground parts of *Lysimachia thyrsiflora* by MRM UPLC-ESI-MS/MS.

compound	concentration ^a μg/mg dry wt	SD	RSD (%)
1	0.60	± 0.247	41.4
2	0.56	± 0.307	54.2
3	0.013	± 0.0123	91.9
4	0.031	± 0.109	34.5
5	0.077	± 0.0288	37.5
Total	1.19	± 0.274	23.1

^aall concentrations were calculated based on peak areas of the [M-H]⁺ selected ion chromatograms relative to the standard α -hederin; each measurement was made in triplicate for six independent plant samples; abbreviations: μg/mg dry wt, μg per mg of dry weight; SD, standard deviation; RSD, relative standard deviation of saponin concentrations measured for six independent plants.

Cytotoxicity of compounds **4** and **5** was evaluated on human cancer cell lines differing in metastatic potential: melanoma BLM and A375, prostate cancer cell DU-145 and PC-3, and glioblastoma astrocytoma (U375), and normal cells of respective origin: skin fibroblasts (HSF) and prostate cells (PNT2). The results obtained (see details in Supplementary Data) showed that for both compounds EC₅₀ values were approximately 30 μg/mL. PC-3 proliferation was inhibited after 48 h at GI₅₀ 24 and 20 μg/mL, respectively. Compared with the reference drug, mitoxantrone, this activity was not significant. Our previous experiments proved that compounds bearing the 13 β ,28-epoxy bridge are much more potent against the cell lines mentioned above [6]. Disc diffusion assay for antifungal activity revealed no significant effect at the concentrations tested.

Experimental

General experimental procedures: Optical rotations were measured in MeOH at 20°C on a P-2000 polarimeter. IR spectra were recorded on a Nicolet iS5 spectrometer. NMR experiments (¹H, ¹³C, DEPT 135, ¹H-¹H DQF-COSY, ROESY, TOCSY, HMBC, HSQC) were performed on a Bruker AVANCE III 600MHz spectrometer (Bruker, Rheinstetten, Germany), using standard pulse sequences, in CD₃OD (99.95%, Sigma-Aldrich, St. Louis, USA). ¹H spectra were recorded at 600.20 MHz, and ¹³C at 150.94 MHz. All chemical shifts (δ) are given in ppm, and TMS was used as an internal standard. Coupling constants are reported in Hz. Spectra were analyzed using MestRec magnetic resonance companion version 4.4 and CARA 1.8.4 (Computer Aided Resonance Assignment). FAB-MS were recorded on a Finnigan MAT 95 mass spectrometer; glycerol as the matrix, Cs ions accelerated at 13keV. HR-ESI-TOF-MS were obtained on a Waters SYNAPT G2-S HDMS spectrometer. CC was carried out on Merck Kieselgel 60 (70 - 230 mesh). Preparative TLC was performed on commercially precoated silica gel G plates (Analtech, 500 μm). Analytical TLC was carried out on Merck silica gel 60 aluminum plates (Merck, Darmstadt, Germany) and the compounds were visualized by spraying with 5% H₂SO₄ in MeOH, followed by heating for 5-10 min at 105°C. All solvents for extraction and chromatographic separation were of analytical grade and purchased from Chempur (Piekary Śląskie, Poland). HPLC analysis of sugar derivatives was carried out on a Dionex apparatus equipped with a PDA 100 UV-VIS detector, on a 250 x 4.6 mm i.d. Hypersil GOLD C-18 (Thermo EC) column at 35°C with isocratic elution with 25% CH₃CN in 50

mM H₃PO₄ for 40 min, at a flow rate of 1 mL/min, and subsequent washing for 15 min. For UV spectroscopy, HPLC-grade acetonitrile, HPLC-grade acetonitrile, methanol and H₃PO₄ were from J.T. Baker (Phillipsburg, USA). Formic acid puriss. p. a. and chloramphenicol ($\geq 98\%$, TLC) were from Sigma-Aldrich (St. Louis, USA). Standard α -hederin (ROTICROM HPLC) was from Roth (Karlsruhe, Germany). Standard sugars, D-glucose, L-glucose, D-arabinose, and L-arabinose were from Sigma, and L-cysteine methyl ester hydrochloride (puriss >99%) and *o*-tolyl-isothiocyanate from Fluka. HPLC grade water was obtained from HLP 5 (HYDROLAB, Poland) apparatus and was filtered through a 0.2 μm filter before use. Bakerbond C18 POLAR+ (1000 mg, 6 mL) cartridges were from J.T. Baker (Phillipsburg, USA).

Plant material: The underground parts of *Lysimachia thyrsiflora* L. were collected from a controlled cultivation in the Garden of Medicinal Plants, Jagiellonian University, Cracow, Poland. Samples were identified by Dr Bożena Muszyńska, Department of Pharmaceutical Botany, Pharmaceutical Faculty, Medical College, Jagiellonian University, Cracow, Poland. A voucher specimen (No. KFG/2011021) is deposited at the Department of Pharmacognosy, Pharmaceutical Faculty, Medical College, Jagiellonian University, Cracow, Poland. The mixture of *L. thyrsiflora* saponins, isolated previously [5,6], was used to develop the UPLC-MS method, to optimize chromatographic conditions, and for the specificity and system suitability tests.

Extraction and isolation: The dried and powdered plant material (940 g) was extracted with CHCl₃ (3 times) and then exhaustively with MeOH containing 0.5% pyridine at 40°C. The combined MeOH extract was concentrated *in vacuo*, to yield a viscous residue (120 g), which was then partitioned between *n*-BuOH and water. The *n*-BuOH-soluble fraction was evaporated to dryness below 45°C and chromatographed on a silica-gel column (Merck Kieselgel 60; 70-230 mesh) with CHCl₃-MeOH-H₂O (23:12:2). Fractions were combined on the basis of TLC (SiO₂, CHCl₃-MeOH-H₂O 23:12:2; sulfuric acid + heating 5-10 min, 105°C) to give 7 major fractions. The pooled saponin fraction with *R_f* values 0.27 and 0.2 (fraction 7, 1500 mg) was further purified by repeated prep. TLC (commercially precoated silica gel G plates; Analtech, 500 microns) using CHCl₃-MeOH-H₂O (23:12:2; 8:7:1). Bands were removed from plates after spraying with water, the silica was extracted with MeOH, and solvent was evaporated *in vacuo* to yield compounds **4** (25 mg) and **5** (38 mg).

Acid hydrolysis: Identification of sugar units was established after total acid hydrolysis of **4** and **5** on a TLC plate with gaseous HCl for 25 min. Two monosaccharides were identified by TLC comparison with authentic samples (CHCl₃-MeOH-H₂O 23:12:2, twice; visualization: aniline phthalate in *n*-BuOH + heating) as glucose and arabinose. Absolute configurations were determined according to a slightly modified method, previously reported by Tanaka *et al.* [12]. Compounds **4** and **5** (5 mg) were dissolved in 1.0 mL of a mixture of 4 M HCl in dioxane and water (1:1; v/v) and heated at 95°C for 3 h. The saponins were extracted with EtOAc (3 x 3.0 mL) and the acid aqueous phases were neutralized with 2M ammonium hydride and concentrated under vacuum. After drying over P₂O₅ for 48 h, the residues were dissolved in anhydrous pyridine (1.0 mL) in oven-dried screw-capped vials purged with argon. Then, L-cysteine methyl ester hydrochloride (5 mg) was added, the vials were again purged with argon, and the mixtures were allowed to react at 60°C for 1 h. Next, 5 μL of *o*-tolyl-isothiocyanate was added and the reaction mixtures were heated at 60°C for an additional 1 h. Samples were directly analyzed by HPLC and the peaks were detected at 14.09 min and 14.9 min for

monosaccharide derivatives of **4**, and at 14.1 min and 15.0 min for monosaccharide derivatives of **5**. The retention times were compared with authentic sugars derivatized according to the same procedure: D-glucose (14.1 min), L-arabinose (14.9 min).

30-O- β -D-Glucopyranosyl-3 β ,16 α ,30-trihydroxy-olean-12-en-28-yl acetate 3-O- β -D-glucopyranosyl-(1 \rightarrow 4)-O-[β -D-glucopyranosyl-(1 \rightarrow 2)]-O- α -L-arabinopyranoside (4**)**

Amorphous white solid.

$[\alpha]_D^{20}$: -4.5 (c 0.4, MeOH).

IR (KBr): 3333 (>OH), 2922 (>CH), 1737 (ester), 1714 (C=O), 1255 and 1033 (>C-O-C) cm^{-1} .

^1H and ^{13}C NMR (CD_3OD , 600 MHz) data of the aglycone moiety and the sugar portion: Tables 1 and 2, respectively.

FAB-MS (negative mode) m/z 1133.4 $[\text{M} - \text{H}]^-$; 971.4 $[\text{M} - \text{H} - 162]^-$, 809.3 $[\text{M} - \text{H} - 162 - 162]^-$

HR-ESI-TOF-MS $[\text{M} + \text{Na}]^+ m/z$ 1157.57 calcd for $\text{C}_{55}\text{H}_{90}\text{O}_{24}\text{Na}$.

Davuricoside L (5**)**

Amorphous white solid.

$[\alpha]_{D20}^{20}$: +13.3 (c 0.4, MeOH).

IR (KBr): 3319 (>OH), 2920 (>CH), 1255 and 1033 (>C-O-C) cm^{-1} .

^1H and ^{13}C NMR (CD_3OD , 600 MHz) data of the aglycone moiety and the sugar portion: Tables 1 and 2, respectively.

FAB-MS (negative mode) m/z 1091.3 $[\text{M} - \text{H}]^-$; 929.3 $[\text{M} - \text{H} - 162]^-$

HR-ESI-TOF-MS $[\text{M} + \text{Na}]^+ m/z$ 1115.56 calcd for $\text{C}_{53}\text{H}_{88}\text{O}_{23}\text{Na}$.

Semiquantitative UPLC-ESI-MS/MS analysis: Preparation of standard solutions: Chloramphenicol (10 mg), as internal standard, was placed in a volumetric flask and the volume brought to 10 mL using MeOH. One mL of this solution was subsequently diluted to make a 100 $\mu\text{g/mL}$ stock solution. α -Hederin (1 mg), as external standard, was added to a volumetric flask and the volume brought to 10 mL with MeOH to make a 100 $\mu\text{g/mL}$ solution. One mL of this solution was subsequently diluted to make a 10 $\mu\text{g/mL}$ stock solution. These solutions were stored at -10°C .

Preparation of calibration samples: To 10 μL of chloramphenicol internal standard stock solution 0 – 150 μL of α -hederin reference stock solution was added and diluted with water to 1 mL making calibration samples with a concentration of chloramphenicol of 1 $\mu\text{g/mL}$ and of α -hederin in the range from 0 – 1.5 $\mu\text{g/mL}$.

Plant samples preparation: Dried, finely ground plant material (0.5 g) from 6 individual plants was extracted with CHCl_3 (2 x 20 mL, for 2 h), and then with MeOH (2 x 20 mL, for 2 h) on a boiling water bath under reflux. The MeOH extracts were combined and the solvent removed under reduced pressure. The residues were redissolved in 8 mL of H_2O , and 2 mL aliquots were cleaned up using SPE C_{18} cartridges, preconditioned with MeOH and H_2O . The cartridge was washed first with 6 mL H_2O , then with 18 mL 30% MeOH. Finally, saponins were eluted with 18 mL MeOH. These fractions were evaporated under reduced pressure, and redissolved in 10 mL MeOH making stock sample solutions. From each stock sample solution, 20 μL , 50 μL , 100 μL and 200 μL were taken, and then 10 μL of the chloramphenicol standard solution was added and brought to 1 mL with H_2O , making a series of sample dilutions. Each dilution was analyzed by UPLC-MS/MS.

Quantification: The UPLC-MS/MS system consisted of a Waters ACQUITY® UPLC® (Waters Corporation, Milford, MA, USA) coupled to a Waters TQD mass spectrometer (electrospray ionization mode ESI-tandem quadrupole). Chromatographic separations were carried out using an Acquity UPLC BEH (bridged

ethyl hybrid) C_{18} column, 2.1×100 mm, and 1.7 μm particle size. The column was maintained at 40°C , and eluted under linear gradient conditions from 95% to 0% of eluent A over 10 min, at a flow rate of 0.3 mL/min. Eluent A: water/formic acid (0.1%, v/v); eluent B: acetonitrile/formic acid (0.1%, v/v). Ten μL aliquots of each sample were injected in triplicate. A Waters TQD mass spectrometer was calibrated for quantitative analysis using α -hederin solution at a concentration of 10 $\mu\text{g/mL}$ at a flow rate 20 $\mu\text{L/min}$, and a mixture of eluent A and B (1:1, v/v) at a flow rate of 0.28 mL/min. Optimized settings were as follows: source temperature 150°C , desolvation temperature 350°C , desolvation gas flow rate 600 L/h, capillary potential -4.00 kV, collision gas flow 0.1 mL/min. Cone potential and collision energy were individually optimized for each transition (Table 3). Nitrogen was used as both nebulizing and drying gas, and argon as collision gas. Saponosides were analyzed using the MRM (Multiple Reaction Monitoring) method. All analytical data were processed using MassLynx V4.1 software (Waters Corporation, Milford, MA, USA).

Method validation: The described method was validated for the determination of α -hederin and the mixture of saponins **1-5** by the UPLC method according to the ICH guidelines [18]. **Specificity:** The solution containing all investigated saponosides (mixture) was analyzed. **System suitability:** The system suitability parameters were defined with respect to peak resolution of examined compounds, using a mixture of **1-5**. **Linearity:** The linearity for α -hederin was assessed by injecting 15 separately prepared solutions covering the range of 0.10 – 1.50 $\mu\text{g/mL}$. The slope of the weighted regression line (weight 1/x), y-intercept, standard deviations of slope and intercept, correlation coefficient, R^2 value and standard error of residuals of the calibration curve were calculated. Next, to determine whether the residuals have normal distribution, the Shapiro-Wilk statistical test was used. **Limit of detection (LOD) and limit of quantification (LOQ):** Based on the standard error of residuals (Se) and the slope (a) of the calibration plots and following the formula $\text{LOD} = 3.3\text{Se}/a$ and $\text{LOQ} = 10\text{Se}/a$, the LOD and LOQ for α -hederin were estimated. **Precision:** The repeatability of the method was checked by a six-fold analysis of the concentration level at 0.7 $\mu\text{g/mL}$ of α -hederin solution. The same protocol was followed for 3 different days to study the intermediate precision of the proposed method. The RSD (%) of the peak area of α -hederin was calculated. Statistical significance of interday differences was tested with ANOVA. **Robustness:** To demonstrate the robustness of the method deliberate small changes of flow rate, content of acetonitrile and column temperature were made around the optimal values. The mobile phase flow rate was 0.30 mL/min; to study the effect of the flow rate on resolution, the flow rate was changed to 0.27 and 0.33 mL/min. The effect of the column temperature was studied at 36°C and 44°C (instead of 40°C), and the mobile phase composition was changed +5% from the initial composition.

Cytotoxicity assay: The human glioblastoma-astrocytoma cell line U375 was cultured in Eagle's minimum essential medium (EMEM) with 2 mM L-glutamine, 1% non-essential amino acids (NEAA), 1 mM sodium pyruvate and 10% fetal bovine serum (FBS). Prostate cancer cells DU-145 and PC3 were cultured in Dulbecco's Modified Eagle's Medium (DMEM-F12 HAM) supplemented with 10% FBS. Malignant melanoma cell line A375 was cultured in DMEM high glucose medium supplemented with 10% FBS and 2 mM L-glutamine, whereas cell line BLM was cultured in RPMI medium supplemented with 10% FBS. Normal prostate cells PNT2 were cultured in RPMI-1640 supplemented with 10% FBS. Human skin fibroblasts (HSF) were cultured in DMEM supplemented with 10% FBS. All reagents were from Sigma-Aldrich, St. Louis, MO,

USA. All cells were cultured in medium containing 100 i.u./mL penicillin, 10 µg/mL streptomycin, and 10 µg/mL neomycin, in standard conditions, a humidified atmosphere at 37°C in the presence of 5% CO₂. Cytotoxicity assay was performed as described previously [10]. In each experiment, for a positive control, mitoxantrone (Ebewe Pharma) was used, at concentrations from 0.1 to 1 µg/mL. For each value measured, 300 cells were analyzed. Each experiment was performed in triplicate. Each variable was expressed as the mean (± SD). The statistical significance was determined using either the Student's *t*-test or the non-parametric Mann-Whitney *U*-test, with *p* < 0.05 considered to indicate significant differences.

Anti-proliferative assay: The assay was performed on prostate cancer cells DU-145 and PC3, and on normal prostate cells PNT2, cultured at conditions described above. Cells were seeded into 24-well plates at an initial density of 7.6×10² cells per well and incubated for 24 h at 37°C. Next, the medium was replaced with either fresh culture medium (control), or with the same medium containing different concentrations (from 0 up to 30 µg/mL) of saponosides **4** and **5**. After incubation for various times (24, 48, 72 h) the medium was removed, the cells trypsinized and the number of cells determined using a cell counter (Coulter Beckman ZS).

Antimicrobial assay: The potential antimicrobial properties of compounds **4** and **5** were investigated by the conventional paper disc diffusion method against the following strains: *Candida*

albicans, *C. glabrata*, *C. krusei*, *C. parapsilosis*, *C. lusitanae*, *Cryptococcus neoformans*, and *Trichophyton mentagrophytes*. All strains were the isolates of the Department of Pharmaceutical Microbiology, Pharmaceutical Faculty, Medical College, Jagiellonian University. Sabouraud broth was used as media for all strains except *C. neoformans*, which was cultured in Casitone broth. Strain suspensions in 0.85% NaCl (100 µL) from 24 h incubations were applied to 9 cm Petri dishes with Sabouraud agar medium. For the compounds being tested, 1 mg was dissolved in 1 mL DMSO. Five µL aliquots of these solutions were applied to paper discs and placed onto the agar plates. The solvent was used as control. The plates were incubated at 30°C for 24 h, and the growth inhibition was assessed based on the dimensions of the lytic zones on the plates.

Supplementary data: Details on the cytotoxicity and antiproliferative assays. ¹H, ¹³C, DEPT, HSQC, HMBC, ROESY, TOCSY, and COSY spectra for compound **4** and **5**.

Acknowledgments - We acknowledge the financial support of grant N N405 297636 from the Ministry of Science and Higher Education, Poland. We wish to thank Dr Bożena Muszyńska for identification of the plant material, and Dr Edward Szneler for recording the NMR spectra. The authors are much indebted to Dr Danuta Trojanowska from the Department of Pharmaceutical Microbiology, Pharmaceutical Faculty, Medical College, Jagiellonian University, for carrying out the antimicrobial activity tests.

References

- [1] Foubert K, Theunis M, Apers S, Vlietinck AJ, Pieters L. (2008) Chemistry, distribution and biological activities of 13,28-epoxy-oleanane saponins from the plant families Myrsinaceae and Primulaceae. *Current Organic Chemistry*, **12**, 629-642.
- [2] Källersjö M, Bergqvist G, Anderberg A. (2000) Generic realignment in primuloid families of the Ericales s.l.: a phylogenetic analysis based on DNA sequences from three chloroplast genes and morphology. *American Journal of Botany*, **87**, 1325-1341.
- [3] Karpova VI, Kintya PK, Chirva VY. (1975) Structure of saponin A from *Naumburgia thyrsoiflora*. *Khimiya Prirodnykh Soedinenii*, **11**, 364-366.
- [4] Kintya PK, Karpova VI, Chirva VY. (1975) Saponin from *Naumburgia thyrsoiflora*. *Khimiya Prirodnykh Soedinenii*, **11**, 520-521.
- [5] Podolak I, Janeczko Z, Galanty A, Michalik M, Trojanowska D. (2007) A triterpene saponin from *Lysimachia thyrsoiflora* L. *Acta Poloniae Pharmaceutica – Drug Research*, **64**, 39-43.
- [6] Podolak I, Koczurkiewicz P, Galanty A, Michalik M. (2013) Cytotoxic triterpene saponins from the underground parts of six *Lysimachia* L. species. *Biochemical Systematics & Ecology*, **47**, 116-120.
- [7] Çaliş I, Şatana ME, Yürüker A, Kelican P, Demirdamar R, Alaçam R, Tanker N, Rüegger H, Sticher O. (1997) Triterpene saponins from *Cyclamen mirabile* and their biological activities. *Journal of Natural Products*, **60**, 315-318.
- [8] Jansakul C, Baumann H, Kenne L, Samuelsson G. (1987) Ardisiacrispin A and B, two utero-contracting saponins from *Ardisia crispa*. *Planta Medica*, **53**, 405-409.
- [9] Dong W, Liu X, Li X, Yang D, Ding L. (2011) A new triterpene saponin from *Androsace integra*. *Fitoterapia*, **82**, 782-785.
- [10] Galanty A, Michalik M, Sędek Ł, Podolak I. (2008) The influence of LTS-4, a saponoside from *Lysimachia thyrsoiflora* L., on human skin fibroblasts and human melanoma cells. *Cellular & Molecular Biology Letters*, **13**, 585-598.
- [11] Mahato SB, Kundu AP. (1994) ¹³C NMR spectra of pentacyclic triterpenoids – a compilation and some salient features. *Phytochemistry*, **37**, 1517-1575.
- [12] Tanaka T, Nakashima T, Ueda T, Tomii K, Kouno I. (2007) Facile discrimination of aldose enantiomers by reversed-phase HPLC. *Chemical & Pharmaceutical Bulletin*, **55**, 899-901.
- [13] Agrawal PK. (1992) NMR spectroscopy in the structural elucidation of oligosaccharides and glycosides. *Phytochemistry*, **31**, 3307-3330.
- [14] De Tommasi N, Piacente S, De Simone F, Pizzi C. (1993) Characterization of three new triterpenoid saponins from *Ardisia japonica*. *Journal of Natural Products*, **56**, 1669-1675.
- [15] Tian JK, Zou ZM, Xu LZ, Tu GZ, Zhang HW, Yang SL, An DG. (2005) Two new triterpene saponins from *Lysimachia davurica*. *Journal of Asian Natural Product Research*, **7**, 601-606.
- [16] Huhman DV, Berhow MA, Sumner LW. (2005) Quantification of saponins in aerial and subterranean tissues of *Medicago truncatula*. *Journal of Agricultural & Food Chemistry*, **53**, 1914-1920.
- [17] Mroczek A, Kapusta I, Janda B, Janiszowska W. (2012) Triterpene saponin content in the roots of red beet (*Beta vulgaris* L.) cultivars. *Journal of Agricultural & Food Chemistry*, **60**, 12397-12402.
- [18] ICH-Q2 (R1) Validation and Analytical Procedures: Text and Methodology. International Conference on Harmonization: Geneva, November 2005, http://www.ich.org/fileadmin/Public_Web_Site/ICH_Products/Guidelines/Quality/Q2_R1/Step4/Q2_R1_Guideline.pdf

Variation of Saponin Contents and Physiological Status in *Quillaja saponaria* Under Different Environmental Conditions

Angélica Grandón S^{a*}, Miguel Espinosa B^a, Darcy Ríos L^a, Manuel Sánchez O^a,
Katia Sáez C^c, Víctor Hernández S^b and José Becerra A^b

^aDepartment of Silviculture, Faculty of Forest Science. Universidad de Concepción. 4070386, Concepción. Chile

^bLaboratory of Chemistry of Natural Products. Faculty of Natural and Oceanographic Sciences. Universidad de Concepción. 4070386, Concepción. Chile

^cDepartment of Statistics, Faculty of Physics and Mathematics Sciences. Universidad de Concepción. 4070386, Concepción. Chile

agrandon@udec.cl

Received: July 24th, 2013; Accepted: August 22nd, 2013

Quillaja saponaria (Quillay), an evergreen tree found in Chile, is one of the main sources of saponins. *Quillaja* saponins have hypocholesterolaemic, anticarcinogenic, antioxidant and pesticidal properties, and are used as adjuvants for vaccines. Samples of Quillay growing at three zones in O'Higgins Region, Chile (Coastal, Central and Mountain zones) were analyzed for content of saponins and physiological status. The results revealed differences in the content of saponins depending on the zone of sample collection. The highest contents were found in samples from the Mountain zone, where the highest saponin contents were accompanied by the lowest foliar nitrogen contents, the highest antioxidant activity and the highest carotenoid contents. The results suggest a physiological and adaptive mechanism of saponins in plants to survive under unfavourable environmental conditions. The results have important implications for a theoretical basis for the design of a reasonable harvest, to avoid the cost of poor quality material, and also to provide a sustainable use and conservation of this important species. Further research on the effects of stress will improve our understanding of the saponins production and their physiological functions in plants, whereas they have generally been studied for their biological and chemical applications.

Keywords: Saponins, Triterpenes, *Quillaja saponaria*, Secondary metabolites, Glycosides.

The Chilean native tree *Quillaja saponaria* Mol. (Quillay), distributed under different geographical and climatic conditions, is one of the most important sources of triterpene saponins [1a,b]. Structurally, the *Quillaja* saponins are referred to as triterpene glycosides, consisting of an aglycone of quillaic acid (hydrophobic region) and several sugars, such as rhamnose, fucose, xylose, arabinose, glucuronic acid and galactose (hydrophilic region) (Figure 1). An extract of *Quillaja* saponins is a heterogeneous mixture, and approximately 70 saponins have been reported [2a].




Figure 1: Basic structure of *Quillaja* saponins. The aglycone is usually the triterpene quillaic acid, with a branched trisaccharide or a disaccharide linked to the 3-position (R^1). The ester linked fucosyl residue at C-28 is substituted by an oligosaccharide in the 2-position (R^2), by a monosaccharide residue or an acyl group in the 3-position (R^3) and by an acyl group in the 4-position (R^4) [2b].

Due to their biological and chemical properties, such as antibacterial, antifungal, insecticidal, hypocholesterolaemic, anticarcinogenic, antioxidant, haemolytic, antioxidant, adjuvants for vaccines, foaming, emulsifying, surfactant and tensoactive [3a,b], they are generally considered to be part of the defensive system of the plants [1a]. *Quillaja* saponins have industrial applications as, for instance, beverage, alimentary, cosmetic, pharmaceutical, medicinal, pesticidal, photographic and mining products [1b,4].

Quillaja saponins are obtained by aqueous extraction of bark (traditional process) and biomass obtained from forest pruning. The industrial use of saponin causes high demand for biomass, but it is also accompanied by sub-utilization of this resource [5] due to the variation of saponin contents in Quillay. This variation has generally been attributed to genetic factors, edaphic conditions, altitude, seasonal effects, age of vegetation, and kind of vegetal tissue [6a]. However, considering the diversity of environments in which Quillay grows, it is possible that the contents of saponins and the physiological status of the trees could be influenced by the surrounding environment. The growing demand for *Quillaja* saponins requires rationalization of the harvesting of existing natural resources to avoid their improper exploitation and sub-utilization.

There has been some research on the effects of abiotic environmental factors on saponin contents, but the majority of published studies have been focused on non woody plants with short life cycles, such as *Panax quinquefolius*, *P. ginseng*, *Asparagus racemosus*, *Glycyrrhiza uralensis*, *Phytolacca dodecandra*, *Bupleurum chinense*, *Clematis chilensis*, *C. hexapetala*, and *Tribulus terrestris*, among others [6b]. With respect to Quillay, over the last 10 years, significant efforts have been made in qualitative and quantitative analysis of the saponins, with the aim of increasing knowledge for commercial applications [1a,3a,b]. However, much less data are available concerning variation of saponin contents in woody species under natural conditions due to the complexity involved, where two or more stressors co-occur and their effects are sometimes additive. Nevertheless, research is necessary on the influence of external conditions on saponin

accumulation in plants that should lead to a better understanding of saponin production in natural conditions. Our study examined the variation of total saponin contents and physiological status in *Quillaja saponaria* growing under different environmental conditions in the O'Higgins Region of Chile (Coastal, Central and Mountain zones). Although these zones are located at the center of the species' native range, studies from these specific regions have not previously been reported. In order to achieve these objectives, the zones were characterized at the edaphic level for available water capacity (AWC), nitrogen (N), organic matter (OM), phosphorus (P) and potassium (K). The trees were analyzed for total saponin contents, and physiological status: photosynthetic pigments, antioxidant activity, nitrogen, phosphorus and potassium.

Edaphic characterization: The edaphic samples from the different zones were significantly different for OM, AWC and P. The Mountain zone showed the highest contents of K, AWC and OM. The content of OM for this zone was 13.9%, 2.9 and 1.7 times higher than the Coastal and Central zones, respectively. The AWC was 22.3% and 16.7% for Mountain and Central zones, respectively, both significantly different when compared with the Coastal zone. K content was significantly higher in the Mountain zone with 655.2 cmol/kg, 2.1 and 3.6 times greater than the Central and Coastal zones, respectively (Table 1). However, in spite of these results, this zone is characterized by presenting harsher environmental conditions than the Coastal and Central zones. The spatio-edaphic pattern of the present study identified three differentiable zones, as shown by PCA. The first and second principal components accounted for 39.9% and 29.6% of the total variation, respectively. Both components explain 70% of the variance. N explained the Coastal zone, N and P the Central zone, and OM, AW and K the Mountain zone. These specific and inherent characteristics of the zone within the Region may explain differences in the physiological status of plants, which is influenced by environmental conditions (Figure 2)

Table 1: Nitrogen (N), phosphorous (P), potassium (K), available water capacity (AWC) and organic matter (OM) from sampling zones of the O'Higgins Region. Mean values \pm S.D.

Variable	Coastal zone	Central zone	Mountain zone	P-value
N (N-NO ₃) (mg/kg)	13.9 \pm 10.0	14.0 \pm 9.0	7.8 \pm 5.2	0.2377
P (mg/kg)	25.3 \pm 6.3	33.6 \pm 21.7	26.1 \pm 9.1	0.4427
K (cmol/ kg)	255.9 \pm 133.0 ^b	315.1 \pm 126.4 ^b	655.2 \pm 266.8 ^a	0.0006
AWC (%)	7.9 \pm 1.6 ^b	16.7 \pm 4.6 ^a	22.3 \pm 13.2 ^a	0.0113
OM (%)	4.8 \pm 1.0 ^b	8.1 \pm 3.9 ^b	13.9 \pm 4.1 ^a	<0.0001

In each column, different letters are significantly different for 95% confidence intervals ($P < 0.05$). Fisher's LSD test.




Figure 2: Biplot of soil samples distributed in Coastal, Central and Mountain zones from O'Higgins region, against values for the first two principal components, PC1 and PC2, for edaphic variables: organic matter (OM), available water (AWC), nitrogen (N), phosphorous (P) and potassium (K). Vectors represent edaphic variables that have a high loading on PC1 and PC2 and that are significantly correlated with the ordination axes ($P < 0.05$).

Physiological status: The nutritional status of *Quillaja* originating from Coastal, Central and Mountain zones was significantly different for N and P. Samples from the Mountain zone showed the lowest N content (0.69%) and the Central zone the highest (1.04%). P was higher in the Coastal zone, with 0.12%, than the Central zone (0.09%). For K content, there was no significant difference among zones (Figure 3A). These results have been consistent across the availability of nutritional resources in each zone. Considering that the O'Higgins Region presents a scarce water pluviometric regime during the months of vegetative development, this contributes to explaining the development and growth of sclerophyll species by means of adaptive and efficient strategies for the use of water and nutritional resources in these difficult Mediterranean climates [7]. Thus the plants can mitigate the physiological process with a trade-off between growth, maintenance, storage, reproduction and defense.

The chlorophyll a and b contents were not significantly different. Carotenoids increased significantly from the Coastal to the Mountain zones, reaching a mean value of 0.34 mg/g, 2.4 and 1.6 times higher than the Coastal and Central zones, respectively (Figure 3B). The highest carotenoid content found in the Mountain zone had been associated with defensive responses in plants under stress conditions, such as water stress and/or radiation; carotenoids have antioxidant properties and a photo-protective function of the photosynthetic system against harmful photooxidative processes [8].




Figure 3: A. Foliar NPK contents in Coastal, Central and Mountain zones. Any variable not having a letter in common is significantly different at $P < 0.05$. Bars represent mean values \pm S.E. B. Pigment concentration: chlorophyll a (Chl a), chlorophyll b (Chl b) carotenoids (car) in Coastal, Central and Mountain zones. Any variable not having a letter in common is significantly different at $P < 0.05$. Bars represent mean values \pm S.E. C. Antioxidant activity in Coastal, Central and Mountain zones. Any zone not having a letter in common is significantly different at $P < 0.05$. Bars represent mean values \pm S.E. D. Saponins production in Coastal, Central and Mountain zones. Any zone not having a letter in common is significantly different at $P < 0.05$. Bars represent mean values \pm S.E.

The antioxidant activity was significantly higher in samples from the Mountain zone compared with those from the Coastal and Central zones (Figure 3C). This implies a defensive mechanism observed in alpine plants subjected to environmental stressors, such as solar radiation and low temperatures, which may activate the formation of chemical reactive oxygen species (e.g., hydrogen peroxide, superoxide and hydroxyl) conducive to oxidative stress [9a]. Under these conditions it has been seen that plants generate antioxidant compounds, intended to entrap radicals harmful to their cells, resulting in a broad adaptation spectrum for the plants [9b]. The high antioxidant activity of Quilajja, over 85%, makes this species interesting to study for the production of antioxidants.

Saponin contents: Saponin contents of Quilajja were different between zones, increasing significantly from the Coastal to the Mountain zone. The highest saponin contents were found in samples from the Mountain zone, where the content was 22.1 mg/g; 1.7 and 1.3 times higher than those of the Coastal and Central zones, respectively (Figure 3D). The saponin contents, found in the range between 1.2% and 2.2% of the dry weight, contrast with values close to 8% of soluble compounds found by San Martín and Briones [6d], and between 6.5% and 15.8% dry weight found by Copaja *et al.* [6c].

Associating saponin contents and physiological status, our study revealed, for the first time, that the highest saponin contents were accompanied by the lowest nitrogen content, and the highest antioxidant activity and carotenoids contents as a synergistic response. This finding showed that higher accumulation of saponins in Quilajja would be stimulated by adverse environmental conditions. Therefore, an evaluation of quality and saponin composition are necessary in future research. The presented results suggest a physiological and adaptive mechanism of saponins in plants to survive under unfavourable environmental conditions.

This study was a first approach and contribution to the knowledge about the variation of saponin contents and physiological status in *Q. saponaria* under different environmental conditions. Further research on the effects of stress will improve our understanding of saponin production and the physiological functions of saponins in plants. Previously, they have generally been studied for their biological and chemical applications.

Experimental

Zones of study: Sampling took place at 3 zones in the O'Higgins Region: Pumanque (Coastal zone, 34° 56' S and 71° 54' 78" W; altitude 180 m), Roblería del Cobre de Loncha National Reserve (Central zone, 34° 12' S and 71° 10' W; altitude 430 m) and Río de los Cipreses National Reserve (Mountain zone, 34° 31' S and 70° 45' W; altitude 1250 m). Pumanque has a Mediterranean climate, with a drought period of 6-7 months and a semiarid period of 1-2 months. Annual average temperature is 14.5°C, seasonal average temperature is 28°C in summer and 3.7°C in winter. The precipitation regimen varies between 500 to 700 mm annually. Roblería del Cobre de Loncha National Reserve presents a warm mild climate, with a prolonged dry season. Maximum annual temperatures oscillate between 25°C to 28°C and the minimum is between 10°C to 8°C. Río de los Cipreses National Reserve has a mild Mountain climate, with 4 months of dry warm weather in summer, and winter with rain and snow precipitation between 800 to 1500 mm annually. The temperature varies between a maximum of 19°C to 21°C and a minimum of 7°C to 5°C, with temperatures under 0°C due to thermal oscillations in winter [10].

Sampling and nutritional analysis of soil: In every zone, 8 randomly distributed soil samples were taken from the surface, at a depth of 20 cm, near trees. One third of the soil sample was used for nutritional analysis, the second third for field capacity, permanent wilting point and available water capacity, and the last third for organic matter content; every soil sample was replicated 3 times. For nutritional analysis, 33 g of dry and sifted soil was analyzed for nitrate (N-NO₃), phosphorous (P) and potassium (K). N-NO₃ was determined through the Kjeldahl and direct Nesslerization Semichrome method. N-NO₃ detection was carried out by spectrophotometric absorbance measurement at a wavelength of 490 nm [11]. P was obtained by Olsen's method, extracting P in 0.5 mol/L NaHCO₃ at pH 8.5. Then, the extract was analyzed by colorimetry using the ethylene blue method with ascorbic acid as reductor, and detected by spectrophotometric absorption at 880 nm. K was extracted in 1 mol/L NH₄C₂H₃O₂ solution at pH 7.0 and detected by Atomic Spectrophotometry Emission (ASE) with either an acetylene-air or propane-air flame, measured at 766.5 nm [12].

Field capacity, permanent wilting point and available water capacity: Field capacity (FC), permanent wilting point (PWP) and percentage of soil available water capacity (AWC) were obtained as follows: The FC was determined by the pressure cooker method [13]. In brief, cylinders filled with soil were hydrated with distilled water for 24 h. Then, they were placed in a pressure cooker under 0.33 and 15 bar for 24 h, in order to obtain wet weight. All samples were later oven dried at 105°C for 24 h to determine dry weight. The relative percentage of AWC at FC and PWP was obtained by difference between wet and dry weight at 0.33 bar and 15 bar, respectively. AWC was obtained by gravimetric weight difference between FC and PWP.

Organic matter content of soil: Organic Matter Content (OM) was determined by Weight Loss on Ignition method (WLOI) [14]. Briefly, an aliquot of 10 g soil was oven dried at 105°C for 24 h to determine dry weight. Then, the samples were calcined in a muffle furnace at 450°C for 24 h, to obtain calcinated weight. Organic matter percentage was obtained by the relational differences of the dry and calcinated soil weight and dry soil weight.

Plant material: In September 2011, 15 trees were sampled in each study zone, with diameters between 70 to 100 cm. For each tree, 10 cm² of bark was extracted at breast height to determine the total saponin contents, and 300 g of foliar biomass was extracted from the middle crown for nutritional analysis, antioxidant activity and pigment quantification.

Foliar nutritional analysis: Nutritional analysis was performed for nitrogen (N), phosphorous (P) and potassium (K). N was obtained through a digestion method; 3 g dried sample was put in H₂SO₄, 98% of density 1.84 Kg/L. The NH₄ concentration was obtained by distillation of NH₃ and manual titration [15]. For P, 3 g of dried sample was passed through a 1 mm sieve. Then it was put in a crucible and placed in a muffle furnace at 500°C for 4-8 h. Once cooled, 10 mL of HCl at 2 mol/L was added and boiled on a hot plate. The content in the crucible was filtered and P detection was determined through colorimetry using the phospho-vanadomolybdate method, measured at 466 nm. K was determined by absorption spectrophotometry and atomic emission with an air acetylene flame measured at 766.5 nm [15].

Pigment quantification: Two g of leaves were extracted in acetone/water (8/2,v/v), and chlorophyll a, chlorophyll b, and carotenoids contents were determined by absorption spectrophotometry at 663 nm, 646 nm and 470 nm, respectively [16].

Antioxidant activity: The antioxidant activity was determined using the DPPH (1,1-diphenyl-2-picrylhydrazyl free radical) scavenging method [17]. Briefly, 2 g of leaves were extracted in 50 mL methanol: water (7/3, v/v) for 24 h at room temperature. The extract was resuspended in ethanol (500 μ L), mixed with deionised water in equal volumes of DPPH (60 μ M), and homogenized in a vortex mixer. Detection was performed by absorbance at 520 nm (20 min) in a spectrophotometer (model UV-VIS 2310). The antioxidant activity, based on the DPPH free radical scavenging ability of the extract, was determined through comparison of the absorbance with the control (100%) containing only DPPH.

Extraction and quantification of saponins: Two g of dried bark was extracted in 50 mL MeOH: H₂O (7/3, v/v) for 24 h at room temperature. The extract was filtered through a 0.2 micron filter and kept frozen at -18°C, before analysis. Saponins were detected by HPLC (Shimadzu, 10 AD-VP, Japan), with a Vydac C-4 column (250 x 4.6 mm) and a 30-45% acetonitrile in water gradient, with 0.15% trifluoroacetic acid. The flow rate was 1 mL min⁻¹ and detection was measured at 210 nm. [6a]. The total saponin contents (dry wt) were determined by comparison with the standard calibration curve, using the standard Quillaja Liquid Ultra

(QL Ultra) of Natural Response S.A. The standard was diluted to different concentrations to obtain the best linearity, with R²=0.9.

Statistical analysis: A parametric variance analysis (ANOVA) was performed for edaphic and physiological variables, every sample being replicated 3 times (n=3). Differences were detected with Tukey multiple comparisons test. The existence of groups of variables by zone was achieved with multivariate analysis of principal components (PCA), which identified groups of samples based on the measured edaphic variables and how these relate to the observed differences between the groups classified by zone. Distributional assumptions of both normality and heterogeneity of variance was analyzed with Shapiro-Wilk and Levene tests, respectively; differences were considered statistically significant if $P < 0.05$ for the test statistic. Analyses were performed with software SAS (SAS Institute Inc, 2003) and Infostat version 2012 [18a,b].

Acknowledgements - The authors would like to thank the project INNOVA CHILE 06FC01IFC-48, Anillo project ACT-38 and Conicyt scholarship. This work is part of a doctoral dissertation of the first author in forestry engineering at the Facultad de Ingeniería Forestal, Universidad de Concepción, Campus Concepción-C.

References

- [1] (a) Güçlü-Üstündag Ö, Mazza G. (2007) Saponins: properties, applications and processing. *Critical Review in Food Science and Nutrition*, **47**, 231-258; (b) San Martín R, Briones R. (1999) Industrial uses and sustainable supply of *Quillaja saponaria* saponins. *Economy Botany*, **53**, 302-311.
- [2] (a) Bankefors J, Nord L, Kenne L. (2010) Multidimensional profiling of components in complex mixtures of natural products for metabolic analysis, proof of concept: Application to *Quillaja* saponins. *Journal of Chromatography B*, **878**, 471-476; (b) Nord L, Kenne L, Jacobsson S. (2001) Multivariate analysis of H NMR spectra for saponins from *Quillaja saponaria* Molina. *Analytica Chimica Acta*, **446**, 199-209.
- [3] (a) Jörg MA, Vera K, Sven BA, Søren B. (2011) Molecular activities, biosynthesis and evolution of triterpenoid saponins. *Phytochemistry*, **72**, 435-457; (b) Roner M, Sprayberry J, Spinks M, Dhanji S. (2007) Antiviral activity obtained from aqueous extracts of the Chilean soapbark tree (*Quillaja saponaria* Molina). *Journal of General Virology*, **88**, 275-285.
- [4] San Martín R, Otero AF, Figueroa M, Escobar V, Cruz A. (2005) Use of *Quillaja* saponins (*Quillaja saponaria* Molina) to control acid mist in copper electrowinning processes Part I. Laboratory scale evaluation. *Hydrometallurgy*, **77**, 163-170.
- [5] CONAF. (2008) Estudio de prefactibilidad para la Industrialización del Quillay para obtención de extractos de saponinas, en la región del Libertador Bernardo O'Higgins. 1° ed. CONAF, Rancagua.
- [6] (a) Kamstrup S, San Martín R, Doberti A, Grande H, Dalsgaard K. (2000) Preparation and characterisation of *Quillaja* saponin with less heterogeneity than Quil-A. *Vaccine*, **18**, 2244-2249; (b) Szakiel A, Paczkowski C, Henry M. (2010) Influence of environmental abiotic factors on the content of saponins in plants. *Phytochemistry Review*, DOI 10.1007/s11101-010-9177-x.6; (c) Copaja SV, Blackburn C, Carmona R. (2003) Variation of saponin contents in *Quillaja saponaria* Molina. *Wood Science and Technology*, **37**, 103-108; (d) San Martín R, Briones R. (2000). Quality control of commercial quillaja (*Quillaja saponaria* Molina) extracts by reverse phase HPLC. *Journal of the Science of Food and Agriculture*, **80**, 2063-2068.
- [7] Cabrera M. (2002) Respuestas ecofisiológicas de plantas en ecosistemas de zonas con clima mediterráneo y ambientes de alta montaña. *Revista Chilena de Historia Natural*, **75**, 625- 637.
- [8] Rozema J, Björn LO, Bornman JF, Gaberscik A, Häder DP, Trost T, Germ M, Klisch M, Gröniger A, Sinha RP, Lebert M, He YY, Buffoni-Hall R, De Bakker NV, Van De Staaij J, Meijkamp BB. (2002) The role of UV-B radiation in aquatic and terrestrial ecosystems—an experimental and functional analysis of the evolution of UV absorbing compounds. *Journal of Photochemistry and Photobiology B: Biology*, **66**, 2-12.
- [9] (a) Blokhina O, Virolainen E, Fagerstedt K. (2003) Antioxidants, oxidative damage and oxygen deprivation stress: a review. *Annals of Botany*, **91**, 179-194; (b) Wildi B, Lutz C. (1996) Antioxidant composition of selected high alpine plant species from different altitudes. *Plant Cell & Environment*, **19**, 138-146.
- [10] Santibañez F, Uribe J. (1993) Ministerio de Agricultura. FIA – CORFO. Atlas Agroclimático de Chile regiones sexta, séptima, octava y novena. Laboratorio de Agroclimatología. Departamento de Ingeniería y Suelos. Universidad de Chile. Santiago. Chile.
- [11] Longeri L, Etchevers J, Venegas J. (1979) Metodología de perfusión para estudios de nitrificación en suelos. *Ciencia e Investigación Agraria*, **6**, 295-299.
- [12] Sadzawka A, Renato G, Mora M, Saavedra N, Carrasco MA, Rojas C. (2000) Métodos de análisis recomendados para los suelos chilenos. INIA. Santiago. Chile.
- [13] Richards L. (1949) Methods of measuring soil moisture tension. *Soil Sciences*, **68**, 95-112.
- [14] Magdoff F. (1996) Soil organic matter fractions and implications for interpreting organic matter test. In *Soil Organic Matter: Analysis and Interpretation* SSSA. Special Publication N° 46. (Magdoff FR, Tabatabai MA, Hanlon Jr EA(Eds). Soil Science Society of America Inc., Madison, Wisconsin, USA, pp. 11-20.
- [15] Sadzawka A, Renato G, Carrasco MA, Mora M. (2004) Métodos de análisis de tejidos vegetales. INIA. Santiago. Chile.
- [16] Lichtenthaler H, Burgstahler R, Buschmann C, Meier, D, Prenzel, U, Schönthal A (1983) Effect of high light and high light stress on composition, function and structure of the photosynthetic apparatus. Volume 3. In: R. Marcelle, H. Clijsters, and M. Van Poucke (eds.). *Effects of stress on photosynthesis-Advances in Agricultural Biotechnology*. Martinus Nijhoff, Dr. W. Junk Publishers: The Hague, The Netherlands.
- [17] Hatano T, Edamatsu R, Mori A, Fujita Y, Yasuhara T, Yoshida T, Okuda T. (1989) Effects of the interaction of tannins with co-existing substances. VI. Effects of tannins and related polyphenols on superoxide anion radical, and on 1,1-diphenyl-picrylhydrazyl radical. *Chemical & Pharmaceutical Bulletin*, **37**, 2016-2021.
- [18] (a) Di Rienzo JA, Casanoves F, Balzarini MG, Gonzalez L, Tablada M, Robledo CW. (2012) Grupo InfoStat, FCA, Universidad Nacional de Córdoba, Argentina. URL <http://www.infostat.com.ar>; (b) SAS Institute Inc. (2003) The SAS System for Windows, Version 9.1. Cary, NC, USA.

New Access to 7,17-seco C₁₉-Diterpenoid Alkaloids via Vacuum Pyrolysis of *N*-Deethyl-8-acetyl Derivatives

Ling Wang, Qi-Feng Chen and Feng-Peng Wang*

Department of Chemistry of Medicinal Natural Products, West China College of Pharmacy, Sichuan University, Chengdu, Sichuan, PR China, 610041

wfp@scu.edu.cn

Received: September 1st, 2013; Accepted: September 24th, 2013

A novel access to 7,17-seco C₁₉-diterpenoid alkaloids via vacuum pyrolysis of *N*-deethyl-8-acetyl derivatives is described.

Keywords: C₁₉-diterpenoid alkaloid, Pyrolysis, 7,17-Seco C₁₉-diterpenoid alkaloid.


The C₁₉-diterpenoid alkaloids are a group of highly oxygenated, complex, natural products displaying many interesting chemical reactions and important biological activities, as well as a synthetic or structurally modified target [1,2].

In our continuing research on the chemistry of diterpenoid alkaloids, we attempted to prepare the pyro-type alkaloid **3** from **2** via pyrolysis under vacuum conditions. Very interestingly, the 7,17-seco compounds **4** and **5**, in addition to the desired compound **6**, were afforded (Scheme 1). This is a novel access to the 7,17-seco C₁₉-diterpenoid alkaloids, different from the known methods. In this paper, we wish to report this new cleavage of the C(7)-C(17) bonds in the *N*-deethyl-8-acetyl derivatives **2** and **8**.


As showed in Scheme 1, treatment of diacetylchasanine (**1**) with NBS at room temperature for 1 h afforded the desired compound **2** in 60% yield. Pyrolysis of compound **2** under vacuum conditions (15 mm Hg, 180-190°C, 30 min) gave the 7,17-seco compounds **4** (20%) and **5** (10%), as well as the pyro compound **6** (22%). It should be pointed out that pyrolysis of **2** under optimum conditions (15 mm Hg, 160-165°C, 6 min) led to the 7,17-seco compound **5** in high yield (80%).

The HR-ESIMS of **4** showed a quasimolecular ion peak at m/z 416.2445 [M+H]⁺ corresponding to the formula C₂₄H₃₃NO₅. The NMR (¹H, ¹³C, and HMQC) data (Table 1) showed the presence of three methoxyl groups (δ_H 3.19, 3.28, 3.30, each 3H, s; δ_C 56.6 q, 57.8 q, 58.8 q), an acetyl group (δ_H 2.02 s; δ_C 171.3s, 21.2 q), a conjugated double bond (δ_C 128.8 d, δ_H 6.28, d, J = 9.6 Hz; δ_C 129.6 d, δ_H 5.87, t, J = 8.8 Hz; δ_C 131.8 d, δ_H 5.23, d, J = 6.4 Hz; δ_C 135.3 s), and an iminium moiety (δ_C 165.5 d, δ_H 7.72, d, J = 1.6 Hz). The molecular formula of **5** was established as C₂₅H₃₇NO₆ according to its HR-ESIMS and ¹³C NMR data. In comparison with compound **4**, the NMR data of compound **5** showed an additional methoxyl group (δ_H 3.25, 3H, s; δ_C 56.1 q), and the absence of a double bond, leading to confirmation of the structure of **5**. The formula C₂₅H₃₅NO₆ of **6** was established by HR-ESIMS. The structure of **6** could be determined easily based on characteristic signals at δ_H 7.39 (1H, s) for H-19, and δ_H 5.70 (1H, d, J = 6.4 Hz) for H-15, and the absence of the 8-OAc group by comparison with **2**.

Similarly, pyrolysis (15 mm Hg, 180-190°C, 18 min) of compound **8** from **7** generated the desired 7,17-seco compound **9** in 66% yield



Scheme 1. The preparation of compounds **2**, **4**, **5** and **6**



Scheme 2. The preparation of compounds **8**, **9**, **11**, and **13**


Table 1: ^1H and ^{13}C NMR data for compound **4** (^1H : 400 MHz, ^{13}C : 100 MHz, CDCl_3).

Position	δ_{C}	δ_{H} Mult ($J=\text{Hz}$)	^1H - ^1H COSY	HMBC ($\text{H}\rightarrow\text{C}$)
1	89.7 d	2.97 dd (11.6, 4.4)	2 α , 2 β	5, 10, 17, 1-OCH ₃
2	24.4 t	1.17 m (α) 1.75 m (β)	1, 2 β , 3 α , 3 β 2 α , 3 β	—
3	37.8 t	1.57 m (β) 1.84 m (α)	2 α , 2 β , 3 α 2 α , 3 β	4 1
4	39.7 s	—	—	—
5	47.3 d	2.34 (overlapped)	6	1, 7, 17, 19
6	80.1 d	4.76 t (8.8)	5, 7	4, 8, 6-OCH ₃
7	131.8 d	5.23 d (6.4)	6	15
8	135.3 s	—	—	—
9	42.7 d	3.08 t (4.4)	10, 14	7, 12, 13, 15
10	44.0 d	2.37 (overlapped)	9, 12 α	8, 14, 17
11	48.3 s	—	—	—
12	34.1 t	2.21 m (α) 2.35 m (β)	10, 13	9, 11, 16 11, 16
13	36.7 d	2.72 m	12 α , 14, 16	9, 10, 15
14	73.0 d	4.86 t (4.0)	9, 13	8, 16, 14-OAc
15	128.8 d	6.28 d (9.6)	16	7, 9, 13, 17
16	129.6 d	5.87 t (8.8)	13, 15	8, 14
17	165.5 d	7.72 d (1.6)	—	5
18	80.6 t	3.15 (a) ABq (8.4) 3.95 (b)	18b 18a	3, 5 3, 19
19	58.2 t	3.42 (a) ABq (18.8) 3.54 (b)	19b 19a	3, 5, 17 3, 17
1-OCH ₃	56.6 q	3.19 s	—	1
6-OCH ₃	57.8q	3.28 s	—	6
18-OCH ₃	58.8 q	3.30 s	—	18
14-OAc	21.2 q	2.02 s	—	—
	171.3s	—	—	—

Table 2: ^{13}C NMR data for compounds **2**, **5**, **6**, **8**, **9**, **11**, and **13** (100 MHz, CDCl_3).

Position	2	5	6	8	9	11	13
1	82.4 d	82.7 d	84.2 d	83.6 d	79.9 d	83.5 d	81.8 d
2	23.3 t	24.4 t	29.6 t	27.8 t	33.9 t	31.4 t	31.6 t
3	28.5 t	31.4 t	39.0 t	72.3 d	72.0 d	72.1 d	71.8 d
4	38.9 s	39.3 s	50.8 s	49.7 s	47.6 s	49.6 s	52.0 s
5	39.0 d	45.8 d	44.0 d	39.5 d	36.9 d	41.1 d	44.0 d
6	82.2 d	89.7 d	85.0 d	80.6 d	82.8 d	80.5 d	83.9 d
7	43.4 d	133.0 d	33.6 d	44.3 d	129.8 d	45.0 d	47.2 d
8	85.8 s	134.4 s	144.7 s	85.5 s	130.3 s	85.2 s	128.4 s
9	53.4 d	43.2 d	49.9 d	55.4 d	44.1 d	54.5 d	55.0 d
10	42.9 d	38.7 d	49.7 d	42.9 d	43.0 d	42.6 d	47.8 d
11	50.4 s	48.1 s	48.5 s	42.8 s	42.6 s	42.8 s	43.5 s
12	28.9 t	32.7 t	25.3 t	31.1 t	29.8 t	34.2 t	36.0 t
13	43.8 d	44.1 d	52.7 d	45.0 d	42.8 d	81.6 s	84.5 s
14	83.7 d	75.7 d	74.1 d	75.5 d	72.9 d	77.2 d	74.6 d
15	37.5 t	37.3 t	118.0 d	37.7 t	132.9 d	39.4 t	117.9 d
16	75.4 d	79.8 d	80.5 d	82.3 d	131.5 d	79.8 d	79.9 d
17	57.0 d	164.6 d	73.3 d	57.9 d	165.3 d	58.2 d	70.8 d
18	79.8 t	80.4 t	78.2 t	74.0 t	71.9 t	73.7 t	72.0 t
19	49.3 t	58.4 t	166.7 d	41.3 t	51.9 t	41.3 t	42.0 t
1-OCH ₃	57.9 q	56.6 q	55.8 q	56.6 q	56.9 q	57.9 q	56.3 q
6-OCH ₃	59.0 q	58.8 q	59.2 q	58.9 q	58.5 q	58.9 q	58.9 q
14-OCH ₃	—	57.7 q	57.6 q	—	—	—	—
16-OCH ₃	56.4 q	56.1 q	56.2 q	56.5 q	—	56.4 q	58.4 q
18-OCH ₃	55.2 q	—	—	54.8 q	58.0 q	55.6 q	57.4 q
3-OAc	—	—	—	170.4 s	170.1 s	170.4 s	170.7 s
	—	—	—	21.5 q	21.1 q	21.2 q	21.8 q
8-OAc	169.1 s	—	—	169.5 s	—	169.6 s	—
	21.1 q	—	—	21.2 q	—	21.1 q	—
13-OAc	—	—	—	—	—	177.7 s	170.1 s
	—	—	—	—	—	21.4 q	21.1 q
14-OAc	170.6s	171.9 s	171.9 s	—	—	—	—
	22.2 q	21.0 q	21.0 q	—	—	—	—
14-OBz	—	—	—	166.1 s	166.4 s	166.1 s	166.7 s
	—	—	—	130.1 s	135.5 s	129.8 s	144.5 s
	—	—	—	133.0 d	129.6 d	133.2 d	132.7 d
	—	—	—	129.6 d \times 2	128.9	129.8	130.1
	—	—	—	128.4 d \times 2	d \times 2	d \times 2	d \times 2
	—	—	—	128.2	128.5	128.2	—
	—	—	—	d \times 2	d \times 2	d \times 2	—

(Scheme 2). However, treatment of compound **11**, bearing the hydroxyl group at C-13, under similar conditions (15 mm Hg, 180–190 °C, 10 min) produced only pyro compound **13** instead of the expected 7,17-seco compound **12** (Scheme 2), indicating that the hydroxyl group at C-13 in **11** was unfavorable for the cleavage of the C(7)-C(17) bond, which was difficult to interpret. The formula $\text{C}_{31}\text{H}_{37}\text{NO}_7$ of **9** was established from HR-ESIMS and ^{13}C NMR data. The NMR spectra of **9** showed the presence of a conjugated double bond (δ_{C} 132.9 d, δ_{H} 6.38, d, $J = 9.2$ Hz; δ_{C} 131.5 d, δ_{H} 5.95, t, $J = 8.8$ Hz; δ_{C} 129.8 d, δ_{H} 5.24, d, $J = 6.4$ Hz; δ_{C} 130.3 s), and an iminium moiety (δ_{C} 165.3 d, δ_{H} 7.78, s). Its structure could be determined easily by comparison of spectral data with those of compound **4**. The molecular formula for **13** was established as $\text{C}_{34}\text{H}_{43}\text{NO}_{10}$ from HR-ESIMS and ^{13}C NMR data. The structure of

**Scheme 3.** A plausible mechanism of pyrolysis of the N-deethyl-8-acetyl derivatives of C_{19} -diterpenoid alkaloids

13 was confirmed by the characteristic signal at δ_{H} 5.73 (1H, d, $J = 6.8$ Hz) for H-15, and the absence of an 8-OAc group in the NMR spectra when compared with those of compound **11**.

A process of cleavage of the C(7)-C(17) bonds in compounds **8** and **11** was postulated (Scheme 3). As shown in Scheme 3, there were two competitive reactions, e. g., cleavage of the C(7)-C(17) bonds via Grob fragmentation ($\text{A}\rightarrow\text{B}$) and elimination ($\text{A}\rightarrow\text{C}$) during pyrolysis. Importantly, the hydroxyl group at C-13 in the substrates, as in **11**, was not helpful for cleavage of the C(7)-C(17) bonds.

In conclusion, although the literature reports many methods of cleavage of the C(7)-C(17) bonds in C_{19} -diterpenoid alkaloids [3–9], most of which lead to complicated products with either low yields or difficult purification, except for the Grob fragmentation- NaBH_4 reduction method developed by us [10,11]. Obviously, the method reported here is simpler and more convenient than those of the Grob fragmentation- NaBH_4 reduction procedure.

Experimental

General methods: Optical rotations were measured in a 1.0 dm cell with a PE-314 polarimeter at $20\pm 1^\circ\text{C}$; IR spectra were recorded on a Nicolet 200 SXV spectrometer; HRMS were obtained with a Bruker BioTOFQ mass spectrometer; ^1H and ^{13}C NMR spectra were acquired on a Varian INOVA-400/54 spectrometer, with TMS as internal standard; silica gel GF254 and H (Qingdao Sea Chemical Factory, China) were used for TLC and CC.

Preparation of compound 2: To a solution of diacetylchamanine (**1**) (7.46 g, 13.90 mmol) in HOAc (100 mL) was added NBS (6.9 g, 38.84 mmol), and the mixture was allowed to stand at room temperature for 3 h prior to being poured into ice water (100 mL). The mixture was basified with conc. NH_4OH solution to pH 9 and extracted with dichloromethane (150 mL \times 3). The extracts were dried (Na_2SO_4) and concentrated to give a crude product, which was subjected to CC (silica gel H, 100 g), using light petroleum/acetone/diethylamine (66:33:1) as eluent, to afford **2** as a white amorphous power (3.87 g, 55%).

Compound 2

$[\alpha]_{\text{D}}^{20}$: +5.0 (c 0.50, CHCl_3).

IR (KBr): 2935, 2824, 1732, 1641, 1243, 1098 cm^{-1} .

^1H NMR (400 MHz, CDCl_3) δ : 1.97, 2.04 (each 3H, s, $\text{OAc}\times 2$), 3.01 (1H, ABq, $J = 8.4$ Hz, H-18), 3.22, 3.26, 3.31, 3.32 (each 3H, s, $\text{OCH}_3\times 4$), 3.58 (1H, ABq, $J = 8.4$ Hz, H-18), 4.06 (1H, d, $J = 5.6$ Hz, H-6), 4.81 (1H, t, $J = 4.8$ Hz, H-14 β).

^{13}C NMR (100 MHz, CDCl_3): Table 2.

HRMS-ESI m/z $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{27}\text{H}_{41}\text{NO}_8$: 508.2910; found: 508.2912.

Preparation of 4, 5, and 6: Compound **2** (570 mg, 1.12 mmol), in a round bottomed flask (50 mL), was heated at 180–190°C under 15 mm Hg for 30 min. After cooling to room temperature, the residue was chromatographed over silica gel H eluting with cyclohexane/acetone/diethylamine (30:1:0.15) to give **4** (a white

amorphous power, 110 mg, 20%), **5** (a white amorphous power, 55 mg, 10%), and **6** (a white amorphous power, 120 mg, 20%).

Compound 4

$[\alpha]_D^{20}$: +16.4 (*c* 0.50, CHCl₃).

IR (KBr): 2925, 1734, 1624, 1399, 1256, 1100 cm⁻¹.

¹H NMR (400 MHz, CDCl₃): Table 1.

¹³C NMR (100 MHz, CDCl₃): Table 1.

HRMS-ESI *m/z* [M + H]⁺ calcd for C₂₄H₃₄NO₅: 416.2437; found: 416.2445.

Compound 5

$[\alpha]_D^{20}$: +24.6 (*c* 0.50, CHCl₃).

IR (KBr): 2927, 2823, 1732, 1637, 1371, 1249, 1103 cm⁻¹.

¹H NMR (400 MHz, CDCl₃) δ: 2.05 (3H, s, OAc), 3.19, 3.25, 3.28, 3.30 (each 3H, s, OCH₃×4), 4.47 (1H, t, *J* = 7.2 Hz, H-6), 4.88 (1H, t, *J* = 4.4 Hz, H-14β), 5.26 (1H, d, *J* = 6.4 Hz, H-7), 7.77 (1H, s, H-17).

¹³C NMR (100 MHz, CDCl₃): Table 2.

HRMS-ESI *m/z* [M + H]⁺ calcd for C₂₅H₃₈NO₆: 448.2699; found: 448.2697.

Compound 6

$[\alpha]_D^{20}$: -7.2 (*c* 0.50, CHCl₃).

IR (KBr): 2935, 2820, 1734, 1639, 1396, 1253, 1100 cm⁻¹.

¹H NMR (400 MHz, CDCl₃) δ: 2.00 (3H, s, OAc), 3.21, 3.28, 3.30, 3.34 (each 3H, s, OCH₃×4), 3.66, 3.75 (2H, ABq, *J* = 8.4 Hz, H-18), 4.20 (1H, d, *J* = 6.4 Hz, H-6), 4.65 (1H, t, *J* = 4.0 Hz, H-14β), 5.70 (1H, d, *J* = 6.4 Hz, H-15), 7.39 (1H, s, H-19).

¹³C NMR (100 MHz, CDCl₃): Table 2.

HRMS-ESI *m/z* [M + H]⁺ calcd for C₂₅H₃₆NO₆: 446.2543; found: 446.2542.

Preparation of compound 8: To a solution of 3-acetyl-13-deoxyindaconitine (**7**) (665 mg, 1.0 mmol) in HOAc (35 mL) was added NBS (576 mg, 3.24 mmol), and the mixture was stirred at room temperature for 1 h prior to being poured into ice water (10 mL). The mixture was basified with conc. NH₄OH solution to pH 9 and extracted with dichloromethane (50 mL × 3). The extracts were dried (Na₂SO₄) and concentrated. CC (silica gel H, 10 g) of the residue, using light petroleum/acetone (2:1) as eluent, afforded compound **8** as a white amorphous power (360 mg, 60%).

Compound 8

$[\alpha]_D^{20}$: +12.8 (*c* 0.50, CHCl₃).

IR (KBr): 3450, 1719, 1637, 1399, 1249, 1103 cm⁻¹.

¹H NMR (400 MHz, CDCl₃) δ: 1.38, 2.07 (each 3H, s, OAc), 3.22, 3.24, 3.26, 3.38 (each 3H, s, OCH₃×4), 4.18 (1H, d, *J* = 6.8 Hz, H-6), 7.43~8.05 (5H, OBz).

¹³C NMR: Table 2.

HRMS-ESI: *m/z* [M + H]⁺ calcd for C₃₄H₄₆NO₁₀: 628.3122; found: 628.3117.

Preparation of compound 9: Compound **8** (50 mg, 0.08 mmol), in a round bottomed flask (10 mL), was heated at 180-190°C under 15 mm Hg for 18 min. After cooling to room temperature, the residue was chromatographed over silica gel H eluting with cyclohexane/acetone (6:1) to give **9** (a white amorphous power, 26 mg, 66%).

References

- [1] Wang FP, Chen QH. (2010) The C₁₉-diterpenoid alkaloids. In *The Alkaloids: Chemistry and Biology*, Vol. 69, Cordell GA (Ed.), Elsevier Science, New York, 1-577.

Compound 9

$[\alpha]_D^{20}$: +48.0 (*c* 0.50, CHCl₃).

IR (KBr): 2934, 2889, 2821, 1718, 1645, 1278, 1102, 715 cm⁻¹.

¹H NMR (400 MHz, CDCl₃) δ: 2.08 (3H, s, OAc), 3.22, 3.23, 3.30 (each 3H, s, OCH₃×3), 3.80 (1H, d, *J* = 18.8 Hz, H-19), 3.03, 4.25 (2H, ABq, *J* = 8.4 Hz, H-18), 4.86 (1H, t, *J* = 7.6 Hz, H-6), 4.92 (1H, dd, *J* = 12.4, 5.2 Hz, H-3), 5.15 (1H, t, *J* = 4.0 Hz, H-14β), 5.24 (1H, d, *J* = 6.8 Hz, H-7), 5.95 (1H, t, *J* = 8.8 Hz, H-16), 6.38 (1H, d, *J* = 9.2 Hz, H-15), 7.78 (1H, s, H-17), 7.40~7.99 (5H, OBz); ¹³C NMR: Table 2.

HRMS-ESI: *m/z* [M + H]⁺ calcd for C₃₁H₃₈NO₇: 536.2648; found: 536.2649.

Preparation of compound 1: To a solution of diacetylandaonitine (**10**) (627 mg, 0.86 mmol) in HOAc (15 mL) was added NBS (528 mg, 2.98 mmol), and the mixture was allowed to stand at room temperature for 3 h prior to being poured into ice water (15 mL). The mixture was basified with conc. NH₄OH solution to pH 9 and extracted with dichloromethane (50 mL × 3). The extracts were dried (Na₂SO₄) and concentrated to give a crude product, which was subjected to CC (silica gel H, 10 g), using cyclohexane/acetone (10:1) as eluent, to afford **11** as a white amorphous power (360 mg, 60%).

Compound 11

$[\alpha]_D^{20}$: +4.8 (*c* 0.50, CHCl₃);

IR (KBr): 3447, 1715, 1637, 1252, 1102 cm⁻¹;

¹H NMR (400 MHz, CDCl₃) δ: 1.27, 2.04, 2.06 (each 3H, s, OAc×3), 3.19, 3.23, 3.27, 3.39 (each 3H, s, OCH₃×4), 4.09 (1H, d, *J* = 6.0 Hz, H-6), 5.07 (1H, dd, *J* = 8.0, 6.0 Hz, H-3), 5.13 (1H, d, *J* = 4.4 Hz, H-14β), 7.44~8.09 (5H, OBz);

¹³C NMR: Table 2;

HRMS-ESI: *m/z* [M + H]⁺ calcd for C₃₆H₄₈NO₁₂: 686.3177; found: 686.3176.

Preparation of 13: Compound **11** (48 mg, 0.07 mmol), in a round bottomed flask (10 mL), was heated at 180-190°C under 15 mm Hg for 10 min. After cooling to room temperature, the residue was chromatographed over silica gel H eluting with cyclohexane/acetone (6:1) to give **13** (a white amorphous power, 26 mg, 56%).

Compound 13

$[\alpha]_D^{20}$: +36.4 (*c* 0.50, CHCl₃).

IR (KBr): 2931, 2821, 1735, 1640, 1245, 1098 cm⁻¹.

¹H NMR (400 MHz, CDCl₃) δ: 2.05, 2.06 (3H, s, OAc×2), 3.22, 3.28, 3.30, 3.46 (each 3H, s, OCH₃×4), 3.00, 4.01 (2H, ABq, *J* = 8.8 Hz, H-18), 4.29 (1H, t, *J* = 7.2 Hz, H-6), 4.89 (1H, dd, *J* = 12.8, 4.8 Hz, H-3), 5.33 (1H, d, *J* = 2.8 Hz, H-14β), 5.24 (1H, d, *J* = 6.8 Hz, H-7), 5.73 (1H, d, *J* = 6.8 Hz, H-15), 7.40~8.14 (5H, OBz).

¹³C NMR: Table 2.

HRMS-ESI: *m/z* [M + H]⁺ calcd for C₃₄H₄₄NO₁₀: 626.2965; found: 626.2969.

Acknowledgement - We are grateful to the National Science Foundation of China (No.81273387) for financial support of this research.

- [2] Wang, FP, Liang XT. (1992) Chemistry of the diterpenoid alkaloids. In *The Alkaloids: Chemistry and Physiology*, Cordell GA (Ed.), Academic Press, New York, Vol. 42, 151–247.
- [3] (a) Wiesner K, Gotz M, Simmons DL, Fowler LR, Bachelor FW, Brown RFC, Biichi G. (1959) The structure of aconitine. *Tetrahedron Letters*, 2, 15-24; (b) Wang FP, Pelletier SW, (1991) Preparation and mechanism of 7,17-seco C₁₉-diterpenoid alkaloids via pyrolysis of their *N*-oxides in diglyme. *Chinese Chemical Letters*, 2, 103-106.
- [4] (a) Edwards OE. (1965) The pyrolysis of diterpenoid ester alkaloids. *Chemical Communications*, 318-319; (b) Wada K, Bando H, Mori T, Wada R, Kanaiwa Y, Amiya T. (1985) Studies on the constituents of *Aconitum* species. III. On the components of *Aconitum subcuneatum* Nakai. *Chemical & Pharmaceutical Bulletin*, 33, 3658-3661; (c) Edwards OE, Fonzes L, Marion L. (1966) The correlation of chasmanine with browniine. *Canadian Journal of Chemistry*, 44, 583-589.
- [5] (a) Pelletier SW, Moore JF, Desai RC, Mody NV, Desai HK. (1982) The pyrodelphinine chromophore. Crystal structures of pyrodelphinine and delphinine. *Journal of Organic Chemistry*, 47, 5290-5297; (b) Wiesner K, Inaba T. (1969) Photoreduction of demethoxy pyrodelphinine. *Journal of American Chemical Society*, 91, 1036-1037.
- [6] (a) Amiya T, Shima T. (1967) The aconite alkaloids. XXXVII. Anhydrodiacetyldecosine (anhydrodiacetyllyucaconine) and its novel hydrogenation. *Bulletin of the Chemical Society of Japan*, 40, 1957-1960; (b) Amiya T, Kanaiwa Y, Nakano N, Shima T. (1978) Some derivatives of anhydrodiacetyldecosine (anhydrodiacetyllyucaconine). *Bulletin of the Chemical Society of Japan*, 51, 248-250.
- [7] (a) Edward OE, Los M, Marion L. (1959) Hydroxylycoctonine. *Canadian Journal of Chemistry*, 37, 1996-2006; (b) Valenta Z, Wright IG. (1960) The structure and chemistry of hydroxylycoctonine. *Tetrahedron*, 9, 284-288; (c) Edwards OE, Marion L, Stewart DKR. (1956) The chemistry of lycoctonine. *Canadian Journal of Chemistry*, 34, 1315-1328.
- [8] (a) Aneja R, Pelletier SW. (1964) The diterpene alkaloids. The structure of heteratisine. *Tetrahedron Letters*, 5, 669-677; (b) Aneja R, Pelletier SW. (1965) The diterpene alkaloids. The pyrolysis and absolute configuration of heteratisine, *Tetrahedron Letters*, 6, 215-223.
- [9] Wang FP, Fan JZ, Jian XX, Li BG. (1999) Modification of norditerpenoids alkaloids: III. Preparation of 7, 17- seco yunnaconitine derivatives via rearrangement of chloroasmine. *Chinese Chemical Letters*, 10, 379-382.
- [10] Wang FP, Yang JS, Chen QH, Yu L, Li BG. (2000) New access to 7,17-seco norditerpenoid alkaloids via reduction of 8-chloro derivatives. *Chemical & Pharmaceutical Bulletin*, 48, 1912-1916.
- [11] Chen QH, Wang FP. (2003) Further study of the synthesis of 7,17-seco norditerpenoid alkaloids. *Journal of Asian Natural Products Research*, 5, 43-48.

Alkaloids from *Boophone haemanthoides* (Amaryllidaceae)Jerald J. Nair^a, Lucie Rárová^b, Miroslav Strnad^{b,c}, Jaume Bastida^d and Johannes van Staden^{a*}^aResearch Centre for Plant Growth and Development, School of Life Sciences, University of KwaZulu-Natal Pietermaritzburg, Private Bag X01, Scottsville 3209, South Africa^bCentre of the Region Haná for Biotechnological and Agricultural Research, Palacký University, Šlechtitelů 11, 78371 Olomouc, Czech Republic^cLaboratory of Growth Regulators, Palacký University & Institute of Experimental Botany AS CR, Šlechtitelů 11, CZ 78371 Olomouc, Czech Republic^dDepartament de Productes Naturals, Facultat de Farmàcia, Universitat de Barcelona, 08028, Barcelona, Spain

rcpgd@ukzn.ac.za

Received: August 21st, 2013; Accepted: September 21st, 2013

In this study, the South African Amaryllid *Boophone haemanthoides* was examined for its phytochemical composition and cytotoxicity. In the process eight alkaloid structures, including the new compound distichaminol, were identified in bulb ethanolic extracts. Of the isolates, lycorine and distichamine exhibited strong activities against human acute lymphoblastic leukemia (CEM), breast adenocarcinoma (MCF7) and cervical adenocarcinoma (HeLa) cells with IC₅₀s ranging from 1.8 to 9.2 µM.

Keywords: Alkaloid, Amaryllidaceae, *Boophone haemanthoides*, Distichaminol.


Southern Africa is a center of prominence for the family Amaryllidaceae, containing roughly a third of the global population of around 1000 species [1]. Its members are known for their horticultural and ornamental appeal with significant impact on the floriculture sector [1]. More importantly, its medicinal properties have long been realized in the traditional medicinal practices of indigenous peoples across the globe [2]. For example, the discovery of galanthamine (**1**) (scheme 1) as an Alzheimer's drug evolved out of the ethnic usage of *Galanthus* species, from which the compound was first isolated, in the Caucasus region of Eastern Europe [3]. In South Africa the Amaryllidaceae has a long history of use for medicinal purposes amongst its indigenous people which can be traced back to its original inhabitants, the San [2a,b].

The chemicals responsible for these medicinal attributes are alkaloids, structural variations of which are unique to the Amaryllidaceae [2]. Six different structural types are discernible for these compounds as represented by galanthamine (**1**), lycorine (**2**), crinine (**3**), homolycorine (**4**), tazettine (**5**) and montanine (**6**) [2]. Other less conspicuous members include degraded, oxidized and truncated variants, such as trisphaeridine (**7**) and ismine (**8**) [2]. All of these compounds are related, as a consequence of their biogenesis, from a common amino acid derived precursor, norbelladine (**9**) [2]. The pharmaceutical value of the Amaryllidaceae is epitomized by the FDA approved status granted to galanthamine (**1**) due to its potent and selective inhibitory interaction with the enzyme acetylcholinesterase (AChE), of significance in the progression of neurodegeneration associated with motor neuron diseases [3]. Following on the commercial success of galanthamine as an Alzheimer's drug, it is anticipated that further clinical candidates will emerge out of the phenanthridones {such as pancratistatin (**10**)} of the lycorine (**2**) series of alkaloids, which are known for their potent and selective antiproliferative properties [4]. Not surprisingly, therefore, AChE inhibition and cytotoxicity are two recurrent themes in the pharmacology of Amaryllidaceae alkaloids [3,4].

Lycorine (**2**) (scheme 1) is the most common alkaloid across the Amaryllidaceae and known for a broad spectrum of biological properties, including antiviral [5a], antifungal [5b], antiparasitic [5c] and anti-inflammatory [5d] activities, as well as antifeedant [5e], cell growth [5f] and AChE [5g] inhibitory effects. Furthermore, it has shown much promise as an antiproliferative agent capable of inhibiting cell division and cell elongation, as well as protein synthesis in eukaryotic cells [5h]. Its chemotherapeutic potential is best underlined by its antitumor effects in a number of cancer cell lines [6]. Furthermore, both *in vitro* and *in vivo* models of leukemia (HL-60) cells support the potency of lycorine as an anticancer agent [6d,e]. Notably, in some cell lines these effects were manifested via the apoptotic mechanistic pathway [6e-g]. In contrast, antiproliferative effects of lycorine were also exerted in apoptosis-resistant cell lines such as OE21 oesophageal cancer and SKMEL-28 melanoma cells [6h].

As mentioned above, the phenanthridone, pancratistatin (**10**), of the lycorine series has shown most promise for clinical development as an anticancer agent [7]. This potential is vindicated by the fact that pancratistatin selectively targeted mitochondria of cancer cells, with minimal effect on normal cells, initiating cell death via the apoptotic pathway, as indicated by early activation of caspase-3, followed by flipping of phosphatidyl serine [7e,f]. A stronger case could be made for commercial development of these targets based on their low interaction with the isoenzyme cytochrome P450 3A4, which is responsible for metabolism of the majority of drugs taken by humans [8]. By contrast, the crinine group, exemplified by crinine (**3**), is a broad and expanding group of compounds also known for diverse biological properties [9].

Given their close structural similarity to the lycorine series of compounds [2], cytotoxic effects have been demonstrated for several members of the crinine group [9b]. Over the past two decades there has been significant interest in diverse research aspects of the Amaryllidaceae, including areas pertaining to its



Scheme 1: Diverse alkaloid structures of the Amaryllidaceae, including major and minor group representatives **1-8**, the common biosynthetic precursor norbelladine **9** as well as the novel compound distichaminol **17** identified in the present study of *Boophone haemanthoides*.

ethnobotany, phytochemistry and pharmacology [2-9]. In addition, synthetic endeavors have targeted some of its bioactive constituents for structure-activity relationship (SAR) study purposes [4,10]. In continuation of these efforts, the South African Amaryllid *Boophone haemanthoides* was here investigated for its alkaloid composition and cytotoxic effects against a mini-panel of cancer cells. Seven alkaloids of the β -crinine series, including crinine (**3**), buphanisine (**11**), buphanidrine (**12**), ambelline (**13**), undulatine (**14**), distichamine (**16**) and distichaminol (**17**), together with lycorine (**2**) (scheme 1) were isolated and identified by a combination of physical and spectroscopic methods. Cytotoxic effects were demonstrated for lycorine and distichamine in acute lymphoblastic leukemia (CEM), breast adenocarcinoma (MCF7) and cervical adenocarcinoma (HeLa) cells with IC_{50} s determined in the range 1.8 to 9.2 μ M, while distichaminol (**17**) was uncovered as a novel alkaloid.

The African genus *Boophone* Herb. is comprised of two known species, *B. disticha* (L.f.) Herb. and *B. haemanthoides* F.M. Leight. *B. disticha* is widely distributed in Africa, ranging from Sudan in the north to the Western Cape Province in the south, while *B. haemanthoides* is a rare and threatened species with a restricted territory within the winter rainfall region of South Africa and parts of southern Namibia [11a]. Both plants are known to be widely used in traditional medicine [11b,c]. Isolation of the alkaloids from bulbs of *B. haemanthoides* is described in the Experimental section. Of the compounds isolated, lycorine (**2**), crinine (**3**), buphanisine (**11**), buphanidrine (**12**), ambelline (**13**) and undulatine (**14**) (scheme 1) are common across most genera of the Amaryllidaceae [2]. By contrast, distichamine (**16**) is unique in that its C-ring double bond is situated at C-2/C-3, as opposed to the usual C-1/C-2 positioning

in most crinine compounds, with concomitant vinylization of the C-3 methoxyl group, as well as oxidation at C-1. Moreover, it has never been found outside the genus *Boophone* and as such occupies a privileged chemosystematic status within the Amaryllidaceae [11d]. A previous phytochemical study made on a wild population of *B. haemanthoides* from the Saldhana Bay area of South Africa revealed the presence of buphanidrine, buphanisine, crinine and distichamine [11d]. In the present study, bulbs of *B. haemanthoides* were collected in the Nieuwoudtville area of the Northern Cape Province of South Africa. The elucidation of the novel structure distichaminol (**17**) was based on its close structural similarity to distichamine (**16**) and buphanamine (**21**), previously described from *B. disticha* [12]. HRMS indicated a mass of 332.1491 g/mol for the $[M+1]^+$ ion of distichaminol, correct for the molecular formula $C_{18}H_{22}NO_5$ and theoretical mass 332.1498 g/mol. Furthermore, its EIMS had the molecular ion $[M]^+$ peak at m/z 331 with a relative abundance of 89%. Diagnostic fragment ions were detected at m/z 316 $[M-CH_3]^+$, 299 $[M-CH_3OH]^+$ and 281 $[M-CH_3OH-H_2O]^+$. FTIR indicated *inter alia* the presence of hydroxyl (3500-3000 cm^{-1}), aryl (1617 cm^{-1}), methoxyl (1039 cm^{-1}) and methylenedioxy (934 cm^{-1}) groups.

The 1H and ^{13}C NMR spectra of distichaminol (**17**) were similar to those of distichamine (**16**) and only differences due to the C-1 hydroxyl group in **17** were significant (Table 1) [12b,c]. The lowfield region (δ 4.4-7.6) of the proton spectrum had four signals resonant for H-10 (δ 7.16, s), the methylenedioxy group (δ 5.86, s), H-2 (δ 4.91, dd, $J=2.4, 1.0$ Hz) and H-1 (δ 4.61, d, $J=2.4$ Hz), respectively. The corresponding carbon resonances were detected at δ 100.6 (d, C-10), 100.9 (t, OCH_2O), 96.8 (d, C-2) and 72.2 (d, C-1) respectively, as determined by HSQC experiments (Table 1). Two

Table 1: 1D and 2D NMR spectroscopic data for distichaminol (**17**).

Position	δ_H	δ_C	COSY	HMBC	NOESY
1	4.61 (d, 2.4)	72.2 (d)	H-2	C-2, C-3, C-4a, C-10a, C-10b, C-11	H-2, H-4a
2	4.91 (dd, 2.4, 1.0)	96.8 (d)	H-1, H-4 α , H-4 β	C-1, C-3, C-4, C-10b	H-1, 3-OCH ₃
3	—	154.9 (s)	—	—	—
4 α	2.05 (ddd, 17.6, 7.0, 1.0)	29.8 (t)	H-2, H-4 β , H-4a	C-2, C-3, C-4a, C-10b	3-OCH ₃ , H-4 β , H-4a
4 β	2.31 (ddd, 17.6, 10.0, 1.0)	29.8 (t)	H-2, H-4 α , H-4a	C-2, C-3, C-4a, C-10b	3-OCH ₃ , H-4 α , H-12 exo
4a	3.26 (dd, 10.0, 7.0)	64.6 (d)	H-4 α , H-4 β	C-1, C-3, C-4, C-10a, C-10b, C-11	H-1, H-4 α , H-6 α
6 α	4.14 (d, 17.3)	57.8 (t)	H-6 β	C-6a, C-7, C-10a, C-12	H-4a, H-6 β , 7-OCH ₃
6 β	3.79 (d, 17.3)	57.8 (t)	H-6 α	C-6a, C-7, C-10a, C-12	H-6 α , 7-OCH ₃ , H-12 $endo$
6a	—	117.0 (s)	—	—	—
7	—	140.3 (s)	—	—	—
8	—	140.5 (s)	—	—	—
9	—	148.7 (s)	—	—	—
10	7.16 (s)	100.6 (d)	—	C-6a, C-8, C-9, C-10a, C-10b	—
10a	—	134.0 (s)	—	—	—
10b	—	48.1 (s)	—	—	—
11 $endo$	2.12 (ddd, 12.5, 9.8, 4.0)	36.3 (t)	H-11 exo , H-12 $endo$, H-12 exo	C-1, C-4a, C-10a, C-10b, C-12	H-11 exo , H-12 $endo$, H-12 exo
11 exo	2.03 (ddd, 12.5, 10.8, 6.4)	36.3 (t)	H-11 $endo$, H-12 $endo$, H-12 exo	C-1, C-4a, C-10a, C-10b, C-12	H-11 $endo$, H-12 $endo$, H-12 exo
12 $endo$	2.76 (ddd, 12.9, 9.0, 5.6)	51.3 (t)	H-11 $endo$, H-11 exo , H-12 exo	C-4a, C-6, C-10b, C-11	H-6 β , H-11 $endo$, H-11 exo , H-12 exo
12 exo	3.45 (ddd, 12.9, 10.0, 3.8)	51.3 (t)	H-11 $endo$, H-11 exo , H-12 $endo$	C-4a, C-6, C-10b, C-11	H-4 β , H-11 $endo$, H-11 exo , H-12 $endo$
3-OCH ₃	3.57 (s)	54.7 (q)	—	C-3	H-2, H-4 α , H-4 β
7-OCH ₃	3.97 (s)	59.5 (q)	—	C-7	H-6 α , H-6 β
OCH ₂ O	5.86 (s)	100.9 (t)	—	C-8, C-9	—

and three bond HMBC correlations linked H-10 to C-6a, C-8, C-9, C-10a and C-10b. The C-1 carbon resonance, indicative of a hydroxy-substituted allylic methine carbon in **17**, is distinguished from that of distichamine (**16**) (δ 201.9, s) in which C-1 possesses an α,β -unsaturated ketone [12c]. Mutual COSY contours between H-1 and H-2 established their vicinal relationship. Furthermore, the small value of the coupling constant between H-1 and H-2 ($J=2.4$ Hz) indicated a dihedral angle approaching 90° for these protons, thus favouring the β -disposition for the C-1 hydroxyl group. By contrast, a J value of 5.6 Hz between H-1 and H-2 reflects α -orientation for the 1-hydroxy group as shown for buphanamine (**21**) [12b]. In addition, allylic coupling was established via COSY between H-2 and the methylene protons H-4 α and H-4 β by the small value of the coupling between them ($J=1.0$ Hz).

The oxygen and nitrogen related proton signals were resonant in the δ 2.6–4.2 region, of which the diastereotopic C-6 protons were immediately apparent based on their chemical shift, multiplicity and coupling constants. As such, H-6 α (δ 4.14, d, $J=17.3$ Hz) was shifted further downfield than H-6 β (δ 3.79, d, $J=17.3$ Hz) due to the former being *cis*-disposed to the nitrogen atom (Table 1). This was also true for H-12 exo (δ 3.45, ddd, $J=12.9, 10.0, 3.8$ Hz) resonating further downfield to H-12 $endo$ (δ 2.76, ddd, $J=12.9, 9.0, 5.6$ Hz). HSQC analysis showed these protons to be attached to methylene carbons which resonated at δ 57.8 (t, C-6) and 51.3 (t, C-12), respectively. The remaining three-proton singlet resonance signals in this region were ascribable to the vinyl C-3 (δ 3.57) and aryl C-7 (δ 3.97) methoxy groups, while the nitrogen related methine proton H-4a was resonant at δ 3.26 (dd, $J=10.0, 7.0$ Hz). As expected, the methylene protons at C-4 and C-11 were the most shielded. An *anti*-periplanar relationship between H-4a and H-4 β was established by the large value of the coupling between them ($J=10.0$ Hz). All eighteen carbons of distichaminol (**17**) were accounted for by ^{13}C and HSQC NMR and multiplicities ascertained by DEPT analysis (Table 1). Furthermore, absolute configuration of the 5,10b-ethano-bridge was arrived at by circular dichroism (CD) measurements which showed that distichaminol was qualitatively similar to β -5,10b-ethanophenanthridine compounds with a maximum at 250 nm and a minimum at 286 nm in its CD curve [13].

A mini-panel of cancer cells comprising CEM, MCF7 and HeLa cells, as well as the normal human fibroblast (BJ) cell line was engaged for the cytotoxicity screen using staurosporine and

galanthamine as positive and negative controls, respectively. As determined by the Calcein AM assay [14], lycorine (**2**) decreased the survival of all three cancer cell lines in a dose-dependent manner after 72 h treatments, with IC₅₀ values of 1.8, 9.2 and 8.9 μM , respectively (Table 2). This result is not surprising given that the cytotoxic properties of lycorine have been widely studied [6]. Similarly, distichamine (**16**) was noticeably active, exhibiting IC₅₀s of 5.1, 2.3 and 4.0 μM against the three cancer cells respectively, in accordance with a previous report [14]. However, both lycorine and distichamine were not selective in their activity since normal BJ cells were also affected to a significant extent by the treatment (IC₅₀ 2.0 and 12.4 μM , respectively), in agreement with earlier findings [6,14]. Furthermore, the β -crinine compounds crinine (**3**), buphanisine (**11**), buphanidrine (**12**) and ambelline (**13**), all of which possess a C-1/C-2 double bond, as well as polar substitution at C-3 α , were inactive against both normal and cancerous cells (IC₅₀s > 50 μM). This was also the case for the 1,2- β -epoxide undulatin (**14**), as well as the new compound distichaminol (**17**) (IC₅₀s > 50 μM against both normal and cancer cell types).

Since the identification of haemanthamine (**19**) as a cytotoxic agent over thirty years ago [5h], alkaloids of the crinine series of the Amaryllidaceae have been the subject of several cytotoxicity based studies. A recent review of the field has revealed that since 1976, seventy-one such compounds have been screened against fifty-four different cancer cell lines [9b]. The review also showed that crinine (**18**), haemanthamine (**19**) and haemanthidine (**20**) are the most common targets in such studies of α -crinanes [9b]. Within the β -series, buphanisine (**11**) and ambelline (**13**) appear routinely across most of the cell lines screened [9b]. The low activities observed here for the β -crinine representatives crinine (**3**), buphanisine (**11**), buphanidrine (**12**), ambelline (**13**) and undulatin (**14**) are in line with previous studies made on various cell lines [9b]. With a few exceptions, α -crinanes in general superseded β -crinanes in terms of potency [9b]. For example, augustine (**15**) was the most active of the β -series with an ED₅₀ of 0.6 $\mu\text{g/mL}$ in both KB (human oral epidermoid carcinoma) and U373 (human glioblastoma astrocytoma) cells [9b].

In relation to the mechanistic basis of these alkaloids, the apoptotic mode of death in cancers has been demonstrated for crinine (**3**), distichamine (**16**), crinine (**18**) and haemanthamine (**19**) [6h,14,15]. For example, McNulty *et al.* showed that up to 90% of rat hepatoma (5123 tc) cells exhibited apoptotic morphology after a 48 h treatment with the α -crinanes crinine and haemanthamine,

Table 2: IC₅₀ (μM) values obtained from Calcein AM assays using the specified cancerous and normal cell lines.

Sample ^a	Cell line, IC ₅₀ (μM) ^{b,c}			
	CEM	MCF7	HeLa	BJ
Galanthamine (1) ^d	> 50	> 50	> 50	> 50
Lycorine (2)	1.8 ± 0.1	9.2 ± 2.1	8.9 ± 2.6	2.0 ± 0.4
Crinine (3)	> 50	> 50	> 50	> 50
Buphanisine (11)	> 50	> 50	> 50	> 50
Buphanidine (12)	> 50	> 50	> 50	> 50
Ambelline (13)	> 50	> 50	> 50	> 50
Undulatine (14)	> 50	> 50	> 50	> 50
Distichamine (16)	5.1 ± 0.6	2.3 ± 0.8	4.0 ± 0.4	12.4 ± 2.2
Distichaminol (17)	> 50	> 50	> 50	> 50
Staurosporine ^d	0.023 ± 0.002	0.064 ± 0.002	0.175 ± 0.007	0.002 ± 0.000

^a All compounds were single entities as determined by TLC, HPLC and NMR analysis. ^b Cells were treated for 72 h with serial concentrations of samples. ^c Values are means of at least three independent experiments performed in triplicate, with standard deviation as indicated. ^d Staurosporine and galanthamine used as positive and negative controls, respectively.

with ED₅₀s determined at 12.5 and 15 μM, respectively [15a]. Interestingly, this activity was seen to be selective as normal human embryonic kidney (293t) cells remained unaffected by the treatment [15a]. Furthermore, haemanthamine (19) was shown to be active against apoptosis-sensitive human oligodendroglioma (Hs683) and mouse melanoma (B16F10) cancer cells with IC₅₀s of 7.0 and 6.8 μM, respectively [6h]. In addition, Berkov *et al.* reported that crinine (3) was active against a mini-panel of cancers comprising SKW-3, HL-60, HI-60/Dox and MDA-MB-231 cells with IC₅₀s ranging from 14.01 to 68.11 μM, with the best activity seen for SKW-3 cells [15d]. Further mechanistic tests via oligonucleosomal DNA fragmentation revealed the apoptotic cell death mode as responsible for its cytotoxicity [15d]. The antiproliferative activity of the rare Amaryllidaceae constituent distichamine (16), capable of inducing caspase-3 activated apoptosis in CEM cells, has recently been described [14].

Essential features of the crinane alkaloid cytotoxic pharmacophore which have come to the fore from these studies may be summarized as follows: i) the modulatory effect of the C-ring double bond (as seen in crinamine and crinine compared with their respective dihydro-analogues [15], as well as in distichamine in which the double bond features at C-2/C-3) [14]; ii) the effect of polar substitution at C-3, indicating that a H-bond acceptor is required at this position (as shown for crinamine, haemanthamine and distichamine, in possession of α-, β- and planar methoxy groups at C-3, respectively) [14,15]; iii) presence of a small substituent at C-11 (H or OH), the function of which does not appear to involve hydrogen bonding but is subject to steric constraints alone (evident for crinine, crinamine, haemanthamine and distichamine) [14,15]; and iv) oxidation at C-1 appears to be pivotal to activity {as in distichamine (16)} [14]. Furthermore, reversal of the polarity of the C-1 center in accommodating a hydroxyl group (as in distichaminol 17) is here shown to be detrimental.

In summary, phytochemical investigation of the rare and threatened southern African Amaryllid *Boophone haemanthoides* has led to the identification of lycorine together with seven members of the crinane series, of which distichaminol was uncovered as a novel compound. This compound is unique within the crinane series of Amaryllidaceae alkaloids in that its C-ring double bond is situated in the unusual C-2/C-3 position (as opposed to the usual C-1/C-2 position), thus accommodating the C-1 hydroxyl as well as the C-3 vinylic methoxyl groups. It is only the second member of the crinane alkaloids identified with such a substitution, the other being distichamine. Distichaminol was shown to be non-toxic to both cancer and normal cells, which contributes further to the understanding of the pharmacophoric requirements for cytotoxic activity as well as apoptosis induction within the crinane series of Amaryllidaceae alkaloids.

Experimental

General: Melting points (uncorrected) were measured on a Gallenkamp melting point apparatus. Optical rotations were determined on a Perkin-Elmer 241 polarimeter installed with a λ₅₈₉ sodium lamp. Circular dichroism (CD) spectra were obtained with a Jasco J-700 spectropolarimeter. IR spectra were measured on a Bio-Rad FTS-40 series spectrometer in dry film. EIMS were run on a Micromass Quattro Ultima spectrometer fitted with a direct injection probe (DIP) with ionization energy set at 70 eV and HRMS (ES) were performed with a Micromass Q-ToF Ultima spectrometer. ¹H, ¹³C NMR, DEPT, COSY, NOESY, HSQC and HMBC spectra were recorded on a Bruker AV400 in CDCl₃ or DMSO-*d*₆, chemical shifts are reported in units of δ (ppm) and coupling constants (*J*) are expressed in Hz. Silica gel Merck KGaA (70-230 mesh) was used for CC and TLC silica gel 60 F₂₅₄ for analyt. and prep. TLC (both Merck KGaA). Gel filtration was carried out on Sephadex LH-20 (Pharmacia). Spots on chromatograms were detected under UV light (254 and 365 nm) and by Dragendorff's reagent stain.

Plant material: Bulbs of *Boophone haemanthoides* were collected during the flowering season in September 2011 in the Nieuwoudtville area of the Northern Cape Province of South Africa. A voucher specimen (Snijman 588 NBG) was authenticated by Dr John C. Manning and deposited at the South African National Botanical Institute (SANBI).

Extraction and isolation of alkaloids: Powdered dried bulbs (784 g) of *Boophone haemanthoides* were extracted by stirring with EtOH for 48 h after which the solvent was evaporated under reduced pressure. The residue (61 g) was dissolved in H₂O (200 mL) and acidified to pH 4 with glacial acetic acid. After removing neutral material with Et₂O, the acidic solution was extracted with 3 x 200 mL CHCl₃ to provide extract A (6.1 g). Basifying the solution to pH 9 with aq. ammonia and extracting again with CHCl₃ (3 x 200 mL) afforded extract B (4.8 g). Finally, a CHCl₃-MeOH (3:2) extraction of the basic solution gave extract C (2.4 g). Extracts A, B and C were combined (13.3 g) after their TLC profiles (run in a 1:1 mixture of EtOAc-MeOH) were seen to be similar and subjected to gravity CC on silica gel by gradient elution with EtOAc and EtOAc-MeOH mixtures at a collection volume of 100 mL per flask. In this manner, 6 pooled fractions were generated after sequential TLC analysis giving FrI (0.71 g, flasks 24-26), FrII (0.81 g, flasks 27-28), FrIII (1.68 g, flasks 29-33), FrIV (2.06 g, flasks 34-41), FrV (0.56 g, flasks 42-50), FrVI (0.67 g, flasks 51-70). Further purification of fractions I-VI via a combination of Sephadex LH-20 CC (MeOH) and PTLC (MeOH-EtOAc/1:1) led to the consecutive isolation of distichamine [6.5 mg, R_f 0.54 (MeOH-EtOAc/1:1)], lycorine (2) [403.7 mg, R_f 0.45 (MeOH-EtOAc/1:1)], undulatine (14) [100.1 mg, R_f 0.44 (MeOH-EtOAc/1:1)], buphanidine (12)

[222.7 mg, R_f 0.39 (MeOH-EtOAc/1:1)], ambelline (**13**) [103.2 mg, R_f 0.38 (MeOH-EtOAc/1:1)], buphanisine (**11**) [82.6 mg, R_f 0.36 (MeOH-EtOAc/1:1)], crinine (**3**) [4.3 mg, R_f 0.33 (MeOH-EtOAc/1:1)] and distichaminol (**17**) [7.8 mg, R_f 0.30 (MeOH-EtOAc/1:1)]. The physical and spectroscopic data for the known compounds closely matched those that have been published [12,16].

Distichaminol (**17**)

MP: 76–78°C.

$[\alpha]_D^{25}$: -43.5 (c 0.2 in CHCl_3).

$\text{CD}[\theta]_D^{25}$: $[\theta]_{250} +2345$, $[\theta]_{286} -1787$ (c 0.2 in CHCl_3).

IR $\nu_{\text{max}}/\text{cm}^{-1}$ (dry film): 3500–3000 (OH), 2926, 1674, 1617 (Ar), 1469, 1376, 1315, 1270, 1211, 1127, 1039 (OCH_3), 934 (OCH_2O).

$^1\text{H NMR}$ (400 MHz, CDCl_3): Table 1.

$^{13}\text{C NMR}$ (100 MHz, CDCl_3): Table 1.

HRMS (Tof MS ES^+): calcd. 332.1498 for $\text{C}_{18}\text{H}_{22}\text{NO}_5$, found 332.1491 g/mol.

LRMS (EI) 70 eV, m/z (rel. int.): 331 $[\text{M}]^+$ (89), 316 $[\text{M}-\text{CH}_3]^+$ (28), 299 $[\text{M}-\text{CH}_3\text{OH}]^+$ (60), 281 $[\text{M}-\text{CH}_3\text{OH}-\text{H}_2\text{O}]^+$ (41), 270 (21), 253 (30), 242 (31), 231 (25), 218 (14), 207 (100), 191 (16), 179 (7), 165 (19), 135 (19), 115 (34).

Cytotoxicity bioassays

Cell cultures: Stock solutions (10 mmol/L) of test compounds were prepared in dimethylsulfoxide (DMSO). Dulbecco's modified Eagle's medium (DMEM), fetal bovine serum (FBS), L-glutamine, penicillin and streptomycin were purchased from Sigma (MO, USA). Calcein AM was obtained from Molecular Probes (Invitrogen Corporation, CA, USA). Test cell lines comprising T-lymphoblastic leukemia CEM, breast carcinoma MCF7, cervical carcinoma HeLa and human fibroblasts BJ were obtained from the American Type Culture Collection (Manassas, VA, USA). All cell

lines were cultured in DMEM medium (Sigma, MO, USA), supplemented with 10% heat-inactivated fetal bovine serum, 2 mmol/L L-glutamine, 10,000U penicillin and 10 mg/mL streptomycin. Cell lines were maintained under standard cell culture conditions (37°C and 5% CO_2 in a humid environment), and were sub-cultured twice or thrice weekly as required according to standard trypsinization procedures [14].

Calcein AM assay: Cell line suspensions containing about 1.0×10^5 cells/mL were placed in 96-well microtiter plates and after 24 h stabilization (time zero), test compounds, which were serially diluted in DMSO, were added (4 x 20 μL aliquots). Control cultures were treated with DMSO alone, such that the final DMSO concentration in the incubation mixtures never exceeded 0.6%. Test compounds were typically evaluated at six 3-fold dilutions and the highest final concentration was generally 50 μM . After 72 h incubation, 100 μL of 2 μM Calcein AM solution (Molecular Probes, Invitrogen, CA, USA) was added and incubation was continued for a further hour. The fluorescence of viable cells was then quantified using a Fluoroskan Ascent instrument (LabSystems, Finland). The percentage of surviving cells in each well was calculated by dividing the intensity of the fluorescence signals from the exposed wells by the intensity of signals from control wells and multiplying by 100. These ratios were then used to construct dose-response curves from which IC_{50} values were calculated. Staurosporine (Sigma-Aldrich, UK) and galanthamine (Teva Czech Industries, Opava, Czech Republic) were used as positive and negative controls, respectively.

Acknowledgment - JJN and JVS thank the University of KwaZulu-Natal for financial assistance. The Centre of the Region Haná for Biotechnological and Agricultural Research (grant No. ED0007/01/01) is gratefully acknowledged by LR and MS.

References

- [1] Meerow AW, Snijman DA. (1998) Amaryllidaceae. In *The Families and Genera of Vascular Plants*. Vol. 3, Kubitzki K. (Ed.). Springer, Berlin, Germany, 83–110.
- [2] (a) Viladomat F, Bastida J, Codina C, Nair JJ, Campbell WE. (1997) Alkaloids of the South African Amaryllidaceae. In *Recent Research Developments in Phytochemistry*. Vol. 1, Pandalai SG. (Ed.). Research Signpost Publishers, Trivandrum, India. 131–171; (b) Nair JJ, Bastida J, Codina C, Viladomat F, Van Staden J. (2013) Alkaloids of the South African Amaryllidaceae: a review. *Natural Product Communications*, 8, 1335–1350; (c) Bastida J, Lavilla R, Viladomat F. (2006) Chemical and biological aspects of *Narcissus* alkaloids. In *The Alkaloids*. Cordell GA. (Ed.). Elsevier, Amsterdam, Netherlands. 87–179.
- [3] (a) Heinrich M, Teoh HL. (2004) Galanthamine from snowdrop- the development of a modern drug against Alzheimer's disease from local Caucasian knowledge. *Journal of Ethnopharmacology*, 92, 147–162; (b) Houghton PJ, Ren Y, Howes M-J. (2006) Acetylcholinesterase inhibitors from plants and fungi. *Natural Product Reports*, 23, 181–199.
- [4] (a) Rinner U, Hudlicky T. (2005) Synthesis of Amaryllidaceae constituents-an update. *Synlett*, 3, 365–387; (b) Ingrassia L, Lefranc F, Mathieu V, Darro F, Kiss R. (2008) Amaryllidaceae isocarboxystiril alkaloids and their derivatives as promising antitumor agents. *Translational Oncology*, 1, 1–13; (c) Kornienko A, Evidente A. (2008) Chemistry, biology and medicinal potential of narciclasine and its congeners. *Chemical Reviews*, 108, 1982–2014; (d) Evidente A, Kornienko A. (2009) Anticancer evaluation of structurally diverse Amaryllidaceae alkaloids and their synthetic derivatives. *Phytochemistry Reviews*, 8, 449–459.
- [5] (a) Hwang YC, Chu JJH, Yang PL, Chen W, Yates MV. (2008) Rapid identification of inhibitors that interfere with poliovirus replication using a cell-based assay. *Antiviral Research*, 77, 232–236; (b) Del Giudice L, Massardo DR, Pontieri P, Wolf K. (2005) Interaction between yeast mitochondrial and nuclear genomes: Null alleles of *RTG* genes affect resistance to the alkaloid lycorine in *Saccharomyces cerevisiae*. *Gene*, 354, 9–14; (c) Mackey ZB, Baca AM, Mallari JP, Apsel B, Shelat A, Hansell EJ, Chiang PK, Wolff B, Guy KR, Williams J, McKerron JH. (2006) Discovery of trypanocidal compounds by whole cell HTS of *Trypanosoma brucei*. *Chemical Biology & Drug Design*, 67, 355–363; (d) Citoglu G, Tanker M, Gamusel B. (1998) Antiinflammatory effects of lycorine and haemanthidine. *Phytotherapy Research*, 12, 205–206; (e) Evidente A, Arrigoni O, Luso R, Calabrese G, Randazzo G. (1986) Further experiments on structure-activity relationships among lycorine alkaloids. *Phytochemistry*, 25, 2739–2743; (f) De Leo P, Dalessandro G, De Santis A, Arrigoni O. (1973) Inhibitory effect of lycorine on cell division and cell elongation. *Plant and Cell Physiology*, 14, 481–486; (g) Nair JJ, Van Staden J. (2012) Acetylcholinesterase inhibition within the lycorine series of Amaryllidaceae alkaloids. *Natural Product Communications*, 7, 959–962; (h) Jimenez A, Santos A, Alonso G, Vazquez D. (1976) Inhibitors of protein synthesis in eukaryotic cells. Comparative effects of some Amaryllidaceae alkaloids. *Biochimica et Biophysica Acta*, 425, 342–348.
- [6] (a) Likhitwitayawuid K, Angerhofer CK, Chai H, Pezzuto JM, Cordell GA, Ruangrunsi N. (1993) Cytotoxic and antimalarial alkaloids from the bulbs of *Crinum amabile*. *Journal of Natural Products*, 56, 1331–1338; (b) Lamoral-Theys D, Decaestecker C, Mathieu V, Dubois J, Kornienko A, Kiss R, Evidente A, Pottier L. (2010) Lycorine and its derivatives for anticancer drug design. *Mini-Reviews in Medicinal Chemistry*, 10, 41–50; (c) Evdokimov NM, Lamoral-Theys D, Mathieu V, Andolfi A, Frolova LV, Pelly SC, Van Otterlo WAL, Magedov IV, Kiss R, Evidente A, Kornienko A. (2011) In search of a cytostatic agent derived from the alkaloid lycorine: synthesis and growth inhibitory properties of lycorine derivatives. *Bioorganic & Medicinal Chemistry*, 19, 7252–7261; (d) Liu J, Hu W, He L, Ye M, Li Y. (2004) Effects of lycorine on HL-60 cells via arresting cell

- cycle and inducing apoptosis. *FEBS letters*, **578**, 245-250; (e) Liu J, Li Y, Tang LJ, Zhang GP, Hu WX. (2007) Treatment of lycorine on SCID mice model with human APL cells. *Biomedicine & Pharmacotherapy*, **61**, 229-234; (f) Li S, Chen C, Zhang H, Guo H, Wang H, Wang L, Zhang X, Hua S, Yu J, Xiao P, Li Y, Liu J, Tang L, Shi Y, Ren W, Hu W. (2007) Apoptosis induced by lycorine in KM3 cells is associated with the G0/G1 cell cycle arrest. *Oncology Reports*, **17**, 377-384; (g) McNulty J, Nair JJ, Bastida J, Pandey S, Griffin C. (2009) Structure-activity studies on the lycorine pharmacophore: a potent inducer of apoptosis in human leukemia cells. *Phytochemistry*, **70**, 913-919; (h) Van Goietsenoven G, Andolfi A, Lallemand B, Cimmino A, Lamoral-Theys D, Gras T, Abou-Donia A, Dubois J, Lefranc F, Mathieu V, Kornienko A, Kiss R, Evidente A. (2010) Amaryllidaceae alkaloids belonging to different structural subgroups display activity against apoptosis-resistant cancer cells. *Journal of Natural Products*, **73**, 1223-1227.
- [7] (a) Pettit GR, Gaddamidi V, Herald DL, Singh SB, Cragg GM, Schmidt JM, Boettner FE, Williams M, Sagawa Y. (1986) Antineoplastic agents. 120. *Pancreaticum littorale*. *Journal of Natural Products*, **49**, 995-1002; (b) Pettit GR, Pettit GR III, Backhaus RA, Boyd MR, Meerow AW. (1993) Cell growth inhibitory isocarboxtyrils from *Hymenocallis*. *Journal of Natural Products*, **56**, 1682-1687; (c) Ingrassia L, Lefranc F, Dewelle J, Pottier L, Mathieu V, Spiegel-Kreinecker S, Sauvage S, El Yazidi M, Dehoux M, Berger W, Van Quaquebeke E, Kiss R. (2009) Structure-activity relationship analysis of novel derivatives of narciclasine (an Amaryllidaceae isocarboxtyril derivative) as potential anticancer agents. *Journal of Medicinal Chemistry*, **52**, 1100-1114; (d) Van Goietsenoven G, Hutton J, Becker J-P, Lallemand B, Robert F, Lefranc F, Pirker C, Vandenbussche G, Van Antwerpen P, Evidente A, Berger W, Prévost M, Pelletier J, Kiss R, Kinzy TG, Kornienko A, Mathieu V. (2010) Targeting of eEF1A with Amaryllidaceae isocarboxtyrils as a strategy to combat melanomas. *The FASEB Journal*, **24**, 4575-4584; (e) McLachlan A, Kekre N, McNulty J, Pandey S. (2005) Pancratistatin: A natural anticancer compound that targets mitochondria specifically in cancer cells to induce apoptosis. *Apoptosis*, **10**, 619-630; (f) Kekre N, Griffin C, McNulty J, Pandey S. (2005) Pancratistatin causes early activation of caspase-3 and the flipping of phosphatidyl serine followed by rapid apoptosis specifically in human lymphoma cells. *Cancer Chemotherapy & Pharmacology*, **56**, 29-38.
- [8] (a) McNulty J, Nair JJ, Singh M, Crankshaw DJ, Holloway AC, Bastida J. (2009) Selective cytochrome P450 3A4 inhibitory activity of Amaryllidaceae alkaloids. *Bioorganic and Medicinal Chemistry Letters*, **19**, 3233-3237; (b) McNulty J, Thorat A, Vurgun N, Nair JJ, Makaji E, Crankshaw DJ, Holloway AC, Pandey S. (2011) Human cytochrome P450 liability studies of *trans*-dihydronarciclasine: a readily available, potent and selective cancer cell growth inhibitor. *Journal of Natural Products*, **74**, 106-108.
- [9] (a) Tram NTM, Titorenkova TV, Bankova VS, Handjieva NV, Popov SS. (2002) *Crinum* L. (Amaryllidaceae). *Fitoterapia*, **73**, 183-208; (b) Nair JJ, Bastida J, Viladomat F, Van Staden J. (2012) Cytotoxic agents of the crinane series of Amaryllidaceae alkaloids. *Natural Product Communications*, **7**, 1677-1688.
- [10] (a) McNulty J, Nair JJ, Sliwinski M, Harrington LE, Pandey S. (2007) Unusual magnesium chloride catalyzed non-Evans *anti*-aldol reactions of an enolizable L-threose derivative. *European Journal of Organic Chemistry*, 5669-5673; (b) McNulty J, Nair JJ, Griffin C, Pandey S. (2008) Synthesis and biological evaluation of fully functionalized *seco*-pancratistatin analogues. *Journal of Natural Products*, **71**, 357-363; (c) McNulty J, Nair JJ, Singh M, Crankshaw DJ, Holloway AC. (2009) Structure-activity studies on *seco*-pancratistatin analogs: potent inhibitors of human cytochrome P450 3A4. *Bioorganic and Medicinal Chemistry Letters*, **19**, 5607-5612; (d) McNulty J, Nair JJ, Singh M, Crankshaw DJ, Holloway AC. (2010) Potent and selective inhibition of human cytochrome P450 3A4 by *seco*-pancratistatin analogs. *Bioorganic and Medicinal Chemistry Letters*, **20**, 2335-2339; (e) McNulty J, Nair JJ, Little JRL, Brennan JD, Bastida J. (2010) Structure-activity relationship studies on acetylcholinesterase inhibition in the lycorine series of Amaryllidaceae alkaloids. *Bioorganic and Medicinal Chemistry Letters*, **20**, 5290-5294; (f) McNulty J, Nair JJ, Singh M, Crankshaw DJ, Holloway AC, Bastida J. (2010) Cytochrome P450 3A4 inhibitory activity studies within the lycorine series of alkaloids. *Natural Product Communications*, **5**, 1195-1200.
- [11] (a) Wrinkle G. (1984) An introduction to the genus *Boophone*. *Herbertia*, **40**, 77-82; (b) Watt JM, Breyer-Brandwijk MG. (1962) *The Medicinal and Poisonous Plants of Southern and Eastern Africa*. Livingston Ltd., Edinburgh; (c) Hutchings A, Scott AH, Lewis G, Cunningham AB. (1996) *Zulu Medicinal Plants: an Inventory*. University of Natal Press, Pietermaritzburg; (d) Nair JJ, Manning JC, Van Staden J. (2012) Distichamine, a chemotaxonomic marker for the genus *Boophone* Herb. (Amaryllidaceae). *South African Journal of Botany*, **83**, 89-91.
- [12] (a) Sandager M, Nielsen ND, Stafford GI, Van Staden J, Jäger AK. (2005) Alkaloids from *Boophone disticha* with affinity to the serotonin transporter in rat brain. *Journal of Ethnopharmacology*, **98**, 367-370; (b) Neergaard JS, Andersen J, Pedersen ME, Stafford GI, Van Staden J, Jäger AK. (2009) Alkaloids from *Boophone disticha* with affinity to the serotonin transporter. *South African Journal of Botany*, **75**, 371-374; (c) Cheesman L, Nair JJ, Van Staden J. (2012) Antibacterial activity of crinane alkaloids from *Boophone disticha* (Amaryllidaceae). *Journal of Ethnopharmacology*, **140**, 405-408.
- [13] (a) Ali AA, Ramadan MA, Frahm AW. (1984) Alkaloid constituents of *Crinum bulbispermum* III: bulbispermimine, a new alkaloid of *Crinum bulbispermum*. *Planta Medica*, **50**, 424-427; (b) Wagner J, Pham HL, Döpke W. (1996) Alkaloids from *Hippeastrum equestre* Herb.-5. Circular dichroism studies. *Tetrahedron*, **52**, 6591-6600; (c) Nair JJ, Campbell WE, Brun R, Viladomat F, Codina C, Bastida J. (2005) Alkaloids from *Nerine filifolia*. *Phytochemistry*, **66**, 373-382.
- [14] Nair JJ, Rárová L, Strnad M, Bastida J, Van Staden J. (2012) Apoptosis-inducing effects of distichamine and narciprimine, rare alkaloids of the plant family Amaryllidaceae. *Bioorganic and Medicinal Chemistry Letters*, **22**, 6195-6199.
- [15] (a) McNulty J, Nair JJ, Codina C, Bastida J, Pandey S, Gerasimoff J, Griffin C. (2007) Selective apoptosis-inducing activity of crinum-type Amaryllidaceae alkaloids. *Phytochemistry*, **68**, 1068-1074; (b) Griffin C, Sharda N, Sood, D, Nair JJ, McNulty J, Pandey S. (2007) Selective cytotoxicity of pancratistatin-related natural Amaryllidaceae alkaloids: evaluation of the activity of two new compounds. *Cancer Cell International*, **7**, 10-16; (c) McNulty J, Nair JJ, Bastida J, Pandey S, Griffin C. (2009) Structure activity studies on the crinane alkaloid apoptosis-inducing pharmacophore. *Natural Product Communications*, **4**, 483-488; (d) Berkov S, Romani S, Herrera M, Viladomat F, Codina C, Momekov G, Ionkova I, Bastida J. (2011) Antiproliferative alkaloids from *Crinum zeylanicum*. *Phytotherapy Research*, **25**, 1686-1692.
- [16] (a) Campbell WE, Nair JJ, Gammon DW, Codina C, Bastida J, Viladomat F, Smith PJ, Albrecht CF. (2000) Bioactive alkaloids from *Brunsvigia radulosa*. *Phytochemistry*, **53**, 587-591; (b) Herrera MR, Machocho AK, Nair JJ, Campbell WE, Brun R, Viladomat F, Codina C, Bastida J. (2001) Alkaloids from *Cyrtanthus elatus*. *Fitoterapia*, **72**, 444-448; (c) Viladomat F, Codina C, Bastida J, Mathee S, Campbell WE. (1995) Further alkaloids from *Brunsvigia josephinae*. *Phytochemistry*, **40**, 961-965.

Supinidine Viridiflorates from the Roots of *Chromolaena pulchella*Mario A. Gómez-Hurtado^{a,b}, J. Martín Torres-Valencia^{a,*}, Rosa E. del Río^b, Gabriela Rodríguez-García^b, Virginia Motilva^c, Sofía García-Mauriño^d, Carlos M. Cerda-García-Rojas^e and Pedro Joseph-Nathan^e^aÁrea Académica de Química, Universidad Autónoma del Estado de Hidalgo, Km 4.5 Carretera Pachuca-Tulancingo, Mineral de la Reforma, Hidalgo 42184, Mexico^bInstituto de Investigaciones Químico-Biológicas, Universidad Michoacana de San Nicolás de Hidalgo, Apartado 137, Morelia, Michoacán 58000, Mexico^cFacultad de Farmacia, Universidad de Sevilla, Profesor García González No. 2, Sevilla 41012, Spain^dFacultad de Biología, Universidad de Sevilla, Profesor García González No. 2, Sevilla 41012, Spain^eDepartamento de Química, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Apartado 14-740, México, D. F., 07000 Mexico

jmartin@uaeh.edu.mx


Received: September 13th, 2013; Accepted: September 18th, 2013

The alkaloid extract from the roots of *Chromolaena pulchella* provided two new pyrrolizidine alkaloids, elucidated as (–)-supinidine triviridiflorate (**1**) and (–)-supinidine diviridiflorate (**2**) based on their physical and spectroscopic properties. Their absolute configuration was determined by chemical correlation with (–)-supinidine (**3**) and (+)-viridifloric acid (**4**).

Keywords: *Chromolaena pulchella*, Asteraceae, Pyrrolizidine alkaloids, Absolute configuration, NMR spectroscopy.

Pyrrolizidine alkaloids (PAs), although widely distributed, are characteristic of certain genera of the Boraginaceae, Leguminosae, and Asteraceae/Compositae families [1]. Many PAs are known to produce hepatic toxicity and there are several records of livestock poisoning [1a,b]. Chemical studies of some *Chromolaena* species (Asteraceae) showed the presence of this class of natural molecules. Thus *N*-oxides of 7-angeloylretronecine, intermidine, licopsamine, echinatine, 3'-acetylretronecine, and supinine have been identified in *C. odorata* [1c]. The present work describes the isolation of the new PAs supinidine triviridiflorate (**1**) and supinidine diviridiflorate (**2**) (Figure 1) from the crude alkaloid extract of the roots of *C. pulchella*, for which we recently reported the isolation of labdanes and *ent*-clerodanes from the aerial parts [1d].

Compound **1**, a pale yellow oil, showed a $[M + 1]^+$ ion at m/z 572.3427 in its HRESI/APCI mass spectrum revealing the molecular formula $C_{29}H_{49}NO_{10} + H$ (calcd m/z 572.3435). The 1H NMR chemical shift values of the alkaloid moiety were in agreement with those for a supinidine type pyrrolizidine ring system [2]. The complete 1H and ^{13}C data, shown in Table 1, gave characteristic signals for a 1,2-unsaturated pyrrolizidine alkaloid with a necic acid esterified at C-9. The 1H NMR spectrum of **1** showed signals assignable to a vinylic proton at δ_H 5.71 (H-2) and an AB system ($J = 13.6$ Hz) at δ_H 4.72 and 4.60 due to protons of the C-9 hydroxymethylene group, whereas the signal for the hydrogen atom attached to bridgehead C-8 was observed at δ_H 4.15. The signals for the methylene groups at C-3 and C-5, bearing the nitrogen atom, were observed at δ_H 3.91 and 3.37, and at 3.11 and 2.53, respectively. Moreover, signals for a trimeric α -isopropyl- α,β -dihydroxybutyric acid residue were observed as quartets at δ_H 5.35 ($J = 6.2$ Hz, H-13), 4.98 ($J = 6.6$ Hz, H-20), and 4.09 ($J = 6.6$ Hz, H-27) showing strong correlation in the COSY spectrum with the doublets at δ_H 1.39 (Me-14), 1.22 (Me-21), and 1.21 (Me-28), respectively. The ^{13}C and APT NMR spectra showed signals for three carbonyl carbon atoms at δ_C 174.0 (C-18), 173.6 (C-25), and



173.3 (C-11), for two vinylic carbons at 136.9 (C-1) and 126.1 (C-2), for three quaternary carbons bearing oxygen atoms at 82.6 (C-26), 81.7 (C-12), and 80.3 (C-19), for seven methine carbons, three of them bearing oxygen atoms at 76.4 (C-13), 71.5 (C-20) and 69.4 (C-27), one bearing the nitrogen atom at 71.5 (C-8), and three owing to the isopropyl groups at 33.8 (C-15), 32.7 (C-29) and 30.9 (C-22), for five methylene carbons, one of them bearing an oxygen atom at 61.9 (C-9), two bearing the nitrogen atom at 61.6 (C-3) and 56.6 (C-5), and two at 30.2 (C-7) and 25.7 (C-6), and for six methyl groups owing to three isopropyl groups at 17.3, 17.0, 16.9, 16.8, 16.6 and 15.3. Esterification at the C-9 position was confirmed by the HMBC correlation between H_2 -9 and the carbonyl group C-11, while the individual assignments for the three acid residues were supported from HMBC correlations of H-13 with C-11 and C-12, of H-20 with C-18 and C-19, and of H-27 with C-25 and C-26.

Compound **2** exhibited in its HRESI/APCI mass spectrum a $[M + 1]^+$ ion at m/z 428.2646 in agreement with the molecular formula $C_{22}H_{37}NO_7 + H$ (calcd m/z 428.2648). The complete 1H and ^{13}C data shown in Table 1 demonstrated high structural similarities

Table 1. ^{13}C and ^1H NMR data for **1** and **2** (100 and 400 MHz, CDCl_3).^a

Position	1		2	
	δ_{C} , mult.	δ_{H} , mult. (<i>J</i> in Hz)	δ_{C} , mult.	δ_{H} , mult. (<i>J</i> in Hz)
1	136.9, C		136.9, C	
2	126.1, CH	5.71, br s	125.7, CH	5.73, br s
3	61.6, CH_2	3.91, br d (15.8)	61.2, CH_2	3.95, br d (15.8)
3'		3.37, dd (15.8, 4.4)		3.40, br d (15.8)
5	56.6, CH_2	3.11, dt (9.9, 5.5)	56.6, CH_2	3.18, dt (10.0, 5.1)
5'		2.53, dt (9.9, 7.0)		2.54, dt (10.0, 7.0)
6	25.7, CH_2	1.80, m	25.7, CH_2	1.81, m
7	30.2, CH_2	1.98, m	29.9, CH_2	2.04, m
7'		1.50, m		1.54, m
8	71.5, CH	4.15, br m	71.5, CH	4.25, br m
9	61.9, CH_2	4.72, br d (13.6)	62.2, CH_2	4.80, br d (13.5)
9'		4.60, br d (13.6)		4.74, br d (13.5)
11	173.3, C		174.0, C	
12	81.7, C		81.2, C	
13	76.4, CH	5.35, q (6.2)	73.9, CH	5.26, q (6.2)
14	13.1, CH_3	1.39, d (6.2)	13.9, CH_3	1.32, d (6.2)
15	33.8, CH	1.98, sept (7.0)	32.6, CH	2.09, sept (7.0)
16	17.0, CH_3	1.05, d (7.0)	16.4, CH_3	0.94, d (7.0)
17	17.3, CH_3	0.96, d (7.0)	17.1, CH_3	0.96, d (7.0)
18	174.0, C		174.3, C	
19	80.3, C		82.4, C	
20	71.5, CH	4.98, q (6.6)	69.2, CH	3.95, q (6.6)
21	14.6, CH_3	1.22, d (6.6)	17.5, CH_3	1.17, d (6.6)
22	30.9, CH	2.09, sept (7.0)	32.3, CH	2.08, sept (7.0)
23	15.3, CH_3	0.86, d (7.0)	16.4, CH_3	0.92, d (7.0)
24	16.9, CH_3	0.74, d (7.0)	16.9, CH_3	0.87, d (7.0)
25	173.6, C			
26	82.6, C			
27	69.4, CH	4.09, q (6.6)		
28	17.2, CH_3	1.21, d (6.6)		
29	32.7, CH	2.09, sept (7.0)		
30	16.8, CH_3	0.94, d (7.0)		
31	16.6, CH_3	1.00, d (7.0)		

^aAssigned by gHMBC and gHMBC.

with the aforementioned PA **1**, implying that compounds **1** and **2** belong to the same class of alkaloids. Compound **2** differed from **1** in the absence of the third α -isopropyl- α,β -dihydroxybutyric acid residue, since in the ^1H NMR spectrum only two methine signals for protons bearing oxygen atoms were observed at δ_{H} 5.26 (q, *J* = 6.2 Hz, H-13), and 3.95 (q, *J* = 6.6 Hz, H-20), whereas in the ^{13}C NMR spectrum only two carbonyl groups signals were observed at δ_{C} 174.0 (C-11), and 174.3 (C-18). As in the case of compound **1**, 2D NMR spectroscopy was employed to completely assign the ^{13}C and ^1H spectra. Alkaline hydrolysis of a mixture of compounds **1** and **2** gave the necine base (–)-supinidine (**3**) [2,3a] and the necic acid residue (+)-viridifloric acid (**4**) [3b].

Experimental

General: Optical rotation, Perkin-Elmer 341 polarimeter; IR, Perkin-Elmer 16F PC IR-FT spectrophotometer using thin films of compounds deposited on a CsI crystal; Low-resolution MS, either Agilent 1100 LC/MSD or Varian Saturn 2000 spectrometers; HRMS, Agilent LCTOF instrument; NMR, JEOL Eclipse 400 spectrometer; CC, Merck silica gel 40; TLC, silica gel 60 precoated glass plates.

Plant material: Specimens of *C. pulchella* (H.B.K.) R.M. King & H. Rob. (Asteraceae) were collected near km 61 of Morelia-Zacapu federal road 15, in the municipality of Constitución, State of Michoacán, México, during October 2005. A specimen (No. 192522) is deposited at the Herbarium of Instituto de Ecología A.

References

- (a) Hartmann T. (1999) Chemical ecology of pyrrolizidine alkaloids. *Planta*, **207**, 483–495; (b) Chen T, Mei N, Fu PP. (2009) Genotoxicity of pyrrolizidine alkaloids. *Journal of Applied Toxicology*, **30**, 183–196; (c) Biller A, Boppré M, Witte L, Hartmann T. (1994) Pyrrolizidine alkaloids in *Chromolaena odorata*. Chemical and chemoeological aspects. *Phytochemistry*, **35**, 615–619; (d) Gómez-Hurtado MA, Torres-Valencia JM, Manríquez-Torres J, del Río RE, Motilva V, García-Mauriño S, Ávila J, Talero E, Cerda-García-Rojas CM, Joseph-Nathan P. (2011) Absolute configuration of labdanes and ent-clerodanes from *Chromolaena pulchella* by vibrational circular dichroism. *Phytochemistry*, **72**, 409–414.
- Logie CG, Grue MR, Liddell JR. (1994) Proton NMR spectroscopy of pyrrolizidine alkaloids. *Phytochemistry*, **37**, 43–109.
- (a) Gruszecka-Kowalik E, Zalkow LH. (1990) Free-radical reactions of retronecine and heliotridine derivatives. The synthesis of (–)-supinidine. *The Journal of Organic Chemistry*, **55**, 3398–3403; (b) Stritzke K, Schulz S, Nishida R. (2002) Absolute configuration and synthesis of β - and δ -lactones present in the pheromone system of the giant white butterfly *Idea leuconoe*. *European Journal of Organic Chemistry*, 388–392.
- Schulz S, Nishida R. (1996) The pheromone system of the male danaine butterfly, *Idea leuconoe*. *Bioorganic & Medicinal Chemistry*, **4**, 341–349.

C., Centro Regional del Bajío, Pátzcuaro, Michoacán, Mexico, where Prof. Jerzy Rzedowski kindly identified the plant material.

Extraction and isolation: Air-dried and powdered roots of *C. pulchella* (870 g) were extracted with MeOH (3.5 L) under reflux for 6 h. Filtration and evaporation of the extract afforded a yellow viscous oil (38.6 g) which gave a positive Dragendorff test. To this product aq. HCl 2% (300 mL) and zinc powder (40 g) were gradually added under stirring for 12 h. The acidic aqueous solution was treated with aq. KOH 5% to obtain a phase with pH 10 and then extracted with CHCl_3 (3 \times 200 mL), dried over anhydrous Na_2SO_4 , filtered, and evaporated to provide 2.6 g of residue. A portion of this (876 mg) was subjected to chromatography on silica gel 60 (20 g) using CHCl_3 –MeOH–diethylamine (98:2:1, 96:4:1, 90:10:1, and 80:20:1). Fractions of 100 mL of each polarity were collected, monitored by TLC, and analyzed by ^1H NMR spectroscopy. The resulting material from each fraction was labelled as A (138 mg), B (354 mg), C (112 mg) and D (116 mg). From fractions A, C, and D fatty materials were isolated. Separation of fraction B (178 mg) by means of preparative TLC using CHCl_3 –MeOH (9:1) as the mobile phase gave pure **1** (45 mg, *R_f* 0.5). Another portion of fraction B (138 mg) was purified by TLC (CHCl_3 –MeOH– H_2O , 200:50:7) affording **2** (20 mg, *R_f* 0.2).

Supinidine triviridiflorate (**1**)

Pale yellow oil; $[\alpha]_{\text{D}}^{20}$: –1.5 (c 3.5, CHCl_3)

IR (film): 3518, 2972, 2937, 2878, 1725, 1454, 1386 cm^{-1} .

^1H and ^{13}C NMR: Table 1.

MS (EI, 70 eV): *m/z* (%) 572 [*M* + 1]⁺ (40), 428 (2), 410 (4), 284 (3), 266 (19), 140 (19), 122 (100), 110 (10), 107 (6), 94 (25), 70 (38); HRESI/APCIMS: *m/z* [*M* + 1]⁺ calcd for $\text{C}_{29}\text{H}_{49}\text{NO}_{10}$ + H: 572.3435; found: 572.3427.

Supinidine diviridiflorate (**2**)

Pale yellow oil; $[\alpha]_{\text{D}}^{20}$: –8.2 (c 2.0, CHCl_3).

IR (film): 3024, 2970, 2945, 1727, 1456, 1389 cm^{-1} .

^1H and ^{13}C NMR: Table 1.

MS (EI, 70 eV): *m/z* (%) = 428 [*M* + 1]⁺ (2), 396 (2), 382 (2), 284 (28), 224 (6), 140 (19), 122 (100); HRESI/APCIMS: *m/z* [*M* + 1]⁺ calcd for $\text{C}_{22}\text{H}_{37}\text{NO}_7$ + H: 428.2648; found: 428.2646.

Hydrolysis of compounds 1 and 2: A mixture of **1** and **2** (40 mg) in MeOH (3 mL) was treated with NaOH (24 mg) in H_2O (0.25 mL) and heated to reflux for 15 min in a micro-wave system working at 100 W. The mixture was treated with H_2O (10 mL) and extracted with CH_2Cl_2 (2 \times 10 mL) to give (–)-supinidine (**3**) (3.5 mg) as a yellow oil, which showed $[\alpha]_{\text{D}}^{20}$ –9.1 (c 0.16, EtOH) lit. $[\alpha]_{\text{D}}^{20}$ –10.4 (c 2.64, EtOH) [3a]. The aq. phase was acidified with 2% HCl and extracted with CH_2Cl_2 (2 \times 10 mL) to give (+)-viridifloric acid (**4**) (24 mg) [3b,4].

Acknowledgments - Partial financial support from Conacyt, Mexico (grant No. U2-80555), from ICYTDF (grant No. 337/2010), and PROMEP-UMSNH (grant No. 332) is acknowledged.

N-Containing Metabolites from the Marine Sponge *Agelas clathrodes*Fan Yang^a, Rui-Hua Ji^b, Jiang Li^c, Jian-Hong Gan^d and Hou-Wen Lin^{a,*}^aDepartment of Pharmacy, Renji Hospital, Shanghai Jiao Tong University School of Medicine, 160 Pujian Road, Shanghai 200127, P.R. China^bDepartment of Anesthesiology, Changzheng Hospital, Second Military Medical University, 415 Fengyang Road, Shanghai 200003, P.R. China^cEastern Hepatobiliary Surgical Hospital, Second Military Medical University, 225 Changhai Road, Shanghai 200438, P.R. China^dCollege of Food Science & Technology, Shanghai Ocean University, 999 Hucheng Huan Road, Shanghai 201306, P.R. China

Fan Yang and Rui-Hua Ji contributed equally to this work.

franklin67@126.com

Received: July 28th, 2013; Accepted: October 1st, 2013

A new bisuracil analogue, 3,3-bis(uracil-1-yl)-propan-1-aminium (**1**), together with four known *N*-containing metabolites (**2–5**), were isolated from the South China Sea sponge *Agelas clathrodes*. Their chemical structures were established on the basis of spectroscopic and spectrometric analysis and comparison with known compounds. Compound **1** is an unusual naturally occurring bisuracil analogue, and compound **2** was isolated from a natural source for the first time. Compounds **2** and **4** exhibit moderate cytotoxicity against cancer cell line SGC7901.

Keywords: *Agelas clathrodes*, Bisuracilyl-substituted, *N*-containing, Marine sponge.

Marine sponges of the genus *Agelas* (order Agelasida, family Agelasidae) have provided a wide variety of natural products, such as glycosphingolipids [1a,b], bromopyrrole alkaloids [1c,d] and 9-*N*-methyladeninium diterpenoids [2a-c]. *A. clathrodes* (Schmidt, 1870) was reported to possess the *N*-containing metabolites, clathrocin [3a], clathramides [3b], (-)-agelasidine A [3c], and dispacamides [3d,e]. As part of an ongoing investigation, studies on *A. clathrodes* led to the isolation of a new bisuracilyl-substituted analogue, 3,3-bis(uracil-1-yl)-propan-1-aminium (**1**), along with four known compounds (**2–5**) (Figure 1).

Compound **1**, an amorphous solid, was deduced as C₁₁H₁₄N₅O₄ by HR-APCI-MS (*m/z* 280.1213 [M]⁺; calcd. 280.1202). The ¹H NMR spectrum displayed signals at δ_H 5.44 (2H, d, *J* = 7.5 Hz) and δ_H 7.38 (2H, d, *J* = 7.5 Hz), which were characteristic of two symmetrical uracil moieties. The symmetry of **1** resulted in a simplified ¹³C NMR spectrum with perfect overlap of the signals. (Table 1), including those of two quaternary carbons, three methines and two methylenes. The uracil units were confirmed by comparing the ¹³C NMR spectral data (δ_C 164.3, 151.5, 142.1 and 100.2) with that in the literature [3f]; the signal intensities of the downfield carbons (δ_C 164.3, 151.5, 142.1 and 100.2) were all higher than those of the resonances at δ_C 72.4, 63.1 and 29.5. The ¹H-¹H COSY spectrum only showed correlation between H-5 (H-5') (δ_H 5.44) and H-6 (H-6') (δ_H 7.38). In the HMBC spectrum, the signal of H-5 correlated with those of C-4 and C-6, and HMBC cross-peaks of H-6 with C-2, C-4 and C-5 were also found. The HMBC correlation from the methine proton H-7 (δ_H 3.42) to C-6, as well as the downfield chemical shift of C-7 (δ_C 72.4) suggested that the uracil moieties were connected to C-7. In addition, the signal of H-7 further correlated with C-9. The downfield chemical shift of C-9 (δ_C 63.1) can be explained by the proximity of the aminium [3g]. On the basis of the foregoing analysis, compound **1** was determined as 3,3-bis(uracil-1-yl)-propan-1-aminium.




Figure 1: Structures of compounds 1–5.

Table 1: ¹H NMR (400 MHz) and ¹³C NMR (100 MHz) data for compound **1** in DMSO-*d*₆.

Position	δ _H (int., mult., <i>J</i> in Hz)	δ _C	HMBC (H→C)
2, 2'		151.5	
4, 4'		164.3	
5, 5'	5.44 (2H, d, 7.5)	100.2	4, 6
6, 6'	7.38 (2H, d, 7.5)	142.1	2, 4, 5
7	3.42 (1H, m)	72.4	6, 9
8	2.56 (2H, ov ^a)	29.5	
9	3.26 (2H, ov)	63.1	7
N-H	10.95 (ov)		

^a Overlapped with solvent signal and deduced from HMQC correlation.

Table 2: Cytotoxic activities of compounds 1–5 on seven human cancer cell lines.

Compound	Human cancer line, IC ₅₀ (μg/mL)						
	A549	H1299	SGC7901	LNCAP	PC3	MDA-231	CT-26
1	26.51	>50	22.73	45.02	34.54	>50	>50
2	29.77	>50	13.24	22.67	26.91	>50	40.5
3	23.86	>50	>50	>50	>50	39.38	25.09
4	>50	>50	16.56	27.16	>50	40.47	27.9
5	>50	>50	23.72	40.9	27.17	>50	>50
5-Fu	7.01	2.5	7.51	8.93	8.17	2.38	3.83

The structures of compounds **2–5** were elucidated by comparison of their ¹H NMR, ¹³C NMR and mass spectra with those in the literature [4]. Compounds **1–5** were assessed for cytotoxic activity (Table 2) against seven human cancer cell lines, including A549 (human alveolar basal epithelial), H1299 (human non-small cell lung carcinoma), SGC7901 (human gastric carcinoma), LNCAP (human prostate cancer), PC3 (human prostate cancer), MDA-231 (human breast cancer), and CT-26 (human colorectal cancer) *in vitro*. Compound **1** exhibited weak cytotoxicity to A549

(IC₅₀ = 26.5 µg/mL) and SGC7901 (IC₅₀ = 22.7 µg/mL). Compounds **2** and **4** displayed moderate cytotoxicity against SGC7901 (IC₅₀ = 13.2 and 16.6 µg/mL).

Experimental

General: Melting point, SGW X-4 melting point apparatus; Optical rotation, JASCO P-1030 polarimeter; HR-APCI-MS, ESI-MS and EI-MS, Q-ToF micro YA019 mass spectrometer; NMR, Bruker AVANCE-400 spectrometer; HPLC, Waters 1525/2996 liquid chromatograph; CC, Sephadex LH-20 (Pharmacia) and YMC ODS-A (50 µm). Fractions were monitored by TLC (HSGF 254, Yantai, China) and compounds were visualized by heating silica gel plates after spraying them with 10% H₂SO₄ in EtOH.

Animal material: Specimens of the marine sponge *Agelas clathrodes* were collected around Yongxing Island and Seven Connected Islets in the South China Sea in June 2007, and were identified (voucher sample No. V107) by Prof. Zhi-Yong Li (Marine Biotechnology Laboratory, Shanghai Jiao Tong University, Shanghai, China).

Extraction and isolation: *A. clathrodes* (240 g, wet weight) was extracted with acetone, CH₂Cl₂, and MeOH (500 mL × 2 each) exhaustively at room temperature. The combined extracts were concentrated under reduced pressure to give a brown gum, which was partitioned between CH₂Cl₂ and H₂O to afford 2.1 g CH₂Cl₂ phase extract. This was subjected to CC on Sephadex LH-20 with CH₂Cl₂/MeOH (1:1) as eluting solvent to afford 3 fractions (*Fr. A-C*). *Fr. A* (670 mg) was subjected to CC on YMC ODS-A (50 µm) to give 6 subfractions (*Fr. A1–Fr. A6*). *Fr. A1* (382.3 mg) was further purified by HPLC (Waters 1525/2996, Symmetry Prep C18, 7 µm, 1.5 mL/min, UV detection at 260 nm) using 10% MeOH/H₂O as eluent to yield compounds **1** (1.6 mg, *t_R* = 14.5 min) and **3** (20.5 mg, *t_R* = 28.9 min). *Fr. A5* (89.6 mg) was separated by HPLC (Waters 1525/2996, Symmetry Prep C18, 7 µm, 60% MeOH/H₂O, 1.5 mL/min, UV detection at 275 nm) to yield compound **4** (13.6 mg, *t_R* = 43.4 min). *Fr. A6* (30.4 mg) was

purified by HPLC (Waters 1525/2996, Symmetry Prep C18, 7 µm, 60% CH₃CN/H₂O, 1.5 mL/min, UV detection at 275 nm) to afford compound **5** (10.7 mg, *t_R* = 27.1 min). Similarly, *Fr. B* (170 mg) was subjected to chromatography repeatedly on Sephadex LH-20 and YMC ODS-A, and further purified by HPLC (Waters 1525/2996, SunFire Silica, 5 µm, 1.5 mL/min, UV detection at 210 and 280 nm) eluting with *n*-hexane/isopropanol (95:5) to give compound **2** (3.7 mg, *t_R* = 21.9 min).

3,3-Bis(uracil-1-yl)-propan-1-aminium (**1**)

White amorphous solid. MP: 107–108°C.

[α]_D²⁵: -2.6 (c 0.5, C₅H₅N).

¹H NMR and ¹³C NMR (DMSO-*d*₆): Table 1.

HR-APCI-MS: *m/z* 280.1213 [M]⁺ (C₁₁H₁₄N₅O₄, calcd. for 280.1202);

Cytotoxicity assay: Cytotoxicity was evaluated as IC₅₀ by using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay, with 5-fluorouracil as positive control. Compounds **1–5** were solubilized in DMSO with the working concentration of test substances ranging from 1 to 100 µg/mL. Cells were inoculated into 96-well plates. After incubation for 24 h, the cells were treated with various concentrations of test substances for 48 h and then incubated with 1 mg/mL MTT at 37°C for 4 h, followed by solubilization in DMSO. The formazan dye products were measured by the absorbance at 490 nm on a microplate reader.

Acknowledgements - This research was supported by the National Natural Science Fund for Distinguished Young Scholars of China (81225023), the National Natural Science Fund of China (Nos. 81302691, 81072573, 81172978, 41106127, and 81001394), the Innovation Program of Shanghai Municipal Education Commission (No. 14YZ037), and partially supported by the Shanghai Subject Chief Scientist (12XD1400200). We are also grateful for the financial support of the National High Technology Research and Development Program of China (863 Projects, Nos. 2011AA09070107 and 2013AA092902).

Reference

- [1] (a) Natori T, Koezuka Y, Higa T. (1993) Agelasphins, novel α-galactosylceramides from the marine sponge *Agelas mauritanus*. *Tetrahedron Letters*, **34**, 5591–5592; (b) Costantino V, Fattorusso E, Imperatore C, Mangoni A. (2004) Glycolipids from sponges. 13.1 Clarhamnoside, the first rhamnosylated α-galactosylceramide from *Agelas clathrodes*. Improving spectral strategies for glycoconjugate structure determination. *The Journal of Organic Chemistry*, **69**, 1174–1179; (c) Appenzeller J, Tilvi S, Martin M T, Gallard J F, El-bitar H, Dau ET, Debitus C, Laurent D, Moriou C, Al-Mourabit A. (2009) Benzoscceprins A and B with a unique benzocyclobutane skeleton and nagelamide S and T from Pacific sponges. *Organic Letters*, **11**, 4874–4877; (d) Yasuda T, Araki A, Kubota T, Ito J, Mikami Y, Fromont J, Kobayashi J. (2009) Bromopyrrole alkaloids from marine sponges of the genus *Agelas*. *Journal of Natural Products*, **72**, 488–491.
- [2] (a) Wu H, Nakamura H, Kobayashi J, Kobayashi M, Ohizumi Y, Hirata Y. (1986) Structures of agelasines, diterpenes having a 9-methyladeninium chromophore isolated from the Okinawan marine sponge *Agelas nakamurai* Hoshino. *Bulletin of the Chemical Society of Japan*, **59**, 2495–2504; (b) Calcul L, Tenney K, Ratnam J, McKerrow J H, Crews P. (2010) Structural variations to the 9-N-methyladeninium diterpenoid hybrid commonly isolated from *Agelas* sponges. *Australian Journal of Chemistry*, **63**, 915–921; (c) Yang F, Hamann MT, Zou Y, Zhang M, Gong X, Xiao J, Chen W, Lin H. (2012) Antimicrobial metabolites from the Paracel Islands sponge *Agelas mauritiana*. *Journal of Natural Products*, **75**, 774–778.
- [3] (a) Morales JJ, Rodriguez AD. (1991) The structure of clathrodin, a novel alkaloid isolated from the Caribbean Sea sponge *Agelas clathrodes*. *Journal of Natural Products*, **54**, 629–631; (b) Cafieri F, Fattorusso E, Mangoni A, Tagliatalata-Scafati O. (1996) Clathramides, unique bromopyrrole alkaloids from the Caribbean sponge *Agelas clathrodes*. *Tetrahedron*, **52**, 13713–13720; (c) Medeiros MA, Lourenco A, Tavares MR, Marcelo Curto MJ, Savluchinske Feio S, Roseiro JC. (2006) (-)-Agelasidine A from *Agelas clathrodes*. *Zeitschrift für Naturforschung. C. A Journal of Biosciences*, **61**, 472–476; (d) Cafieri F, Carnuccio R, Fattorusso E, Tagliatalata-Scafati O, Vallefucio T. (1997) Anti-histaminic activity of bromopyrrole alkaloids isolated from Caribbean *Agelas* sponges. *Bioorganic & Medicinal Chemistry Letters*, **7**, 2283–2288; (e) Cafieri F, Fattorusso E, Mangoni A, Tagliatalata-Scafati O. (1996) Dispacamides, anti-histamine alkaloids from Caribbean *Agelas* sponges. *Tetrahedron Letters*, **37**, 3587–3590; (f) Narender T, Khaliq T, Srivastava M. (2007) Naturally occurring 1, 1'-trimethylenebisuracil from the marine sea hare *Dolabella auricularia*. *Natural Product Communications*, **2**, 71–73; (g) Alam N, Bae BH, Hong J, Lee CO, Shin BA, Im KS, Jung JH. (2001) Additional bioactive Lyso-PAF congeners from the sponge *Spirastrella abata*. *Journal of Natural Products*, **64**, 533–535.
- [4] (a) Ang KH, Prager RH, Williams CM. (1995) The chemistry of 5-oxodihydroisoxazoles. XIII. Reactions of the imino carbene derived from photolysis of ethyl 5-oxo-2-phenyl-2, 5-dihydroisoxazole-4-carboxylate. *Australian Journal of Chemistry*, **48**, 567–575; (b) Koenig GM, Wright AD, Linden A. (1998) Antiplasmodial and cytotoxic metabolites from the Maltese sponge *Agelas oroides*. *Planta Medica-Natural Products and Medicinal Plant Research*, **64**, 443–447; (c) Lindel T, Hoffmann H, Hochgürtel M, Pawlik JR. (2000) Structure–activity relationship of inhibition of fish feeding by sponge-derived and synthetic pyrrole–imidazole alkaloids. *Journal of Chemical Ecology*, **26**, 1477–1496; (d) Morikawa T, Xie H, Matsuda H, Wang T, Yoshikawa M. (2006) Bioactive constituents from Chinese natural medicines. XVII. Constituents with radical scavenging effect and new glucosyloxybenzyl 2-isobutylmalates from *Gymnadenia conopsea*. *Chemical & Pharmaceutical Bulletin*, **54**, 506–513.

Two New Compounds and Anti-complementary Constituents from *Amomum tsao-ko*

Jiahong Jin, Zhihong Cheng and Daofeng Chen*

Department of Pharmacognosy, School of Pharmacy, Fudan University, Shanghai 201203, People's Republic of China

dfchen@shmu.edu.cn

Received: August 29th, 2012; Accepted: October 8th, 2013

Two new compounds, (2*R*,3*R*,4*R*)-3',5'-dimethoxy-3,4,7,4'-tetrahydroxy-flavan (**1**) and 2-(4-hydroxy-3-methoxybenzoyl)-4-methoxy-benzaldehyde (**2**), together with 35 known phenolic compounds were obtained from the fruits of *Amomum tsao-ko*. Structures of the new compounds were elucidated on the basis of spectroscopic means, including 2D NMR, and high-resolution MS analysis. The isolated compounds were tested *in vitro* for their complement-inhibitory properties against the classical pathway (CP) and alternative pathway (AP). The results showed that 14 compounds exhibited anti-complementary activities against the CP and AP with CH₅₀ values of 0.42 - 4.43 mM and AP₅₀ values of 0.53 - 1.51 mM. Preliminary mechanism studies showed that 1,7-bis(4-hydroxyphenyl)-4(*E*)-hepten-3-one (**8**) blocked C1q, C2, C3, C4, C5 and C9 components of the complement system, and hydroquinone (**15**) acted on C1q, C2, C3, C5 and C9 components.

Keywords: *Amomum tsao-ko*, Zingiberaceae, Complement inhibitor, Phenols, Diarylheptanoids, Benzenediols.

When activated inappropriately, the complement system may evoke pathologic reactions in a variety of inflammatory and degenerative diseases such as systemic lupus erythematosus (SLE), rheumatoid arthritis (RA), as well as acute respiratory distress syndrome (ARDS) [1, 2]. Therefore, inhibition of individual complement is a promising approach for the prevention and treatment of these diseases and numerous natural products have been reported to possess anti-complementary effect [3-5].

Amomum tsao-ko Crevost et Lemaire (Zingiberaceae) is an annual plant widely distributed in tropical regions of Asia. Its fruits are used under the name of "Cao-Guo" (Fructus Tsaoko) in traditional Chinese medicine (TCM) for the treatment of stomach disorders and infection of the throat, as well as a food flavor enhancer [6, 7]. In our effort to search for anti-complementary agents from TCMs, the ethanolic extract of *A. tsao-ko* fruits was found to show anti-complementary activity (CH₅₀: 0.75 ± 0.06 mg/mL, AP₅₀: 1.89 ± 0.35 mg/mL). Bioactivity-directed fractionation and isolation was thus performed with the ethanolic extract of *A. tsao-ko* fruits and led to the isolation of two new compounds, (2*R*,3*R*,4*R*)-3',5'-dimethoxy-3,4,7,4'-tetrahydroxy-flavan (**1**) and 2-(4-hydroxy-3-methoxybenzoyl)-4-methoxy-benzaldehyde (**2**) (Figure 1), together with 35 known ones (**3-37**). The isolates were tested for their *in vitro* anti-complementary activities against both the CP and AP, and the targets of the two most active compounds (**8** and **15**) on the complement activation cascade were also investigated.

Compound **1** was obtained as a yellowish amorphous powder. Its molecular formula was determined as C₁₇H₁₈O₇ by HR-ESI-MS (*m/z* 357.0943 [*M* + Na]⁺). Taken together with three oxygenated carbon signals at δ_C 86.6 (C-2), 72.2 (C-3) and 72.1 (C-4) (Table 1), it is easily speculated that **1** possesses a flavan 3,4-diol skeleton, as is the case with melacacidin (2,3-*cis*-3,4-*cis*-flavan-3,3',4,4',7,8-hexaol) [8]. The key difference between **1** and melacacidin was that a symmetrical structure [δ_H 3.82 (3H, s, 3'-OCH₃), 3.81 (3H, s, 5'-OCH₃) and δ_H 6.68 (2H, s, H-2', 6')] was observed in ring B of **1**, while melacacidin possesses 3,4-disubstituted groups in ring B. The HMBC correlations between δ_H 6.68 (1H, s, H-2') in ring B





Figure 1: Structures of **1** and **2**.

and δ_C 86.6 (C-2) and δ_C 72.2 (C-3), and between δ_H 6.78 (1H, d, *J* = 8.3 Hz, H-5) and δ_C 72.2 (C-3) confirmed the flavanol skeleton of **1** (Figure 2A). The locations of methoxy groups at C-3'/5' and hydroxyl at C-4' were also verified by the HMBC correlations between δ_H 3.82 (3H, s, 3'-OCH₃) and δ_C 148.6 (C-3') and δ_C 149.2 (C-4'), and between δ_H 7.52 (1H, s, 4'-OH) and δ_C 149.2 (C-4'), respectively. The evidence for assignment of δ_H 7.14 (1H, s, 7-OH) was that the X-part [δ_H 6.98 (1H, d, *J* = 2.0 Hz, H-8)] of the ABX system showed HMBC correlations with δ_C 110.5 (C-6), 147.5 (C-9) and δ_C 119.5 (C-10). The absolute configuration of **1** was determined as 2*R*,3*R*,4*R* by the negative Cotton effects at 240 nm and 274 nm in its CD spectrum [9], as well as the ROESY correlations of H-2/H-3, and H-3/H-4 (Figure 2A). Therefore, the structure of compound **1** was determined as (2*R*,3*R*,4*R*)-3',5'-dimethoxy-3,4,7,4'-tetrahydroxy-flavan.

Compound **2** was obtained as colorless needles (PE-acetone, 5:1). The molecular formula was determined as C₁₆H₁₄O₅ by HR-ESI-MS (*m/z* 309.0736 [*M* + Na]⁺). The ¹³C NMR spectrum (Table 1) showed 12 carbon signals for two benzene rings (δ_C 153.1, 151.7, 148.2, 147.5, 128.9, 128.7, 126.2, 123.5, 115.4, 114.9, 111.1 and 110.7), one carbonyl at δ_C 196.3, one aldehyde at δ_C 191.2, and two overlapped methoxy groups (δ_C 55.6). The IR spectrum also showed the typical absorption band at 1612 cm⁻¹ for benzene rings. The ¹H NMR spectrum of **2** (Table 1) showed two sets of ABX system proton signals at δ_H 6.85 (1H, d, *J* = 8.3 Hz), 7.42 (1H, d, *J* = 2.0 Hz) and 7.48 (1H, dd, *J* = 8.3, 2.0 Hz), as well as signals at δ_H 6.95 (1H, d, *J* = 8.3 Hz), 7.36 (1H, d, *J* = 2.0 Hz) and 7.40 (1H, dd,

Table 1: ^1H - and ^{13}C -NMR spectral data of compounds **1**^a and **2**^b (δ in ppm, J in Hz)

Position	1		2	
	δ_{H} (multiplicity)	δ_{C}	δ_{H} (multiplicity)	δ_{C}
1				148.2
2	4.66 (d, $J=4.3$)	86.6		128.7
3	4.20 (dd, $J=4.3, 6.1$)	72.2	7.36 (d, $J=2.0$)	110.7
4	3.79 (d, $J=6.1$)	72.1		153.1
5	6.78 (d, $J=8.3$)	134.1	7.40 (dd, $J=8.3, 2.0$)	126.2
6	6.82 (dd, $J=8.3, 2.1$)	110.5	6.95 (d, $J=8.3$)	115.4
7		152.6		
8	6.98 (d, $J=2.1$)	112.8		
9		147.5		
10		119.5		
1'		136.2		128.8
2'	6.68 (br. s)	104.3	7.42 (d, $J=2.0$)	111.1
3'		148.6		147.5
4'		149.2		151.7
5'		148.6	6.85 (d, $J=8.3$)	114.9
6'	6.68 (br. s)	104.3	7.48 (dd, $J=8.3, 2.0$)	123.5
1-CHO			9.75 (s)	191.2
4-OCH ₃			3.82(s)	55.6
7-OH	7.14 (s)			
1'-COAr				196.3
3'-OCH ₃	3.82 (s)	56.5	3.79 (s)	55.6
4'-OH	7.52 (s)		10.05 (br. s)	
5'-OCH ₃	3.81 (s)	56.5		


^a Measured in acetone- d_6 at 400 MHz for ^1H NMR and at 100 MHz for ^{13}C NMR.^b Measured in DMSO- d_6 at 400 MHz for ^1H NMR and at 100 MHz for ^{13}C NMR.**Figure 2:** Key HMBC and ROESY correlations of **1** (A) and **2** (B).

2.0 Hz). The X-part of the ABX system in the A ring was deduced to be located at C-3 based on the HMBC correlations from δ_{H} 7.36 (1H, d, $J=2.0$ Hz, H-3) to δ_{C} 148.2 (C-1), 153.1 (C-4), 126.2 (C-5) and 196.3 (C-1') (Figure 2B). The methoxy group in the A ring was thus assigned at C-4 by the ROESY correlation between δ_{H} 7.36 (1H, d, $J=2.0$ Hz, H-3) and the methyl signal at δ_{H} 3.79 (s, 3'-OCH₃). The aldehyde was assigned at C-1 by the ROESY correlation between the B-part of ABX system at δ_{H} 6.95 (1H, d, $J=8.3$ Hz, H-6) and the aldehyde proton at δ_{H} 9.75 (1H, s, 1-CHO). Similarly, as shown in Figure 2B, the other ABX system and the locations of methoxy and hydroxyl in ring B were assigned. These two benzene rings were connected through one carbonyl group (δ_{C} 196.3, 1'-COAr), as indicated by the HMBC correlations between δ_{H} 7.36 (1H, d, $J=2.0$ Hz, H-3), 7.42 (1H, d, $J=2.0$ Hz, H-2'), 7.48 (1H, dd, $J=8.3, 2.0$ Hz, H-6') and the carbonyl group (δ_{C} 196.3, 1'-COAr) (Figure 2B). Hence, the structure of compound **2** was established as 2-(4-hydroxy-3-methoxybenzoyl)-4-methoxybenzaldehyde.

The 35 known phenolic compounds were identified by comparison of their ^1H , ^{13}C NMR and MS data with those reported in the literatures [10-20] as 4'-hydroxy-4-methoxychalcone (**3**), 4'-hydroxy-2'-methoxychalcone (**4**), 4,4'-dimethoxychalcone (**5**), 1,3-dimethoxybenzene (**6**), 4',7-dihydroxy-3',6-diprenylflavone (**7**), 1,7-bis(4-hydroxyphenyl)-4(*E*)-hepten-3-one (**8**), 4-hydroxy-2,5-dimethoxy-benzaldehyde (**9**), 3',7-dihydroxy-4'-methoxy-flavan (**10**), 2-methoxy-hydroquinone (**11**), 2',4'-dihydroxy-4-methoxy-chalcone (**12**), 5-indancarbaldehyde (**13**), 4-methoxybenzaldehyde (**14**), hydroquinone (**15**), 4-indancarbaldehyde (**16**), (-)-catechin

Table 2: Anti-complementary activities of the compounds from *A. tsao-ko* through the classical pathway (CH₅₀) and alternative pathway (AP₅₀) (mean \pm SD, $n=3$)

Compounds	CH ₅₀ (mM)	AP ₅₀ (mM)
4'-Hydroxy-2'-methoxychalcone (4)	0.96 \pm 0.15	NA ^a
1,7-Bis(4-hydroxyphenyl)-4(<i>E</i>)-hepten-3-one (8)	0.42 \pm 0.15	0.66 \pm 0.11
2-Methoxy-hydroquinone (11)	0.62 \pm 0.09	0.86 \pm 0.13
2',4'-Dihydroxy-4-methoxychalcone (12)	2.69 \pm 0.42	NA
5-Indancarboxaldehyde (13)	1.33 \pm 0.15	1.51 \pm 0.38
Hydroquinone (15)	0.55 \pm 0.11	0.53 \pm 0.15
4-Hydroxy-2'-methoxychalcone (23)	4.43 \pm 1.26	NA
Tsaokoarylone (25)	0.84 \pm 0.07	0.72 \pm 0.16
6,7-Dihydroxyindan-4-carbaldehyde (28)	0.66 \pm 0.14	0.96 \pm 0.13
3-Methoxy-catechol (31)	0.56 \pm 0.13	0.54 \pm 0.10
2-Methoxy-resorcinol (32)	0.64 \pm 0.14	0.58 \pm 0.14
4-(2-Hydroxypropyl)phenol (34)	0.93 \pm 0.14	1.26 \pm 0.18
Catechol (36)	0.58 \pm 0.13	0.57 \pm 0.15
4-Methoxy-catechol (37)	0.69 \pm 0.16	0.64 \pm 0.08
Heparin ^b	40 \pm 14 ^c	97 \pm 19 ^c

^a NA: Not active. ^b Positive control. ^c $\mu\text{g/mL}$.**Figure 3:** Targets of 1,7-bis(4-hydroxyphenyl)-4(*E*)-hepten-3-one (BH, A) and hydroquinone (HQ, B) on the complement activation cascade. BH-, HQ-treated sera were mixed with various complement-depleted (C-depleted) sera and the capacity of these C-depleted sera to restore hemolytic capacity in the CP was estimated by adding sheep antibody-sensitized erythrocytes. Cont., complement control group. Results are expressed as hemolytic percentages. Data are expressed as mean \pm SD ($n=3$).

(**17**), anisole (**18**), 2',4,4'-trimethoxychalcone (**19**), 4-(1-hydroxypropyl)phenol-ethyl-4-hydroxy-(*S*)-benzenemethanol (**20**), abyssinoflavanone VII (**21**), 4-hydroxy-4'-methoxychalcone (**22**), 4-hydroxy-2'-methoxychalcone (**23**), 2-methoxy-benzaldehyde (**24**), tsaokoarylone (**25**), 3-methoxy-benzaldehyde (**26**), 3-hydroxy-4-methoxybenzaldehyde (**27**), 6,7-dihydroxy-4-indancarbaldehyde (**28**), 3-methoxy-4-hydroxy-benzaldehyde (**29**), 6-hydroxy-4-aldehydeindene (**30**), 3-methoxy-catechol (**31**), 2-methoxy-resorcinol (**32**), 3,5-dihydroxybenzoic acid (**33**), 4-(2-hydroxypropyl)phenol (**34**), 3-hydroxybenzoic acid (**35**), catechol (**36**), and 4-methoxy-catechol (**37**). Except for compounds **8**, **14**, **15**, **17**, **18**, **25**, **29** and **36**, all the other compounds were obtained from this species for the first time.

The isolated compounds (purity > 90%, by HPLC analysis), except **33** and **35**, were evaluated for their anti-complementary activities against both CP and AP. As shown in Table 2, fourteen compounds possessed anti-complementary effects against the CP with CH₅₀ values of 0.42 - 4.43 mM. On the AP, eleven isolates were found

active, with AP_{50} values of 0.53 - 1.51 mM. Compounds **8**, **15**, **31**, **32**, **36** and **37** showed good activities against both CP and AP with CH_{50} and AP_{50} values less than 0.7 mM. Derivatives of chalcone (**4**, **12** and **23**) were inactive against AP while they demonstrated inhibitory activity against CP. As the two most active compounds against the CP and AP respectively, 1,7-bis(4-hydroxyphenyl)-4(*E*)-hepten-3-one (**8**) and hydroquinone (**15**) were selected for the preliminary mechanism study. When treated with 1,7-bis(4-hydroxyphenyl)-4(*E*)-hepten-3-one (**8**), hemolytic percentages of the C1q-, C2-, C3-, C4-, C5- and C9-depleted sera were $5.04 \pm 1.03\%$, $8.33 \pm 2.98\%$, $9.46 \pm 3.00\%$, $8.36 \pm 3.78\%$, $8.21 \pm 2.43\%$ and $9.10 \pm 2.19\%$, respectively (Figure 3A), indicating that **8** interacted with C1q, C2, C3, C4, C5 and C9. Meanwhile, hydroquinone (**15**) acted on C1q, C2, C3, C5 and C9 components (Figure 3B).

Experimental

Reagents: Sheep erythrocytes were collected in Alsevers' solution. Normal human serum (NHS) was obtained from healthy male donors (average age of 20 years). Rabbit erythrocytes were obtained from the ear vein of New Zealand white rabbits. Heparin (sodium salt, 160 IU/mg) was purchased from Shanghai Aizite Biotech Co. Ltd. (Shanghai, China). Veronal buffer saline (VBS, pH 7.4) contained 0.5 mM Mg^{2+} and 0.15 mM Ca^{2+} (VBS $^{2+}$), and VBS 5 mM Mg^{2+} and 8 mM EGTA (VBS-Mg-EGTA). Anti-C1q, Human (Goat); Anti-C2, Human (Goat); Anti-C5, Human (Rabbit) and Anti-C9, Human (Goat) were purchased from Merck Biosciences (Darmstadt, German). Anti-C3, Human (Goat) and Anti-C4, Human (Goat) were purchased from Shanghai Sun Biotech Co. Ltd. (Shanghai, China).

Plant material: The fruits of *A. tsao-ko* were purchased from Huayu Materia Medica Co., Ltd. (Shanghai, China) in October 2007. The plant material was verified by Dr Daofeng Chen. A voucher specimen (DFC-CG-H2003050607) has been deposited in the Herbarium of Materia Medica, Department of Pharmacognosy, School of Pharmacy, Fudan University, Shanghai, China.

Extraction and isolation: The air-dried fruits (15.0 kg) were extracted with 95% ethanol (10 L \times 5 times) at room temperature. The extracts were combined and concentrated under reduced pressure, and the resulting residue (319.2 g) was resuspended in 2.0 L of water and successively partitioned with PE (60-90°C), ethyl acetate, and *n*-butanol (each 2 L \times 4 times). The *n*-butanol fraction was found to show the highest activity. Therefore, the active *n*-butanol fraction was subjected to AB-8 macroporous resin chromatography eluted with 30%, 50%, 70% and 90% aqueous ethanol to afford 4 subfractions. The subfractions obtained with 50% and 70% ethanol were found to show obvious anti-complementary activities and were thus further chromatographed over a silica gel column (CC).

The 50% EtOH subfraction (21.3 g) was subjected to CC (silica gel, PE-acetone, from 100:0 to 1:1) to yield F-1 - F-8. F-2 (1.7 g) was separated by CC (silica gel, PE-ethyl acetate, from 20:1 to 10:1) and then purified over Sephadex LH-20 (chloroform-methanol, 1:1) to afford **13** (21.2 mg). F-3 (1.2 g) was purified over CC (silica gel, PE-ethyl acetate, 10:1) to give **16** (12.7 mg), **22** (8.0 mg), **26** (4.3 mg), **34** (10.2 mg), **9** (14.3 mg) and **10** (5.8 mg). F-5 (2.5 g) was loaded onto a column of silica gel and eluted with PE-ethyl acetate, from 10:1 to 5:1, and then subjected to Sephadex LH-20 (chloroform-methanol, 1:1) and preparative TLC (chloroform-methanol, 20:1) to afford **2** (15.4 mg), **3** (4.4 mg), **4** (3.9 mg), **14** (4.1 mg), **21** (12.1 mg) and **23** (16.6 mg). F-6 (1.1 g) was purified by CC (silica gel, PE-acetone, from 5:0 to 5:1) and then subjected to

gel filtration over Sephadex LH-20 (chloroform-methanol, 1:1) and preparative TLC (chloroform-methanol, 15:1) to afford **31** (8.7 mg), **32** (12.1 mg), **36** (9.5 mg), **27** (3.3 mg), **30** (4.4 mg) and **35** (3.9 mg). F-7 (1.8 g) was purified by CC (silica gel, PE-acetone, from 10:1 to 5:1) and then subjected to preparative TLC (chloroform-methanol, 10:1) to furnish **11** (15.3 mg), **15** (6.8 mg), **33** (9.3 mg) and **37** (13.8 mg).

The 70% EtOH subfraction (13.7 g) was subjected to CC (silica gel, PE-acetone, from 100:0 to 1:1) to yield F-9 - F-16. F-11 was further separated by CC (silica gel, chloroform-methanol, 10:1) and then purified over Sephadex LH-20 (chloroform-methanol, 1:1) and preparative TLC (chloroform-methanol, 15:1) to give **20** (9.9 mg), **24** (9.3 mg) and **25** (17.9 mg). F-12 (1.5 g) was chromatographed over Sephadex LH-20 (methanol) and then separated by CC (silica gel, chloroform-methanol, 15:1) to yield **1** (5.3 mg), **7** (5.1 mg), **8** (7.9 mg) and **18** (6.3 mg). F-13 (1.6 g) was chromatographed over a column of silica gel, eluting with chloroform-methanol, 15:1, and then purified over Sephadex LH-20 (methanol) to afford **12** (7.2 mg), **17** (7.7 mg), **28** (5.3 mg) and **29** (10.9 mg). F-14 (2.7 g) was purified by CC (silica gel, chloroform-methanol, 7:1) and then purified over Sephadex LH-20 (chloroform-methanol, 1:1) to afford **5** (8.3 mg), **6** (5.9 mg) and **19** (6.3 mg).

(2*R*,3*R*,4*R*)-3',5'-Dimethoxy-3,4,7,4'-tetrahydroxy-flavan (**1**)

Pale yellow powder.

MP: 182-184°C.

$[\alpha]_D^{25}$: -26.8 (*c* 0.02, acetone).

IR (KBr) ν_{max} : 3420, 2955, 2920, 2850, 1634, 1558, 1446, 1384, 1303 cm^{-1} .

UV (acetone): λ_{max} (log ϵ): 223 (5.2), 255 (3.1, sh) nm.

CD (*c* 0.1, methanol): nm($\Delta\epsilon$): 240 (-32.1), 274 (-11.4).

1H NMR and ^{13}C NMR (acetone- d_6): Table 1;

HR-ESI-MS: m/z 357.0943 ($[M + Na]^+$, $C_{17}H_{18}O_7Na^+$, calc. 357.0945).

2-(4-Hydroxy-3-methoxybenzoyl)-4-methoxy-benzaldehyde (**2**)

Colorless needles (PE-acetone, 5:1).

MP: 166-167°C.

IR (KBr) ν_{max} : 3416, 1721, 1612, 1462, 1383, 1272 cm^{-1} .

UV (acetone): λ_{max} (log ϵ): 223 (4.6), 250 (2.7, sh) nm.

1H NMR and ^{13}C NMR (DMSO- d_6): Table 1.

HR-ESI-MS: m/z 309.0736 ($[M + Na]^+$, $C_{16}H_{14}O_5Na^+$, calc. 309.0734).

Anti-complementary activity through the classical pathway: The anti-complementary activities through the classical pathway (CP) were measured as described previously [21]. Briefly, sensitized erythrocytes (EAs) were prepared by incubation of sheep erythrocytes (4.0×10^8 cells/mL) with equal volumes of rabbit anti-sheep erythrocyte antibody in VBS $^{2+}$. Samples and heparin (positive control) were individually dissolved in VBS $^{2+}$. Normal human serum (NHS) was used as the complement source. The 1:80 diluted serum of Guinea pig was chosen to give sub-maximal lysis in the absence of complement inhibitors. Serial dilutions of the test samples (100 μ L) were preincubated with a mixture of 100 μ L NHS and 200 μ L VBS $^{2+}$ at 37°C for 10 min, followed by adding 200 μ L EA to the mixture and co-incubating at 37°C for 30 min. The different assay controls were incubated under the same conditions: (1) vehicle control, 200 μ L EAs in 400 μ L VBS $^{2+}$; (2) control 100% lysis, 200 μ L EAs in 400 μ L water; (3) sample control, 100 μ L dilution of each sample in 500 μ L VBS $^{2+}$. After reaction, the resulting mixture was centrifuged immediately, and the optical density of the supernatant was then measured at 405 nm with a spectrophotometer (Wellscan MK3, Labsystems Dragon). The

absorbance of sample (As), sample control (Asc) and 100% lysis control (Al) were obtained.

Anti-complementary activity through the alternative pathway: The anti-complementary activities through the classical pathway (CP) and alternative pathway (AP) were measured according to Klerx's method [22]. Briefly, each sample was dissolved in EGTA-VB, and serial dilutions of the samples were prepared by adding appropriate volumes of EGTA-VB. After pre-incubation of each sample (150 μ L) with 1:10 diluted NHS (150 μ L) at 37°C for 10 min, 200 μ L rabbit erythrocytes (ERs 3.0×10^8 cells/mL) were added, followed by a second incubation at 37°C for 30 min. After reaction, the resulting mixture was centrifuged immediately, and the optical density of the supernatant was then measured at 405 nm.

Identification of the targets on the complement activation cascade: Tests to identify the targets of the complement activation cascade were conducted according to Xu's method [5]. Various dilutions of each antiserum were incubated with the same volume of NHS (1:10, v/v) at 37°C for 15 min. After centrifugation, the supernatant (200 μ L) was incubated with 200 μ L VBS²⁺ and 200 μ L EAs, and cell lysis was measured. The antiserum dilution against the NHS hemolytic capacity was then determined. The optimal dilutions (1:1 for C3 and C4; 1:32 for C5, and 1:64 for C1q, C2, and C9, v/v) were incubated with the same volume of NHS (1:10, v/v) at 37°C for 15 min, followed by centrifugation, and the supernatants were collected and stored as complement-depleted (C-depleted) sera in aliquots at -70°C before use in hemolytic assays.

Capacity of depleted sera to lyse EAs through the CP was assessed in the presence or absence of sample-treated NHS. Sample-treated NHS was obtained by incubating an optimally diluted sample with an equal volume of 1:10 (v/v) diluted NHS at 37°C for 10 min. The examined concentrations of 1,7-bis(4-hydroxyphenyl)-4(*E*)-hepten-3-one (**8**) and hydroquinone (**15**), just sufficient to cause complete loss of hemolytic activity of 1:10 diluted NHS, were 1.12 mM and 1.24 mM. For the target complement group (the assay of capacity of various depleted sera to restore the hemolytic capacity of sample-treated serum), 200 μ L EAs and 200 μ L individual depleted sera of C1q, C2, C3, C4, C5 or C9 were added to 200 μ L sample-treated NHS and the mixture was incubated at 37°C for 30 min. After centrifugation and measurement of the optical density of the supernatant, the percentage of hemolysis was calculated. For the assay of the individual depleted serum group, C-depleted sera were directly incubated with EAs under the same conditions, and the hemolytic activities were calculated. The controls: (1) vehicle control: 200 μ L EAs in 400 μ L VBS²⁺; (2) 100% lysis: 200 μ L EAs in 400 μ L water; (3) complement control: 100 μ L NHS (1:10, v/v) and 200 μ L EAs in 300 μ L VBS²⁺; and (4) sample control, 100 μ L sample in 500 μ L VBS²⁺, were incubated under the same conditions.

Acknowledgments - This work was supported by grants from the National Natural Science Foundation for Excellent Youth (30925042), the State Key Program for Innovative Drugs from the Ministry of Science and Technology (2009ZX09502-013 and 2009ZX09301-011).

References

- [1] Makridea SC. (1998) Therapeutic inhibition of the complement system. *Pharmacological Reviews*, **50**, 59-87.
- [2] Lustep HL, Clark WM. (2001) Current status of neuroprotective agents in the treatment of acute ischemic stroke. *Current Neurology and Neuroscience Reports*, **1**, 13-18.
- [3] Peiris JSM, Chu CM, Cheng VCC, Chan KS, Hung IFN, Poon LLM. (2003) Clinical progression and viral load in a community outbreak of coronavirus-associated SARS pneumonia: a prospective study. *The Lancet*, **361**, 1767-1772.
- [4] Kirschfink M. (1997) Controlling the complement system in inflammation. *Immunopharmacology*, **8**, 51-62.
- [5] Xu H, Zhang YY, Zhang JW, Chen DF. (2007) Isolation and characterization of an anti-complementary polysaccharide D3-S1 from the roots of *Bupleurum smithii*. *International Immunopharmacology*, **2**, 175-180.
- [6] Teresita SM, Hiroe K, Masashi H, Nakatani N. (2000) *Amomum tsao-ko* and their radical scavenging and antioxidant activities. *Journal of the American Oil Chemists' Society*, **77**, 667-673.
- [7] Peng JM, Ma J, Zhang LX. (2006) Review of progress in *Amomum tsao-ko*. *Chinese Traditional Patent Medicine*, **7**, 1036-1038.
- [8] Lai YF. (1985) Facile self-condensation of melacacidin: a demonstration of the reactivity of the pyrogallol A-ring. *Journal of the Chemical Society, Chemical Communications*, 1273-1274.
- [9] Daneel F, Jannie PJM, Desmond S, Larry AW. (2004) Circular dichroic properties of flavan-3,4-diols. *Journal of Natural Products*, **67**, 174-178.
- [10] Satyanarayana M, Tiwari P, Tripathi BK, Srivastava AK, Pratap R. (2004) Synthesis and antihyperglycemic activity of chalcone based aryloxypropanolamines. *Bioorganic Medicinal Chemistry*, **12**, 883-889.
- [11] Bai LP, Jiang H, Kang TG, Zhang HM. (2005) Chemical constituents from fruits of *Amorpha fruticosa*. *Nature Medicine*, **58**, 275-277.
- [12] Pouchert C. (1992) Aldrich Library of ¹³C and ¹H FT NMR Spectra, Vol. 2. Elsevier Publishers, Amsterdam, Netherlands. 183.
- [13] Kiichiro K, Sachio D, Yukio H. (1992) New prenylflavones and dibenzoylmethane from *Glycyrrhiza inflata*. *Journal of Natural Products*, **55**, 1197-1203.
- [14] Nomura M, Tokoroyama T, Kubota T. (1981) Biarylheptanoids and other constituents from wood of *Alnus japonica*. *Phytochemistry*, **20**, 1097-1104.
- [15] Zhang M, Zhang CF, Wang ZT. (2005) Chemical constituents of *Ligularia pleurocaulis*. *Acta Pharmacologica Sinica*, **40**, 529-532.
- [16] Ponce MA, Scervino JM, Erra-Balsells R, Ocampo JA, Godeas AM. (2004) Flavonoids from shoots and roots of *Trifolium repens* (white clover) grown in presence or absence of the arbuscular mycorrhizal fungus *Glomus intraradices*. *Phytochemistry*, **65**, 1925-1930.
- [17] Yang H, Wang D, Tong L, Cai BC. (2007) Chemical constituents of Tibetan medicine *Oxytropis falcata* (L). *Journal of Chinese Pharmaceutical Sciences*, **38**, 1458-1460.
- [18] Marcinkiewicz S, Green J. (1963) Paper chromatography and chemical structure. IV. Intramolecular hydrogen bonding. *Journal of Chromatography A*, **10**, 184-189.
- [19] Lee HS. (2004) Acaricidal activity of constituents identified in *Foeniculum vulgare* fruit oil against *Dermatophagoides* spp. (Acari: Pyroglyphidae). *Journal of Agricultural and Food Chemistry*, **52**, 2887-2889.
- [20] Wang SJ, Pei YH. (2000) Studies on chemical constituents of leaves of *Betula platyphylla* Suk. *Journal of Shenyang Pharmaceutical University*, **17**, 256-257.
- [21] Mayer MM. (1961) *Complement and complement fixation*. 2ed. *Experimental Immunochemistry*. Kabat EA, Mayer MM. (Eds). Springfield, Illinois, USA, 133-240.
- [22] Klerx JP, Beukelman CJ, Van DH, Willers JM. (1983) Microassay for colorimetric estimation of complement activity in guinea pig, human and mouse serum. *Journal of Immunological Methods*, **63**, 215-220.

Antiangiogenic Activity of Flavonoids from *Melia azedarach*Shigenori Kumazawa^{a,*}, Satomi Kubota^a, Haruna Yamamoto^a, Naoki Okamura^a, Yasumasa Sugiyama^b, Hirokazu Kobayashi^a, Motoyasu Nakanishi^c and Toshiro Ohta^a^aDepartment of Food and Nutritional Sciences, University of Shizuoka, 52-1 Yada, Suruga-ku, Shizuoka 422-8526, Japan^bBioorganic Chemistry Research Group, Faculty of Fisheries, Kagoshima University, 4-50-20 Shimoarata, Kagoshima 890-0056, Japan^cInnovation and Joint Research Center, Shizuoka University, 3-5-1 Johoku, Naka-ku, Hamamatsu, Shizuoka 432-8561, Japan

kumazawa@u-shizuoka-ken.ac.jp

Received: August 30th, 2013; Accepted: September 19th, 2013

Three flavonoid glycosides, **1** (rutin: quercetin 3-*O*-rutinoside), **2** (kaempferol 3-*O*-robinobioside) and **3** (kaempferol 3-*O*-rutinoside) were isolated from the subcritical water extracts of *Melia azedarach* leaves. Strong antiangiogenic activity of these compounds was observed in the *in vivo* assay using the chorioallantoic membrane (CAM) from growing chick embryos.

Keywords: Angiogenesis, Flavonoid, Subcritical extraction, *Melia azedarach*, Quercetin, Kaempferol.

Angiogenesis is the process of new endothelial blood vessel development. It is essential for the growth, invasion and metastasis of most solid tumors and has become a valuable pharmacological target for cancer prevention and treatment [1a,b]. We performed *in vivo* experimental angiogenesis screening of plant extracts using the chorioallantoic membrane (CAM) from growing chick embryos [2]. We found that the leaf extracts of *Melia azedarach* L. (Meliaceae) have strong antiangiogenic activities. In particular, the activity of the leaf extracts obtained by using subcritical water was stronger than that of the extracts by using water or alcohol under atmospheric pressure (data not shown).

At present, subcritical water extraction is used for a variety of applications across various fields of study. Studies have shown its use in extracting phenolic compounds from grape pomace and potato [3a,b]. Recently, subcritical water has gained interest as an alternative solvent for the extraction of natural active compounds. We utilized this technique to identify effective antiangiogenesis compounds from medicinal plants.

M. azedarach is a large evergreen tree that is distributed from southeastern Asia to northern Australia. Previous phytochemical and pharmacological studies on the roots of the plant have revealed that it can yield limonoids with antifeedant effects [4a,b]. Furthermore, steroids, with cytotoxic effects, and flavonoids, with antioxidant activities, have been isolated from the leaves [5a,b]. However, to our knowledge, compounds exhibiting antiangiogenesis activities have not been isolated from *M. azedarach*.

Bioactivity-guided isolation was performed through the antiangiogenic activity of the subcritical water extracts of *M. azedarach* leaves by using the CAM assay. The isolation procedures are described in the Experimental section. We isolated and identified three flavonoid glycosides by NMR and MS analysis, **1** (rutin: quercetin 3-*O*-rutinoside), **2** (kaempferol 3-*O*-robinobioside) and **3** (kaempferol 3-*O*-rutinoside), as the active compounds from the leaf extracts of *M. azedarach*.




Figure 1: Effects of **1**–**3** on angiogenesis of the chorioallantoic membrane (CAM) *in vivo*. Formation and inhibition of the CAM vascular network when treated with the MeOH control (**a**), or 50 µg of **1** (**b**), **2** (**c**) and **3** (**d**) are shown.

Compound **1** is the main flavonoid in various plants and has been reported to possess a variety of pharmacological activities [6]. It has been previously identified from leaf extracts of *M. azedarach* [7]. Compound **2** has been isolated from leaves of *Alternanthera brasiliana* and shown to exhibit inhibitory effects on lymphocyte proliferation *in vitro* [8]. It has also been identified from *Gynura formosana* Kiamra [9]. However, to date, no studies have reported the isolation of **2** from *M. azedarach*. Compound **3** has been isolated from the leaves of *M. azedarach* and reported to possess antioxidant activity [7].

Figure 1 shows the results of the antiangiogenic activities of **1**–**3** (50 µg/egg) using the CAM assay. In eggs treated with **1**–**3** (Figure 1b–d), the avascular zone was observed inside and around the silicon ring on the CAM surface. This result indicates that **1**–**3** inhibited angiogenesis, and the subsequent formation of the vascular network. Although **1**–**3** are known flavonoid glycosides, their antiangiogenic activity is reported here for the first time. We

also tested the antiangiogenic activity of **1**~**3** *in vitro* by using human umbilical vein endothelial cells (HUVECs), because we had observed the *in vitro* activity of quercetin and kaempferol, which are the aglycones of **1**~**3** [10]. However, **1**~**3** did not show any sign of the anticipated *in vitro* antiangiogenic activity (data not shown). It is reported that dietary flavonoid glycosides are subjected to deglycosidation by enterobacteria for absorption in the intestine [11]. However, the metabolism of flavonoid glycosides on the CAM surface was unclear, and hydrolysis of **1**~**3** could not be confirmed in the present study.

This study suggests that the glycosides from the leaves of *M. azedarach* exhibit antiangiogenic activity *in vivo*. These findings further extend the potential pharmacological effects of *M. azedarach* and could demonstrate its usefulness in cancer prevention and treatment.

Experimental

General: NMR experiments were performed on a Bruker BioSpin AVANCE III 400 MHz spectrometer (Billerica, MA, USA). Chemical shifts (δ in ppm) were referenced to the carbon (δ_C 49.0) and residual proton (δ_H 3.30) signals of CD₃OD. Low-resolution electrospray ionization (ESI) MS were recorded on an LCG mass spectrometer (Thermo Fisher Scientific, Waltham, MA, USA).

Plant material: The leaves of *M. azedarach* were collected from Mima, Tokushima Prefecture, Japan, in July 2009. The plant was identified by Hideto Miyamoto (Medicinal Botanical Garden, University of Shizuoka, Japan). A voucher specimen was deposited at the University of Shizuoka, Japan.

References

- [1] (a) Folkman J. (1995) Angiogenesis in cancer, vascular, rheumatic and other disease. *Nature Medicine*, **1**, 27-31; (b) Carmeliet P. (2003) Angiogenesis in health and disease. *Nature Medicine*, **9**, 653-660.
- [2] Tsuchiya I, Hosoya T, Usida M, Kunimasa K, Ohta T, Kumazawa S. (2013) Nymphaeol-A isolated from Okinawan propolis suppresses angiogenesis and induces caspase-dependent apoptosis via inactivation of survival signals. *Evidence-Based Complementary and Alternative Medicine*, 2013: 826245. doi: 10.1155/2013/826245. Epub 2013 Apr 24.
- [3] (a) Monrad JK, Howard LR, King JW, Srinivas K, Mauromoustakos A. (2010) Subcritical solvent extraction of anthocyanins from dried red grape pomace. *Journal of Agricultural and Food Chemistry*, **58**, 2862-2868; (b) Singh PP, Saldaña MDA. (2011) Subcritical water extraction of phenolic compounds from potato peel. *Food Research International*, **44**, 2452-2458.
- [4] (a) Nakatani M, Huang RC, Okamura H, Iwagawa T, Tadera K, Naoki H. (1995) Three new antifeeding meliacarpinins from Chinese *Melia azedarach* Linn. *Tetrahedron*, **51**, 11731-11736; (b) Carpinella MC, Defago MT, Valladares G, Palacios SM. (2003) Antifeedant and insecticide properties of a limonoid from *Melia azedarach* (Meliaceae) with potential use for pest management. *Journal of Agricultural and Food Chemistry*, **51**, 369-374.
- [5] (a) Wu SB, Ji YP, Zhu JJ, Zho Y, Xia G, Hu YH, Hu JF. (2009) Steroids from the leaves of Chinese *Melia azedarach* and their cytotoxic effects on human cancer cell lines. *Steroids*, **74**, 761-765; (b) Salib JY, Michael HN, El-Nogoumy SI. (2008) New lactoyl glycoside quercetin from *Melia azedarach* leaves. *Chemistry of Natural Compounds*, **44**, 13-15.
- [6] Okamoto T. (2005) Safety of quercetin for clinical application (Review). *International Journal of Molecular Medicine*, **16**, 275-278.
- [7] Brochado CO, Almeida AP, Barreto BP, Costa LP, Ribeiro LS, Pereira RLC, Gonçalves Koatz VL, Costa SS. (2003) Flavonol robinobiosides and rutinoides from *Alternanthera brasiliana* (Amaranthaceae) and their effects on lymphocyte proliferation *in vitro*. *Journal of Brazilian Chemical Society*, **14**, 449-451.
- [8] Hou WC, Lin RD, Lee TH, Huang YH, Hsu FL, Lee MH. (2005) The phenolic constituents and free radical scavenging activities of *Gynura formosana* Kiamra. *Journal of the Science of Food and Agriculture*, **85**, 615-621.
- [9] Ahn MR, Kunimasa K, Kumazawa S, Nakayama T, Kaji K, Uto Y, Hori H, Nagasawa H, Ohta T. (2009) Correlation between antiangiogenic activity and antioxidant activity of various components from propolis. *Molecular Nutrition and Food Research*, **53**, 643-651.
- [10] Murota K, Terao J. (2003) Antioxidative flavonoid quercetin: implication of its intestinal absorption and metabolism. *Archives of Biochemistry and Biophysics*, **417**, 12-17.

Extraction and isolation: Dried leaves of *M. azedarach* (2.4 kg) were extracted with subcritical water (100°C, 6 MPa, 5 min) to give a crude extract (2.2 g). This was mixed with 1 g of silica gel (Merck, Silica gel 60, Germany) and concentrated to yield dried silica gel, which was placed on the top of a silica gel column (320 mm × 30 mm i.d.) and eluted with *n*-hexane, EtOAc and MeOH to give 19 fractions (frs.): frs. 1~6, *n*-hexane/EtOAc (1:1) eluate; frs. 7~12, EtOAc/MeOH (1:1) eluate; and frs. 13~19, MeOH eluate. A total of 200 mL of each fraction was collected and evaluated by the CAM assay. Fr. 7, which showed the highest activity, was rechromatographed by preparative HPLC on a 250 mm × 20 mm i.d. ODS column (Shiseido, Tokyo, Japan) with 0.1% TFA in CH₃CN-H₂O (18:82) at a 10 mL/min flow rate, thus yielding the active compounds **1** (2.4 mg), **2** (2.9 mg) and **3** (4.1 mg).

CAM assay: The CAM assay was performed as previously described [2]. In brief, fertilized chicken eggs were incubated at 37°C. On incubation day 3, a small window was opened in the shell, and 4 mL of the albumen was removed and further incubated. After 5-days, CAM was treated with various doses of the samples and incubated at 37°C for another 2 days. An appropriate volume of white emulsion was injected into CAM to clearly visualize the vascular network. Observations of the vascular networks were carried out to evaluate the antiangiogenic activity. Retinoic acid (5 nmol/egg) was used as the positive control.

Acknowledgments – We thank K. Kudo of Tohzai Chemical Industry Co., Ltd. for preparing the subcritical water extracts of the leaves of *M. azedarach*.

Application of Mixture Analysis to Crude Materials from Natural Resources (IV)^[1(a-c)]: Identification of *Glycyrrhiza* Species by Direct Analysis in Real Time Mass Spectrometry (II)

Eriko Fukuda^a, Yoshihiro Uesawa^b, Masaki Baba^a and Yoshihito Okada^{a*}

^aDepartment of Natural Medicine and Phytochemistry, Meiji Pharmaceutical University, 2-522-1 Noshio, Kiyose, Tokyo 204-8588, Japan

^bDepartment of Clinical Pharmaceutics, Meiji Pharmaceutical University, 2-522-1 Noshio, Kiyose, Tokyo 204-8588, Japan

y-okada@my-pharm.ac.jp

Received: May 14th, 2013; Accepted: September 3rd, 2013

In order to identify *Glycyrrhiza* species by chemical fingerprinting, the bark of the roots and stolons of *Glycyrrhiza uralensis* Fischer and *G. glabra* Linné were analyzed using DART (Direct Analysis in Real Time)-MS. The characteristic peaks of each species were determined statistically by volcano plot. This summarizes the relationship between the *p*-values of a statistical test and the magnitude of the difference in values of the samples in the groups. In this experiment, peaks that had a *p* value <0.05 in the *t* test and ≥ 2 absolute difference were defined as characteristic. As a result, characteristic peaks of *G. uralensis* were found at *m/z* 299, 315, 341, and 369. In contrast, characteristic peaks of *G. glabra* were found at *m/z* 323, 325, 337, 339, and 391. In conclusion, we found several characteristic peaks to distinguish *G. uralensis* and *G. glabra* by DART-MS using volcano plot. This method can be applied to identify the *Glycyrrhiza* species.

Keywords: Direct Analysis in Real Time-MS (DART-MS), *Glycyrrhiza uralensis*, *Glycyrrhiza glabra*, Volcano plot, Mixture analysis.

The roots and stolons of some *Glycyrrhiza* species have been used as pharmaceuticals. In the current edition of the Japanese Pharmacopoeia (JP XVI), *G. uralensis* Fischer and *G. glabra* Linné are defined as original plant sources of “Glycyrrhizae Radix”. Meanwhile, in the Chinese Pharmacopoeia, *G. inflata* Batalin is also defined [2]. These three species contain glycyrrhizic acid as a main active pharmaceutical ingredient, but other components are also medicinal. Phenolic constituents such as flavonoids and coumarins vary according to the species. Glycycoumarin, glabridin, and licochalcone A were reconfirmed as species-specific typical constituents of *G. uralensis*, *G. glabra*, and *G. inflata*, respectively [3]. Glycycoumarin is an antispasmodic ingredient [4]. Chandrasekaran *et al.* reported that *G. glabra* (almost devoid of glycyrrhizin) exhibits anti-inflammatory properties, which could be influenced in part by glabridin and isoliquiritigenin [5]. It is possible that the differences in ingredients, especially phenolic components, influence medicinal effects. Therefore, species identification is required. Identification presupposes the possibility that the appropriate species can be proposed as the situation demands in Kampo medicine.

In a previous study, we reported the identification of *G. inflata* by DART-MS. DART (Direct Analysis in Real Time) is a novel mass spectrometric ion source [6]. This method provides [M+H]⁺ cation adducts in positive mode, resulting in fewer cleavage reactions. This point is helpful to identify the target substance in mixture states. Therefore, it is often used for qualitative analysis [7-9]. The powder of the roots and stolons of *G. inflata* can be differentiated from those of *G. uralensis* and *G. glabra* by detection of the peak that originated mainly from the [M+H]⁺ ion of licochalcone A [1a]. On the other hand, glycycoumarin and glabridin were not identified by DART-MS analysis of the powder of the roots and stolons of *G. uralensis* and *G. glabra* under the same conditions. [1a] Compared with licochalcone A in *G. inflata*, the amounts of glycycoumarin

and glabridin in the other two species are very small and below detectable limits. Thus, it is not enough to show identification of *G. uralensis* and *G. glabra* by detection of one particular peak. In order to discriminate between the two species, several characteristic peaks for each species were selected by multivariate analysis. Multivariate analysis using the DART-MS data has also been developed by means of chemometric classification [10] and identification of unknown samples [11].

According to Kuwajima *et al.*, nonglycoside flavonoid constituents were mostly lost when root bark is removed [12]. The root bark of *Glycyrrhiza* species is thought to be rich in nonglycosidic flavonoids and have signature substances that are easy to detect. Therefore, the root bark of *G. uralensis* and *G. glabra* were analyzed by DART-MS. The current research examined seven samples of *Glycyrrhiza*, four of *G. uralensis*, and three of *G. glabra*, as shown in Table 1. Species identification had been determined by HPLC, based on modified conditions of Shibano *et al.* [13], before DART-MS analysis.

Table1: sample list.

Sample No.	Place of production	Species identification
Gly-3	Japan	<i>G. glabra</i>
Gly-5	Japan	<i>G. uralensis</i>
Gly-17	Kazakhstan	<i>G. glabra</i>
Gly-18	Uzbekistan	<i>G. glabra</i>
Gly-19	China	<i>G. uralensis</i>
Gly-20	Russia	<i>G. uralensis</i>
Gly-21	Mongolia	<i>G. uralensis</i>

Root bark was taken by a molding plane from three different points per sample. The root bark of each test sample was placed between the ion source and mass spectrometer detector for measurement, and subjected to streaming helium gas for 48 sec.




Figure 1A: DART-MS of the bark of *G. uralensis*.Figure 1B: DART-MS of the bark of *G. glabra*.

Figure 2: The volcano plot for the 43 peaks from the DART-MS data of the bark of licorice roots and stolons.

The x-axis is the difference of the average of both groups, and the y-axis is the p -value in $-\log_{10}p$ scale computed. Each shape represents one of 43 peaks. Using two absolute values of difference and a p -value of 0.05 as the cutoff threshold, four peaks from the upper left region (characteristic for *G. uralensis*; represented as \blacktriangle) and five peaks from the upper right region (characteristic for *G. glabra*; represented as \bullet) were selected (See Table 2).

Typical spectral data of the bark of *G. uralensis* and *G. glabra* are shown in Figure 1. In the range of m/z 280–450, where it is thought that secondary metabolites are greatly detected, the ions selected for the statistical analysis had abundances of at least 20% relative to the most abundant ion in that mass spectral range (20% threshold). The relative intensity of each peak was then calculated. In order to compare each sample, the data were divided by the standard deviation (SD) in the sample of the same individual, which standardizes the data ($SD=1$). Then, 128 out of 171 peaks which had any peak-appearance in each sample were removed. Therefore, 43 peaks were used for the analysis.

The volcano plot summarizes the relationship between the p -values of a statistical test and the magnitude of the difference values of the samples in the groups. It is a scatter plot of the fold-change or the difference value versus the p -value (in $-\log_{10}p$ scale). P -value was calculated by the Student's t -test. The $-\log_{10}p$ -values are plotted on the y-axis. As the intensity difference of the peaks is within normal distribution, we selected the difference values of the two for the x-axis. Value of difference is calculated as [(average of *G. glabra*) - (average of *G. uralensis*)].

Figure 2 is a volcano plot of the DART-MS data of the bark of licorice. The values of difference and p values between these two conditions were plotted for each peak. Peaks landing on the upper left region and the upper right region have a small p -value with a large absolute difference. The peaks that had a p value <0.05 in the t test and ≥ 2 absolute differences were selected as characteristic.

The volcano plot showed that peaks for m/z 299, 315, 341, and 369 are signature peaks of *G. uralensis*, and m/z 323, 325, 337, 339, and 391 of *G. glabra*. The peaks for m/z 369 and 325 were identified by LC/ESI/MS as $[M+H]^+$ ions of glycycomarin and glabridin, respectively.

In the case of *G. uralensis*, four characteristic peaks are found. In 9 out of 12 *G. uralensis* samples (75%), three or more of these characteristic peaks had relative intensities $>20\%$ of original DART-MS data. However, peaks that were characteristic of *G. glabra* were also detected in some *G. uralensis* samples, for example Gly19-2. (Table 3)

In the case of *G. glabra*, five characteristic peaks are found. In 7 out of 9 samples (77%), three or more characteristic peaks had relative intensities greater than 20% of original data. However, peaks that were characteristic of *G. uralensis* were also detected in some *G. glabra* samples, for example Gly17-2 (Table 3).

Although several unidentified samples remained, the detection of characteristic peaks is useful for species identification.

Gly3-1 and Gly3-3 did not detect typical peaks enough. Since they have high intensity peaks, most of the peaks under the threshold value would have been eliminated from the statistical analysis.

It is important to note that the statistical analysis performed in this study was only performed on peaks that had abundances of at least 20% relative to the most abundant ion in a given mass spectral range.

Table 2: The *p*-value and the absolute difference value for each of the nine peaks extracted.

<i>m/z</i>	299*	315*	341*	369*	323**	325**	337**	339**	391**
<i>p</i> value	2.07E-05	0.004087	0.033929	0.000354	0.02424	0.000605	0.000839	0.00479	0.005785
$-\log_{10} p$	4.683174	2.388613	1.469428	3.450702	1.615475	3.217991	3.076434	2.319649	2.23766
value of difference	-5.51027	-3.2431	-3.60396	-2.65802	2.391156	2.59533	3.276268	3.548244	2.367483

*Characteristic for *G. uralensis* ** Characteristic for *G. glabra*.*P*-values calculated based on statistical *t*-test.Value of difference calculated as [(average of *G. glabra*)-(average of *G. uralensis*)]**Table 3:** Calculated relative intensity data of each characteristic peak and detection of the number of peaks.

<i>m/z</i>	299*	315*	341*	369*	323**	325**	337**	339**	391**	Own species character detection	Inverse side character detection
<i>G. uralensis</i>											
Gly5-1	33.24	0	100	15.04	14.58	0	0	0	0	2/4	0/5
Gly5-2	100	12.31	20.50	55.82	3.97	0	0	9.17	0	3/4	0/5
Gly5-3	100	52.12	17.59	55.68	0	0	0	9.11	0	3/4	0/5
Gly19-1	56.36	100	5.27	34.10	0	0	3.72	4.69	0	3/4	0/5
Gly19-2	100	31.66	62.54	57.01	0	0	22.91	19.92	0	4/4	1/5
Gly19-3	100	28.21	38.96	61.31	0	0	0	14.67	0	4/4	0/5
Gly20-1	100	69.85	99.78	35.70	12.34	0	0	38.59	11.29	4/4	1/5
Gly20-2	100	86.93	71.77	33.13	11.07	0	0	21.51	12.11	4/4	1/5
Gly20-3	73.15	70.90	48.91	39.54	0	0	65.96	25.18	0	4/4	2/5
Gly21-1	3.91	3.89	100	8.57	28.08	0	0	13.50	2.62	1/4	1/5
Gly21-2	13.78	15.29	100	6.97	44.04	0	6.47	6.11	11.02	1/4	1/5
Gly21-3	100	84.84	51.43	48.44	0	0	14.76	13.35	0	4/4	0/5
<i>G. glabra</i>											
Gly3-1	0	0	0	0	0	0	0	0	0	0/5	0/4
Gly3-2	0	0	0	0	0	0	51.85	100	25.10	3/5	0/4
Gly3-3	0	0	0	0	0	0	0	100	0	1/5	0/4
Gly17-1	0	0	23.07	0	100	50.58	83.03	42.46	18.90	4/5	1/4
Gly17-2	2.54	0	4.31	2.57	58.58	42.18	70.27	43.83	12.46	4/5	0/4
Gly17-3	0	0	0	0	99.60	48.36	100	49.26	31.28	5/5	0/4
Gly18-1	0	0	100	0	31.70	46.03	39.31	0	87.29	4/5	1/4
Gly18-2	0	0	44.40	0	62.63	56.74	78.67	38.28	79.51	5/5	1/4
Gly18-3	0	0	40.94	0	48.37	100	56.67	63.73	93.37	5/5	1/4

*Characteristic for *G. uralensis* ** Characteristic for *G. glabra*. (20% threshold)In the case of *G. uralensis*, there are four characteristic peaks (*m/z* 299, 315, 341, 369). The number of peaks (*m/z* 299, 315, 341, 369) with relative intensities >20% in each *G. uralensis* sample (Gly5, 19, 20, 21 series) is shown in the "Own species character detection" column.On the other hand, in *G. uralensis*, the number of peaks present characteristic for *G. glabra* with more than 20% relative intensity is shown in "Inverse side character detection" column.In the case of *G. glabra*, five peaks were found (*m/z* 323, 325, 337, 339, 391). The number of peaks (*m/z* 323, 325, 337, 339, 391) with relative intensities >20% in each *G. glabra* sample (Gly3, 17, 18 series) is shown in the "Own species character detection" column.On the other hand, in *G. glabra*, the number of peaks present characteristic for *G. uralensis* at more than 20% relative intensity is shown in "Inverse side character detection" column.

In conclusion, we found several characteristic peaks to identify *G. uralensis* and *G. glabra* by DART-MS using a statistical method; this can be applied to identify these two *Glycyrrhiza* species.

Experimental

Plant material: The roots and stolons of *G. uralensis* Fischer (Hokkaido, Japan) and *G. glabra* Linné (Hokkaido, Japan) were kindly supplied and identified by Dr Toshiro Shibata, the Research Center for Medicinal Plant Resources in Nayoro, Japan. Voucher specimens were deposited at the Dept. of Natural Medicine and Phytochemistry, Meiji Pharmaceutical University, Tokyo, Japan (No.07G0003, No.07G0002, respectively).

The other roots and stolons of *G. uralensis* (China, Russia and Mongolia) and *G. glabra* (Uzbekistan and Kazakhstan) were kindly supplied by Mr Fujio Kanai, Kanai Tokichi Shoten Co., Ltd. Voucher specimens were deposited at the Dept. of Natural Medicine and Phytochemistry, Meiji Pharmaceutical University, Tokyo, Japan (No.11G0001, No. 11G0002, No. 11G0003, No.11G0004, No.11G0005 respectively).

Species identification by HPLC: All samples were identified by HPLC analysis under similar conditions as those described by Shibano *et al.* [13]. A 5C₁₈-AR-II (i.d. 4.6 x 150 mm, Nacalai Tesque Inc.) column was used. HPLC analysis was carried out on D-7000 type software; column oven: L-7300, Diode Array Detector;

L-7450H, pump; L-7100 (HITACHI). The presence of species specific substances was confirmed: *G. uralensis*: glycycomarin, and *G. glabra*: glabridin [3].

Chemicals: All chemicals were analytical reagent grade. Glabridin was purchased from Wako Pure Chemical Industries Ltd. (Osaka, Japan), and glycycomarin was kindly provided by Dr Makio Shibano (Osaka University of Pharmaceutical Science, Japan).

DART mass spectrometry: All experiments were performed using a JMS-T100TD (JEOL Ltd., Japan) orthogonal TOF-MS with a DART ion source, which used helium gas at a flow rate of approximately 2.5 L/min. The gas heater was set at 250°C. The mass spectrometer was operated in positive mode. Orifice 1, ring lens and orifice 2 potentials were set at 50, 10, and 5 V, respectively. The detector voltage was set to 2500 V, and the ion guide potential was 500 V. Data acquisition was taken from *m/z* 50 to 500.

LC/ESI/MS: All samples were analyzed by LC/ESI/MS to confirm that peaks for *m/z* 369 and 325 in the DART-MS were glycycomarin and glabridin, respectively. HPLC analysis samples were diluted and injected into the LC/ESI/MS system {LCMS-2010EV system with an ESI probe (Shimadzu Co. Ltd.)}; column, COSMOSIL 3C₁₈-EB (i.d. 2.0 x 150 mm, Nacalai Tesque Inc.); guard column, OPTI-GUARD 1 mm C₁₈ (Lab Lab Company Co. Ltd.). The heat block and CDL temperatures were 200°C and 250°C,

respectively; nebulization gas, nitrogen (flow rate, 1.5 L/min); solvent, water (0.1% formic acid) and acetonitrile (0.1% formic acid) gradient of 30% to 80% acetonitrile(0.1% formic acid) in 55 min.; flow rate, 0.2 mL/min.

Statistical analysis: The peak tables of DART-MS were imported into Microsoft Excel 2007.

Acknowledgments - The authors would like to express appreciation to Dr Toshiro Shibata (Research Center for Medicinal Plant Resources) for providing authentic *G. uralensis* and *G. glabra*. Appreciation is expressed to Mr Fujio Kanai (Kanai Tokichi Shoten Co., Ltd.) for providing *G. uralensis* and *G. glabra*. Additionally, they are grateful to Makio Shibano (Osaka University of Pharmaceutical Science) for providing glycycomarin.

References

- [1] Part IV of the series: Application of mixture analysis to crude materials from natural resources. (a) Part I: Fukuda E, Baba M, Iwasaki N, Uesawa Y, Arifuku K, Kamo O, Tsubono K, Okada Y. (2010) Identification of *Glycyrrhiza* species by direct analysis in real time mass spectrometry. *Natural Product Communications*, **5**, 1755-1758; (b) Part II: Fukuda E, Yoshida M, Baba M, Uesawa Y, Suzuki R, Kamo O, Koji Tsubono K, Arifuku K, Yatsunami K, Okada Y. (2011) Application to classification of mulberry leaves using multivariate analysis of proton NMR metabolomic data. *Natural Product Communications*, **6**, 1621-1625; (c) Part III: Fukuda E, Baba M, Uesawa Y, Kamo O, Arifuku K, Tsubono K, Okada Y. (2012) Application of mixture analysis to crude materials from natural resources (III): NMR spectral studies to analyze chalcones from *Angelica keiskei*. *Natural Product Communications*, **7**, 21-23.
- [2] Kondo K, Shiba M, Yamaji H, Morota T, Zhengmin C, Huixia P, Shoyama Y. (2007) Species identification of licorice using nrDNA and cpDNA genetic markers. *Biological & Pharmaceutical Bulletin*, **30**, 1497-1502.
- [3] Hayashi H, Hosono N, Kondo M, Hiraoka N, Ikeshiro Y, Shibano M, Kusano G, Yamamoto H, Tanaka T, Inoue K. (2000) Phylogenetic relationship of six *Glycyrrhiza* species based on rbcL sequences and chemical constituents. *Biological & Pharmaceutical Bulletin*, **23**, 602-606.
- [4] Nagai H, Yamamoto Y, Sato Y, Akao T, Tani T. (2006) Pharmaceutical evaluation of cultivated *Glycyrrhiza uralensis* roots in comparison of their antispasmodic activity and glycycomarin contents with those of licorice. *Biological & Pharmaceutical Bulletin*, **29**, 2442-5.
- [5] Chandrasekaran CV, Deepak HB, Thiagarajan P, Kathiresan S, Sangli GK, Deepak M, Agarwal A. (2011) Dual inhibitory effect of *Glycyrrhiza glabra* (GutGard™) on COX and LOX products. *Phytomedicine*, **18**, 278-274.
- [6] Cody BR, Laramée AJ, Durst DH. (2005) Versatile new ion source for the analysis of materials in open air under ambient conditions. *Analytical Chemistry*, **77**, 2297-2302.
- [7] Kim JK, Jang YP. (2009) Direct analysis of curcumin in turmeric by DART-MS. *Phytochemical Analysis*, **20**, 372-377.
- [8] DeRoo CS, Armitage RA. (2011) Direct identification of dyes in textiles by direct analysis in real time-time of flight mass spectrometry. *Analytical Chemistry*, **83**, 6924-6928.
- [9] Li Y. (2012) Confined direct analysis in real time ion source and its applications in analysis of volatile organic compounds of *Citrus limon* (lemon) and *Allium cepa* (onion). *Rapid Communications in Mass Spectrometry*, **26**, 1194-1202.
- [10] Lee SM, Kim HJ, Jang YP. (2011) Chemometric classification of morphologically similar Umbelliferae medicinal herbs by DART-TOF-MS fingerprint. *Phytochemical Analysis*, **23**, 508-512.
- [11] Lancaster C, Espinoza E. (2012) Analysis of select *Dalbergia* and trade timber using direct analysis in real time and time-of-flight mass spectrometry for CITES enforcement. *Rapid Communications in Mass Spectrometry*, **26**, 1147-1156.
- [12] Kuwajima H, Taneda Y, Chen WZ, Kawanishi T, Hori K, Taniyama T, Kobayashi M, Ren J, Kitagawa I. (1999) Variation of chemical constituents in processed licorice roots: Quantitative determination of saponin and flavonoid constituents in bark removed and roasted licorice roots. *Yakugaku Zasshi*, **119**, 945-955.
- [13] Shibano M, Henmi A, Matsumoto Y, Kusano G, Miyase T, Hatakeyama Y. (1997) Studies on the index compounds for HPLC analysis of *Glycyrrhiza uralensis*. *Heterocycles*, **45**, 2053-2060.

Comparison of Total Phenolic Content, Scavenging Activity and HPLC-ESI-MS/MS Profiles of Both Young and Mature Leaves and Stems of *Andrographis paniculata*

Lee Suan Chua^{a*}, Ken Choy Yap^a and Indu Bala Jaganath^b

^aMetabolites Profiling Laboratory, Institute of Bioproduct Development, Universiti Teknologi Malaysia, 81310 UTM Skudai, Johor Bahru, Johor, Malaysia

^bBiotechnology Research Centre, Malaysian Agricultural Research and Development Institute, 50774 Kuala Lumpur, Malaysia

chualeesuan@utm.my

Received: March 22nd, 2013; Accepted: September 15th, 2013

The total phenolic content and radical scavenging activity of *Andrographis paniculata* has been investigated to estimate the amount of phenolic compounds and diterpene lactones, respectively in the plant extracts. The stem extracts exhibited higher total phenolic content and scavenging activity than those of the leaf extracts from both young and mature plants. A range of 19.6–47.8 mg extract of *A. paniculata* from different parts of the plant is equivalent to the scavenging activity exhibited by one mg of standard Trolox. HPLC-ESI-MS/MS was also used to identify simultaneously the phytochemicals from the leaves and stems of both young and mature plant samples. Of the identified compounds, seven of the sixteen diterpene lactones, three of the six flavonoids, five of the six phenolic acids and two cyclic acids are reported here for the first time for this species. Multivariate statistical approaches such as Hierarchical Component Analysis (HCA) and Principle Component Analysis (PCA) have clustered the plant extracts into the leaf and stem groups, regardless of plant age. Further classification based on the phytochemical profiles revealed that mostly phenolic acids and flavonoids were from the young leaf extracts, and diterpenoids and their glycosides from the mature leaf extracts. However, the phytochemical profiles for the stems of both young and mature plants were not significantly different as presented in the dendrogram of HCA and the score plot of PCA. The marker for mature plants might be the m/z 557 ion (dihydroxyl dimethyl 19-[(β -D-glucopyranosyl)oxy]-19-oxo-ent-labda-8(17),13-dien-16,15-olide), whereas the m/z 521 ion (propyl neoandrographolide) could be the marker for leaf extracts.

Keywords: *Andrographis paniculata*, HPLC-ESI-MS/MS, Diterpene lactones, Polyphenols, Phenolic acids.

Andrographis paniculata Nees, family Acanthaceae, is an annual herbaceous plant, commonly known as “King of Bitters”. It is widely grown in the tropical areas of Southeast Asia. The aerial parts (leaf and stem) of the plant have been used widely for centuries as a folklore remedy in traditional Chinese and Ayurvedic medicines for a broad range of ailments, including liver diseases, gastric disorders and the common cold [1]. This plant is the predominant constituent for at least 26 Ayurvedic formulations, as recorded in the Indian Pharmacopoeia [2-4].

Recently, many pharmacological studies have been carried out on the extracts of *A. paniculata*, mainly due to its wide application in the treatment of various infectious diseases and immune disorder symptoms. A comprehensive review of the phytochemistry, pharmacology and clinical uses of *A. paniculata* has been reported by Perumal Samy *et al.* [5]. They also reported that *A. paniculata* has very low toxicity to animals and plants. Diterpenoid lactones and flavonoids are the main chemical constituents believed to be responsible for the biological activities of the plant [6]. Till now, 50 labdane diterpenoids and 30 flavonoids have been reported from this plant [7-10]. Andrographolide is the prime active constituent, with a bitter taste, maximally found in the leaves (> 2%, w/w) [11,12]. Most phytochemical studies have been focused on the aerial part of the plant. Radhika *et al.* [13] reported three flavones from the stem of *A. paniculata*. To the best of our knowledge, no study has been carried out to differentiate the phytochemical profiles of the leaves and stems of both young and mature plants. The presence of diterpenoids, polyphenols and their derivatives in different parts of the plant at different harvest ages has not been simultaneously analyzed and compared so far.

Numerous analytical methods have been reported for the determination of andrographolide and its derivatives in the plant extracts. Of these methods, high performance liquid chromatography has been reported as the most accurate approach, since the spectrophotometric method suffers from the instability of red colored substance formation [14], and the gravimetric method gives an over estimation of andrographolide due to the co-solubility of other substances in ethyl acetate [15]. In the present study, a hyphenated system consisting of high performance liquid chromatography coupled to tandem mass spectrometry was used to determine simultaneously the major phytochemicals in the methanolic extract of *A. paniculata*. The complex chemical composition of the extract was chromatographically separated in a column by a gradient profile of a binary solvent system and then ionized by an electrospray ionization source before fragmentation and detection by a mass analyzer.

Besides the use of the high throughput analytical tool, the plant extracts were usually subjected to total phenol and antioxidant property assays. The total phenolic content is widely applied for the quantitative estimation of polyphenols such as flavones, flavanones and flavonols. The presence of diterpene lactones could be estimated from the free radical scavenging activity assay. The antiradical mechanism of diterpene lactones, particularly neoandrographolide, has been reported because of the presence of allylic hydrogens on the unsaturated lactone ring [16]. Therefore, diterpene lactones could scavenge free radicals by donating the allylic hydrogen atoms from its aglycone, either by homolytic cleavage or by deprotonation-oxidation mechanisms.

Table 1 shows the results of the TPC and DPPH activity of the leaves and stems of both young and mature plant samples. The stems had higher TPC values than the leaf samples, with the stems of the young plants exhibiting the highest TPC value of 1.64 μg GAE/mg extract. In line with the TPC results, the stem extracts also exhibited higher free radical scavenging activity than the leaf extracts. Therefore, the scavenging activity of the plant extracts might be attributed to the presence of phenolic compounds and diterpene lactones [16,17]. The scavenging activity expressed by the stem extracts was about two-fold higher than that of the leaves of mature plants. However, there was only a small degree of difference in scavenging activity between the leaves and stems of young plants. This observation also agreed with the TPC of the leaves and stems of young plants, where the degree of difference in the TPC value of both extracts was relatively small compared with the mature plant extracts.

Table 1: Total phenolic content and scavenging activities of different parts of *A. paniculata*.

^a Extract	^b TPC		^c DPPH		Scavenging activity
	(μg GAE/mg extract)		(μg TE/mg extract)		index
	mean	^d SD	mean	^d SD	(mg extract/mg Trolox)
YL	1.43	0.0032	36.00	0.0000	28.5
YS	1.64	0.0280	40.89	0.3849	25.2
ML	1.13	0.0056	20.89	1.3878	47.8
MS	1.51	0.0000	52.89	0.3849	19.6

^aYL, YS, ML and MS are denoted for young leaf, young stem, mature leaf and mature stem, respectively.

^bTotal phenolic content in μg of gallic acid equivalents in a mg of extract.

^cDPPH scavenging activity in μg of Trolox equivalent in a mg of extract.

^dStandard deviation of triplicate data.

The capacity of scavenging activity of plant extracts was also compared with the standard chemical Trolox. The comparison was carried out by using the scavenging activity index, which is defined as the amount of plant extract required to exhibit the same scavenging activity as a milligram of Trolox. It means that the lower the value of the index, the higher the scavenging activity of the plant extract. The scavenging activity index of the leaves and stems from mature plant extracts was the highest (47.8) and lowest (19.6) value, respectively. Therefore, the stems of mature plants exhibited the highest scavenging activity (52.89 μg TE/mg extract), and the leaves of mature plants the lowest (20.89 μg TE/mg extract) among the plant extracts.

A high sensitivity hyphenated liquid chromatograph integrated with a tandem mass spectrometer was used to identify phytochemicals from different parts of the methanolic *A. paniculata* extracts. Sixteen bicyclic diterpenoids, six flavonoids (2 flavanones and 4 flavones), and eight cyclic and phenolic acids were detected in this study. The compounds and their fragment ions are tabulated in Table 2. The presence of these metabolites was confirmed by referring the data to the theoretical fragments generated by MS Fragmenter 12.0 and the spectral data from literature values [18].

Table 2 also shows that bicyclic diterpenoids were mostly detected in the leaves, in particular the higher molecular weight glycosidic diterpenes from the leaves of mature plants. However, cyclic and phenolic acids were mostly detected from the leaves of young plants. It is interesting to note that andrographolide was only detected in the leaf extracts. This finding was not in line with the previous result reporting the existence of andrographolide in all parts of the plant, maximally in the leaves [19]. The contradictory finding might be due to geographical and climatic variation of plant origin. However, Arpini *et al.* [20] reported no significant difference between the chemical fingerprints of the aerial parts of plants from India and China.

In addition to the detection of nine known bicyclic diterpenoids, seven other diterpenoidal derivatives were detected from the plant grown in Malaysia (Table 2). Of the flavonoids, three (flavones) are reported for the first time for *A. paniculata*. Seven cyclic and phenolic acids are also reported for the first time for *A. paniculata*. However, the presence of these compounds needs to be confirmed, for example by nucleus magnetic resonance spectroscopy.

Hierarchical clustering analysis (HCA) was carried out based on the presence and absence of phytochemicals identified from the plant extracts, without considering the peak signal intensity of the metabolites (Figure 2). This kind of chemometric evaluation has been used by researchers from India who revealed that it is a useful complementary technique for quality determination of *A. paniculata* from different origins [21]. Significantly, the leaf extracts of both young and mature plants were in the same cluster, whereas the stem extracts were in another (Figure 2a). This observation could be explained by high similarity in the phytochemical profile of the same part of the plant, even at different plant ages. However, the dendrogram of variables segregated the identified phytochemicals into three major groups, as presented in Figure 2b. Mostly phenolic acids and flavonoids were clustered in the same group, whereas diterpenoids and their glycosides were grouped in the same class. Another group of metabolites consisted of the mixture of diterpenoids and phenols. The phytochemicals in the same class most probably are involved in a similar pathway of secondary metabolism. This is because these compounds are important secondary metabolites for plant defense. Both mevalonic acid (MVA) and methyl erythritol phosphate (MEP), and deoxyxylulose (DXP) pathways have been suggested for the biosynthetic routes to the production of andrographolide [22].

Another unsupervised multivariate analysis, principle component analysis (PCA), has been used to reduce the multidimensional mass spectral data into 12 principle components using Pareto scaling. The setting parameters were minimum spectral peak width, 0.3 Da; mass tolerance, 0.1 Da and retention time tolerance, 0.1 min. The first four components were covered for 51.1% of the total variation. In line with the pattern recognition result of HCA, the leaf and stem extracts were significantly clustered into two major groups; the positive and negative regions, respectively in the first principle component (PC1) of the score plot, regardless of the plant age (Figure 3a). In the positive region of the second principle component (PC2), the young leaf extracts were further separated from the mature leaf extracts. Thus, the plant extracts divided into three groups, namely young leaf extracts, mature leaf extracts, and young and mature stem extracts. This classification agreed with the grouping based on HCA for the identified metabolites (Figure 2b). The first group, consisting of phenolic acids and flavonoids, might be attributed to the young leaf extracts. The second group, consisting of diterpenoids and their glycosides, might be due to the mature leaf extracts. Most probably, the third group, which consisted of a diterpenoid and phenol mixture, was due to the metabolites extracted from the stems of both young and mature plants. The observation also shows that the constituents of the stem extracts share the nearest similarity in their phytochemical profiles, regardless of plant age. From the loading plot (Figure 3b) and the “compare” function of ACD software under the module of SpecManager 12.01, the marker ion for the mature plant might be m/z 557 (dihydroxyl dimethyl 19-[(β-D-glucopyranosyl)oxy]-19-oxo-ent-labda-8(17),13-dien-16,15-olide), whereas the m/z 521 ion (propyl neoandrographolide) could be the marker for the leaf extracts.

Table 2: Bicyclic diterpenoids, flavonoids, phenolic acids and cyclic acids detected from the methanolic leaf and stem extracts of both young and mature *Andrographis paniculata*.

Compound Name	Formula	MW (g/mol)	Rt (min)	Fragment ion at negative mode (m/z)	^a YL	^b YS	^c ML	^d MS	References
Bicyclic diterpenoids (M)									
3,19-Dihydroxy-14,15,16-trinor-ent-labda-8(17),11-diene-13-oic acid	C ₁₇ H ₂₈ O ₄	294	13.24	293/275(-H ₂ O)/249(-C ₃ H ₅)	-	-	√	-	[9]
Andrographolactone	C ₂₀ H ₃₂ O ₂	296	13.80	295/277(-H ₂ O)/251(-CO ₂)/233(-H ₂ O)/195/171/165	-	√	√	√	[26]
14-Deoxy-11,12 dihydroandrographolide	C ₂₀ H ₃₂ O ₄	332	7.57	331/303(-CO)/287(-CO ₂)/283/269/255/239/213/197/171/145/108/69	√	√	√	√	[7, 27, 28]
3,19-Dihydroxy-ent-labda-8(17),12-dien-16,15-olide	C ₂₀ H ₃₀ O ₄	334	8.74	333/313/285(-CO)/269(-CO ₂)/257/241/69	√	-	√	-	[9]
Andrographolide	C ₂₀ H ₃₀ O ₅	350	3.49	349/331(-H ₂ O)/305(-CO ₂)/303(-CO ₂ H ₂)/287(-2OCH ₃)/283/273(-2CH ₃ -CO)/265/253/243/239/245/83/67/55	√	-	√	-	[7, 27, 28]
14-Deoxy-11-hydroxyandrographolide	C ₂₀ H ₃₀ O ₅	350	6.41	349/331(-H ₂ O)/305(-CO ₂)/301(-2CH ₃)/287(-2OCH ₃)/273/267/255/253/239/213/199/171/145/133/108/83/69	-	√	-	-	[7, 29]
3,18,19-Trihydroxy-ent-labda-8(17),13-diene-16,15-olide	C ₂₀ H ₃₀ O ₅	350	2.91	349/331(-H ₂ O)/305(-CO ₂)/301(-2CH ₃)/299/287(-2OCH ₃)/283/265/253/239/217/215/199/171/145/83/67/55	√	-	√	-	[9]
Methyl methoxyl andrographolide	C ₂₁ H ₃₂ O ₇	396	7.19	395(+CO+H ₂ O)/377(+CO)/349(M-H)/331(-H ₂ O)/303(-CO)/287(-CO ₂)/255/239/213	√	√	√	√	The present study
14-Deoxy-11-hydroandrographolide dimer glycoside	C ₄₀ H ₆₀ O ₁₄	844	5.87	843(2M+Glu-2H ₂ O)/349(M-H)/331(-H ₂ O)/287(-CO ₂)/267/239/213	-	-	√	-	The present study
Neoandrographolide	C ₂₀ H ₃₀ O ₈	480	9.13	479/317(-Glu)/289(-CO)/161(Glu-H)/113/101/85/59/57	√	√	√	-	[7, 27]
Propyl neoandrographolide	C ₂₃ H ₄₀ O ₈	522	10.13	521(+C ₃ H ₆)/479(M-H)/461(-H ₂ O)/413/401/317(-Glu)/289(-CO)/161(Glu-H)/101/85/59	√	-	√	-	The present study
Methyl methoxyl neoandrographolide	C ₂₃ H ₄₀ O ₉	526	9.74	525(+CH ₃ +OCH ₃)/479(M-H)/317(-Glu)/161(Glu-H)	√	√	√	√	The present study
Propyl neoandrographolide dimer	C ₄₀ H ₆₀ O ₁₆	1132	10.32	1131(+2CO ₂)/1043(2M-H)/521(M-H)/479(-C ₃ H ₆)/461(-H ₂ O)/413/317(-Glu)/161(Glu-H)/143/59	-	-	√	-	The present study
14-Deoxyandrographiside	C ₂₀ H ₄₀ O ₉	496	8.57	495/333(-Glu)/303(-2CH ₃)/285(-H ₂ O)/161(Glu-H)/113/101/85	-	-	√	-	[27, 29, 30]
Methyl methoxyl 14-deoxyandrographiside	C ₂₁ H ₄₂ O ₁₁	542	8.71	541(+OCH ₃ +CH ₃)/495(M-H)/333(-Glu)/285(-2CH ₃ -H ₂ O)/161(Glu-H)/113/101	√	-	√	√	The present study
Dihydroxyl dimethyl 19-[(β-D-glucopyranosyl)oxy]-19-oxo-ent-labda-8(17),13-dien-16,15-olide	C ₂₈ H ₄₆ O ₁₁	558	6.42	557(+2OH+2CH ₃)/493(M-H)/449(-CO ₂)/331(M-Glu)/303(-CO)/287(-CO ₂)/161(Glu-H)/101	-	-	√	√	The present study
Flavavone and Flavones									
2'-Hydroxy-2,4',6'-trimethoxychalcone	C ₁₈ H ₁₈ O ₅	314	11.23	313/298(-CH ₃)/283(-CH ₃)/255(-CO)/211(-CO ₂)/183(-CO)/165/137	-	-	-	√	[31]
7-O-Methyldihydrowogonin	C ₁₇ H ₁₆ O ₅	299	15.10	299/281(-H ₂ O)/253(-CO)/169	√	√	-	√	[32]
Dihydroskullcapflavone I	C ₁₇ H ₁₆ O ₆	316	12.96	315/297(-H ₂ O)/229/155/113	-	-	-	√	[31]
7-O-Methylwogonin	C ₁₇ H ₁₆ O ₅	298	14.30	297/279(-H ₂ O)/261(-H ₂ O)/183(-benzyl)	√	√	√	√	[33]
Apigenin-7-O-glucoronide	C ₂₁ H ₁₈ O ₁₁	446	7.01	445/269(-Gln)/225(-CO ₂)/183/149/117/85	-	-	√	-	The present study
Luteolin-7-O-glucoronide	C ₂₁ H ₁₈ O ₁₂	462	7.40	461/285(-Gln)/255(-OCH ₃)/241(-CO ₂)/213(-CO)/187/145	-	-	√	√	The present study
Diosmetin-7-glycoside	C ₂₁ H ₁₈ O ₁₂	462	9.10	461/299(-Glu)/269(-2CH ₃)/241(-CO)/184/165/137	-	√	-	-	The present study
Cyclic and Phenolic acids									
Dihydroxyl glucosyl cyclohexane	C ₁₇ H ₂₆ O ₄	294	1.16	293/131(-Glu)/113(-H ₂ O)/85(-CO)/59	√	-	√	√	The present study
Quinic acid	C ₇ H ₁₂ O ₆	192	1.55	191/173(-H ₂ O)/127(-CH ₂ -OCH ₃)/108/93/87/85/67	√	√	√	√	The present study
Coumaroylquinic Acid	C ₁₆ H ₁₈ O ₈	338	1.74	337/191(quinic acid)/173(-H ₂ O)/163(coumaric acid)/119/93/85	√	-	-	-	The present study
Caffeoylquinic Acid (Chlorogenic acid)	C ₁₆ H ₁₈ O ₉	354	1.75	353/191(quinic acid)/179(caffeic acid)/173(-H ₂ O)/135/127/93/85	-	-	-	√	[34]
Feruloylquinic acid	C ₁₇ H ₂₀ O ₉	368	7.35	367/337(-COH ₂)/323(-CO ₂)/305(-H ₂ O)/293(-COH ₂)/257(-2H ₂ O)/241/229/217/69	-	√	-	-	The present study
Caffeic glycoside	C ₁₅ H ₁₈ O ₉	342	0.97	341/179 (-Glu)/161(Glu-H)/143(-H ₂ O)/113(-2CH ₃)/101/59	√	-	√	-	The present study
Hydroxyl cyclohexyl chlorogenic acid	C ₂₂ H ₃₂ O ₁₀	452	1.36	451/405(-COH ₂)/353(chlorogenic acid)/345/191(quinic acid)/179(caffeic acid)/165/139/59	-	-	-	√	The present study
Chlorogenic glycoside	C ₂₂ H ₃₂ O ₁₄	516	6.80	515/353(-Glu)/255/203/191(quinic acid)/179(caffeic acid)/173(-H ₂ O)/135/137/93/67	√	-	-	-	The present study

^aYL : young leaf extracts; ^bYS: young stem extracts; ^cML: mature leaf extracts; ^dMS: mature stem extracts.

The simultaneous identification of plant metabolites using the high throughput HPLC-ESI-MS/MS approach has revealed that different parts of *A. paniculata* have different phytochemical profiles. The difference could also be seen for the same plant part at different plant ages. However, the difference was not significant for the stems of young and mature plant extracts based on HCA and PCA. Besides that, a difference has also been proven in total phenolic content and free radical scavenging activity of the plant extracts.

The stem extracts of the plant were reported to have higher total phenols and radical scavenging activity than the leaf extracts. The degree of difference became significant as the plant grew. More glycosidic diterpenoid lactones were detected in the leaves of the mature plant extract. This observation was in line with the previous findings that the antioxidant property of flavonol aglycones was higher than their glycosides [23, 24].




Figure 2: Hierarchical clustering analysis of plant extracts (a) and identified metabolites (b).




Figure 3: Score (a) and loading (b) plots of mass spectral data from various parts of plant extracts. YL: young leaves; YS: young stems; ML: mature leaves; MS: mature stems.

Experimental

Chemicals and reagents: HPLC-grade of acetonitrile and methanol were obtained from Fisher Scientific (Pittsburg, USA). Ammonium formate and formic acid were purchased from Fluka (Buchs, Switzerland). Folin-Ciocalteu reagent and 1,1-diphenyl-2-picrylhydrazyl (DPPH) were sourced from Sigma-Aldrich (St. Louis, MO). Gallic acid (98%) and 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox, 97%) were obtained from Acros Organics (Pittsburgh, PA). 18.2 Megaohm^{cm} water was produced from a Barnstead NANOpure Diamond water purification system (State of Illinois, USA).

Plant material: Wild plant samples of *A. paniculata* were collected and authenticated by the Malaysian Agricultural Research Development Institute (MARDI), Selangor, Malaysia. The voucher specimen (MARDI 4770) has been deposited in the Herbarium of MARDI. The plant samples were cleaned and segregated into young (< 4 weeks old) and mature plants (> 6 weeks old). The mature plants usually start flowering when 6 to 8 weeks old. The leaves and stems were separated and air-dried indoors at room temperature for one day. The semi-dried plant samples were pulverized in liquid nitrogen and further freeze-dried for another 2 days at -80°C.

Methanolic extraction of plant samples: The dried and finely powdered leaves (500 mg) and stems (500 mg) of *A. paniculata* were extracted with methanol: water (1:1, 10 mL) using a shaker at

room temperature (25°C). After 60 min of extraction, the extracts were filtered and stored at -20°C for subsequent analysis.

Total phenolic content: The total phenolic content (TPC) of the extracts was analyzed using Folin-Ciocalteu reagent, based on the method described by Chua *et al.* [25], with minor modification. The extracts (1.5 mg/mL) were mixed with 1500 µL of Folin-Ciocalteu reagent. Sodium carbonate (100 g/L; 1200 µL) was added to the mixture after 5 min. The solution was shaken thoroughly and incubated for 2 h at 30°C, before the absorbance was measured at 765 nm using a UV-Vis spectrophotometer (Perkin-Elmer Lambda 25, Waltham, MA). Gallic acid (0-200 µg/mL) was used as a standard for calibration curve preparation. The TPC was expressed as µg of gallic acid equivalents (GAE) in mg of dry plant extract. All assays were carried out in triplicate.

Free radical-scavenging activity by DPPH: The antioxidant property of *A. paniculata* extracts was determined by the DPPH (1,1-diphenyl-2-picrylhydrazyl) assay. The plant extracts (1.5 mg) were dissolved in 1 mL methanol and the solutions (0.1 mL) were added to 3 mL of DPPH• solution (1 mM). The absorbance was measured at 515 nm after 30 min of incubation at 30°C. Trolox was used as positive control and prepared in a series of concentrations ranging from 0 – 600 µg/mL for a calibration curve. The ability to scavenge DPPH• was calculated using Eq. (1), where W is the dry weight of either Trolox (standard chemical) or plant extract, and A_{control} and A_{sample} are the absorbance of control and sample, respectively. The equation was used to calculate the scavenging activity index. The experiment was performed in triplicate.

$$\text{Scavenging activity (mg}^{-1}\text{)} = [(A_{\text{control}} - A_{\text{sample}}) / A_{\text{control}}] / W \quad (1)$$

HPLC-ESI-MS/MS: A hyphenated system consisted of a high performance liquid chromatograph (Agilent 1200; Agilent Technologies Inc., Santa Clara, CA) and a triple quadrupole-linear ion trap tandem mass spectrometer (Applied Biosystems 3200 Q TRAP; Life Technologies Corporation, Carlsbad, CA) was used for phytochemical separation and identification, respectively. The phytochemicals were separated using a C18 reserved phase Aqua column (2.1 x 50 mm, 5µm) from Phenomenex, Torrance, CA, and then ionized by an electrospray ionization source (ESI) before detection. The mobile phase was a binary solvent system consisting of solvent A (water with 0.1% formic acid and 5 mM ammonium formate) and solvent B (acetonitrile with 0.1% formic acid and 5 mM ammonium formate). The gradient was: 0-20 min, 10-90 % B; 20-25 min, 90 % B; 25.1-30 min, 10 % B for column equilibration before the next run. The total runtime was 30 min. The flow rate was 0.2 mL/min and the injection volume was 20 µL. All samples were filtered through a 0.2-µm nylon membrane filter prior to injection. The mass spectra were acquired from m/z 100–1200 with a 20-ms ion accumulation time. All mass spectrometric data were acquired in negative ionization mode. The capillary and voltage of the ESI source were maintained at 400°C and 4.5 kV, respectively. All other parameters were as follows: nitrogen was used as ion source gas for nebulisation, 40 psi; for drying solvent, 40 psi; curtain gas, 10 psi; collision gas, high; declustering potential, 40 V, and collision exit energy, 10 V. The scan rate was 1000 amu/s. Data acquisition and data processing were performed using Analyst 1.4.2. The scan mode of enhanced mass spectra (EMS) was used to screen the sample profile. Enhanced product ion (EPI) scan was used to determine the characteristic ions and to confirm the presence of aglycone peaks.

Data processing and interpretation: The data processing software, MarkerView 1.2 (Applied Biosystems/MDS Sciex), was applied to

perform sample classification on triplicate mass spectra of different parts of the plants at different harvest ages by using principle component analysis (PCA). Hierarchical clustering analysis (HCA) was carried out to classify the plant extracts based on sample extracts, as well as identified phytochemicals using R version

2.11.1. The module of MS Fragmenter 12.0 and SpecManager 12.01 in Advanced Chemistry Development (ACD, Toronto, Canada) software was used to predict compound fragmentation and for data mining, respectively.

References

- [1] Radhika P, Lakshmi KR. (2010) Antimicrobial activity of the chloroform extracts of the root and the stem of *Andrographis paniculata* Nees. *International Research Journal of Microbiology*, **1**, 37-39.
- [2] Mishra SK, Sangwan NS, Sangwan RS. (2007) *Andrographis paniculata* (Kalmegh): A review. *Pharmacognosy Reviews*, **1**, 283-298.
- [3] Raini AP, Kumar A, Pareek SK. (2007) HPTLC analysis of hepatoprotective diterpenoid andrographolide from *Andrographis paniculata* Nees (Kalmegh). *Indian Journal of Pharmaceutical Sciences*, **69**, 473-475.
- [4] Kapil A, Koul IB, Banarjee SK, Gupta BD. (1993) Antihepatotoxic effects of major diterpenoid constituents of *Andrographis paniculata*. *Biochemical Pharmacology*, **46**, 182-185.
- [5] Perumal Samy R, Thwin MM, Gopalakrishnakone P. (2007) Phytochemistry, pharmacology and clinical use of *Andrographis paniculata*. *Natural Product Communications*, **2**, 607-618.
- [6] Tang W, Eisenbrand G. (1992) *Chinese drugs of plant origin: Chemistry, pharmacology and use in traditional and modern medicine*. Springer-Verlag, Berlin.
- [7] Matsuda T, Kuroyanagi M, Sugiyama S, Umehara K, Ueno A, Nishi K. (1994) Cell differentiation inducing diterpenes from *Andrographis paniculata* Nees. *Chemical and Pharmaceutical Bulletin*, **42**, 1216-1225.
- [8] Koteswara Rao Y, Vimalamma G, Rao CV, Tzeng YM. (2004) Flavonoids and andrographolides from *Andrographis paniculata*. *Phytochemistry*, **65**, 2317-2321.
- [9] Chen LX, Qiu F, Wei H, Qu GX, Yao XS. (2006) Nine new ent-labdane diterpenoids from the aerial parts of *Andrographis paniculata*. *Helvetica Chimica Acta*, **89**, 2654-2664.
- [10] Wu TS, Chern HJ, Damu AG, Kuo PC, Su CR, Lee EJ, Teng CM. (2008) Flavonoids and ent-labdane diterpenoids from *Andrographis paniculata* and their antiplatelet aggregatory and vasorelaxing effects. *Journal of Asian Natural Products Research*, **10**, 17-24.
- [11] Pandey MK. (2011) Physicochemical standardization of *Andrographis paniculata* (Nees): An Ayurvedic drug. *International Journal of Pharmaceutical Research and Development*, **3**, 81-89.
- [12] Sharma A, Lal K, Handa SS. (1992) Standardization of the Indian crude drug Kalmegh by high pressure liquid chromatographic determination of andrographolide. *Phytochemical Analysis*, **3**, 129-131.
- [13] Radhika P, Prasad YR, Lakshmi KR. (2010) Flavones from the stem of *Andrographis paniculata* Nees. *Natural Product Communications*, **5**, 59-60.
- [14] Maiti PC, Kanji SK, Chatterjee R. (1959) Studies in Kalmegh extract. *Indian Journal of Pharmacology*, **21**, 169-171.
- [15] Indian Pharmacopoeia Committee (1955) *Pharmacopoeia of India*: (the Indian pharmacopoeia). Manager of Publications, Delhi, 1st ed.
- [16] Kamdem RE, Sang S, Ho CT. (2002) Mechanism of the superoxide scavenging activity of neoandrographolide – A natural product from *Andrographis paniculata* Nees. *Journal of Agricultural and Food Chemistry*, **50**, 4662-4665.
- [17] Ojha SK, Nandave M, Kumari S, Arya DS. (2009) Antioxidant activity of *Andrographis paniculata* in ischemic myocardium of rats. *Global Journal of Pharmacology*, **3**, 154-157.
- [18] Pellati F, Orlandini G, Pinetti D, Benvenuti S. (2011) HPLC-DAD and HPLC-ESI-MS/MS methods for metabolite profiling of propolis extracts. *Journal of Pharmaceutical and Biomedical Analysis*, **55**, 934-948.
- [19] Jarukamjorn K, Nemoto N. (2008) Pharmacological aspects of *Andrographis paniculata* on health and its major diterpenoid constituent andrographolide. *Journal of Health Science*, **54**, 370-381.
- [20] Arpini S, Fuzzati N, Giori A, Martino E, Mombelli G, Pagni L, Ramaschi G. (2008) HPLC-DAD-MS fingerprint of *Andrographis paniculata* (Burn. F.) Nees (Acanthaceae). *Natural Product Communications*, **3**, 1981-1984.
- [21] Sharma SN, Jha Z, Sharma DK. (2011) Chemometrics evaluation of the herbal drug *Andrographis paniculata*. *Natural Product Communications*, **6**, 1929-1932.
- [22] Srivastava N, Akhila A. (2010) Biosynthesis of andrographolide in *Andrographis paniculata*. *Phytochemistry*, **71**, 1298-1304.
- [23] Arora A, Nair MG, Strasburg GM. (1998) Structure-activity relationships for the antioxidant activities of a series of flavonoids in a liposomal system. *Free Radical Biology and Medicine*, **24**, 1355-1363.
- [24] Sanbongi C, Osakabe N, Natsume M, Takizawa T, Gomi S, Osaka T. (1998) Antioxidant polyphenols isolated from *Theobroma cacao*. *Journal of Agricultural and Food Chemistry*, **46**, 454-457.
- [25] Chua LS, Abdul Latiff N, Lee SY, Lee CT, Sarmidi MR, Abdul Aziz R. (2011) Flavonoids and phenolic acids from *Labisia pumila* (Kacip Fatimah). *Food Chemistry*, **127**, 1186-1192.
- [26] Wang GC, Wang Y, Williams ID, Sung HHY, Zhang XQ, Zhang DM, Jiang RW, Yao XS, Ye WC. (2009) Andrographolactone, a unique diterpene from *Andrographis paniculata*. *Tetrahedron Letters*, **50**, 4824-4826.
- [27] Radhika P, Prasad YR, Sowjanya K. (2012) A new diterpene from the leaves of *Andrographis paniculata* Nees. *Natural Product Communications*, **7**, 485-486.
- [28] Hapuarachchi SD, Ali Z, Abe N, Sugandhika ST, Sandun ST, Khan IA. (2013) Andrographidine G, a new flavones glucoside from *Andrographis paniculata*. *Natural Product Communications*, **8**, 333-334.
- [29] Kulyal P, Tiwari UK, Shukla A, Gaur AK. (2010) Chemical constituents isolated from *Andrographis paniculata*. *Indian Journal of Chemistry*, **49B**, 356-359.
- [30] Sule A, Ahmed QU, Samah OA, Omar MN, Hassan NM, Kamal LZM, Yarmo MA. (2011) Bioassay guided isolation of antibacterial compounds from *Andrographis paniculata*. *American Journal of Applied Sciences*, **8**, 525-534.
- [31] Kishore PH, Reddy MVB, Reddy MK, Gunasekar D, Caux C, Bodo B. (2003) Flavonoids from *Andrographis lineate*. *Phytochemistry*, **63**, 457-461.
- [32] Gupta KK, Taneja SC, Dhar KL. (1996) Flavonoid glycoside of *Andrographis paniculata*. *Indian Journal of Chemistry*, **35B**, 512-513.
- [33] Kuroyanagi M, Sato M, Ueno A, Nishi K. (1987) Flavonoids from *Andrographis paniculata*. *Chemical and Pharmaceutical Bulletin*, **35**, 4429-4435.
- [34] Satyanarayana D, Mythirayee C, Krishnamurthy V. (1978) Polyphenols of *Andrographis paniculata* Nees. *Leather Science*, **25**, 250-251.

Xanthenes from aerial parts of *Hypericum laricifolium* Juss.Irama Ramírez-González^{a,*}, Juan Manuel Amaro-Luis^b and Ali Bahsas^b^aInstituto de Investigaciones, Facultad de Farmacia y Bioanálisis, Universidad de Los Andes, Mérida^bLaboratorio de Productos Naturales, Departamento de Química, Facultad de Ciencias, Universidad de Los Andes, Mérida, República Bolivariana de Venezuela

irama@ula.ve

Received: February 26th, 2013; Accepted: July 20th, 2013

From the aerial parts of *Hypericum laricifolium* Juss., twelve compounds were isolated and identified. They were the xanthenes: 1-hydroxy-7-methoxy-xanthone (**1**), 1,7-dihydroxy-xanthone (**2**), 2-hydroxy-xanthone (**3**), 6-deoxyisojacareubin (**4**), 1,3-dihydroxy-6-methoxy-xanthone (**6**), and 1,5,6-trihydroxy-7-methoxy-xanthone (**7**), together with β -sitosterol, betulinic acid, vanillic acid, isoquercitrin and a mixture of quercetin and isorhamnetin. All the compounds were characterized by spectroscopic and mass spectrometric methods, and by comparison with literature data. This is the first report on the presence of xanthenes in *H. laricifolium*. 1,3-Dihydroxy-6-methoxy-xanthone has been previously synthesized, but this is the first report of its isolation from a natural source.

Keywords: *Hypericum laricifolium*, Guttiferae, Xanthenes, Flavonoids, 1,3-Dihydroxy-6-methoxy-xanthone.

The genus *Hypericum*, family Guttiferae, contains about 475 species, occurring world-wide, which have been included in 36 taxonomic sections on the basis of morphological characters [1]. We have undertaken a phytochemical study of *H. laricifolium* Juss., a sclerophyllous shrub abundant in moor habitats and whose area of distribution extends from western Venezuela to north Peru [2]. In Venezuela, it is traditionally known as “huesito” and is distributed mainly in Mérida State [3]. Several ethnobotanical uses and medicinal properties are attributed to this plant; thus, in Peru and Colombia its flowers are used as a source of natural dyes [4], and in Ecuador the leaves are used for cattle food, the stems to make charcoal, and the wood to build houses, and farming instruments [5]. In Andean traditional medicine, *H. laricifolium* is recognized for its antibacterial, antiviral and antifungal activity [6], and by its efficacy in the treatment of gastrointestinal affections [7] and psychosomatic disorders [8]. A decoction of this plant is drunk to treat cold as well as for treating skeletal pain and as a tranquilizer [5]. To the best of our knowledge, there is only one report dealing with phytochemical studies of this species, collected in Ecuador, in which triterpenes, flavonoids and other phenolic compounds were isolated [9].

Here, we present the results of a chemical study of the aerial parts of *H. laricifolium* gathered in Piedras Blancas moor (Mérida State, Venezuela). Chromatographic separation of the dichloromethane extract resulted in the isolation of compounds **1**, **2**, **3** and **4**, as well as β -sitosterol, betulinic acid and vanillic acid. Compound **1** was identified as 1-hydroxy-7-methoxy-xanthone by comparison of its NMR spectroscopic and mass spectrometric data with those reported in the literature [10a-c]. This xanthone has been reported earlier from other *Hypericum* species [10a-10c]. Spectral data of compound **2** were very similar to those of compound **1**. The only notable differences were the absence of signals attributed to the C-1 methoxyl group and the appearance, in the ¹H NMR spectrum, of a new singlet at δ_H : 9.06, assigned to an additional hydroxyl proton (OH-7). Detailed analysis of its 2D-NMR spectrum led to the conclusion that **2** is 1,7-dihydroxy-xanthone, which has been previously isolated from several *Hypericum* species [10a,c, 11a,b]. The compound **3**, based on ¹H, ¹H-COSY and HMBC data,

was identified as 2-hydroxy-xanthone [10a, 11b, 12]. Compound **4** was identified from its NMR and mass spectrometric data as 6-deoxyisojacareubin. To the best of our knowledge this xanthone has been isolated previously from only two *Hypericum* species: *H. japonicum* [11a,13] and *H. wightianum* [14]. Treatment of **4** with AcO₂/pyridine yielded 1,5-di-O-acetyl-6-deoxyisojacareubin (**5**), which was identified on the basis of its spectral data. From the dichloromethane extract were also isolated betulinic and vanillic acids; both have been previously reported in *Hypericum* species [14-16]. From the methanol extract, the flavonoids isoquercitrin, quercetin and isorhamnetin were isolated and identified by comparison of their spectroscopic data with those reported in the literature. These three flavonoids have been isolated from other *Hypericum* species [10b,c, 17a,b].

Finally, xanthenes **6** and **7** were also isolated from this extract. Analysis of ¹H NMR and HMBC data identified compound **6** as 1,3-dihydroxy-6-methoxy-xanthone and compound **7** as 1,5,6-trihydroxy-7-methoxy-xanthone. Compound **6** has been previously synthesized [18], but, up to now, has not been reported as a natural product. It is particularly interesting to emphasize that xanthone **6** is a powerful inductor of apoptosis in cells transfected with human multidrug-resistance protein 1 (MRP1) [34]. Compound **7** has been reported previously from *Harungana madagascarensis* Lam ex Poir [19a] and *Hypericum ascyron* L. [19b].

Experimental

Plant material: *Hypericum laricifolium* Juss. was collected at paramo Piedras Blancas, Municipio Autónomo Miranda, Estado Mérida, Venezuela, in December 2008. A voucher specimen (J. M. Amaro & I. Ramírez, N° 1645) was deposited at Herbarium MERF of the University of Los Andes (ULA).

General: Melting points, Fisher-Johns apparatus; UV, Perkin-Elmer, Lambda 3B spectrophotometer; IR, Perkin-Elmer FT-1725X spectrophotometer; 1D and 2D-NMR, Bruker-Avance DRX400; EI MS, Hewlett-Packard 5930A; TLC, silica gel PF 254; CC, silica gel 60 (70-230 mesh.).

Extraction: Dried leaves and flowers, finely pulverized (≈ 5.7 Kg), were exhaustively extracted with dichloromethane at room temperature. The extract obtained was concentrated *in vacuo* to afford brown oil (≈ 460 g). Subsequently, plant material was further extracted under reflux in a Soxhlet apparatus with methanol and by concentrating the methanolic solution a new extract (≈ 750 g) was obtained.

Isolation of the constituents: The dichloromethane extract was subjected to vacuum CC on silica gel eluted with *n*-hexane, dichloromethane and acetone, and binary mixtures of these solvents, in order of increasing polarity. Fractions of 1 L were collected, which were concentrated *in vacuo*, analyzed by TLC and suitably combined; 20 fractions were obtained (A-T). Rechromatography of C (≈ 27 g) on a silica gel column eluted with mixtures of *n*-hexane-acetone (17:3) gave β -sitosterol (≈ 32 mg), betulinic acid (≈ 45 mg) and compound **1** (≈ 38 mg), which precipitated as an amorphous solid, homogeneous by TLC. Fraction D (≈ 64 g) was rechromatographed on a silica gel column, and fractions eluted with *n*-hexane-acetone (1:1; 2:3) were combined, concentrated *in vacuo* and passed through a column of Sephadex LH-20 affording compounds **2** (≈ 42 mg) and **4** (≈ 85 mg).

Concentration of fraction E (≈ 45 g) provided a solid residue that was partially purified by flash chromatography on silica gel

developed with *n*-hexane-acetone (4:1). Final purification, which was carried out by preparative TLC on silica gel plates eluted with *n*-hexane-acetone (4:1), gave pure compound **3** (≈ 35 mg). Rechromatography of fraction F (≈ 18 g) on a silica gel column, using *n*-hexane-acetone (2:3) as eluant, afforded pure vanillic acid, which was crystallized from methanol as white needles (≈ 22 mg). A sample of solid, crude, methanol extract (≈ 30 g) was redissolved in methanol and a portion of this, which remained insoluble (≈ 5.5 g), proved to be a mixture of quercetin and isorhamnetin, which were identified by means of ^1H NMR analysis. The soluble portion was concentrated and subjected to chromatography on a Sephadex LH-20 column eluted with methanol. Twenty fractions were collected: Fraction 6, after concentration to dryness, gave a residue (≈ 65 mg) that was crystallized from methanol and identified by NMR as quercitrin (≈ 65 mg). In the same way, concentration of fractions 7 and 8 provided two different yellow solids, **6** (≈ 8 mg) and **7** (≈ 15 mg), respectively.

Acknowledgments - The authors are grateful to the Venezuelan Ministry of Popular Power for Science, Technology and Innovation (MCTI), "Science Mission" Program for financial support (Grant N° 2008000937). We also would like to thank Prof Angel G. Ravelo, Instituto Universitario de Bio-Orgánica, Universidad de La Laguna, Tenerife, Spain, for the EIMS.

References

- [1] Robson NKB. (2003). *Hypericum* botany. In *The Genus Hypericum*. Ernst E. (Ed.). Taylor and Francis, New York, pp 1-22.
- [2] Crockett S, Eberhardt M, Kunert O, Schühly W. (2010) *Hypericum* species in the Páramos of Central and South America: a special focus upon *H. irazuense* Kuntze ex N. Robson. *Phytochemistry Reviews*, **9**, 255-269.
- [3] Fariñas MR, Lázaro N, Monasterio M. (2008) Ecología comparada de *Hypericum laricifolium* Juss. y de *H. juniperinum* Kunth en el valle fluvio-glacial del páramo de Mucubají, Mérida, Venezuela. *Ecotropicos*, **21**, 75-88.
- [4] Antúnez de Mayolo KK. (1989) Peruvian natural dye plants. *Economic Botany*, **43**, 181-191.
- [5] Aguilar Z, Ulloa C, Hidalgo P. (2009) *Guía de Plantas Útiles de los Páramos de Zuleta, Ecuador*. Proyecto de Manejo y Aprovechamiento Sustentable de Alpacas en los Páramos de Zuleta. PPA-EcoCiencia. Quito.
- [6] Bussmann RW, Malca-García G, Glenn A, Sharon D, Chait G, Díaz D, Pourmand K, Jonat B, Somogy S, Guardado G, Aguirre C, Chan R, Meyer K, Kuhlman A, Townesmith A, Effio-Carbajal J, Frías-Fernández F, Benito M. (2010) Minimum inhibitory concentrations of medicinal plants used in Northern Peru as antibacterial remedies. *Journal of Ethnopharmacology*, **132**, 101-108.
- [7] Ansaloni R, Wilches I, León F, Orellana A, Peñaherrera E, Tobar V, de Witte P. (2010) Estudio preliminar sobre plantas medicinales utilizadas en algunas comunidades de las provincias de Azuay, Cañar y Loja, para afecciones del aparato gastrointestinal. *Revista Tecnológica ESPOL-RTE*, **23**, 89-97.
- [8] Bussmann RW, Glenn A, Sharon D. (2010) Healing the body and soul: traditional remedies for "magical" ailments, nervous system and psychosomatic disorders in Northern Peru. *African Journal of Pharmacy and Pharmacology*, **4**, 580-629.
- [9] El-Seedi HR, Ringbom T, Torsell K, Bohlin L. (2003) Constituents of *Hypericum laricifolium* and their cyclooxygenase (COX) enzyme activities. *Chemical & Pharmaceutical Bulletin*, **51**, 1439-1440.
- [10] (a) Gunatilanka AAL, de Silva AMYJ, Sotheeswaran S. (1982) Studies on medicinal and related plants of Sri Lanka. Part 6. Minor xanthenes of *Hypericum mysorensense*. *Phytochemistry*, **21**, 1751-1753; (b) Guo Ch, Zheng Q-M, Zheng H-Ch. (2005) The chemical constituents of *Hypericum sampsonii*. *Pharmaceutical Care and Research*, **5**, 341-344; (c) Chae S, Lee SY, Kim JS, Bae K, Kim SK, Kang SS. (2006) Constituents from *Hypericum ascyron*. *Korean Journal of Pharmacognosy*, **37**, 162-168.
- [11] (a) Wu Q-L, Wang Sh-P, Du L-J, Yang J-Sh, Xiao P-G. (1998) Xanthenes from *Hypericum japonicum* and *H. henryi*. *Phytochemistry*, **49**, 1395-1402; (b) Hong D, Yin F, Hu L-H, Lu P. (2004) Sulfonated xanthenes from *Hypericum sampsonii*. *Phytochemistry*, **65**, 2595-2598.
- [12] Rath G, Potterat O, Mavi S, Hostettmann K. (1996) Xanthenes from *Hypericum roeperianum*. *Phytochemistry*, **43**, 513-520.
- [13] Fu P, Li T, Liu R, Zhang W, Chen H. (2004) Xanthenes from the whole plant of *Hypericum japonicum*. *Natural Product Research Development*, **16**, 511-513.
- [14] Tao S, Wu F. (2004) Studies on chemical constituents of *Hypericum wightianum*. *Natural Product Research Development*, **16**, 26-27.
- [15] Zofou D, Kowa TK, Wabo HK, Ngemenya MN, Tane P, Titanji VPK. (2011) *Hypericum lanceolatum* (Hypericaceae) as a potential source of new anti-malarial agents: a bioassay-guided fractionation of the stem bark. *Malaria Journal*, **10**, 167-173.
- [16] Abreu IN, Porto ALM, Marsaioli AJ, Mazzafera P. (2004) Distribution of bioactive substances from *Hypericum brasiliense* during plant growth. *Plant Science*, **167**, 949-954.
- [17] (a) Schulz HU, Schurer M, Bassler D, Weiser D. (2005) Investigation of the bioavailability of hypericin, pseudohypericin, hyperforin and the flavonoids quercetin and isorhamnetin following single and multiple oral dosing of a *Hypericum* extract containing tablet. *Arzneimittelforschung*, **55**, 15-22; (b) Appleton J. (2010) Evaluating the bioavailability of isoquercetin. *Natural Medicine Journal*, **2**, 1-6.
- [18] Genoux-Bastide E, Lorendeau D, Nicolle E, Yahiaoui S, Magnard S, Di Pietro A, Baubichon-Cortay H, Boumendjel A. (2011) Identification of Xanthenes as selective killers of cancer cells overexpressing the ABC transporter MRP1. *Chem Med Chem*, **6**, 1478-1484.
- [19] (a) Iinuma M, Tosa H, Ito T, Tanaka T, Aqil M. (1995) Two prenylated anthrones in *Harungana madagascariensis*. *Phytochemistry*, **40**, 267-270; (b) Hu L-H, Yip S-Ch, Sim K-Y. (1999) Xanthenes from *Hypericum ascyron*. *Phytochemistry*, **52**, 1371-1373.

A New Xanthone from the Pericarp of *Garcinia mangostana*Manqin Fu^a, Samuel X. Qiu^b, Yujuan Xu^a, Jijun Wu^a, Yulong Chen^a, Yuanshan Yu^a and Gengsheng Xiao^{a,*}^aGuangdong Open Access Laboratory of Agri-food Processing, Sericulture and Agri-food Research Institute, Guangdong Academy of Agricultural Sciences, Guangzhou 510610, P.R. China^bKey Laboratory of Plant Resources Conservation and Sustainable Utilization, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, P.R. China

fumanqin84@126.com


Received: August 8th, 2013; Accepted: September 21st, 2013

A new prenylxanthone, garcimangostanol (**1**), was isolated from the EtOAc-soluble partition of the ethanol extract of the pericarp of *Garcinia mangostana* L., along with three known compounds, namely 8-deoxygartanin (**2**), 1-isomangostin (**3**), and garcinone C (**4**). The structure of compound **1** was elucidated on the basis of its 1D, 2D NMR and MS data. Compounds **1-4** exhibited either significant or moderate cytotoxicity against MCF-7, A549, Hep-G2 and CNE human cancer cell lines *in vitro* with IC₅₀ values from 4.0 ± 0.3 to 23.6 ± 1.5 μM by MTT colorimetric assay.

Keywords: *Garcinia mangostana*, Xanthone, Cytotoxicity.

Garcinia mangostana L. (family Clusiaceae), commonly known as mangosteen, is widely distributed in Indonesia, Thailand, Myanmar and some other Southeast Asian countries. Its pericarp has long been used as anti-inflammatory agent, astringent and indigenous medicine for treatment of skin infections, wounds and diarrhea in Southeast Asia [1]. Earlier chemical studies on mangosteen pericarp revealed a series of xanthone derivatives, which showed antioxidant, anti-inflammatory, antifungal, and antitumor activities [2]. In this article, we report the isolation and structural elucidation of a new prenylated xanthone (**1**), along with three known xanthones: 8-deoxygartanin (**2**) [3a], 1-isomangostin (**3**) [3b], and garcinone C (**4**) [3c] from the EtOAc-soluble partition of the ethanol extract of the pericarp of *G. mangostana*. In addition, the cytotoxicity is also described of compounds **1-4** against human breast carcinoma (MCF-7), human lung cancer (A549), human hepatoma (Hep-G2) and human carcinoma of nasopharynx (CNE) cell lines, tested by MTT colorimetric assay [4].

Garcimangostanol (**1**), a yellow amorphous powder, has the molecular formula C₁₉H₁₈O₆, as deduced from the HREI mass spectrum (*m/z* 342.1099 [M]⁺) and ESIMS (*m/z* 343 [M + H]⁺), as well as the NMR spectral data (Table 1). The ¹H NMR spectrum indicated the characteristic signal pattern of a prenyl moiety at δ 1.64 (3H, s, H-4'), 1.78 (3H, s, H-5'), 3.34 (2H, d, *J* = 7.6 Hz, H-1'), and 5.22 (1H, m, H-2'), which was further supported by the ¹³C NMR spectral signals at δ 17.8 (C-4'), 21.8 (C-1'), 25.8 (C-5'), 123.1 (C-2'), and 131.7 (C-3'). The ¹H NMR spectrum also revealed an intramolecularly hydrogen-bonded hydroxyl group at δ 13.31 (1H, s, OH-1), an aromatic proton at δ 6.60 (1H, s, H-4), two sets of coupled doublets at δ 7.01 (1H, d, *J* = 8.8 Hz, H-7) and 7.65 (1H, d, *J* = 8.8 Hz, H-8), and a methoxyl group [δ 4.00 (3H, s, 3-OCH₃)]. The ¹H and ¹³C NMR chemical shifts of **1** were closely similar to those of dulxanthone-A [5], except that an aromatic proton at C-2 and a prenyl group at C-4 in dulxanthone-A were replaced by an aromatic proton at C-4 and a prenyl group at C-2 in compound **1**. This structure assignment for **1** was confirmed by its HSQC and HMBC spectra (Figure 1b). The HMBC correlations from H-4 to C-4a (δ 157.0), C-9a (δ 103.3), C-2 (δ 112.0), and C-3 (δ 164.9), from H-1' (δ 3.34) to C-2', C-3', C-1 (δ 160.3), C-2, and C-3, and from 1-OH (δ 13.31) to C-1, C-2, and C-9a were indicative of the

**Figure 1:** (a) Structures of compounds **1-4**; (b) Key HMBC correlations of compound **1**.**Table 1:** ¹H and ¹³C NMR spectroscopic data of compound **1** in acetone-*d*₆

position	δ _H	δ _C
1		160.3
2		112.0
3		164.9
4	6.60(s)	90.7
5		133.3
6		152.1
7	7.01(d, <i>J</i> = 8.8)	113.6
8	7.65(d, <i>J</i> = 8.8)	117.3
9		181.3
4 ^a		157.0
8 ^a		114.8
9 ^a		103.3
10 ^a		147.0
1'	3.34(d, <i>J</i> = 7.6)	21.8
2'	5.21-5.23(m)	123.1
3'		131.7
4'	1.64(s)	17.8
5'	1.78(s)	25.8
3-OMe	4.00(s)	56.6
1-OH	13.31(s)	

Table 2: Cytotoxicity (IC₅₀, μM) of compounds **1-4** against tumor cell lines^a

Compounds	MCF-7	A549	Hep-G2	CNE
1	14.1 ± 1.1	11.9 ± 0.8	13.5 ± 1.0	4.2 ± 0.2
2	16.8 ± 1.4	15.1 ± 1.2	12.7 ± 0.5	4.0 ± 0.3
3	18.6 ± 1.6	23.6 ± 1.5	15.6 ± 1.2	5.7 ± 0.3
4	7.1 ± 0.5	4.9 ± 0.8	4.3 ± 0.1	6.6 ± 0.2
Doxorubicin ^a	3.4 ± 0.1	3.7 ± 0.2	3.5 ± 0.1	2.3 ± 0.1

^aEach value represents mean ± SD, and doxorubicin was used as a positive control.

connectivity of the aromatic proton at C-4 and the prenyl group at C-2. Thus, the structure of **1** was determined as 1,5,6-trihydroxy-2-(3-methylbut-2-enyl)-3-methoxyxanthone.

Compounds **1-4** were tested for their cytotoxicity against MCF-7, A549, Hep-G2 and CNE cell lines by MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) colorimetric assay, in which

doxorubicin was used as a positive control. The IC_{50} values of compounds **1-4** on the viability of the test cancer cell lines after 72 h of incubation are presented in Table 2. The results showed that compounds **1-3** were moderately cytotoxic against MCF-7, A549, and Hep-G2 cell lines with IC_{50} values from 11.9 ± 0.8 to 23.6 ± 1.5 μ M, and significantly cytotoxic against CNE cells with IC_{50} values of 4.2 ± 0.2 , 4.0 ± 0.3 , and 5.7 ± 0.3 μ M, respectively. Moreover, compound **4** showed strong cytotoxicity against all the tested cell lines with IC_{50} values from 4.3 ± 0.1 to 7.1 ± 0.5 μ M.

Some structural features apparently contribute to the potent inhibitory activity of xanthenes, such as isopentenyl cyclization lowered the inhibitory activity (compound **3**), and the hydrated form, 3-hydroxy-3-dimethyl butanyl (compound **4**), enhanced the inhibitory activity. The remarkable cancer cell anti-proliferative activities of compounds **1-4** has shed some light on the active ingredients and action mechanism supportive of the beneficial properties and rationale for its use as a folk medicine.

Experimental Section

General: The following instruments were used to obtain physical data: Bruker DRX-400 spectrometer (400 MHz for 1H NMR, 100 MHz for ^{13}C NMR); Perkin-Elmer Lambda 35 UV-vis, WQF-410 FT-IR spectrophotometer; MDS SCIEX API 2000 LC/MS/MS, MAT95XP mass spectrometer; XT-4 Boettius micromelting point apparatus; CC, RP-C₁₈, Sephadex LH-20; TLC: pre-coated silica gel GF254; Doxorubicin and MTT were purchased from Sigma Chemical Co. (St. Louis, MO).

Plant material: The pericarp of *G. mangostana* was obtained by peeling the fruits bought from a local grocery store in Fengyang, China, which were imported from Thailand in August 2012. The plant material was identified by Prof. Yulian Wang (Anhui Science and Technology University, China). A voucher sample was deposited at the herbarium of Anhui Science and Technology University.

Extraction and isolation: The powdered dry pericarps of *G. mangostana* (2 kg) were extracted 3 times with 95% ethanol (5 L \times 3) at room temperature, for 3 days each. After evaporation of the solvent *in vacuo*, the combined crude ethanolic extract (120 g) was partitioned with EtOAc (2 L \times 3) to afford the EtOAc (50 g) extract. This was subjected to a silica gel CC and eluted with light petroleum-acetone with increasing polarities (10:1-1:1) to yield 8 fractions (1-8). Fraction 3 (4.5 g) was subjected to silica gel CC

using light petroleum-EtOAc (8:1-1:1) to afford **1** (15 mg). Fraction 4 (2.4 g) was subjected to Sephadex LH-20 CC and eluted with MeOH to obtain **2** (25 mg). Fraction 5 (3.5 g) was separated by ODS CC eluting with 80% MeOH, followed by silica gel using light petroleum-acetone (90:10-50:50) to yield **3** (12 mg) and **4** (30 mg).

Garcimangostanol (1)

Yellow amorphous powder.

MP: 183 - 185°C.

UV(MeOH): λ_{max} (log ϵ): 254 (4.31), 284 (3.96), 328 (4.12) nm.

IR (KBr): 3545, 1645, 1587 cm^{-1} .

1H NMR and ^{13}C NMR: Table 1.

MS ((+)-ESI): m/z = 343 $[M+H]^+$.

HR-EI-MS: 342.1099 (100) (calcd. 342.1098 for C₁₉H₁₈O₆, $[M]^+$).

Cell culture: MCF-7, A549, Hep-G2, and CNE cell lines were cultured as monolayers in nutrient RPMI 1640 medium supplemented with streptomycin (100 μ g mL⁻¹), penicillin (100 U mL⁻¹), and 10% heat inactivated (56 °C) fetal bovine serum (FBS) at 37°C in a 5% CO₂ humidified incubator.

Cytotoxicity assay: Cells were plated at 1×10^4 cell/well in 96-well microtiter plates and allowed to adhere overnight. Stock solutions of compounds (10 mM) were dissolved in dimethylsulfoxide (DMSO). The cells were treated in triplicate with various concentrations (50, 25, 12.5, 6.25, 3.125, 0 μ M) of test samples, and doxorubicin as the positive control for 48 h. At the end of the incubation time, 20 μ L MTT reagent (5 mg/mL) was added to each well for 4 h, and the resulting crystals were dissolved in DMSO. Absorbance (A) was measured in an ELISA plate reader (TECAN, Austria) at a wavelength of 570 nm. Data were calculated as a percentage of inhibition using the following formula: Inhibitory ratio (%) = $(A_{control} - A_{sample}) / (A_{control} - A_{blank}) \times 100\%$. A_{sample} , $A_{control}$ and A_{blank} refer to the absorbance of the sample, control (cells grown only in nutrient medium) and blank (no cells), respectively. IC_{50} concentration was defined as the concentration of an agent inhibiting cell survival by 50%, compared with a vehicle-treated control.

Acknowledgement - This work was jointly supported by the National Science and Technology Support Program (No. 2012BAD31B03) and Guangdong Science and Technology Program (No. 2010B080100026).

References

- [1] (a) Chairungsilerd N, Takeuchi K, Ohizumi Y, Nozoe S, Ohta T. (1996) Mangostanol, a prenylxanthone from *Garcinia mangostana*. *Phytochemistry*, **43**, 1009-1102; (b) Suksamrarn S, Suwannapoch N, Ratananukul P, Aroonlerk N, Suksamrarn A. (2002) Xanthenes from the green fruit hulls of *Garcinia mangostana*. *Journal of Natural Products*, **65**, 761-763; (c) Zhang Y, Song Z, Hao J, Qiu S, Xu Z. (2010) Two new prenylated xanthenes and a new prenylated tetrahydroxanthone from the pericarp of *Garcinia mangostana*. *Fitoterapia*, **81**, 595-599.
- [2] (a) Jung HA, Su BN, Keller WJ, Mehta RG, Kinghorn AD. (2006) Antioxidant xanthenes from the pericarp of *Garcinia mangostana* (Mangosteen). *Journal of Agricultural and Food Chemistry*, **54**, 2077-2082; (b) Gopalakrishnan G, Balaganesan B. (2000) Two novel xanthenes from *Garcinia mangostana*. *Fitoterapia*, **71**, 607-609; (c) Chin YW, Shin E, Hwang BY, Lee MK. (2011) Antifibrotic constituents from *Garcinia mangostana*. *Natural Product Communications*, **6**, 1267-1268.
- [3] (a) Nguyen LHD, Vo HT, Pham HD, Connolly JD, Harrison LJ. (2003) Xanthenes from the bark of *Garcinia merguensis*. *Phytochemistry*, **63**, 467-470; (b) Mahabusarakam W, Wiriyaichitra P, Taylor WC. (1987) Chemical constituents of *Garcinia mangostana*. *Journal of Natural Products*, **50**, 474-478; (c) Sen AK, Sarkar KK, Mazumder PC, Banerji N, Uusvuori R, Hase TA. (1982) The structures of garcinones a, b and c: three new xanthenes from *Garcinia mangostana*. *Phytochemistry*, **21**, 1747-1750.
- [4] Mosmann T. (1983) Rapid colorimetric assay for cellular growth and survival application to proliferation and cytotoxicity assays. *Journal of Immunological Methods*, **65**, 55-63.
- [5] Ito C, Miyamoto Y, Nakayama M, Kawai Y, Rao KS, Furukawa H. (1997) A novel depsidone and some new xanthenes from *Garcinia* species. *Chemical & Pharmaceutical Bulletin*, **45**, 1403-1413.

Isolation of a Phomoxanthone A Derivative, a New Metabolite of Tetrahydroxanthone, from a *Phomopsis* sp. Isolated from the Mangrove, *Rhizophora mucronata*

Yoshihito Shiono^a, Takehiro Sasaki^a, Fumiaki Shibuya^a, Yukito Yasuda^a, Takuya Koseki^a and Unang Supratman^b

^aDepartment of Food, Life, and Environmental Science, Faculty of Agriculture, Yamagata University, Tsuruoka, Yamagata 997-8555, Japan

^bDepartment of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Padjadjaran, Jatinangor 45363, Indonesia

yshiono@tds1.tr.yamagata-u.ac.jp

Received: July 16th, 2013; Accepted: September 14th, 2013

From the rice culture of *Phomopsis* sp. IM 41-1, isolated from the mangrove plant, *Rhizophora mucronata*, two dimeric tetrahydroxanthones, phomoxanthone A (**1**) and 12-*O*-deacetyl-phomoxanthone A (**2**) were obtained. Their structures were determined by spectroscopic and spectrometric methods. Both compounds exhibited antimicrobial activities.

Keywords: Endophyte, Mangrove, *Phomopsis* sp., Phomoxanthone, Antimicrobial activity.

The interaction of endophytes, fungi that exist in a symbiotic relationship within the intracellular space of plants without causing apparent harm to their host plants, has received considerable attention. Endophytes produce bioactive substances that are known to enhance growth of the host plant and provide protection against pathogens and herbivores. These fungi have proven to be a promising source for the production of structurally novel and pharmacologically active secondary metabolites [1]. Our previous investigation of endophytic fungi isolated from plants in Japan resulted in the isolation of some bioactive and structurally unique metabolites [2]. In our continuing efforts to discover biologically active compounds among the secondary metabolites of endophytes, we have investigated the fungal strain *Phomopsis* sp. IM 41-1 isolated from *Rhizophora mucronata*, a mangrove plant collected from a forest in Muara Angke, Jakarta, Indonesia. *R. mucronata* (local name: Bakau Hitam) has found use in folk medicine for treatment of diarrhea, dysentery, fever, malaria, and leprosy [3]. Examination of the rice culture of *Phomopsis* sp. IM 41-1 resulted in the isolation of two dimeric xanthone-type compounds, **1** and **2**. In this report, we describe the characteristics and fermentation products of the producing strain, as well as the isolation, structure elucidation and biological characterization of the two compounds.

Steamed unpolished rice was used as the substrate for the cultivation of the endophyte, *Phomopsis* sp. IM 41-1. The culture medium was extracted with methanol. The extract was concentrated, and the aqueous residue extracted with ethyl acetate. The EtOAc layer was subjected to silica gel column chromatography eluting with a *n*-hexane–EtOAc gradient. Purification of the compounds was guided by their characteristic coloration on TLC. Further chromatographic studies yielded two pure compounds: phomoxanthone A (**1**) [4] and a new phomoxanthone A derivative (**2**).

Compound **2** was obtained as a yellow powder, and its molecular formula was determined to be C₃₆H₃₆O₁₅ by HRESITOFMS. The IR




Figure 1: Structures of compounds **1** and **2**.

spectrum of **2** exhibited absorptions due to hydroxyl, carbonyl, and aromatic functions, whereas the UV spectrum showed the characteristic absorption pattern of a phomoxanthone derivative (λ_{max} 340 and 221 nm). The NMR spectrum of **2** also showed two sets of signals at the same position that were either completely overlapped or separated. The ¹³C NMR and DEPT experiments revealed the presence of 36 carbons: four methylene, eight methine, five methyl, and 19 quaternary (sp² carbons also included three esters and two carbonyls). The ¹H NMR spectral data revealed the presence of five methyl groups [δ_{H} 1.07 (total 6H, d, J = 6.0 Hz, Me-11, 11') and 2.10 (total 9H, s, Me-14, 14' and 16')], of which three were singlets and two doublets; four aromatic protons [δ_{H} 6.47 (1H, d, J = 8.0 Hz, H-2), 6.51 (1H, d, J = 8.0 Hz, H-2'), 7.40 (1H, d, J = 8.0 Hz, H-3), 7.41 (1H, d, J = 8.0 Hz, H-3')], two oxymethylenes [δ_{H} 3.55 (1H, d, J = 12.2 Hz, H-12), 4.10 (1H, d, J = 12.2 Hz, H-12), 4.18 (1H, d, J = 12.5 Hz, H-12'), 4.50 (1H, d, J = 12.5 Hz, H-12)], two oxymethines [δ_{H} 5.56 (1H, s, H-5) and 5.75 (1H, s, H-5')] and two exchangeable downfield protons [δ_{H} 11.8 (1H, s, OH-1), 11.9 (1H, s, OH-1'), 13.1 (1H, s, OH-8) and 14.1 (1H, s, OH-8')]. These signals were assigned based on the COSY spectrum and compared with the ¹H NMR signals of **1**. These results suggested that **2** was a nonsymmetrical dimer similar to **1**. Analysis of the 1D and 2D NMR data of **2** suggested that it

Analysis of the 1D and 2D NMR data of **2** suggested that it consisted of one monomer identical to that of **1**, but with the other one lacking the substitution pattern of the acetoxy moiety. The plane structure of **2** was determined as shown in Fig. 2 by HMBC analysis. NOE experiments (Fig. 2) exhibited key correlations from H₃-11 to H-5, from H-12 to H-6, from H₃-11' to H-5' and from H-12' to H-6', implying the same relative configuration as **1**. Finally, the ¹H and ¹³C NMR spectra of **1** corresponded with those of the acetylated product obtained by acetylation of **2** with Ac₂O/pyridine, providing further evidence for the structure of **2**. The relative configuration at C-5, C-6, and C-10a in the acetylation product of **2** was considered to be the same as that in **1**, because protons and carbons at C-5, C-6, C-7, and C-12 in the acetylation product of **2** had chemical shifts and coupling constants quite similar to those of **1**. Based on these results, the structure assigned to **2** was 12-*O*-deacetyl-phomoxanthone A (**2**) (Fig. 1). In addition, the possibility of **2** being an artifact of **1** produced during the separation could be excluded since **1** was stable for one week in MeOH and silica gel.




Figure 2: Selected HMBC and NOE correlations observed for **2**.

The activities of compounds **1** and **2** against Gram-positive and Gram-negative bacteria and fungal strains were evaluated using the agar diffusion method (Table 1). Both compounds exhibited moderate activity against *Botrytis cinerea*, *Sclerotinia sclerotiorum*, *Diaporthe medusae*, and *Staphylococcus aureus* at a concentration of 30 µg/disk. Both compounds lacked activity against the Gram-negative bacterium *Pseudomonas aeruginosa*. Although the synthetic products are very effective in the assay system, the antimicrobial activity of compounds **1** and **2** produced by *Phomopsis* sp. IM 41-1 could play an important role in the protection of the host plant from degradation and disease caused by pathogens.

Table 1: Antimicrobial activities of compounds **1** and **2**

Microorganisms	1 ^a	2 ^a	Control ^b
<i>Trichoderma harzianum</i> NBRC 33016	ND	ND	15
<i>Verticillium dahliae</i> Klebahn NBRC 9470	ND	ND	20
<i>Sclerotinia sclerotiorum</i> de Bary NBRC 103652	13	12	15
<i>Diaporthe medusae</i> Nitschke NBRC 30895	11	ND	10
<i>Botrytis cinerea</i> Persoon NBRC 100717	11	11	20
<i>Staphylococcus aureus</i> NBRC 13276	9	9	32
<i>Pseudomonas aeruginosa</i> ATCC 15442	ND	ND	18

^aDiameter of the inhibition areas (mm) using the plate diffusion assay (30 µg of each tested compound soaked in a 8 mm filter disk). ^bThe positive control, thiabendazole was used for *T. harzianum*, *V. dahliae*, *S. sclerotiorum*, *D. medusae*, and *B. cinerea*, penicillin G (1.3 µg/disk) for *S. aureus*, and chloramphenicol (10 µg/disk) for *P. aeruginosa*. ND: not detectable.

Previous published reports have investigated biological activities of these compound types, including antimicrobial, antimalarial,

antitubercular and cytotoxic activities. Although phomoxanthones A and B exhibited some activity, *O*-5,12,5',12'-tetradecetyl-phomoxanthone A did not show any in these tests [4, 5]. These previous reports suggested that the acetylated moieties had a significant effect on biological activities, whereas the activity of **1** was linked to its lipophilicity. In the present study, the differences in the antimicrobial activities between **1** and **2** were small. Further pharmacological studies of **1** and **2** are currently in progress to better reveal the effect of the acetylation.

Experimental

General: Optical rotation values were measured with a Horiba SEPA-300 polarimeter, and IR and UV spectra were respectively recorded with Jasco J-20A, Shimadzu UV mini-1240, and Jasco J-20A spectrophotometers. Mass spectra were obtained with Jeol JMS-700 and Synapt G2 mass spectrometers. NMR data were recorded on a Jeol EX-400 spectrometer at 400 MHz for ¹H and 100 MHz for ¹³C. Chemical shifts are given on a δ (ppm) scale with TMS as an internal standard. ¹H, ¹³C, DEPT, COSY, HMQC and HMBC spectra were recorded using standard Jeol standard pulse sequences.

Isolation procedure: The fungal strain *Phomopsis* sp. IM 41-1 was cultivated on sterile steamed unpolished rice (total 1000 g, 20 g / Petri dish x 50) at 25°C for 4 weeks. The rice was extracted with MeOH (5 L), and the extract concentrated. The resulting aqueous concentrate (0.5 L) was partitioned with *n*-hexane (0.5 L), and EtOAc (1.5 L). Each fraction was concentrated to dryness to yield the *n*-hexane (600 mg), EtOAc (500 mg) and aqueous soluble residues (1.5 g). Purification of the EtOAc soluble fraction was monitored on TLC plates by the characteristic intense blue coloration with 10% vanillin in H₂SO₄. The EtOAc soluble fraction was chromatographed on a silica gel column using stepwise elution with *n*-hexane: EtOAc (100:0:100) and EtOAc: MeOH (50:50:100), respectively, to afford fractions 1-1 to 1-13. Fractions 1-3 and 1-4 were combined and further chromatographed on a silica gel column using CHCl₃: EtOAc (80:20) to yield phomoxanthone A (**1**, 30.0 mg) and 12-*O*-deacetyl-phomoxanthone A (**2**, 19.0 mg).

12-*O*-Deacetyl-phomoxanthone A (**2**)

Yellow amorphous powder.

[α]_D²⁰: +40.0 (c 0.65, CHCl₃).

UV/Vis (MeOH) nm (log ε): 340 (4.2), 221 (4.0).

IR (KBr): 3421, 2927, 1747, 1608, 1437, 1373, 1230, 1049, 883 cm⁻¹.

¹³C NMR (100 MHz, CDCl₃): δ_C 17.5 (Me-11)^a, 17.6 (Me-11')^a, 20.7 (C-14)^b, 20.8 (C-14')^b, 20.9 (C-16)^b, 27.6 (C-6), 27.7 (C-6'), 33.3 (C-7, 7'), 65.2 (C-12), 65.5 (C-12'), 70.2 (C-5), 70.4 (C-5'), 80.4 (C-10a')^c, 82.4 (C-10a)^c, 100.3 (C-8a)^d, 100.8 (C-8a')^d, 106.3 (C-9a)^e, 106.4 (s, C-9a')^e, 107.9 (C-2), 108.2 (C-2'), 117.7 (C-4), 117.9 (C-4'), 140.1 (C-3, 3'), 157.1 (C-4a)^f, 157.4 (C-4a')^f, 159.3 (C-1), 159.5 (C-1'), 170.3 (C-13)^g, 170.4 (C-13')^g, 170.6 (C-15')^g, 177.8 (C-8), 177.9 (C-8') and 187.8 (s, C-9, 9').

¹H NMR (400 MHz, CDCl₃): δ_H 1.07 (total 6H, d, *J* = 6.0 Hz, Me-11, 11'), 2.10 (total 9H, s, Me-14, 14', 16'), 2.36-2.47 (overlapped signals, H-6, 6', 7, 7'), 3.55 (1H, m, H-12), 4.10 (1H, d, *J* = 12.2 Hz, H-12), 4.18 (1H, d, *J* = 12.5 Hz, H-12'), 4.50 (1H, d, *J* = 12.5 Hz, H-12'), 5.56 (1H, s, H-5'), 5.75 (1H, s, H-5), 6.47 (1H, d, *J* = 8.0 Hz, H-2), 6.51 (1H, d, *J* = 8.0 Hz, H-2'), 7.40 (1H, d, *J* = 8.0 Hz, H-3), 7.41 (1H, d, *J* = 8.0 Hz, H-3'), 11.8 (1H, s, OH-1), 11.9 (1H, s, OH-1'), 14.0 (1H, s, OH-8) and 14.1 (1H, s, OH-8').

^{a-g} Assignments may be interchangeable.

Acetylation of **2:** 12-*O*-Deacetyl-phomoxanthone A (**2**, 5 mg) in pyridine (2 mL) was acetylated with acetic anhydride (2 mL) at

water and extracted with EtOAc (5 mL x 3). The organic layer was washed with saturated NaCl and dried over Na₂SO₄ and concentrated *in vacuo* to give a residue, which was purified by silica gel CC to yield the acetate (4 mg) as an amorphous substance. The ¹H and ¹³C NMR data of this compound were identical to those obtained for natural phomoxanthone A (**1**).

¹H NMR (400 MHz, CDCl₃): δ_H 1.90 (6H, s, Me-16, 16'), 1.00 (6H, d, *J* = 5.6 Hz, Me-11, 11'), 2.05 (6H, s, Me-14, 14'), 2.34 (4H, m, H-6, 6', 7, 7'), 2.45 (2H, m, H-7, 7'), 4.17 (2H, d, *J* = 12.7 Hz, H-12, 12'), 4.26 (2H, d, *J* = 12.7 Hz, H-12, 12'), 5.38 (2H, s, H-5, 5'), 6.57 (2H, d, *J* = 8.6 Hz, H-2), 7.34 (2H, d, *J* = 8.6 Hz, H-3, 3'), 11.52 (2H, s, OH-1, 1'), 14.10 (2H, s, OH-8, 8').

¹³C NMR (100 MHz, CDCl₃): δ_C 17.5 (C-11, 11'), 20.4 (C-16, 16'), 20.6 (C-14, 14'), 27.8 (C-6, 6'), 33.3 (C-7, 7'), 64.5 (C-12, 12'), 70.3 (C-5, 5'), 80.3 (C-10, 10'), 100.1 (C-8a, 8a'), 106.0 (C-9a,

9a'), 109.3 (C-2, 2'), 115.3 (C-4, 4'), 141.2 (C-3, 3'), 153.5 (C-4a, 4a'), 161.5 (C-1, 1'), 169.6 (C-13, 13'), 170.2 (C-15, 15'), 177.4 (C-8, 8'), 187.7 (C-9, 9').

HR-ESI-TOF-MS: *m/z* [M+Na]⁺ calcd for C₃₈H₃₈O₁₆Na⁺: 773.2058; found: 773.2040.

Antimicrobial activity: The test microorganisms were *Pseudomonas tolaasii* NBRC 103163, *Trichoderma harzianum* NBRC 33016, *Verticillium dahliae* Klebahn NBRC 9470, *Botrytis cinerea* Persoon NBRC 100717, *Sclerotinia sclerotiorum* (Libert) de Bary NBRC 103652, *Diaporthe medusaea* Nitschke NBRC 30895, *Staphylococcus aureus* NBRC 13276 and *Pseudomonas aeruginosa* ATCC 15442. Antimicrobial assays were carried out by the paper disk diffusion method using a published protocol [6].

References

- [1] Gunatilaka AAL. (2006) Natural products from plant-associated microorganisms: Distribution, structural diversity, bioactivity, and implications of their occurrence. *Journal of Natural Products*, **69**, 509-526.
- [2] Shiono Y, Kosukegawa A, Koseki T, Murayama T, Kwon E, Uesugi S, Kimura K. (2012) A dimeric pyrrocidine from *Neonectria ramulariae* is an inhibitor of prolyl oligopeptidase. *Phytochemistry Letters*, **5**, 91-95.
- [3] Duke NC, Pinzon ZS. (1992) The genus *Rhizophora* seedling from leaf scar nodes: a technique for studying recruitment and growth in mangrove forest. *Biotropica*, **24**, 173-186.
- [4] Isaka M, Jaturapat A, Rukseree K, Danwisetkanjana K, Tanticharoen M, Thebtaranonth Y. (2001) Phomoxanthones A and B, novel xanthone dimers from the endophytic fungus *Phomopsis* species. *Journal of Natural Products*, **64**, 1015-1018.
- [5] Rukachaisirikul V, Sommart U, Phongpaichit S, Sakayaroj J, Kirtikara K. (2008) Metabolites from the endophytic fungus *Phomopsis* sp. PSU-D15. *Phytochemistry*, **69**, 783-787.
- [6] Shiono Y, Murayama T, Takahashi K, Okada K, Katohda S, Ikeda M. (2005) Three oxygenated cyclohexenone derivatives, produced by an endophytic fungus. *Bioscience, Biotechnology and Biochemistry*, **69**, 287-292.

Anti-allergic Inflammatory Effects of Cyanogenic and Phenolic Glycosides from the Seed of *Prunus persica*

Geum Jin Kim^a, Hyun Gyu Choi^a, Ji Hyang Kim^a, Sang Hyun Kim^b, Jeong Ah Kim^{c*} and Seung Ho Lee^{a*}

^aCollege of Pharmacy, Yeungnam University, Gyongsan, Gyungbuk, 712-749 Korea

^bSchool of Medicine, Kyungpook National University, Daegu, 702-701 Korea

^cCollege of Pharmacy, Research Institute of Pharmaceutical Sciences, Kyungpook National University, Daegu, 702-701 Korea

jkim6923@knu.ac.kr (J.A.Kim), seungho@yu.ac.kr (S.H.Lee)

Received: May 6th, 2013; Accepted: September 21st, 2013

A methanol extract of the seed of *Prunus persica* (Rosaceae) was found to inhibit histamine release in human mast cells. Activity-guided fractionation of the methanol extract yielded three cyanogenic glycosides (**1-3**) and other phenolic compounds (**4-8**). To evaluate their anti-allergic and anti-inflammatory activities, the isolates (**1-8**) were tested for their inhibitory effects on histamine release and on the gene expressions of tumor necrosis factor (TNF)- α and interleukin (IL)-6 in human mast cells. Of these, phenolic glycosides **7** and **8** suppressed histamine release and inhibited the pro-inflammatory cytokines TNF- α and IL-6. These results suggest that isolates from *P. persica* are among the anti-allergic inflammatory principles in this medicinal plant.

Keywords: *Prunus persica*, Cyanogenic glycosides, Phenolic glycosides, Anti-allergic inflammatory activities.

Allergic inflammation affects target cells, such as epithelial, fibroblast, vascular, and airway smooth muscle cells, which become an important source of inflammatory mediators [1a]. Mast cells appear to play a critical role in initiating allergic inflammation, since they are directly activated by allergens via an immunoglobulin E (IgE)-dependent mechanism [1a,1b]. Activated mast cells lead to degranulated histamine, and release of pro-inflammatory and chemotactic cytokines [1b]. Several proinflammatory cytokines have been implicated in allergic diseases, including interleukin (IL)-1 β , IL-6, TNF (tumor necrosis factor)- α , and granulocyte-macrophage colony-stimulating factor (GM-CSF), and these may be important for amplifying allergic inflammatory responses [1a,1b].

Prunus persica (L.) Batsch (Rosaceae) is a deciduous tree that bears an edible fruit. Its seeds are used in traditional medicine to treat amenorrhea and rheumatoid arthritis [2a]. Phytochemical studies on this plant have reported various cyanogenic glycosides, glycerides, sterols, and fatty acids [2b-2d]. Furthermore, pharmacological studies on the seeds of *P. persica* have shown they have multiple activities, which include anti-cancer, anti-oxidant, hepatoprotective, antinociceptive, and anti-coagulant, and inhibitory effects on platelet aggregation [2d-2f]. In a previous study, we described the anti-allergic inflammatory effect of the ethanol extract of fruits of *P. persica* in human mast cells [3]. Activity-guided fractionation of the methanol extract led to the isolation of three cyanogenic glycosides, prunasin (**1**), amygdalin (**2**), neoamygdalin (**3**) and five phenolic compounds, dehydroconiferyl alcohol (**4**), dehydroconiferyl alcohol-4'-*O*- β -D-glucopyranoside (**5**), dehydroconiferyl alcohol-10-*O*- β -D-glucopyranoside (**6**), vanilloside (**7**) and lacticolorin (**8**). Compounds **5**, **7** and **8** were isolated for the first time from this plant.

It is known that the inhibitory effect of mast cells on histamine release plays an important role in the anti-allergic effect [1a]. Therefore, we examined *in vitro* histamine release inhibitory activity using assays of compound 48/80-induced histamine release from human mast cells and pro-inflammatory cytokines inhibitory activity. As shown in Figure 2, compared with the




Figure 1: Chemical structures of compounds **1-8** isolated from *Prunus persica*.




Figure 2: Anti-allergic inflammatory effects of compounds **1-8** isolated from *P. persica* on human mast cells. HMC-1 cells were pre-incubated with compounds **1-8** for 30 min prior to incubation with PMA (20 nM) and calcium ionophore A23187 (1 μ M). (A) Effect of compounds **1-8** on the expression of pro-inflammatory cytokines. Extraction and analysis of mRNA was performed as described in Experimental. (B) Effect of compounds **1-8** on histamine release in mast cells. Histamine content was measured using a spectrometer. Each bar represents the mean \pm SEM of three independent experiments. *Significant difference at $P < 0.05$.

phorbol-12-myristate 13-acetate plus calcium ionophore A23187 (PMACI) control, compounds **4**, **5**, **7** and **8** significantly inhibited histamine release and pro-inflammatory cytokines.

Experimental

Plant material: Seeds of *Prunus persica* were purchased from Yak-Ryung-si market in Daegu, Korea and were identified by Dr Seung Ho Lee at the College of Pharmacy, Yeungnam University. A voucher specimen (YU00163) has been deposited in the Natural Product Laboratory at the College of Pharmacy, Yeungnam University, Korea.

Extraction and isolation: The seeds of *P. persica* (1.2 kg) were pulverized and extracted with MeOH (5.0 L \times 3) for one week at room temperature. The solution was evaporated to dryness to obtain 85.0 g of extract. This was loaded onto a silica gel column and eluted with a step-wise gradient mixture of ethyl acetate (EtOAc) and MeOH (1:0 to 0:1) to yield 13 sub-fractions (Fr.3-1 ~ Fr.3-9). Fr.3 (678.7 mg) was further divided into 9 sub-fractions (Fr.1-1 ~ Fr.1-9) using silica gel chromatography and eluting with a step-wise gradient mixture of methylene chloride (MC) and acetone (100:0 to 40:60). Fr.3-5 (75.9 mg) was purified by preparative HPLC (MeOH-H₂O, 3:7 to 7:3; 6 mL/min) to afford (-) dehydroconiferyl alcohol (**4**) (t_R 29.8 min, 4.5 mg) [4a]. Fr.4 (8.5 g) was loaded onto silica gel and eluted with a mixture of MC and MeOH (100:0 to 85:15) to yield 6 sub-fractions (Fr.4-1 ~ Fr.4-6). Purification of Fr.4-2 (113.5 mg) using preparative HPLC (MeOH-H₂O, 40:60 to 65:35; 6 mL/min) yielded (-) lacticolorin (**8**) (t_R 37.4 min, 3.0 mg) [4f]. Fr.5 (2.1 g) was separated on silica gel using a gradient mixture of MC and MeOH (10:0 to 5:5) to afford 9 sub-fractions (Fr.5-1 ~ Fr.5-9). Fr.5-7 (139.9 mg) was further purified by preparative HPLC with a gradient mixture of MeOH and H₂O (from 1:9 to 5:5, 6 mL/min) to afford (+) neoamygdalin (**3**, t_R 23.2 min 69.4 mg) [4b]. Fr.5-8 (755.5 mg) was also purified by preparative HPLC (MeOH-H₂O, 3:7 to 4:6; 6 mL/min) to give (-) dehydroconiferyl alcohol-10-*O*- β -D-glucopyranoside (**6**, t_R 30.8 min, 1.2 mg) [4d]. Fr.6 (15.2 g) was subjected to silica gel vacuum liquid chromatography, and eluted with a gradient mixture of CHCl₃/MeOH/H₂O [9:1:0.05, 9:1.5:0.05, 9:2:0.1, 9:3:0.1, 0:10:0 (each 2 L)] to yield (-) amygdalin (**2**, 3.0 g) [4b] and 4 sub-

fractions (Fr.6-1 ~ Fr.6-4). Fr.6-3 was fractionated by silica gel chromatography using a gradient of CHCl₃ and MeOH (10:0 to 7:3) to yield (-) prunasin (**1**, 50.0 mg) [4b] and 7 sub-fractions (Fr.6-3-1 ~ Fr.6-3-7). Fr.6-3-7 (458.6 mg) was further purified by preparative HPLC (MeOH-H₂O, 1:9 to 6:4; 6 mL/min) to afford (-) vanilloside (**7**, t_R 27.9 min, 5.7 mg) [4e] and (-) dehydroconiferyl alcohol-4'-*O*- β -D-glucopyranoside (**5**, t_R 57.9 min, 21.2 mg) [4c].

Histamine assay: The histamine content was measured by the *o*-phthalaldehyde spectrofluorometric procedure, as previously described [1b]. The serum was separated from the released histamine by centrifugation at 400 \times g for 5 min at 4°C. The fluorescent intensity was measured at an excitation wavelength of 355 nm and an emission wavelength of 450 nm using a Perkin-Elmer Fluorescence Spectrometer LS-50B (Norwalk, CT, USA).

Real-time polymerase chain reaction (PCR): Quantitative real-time PCR was carried out using the Thermal Cycler Dice TP850 (Takarabio Inc., Shiga, Japan) according to the manufacturer's protocol. Total RNA was isolated from ear tissues of each group. The conditions for PCR were similar to those previously described. Briefly, 2 μ L of cDNA (100 ng), 1 μ L of sense and antisense primer solution (0.4 μ M), 12.5 μ L of SYBR Premix Ex Taq (Takarabio Inc.), and 9.5 μ L of D₂O were mixed together to obtain a final 25 μ L reaction mixture in each reaction tube. The primers used for PCR were (IL-6 F 5'-AAAGAGGCACTGGCAGAAAA, R 5'-ATCTGAGGTGCCCATGCTAC, TNF- α F 5'-CCTGTAGCCCACGTCGTAGC, R 5'-TTGACCTCAGCGCTGAGTTG). The amplification conditions were 10 s at 95°C, 40 cycles of 5 s at 95°C and 30 s at 60°C, 15 s at 95°C, 30 s at 60°C, and 15 s at 95°C. The normalization and quantification of mRNA expression was performed using the TP850 software supplied by the manufacturer.

Statistical analysis: Statistical analyses were performed using SAS statistical software (SAS Institute, Cary, NC, USA). Treatment effects were analyzed using ANOVA followed by Dunnett's multiple range tests. Significance was set at $P < 0.05$.

Acknowledgments - This research was supported by Kyungpook National University Research Fund, 2012.

References

- [1] (a) Barnes PJ. (2011) Pathophysiology of allergic inflammation. *Immunological Reviews*, **242**, 31–50; (b) Kim SH, Jun CD, Suk K, Choi BJ, Lim H, Park S, Lee SH, Shin HY, Kim DK, Shin TY. (2006) Gallic acid inhibits histamine release and pro-inflammatory cytokine production in mast cells. *Toxicological Sciences*, **91**, 123–131.
- [2] (a) Frank SSJ. (1998) Amygdalin in *Prunus* leaves. *Phytochemistry*, **47**, 1537–1538; (b) Kosuge T, Ishida H, Ishii M. (1985) Studies on active substances in the herbs used for Oketsu ("Stagnant Blood") in Chinese medicine. II. On the anticoagulative principle in Persicae semen. *Chemical & Pharmaceutical Bulletin*, **33**, 1496–1498; (c) Fukuda T, Ito H, Mukainaka T, Tokuda H, Nishino H, Yoshida T. (2003) Anti-tumor promoting effect of glycosides from *Prunus persica* seeds. *Biological & Pharmaceutical Bulletin*, **26**, 271–273; (d) Wu H, Shi J, Xue S, Kakuda Y, Wang D, Jiang Y, Ye X, Li Y, Subramanian J. (2011) Essential oil extracted from peach (*Prunus persica*) kernel and its physicochemical and antioxidant properties. *LWT-Food Science and Technology*, **44**, 2032–2039; (e) Hwang HJ, Kim P, Kim CJ, Lee HJ, Shim I, Yin CS, Yang Y, Hahn DH. (2008) Antinociceptive effect of amygdalin isolated from *Prunus armeniaca* on formalin-induced pain in rats. *Biological & Pharmaceutical Bulletin*, **31**, 1559–1564; (f) Yun-Choi HS, Kim, SO, Kim JH, Lee JR, Cho HI. (1985) Modified smear method for screening potential inhibitors of platelet aggregation from plant sources. *Journal of Natural Products*, **48**, 363–370.
- [3] Shin TY, Park SB, Yoo JS, Kim IK, Lee HS, Kwon TK, Kim MK, Kim JC, Kim SH. (2010) Anti-allergic inflammatory activity of the fruit of *Prunus persica*: role of calcium and NF- κ B. *Food and Chemical Toxicology*, **48**, 2797–2802.
- [4] (a) Chin YW, Chai HB, Keller WJ, Kinghorn AD. (2008) Lignans and other constituents of the fruits of *Euterpe oleracea* (Acai) with antioxidant and cytoprotective activities. *Journal of Agricultural and Food Chemistry*, **56**, 7759–7764; (b) Anthony AR. (1990) ¹H and ¹³C NMR analysis of D-amygdalin: oligosaccharide assignment and sequencing. *Magnetic Resonance in Chemistry*, **28**, 765–773; (c) Wang M, Li J, Rangarajan M, Shao Y, LaVoie EJ, Huang TC, Ho CT. (1998) Antioxidative phenolic compounds from sage (*Salvia officinalis*). *Journal of Agricultural and Food Chemistry*, **46**, 4869–4873; (d) Abe F, Yamauchi T. (1986) Lignans from *Trachelospermum asiaticum* (Tracheolospermum) II. *Chemical and Pharmaceutical Bulletin*, **34**, 4340–4345; (e) Ida Y, Satoh Y, Ohtsuka M, Nagasao M, Shoji J. (1993) Phenolic constituents of *Phellodendron amurense* Bark. *Phytochemistry*, **35**, 209–215; (f) Perold GW, Beylis P, Howard AS. (1973) Metabolites of proteaceae. Part VII. Lacticolorin, a phenolic glucoside ester, and other metabolites of *Protea laticolor* Salisb. *Journal of the Chemical Society, Perkin Transaction I*, **6**, 638–643.

Isolation, Synthesis and Biological Evaluation of Phenylpropanoids from the Rhizomes of *Alpinia galanga*

Sumit S Chourasiya^{a,b}, Eppakayala Sreedhar^a, K. Suresh Babu^{a*}, Nagula Shankaraiah^b, V. Lakshma Nayak^c, S. Ramakrishna^c, S. Sravani^d and M.V. Basaveswara Rao^e

^aDivision of Natural Product Chemistry, CSIR- Indian Institute of Chemical Technology, Hyderabad-500007, India

^bDepartment of Medicinal Chemistry, National Institute of Pharmaceutical Education and Research (NIPER), Hyderabad-500037, India

^cDivision of Medicinal Chemistry & Pharmacology Division, CSIR-Indian Institute of Chemical Technology, Hyderabad-500 607, India

^dPharmacognosy Division, Manipal University, Manipal-576 104, India

^eDepartment of Chemistry, Krishna University, Machilipatnam, India

suresh@iict.res.in

Received: July 17th, 2013; Accepted: October 4th, 2013

Bioactivity guided investigation of the DCM: MeOH (1:1) extract from the rhizomes of *Alpinia galanga* led to the isolation of phenylpropanoids (**1-9**) and their structures were established by ¹H NMR, ¹³C NMR, IR and LC-MS/MS. These compounds have been evaluated for their *in vitro* anticancer activity against the human cancer cell lines A549 (lung cancer), Colo-205 (colon cancer), A431 (skin cancer), NCI H460 (lung cancer), PC-3 (prostate cancer), and HT-29 (colon cancer). Compounds **4** and **9** showed potent anticancer activity (ranging from 1.3–19.7 µg/mL) against all the tested cancer cell lines. In addition, an asymmetric synthesis of acetoxychavicol acetate (**1**) and *trans*-*p*-coumaryl alcohol (**4**) has been accomplished in six steps starting from readily available *p*-hydroxybenzaldehyde for the first time. Grignard reaction and Sharpless kinetic resolution reactions were utilized as the key steps to install the basic core.

Keywords: *Alpinia galanga*, Phenylpropanoids, Cytotoxic activity, Stereoselective synthesis.

Cancer is one of the most serious threats against human health in the world. Over the past few decades, extensive research has led to the development of a plethora of chemotherapeutic agents; however, none of these agents are capable of completely eliminating cancer [1]. The limitations of current anticancer drugs, increased incidence, and rapid development of drug resistance [2] have highlighted the need for the discovery of new anticancer agents, preferably with novel mechanisms of action. In this respect, the plant kingdom is now recognized as a source of promising chemopreventive agents [3]. They can be viewed as evolved privileged structures with potent biological activities and biologically pre-validated leads which have played a highly significant role in the discovery and development of new drugs for the treatment of human diseases. This is especially true in the cancer field, where about half the drugs are of natural origin and include two of the most important agents, taxol and taxotere.[4]

As part of pharmacological-phytochemical integrated studies of medicinal plants from the Indian flora [5], we are investigating chemical constituents of plants belonging to the Zingiberaceae family, as well as their cytotoxic activity [6]. In a continuing investigation, phytochemical studies were carried out on *Alpinia galanga*, a tree that grows commonly in the southern part of India, where it is popularly known as Kulanjan. This plant is reported to exhibit antileishmanial [7], antifungal [8], hypoglycemic [9], gastro-protective [10], and anticancer activities [11].

As part of an ongoing effort to discover potential leads from Indian medicinal plants for development of anticancer therapeutics, we observed potent, significant, cytotoxic activity in the DCM: methanol (1:1) extract of *Alpinia galanga* (L.) Willd. The active




Figure 1: Isolated compounds from the rhizomes of *Alpinia galanga*.

extract was subjected to bioassay-guided separation that resulted in the isolation of phenyl propanoid derivatives (**1-9**).

Herein, we report the isolation, structure elucidation and cytotoxicity evaluation of phenylpropanoids (**1-9**), along with the synthesis of *trans*-*p*-coumaryl alcohol (**4**) and acetoxychavicol acetate (**1**) by using the Grignard reaction and Sharpless kinetic resolution as the key steps. In addition, we also prepared a series of ester analogues by using different acids and assessed them for their cytotoxic activity. To the best of our knowledge, this is the first report on the asymmetric synthesis of phenylpropanoid ester derivatives

The concentrated DCM: MeOH (1:1) extract was chromatographed on silica gel, and the resultant fractions were subjected to bioassay for cytotoxic activity against a panel of cancer cell lines. Repeated column chromatography of the bioactive fractions resulted in the isolation of nine compounds. The compounds were identified as (1'S)-1'-acetoxychavicol acetate (**1**) [12,13], *p*-coumaryl diacetate (**2**) [14], (1'S)-1'-acetoxyeugenol acetate (**3**) [12,13], *trans*-*p*-coumaryl alcohol (**4**) [10], *p*-hydroxybenzaldehyde (**5**) [14],


1'S-1'-hydroxychavicol acetate (**6**) [10], *p*-methyl benzaldehyde (**7**), *trans-p*-hydroxycinnamyl acetate (**8**) [10], and *trans-o*-hydroxycinnamaldehyde (**9**) from ^1H and ^{13}C NMR data comparison with those reported in literature.

The isolated phenylpropanoids (**1-9**) were evaluated for their anticancer activity against a panel of six human cancer cell lines, A549 (lung cancer), Colo-205 (colon cancer), A431 (skin cancer), NCI H460 (lung cancer), PC-3 (prostate cancer), and HT-29 (colon cancer) by employing MTT assay [6]. Doxorubicin was used as the reference drug. The results, summarized in Table 1, are expressed as IC_{50} values. The *in vitro* screening results revealed that these compounds exhibit promising anticancer activity with IC_{50} values ranging from 1.9 to 13.5 $\mu\text{g/mL}$ against different cancer cell lines. The most active compounds, **4** and **9**, gave IC_{50} values ranging from 1.8–13.4 and 5.6–19.9 $\mu\text{g/mL}$, respectively.


Table 1: Cytotoxicity (IC_{50} , $\mu\text{g/mL}$) of the isolated compounds (**1-9**).

Compounds	Cell lines IC_{50} $\mu\text{g/mL}$					
	A549	Colo-205	A431	NCI H460	PC-3	HT-29
1	77.6	50.1	19.9	60.5	31.6	33.1
2	172.3	128.0	100.4	83.1	100.0	146.8
3	79.1	108.1	91.2	97.2	72.4	42.6
4	1.9	3.4	1.3	4.4	13.5	4.4
5	39.9	14.8	12.9	21.4	30.9	25.7
6	58.9	77.8	95.5	66.0	56.0	36.3
7	39.8	37.1	33.1	37.1	22.4	47.5
8	115.2	34.9	46.8	59.1	21.4	165.5
9	9.8	14.8	19.9	5.2	17.7	5.6
Doxorubicin	1.1	1.2	0.37	0.4	1.4	1.1

Synthesis of **4 and its derivatives:** Prompted by the biological properties of **4**, the present study was devised to synthesize new derivatives by using the above lead compounds and investigating their cytotoxic properties. As shown in Scheme 1, the synthesis began with 4-hydroxy benzaldehyde **11**, which was converted to its tosyl ester [**15**] **12** in 71% yield. Grignard reaction of **12** in the presence of vinyl magnesium bromide furnished racemic alcohol **13**, the tosyl deprotection [**15**] of which, in the presence of methanol under reflux conditions, yields 4-hydroxy phenyl methyl ether phenylpropanoid core **4** via allylic rearrangement[16]. Using this compound as a starting material, we made several derivatives by esterifying [17] with aliphatic and aromatic acids in the presence of EDCI and DMAP pyridine to afford corresponding esters **15a-h**.



Scheme 1: (a) tosyl chloride, TEA, r.t., 2 h, 71%; (b) vinyl magnesium bromide, THF, 0°C, 45 min, 60%; (c) methanol, K_2CO_3 , reflux, 2 h, 60%; (d) different acids, DMAP, EDCI, pyridine, r.t., 12 h, 65–85%.



Scheme 2: (a) TBDMSCl, imidazole, CH_2Cl_2 , r.t., 1 h, 91%; (b) vinyl magnesium bromide, THF, 0°C, 45 min, 91%; (c) (-) DIPT, $\text{Ti}(\text{iOPr})_4$, DCM, -20°C, 12 h, 51%; (d) TBAF, THF, 0°C, 1 h, 68%; (e) acetic anhydride, TEA, CH_2Cl_2 , r.t., 1 h, 70%; (f) acetic anhydride, TEA, CH_2Cl_2 , r.t., 30 min, 70% (for **22a**); (g) propionic anhydride, TEA, CH_2Cl_2 , r.t., 30 min, 73% (for **22b**); (h) tiglic acid, DMAP, EDCI, pyridine, r.t., 16 h, 74% (for **22c**).

Stereoselective synthesis of **22a-c and **21**:** Further, we have also extended our efforts to the synthesis of **21** and **22a-c**. As shown in Scheme 2. Synthesis began with 4-hydroxy benzaldehyde (**11**), which was converted to its TBDMS ether **16** [16] in 91% yield. Grignard reaction [16] of **16** with vinyl magnesium bromide furnished racemic alcohol **17**, which was resolved by Sharpless kinetic resolution [18] to give the required *S* alcohol **18** and epoxide **19**. The deprotection of the TBDMS group [16] in **17** and **18** with TBAF afforded phenylpropanoid cores **20** and **10**. Finally, the phenylpropanoid core **20** was coupled [17] with acetic acid in the presence of TEA to afford **21** in 71% yield.

Selective acetylation of the phenolic hydroxyl of **10** leading to monoacetate **22a-c** was accomplished with acetic acid and propanoic acid in the presence of triethylamine to afford **22a** (70%) and **22b** (73%), and with tiglic acid in the presence of EDCI and DMAP in pyridine to afford **22c** (74%).

Biological activity: All the synthesized derivatives were tested *in vitro* against a panel of cancer cell lines [6]. To our disappointment, introduction of the ester moiety at C-4 (phenolic OH) was not a good site to modify since none of the derivatives showed any activity. The lack of activity of the synthetic derivatives cannot be explained simply by reaction mechanism. Either the binding affinity or selectivity of the structures of the compounds to the reactive site of the target (enzyme) must play a key role. At this point, it may be that the hydroxyl group remains the optimum functional group to exhibit activity. These early conclusions point the way for further more focused studies aimed at the design and synthesis of more potent and selective analogues as biological tools and potential drug candidates.

Conclusion: In conclusion, we have isolated nine phenylpropanoid derivatives and their cytotoxic properties were investigated. The results of our study indicated that (*E*)-3-methoxyprop-1-enyl phenol (**4**) showed potent activity comparable with the standard drug doxorubicin. Based on these results, we also synthesized (1'S)-1'-acetoxychavicol acetate (**1**) and *trans-p*-coumaryl alcohol (**4**), and their derivatives using the Grignard and Sharpless asymmetric reactions as key steps.

Experimental

General: Melting points were recorded on a Fisher Johns apparatus and are uncorrected. FABMS were obtained from a VG Auto spec-M instrument, IR spectra from a Nicolet spectrometer, and ^1H NMR

and ^{13}C NMR spectra from Varian 200 and 400 MHz and Bruker 300 MHz spectrometers, using TMS as internal standard. The solvents used were all of AR grade and were distilled under positive pressure in a dry nitrogen atmosphere where necessary. TLC was performed on Merck silica gel 60 F₂₅₄ plates. Visualization was by spraying with 5% H₂SO₄ solution followed by heating. CC was performed on silica gel (60-120 mesh) purchased from Merck, Mumbai, India.

Plant material: The rhizomes of *Alpinia galanga* was collected from the forest of Tirumala in Chittoor Dist. (Andhra Pradesh, India) in January 2010 and identification was made by Prof. Dr K. Madhava Chetty, Department of Botany, Sri Venkateswara University, Tirupati. A voucher specimen of the plant is deposited in the herbarium, Department of Botany, with an accession number 546.

Extraction and isolation: The active chloroform extract (1 g) of *Alpinia galanga* was fractionated by CC with ethyl acetate-*n*-hexane. Initially, the column was started with 1% ethyl acetate: *n*-hexane, and then with increasing polarity of solvent (2%, 4%, 8%, 10%, 15%, 20%, 25%, and 30%); 9 compounds were obtained: (S)-4-(1-acetoxyallyl)phenyl acetate (**1**) (20 mg) [12,13], (E)-3-(4-acetoxyphenyl)allyl acetate (**2**) (25 mg) [14], (S)-1-(4-acetoxy-3-methoxyphenyl)allyl acetate (**3**) (20 mg) [12,13], (E)-4-(3-methoxyprop-1-en-1-yl)phenol (**4**) (20 mg) [10] (50 mg), 4-hydroxybenzaldehyde (**5**) (15 mg) [14], (S)-4-(1-hydroxyallyl)phenyl acetate (**6**) (18 mg) [10], 4-methylbenzaldehyde (10 mg) (**7**), (E)-3-(4-hydroxyphenyl)allyl acetate (**8**) (22 mg) [10], and (E)-3-(2-hydroxyphenyl)acrylaldehyde (26 mg) (**9**).

4-(1-Hydroxyallyl) phenyl 4-methylbenzenesulfonate (12): The 4-hydroxybenzaldehyde (**11**) (10 g, 80.96 mmol) was dissolved in DCM (100 mL). To the stirred solution, TEA (8.27g, 81.96 mmol) was added at 0°C, followed by *p*-toluene sulfonyl chloride (15.57g, 81.96 mmol). Then the reaction temperature was raised to 25°C and stirred for 2 h. The reaction was diluted with water (50 mL) and the layers separated. The organic layer was further washed with water (2×30 mL), followed by saturated aq. NaHCO₃ (30 mL) and brine (30 mL). The combined organic layers were dried with Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by flash chromatography using Hex: EA (85:15) to give tosylated *p*-hydroxybenzaldehyde (**12**).

White crystalline solid.

Yield: 71%.

^1H NMR (300 MHz, CDCl₃): δ 2.46 (s, 3H, CH₃), 7.16 (d, 2H, J = 8.4 Hz, Ar H), 7.32 (d, 2H, J = 8.12 Hz, Ar H), 7.71 (d, 2H, J = 8.3 Hz, Ar H), 7.83 (d, 2H, J = 8.4 Hz, Ar H), 9.96 (s, 1H, CHO).

^{13}C NMR (75 MHz, CDCl₃): δ 21.6 (CH₃), 123.0 (C), 128.4 (C), 129.9 (C), 131.2 (C), 131.9 (C), 134.7 (C), 145.8 (C), 153.8 (C), 190.6 (C=O).

ESI-MS: m/z 304.0 (M⁺)

4-(1-Hydroxyallyl)phenyl 4-methylbenzenesulfonate (13): To a solution of **12** (7 g, 25.37 mmole) in dry THF (70 mL) under a nitrogen atmosphere at 0°C was added, dropwise, 30.43 mL of a 1M solution of vinyl magnesium bromide in THF. The reaction was stirred for 45 min. at r.t. After completion (TLC), the reaction was quenched with saturated NH₄Cl solution at 0°C and extracted with ethyl acetate (2×50 mL). The combined organic layers were washed with brine, dried over Na₂SO₄ and the solvent was evaporated to dryness. The crude product was purified by CC using Hex: EA (90:10) as eluent to give vinyl alcohol **13**.

Light yellow viscous liquid.

Yield: 60%.

^1H NMR (300 MHz, CDCl₃): δ 2.46 (s, 3H, CH₃), 5.17 (dd, 2H, J = 10.4, 4.5 Hz, =CH₂), 5.31 (d, 1H, J = 17.3 Hz, CH-OH), 5.89-6.0 (m, 1H, CH=), 6.94 (d, 2H, J = 8.3 Hz, ArH), 7.25-7.31 (m, 4H Ar H), 7.70 (d, 2H, J = 8.3 Hz, Ar H).

^{13}C NMR (75 MHz, CDCl₃): δ 21.8 (CH₃), 74.3 (C-OH), 115.8 (C), 122.3 (C), 127.4 (C), 128.4 (C), 129.6 (C), 139.7 (=CH₂), 141.8 (CH=), 145.6 (C), 150.8 (C).

ESI-MS: m/z 304.0 (M⁺).

(E)-4-(3-Methoxyprop-1-en-1-yl) phenol (4): To a stirred solution of vinyl alcohol **13** (0.1 g, 0.3486 mmol) in methanol (bottle grade 5 mL) was added K₂CO₃ (0.24 g, 1.745 mmol), and the mixture was heated at reflux for 2 h. The reaction mixture was cooled to r.t., and checked by TLC. The mixture was acidified with 1 N HCl until the pH of the mixture reached 2. The combined aq/methanol solution was extracted with ethyl acetate (2×15 mL). The organic extracts were washed with brine (10 mL), dried with Na₂SO₄, filtered and concentrated *in vacuo*. The crude product was purified by CC using Hex: EA (80:20) to give **4**.

Light yellow viscous liquid.

Yield: 60%.

^1H NMR (300 MHz, CDCl₃): δ 3.36 (s, 3H, OCH₃), 4.05 (d, 2H, J = 6.0 Hz, CH₂-OMe), 5.67 (brs, 1H, OH), 6.01-6.11 (m, 1H =CH), 6.48 (d, 1H, J = 15.8 Hz, CH=), 6.73 (d, 2H, J = 8.4 Hz, Ar H), 7.19 (d, 2H, J = 8.4 Hz, Ar H).

^{13}C NMR (75 MHz, CDCl₃): δ : 57.6 (OCH₃), 73.3 (CH₂-O), 115.5 (C), 122.8 (C), 127.8 (C), 129.0 (C), 132.9 (C), 155.6 (C).

EI-MS: m/z 1640.0 (M⁺).

General procedure for the preparation of compounds 15a-15h:

EDCI (140 mg, 0.35 mmol) and DMAP (4.45 mg, 0.036 mmol) were added to a solution of tiglic acid in pyridine at 0°C. The reaction mixture was stirred at room temperature to dissolve the EDCI completely. After complete dissolution, a solution of compound **4** (0.04 g, 0.1829 mmol) in pyridine was added. The reaction was monitored by TLC to check the completion of the reaction (16 h). The reaction mixture was concentrated to remove pyridine. The reaction was quenched with aqueous 1 N HCl and extracted with DCM (2 × 20 mL). The product was purified by CC using *n*-hexane-EA (85:15) to give **15a-15h** in good yields.

(E)-4-(3-Methoxyprop-1-en-1-yl)phenyl acetate (15a)

Pale yellow liquid.

Yield: 72%.

IR (KBr): 2926, 1760, 1602, 1197 cm⁻¹.

^1H NMR (300 MHz, CDCl₃): δ 2.26 (s, 3H, CH₃CO), 3.36 (s, 3H, OCH₃), 4.05 (d, 2H, J = 6.0 Hz, CH₂O), 6.05-6.20 (m, 1H, =CH), 6.48 (d, 1H, J = 15.8 Hz, CH=) 7.00 (d, 2H, J = 7.9 Hz, Ar H), 7.34 (d, 2H, J = 7.9 Hz, Ar H).

^{13}C NMR (75 MHz, CDCl₃): δ 21.0 (CH₃), 57.8 (OCH₃), 72.8 (CH₂O), 115.5 (C), 121.6 (C), 126.1 (C), 127.2 (CH), 134.3 (CH), 155.0 (C), 168.7 (C=O).

ESI-MS: C₁₂H₁₄O₃, m/z 206.0 (M⁺), 246 (M⁺+40).

(E)-4-(3-Methoxyprop-1-en-1-yl)phenyl propionate (15b)

Pale yellow liquid.

Yield: 82%.

IR (KBr): 2922, 1758, 1598, 1346 cm⁻¹.

^1H NMR: (300 MHz, CDCl₃): δ 1.26 (t, 3H, J = 7.9 Hz, CH₃), 2.56 (q, 2H, J = 7.5 Hz, CH₂CH₃), 3.36 (s, 3H, OCH₃), 4.05 (d, 2H, J = 6.0 Hz, -CH₂), 6.05 (m, 1 H, =CH), 6.48 (d, 1H, J = 15.8 Hz, CH=), 6.73 (d, 2H, J = 8.4 Hz, Ar H), 7.19 (d, 2 H, J = 8.4 Hz, Ar H).

^{13}C NMR (75 MHz, CDCl_3): δ 8.93 (CH_3), 27.6 (CH_2), 57.8 (OCH_3), 72.8 (CH_2O), 121.5 (C), 123.6 (C), 126.1 (C), 127.2 (CH), 130.5 (CH), 148.3 (C), 172.8 (C=O).
ESI-MS: $\text{C}_{13}\text{H}_{16}\text{O}_3$, m/z 220.0 (M^+), 243 ($\text{M}^+ + 23$).

(E)-4-(3-Methoxyprop-1-en-1-yl)phenyl 3-methylbut-2-enoate (15c)

Pale yellow liquid.

Yield: 80%.

IR (KBr): 2924, 1739, 1601, 1123 cm^{-1} .

^1H NMR (300 MHz, CDCl_3): δ 2.0 (s, 3H, CH_3), 2.21 (s, 3H, CH_3), 3.36 (s, 3H, OCH_3), 4.05 (d, 2H, $J = 6.0$ Hz, CH_2O), 5.88 (s, 1H, $=\text{CHCO}$), 6.10–6.20 (m, 1 H, $=\text{CH}$), 6.48 (d, 1H, $J = 15.8$ Hz, $\text{CH}=\text{}$), 7.03 (d, 2H, $J = 8.4$ Hz, Ar H), 7.35 (d, 2 H, $J = 8.4$ Hz, Ar H).

^{13}C NMR (75 MHz, CDCl_3): δ : 27.1 (CH_3), 29.4 (CH_3), 58.0 (OCH_3), 72.3 (CH_2O), 115.1 (C), 122.0 (C), 125.9 (CH), 131.5 (CH), 155.4 (C), 162.9 (C=O).

ESI-MS: $\text{C}_{15}\text{H}_{18}\text{O}_3$, m/z 246.0 (M^+), 269 ($\text{M}^+ + 23$).

(E)-4-(3-Methoxyprop-1-en-1-yl)phenyl pent-4-enoate (15d)

Pale yellow liquid.

Yield: 85%.

IR (KBr): 2925, 1756, 1601, 1167 cm^{-1} .

^1H NMR (300 MHz, CDCl_3): δ 2.49 (q, 2H, $J = 6.8$ Hz, $=\text{CH-CH}_2$), 2.63 (t, 2H, $J = 7.8$ Hz, $\text{CH}_2\text{-CO}$), 3.36 (s, 3H, OCH_3), 4.05 (d, 2H, $J = 5.85$ Hz, $\text{CH}_2\text{-CH}_3$), 5.06 (dd, 2H, $J = 10.74$ Hz, $\text{CH}_2=\text{}$), 5.87–5.92 (m, 1H, $\text{CH}_2=\text{CH-}$), 6.18–6.25 (m, 1H, $=\text{CH}$), 6.54 (1H, d, $J = 16.5$ Hz, $\text{CH}=\text{}$), 6.9 (d, 2H, $J = 7.8$ Hz, Ar H), 7.35 (d, H, $J = 8.7$ Hz, Ar H).

^{13}C NMR (75 MHz, CDCl_3): δ 29.6 (CH), 33.5 (CH), 57.9 (OCH_3), 72.8 (OCH_2), 115.8 (C), 121.5 (C), 126.1 (C), 127.2 (CH), 131.4 (CH), 134.4 (CH), 136.2 (CH), 150.3 (C), 171.3 (C=O).

ESI-MS: m/z 246.0 (M^+), 269 ($\text{M}^+ + 23$).

(E)-4-(3-Methoxyprop-1-en-1-yl)phenyl hex-5-enoate (15e)

Pale yellow liquid.

Yield: 75%.

IR (KBr): 2923, 1759, 1506, 1166 cm^{-1} .

^1H NMR (300 MHz, CDCl_3): δ 1.84 (q, 2H, $J = 7.5$ Hz, $\text{CH}_2=\text{CH-CH}_2$), 2.18–2.28 (m, 2H, $-\text{CH}_2\text{CH}_2\text{CO}$), 2.54 (t, 2H, $J = 7.5$, $\text{CH}_2\text{-CO}$), 3.36 (s, 3H, $-\text{CH}_3$), 4.05 (dd, 2H, $J = 1.5$, 0.85 Hz, $-\text{CH}_2\text{-OCH}_3$), 5.76 (m, 1 H, $\text{CH}_2=\text{CH}$), 6.18 (m, 1H, $=\text{CH}$), 6.54 (1H, d, $J = 16.5$ Hz, $\text{CH}=\text{}$), 7.0 (d, 2H, $J = 8.3$ Hz Ar H), 7.35 (d, 2H, $J = 8.3$ Hz, Ar H).

^{13}C NMR (75 MHz, CDCl_3): δ : 23.9 (CH_2), 32.9 (CH_2), 33.5 (CH_2), 57.9 (OCH_3), 72.8 (OCH_2), 121.5 (C), 126.1 (C), 127.3 (CH), 131.3 (CH), 134.0 (CH), 137.2 (CH), 150.3 (C), 171.3 (C=O).

ESI-MS: m/z 260.0 (M^+), 283 ($\text{M}^+ + 23$).

(E)-4-(3-Methoxyprop-1-en-1-yl)phenyl octanoate (15f)

Pale yellow liquid.

Yield: 85%.

IR (KBr): 2925, 1761, 1607, 1215 cm^{-1} .

^1H NMR (300 MHz, CDCl_3): δ 0.9 (t, 3H, $J = 4.5$ Hz, CH_3), 1.29–1.33 (m, 6H, $-\text{CH}_2$), 1.61–1.7 (m, 2H, $-\text{CH}_2$), 2.25–2.30 (m, 2H, $-\text{CH}_2$), 2.50–2.55 (m, 2H, $\text{CH}_2\text{-CO}$), 3.36 (s, 3H, OCH_3), 4.05 (d, 2H, $J = 6.0$ Hz $\text{CH}_2\text{-OCH}_3$), 6.18 (m, 1H, $=\text{CH}$), 6.54 (1H, d, $J = 16.5$ Hz $\text{CH}=\text{}$), 7.0 (d, 2H, $J = 8.3$ Hz Ar H), 7.35 (d, 2 H, $J = 8.3$ Hz Ar H).

^{13}C NMR (75 MHz, CDCl_3): δ 14.0 (CH_3), 22.5 (CH_2), 24.8 (CH_2), 28.8 (CH_2), 29.6 (CH_2), 31.6 (CH_2), 34.5 (CH_2), 57.9 (OCH_3), 72.8 (OCH_2), 121.5 (C), 125.9 (C), 127.3 (C), 131.3 (CH), 134.4 (CH), 150.3 (C), 171.3 (C=O).

ESI-MS: m/z 290.0 (M^+), 313 ($\text{M}^+ + 23$).

(E)-4-(3-Methoxyprop-1-en-1-yl)phenyl 4-fluorobenzoate (15g)

Pale yellow liquid.

Yield: 82%.

IR (KBr): 2925, 1735, 1601, 1265 cm^{-1} .

^1H NMR (300 MHz, CDCl_3): δ 3.36 (s, 3H, OCH_3), 4.05 (d, 2H, $J = 6.0$ Hz, $\text{CH}_2\text{-OCH}_3$), 6.12–6.25 (m, 1 H, $=\text{CH}$), 6.40 (d, 1H, $J = 16.0$ Hz, $\text{CH}=\text{}$), 6.59 (d, 2H, $J = 16.0$ Hz, Ar H), 6.83 (d, 2H, $J = 16.0$ Hz, Ar H), 7.09 (d, 2 H, $J = 8.3$ Hz, Ar H), 7.83 (d, 2H, $J = 16.0$ Hz, Ar H).

^{13}C NMR (75 MHz, CDCl_3): δ 57.9 (OCH_3), 72.9 (OCH_2), 121.6 (C), 126.2 (C), 127.4 (C), 131.2 (C), 131.3 (CH), 131.8 (CH), 150.0 (C), 164.1 (C=O), 167.8 (C).

ESI-MS: m/z 286.0 (M^+), 287 ($\text{M}^+ + 1$).

4-((E)-3-Methoxyprop-1-en-1-yl)phenyl cinnamate (15h)

Pale yellow liquid.

Yield: 74%.

IR (KBr): 2921, 1728, 1601, 1137 cm^{-1} .

^1H NMR (300 MHz, CDCl_3): δ 3.34 (s, 3H, OCH_3), 4.06 (d, 2H, $J = 5.7$ Hz, $-\text{CH}_2\text{-OCH}_3$), 6.38 (d, 1H, $J = 5.2$ Hz, $\text{CH}=\text{}$), 6.57 (d, 1H, $J = 5.2$ Hz, $=\text{CH}$), 7.09 (d, 1H, $J = 5.2$ Hz, $\text{CH}=\text{CH-C=O}$), 7.3–7.48 (m, 10H Ar H).

^{13}C NMR (75 MHz, CDCl_3): δ 57.8 (CH_3), 72.8 (OCH_2), 117.2 (C), 121.6 (C), 126.6 (C), 127.3 (C), 127.9 (C), 128.2 (C), 128.8 (C), 130.3 (CH), 131.3 (CH), 134.2 (CH), 144.5 (CH), 146.5 (C), 150.1 (C), 165.2 (C=O).

ESI-MS (m/z): 317 ($\text{M}^+ + 23$).

4-((Tert-butylidimethylsilyl)oxy)benzaldehyde (16): About 10 g of *p*-hydroxybenzaldehyde (**11**) was dissolved in 100 mL of dry DCM and to the stirred solution was added imidazole portion wise at 0°C . After being stirred for 20 min at 0°C , TBDMSCl was added and the mixture was stirred for 1 h at r.t. The reaction was completed in 1 h. Then the reaction mixture was quenched with cold water and the compound extracted with chloroform. The organic layer was washed with brine and dried over sodium sulfate. The organic layers were combined and evaporated in a rotary evaporator. The crude product was purified by silica gel CC using *n*-hexane: EtOAc (85:15) to give **16** [16] as a white crystalline solid (91%).

1-4-((tert-butylidimethylsilyl)oxy)phenyl}prop-2-en-1-ol (17): To a solution of **16** (10 g, 42.37 mmole) in dry THF (100 mL) under a nitrogen atmosphere at 0°C was added, dropwise, 50.48 mL of a 1M solution of magnesium bromide in THF. The reaction was allowed to stir for 45 min at r.t. After completion (TLC), the reaction was quenched with saturated NH_4Cl solution at 0°C and extracted with ethyl acetate (2 \times 50 mL). The combined organic layers were washed with brine, dried over Na_2SO_4 and the solvent was evaporated to dryness. The crude product was purified by CC using Hex: EA (90:10) as eluent to give vinyl alcohol (**17**) [16] as light yellow viscous liquid (60%).

4-(1-Hydroxyallyl) phenol (20): To a stirred solution of **17** (2 g, 6.8 mmol) in LR grade THF at 0°C was added TBAF, dropwise, and the reaction was stirred for 1 h. The progress of the reaction was monitored by TLC. After completion (1 h), the reaction mixture was concentrated to remove THF and the oil was dissolved in EA and extracted with 1 N HCl (20 mL), washed with brine, dried over Na_2SO_4 and concentrated. The product was purified by silica gel CC using *n*-hexane: EtOAc (90:10) to give **20** [16] as a yellow oily liquid (68%).

4-(1-Acetoxyallyl) phenyl acetate (21): To a solution of **20** (1 g, 6.5 mmol) in DCM (10 mL), under a nitrogen atmosphere at r.t., was added, dropwise, triethylamine (2.32 mL, 16.392 mmol), followed by acetic anhydride (1.52 mL, 16.2 mmol). The reaction mixture

was stirred for 1 h at 0°C. After completion (TLC), the reaction was quenched with ice at r.t., extracted with CHCl₃ (2× 25 mL), the combined organic layers washed with brine, dried over Na₂SO₄ and the solvent evaporated to dryness. The crude product was purified by CC using Hex: EA (90:10) to give compound **21** [16] as a light yellow oily liquid (70%).

(S)-1-(4-((Tert-butyldimethylsilyl)oxy}phenyl)prop-2-en-1-ol (18) and (1S)-(4-((tert-butyldimethylsilyl)oxy}phenyl)(oxiran-2-yl)methanol (19): Completely evacuated, flame dried, powdered 4 Å molecular sieves (10 g) and (-)-DIPT (2.10 g, 9.38 mmol) were placed in a 250 mL two necked r.b flask and dissolved in dry DCM (40 mL). To the stirred mixture a solution of Ti (iOPr)₄ (2.22 g, 7.82 mmol) was added at -20°C, followed by (1.44 g, 9.38 mmol) cumene hydroperoxide at an interval of 20 min. After being stirred for 20 min, a solution of compound **17** (4.12 g, 15.64 mmol) in DCM (10 mL) was added, dropwise. The reaction mixture was allowed to stir for 12 h. The reaction was monitored by TLC. After completion (12 h), the reaction was quenched by a solution containing 4 g of NaOH dissolved in 40 mL of saturated NaCl at 0°C. Stirring continued for 3 h at r.t. before the mixture was filtered through Celite and the filtrate concentrated. The product was purified by silica gel CC using *n*-hexane: EtOAc (90:10) to give **18** and *n*-hexane: EtOAc (75:25) to give **19**.

Compound 18

Syrupy liquid.

Yield: 48%.

[α]_D²⁰: -15 (c 0.1, CHCl₃).

¹H NMR (300 MHz, CDCl₃): δ 0.18 (s, 6H, dimethyl), 0.98 (s, 9H, ter butyl), 5.09-5.18 (m, 2H, =CH₂), 5.31 (d, 1H, *J* = 16.9 Hz, -CHOH), 5.94-6.05 (m, 1H, CH=), 6.90 (d, 2H, *J* = 8.4 Hz, Ar H), 7.17 (d, 2H, *J* = 8.4 Hz, Ar H).

¹³C NMR (75 MHz, CDCl₃): δ -4.5 (C-Si), 17.8 (CH₃Si), 25.6 (CH₃C), 75.1 (C-OH), 114.3 (C), 120.0 (C), 127.5 (C), 135.2 (CH₂), 140.3 (CH), 154.8 (C).

ESI-MS (*m/z*): 287 (M⁺+23).

Compound 19

A syrupy liquid.

Yield: 47%.

[α]_D²⁰: -100 (c 0.1, CHCl₃).

¹H NMR (300 MHz, CDCl₃): δ 0.18 (s, 6H, CH₃), 0.98 (s, 9H, ter butyl), 2.70-2.73 (m, 1 H, CH-O), 2.90-2.93 (m, 1 H, CH-O), 3.13-3.16 (m, 1H, CH-O), 4.84 (d, 1H, *J* = 2.26 Hz, CH-OH), 6.73 (d, 2H, *J* = 8.4 Hz, Ar H), 6.9 (d, 2 H, *J* = 8.4 Hz, Ar H).

¹³C NMR (75 MHz, CDCl₃): δ: -4.69 (C-Si), 18.14 (CH₃), 25.6 (CH₃), 43.5 (-CH-O), 70.46 (CH-OH), 120.1 (C), 127.6 (C), 132.0 (C), 155.6 (C).

ESI-MS: C₁₅H₂₄O₃Si, *m/z* 281 (M⁺+1), *m/z* 303 (M⁺+23).

(S)-4-(1-Hydroxyallyl) phenol (10): To a stirred solution of compound **18** (2g, 6.8 mmol) in LR grade THF at 0°C was added TBAF, dropwise, and the reaction was stirred for 1 h. After completion of the reaction (TLC), the mixture was concentrated to remove THF, and the oil was dissolved in EA and extracted with 1 N HCl (20 mL), washed with brine, dried over Na₂SO₄ and concentrated. The product was purified by silica gel CC using *N*-hexane: EtOAc (90:10) to give **10**.

Yellow oily liquid.

Yield: 68%.

[α]_D²⁰: -15 (c 0.1, CH₃OH).

¹H NMR (300 MHz, CDCl₃): δ 5.0-5.18 (m, 2H, =CH₂), 5.25 (d, 1H, *J* = 15.6 Hz, CH-OH), 5.98-6.1 (m, 1 H, CH=), 6.73 (d, 2H, *J* = 8.4 Hz, Ar H), 7.19 (d, 2 H, 8.4 Hz, Ar H).

¹³C NMR (75 MHz, CDCl₃): δ: 64.6 (CH), 106.4 (C), 113.6 (C), 119.1 (CH₂), 124.6 (CH), 148.5 (C).

(S)-4-(1-Acetoxyallyl)phenyl acetate (23): To a stirred solution of compound **10** (0.05g, 0.33 mmol) in DCM (10 mL) under nitrogen at r.t. was added triethylamine (2.32 mL, 16.392 mmol), followed by acetic anhydride (51.48 mg, 0.39 mmol), dropwise. The reaction mixture was stirred for 1 h at 0°C. After completion (TLC), the reaction was quenched with ice at r.t. and extracted with CHCl₃ (2×10 mL). The combined organic layers were washed with brine, dried over Na₂SO₄ and the solvent was evaporated to dryness. The crude product was purified by CC using Hex: EA (90:10) to give compound **23** [16] as a light yellow oily liquid (70%).

General procedure for the preparation of compounds 22a & 22b:

To a stirred solution of compound **10** (0.05 g, 0.33 mmol) in DCM (10 mL) under a nitrogen atmosphere at r.t. was added triethylamine (2.32 mL, 16.392 mmol). To this solution, acetic anhydride (51.48 mg, 0.39 mmol) was added, dropwise. The reaction mixture was allowed to stir for 30 min. at 0°C to achieve selective acylation of the phenolic OH. After completion (TLC), the reaction was quenched with ice at r.t. and extracted with CHCl₃ (2×10 mL). The combined organic layers were washed with brine, dried over Na₂SO₄ and the solvent evaporated to dryness. The crude product was purified by CC using Hex: EA (90:10) to give compounds **22a** and **22b**.

(S)-4-(1-Hydroxyallyl) phenyl acetate (22a)

Light yellow oily liquid.

Yield: 70%.

[α]_D²⁰: -65 (c 0.1, CHCl₃).

IR (KBr): 3019, 1765, 1741, 1606 cm⁻¹.

¹H NMR (300 MHz, CDCl₃): δ 2.29 (s, 3H, OAc), 5.22-5.31 (dd, 2H, =CH₂), 5.3 (d, 1H, CH-OH), 5.9-6.02 (m, 1H, CH=) 7.04 (d, 2H, *J* = 8.4, Ar H), 7.3 (d, 2H, *J* = 8.4 Hz, Ar H).

¹³C NMR (75 MHz, CDCl₃): δ 21.0 (CH₃), 75.14 (CH-OH), 116.9 (C), 121.5 (C), 128.3 (C), 135.9 (CH₂), 136.3 (CH), 150.3 (C), 169.2 (C=O).

ESI-MS: C₁₁H₁₂O, *m/z* 192 (M⁺).

(S)-4-(1-Hydroxyallyl)phenyl propionate (22b)

Light yellow oily liquid.

Yield: 73%.

[α]_D²⁰: -105 (c 0.1, CHCl₃).

IR (KBr): 2922, 1758, 1729, 1640, 1346 cm⁻¹.

¹H NMR (300 MHz, CDCl₃): δ 1.09 (t, 3H CH₃), 2.53 (q, 2H, CH₂), 5.22-5.31 (dd, 2H, =CH₂), 5.32 (d, 1H, *J* = 17.34 Hz, CH-OH), 5.90-6.10 (m, 1H, CH=), 7.04 (d, 2H, *J* = 8.4, Ar H), 7.3 (d, 2H, *J* = 8.4 Hz, Ar H).

¹³C NMR (75 MHz, CDCl₃): δ 9.5 (CH₃), 21.04 (CH₂), 75.1 (CH-OH), 116.9 (C), 121.5 (C), 128.3 (C), 135.9 (CH₂), 136.3 (CH), 150.3 (C), 169.2 (C=O).

ESI-MS: C₁₂H₁₄O₃, *m/z* 206 (M⁺).

(S, Z)-4-(1-Hydroxyallyl)phenyl 2-methylbut-2-enoate (22c): To a stirred solution of tiglic acid (1.09 mmol) in pyridine at 0°C, EDCI (140 mg, 0.35 mmol) and DMAP (4.45 mg, 0.036 mmol) were added. The reaction mixture was stirred at room temperature and, after complete dissolution of EDC, a solution of compound **10** (0.04 g, 0.1829 mmol) in pyridine was added. The reaction was allowed to stir for 16 h and then the mixture was concentrated to remove pyridine. The reaction was quenched with aqueous 1 N HCl and extracted with DCM (2 × 20 mL). The combined organic layers were washed with brine, dried over Na₂SO₄ and evaporated to

dryness. The product was purified by CC using *n*-hexane: EtOAc (85:15) to give **22c**.

Yellow liquid.

Yield: 74%.

$[\alpha]_D^{20}$: -25 (*c* 0.1, CHCl₃).

IR (KBr): 3445, 2922, 1729, 1640, 1346 cm⁻¹;

¹H NMR (300 MHz, CDCl₃): δ 1.90 (d, 3H, *J*=7.1 Hz, CH₃), 1.94 (s, 3H, CH₃), 2.15 (brs 1H OH), 5.31 (dd, 2H, =CH₂), 5.32 (d, 1H, *J*=17.3 Hz, CH-OH), 7.04 (d, 2H, *J*=8.4, Ar H), 7.3 (d, 2H, *J*=8.4 Hz, ArH).

¹³C NMR (75 MHz, CDCl₃): δ 12.1 (CH₃), 14.6 (CH₃), 74.8 (CH-OH), 114.7 (C), 115.2 (C), 121.7 (C), 127.3 (C), 127.4 (CH), 139.4 (CH), 150.4 (C), 166.5 (C=O).

ESI-MS: C₁₄H₁₆O₃, *m/z* 232 (M⁺), 255 (M⁺+23).

Acknowledgements - The authors thank Dr J. S. Yadav, Director, IICT, for his constant encouragement. The authors also thank Dr Ahmed Kamal, Project Director, NIPER- Hyderabad for his help during the research. Financial assistance from the Ministry of Chemical and Fertilizer in the form of a fellowship is gratefully acknowledged.

References

- [1] Cragg GM, Grothaus PG, Newman DJ. (2009) Impact of natural products on developing new anti-cancer agent. *Chemical Reviews*, **109**, 3012-3043.
- [2] Sreedhar AS, Csermely P. (2004) Heat shock protein in the regulation of apoptosis: New strategies in tumour therapy. *Pharmacology & Therapeutics*, **101**, 227-257.
- [3] Cragg GM, Newman DJ. (2005) Plants as source of anticancer agent. *Journal of Ethnopharmacology*, **100**, 72-79
- [4] Newman DJ, Cragg GM, Snader KM. (2003) Natural products as sources of new drugs over the period 1981-2002. *Journal of Natural Products*, **66**, 1022-1037.
- [5] (a) Babu KS, Babu TH, Srinivas PV, Kishore KH, Murthy USN, Rao JM. (2006) Synthesis and biological evaluation of novel C (7) modified chrysin analogues as antibacterial agents. *Bioorganic & Medicinal Chemistry Letters*, **16**, 221-224; (b) Babu KS, Babu TH, Srinivas PV, Sastry BS, KH. Kishore, Murthy USN, Rao JM. (2005) Synthesis and *in vitro* study of novel 7-*O*-acyl derivatives of oroxylin A as antibacterial agents. *Bioorganic & Medicinal Chemistry Letters*, **15**, 3953-3956 ;(c) Babu TH, Manjulatha K, Hymavathi A, Tiwari AK, Purohit MG, Rao JM. (2010) Gastroprotective flavonoid constituents from *Oroxylum indicum* Vent. *Bioorganic & Medicinal Chemistry Letters*, **20**, 117-120.
- [6] (a) Reddy PP, Ranga Rao R, Shashidhar J, Sastry BS, Madhusudana Rao J, Suresh Babu K. (2009) Phytochemical investigation of labdane diterpenes from the rhizomes of *Hedychium spicatum* and their cytotoxic activity. *Bioorganic & Medicinal Chemistry Letters*, **19**, 6078-6081 and references cited therein; (b) Suresh Kumar G, Prabhakar Reddy P, Shaik TB, Kalivendi SV, Babu KS. (2010) Two new cytotoxic labdane diterpenes from the rhizomes of *Hedychium coronarium*. *Bioorganic & Medicinal Chemistry Letters*, **20**, 7544-7548.
- [7] Singh KR, Dey CS, Sharma SS, Bhutani KK, Singh IP. (2010) Antileishmanial phenylpropanoids from *Alpinia galanga* (Linn.) Wild. *Indian Journal of Experimental Biology*, **48**, 314-315.
- [8] Phongpaichit S, Subhadhirasaku S, Wattanapiromsaki C. (2005) Antifungal activities of extracts from Thai medicinal plants against opportunistic fungal pathogens associated with AIDS patients. *Mycoses*, **48**, 333-338.
- [9] Akhtar MS, Khan MA, Malik MT. (2002) Hypoglycemic activity of *Alpinia galanga* rhizome and its extracts in rabbits. *Fitoterapia*, **73**, 623-628.
- [10] Matsuda H, Pongpiriyadacha Y, Morikawa T, Ochi M, Yoshikawa M. (2003) Gastroprotective effects of phenylpropanoids from the rhizomes of *Alpinia galanga* in rats: structural requirements and mode of action. *European Journal of Pharmacology*, **471**, 59-67.
- [11] Moffat J, Hashimoto M, Kojima A, Kennedy DO, Murakami A, Koshimizu K, Ohigasha H, Matsui-Yuasa I. (2000) Apoptosis induced by 1'-acetoxychavicol acetate in Ehrlich ascites tumour cell is associated with modulation of polyamine metabolism and caspase-3 activation. *Carcinogenesis*, **21**, 2151-2157.
- [12] Janseen AM, Scheffer JJ. (1985) Acetoxychavicol acetate, an antifungal component of *Alpinia galanga*. *Planta Medica*, **51**, 507-511.
- [13] Mitsui S, Kobayashi S, Nagahori H, Ogiso A. (1976) Constituents from the seeds of *Alpinia galanga* Wild and their Anti-ulcer activities. *Chemical and Pharmaceutical Bulletin*, **24**, 2377-2382.
- [14] Noro T, Sekia T, Katoh M, Oda Y, Miyase T, Kuruyanagi M, Ueno A, Ukishima S. (1988) Inhibitor of xanthine oxidase from *Alpinia galanga*. *Chemical and Pharmaceutical Bulletin*, **36**, 244-248.
- [15] Raji Reddy Ch, Rao NN. (2010) Total synthesis of a diarylheptanoid, rhoiptelol B. *European Journal of Organic Chemistry*, 345-351.
- [16] Azuma H, Miyasaka K. (2006) Lipase-catalyzed preparation of optically active 1'-acetoxychavicol acetates and their structure-activity relationships in apoptotic activity against human leukemia HL-60 cells. *Bioorganic & Medicinal Chemistry*, **14**, 1811-1818.
- [17] Sreedhar E, Kumar RSC, Reddy GV, Robinson A, Babu KS, Rao JM, Srinivas PV. (2009) The first total synthesis of neohelmantins A-D, amino derivatives of the 1,2-dihydroxypropane core and biological evaluation. *Tetrahedron Asymmetry*, **20**, 440-448.
- [18] Gao Y, Hanson RM, Klunder JM, Ko SY, Masamune H, Sharpless, KB. (1987) Catalytic asymmetric epoxidation and kinetic resolution: Modified procedures including *in situ* derivatization. *Journal of American Chemical Society*, **109**, 5765-5780.

Rosmarinic Acid Interaction with Planktonic and Biofilm *Staphylococcus aureus*

Livia Slobodníková^a, Silvia Fialová^b, Helena Hupková^a and Daniel Grančai^b

^aComenius University in Bratislava, Faculty of Medicine, Institute of Microbiology, Sasinkova 4, 811 08 Bratislava, Slovakia

^bComenius University in Bratislava, Faculty of Pharmacy, Department of Pharmacognosy and Botany, Odbojárov 10, 832 32 Bratislava, Slovakia

fialova@pharm.uniba.sk

Received: August 26th, 2013; Accepted: September 23rd, 2013

The subject of study was the evaluation of antibacterial activities of rosmarinic acid (RA) on clinical *Staphylococcus aureus* strains obtained from catheter-related infections. Minimal inhibitory (MIC) and minimal bactericidal concentrations (MBC) of RA were tested by broth microdilution assay. Biofilm-eradication activity was detected on 24-hour biofilm in microtiter plates using a regrowth technique; activity on biofilm formation was measured by a microtiter plate method after RA application to bacterial samples after 0, 1, 3 and 6 hours of biofilm development. RA had antimicrobial activity on all tested strains in concentrations from 625 to 1250 $\mu\text{g}\cdot\text{mL}^{-1}$ (MICs equal to MBCs). No biofilm-eradication activity on 24-hour biofilm was observed in the tested range of concentrations (from 156 to 5000 $\mu\text{g}\cdot\text{mL}^{-1}$). Subinhibitory RA concentrations suppressed the biofilm production, when applied at early stages of its development. Concentrations lower than subinhibitory stimulated the biofilm mass production in a concentration- and time-dependent manner. Considering our results, RA could be a candidate for a topical antimicrobial agent with killing activity on planktonic forms of bacteria and suppressing activity in the early stages of biofilm development, but probably not for the therapy of catheter-related infections as a sole agent.

Keywords: Rosmarinic acid, Antibacterial activity, *Staphylococcus aureus*, Catheter-related infections, Biofilm.

Phenolic acids are important phenols produced in plants, for which they have diverse functions including participation in plant-microbe interaction or symbiosis [1]. One of the most common phenolic acids, widely found in the Lamiaceae family, is rosmarinic acid (RA), an ester of caffeic acid and 3, 4-dihydroxyphenylacetic acid, synthesized in the plant tissue from the amino acids L-phenylalanine and L-tyrosine [2]. RA is an important secondary metabolite in many medical plants that have been used in traditional medicine for more than 2000 years. It is the main bioactive compound in some of them [3] and many plant extracts containing RA are used as phytotherapeutical preparations [4].

Natural polyphenols and related substances could play an important role in the human defense system, e.g. as antimicrobial, antioxidant, or anticancer agents [2,4]. RA and its derivatives possess promising biological activities, such as neuroprotective, cardioprotective and chemopreventive effects, as well as antioxidant, astringent, anti-inflammatory and antimicrobial activity [4-7]. RA is a biologically active, non-toxic substance with antioxidant activity stronger than that of α -tocopherol (vitamin E) [4]. The anti-inflammatory activity of RA was proved by its inhibition of lipoxygenase and cyclooxygenase, which resulted in the inhibition of the arachidonic acid pathway [2].

Antibacterial activity of RA has been shown against Gram-positive and certain Gram-negative bacteria [8,9]; however, up till now there are no available data either about RA activity on clinical bacterial strains from hospital infections, or about its interaction with biofilm forms of bacterial growth. *Staphylococcus aureus*, a frequent skin colonizer, is one of the most important pathogens with respect to catheter-related infections, characteristic of biofilm formation [10]. The increasing frequency of polyresistant, mostly nosocomial strains, necessitates research on new antimicrobial agents, active both against planktonic and biofilm forms of microbial growth; to

such agents may belong substances of herbal origin as well. This study is, therefore, devoted to research of RA *in vitro* activity on clinical *S. aureus* strains from catheter-related infections.

Antimicrobial activity of rosmarinic acid on planktonic bacteria:

The suppressive activity of RA on several microbial species has already been published. Previous works on antimicrobial activity of RA were provided using both Gram-positive and Gram-negative bacteria. Mencherini *et al.* detected major bacteriostatic effects of RA against the Gram-positive bacteria, *S. aureus* and *S. epidermidis*. In the case of Gram-negative bacteria, yeasts, and molds, the MICs were 20-times higher [3]. Klancnik *et al.* examined the antimicrobial activity of RA against *Bacillus cereus*, *S. aureus*, *Salmonella infantis*, *Campylobacter jejuni* and *C. coli* (MICs on agar dilution were 6 mg/mL, 10 mg/mL, 1.25 mg/mL, 1.25 mg/mL, and 1.25 mg/mL, respectively) [11]. In our study, the MIC and MBC of RA were measured by a standardized broth microdilution method, using both susceptible and polyresistant *S. aureus* clinical isolates, including MRSA strains. Without respect to the antimicrobial susceptibility patterns of the strains, RA proved active against all of them, in concentrations from 625 to 1250 $\mu\text{g}\cdot\text{mL}^{-1}$ (Table 1). The MIC values were equal to the MBCs.



Antibiofilm activity of rosmarinic acid:

Microorganisms growing in biofilm are protected against many external factors, including antimicrobial agents [12]. The minimal biofilm eradicating concentrations of many antibiotics can be several times higher than MICs, measured in the planktonic form of growth, which may disable the clinical use of many of these antibiotics in the therapy of biofilm-related infections [13]. Even if many data are available on biofilm-inhibitory activity of various plant extracts, or molecules of plant origin [14-17], data about the real biofilm eradicating activity (i.e. total killing of bacteria in already established biofilm) are missing. Data about RA interaction with the biofilm form of





Table 1: Rosmarinic acid activity against *S. aureus* strains (tested in Mueller-Hinton broth).

Strain	Origin	Antimicrobial susceptibility						RA MIC/MBC [$\mu\text{g}\cdot\text{mL}^{-1}$]
		OXA	GEN	CIP	VAN	COT	MLSB	
1	CVC	R	S	R	S	S	+	625
2	CVC	R	S	R	S	S	+	625
3	CVC	S	S	R	S	S	-	625
4	CVC	R	S	R	S	S	+	625
5	CVC	S	S	R	S	S	+	625
6	CVC	R	S	R	S	S	+	625
7	CVC	R	R	R	S	S	+	625
8	CVC	S	S	R	S	S	+	625
9	WDC	S	S	S	S	S	+	625
10	WDC	S	S	S	S	S	-	1250
11	WDC	S	S	S	S	S	-	625
12	WDC	S	S	S	S	S	-	1250
13	WDC	S	S	S	S	S	-	1250
14	RSC	R	S	R	S	R	+	1250
15	RSC	R	S	R	S	S	+	625
16	RSC	S	S	S	S	S	+	1250
ATCC 43300	R							625
ATCC 29213	S							625

CVC – central venous catheter; WDR - wound drainage catheter; RSC - respiratory suction catheter; OXA – oxacillin; GEN – gentamicin; CIP – ciprofloxacin; VAN – vancomycin; COT – cotrimoxazole; MLSB – resistance to macrolides, lincosamides and streptogramins-B; RA – rosmarinic acid; MIC – minimal inhibitory concentration; MBC – minimal bactericidal concentration; S – susceptible; R - resistant

**Figure 1:** Impact of rosmarinic acid on biofilm formation by *Staphylococcus aureus* after 24 hours cultivation (strain 11); the values are mean \pm SD, * $P < 0.05$, versus control (bacteria in medium alone).**Figure 2:** Impact of rosmarinic acid on biofilm formation by *Staphylococcus aureus* after 24 hours cultivation (strain 13); the values are mean \pm SD, * $P < 0.05$, versus control (bacteria in medium alone).

bacteria in the available literary sources are lacking as well. Therefore, the next step in our study was detection of RA antibiofilm activity. Despite our expectations, we could not see any biofilm eradication activity of RA in our test system (24 hour biofilm on the surface of polystyrene microtiter wells; detection of regrowth of the surviving bacteria), even when the highest RA concentration used in the study ($5000 \mu\text{g}\cdot\text{mL}^{-1}$) exceeded the detected MICs (from 625 to $1250 \mu\text{g}\cdot\text{mL}^{-1}$; Table 1) by two- to three-times.

**Figure 3:** Concentration- and time-dependent activity of rosmarinic acid on biofilm formation by *Staphylococcus aureus* (strain 11).**Figure 4:** The corresponding ethanol solvent activity on biofilm formation by *Staphylococcus aureus* (strain 11).**Figure 5:** Concentration- and time-dependent activity of rosmarinic acid on biofilm formation by *Staphylococcus aureus* (strain 13).**Figure 6:** The corresponding ethanol solvent activity on biofilm formation by *Staphylococcus aureus* (strain 13).

Various plant extracts or pure substances of herbal origin have been screened for biofilm-reducing activity (for review see Taraszkiewicz *et al.* [16]), with several promising results. The biofilm suppressing activity may be mediated by interfering with microbial attachment mechanisms [18], by interfering with autoinducer molecules or their receptors (quorum quenching activity) [19], or by specific suppression of genetic determinants of biofilm matrix synthesis [15]. RA impact on biofilm formation in our study was tested in very rich medium supporting the biofilm production (Tryptose-Soy Broth supplemented with 1% glucose – TSB-S). As the composition of medium substantially influences the measured MIC values, one- to two-dilutions higher MICs were achieved in this richer medium (from 2500 to 5000 $\mu\text{g.mL}^{-1}$; Figures 1 and 2 for strains 11 and 13) in comparison with MIC values obtained in the Mueller-Hinton medium (MH), used for MIC, MBC, and biofilm-eradication activity testing (from 625 to 1250 $\mu\text{g.mL}^{-1}$; Table 1). Unfortunately, MH medium is not effective in supporting biofilm production, so it was not possible to use it in the testing of the RA biofilm-reducing activity. The sub-inhibitory concentrations of RA (MIC/2, i.e. 1250 $\mu\text{g.mL}^{-1}$ or 2500 $\mu\text{g.mL}^{-1}$, in TSB-S) showed a mild biofilm-suppressing activity, but the concentrations lower than sub-inhibitory (\leq MIC/4, i.e. \leq 625 $\mu\text{g.mL}^{-1}$ or \leq 1250 $\mu\text{g.mL}^{-1}$ in TSB-S) stimulated biofilm mass production (Figures 1 and 2).

When evaluating the role of the time interval between seeding the microtiter wells with bacteria and the moment of RA application, a time-, concentration- and strain-dependent pattern was observed (Figures 3 and 5). Generally, the highest biofilm-inhibitory activity was observed at the very early stages of biofilm production (1 to 3 hours). On the other hand, the highest biofilm-stimulating activity was observed when applied to 6-hour biofilm (Figures 3 and 5). An exception was strain 13 at the application time of 6 hours, when the highest stimulatory effect was observed in higher concentrations (2500 $\mu\text{g.mL}^{-1}$ in TSB-S) (Figure 5). This strain had a higher MIC (1250 $\mu\text{g.mL}^{-1}$ in MH medium, and 5000 $\mu\text{g.mL}^{-1}$ in TSB-S) and probably more developed biofilm after 6-hour cultivation than strain 11 (Figures 3 and 5).

Except for the biofilm stimulation by the RA ethanol solution in low concentrations, a stimulatory activity was detected by the ethanol solvent itself (Figures 4 and 6), but lower than in the case of RA. This phenomenon has already been described by several authors, who detected biofilm stimulation by low concentrations of antibiotics [20], alcohol [21], or even plant extracts [22, 23]. It seems that RA also belongs to the xenobiotic class, which can induce the bacterial defense response to environmental stress if applied in rather low concentrations. It is also possible, that ethanol and RA acted synergistically in the process of biofilm building by the *S. aureus* strains used in our study.

Rosmarinic acid in the present study showed effective killing activity on all of the tested *S. aureus* strains and decreased the biofilm production at sub-inhibitory concentrations, when applied at the early stages of its formation. Based on the obtained results, RA could be a candidate for a potential antimicrobial agent with killing activity on planktonic forms of bacteria and suppressing activity in the early stages of biofilm development. However, lack of RA effect on already developed biofilm does not support our hypothesis about its use in the therapy of catheter-related infections as a sole agent.

Experimental

Bacterial strains: Sixteen clinical *S. aureus* strains isolated from infections associated with central venous, wound, and respiratory

suction catheters were used in the study (Table 1). Two collection strains (ATCC 29213 and ATCC 43300) were included as well. The strains were preserved in Brain-Heart Infusion with 20% glycerol at -20°C . Before analysis, they were inoculated on blood agar and cultivated overnight at 37°C .

Antimicrobial susceptibility: The antimicrobial susceptibility to oxacillin, gentamicin, ciprofloxacin, vancomycin, cotrimoxazole, erythromycin and clindamycin was tested by the disk-diffusion method [24], and interpreted according to the CLSI guidelines [25]. Commercial antibiotic discs (OXOID, UK) and Mueller-Hinton agar medium (OXOID, UK) were used in the tests. Resistance to oxacillin/methicillin was confirmed by the Penicillin Binding Protein 2a (PBP2a) latex agglutination test (OXOID, UK).

Rosmarinic acid: Pure RA (Sigma-Aldrich, Germany) was dissolved in 50% ethanol, diluted 1:1 with twice concentrated medium, and afterwards serial geometric dilutions from 5000 to 156 $\mu\text{g.mL}^{-1}$ were prepared. Mueller-Hinton Broth (OXOID, UK) was used for antimicrobial activity testing on planktonic bacteria and for biofilm eradication activity, and Tryptose-Soy Broth (OXOID, UK) supplemented with 1% of glucose for detection of impact on biofilm production. The final concentrations of solvent (ethanol) in the samples ranged from 12.5 to 0.8%.

Rosmarinic acid antimicrobial activity: The planktonic form of bacterial growth was tested for minimal inhibitory (MIC) and minimal bactericidal concentrations (MBC) of RA by broth microdilution assay according to CLSI [26].

Biofilm eradication activity: The antibiofilm activity of RA was tested on 24 h biofilm prepared on the surface of microtiter wells in Tryptose-Soy Broth supplemented with 1% glucose according to Stepanovic *et al.* [27]. The medium containing biofilm-free bacteria was discarded and the wells were washed 3-times with PBS in order to remove bacteria not associated with biofilm. Rosmarinic acid in concentrations $>$ MBC was added to washed biofilms in the wells. After 24 h incubation at 35°C , the RA was drained out, the wells washed and fresh medium added. The biofilm eradication activity was detected by the regrowth method [28], after additional 24 h cultivation at 35°C .

Impact on biofilm formation: The activity of RA on biofilm formation was tested in microtiter plate wells in Tryptose-Soy Broth with 1% of glucose. Rosmarinic acid was added to the bacteria in the microtiter plate wells at 0, 1, 3, and 6 h after inoculation. The intensity of biofilm production was measured spectrophotometrically at 570 nm (MRX Microplate Reader, DYNEX Technologies, USA) after fixation with methanol, staining with 0.5% crystal violet and elution with 99% ethanol [27]. Medium free bacteria was used as a blank sample. All tests were performed in 3 parallel samples. In each test run, samples containing the solvent in corresponding concentrations were tested. Bacterial suspensions in RA- and solvent-free medium were used as controls.

Statistical analysis: The results were statistically evaluated by the unpaired t-test, using two-tailed *P* value.

Acknowledgments - This study was supported by the grant of the Slovak Ministry of Education VEGA 1/0564/11 (Microbial biofilm on abiotic surfaces in patients body - its forming characteristics, influence, and possibilities of elimination) and VEGA1/0059/11 (Screening of biological activities of selected plant species and the identification of their active constituents) and the grant of the Faculty of Pharmacy FAF UK 18/2013 (Screening of secondary

metabolites and determination of anti-glycation and antimicrobial activity of selected species of *Mentha* L. and *Lycopus* L. leaves.

The authors also thank Dr Viera Žufková for English emendation.

References

- [1] Santi MM, Dipjyoti C, Satyari D. (2010) Phenolic acids act as signaling molecules in plant-microbe symbioses. *Plant Signaling & Behavior*, **5**, 359-368.
- [2] Petersen M. (2013) Rosmarinic acid: new aspects. *Phytochemistry Reviews*, **12**, 207-227.
- [3] Mencherini T, Picerno P, Scesa C, Aquino R. (2007) Triterpene, antioxidant, and antimicrobial compounds from *Melissa officinalis*. *Journal of Natural Products*, **70**, 1889-1894.
- [4] Holzmanova V. (1996) Rosmarinic acid and its biological activity, *Chemical Papers*, **90**, 486-496.
- [5] Bulgakov VP, Inyushkina YV, Fedoreyev SA. (2012) Rosmarinic acid and its derivatives: biotechnology and applications. *Critical Reviews in Biotechnology*, **32**, 203-217.
- [6] Tang GY, Li XJ, Zhang HY. (2008) Antidiabetic components contained in vegetables and legumes. *Molecules*, **13**, 1189-1194.
- [7] Xiao C, Xinxin C, Jiakang H, Lanxiang J, Miaomiao W, Qingjun C, Mingfeng G, Xianxing X, Xuming D, Jiakang H. (2012) Effects of a natural prolyl oligopeptidase inhibitor, rosmarinic acid, on lipopolysaccharide-induced acute lung injury in mice. *Molecules*, **17**, 3586-3598.
- [8] Moreno S, Scheyer T, Romano CS, Vojnov AA. (2006) Antioxidant and antimicrobial activities of rosemary extracts linked to their polyphenol composition. *Free Radical Research*, **40**, 223-231.
- [9] Cetin-Karaca H. (2011) Evaluation of natural antimicrobial phenolic compounds against foodborne pathogens. Master's Thesis. University of Kentucky, Kentucky, USA Paper 652. Available online: http://uknowledge.uky.edu/gradschool_theses/652 (accessed on 10 June 2013).
- [10] Cramton SE, Götz F. (2004) Biofilm development in *Staphylococcus*. In *Microbial biofilms*. Ghannoun M, O'Toole GA. (Eds). ASM Press, Washington, DC, USA, 64-84.
- [11] Klančnik A, Piskernik S, Jersek B, Smole Mozina S. (2010) Evaluation of diffusion and dilution methods to determine the antibacterial activity of plant extracts. *Journal of Microbiological Methods*, **81**, 121-126.
- [12] Donlan RM, Costerton JW. (2002) Biofilms: Survival mechanisms of clinically relevant microorganisms. *Clinical Microbiology Reviews*, **15**, 167-193.
- [13] Kotulova D, Slobodnikova L. (2010) Susceptibility of *Staphylococcus aureus* biofilms to vancomycin, gentamicin and rifampin. *Epidemiologie, mikrobiologie, imunologie*, **59**, 80-87.
- [14] Carneiro VA, Santos HS, Arruda FVS, Bandeira PN, Albuquerque MR, Pereira MO, Henriques M, Cavada BS, Teixeira EH. (2011) Casbane diterpene as a promising natural antimicrobial agent against biofilm-associated infections. *Molecules*, **16**, 190-201.
- [15] Lee JH, Park JH, Cho HS, Joo SW, Cho MH, Lee J. (2013) Anti-biofilm activities of quercetin and tannic acid against *Staphylococcus aureus*. *Biofouling*, **29**, 491-499.
- [16] Taraszkiewicz A, Fila G, Grinholc M, Nakonieczna J. (2013) Innovative strategies to overcome biofilm resistance. *BioMed Research International*, **1**, Article ID 150653, 1-13. Available online: <http://dx.doi.org/10.1155/2013/150653/> (accessed on 10 June 2013).
- [17] Quave CL, Plano LR, Pantuso T, Bennett BC. (2008) Effects of extracts from Italian medicinal plants on planktonic growth, biofilm formation and adherence of methicillin-resistant *Staphylococcus aureus*. *Journal of Ethnopharmacology*, **118**, 418-428.
- [18] Wojnicz D, Kucharska AZ, Sokol-Letowska A, Kicia M, Tichaczek-Goska D. (2012) Medicinal plants extracts affect virulence factors expression and biofilm formation by the uropathogenic *Escherichia coli*. *Urological Research*, **40**, 683-697.
- [19] Estrela AB, Abraham WR. (2010) Combining biofilm-controlling compounds and antibiotics as a promising new way to control biofilm infections. *Pharmaceuticals*, **3**, 1374-1393.
- [20] Strelkova EA, Zhurina MV, Plakunov VK, Belyaev SS. (2012) Stimulation of biofilm formation by antibiotics. *Microbiology*, **81**, 259-262.
- [21] Knobloch JK, Bartscht K, Sabottke A, Rohde H, Feucht HH, Mack D. (2001) Biofilm formation by *Staphylococcus epidermidis* depends on functional RsbU, an activator of the sigB operon: differential activation mechanisms due to ethanol and salt stress. *Journal of Bacteriology*, **183**, 2624-2633.
- [22] Beauregard PB, Chai Y, Vlamakis H, Losick R, Kolter R. (2013) *Bacillus subtilis* biofilm induction by plant polysaccharides. *Proceedings of the National Academy of Sciences of the United States of America*, **110**, 1621-1630.
- [23] Vacheva A, Mustafa B, Staneva J, Marhova M, Kostadinova S, Todorova M, Ivanova R, Stoitsova S. (2011) Effects of extracts from medicinal plants on biofilm formation by *Escherichia coli* urinary tract isolates. *Biotechnology & Biotechnological Equipment*, **25**, Suppl., 92-97.
- [24] CLSI. (2009) Performance standards for antimicrobial disk susceptibility tests; Approved standard - Tenth edition. CLSI Document M2-A10. Clinical and Laboratory Standards Institute. Wayne, Pennsylvania, USA.
- [25] CLSI. (2010) Performance standards for antimicrobial susceptibility testing: 20th informational supplement. CLSI document M100-S20. Clinical and Laboratory Standards Institute. Wayne, Pennsylvania, USA.
- [26] CLSI. (2009) Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; Approved standard - Eighth edition. CLSI Document M07-A8. Clinical and Laboratory Standards Institute. Wayne, Pennsylvania, USA.
- [27] Stepanovic S, Vukovic D, Hola V, di Bona Ventura G, Djukic S, Cirkovic I, Ruzicka F. (2007) Quantification of biofilm in microtiter plates: overview of testing conditions and practical recommendations for assessment of biofilm production by staphylococci. *Acta Pathologica, Microbiologica et Immunologica Scandinavica [B]*, **115**, 891-899.
- [28] Cernohorska L, Votava M. (2004) Determination of minimal regrowth concentration (MRC) in clinical isolates of various biofilm-forming bacteria. *Folia Microbiologica*, **49**, 75-78.

New Butenolide and Pentenolide from *Dysidea cinerea*

Phan Van Kiem^{a,*}, Nguyen Xuan Nhiem^a, Ngo Van Quang^c, Chau Van Minh^a, Nguyen Hoai Nam^a, Nguyen Thi Cuc^a, Hoang Le Tuan Anh^a, Bui Huu Tai^a, Pham Hai Yen^a, Nguyen Xuan Cuong^a, Nguyen Phuong Thao^a, Nguyen Thi Hoai^d, Nan Young Kim^b, Seon Ju Park^b and Kim Seung Hyun^{b,*}

^aInstitute of Marine Biochemistry, Vietnam Academy of Science and Technology (VAST), 18 Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam

^bCollege of Pharmacy, Yonsei Institute of Pharmaceutical Sciences, Yonsei University, Incheon 406-840, Korea

^cInstitute of Chemistry, VAST, 18 Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam

^dFaculty of Pharmacy, Hue University of Pharmacy and Medicine, Hue, Vietnam

phankiem@yahoo.com; kimsh11@yonsei.ac.kr

Received: September 26th, 2013; Accepted: October 10th, 2013

Two new butenolide and pentenolide derivatives, dysideolides A-B, were isolated from the marine sponge *Dysidea cinerea*. Their structures were determined by the combination of spectroscopic and chemical methods, including 1D- and 2D-NMR spectroscopy, and CD spectra, as well as by comparing with the NMR data reported in the literature.

Keywords: *Dysidea cinerea*, Dysideidae, Dysideolide A, Dysideolide B, Marine sponge, Circular dichroism.

The marine sponge *Dysidea cinerea* Keller (Dysideidae) is distributed in the Red Sea, Western India, and Zanzibar. Marine sponges are a prolific source of anti-HIV proteins and study on chemical components of *D. cinerea* has led to the isolation of avarol and avarone derivatives [1a]. Several biological activities, such as anti-cytotoxic activity and inhibition of human immunodeficiency virus type 1 reverse transcriptase [1b,c], have also been reported. In the course of study of the chemical constituents of *D. cinerea*, butenolide and pentenolide derivatives were isolated (Figure 1).

Compound **1** was isolated as colorless oil and its molecular formula was determined as C₁₃H₂₂O₃ from the ion at *m/z* 225.1490 [M-H]⁻ in the HR ESI MS (Calcd. for C₁₃H₂₁O₃: 225.1496). The ¹H NMR spectrum of **1** showed: one olefinic proton at δ_H 5.84, two oxymethine protons at δ_H 3.86 and 4.91, and one tertiary methyl group at δ_H 2.10. The ¹³C NMR and DEPT spectra of **1** revealed 13 carbon signals, including one carbonyl (δ_C 176.3), one quaternary (δ_C 170.6), three methine (δ_C 70.0, 88.9, and 118.0), six methylene (δ_C 23.7, 27.2, 30.4, 30.6, 33.0, and 34.9), and two methyl carbons (δ_C 14.0 and 14.4) (Table 1). The ¹H and ¹³C NMR data of **1** were very similar to those of 4,10-dihydroxy-10-methyl-dodec-2-en-1,4-olide, except for the position of the hydroxyl group at C-5 [2]. The long range correlations between H-4 (δ_H 4.91) and C-1 (δ_C 176.3), C-2 (δ_C 118.0), C-3 (δ_C 170.60, and 3-Me (δ_C 14.0); between 3-Me (δ_H 2.10) and C-2 (δ_C 118.0), C-3 (δ_C 170.6), and C-4 (δ_C 88.9) in the HMBC spectrum (Figure 2) suggested that compound **1** had an α,β-unsaturated lactone and the methyl group was located at C-3. Furthermore, the hydroxyl group at C-5 was confirmed by the HMBC correlation between H-5 (δ_H 3.86) and C-3 (δ_C 170.6), C-4 (δ_C 88.9), and C-6 (δ_C 34.9), and by COSY correlations between H-4 (δ_H 4.91) and H-5 (δ_H 3.86); and between H-5 (δ_H 3.86) and H-6 (δ_H 1.62). The absolute configuration of **1** was determined by circular dichroism. The CD spectrum of **1** showed a positive peak at 207 nm, consistent with those of compound **1a** (a positive peak at 203 nm) [3a]. Therefore, the stereochemistry at C-4 of **1** was proved to be *R*. The remaining configuration at C-5 was determined by the advanced Mosher's method [3b]. Treatment of **1** with (*S*)- and




Figure 1: Structures of compounds **1** – **2** and reference compounds.


(*R*)-*R*-methoxy-*R*-(trifluoromethyl)phenylacetyl (MTPA) chlorides using catalytic DMAP afforded (*S*)- and (*R*)-MTPA esters (**1s** and **1r**), respectively. Analysis of the proton chemical shift differences between its (*S*)- and (*R*)-MTPA esters showed negative Δδ_H signs for H-2 (-0.03), H-4 (-0.06), and 3-Me (-0.03), and positive signs for H-5 (+0.03), and H-12 (+0.02). This distribution of Δδ signs enabled assignment of the *R* configuration to C-5. Consequently, compound **1** was elucidated to be (4*R*,5*R*)-4,5-dihydroxy-3-methyldodec-2-en-1,4-olide, and named dysideolide A.

Compound **2** was isolated as colorless oil and its molecular formula was determined to be C₁₃H₂₂O₃ by the HRESIMS ion at *m/z* 225.1491 [M-H]⁻ (Calcd. for C₁₃H₂₁O₃: 225.1496). The ¹H NMR spectrum of **2** showed: one olefinic proton at δ_H 5.78, two oxymethine protons at δ_H 4.07 and 4.22, and one tertiary methyl group at δ_H 2.03. The ¹³C NMR and DEPT spectra of **2** revealed 13 carbon signals, including one carbonyl (δ_C 163.5), one quaternary (δ_C 158.9), three methine (δ_C 69.4, 81.8, and 116.7), six methylene (δ_C 22.6, 24.9, 29.1, 29.3, 31.7, and 32.7), and two methyl carbons (δ_C 14.1 and 19.1). The ¹H- and ¹³C NMR data of **2** were very similar to those of dysideolide A, except for the butenolide group replaced by a pentenolide group. The HMBC correlations between H-5 (δ_H 4.22) and C-3 (δ_C 158.9), C-4 (δ_C 69.4), and C-7 (δ_C 24.9); between 3-Me (δ_H 2.03) and C-2 (δ_C 116.7), C-3 (δ_C 158.9), and C-4 (δ_C 69.4) suggested that compound **2** had an α,β-unsaturated-δ-lactone and the methyl group was at C-3. In addition, the hydroxyl

Table 1: NMR spectral data for compounds 1–2.

Pos.	1		2	
	$\delta_C^{a,c)}$	$\delta_H^{a,d)}$ (mult., J in Hz)	$\delta_C^{b,c)}$	$\delta_H^{b,d)}$ (mult., J in Hz)
1	176.3	-	163.5	-
2	118.0	5.84 (s)	116.7	5.78 (s)
3	170.6	-	158.9	-
4	88.9	4.91 (s)	69.4	4.07 (d, 7.6)
5	70.0	3.86 (t, 6.4)	81.8	4.22 (dt, 3.6, 7.6)
6	34.9	1.62 (m)	32.7	1.67 (m), 1.75 (m)
7	27.2	1.37, 1.51	24.9	1.41, 1.55
8	30.6	1.31	29.3	1.27
9	30.4	1.31	29.1	1.27
10	33.0	1.29	31.7	1.23
11	23.7	1.30	22.6	1.25
12	14.4	0.88 (t, 6.5)	14.1	0.86 (t, 6.5)
3-Me	14.0	2.10 (s)	19.1	2.03 (s)

^{a)}CD₃OD, ^{b)}CDCl₃, ^{c)}100 MHz, ^{d)}400 MHz. Assignments were made by HMQC, HMBC, and COSY experiments.

**Figure 2:** Key HMBC and COSY correlations of 1–2.

group at C-4 was confirmed by the HMBC correlations between H-4 (δ_H 4.07) and C-2 (δ_C 116.7), C-5 (δ_C 81.8), and C-6 (δ_C 32.7), and COSY correlations between H-4 (δ_H 4.07) and H-5 (δ_H 4.22) and H-6 (δ_H 1.67 and 1.75). The absolute configuration at C-5 of **2** was fixed by CD measurements and comparison with **2a** [3c]. The CD spectrum of **2** showed a positive peak at 219 nm, while compound **2a** showed a negative peak at 221 nm (5*S* configuration) [3c]. Based on this evidence, the stereochemistry at C-5 was proved to be *R*. Moreover, the coupling constant between H-4 and H-5 in the hexacyclic ester of **2**, $J_{4,5} = 7.6$ Hz, confirmed the *threo* of the both protons H-4 and H-5 by comparing coupling constant of two *erythro* protons of **2a**, $J_{4,5} = 3.0$ Hz [3c] and *threo* protons of **2b**, $J_{4,5} = 8.8$ Hz [3d]. Based on the above evidence, compound **2** was elucidated as (4*S*,5*R*)-4,5-dihydroxy-3-methyldeca-2-en-1,5-olide, and named dysideolide B.

Experimental

General: The NMR spectra were recorded on an Agilent 400-MR spectrometer using TMS as internal standard. The HRESIMS were obtained using an AGILENT 6550 iFunnel Q-TOF LC/MS system. Optical rotations were determined on a Jasco DIP-370 automatic polarimeter. Circular dichroism spectra were determined on a ChirascanTM CD spectrometer. Preparative HPLC was carried out using an AGILENT 1200 HPLC system. CC was performed using either silica-gel (Kiesel gel 60, 70–230 mesh and 230–400 mesh, Merck) or YMC RP-18 resin (30–50 μ m, Fuji silica Chemical Ltd.), and TLC by using pre-coated silica-gel 60 F254 (0.25 mm, Merck) and RP-18 F254S plates (0.25 mm, Merck).

References

- (a) Hirsch S, Rudi A, Kashman Y, Loya Y. (1991) Newavarone and avarol derivatives from the marine sponge *Dysidea cinerea*. *Journal of Natural Products*, **54**, 92–97; (b) Rudi A, Yosief T, Schleyer M, Kashman Y. (1999) Bilosipens A and B: Two novel cytotoxic sesterpenes from the marine sponge *Dysidea cinerea*. *Organic Letters*, **1**, 471–472; (c) Loya S, Hizi A. (1990) The inhibition of human immunodeficiency virus type 1 reverse transcriptase by avarol and avarone derivatives. *FEBS Letters*, **269**, 131–134.
- Mukku VJR, Speitling M, Laatsch H, Helmke E. (2000) New butenolides from two marine *Streptomyces*. *Journal of Natural Products*, **63**, 1570–1572.
- (a) Gawronski JK, van Oeveren A, van der Deen H, Leung CW, Feringa BL. (1996) Simple circular dichroism method for the determination of absolute configuration of 5-substituted 2(5H)-furanones. *The Journal of Organic Chemistry*, **61**, 1513–1515; (b) Nhiem NX, Kiem PV, Minh CV, Kim N, Park S, Lee HY, Kim ES, Kim YH, Kim S, Koh Y-S, Kim SH. (2013) Diarylheptanoids and flavonoids from *Viscum album* inhibit LPS-stimulated production of pro-inflammatory cytokines in bone marrow-derived dendritic cells. *Journal of Natural Products*, **76**, 495–502; (c) Yu YM, Yang JS, Peng CZ, Cao V, Cong PZ, Zou ZM, Lu Y, Yang SY, Gu YC. (2009) Lactones from *Angiopteriscaudatiformis*. *Journal of Natural Products*, **72**, 921–924. (d) Buchanan MS, Hashimoto T, Takaoka S, Asakawa Y. (1995) (+)-Osmundalactone, γ -lactones and spiromentins from the fungus *Paxillus atrotomentosus*. *Phytochemistry*, **40**, 1251–1257.

Sponge material: *D. cinerea* was collected in Lang Co beach, Vietnam during August 2011, and identified by Dr Do Cong Thung, Institute of Marine Resources and Environment. A voucher specimen (DC1108) was deposited at the Herbarium of the Institute of Marine Biochemistry, VAST.

Extraction and isolation: Freeze dried tissue of *D. cinerea* (1.0 kg) was cleaned to remove sodium chloride and then extracted 3 times with MeOH under reflux for 15 h to yield 60 g of a dark solid extract. This was suspended in water and partitioned with CHCl₃ to obtain CHCl₃ (DC1, 20.0 g) and water extracts (DC2, 40 g) after removal of the solvents *in vacuo*. The DC1 extract (20.0 g) was chromatographed on a silica gel column eluting with a gradient of *n*-hexane – acetone (40 : 1 \rightarrow 0 : 1, v/v) to obtain 5 sub-fractions, DC1A (3.2 g), DC1B (4.5 g), DC1C (3.3g), DC1D (3.1 g), and DC1E (5.0 g). The sub-fraction DC1B (4.5 g) was chromatographed on a silica gel column eluting with *n*-hexane – EtOAc (10 : 1, v/v) to give 3 smaller fractions, DC1B1 (1.0 g), DC1B2 (1.5 g), and DC1B3 (1.7 g). Fraction DC1B2 was chromatographed on an YMC RP-18 column eluting with MeOH – water (5 : 1, v/v) to give 3 fractions, DC1B2A (450 mg), DC1B2B (200 mg), and DC1B2C (400 mg). Fraction DC1B2B was chromatographed by HPLC using J'sphere ODS H-80, 250 mm \times 20 mm, and 35% acetonitrile in water at a flow rate of 3 mL/min to yield **1** (7.0 mg) and **2** (6.0 mg).

Dysideolide A (1)

Colorless oil.

$[\alpha]_D^{25}$: +45 (*c* 0.1, MeOH);

CD $[\theta]_{207}^{25}$: $+2.2 \times 10^4$ (*c* 5.0×10^{-4} , MeOH).

¹H and ¹³C NMR (CD₃OD): Table 1.

HRESIMS found *m/z*: 225.1490 [M – H][–] (Calcd. for C₁₃H₂₁O₃: 225.1496).

(S)-MTPA ester of 1 (1s)

¹H NMR (500 MHz, CDCl₃): δ_H 5.86 (H-2), 4.79 (H-4), 4.61 (H-5), 0.89 (H-12), and 2.13 (3-Me).

(R)-MTPA ester of 1 (1r)

¹H NMR (500 MHz, CDCl₃): δ_H 5.89 (H-2), 4.85 (H-4), 4.58 (H-5), 0.87 (H-12), and 2.16 (3-Me).

Dysideolide B (2)

Colorless oil.

$[\alpha]_D^{25}$: +35 (*c* 0.1, MeOH).

CD $[\theta]_{219}^{25}$: $+2.2 \times 10^4$ (*c* 5.0×10^{-4} , MeOH).

¹H and ¹³C NMR (CDCl₃, 400 MHz): Table 1.

HRESIMS found *m/z*: 225.1491 [M – H][–] (Calcd. for C₁₃H₂₁O₃: 225.1496).

Acknowledgments - This study was financially supported by Vietnam 02/2011/HD-NCCBUD Project and partially supported by the National Research Foundation of Korea (NRF).

A New Cyclopeptide from Endophytic *Streptomyces* sp. YIM 64018Xueqiong Yang^{a,1}, Yabin Yang^{a,1}, Tianfeng Peng^a, Fangfang Yang^a, Hao Zhou^a, Lixing Zhao^b, Lihua Xu^b and Zhongtao Ding^{a,*}^aKey Laboratory of Medicinal Chemistry for Natural Resource, Ministry of Education, School of Chemical Science and Technology, Yunnan University, Kunming 650091, China^bYunnan Institute of Microbiology, Yunnan University, Kunming 650091, China¹The authors contributed equally to this paper


ztding@ynu.edu.cn

Received: September 25th, 2013; Accepted: October 17th, 2013


One new cyclopeptide, cyclo(L-Phe-L-Ala-L-Phe-Gly), named as vinaceuline (**1**) and three known cyclodipeptides, cyclo (Phe-Gly), cyclo (Phe-4-hydroxyl-Pro) and cyclo (Phe-Ile) were isolated from broth culture of endophytic *Streptomyces* YIM 64018 associated with *Paraboea sinensis*. The planar structure of the new compound was assigned on the basis of 1D and 2D NMR spectroscopic techniques, while the absolute configurations of the amino acid residues were determined by application of the advanced Marfey method. Cyclotetrapeptides are rarely found as *Streptomyces* metabolites.

Keywords: *Streptomyces* sp., Cyclopeptide, Vinaceuline.

Streptomyces are believed to be a rich source of new and useful compounds, and up to 40% of known microbial metabolites are derived from this group. Endophytes are also considered as important natural sources of a variety of secondary metabolites [1,2]. New cyclopeptides with antimicrobial and anticancer activities have been reported from different bacterial genera [3,4]. In this work, chemical investigation of the endophytic *Streptomyces* sp. (YIM64018) was carried out for its antimicrobial activity against *Escherichia coli*. One new cyclopeptide and three known cyclodipeptides were isolated, and their structures were determined as cyclo(L-Phe-L-Ala-L-Phe-Gly) (**1**), named as vinaceuline, cyclo (Phe-Gly)[5], cyclo (Phe-4-hydroxyl-Pro)[5] and cyclo (Phe-Ile)[6] (Figure 1).

Figure 1: Structure of compound **1**.

The molecular formula of compound **1** was determined as C₂₃H₂₆N₄O₄ from its DEPT spectrum and HRESI-MS (*m/z* 423.2069 for [M+H]⁺), with 13 degrees of unsaturation. The ¹³C NMR spectrum showed the presence of four amide CO at δ 168.8, 168.2, 167.1 and 166.5, three CH at δ 57.1, 56.8 and 51.1, three CH₂ at δ 44.8, 40.6 and 40.3, and one CH₃ at δ 20.5. The ¹H NMR spectrum showed the presence of four amide NH at δ 9.35 (1H, br s), 9.18 (2H, br s), and 9.06 (1H, br s), three methine protons at δ 4.63 (1H, br s), 4.58 (1H, br s), and 4.17 (1H, t, *J* = 5.7 Hz), three methylene signals at δ 3.95, 3.52 (2H), 3.45 (2H), and 3.28 (2H), one methyl group at δ 1.09 (3H, d, *J* = 7.0 Hz). These facts indicated that **1** is a cyclotetrapeptide. The amino acid units were identified by ¹H-¹H COSY, HSQC and HMBC experiments as two phenylalanine groups (Phe), one alanine group (Ala) and one glycine group (Gly) (Figure 2).

Figure 2: Key HMBC and ¹H-¹H COSY correlations of compound **1**.

The sequence of the amino acid residues in **1** was established by analysis of the HMBC (Figure 2) and ROESY data. The HMBC spectrum showed correlations between δ c 168.2 (Phe (1)-CO) and δ H 3.95 (α H- Gly), δ c 168.8 (Ala-CO) and δ H 9.18 NH (Phe (1)), and δ c 167.1 (Phe (2)-CO) and δ H 4.17 (α H- Ala). The ROESY spectrum also revealed correlations between δ H 9.35 NH {(Phe (2))} and 9.06 NH (Gly), and 9.18 NH (Ala). This evidence indicated the presence of a tetrapeptide fragment as -Gly-Phe-Ala-Phe-. The appearance of fragments of Gly-Phe or Phe-Gly, Phe-Ala or Ala-Phe in ESI-MS analyses at *m/z* 205 [M+H]⁺ and 219 [M+H]⁺ also confirmed this proposed linkage pattern. To determine the absolute configurations of the amino acid constituents in **1**, the advanced Marfey method was applied [7]. All derivatives were identified by their retention times and molecular weights. The configuration of the Phe unit was determined by comparing the retention times of LD-derivatized hydrolysate (27.94 min and 39.49 min) with the L-derivatized hydrolysate (27.86 min). The configuration of the Ala unit was determined by comparing the retention times of LD-derivatized hydrolysate (14.25 min and 20.65 min) with the L-derivatized hydrolysate (14.52 min). Therefore the structure of compound **1** was determined as cyclo (Gly-L-Phe-L-Ala-L-Phe) and was named vinaceuline (**1**).

Three known compounds were determined as cyclo (Phe-Gly), cyclo (Phe-4-hydroxyl-Pro) and cyclo (Phe-Ile) by spectroscopic and ESIMS analysis, and by comparison with references [5-6].

Experimental

General experimental procedures: Silica gel (100-200, 200-300 mesh; Qingdao Marine Chemical Group Co.) and Sephadex LH-20 (GE Healthcare Co) were used for CC. 1D and 2D NMR spectra were obtained on Bruker DRX-600 MHz and Bruker DRX-500 MHz instruments with TMS as internal standard. MS were recorded with Agilent G3250AA, AutoSpec Premier P776 and Waters Xevo TQ-S mass spectrometers.

Biological material and cultivation of actinomycetic strain: The bacterial strain was isolated using trehalose-proline medium from the plant *Paraboea sinensis*, and was assigned as *Streptomyces vinaceus* by characterization and complete 16S rRNA gene sequence. The strain has been preserved at Yunnan Institute of Microbiology, Yunnan University, China. This bacterium was cultivated on a 40 L scale using 1L Erlenmeyer flasks containing 250 mL of the seed medium (yeast extract 0.4%, glucose 0.4%, malt extract 0.5%, decavitamin 0.01%, pH 7) for 3 days and the fermentation medium (soluble starch 1%, glucose 1%, peptone 0.5%, NaCl 0.4%, 0.05% K₂HPO₄, MgSO₄·7H₂O 0.05%, CaCO₃ 0.2%, pH 7.0) for 5 days at 28°C on a rotary shaker (250 rpm).

Extraction and isolation of compounds: The fermentation broth (40L) containing *Streptomyces* sp. YIM 64018 was filtered, the filtrate extracted with EtOAc and the solvent removed under vacuum. The EtOAc extract (11 g) was separated into 4 fractions by CC on silica gel (200–300 mesh), eluting stepwise with CHCl₃/MeOH gradient (CHCl₃, CHCl₃/MeOH: 10:1, v/v, CHCl₃/MeOH: 3:1, v/v, MeOH). The second fraction (eluted with CHCl₃/MeOH: 10:1, v/v) was separated by CC on silica gel eluting stepwise with a CHCl₃/MeOH gradient (from 30:1 to 2:1, v/v) to produce 5 fractions (Fr.1~Fr.5). Fr.2 was further purified by repeated CC on silica gel eluting stepwise with a CHCl₃/MeOH gradient (from 20:1 to 1:1, v/v) and Sephadex LH-20 (MeOH) to afford **1** (4.6 mg), cyclo (Phe-Gly) (6.2 mg), cyclo (Phe-4-hydroxyl-Pro) (2.7 mg), and cyclo (Phe-Ile) (5.1mg).

Acid hydrolysis and advanced Marfey analysis: A 1.0 mg sample of compound **1** was subjected to acid hydrolysis at 110°C for 24 h with 6 N HCl (1.0 mL), and then the hydrolysates were evaporated to dryness and resuspended in acetone (100 µL). To each half portion (50 µL) was added 20 µL NaHCO₃ (1 M) and either 100 µL 1-fluoro-2,4-dinitrophenyl-5-L-leucinamide or 1-fluoro-2,4-dinitrophenyl-5-D-leucinamide (L-FDLA or D-FDLA, 1% in acetone), and the mixture was heated at 40°C for 1.5 h. The reaction

mixtures were cooled, neutralized with 2 N HCl (10 µL), dried and dissolved in 50% aqueous MeCN. About 5 µL of each solution of FDLA derivatives was analyzed by LC/MS.

The analysis of the L- and LD-FDLA (mixture of D- and L-FDLA) derivatives was performed using an Agilent Eclipse XDB-C₁₈ column (4.6×150 mm, 5 µm) maintained at 40°C. Aqueous MeCN containing 0.01 M methanoic acid was used as mobile phase with a linear gradient elution mode (MeCN, 30-50% for 50 min) at a flow rate of 1.0 mL/min. A Waters Xevo TQ-S mass spectrometer was used for detection in ESI (negative) mode. The capillary voltage was kept at 2.5 KV, and the ion source at 350°C. Nitrogen gas was used as a sheath gas at 400 L/h. A mass range of m/z 100-1000 was scanned in 0.2 s.

Compound 1

white crystals (CHCl₃).

[α]_D²⁵ 21.3 (c 0.1, DMSO).

¹H NMR (C₅D₅N, 500 MHz) δ: 9.35 (1H, br s, Phe(2)-NH), 9.18 (1H, br s, Phe(1)-NH), 9.18 (1H, br s, Ala-NH), 9.06 (1H, br s, Gly-NH), 7.44 (2H, d, J = 8.0 Hz, Phe (1)-H-2,6), 7.43 (2H, d, J = 8.0 Hz, Phe (2)-H-2, 6), 7.25 (2H, d, J = 8.0 Hz, Phe (1)-H-3, 5), 7.23 (2H, d, J = 8.0 Hz, Phe (2)-H-3, 5), 7.22 (1H, d, J = 8.0 Hz, Phe (1)-H-4), 7.20 (1H, d, J = 8.0 Hz, Phe (2)-H-4), 4.63 (1H, br s, Phe (1)-αH), 4.58 (1H, br s, Phe (2)-αH), 4.17 (1H, q, J = 5.7Hz, Ala-αH), 3.95 (1H, dd, J = 3.0, 17.0 Hz, Gly-αH), 3.52 (1H, m, Gly-αH), 3.45 (2H, m, Phe (1,2)-βH), 3.28 (2H, m, Phe (1,2)-βH), 1.09 (3H, d, J = 7.0Hz, Ala-βH).

¹³C NMR (C₅D₅N, 125 MHz) δ: 168.8 (Ala-CO), 168.2 (Phe(1)-CO), 167.1 (Phe(2)-CO), 166.5 (Gly-CO), 136.9 (Phe (1)-C-1), 136.7 (Phe (2)-C-1), 130.7 (Phe(1)-C-2,6), 130.5 (Phe(2)-C-2,6), 128.5 (Phe(1)-C-3,5), 128.5 (Phe(2)-C-3,5), 127.0 (Phe(1)-C-4), 126.9 (Phe(2)-C-4), 57.1 (Phe (1)-αC), 56.8 (Phe (2)-αC), 51.1 (Ala-αC), 44.8 (Gly-αC), 40.6 (Phe (1)-βC), 40.3 (Phe (2)-βC), 20.5 (Ala-βC).

HRESI-MS: m/z [M+H]⁺ calcd for C₂₃H₂₇N₄O₄: 423.2026; found: 423.2069.

Acknowledgments - This project was supported by the State Program of National Natural Science Foundation of China (No. 81060263; No. 81360480). The advanced Marfey method was conducted with the guidance of Prof. Ninghua Tan and Miss Huiming Xu in Kunming Institute of Botany, the Chinese Academy of Sciences, P. R. China.

References

- [1] Genilloud O, González I, Salazar O, Martín J, Tormo JR, Vicente F. (2011) Current approaches to exploit actinomycetes as a source of novel natural products. *Journal of Industrial Microbiology & Biotechnology*, **38**, 375–389.
- [2] Zhang Y, Han T, Ming Q, Wu L, Rahman K, Qin L. (2012) Alkaloids produced by endophytic fungi: a review. *Natural Product Communications*, **7**, 963–968.
- [3] Romano A, Vitullo D, Di Pietro A, Lima G, Lanzotti V. (2011) Antifungal lipopeptides from *Bacillus amyloliquefaciens* strain BO7. *Journal of Natural Products*, **74**, 145–151.
- [4] Kanoh K, Matsuo Y, Adachi K, Imagawa H, Nishizawa M, Shizuri Y. (2005) Mechercharmycins A and B, cytotoxic substances from marine-derived *Thermoactinomyces* sp. YM3-251. *The Journal of Antibiotics*, **58**, 289–292.
- [5] Li DH, Gu QQ, Zhu WM, Liu HB, Fang YC, Zhu TJ. (2005) Antitumor components from marine actinomycete 11014 I. cyclic dipeptides. *Chinese Journal of Antibiotics*, **30**, 449–452.
- [6] Wang YC, Zhou J, Tan NH, Ding ZT, Jiang X. (1999) Cyclic dipeptides from *Schizandra chinensis* and their syntheses. *Acta Pharmaceutica Sinica*, **34**, 19–22.
- [7] Fujii K, Ikai Y, Oka H, Suzuki M, Harada K. (1997) A nonempirical method using LC/MS for determination of the absolute configuration of constituent amino acids in a peptide: combination of Marfey's method with mass spectrometry and its practical application. *Analytical Chemistry*, **69**, 5146–5151.

Involvement of Trypsin-Digested Silk Peptides in the Induction of RAW264.7 Macrophage Activation

Kyoung-Ho Pyo^a, Min-Ki Kim^a, Kwang-Soon Shin^b, Hyang Sook Chun^c and Eun-Hee Shin^{a,d,*}

^aDepartment of Parasitology and Tropical Medicine, Seoul National University College of Medicine and Institute of Endemic Diseases, Seoul National University Medical Research Center, Seoul 110-799, Republic of Korea

^bDepartment of Food science and Biotechnology, Kyonggi University, Suwon-si, 443-760, Republic of Korea

^cDepartment of Food Science & Technology, Chung-Ang University, Ansong 456-756, South Korea

^dSeoul National University Bundang Hospital, Seongnam 463-707, Republic of Korea

ehshin@snu.ac.kr

Received: April 4th, 2013; Accepted: August 16th, 2013

The activation of macrophages by trypsin-digested silk peptides was investigated by considering CD11b and CD40 expression in the RAW264.7 cell, a murine macrophage. Silk protein hydrolysates were digested with trypsin, following by centrifugal purification using the Centrprep 30k concentrator. Trypsin-digested total silk peptides and its centrifugal fractions were tested for macrophage activation *in vitro*. The functional peptide of fractionated silk peptides was examined by LC/MS/MS analysis. Trypsin-digested and fractionated silk peptides of more than 30 kDa induced an increase in the activation markers CD11b and CD40 in RAW264.7 cells. These results are supported by morphological changes reflecting an increase in the number of dendrites in activated cells. The fractionated silk peptides examined by LC/MS/MS contained partial peptides of *Bombyx mori* fibroin. These results suggest that the activation of RAW264.7 macrophages may be induced not by sericin-derived peptides but by fibroin-derived ones.

Keywords: Silk peptide, Trypsin, Fibroin, Immune response, Macrophage, CD11b, CD40.

Bombyx mori is a domesticated silkworm species, and its cocoon silk has been an important textile source [1a]. In recent years, there has been a renewal of interest in the usefulness of silk as a bioactive material [1a,b]. Silk protein produced by *B. mori* consists of two major proteins: fibroin and sericin [1c]. Previous studies have emphasized the biological usefulness of sericin because of its moisture-absorbing, antioxidant, and wound-healing properties [1d, 2a]. In contrast, previous studies have focused on fibroin for its use in tissue engineering scaffolds, wound dressings, and vascular grafts [1d,2b,c]. Recent studies have demonstrated some bio-efficacy of silk fibroin, including its ability to reduce blood pressure, its anti-tumor activity, and for treatment of atopic dermatitis [3,4a,b]. In addition, silk protein has been used to enhance the immune response, ameliorate the progression of skin lesions resembling atopic dermatitis, and facilitate insulin-releasing activity through the induction of β -cell activity in C57BL/KsJ-db/db mice [4b,c,5a].

Recently, an increase in the immune response in mice treated with silk protein has been demonstrated, namely an antiprotozoal effect on *Toxoplasma gondii*, a zoonotic protozoan [4c]. The protective immune response to *T. gondii* usually entails the activation of macrophages as well as T cells [5b]. A macrophage is an antigen-presenting cell (APC) and plays an important role in innate immunity and helper T-cell immune responses [5c,6]. The activation of macrophages to microbial ligands results in an increase in surface molecules such as CD11b and CD40 [5c,6]. CD11b is an activation marker of macrophages and is expressed on the surface of monocytes, granulocytes, macrophages, and natural killer cells. CD40 is a co-stimulatory molecule in APCs and is an essential mediator of T-cell activation for the secretion of IL-12 and TNF- α [5c,6]. In this regard, the present study identifies the specific fraction in silk peptides not only for the activation of macrophages as effector cells for an innate immune response but also APCs for an immune response mediated by T cells. For this purpose, this

study examines two biomarkers on the surface of the RAW264.7 cell, a murine macrophage —CD11b and CD40—which increase during the activation of macrophages.

With LPS treatment, the expression of CD11b and CD40 in RAW264.7 cells increased by 58.9% and 86.6%, respectively (Figure 1A and 1F). For an analysis of the synergistic effect of LPS and silk peptides (TSP) on the expression of CD11b and CD40, LPS and TSP were co-treated during cell culture. The results indicate that CD11b expression was slightly lower in co-treated cells (49.7%) than in LPS (58.9%), suggesting no synergetic effect of LPS and TSP on CD11b expression. By contrast, TSP treatment induced a 24.2% increase in CD11b expression in RAW264.7 cells. Similarly, FSP>30 treatment induced a 26.3% increase in CD11b expression, and FSP<30 treatment, a 3.8% increase. In terms of CD40 expression, TSP treatment induced a 30.9% increase in CD40 expression, and FSP>30 treatment, a 36.5% increase. LPS treatment induced an 86.6% increase in CD40 expression, and co-treatment of LPS and TSP, a 77.9% increase. There was no synergistic effect of LPS and TSP, and FSP<30 had no effect on CD40 expression (Figure 1F-J).

For an analysis of morphological changes in RAW264.7 cells after silk peptide treatment, the cells were cultured with either peptides (TSP, FSP>30, or FSP<30) or LPS for 24 h in six-well plates (Figure 2). When RAW264.7 cells were not stimulated, they had no dendrites, which increase in activated macrophages (Figure 2A). However, those cells treated with LPS had many dendrites at 24 h after culture (Figure 2B). Similarly, the addition of either TSP or FSP>30 induced the production of many dendrites (Figure 2D and 2E).

By contrast, FSP<30 treatment did not induce the formation of dendrites (Figure 2F), and the morphology was similar to that of normal RAW264.7 cells (Figure 2A).




Figure 1: CD11b and CD40 expression in RAW264.7 cells after treatment with silk peptides. Activation markers of macrophages—CD11b and CD40—were examined through a FACS analysis after cultivation for 24 h. (A & F) Designated LPS stimulation; (B & G) LPS + TSP; (C & H) TSP only; (D & I) FSP>30; (E & J) FSP<30.




Figure 2: Morphological changes in RAW264.7 cells after treatment with silk peptides. RAW264.7 cells were incubated for 24 h with LPS, LPS+TSP, FSP>30, or FSP<30. Then the cells were observed under a microscope for dendrites in RAW264.7 cells treated with each of the aforementioned cell stimulants. Arrows indicate macrophage dendrites. Cell images were taken at original magnification $\times 400$.




Figure 3: Verification of residual enzyme activity after the digestion of silk protein hydrolysates with trypsin. Silk protein hydrolysates were digested with trypsin, followed by enzyme inactivation at 95°C for 30 min. For an analysis of residual trypsin activity, BSA was added to the digestion mixture. In addition, proteolysis samples were loaded on a 15% polyacrylamide gel. M: Marker; Lane 1: BSA; Lane 2: BSA+Trypsin; Lane 3: BSA+Heat-inactivated trypsin; Lane 4: BSA+TSP (1:1); Lane 5: BSA+TSP (1:2); Lane 6: BSA+TSP (1:5).

Trypsin was used for the proteolysis of silk protein to augment the absorption of ingested proteins. For an analysis of residual trypsin activity, BSA was used (Figure 3). Lane 1 shows intact BSA (63 kDa), and lane 2, trypsin-digested BSA. Lane 3 shows undegraded BSA when heat-inactivated trypsin was added. Here, heat inactivation reduced the proteolytic activity of trypsin.

For an analysis of residual trypsin activity after the proteolysis of silk protein, the reaction product of silk protein with trypsin was mixed with BSA for further proteolysis in the ratio of 1:1, 1:2, or 1:5 for BSA to the reaction mixture of trypsin and silk peptides. The results indicate no further proteolysis of BSA in the reaction mixture of trypsin and silk peptides (lanes 4, 5, and 6). This implies that there was no residual trypsin activity in trypsin-digested silk peptides and thus there was no relationship between residual trypsin

Table 1: LC-MS/MS analysis of silk peptide fraction of FSP>30.

LC-MS/MS data	Sequences	score
Fibroin [Bombyx mandarina]	DASGAVIEEITTK	90
Putative reading frame (fibroin) [Bombyx mori]	DASGAVIEEITTK	89
Aspartic endopeptidase Pep1	MVVFVKVTAAVFLGATIASAAPPTK	80
Hypothetical protein	KKVSLNVWLALK	78
PREDICTED: similar to hCG1980490	EAGPPR	74
PREDICTED: similar to hCG1980490	ILQLPGAPDCPLRLVLAKGAEGEEAGSLP	74
Fibroin	WVTQMLEGNGSVSLLR	73
Fibroin	DIDDGK	73
Fibroin	SIAILNVQEILK	73
CRISPR system CASCADE complex protein CasC	QKALEVVKALGLK	71
Secretion protein HlyD	RGGGASGVHGASGARAASGAHGSGASQ	70
	QAQP	70

activity and the activation of RAW264.7 cells. The silk peptide fraction of FSP>30 was concentrated using Centriprep 30K and analyzed by LC-MS/MS. Peptide sequences were aligned and counted by plot scores by using MASCOT (Matrix Science, www.matrixscience.com). Ten peptides were selected from the list of top hits (Table 1). The scores ranged from 70 to 90, and other putative peptides such as Pep1, hCG, CasC, and HlyD as well as fibroin-derived peptides were also detected (Table 1). However, because the results of the MALDI-TOF analysis indicate a portion of some peptides, *B. mori* fibroin-derived sequences DIDDGK and SIAILNVQEILK were considered as a main ingredient of FSP>30. Here no sericin fragment was found in silk peptides of FSP>30. Although this result does not clearly suggest that the two targeted sequences are functional peptides, fibroin-derived peptides may be major peptides in FSP>30.

Previous studies have emphasized the usefulness of silk protein in textiles [1a]. However, there has been some concern over its biomedical and biomaterial applications such as cosmetics and functional foods because of its safety and functional characteristics [1b-d,2a,b, 3,4a-c,5a]. In this regard, silk protein has been known to have a favorable effect on diabetes, blood pressure, anti-tumor activity, skin lesions resembling atopic dermatitis, and antiprotozoal activity [3,4a-c,5a]. A recent study examined the ability of silk protein to induce immunity to infection or enhance the immune system by activating immune cells [4c]. When BALB/c mice ingested silk protein hydrolysates, they were immune to *T. gondii*, an intracellular protozoan, and this protective immunity was characterized by increases in CD4⁺- and CD8⁺-T cells, and a sharp increase in IFN- γ [4c]. These results suggest that silk protein can be used as a natural compound for enhancing immunity [4c]. As a result, there has been growing interest in the role of silk protein as a functional material for various applications in the biological, biomedical, and biotechnology fields.

Silk protein consists of fibroin and sericin, which have distinct characteristics and biological functions [7]. Fibroin is a glycoprotein composed of two equimolar protein subunits of 370 kDa and 25 kDa covalently linked by disulfide bonds [7]. Sericin contains 18 amino acids, including essential amino acids. Sericin accounts for approximately 20-30 % of total cocoon weight [7], and, therefore, *B. mori* silk protein hydrolysates prepared from CaCl_2 treatment contain 70-80% fibroin and 20-30% sericin. In a recent study, enzymatic degradation was used to extract bioactive peptides from silk protein [7,8,9a]. Collagenase IA, α -Chymotrypsin, Protease XIV, Protease P, Alkylase, Alkaline protease, Trypsin, Protease N, Alcalase 2.4L, As 1.398, and Neutrase have been used for the enzymatic degradation of silk protein. Among these, trypsin is a serine protease found in the digestive system of many vertebrates, where it hydrolyses proteins [9b]. Trypsin has been found to cleave the Arg-Gly bond toward the insulin β -chain at pH 8.0 [10a]. Although the enzymatic hydrolysis of silk protein has been attempted using various enzymes, trypsin is a safe endogenous enzyme used in the food industry. Tryptic hydrolysis has been well analyzed in the food science field to improve the quality of food products [10b]. In the present study, trypsin was used to increase the absorption of silk peptides by pre-performing the digestion process in the digestive tract. The results of the computer-based cleavage site prediction of endopeptidases with PeptideCutter (<http://www.expasy.org/tools/peptidecutter/>) [11a] indicate that the tryptic hydrolysis of fibroin produced peptides of various sizes from 146 Da to 325,525 Da. By contrast, the tryptic hydrolysis of sericin produced peptides ranging from 146 Da to 7,065 Da (data not shown). For the identification of the size-based fraction of silk peptides inducing an immune response, the fraction based on 30 kDa was obtained using the Centriprep 30k concentrator. A recent study found an increase in the immune enhancement of silk protein hydrolysates, namely an increase in T-cell responses [4c]. More specifically, the initial immune response mediated by T cells was induced and accelerated by APCs such as macrophages and dendritic cells. Macrophages play a critical role in innate as well as acquired immunity and can induce protective cellular immunity to *T. gondii* infection. In this regard, we used the RAW264.7 cell, a murine macrophage, to investigate the silk peptide fraction and examine the immune response because trypsin-digested silk peptides can activate macrophages and increase the expression of specific activation markers in macrophages.

LPS can activate macrophages through TLR4 (Toll-like receptor 4) [11b]. Macrophages activated with LPS can increase the expression of CD11b and CD40. In this regard, RAW264.7 cells can express the activation markers CD11b and CD40 in response to LPS [5a,b,6]. In the present study, LPS alone, and LPS with TSP induced a similar increase in CD11b and CD40 expression in RAW264.7 cells. This indicates no synergistic effect of silk peptides and LPS on CD11b and CD40 expression. TSP and FSP>30 induced 24.2% and 26.3% increases, respectively, in CD11b expression. However, FSP<30 induced only a negligible (3.8%) increase in CD11b expression. Similarly, TSP and FSP>30 induced 30.9% and 36.5% increases, respectively, in the expression of CD40, a co-activation marker. Morphological changes in RAW264.7 cells reflected the result in CD11b and CD40 expression. That is, when RAW264.7 cells were not stimulated, they had no dendrites, which generally increase in activated macrophages. However, RAW264.7 cells with LPS treatment had many dendrites at 24 h after treatment. Similarly, TSP and FSP>30 induced the production of many dendrites. However, FSP<30 did not induce dendrite formation. These results indicate that TSP and FSP>30 may activate macrophages through a mechanism different

from that of LPS because of a lack of any synergistic effect of LPS and TSP. When macrophages are activated, surface molecules such as TLRs, integrin, and activating receptors increase to recognize pathogens, and there is an increase in cell motility [11b]. Although the mechanism underlying the activation of RAW264.7 cells by silk peptides remains unclear, this study's results suggest that silk peptides may activate cells not through TLR 4 but through other activation receptors. As another mechanism, trypsin is a factor that can activate PAR2 (protease-activated receptor 2) and induce superoxide anion production through degranulation by eosinophils [11c]. However, trypsin activity was checked in this study through a BSA degradation test, and the endotoxin level was determined using the LAL test kit (data not shown). The results demonstrate no relationship between both these factors and the immune activation of RAW264.7 cells. In addition, immune activation by TSP treatment was consistent with the effect of FSP>30. Further, according to the LC-MS/MS analysis, FSP>30 contained fibroin-derived peptides. As discussed earlier, the tryptic degradation of sericin was divided into small peptides of less than 30 kDa, and the results for anti-tumor activity and blood pressure [3,4a] suggests that fibroin, a major ingredient in FSP>30, may be a key factor in the activation of RAW264.7 cells. The present study suggests that the activation of RAW264.7 macrophages may be induced not by sericin-derived peptides but by fibroin-derived ones. In this regard, future research should precisely identify the active peptides.

Experimental

Preparation of trypsin-digested silk peptides: Hydrolyzed silk protein was prepared from cocoons of the silkworm *B. mori* [4c]. The hydrolysates were sterilized by boiling at $90^\circ\text{C} \pm 3^\circ\text{C}$. For the additional trypsin digestion of the hydrolysates, trypsin was dissolved in DW and reacted at 37°C for 12 h with the hydrolysates by adjusting at the final trypsin concentration of 0.25%. After the digestion, the trypsin enzyme was inactivated by boiling at 95°C for 30 min. Trypsin-digested total silk peptides (TSP) were concentrated and freeze-dried.

Cell culture and the in vitro treatment of silk peptides: RAW264.7 cells were obtained from ATCC and maintained in DMEM (WelGENE Inc., Daegu, Korea) with 10 % FBS (Gibco, Carlsbad, CA, USA), 1% antibiotics/antimycotics (Gibco), 10 mM HEPES, 2 mM L-glutamine, 2 mM sodium bicarbonate, and 5×10^{-5} M 2-mercaptoethanol. Cultured RAW264.7 cells were counted and seeded at 1.0×10^6 cells/well onto six-well plates. For the *in vitro* experiment, 5 experimental groups were considered: LPS, LPS + trypsin-digested total silk peptides (TSP), TSP, fractionated silk peptides of more than 30 kDa (FSP>30), and fractionated silk peptides of less than 30 kDa (FSP<30). The concentration of LPS was 10 $\mu\text{g/mL}$, and that of the other stimulants 0.83 mg/mL for both TSP and fractionated silk proteins (FSP>30 and FSP<30). RAW264.7 cells were cultured *in vitro* for 24 h and then harvested using cold PBS to prevent cell activation by chemical or mechanical stimulation.

Fluorescence-activated cell sorter analysis: Harvested RAW264.7 cells were washed with DMEM without FBS by centrifugation at 1,200 rpm at 4°C for 5 min. The cell pellet was suspended in FACS (fluorescence-activated cell sorter) buffer (1% BSA and 0.1% NaN_3). An anti-CD16/32 antibody (eBioscience, San Diego, USA) was applied and incubated for 10 min to block Fc receptors. FITC-conjugated anti-mouse CD11b (eBioscience) and PE-conjugated anti-mouse CD40 (eBioscience) were used for the FACS staining of RAW264.7 cells. Both anti-CD11b-FITC and anti-CD40-PE were diluted to 1: 100 with FACS buffer. The number of stained cells

among 2×10^4 cells was determined by FL-1 (FITC) and FL-2 (PE) filters in the FACSCalibur flow cytometer (BD, USA).

Fractionation of silk peptides using Centriprep 30K: One hundred mg of the silk peptide (TSP) was dissolved in 10 mL of DDW and fractionated by the Centriprep 30K concentrator (Millipore, Bedford, MA, USA). Ultrafiltration was performed at $1,500 \times g$ at 4°C for 10 min. The concentrated samples were dissolved in DDW up to 10 mL and centrifuged again for 5 min. This process was repeated, and the final sample (FSP>30) was sterilized using a $0.45 \mu\text{m}$ syringe-driven filter unit (Millipore, MA, USA). Small peptides (FSP<30) were collected from the Centriprep filtered fraction and filtered using a $0.45 \mu\text{m}$ syringe filter (Millipore).

Morphological observation of RAW264.7 cells activated by silk peptides: RAW264.7 cells were cultured with LPS, LPS+TSP, TSP, FSP>30, or FSP<30 in six-well plates. After 24 h, the morphology of the cells was observed using a microscope equipped with a video camera. The cells were observed at $\times 400$ magnification.

Residual enzyme activity after the trypsin proteolysis of silk peptides: Silk protein hydrolysates were digested with trypsin (Novozymes, Bagsvaerd, Denmark) at 37°C for 12 h for proteolysis of silk protein. The reaction was stopped by incubating at 95°C for 30 min. For the determination of residual enzyme activity at the end of the enzyme reaction, the reaction product between silk protein and trypsin was incubated with $15 \mu\text{g}$ of BSA at 37°C for 12 h for the final reaction ratio of 1:1, 1:2, or 1:5 in BSA to the silk peptide reaction product. The final reaction sample was loaded on a 15% SDS-polyacrylamide gel based on SDS-PAGE.

In-gel digestion for LC/MS/MS: Ten μg of the protein sample was separated by 12% SDS-PAGE (mini-PROTEAN, Bio-Rad, Richmond, CA, USA), and the gel was stained with Coomassie Brilliant Blue R-250. In-gel digestion was conducted based on prescribed methods [12]. The gel was fractionated into 6 parts according to their molecular weight. Each part of the gel was digested with trypsin ($0.1 \mu\text{g}$) for 16 h at 37°C after the reduction and alkylation of cysteine in the protein. Digested peptides were extracted using an extraction solution (50 mM ammonium bicarbonate, 50% acetonitrile, and 5% trifluoroacetic acid). Digested peptides were resolved in $10 \mu\text{L}$ of sample buffer containing 0.02% formic acid and 0.5% acetic acid.

1-DE and LC-MS/MS using LCQ mass spectrometry: Peptide samples ($10 \mu\text{L}$) were concentrated using a MGU30-C18 trapping column (LC Packings, Sunnyvale, CA). Then the peptides were eluted from the column and directed onto a $10 \text{ cm} \times 5 \mu\text{m}$ i.d. C18 reverse phase column at a flow rate of 120 nL/min . For an additional analysis, peptides were eluted by a gradient of 0–65% acetonitrile for 120 min. Each full MS (m/z range of 300 to 2,000) scan was followed by 3 MS/MS scans of the most abundant precursor ions in the MS (with dynamic exclusion enabled). For protein identification, MS/MS spectra were searched by MASCOT (Matrix science, www.matrixscience.com) and determined by the database of the *B. mori* genome sequence.

Acknowledgments - This work was supported by Worldway Co. Ltd., Korea (grant no. 800-20100530).

References

- (a) Sutherland TD, Young J, Weisman S, Hayashi CY, Merritt DJ. (2010) Insect silk: One name, many materials. *Annual Review Entomology*, **55**, 171-188; (b) Horan RL, Antle K, Collette AL, Wang Y, Huang J, Moreau JE, Volloch V, Kaplan DL, Altman GH. (2005) *In vitro* degradation of silk fibroin. *Biomaterials*, **26**, 3385-3393; (c) Santin M, Motta A, Freddi G, Cannas M. (1999) *In vitro* evaluation of the inflammatory potential of the silk fibroin. *Journal of Biomedical Materials Research*, **46**, 382-389; (d) Tsubouchi K, Igarashi Y, Takasu Y, Yamada H. (2005) Sericin enhances attachment of cultured human skin fibroblasts. *Bioscience Biotechnology Biochemistry*, **69**, 403-405.
- (a) Teramoto H, Kameda T, Tamada Y. (2008) Preparation of gel film from *Bombyx mori* silk sericin and its characterization as a wound dressing. *Bioscience Biotechnology Biochemistry*, **72**, 80375-1-8; (b) Jin H, Chen J, Karageorgiou V, Altman GH, Kaplan DL. (2004) Human bone marrow stromal cell responses on electrospun silk fibroin mats. *Biomaterials*, **25**, 1039-1047; (c) Yamada H, Nakao H, Takasu Y, Tsubouchi K. (2001) Preparation of undegraded native molecular fibroin solution from silkworm cocoons. *Materials Science and Engineering: C*, **14**, 41-46.
- Igarashi K, Yoshioka K, Mizutani K, Miyakoshi M, Murakami T, Akizawa T. (2006) Blood pressure-depressing activity of a peptide derived from silkworm fibroin in spontaneously hypertensive rats. *Bioscience Biotechnology Biochemistry*, **70**, 517-520.
- (a) Byun EB, Sung NY, Kim JH, Choi J, Matsui T, Byun MW, Lee JW. (2010) Enhancement of anti-tumor activity of gamma-irradiated silk fibroin via immunomodulatory effects. *Chemico-Biological Interactions*, **186**, 90-95; (b) Ikegawa Y, Sato S, Lim G, Hur W, Tanaka K, Komori M, Takenaka S, Taira T. (2012) Amelioration of the progression of an atopic dermatitis-like skin lesion by silk peptide and identification of functional peptides. *Bioscience, Biotechnology and Biochemistry*, **76**, 110748-1-5; (c) Moon JH, Pyo KH, Jung BK, Chun HS, Chai JY, Shin EH. (2011) Resistance to *Toxoplasma gondii* infection in mice treated with silk protein by enhanced immune responses. *Korean Journal of Parasitology*, **49**, 303-308.
- (a) Jung EY, Lee HS, Lee HJ, Kim JM, Lee KW, Suh HJ. (2010) Feeding silk protein hydrolysates to C57BL/6J-db/db mice improves blood glucose and lipid profiles. *Nutrition Research*, **30**, 783-790; (b) Fukao T, Frucht DM, Yap G, Gadina M, O'Shea JJ, Koyasu S. (2001) Inducible expression of Stat4 in dendritic cells and macrophages and its critical role in innate and adaptive immune responses. *Journal of Immunology*, **166**, 4446-4455; (c) Berghaus LJ, Moore JN, Hurley DJ, Vandenplas ML, Fortes BP, Wolfert MA, Boons GJ. (2010) Innate immune responses of primary murine macrophage-lineage cells and RAW 264.7 cells to ligands of Toll-like receptors 2, 3, and 4. *Comparative Immunology, Microbiology & Infectious Disease*, **33**, 443-454.
- Fearon D, Locksley R. (1996) The instructive role of innate immunity in the acquired immune response. *Science*, **272**, 50-54.
- Padamwar M, Pawar AP. (2004) Silk sericin and its applications: A review. *Journal of Scientific & Industrial Research*, **63**, 323-329.
- Li M, Ogiso M, Minoura N. (2003) Enzymatic degradation behavior of porous silk fibroin sheets. *Biomaterials*, **24**, 357-365.
- (a) Wu JH, Wang Z, Xu SY. (2008) Enzymatic production of bioactive peptides from sericin recovered from silk industry wastewater. *Process Biochemistry*, **43**, 480-487; (b) Polgar L. (2005) The catalytic triad of serine peptidases. *Cellular and Molecular Life Sciences*, **62**, 2161-2172.
- (a) Corey DR, Craik CS. (1992) An investigation into the minimum requirements for peptide hydrolysis by mutation of the catalytic triad of trypsin. *Journal of the American Chemical Society*, **114**, 1784-1790; (b) Bucinski A, Karamac M, Amarowicz R, Pegg RB. (2008) Modeling the tryptic hydrolysis of pea proteins using an artificial neural network. *LWT-Food Science and Technology*, **41**, 942-945.
- (a) Piippo M, Lietzen N, Nevalainen OS, Salmi J, Nyman TA. (2010) Pripper: Prediction of caspase cleavage sites from whole proteomes. *BMC Bioinformatics*, **11**, 320-328; (b) Iwasaki A, Medzhitov R. (2004) Toll-like receptor control of the adaptive immune responses. *Nature Immunology*, **5**, 987-995; (c) Miike S, McWilliam AS, Kita H. (2001) Trypsin induces activation and inflammatory mediator release from human eosinophils through protease-activated receptor-2. *Journal of Immunology*, **167**, 6615-6622.
- Kim YH, Cho K, Yun SH, Kim JY, Kwon KH, Yoo JS, Kim SI. (2006) Analysis of aromatic catabolic pathways in *Pseudomonas putida* KT 2440 using a combined proteomic approach: 2-DE/MS and cleavable isotope-coded affinity tag analysis. *Proteomics*, **6**, 1301-1318.

Low-Volatile Lipophilic Compounds in Needles, Defoliated Twigs, and Outer Bark of *Pinus thunbergii*

Alexander V. Shpatov^a, Sergey A. Popov^{a*}, Olga I. Salnikova^a, Ekaterina A. Khokhrina^a, Emma N. Schmidt^a and Byung Hun Um^b

^aNovosibirsk Institute of Organic Chemistry, Lavrentyev Ave. 9, Novosibirsk, 630090, Russia

^bNatural Products Research Center, Korea Institute of Science and Technology – Gangneung Institute, 290 Daejeon-dong, Gangneung 210-340, Gangwon-do, Republic of Korea

spopov@nioch.nsc.ru

Received: August 7th, 2013; Accepted: September 19th, 2013

Despite a long history of the use of *Pinus thunbergii* for technical, medicinal, agricultural, and other purposes, the composition of low-volatile metabolites in the used parts of the plant has been poorly investigated. We report here on the distribution of lipophilic extractive compounds in different parts of the shoot system (needles, defoliated twigs, outer bark) of *P. thunbergii* studied by GC/MS. The highest and lowest contents of lipophilic substances were found in defoliated twigs and in outer bark correspondingly. Acid compounds in the extract of needles comprised mainly labdane type diterpenoids (*trans*-communic acid), while in the extracts of defoliated twigs and outer bark the acids were represented predominantly by abietane type compounds (neoabietic, dehydroabietic, abietic, levopimaric and palustic acids). The major neutral components of the extract of needles were 10-nonacosanol, labdanoids (18-hydroxy-13-*epi*-manoyl oxide, *trans*-communol), and β -sitosterol. In the case of the extract of defoliated twigs, labdanoids (18-hydroxy-13-*epi*-manoyl oxide, *trans*-communol, 13-*epi*-torulosol), serratane triterpenoids (3 β -methoxyserrat-14-en-21-one), and β -sitosterol were the main neutral constituents, whereas serratane triterpenoids (3 β -methoxyserrat-14-en-21-one) alone dominated among the neutral compounds of the outer bark extract. Most of the neutral components and the labdane type acids were detected for the first time in organs and tissues of *P. thunbergii*. The distribution of lipophilic metabolites in the studied parts of *P. thunbergii* shoot system may be applied for chemotaxonomy purposes. Diversified accumulation of extractive substances in different organs of the plant should be taken into account for isolation of specific components from the pine raw material.

Keywords: *Pinus thunbergii*, Phytochemical study, Extractive compounds, Diterpenes, Triterpenes, Labdanoids, Serratane triterpenoids, GC/MS.

Pinus thunbergii Parl. (fam. Pinaceae, gen. *Pinus*, subgen. *Pinus*, sect. *Pinus*, subsect. *Pinus*) is a two-needle pine species naturally growing in coastal areas of southern Korea and Japan up to elevations of about 1000 m. Various parts of the plant have been used in traditional and folk medicines and cuisine. Needles and sprouts of *P. thunbergii* are used as ingredients in preparation of a healthy drink preventing arteriosclerosis, hypertension, and aging [1]. Water soluble compounds from the bark of the plant showed antioxidant activity [2], as well as skin whitening and wrinkle smoothing effects [3]. Ethanol and methanol extracts of the pine needles were shown to inhibit α -amylase [4] and be active against influenza viruses [5]. Essential oil from needles possesses antimicrobial activity [6], while essential oils from bark, needles, and wood exhibit growth inhibitory activity to microalgae [7]. Among all the *P. thunbergii* extractives, only the composition of the needle essential oil has been thoroughly investigated [8,9]. However, high-volatile essential oil components, predominantly mono- and sesquiterpenoids, comprise a minor part of the pines' lipophilic extractives [9,10]. Certain data on extractive compounds from different parts of *P. thunbergii* have been reported. The chemical composition of the pine oleoresin, including mono-, sesqui-, and diterpenoids, was briefly characterized [11]. Resin acids from the plant cone extracts were studied several decades ago [12]. Some leaf wax components, estolides, were isolated and their structures elucidated [13], and the structures of polyphenols from the pine needles were determined [14]. A number of phenolic compounds were identified in wood and bark of *P. thunbergii* [15]. Total phenolic and proanthocyanidin contents of the water extract of pine bark was ascertained [2]. The diterpene hydrocarbon, cembrene (thunbergene), was first isolated from the root turpentine oil of *P. thunbergii* [16]. Data on the high-volatile components and

several groups of extractive compounds of *P. thunbergii* are available, but the composition of the low-volatile lipophilic metabolites in needles, twigs, and especially bark of the plant remains generally undisclosed.

The goal of this work was a comparative study of the low-volatile lipophilic components of the needles, defoliated twigs, and outer bark of *P. thunbergii* from a single natural population, employing a GC/MS method. The distribution of lipophilic compounds in the shoot system of *P. thunbergii* will help to elaborate isolation methods for valuable phytochemicals from the plant raw material. On the other hand, chromatographic profiling of low-volatile secondary metabolites could be useful for chemotaxonomy of the genus *Pinus*.

Table 1: Yields of extracts from needles (N), defoliated twigs (T), and outer bark (B) of *P. thunbergii*, yields of fractions of acids (A) and neutral compounds (NC) from the extracts, and ratio of the fraction yields.

	Extract ^a	Acids (A) ^b	Neutral compds. (NC) ^b	Ratio of acids/neutral compds. (A/NC)
Needles [9]	0.3 (essential oil) ^c			
Needles (N)	3.7	64.8	23.8	2.7
Defoliated twigs (T)	5.3	62.9	18.1	3.5
Outer bark (B)	1.1	31.1	31.9	1.0

^aYields (% w/w) of extracts relative to the air-dried raw material. ^bYields (% w/w) of fractions relative to the extract. ^cYield (% w/w) of essential oil relative to the fresh (8.5% moisture) raw material.

Air-dried needles, defoliated twigs, and outer bark of *P. thunbergii* were exhaustively extracted with *n*-hexane. The obtained extracts were fractionated according to the general procedure [17] into neutral compounds (NC) and acids (A). Yields of the extracts and the resulting fractions are shown in Table 1. The yields of

The major component of the acid fraction from the needle extract (**N-A**) was the labdanoid, *trans*-communic acid (43.8%), while contents of other diterpene acids were considerably lower {dehydroabietic (7.2%), abietic (5.9%), imbricatoloic (5.1%) acids}. The total percentage of fatty acids was *ca.* 4%. On the contrary, in the analogous fraction from the defoliated twig extract (**T-A**) the main acids were of the abietane type {neoabietic (19.0%), levopimaric together with palustric (19.0%), and dehydroabietic acids (10.1%)}. The content of labdane type acids {*trans*-communic (4.7%), imbricatoloic (3.0%), and imbricatoloic (2.7%) acids} in this fraction was significantly inferior to that of fraction **N-A**. The fraction **T-A** was richer in fatty acids (*ca.* 7%) than **N-A**. Analogous to fraction **T-A**, the major components among the acids from the extract of the outer bark (**B-A**) were abietanoids {dehydroabietic (30.5%) and abietic (8.0%) acids}, while very minor amounts of labdane type acids {dihydroagathic acid together with dihydroagathic acid monomethyl ester (0.4%)} were detected. Fraction **B-A** contained a significant amount of fatty acids (*ca.* 23%), the dominant one being lignoceric acid (9.1%). In the order of the fractions **N-A**, **T-A**, and **B-A**, the percentages of labdane type acids (*trans*-communic, imbricatoloic, imbricatoloic, and dihydroagathic acids) decreased, while the percentages of dehydroabietic, abietic, pimaric, and isopimaric acids increased. The maximal amounts of neoabietic and levopimaric together with palustric acids were found in fraction **T-A** compared with the other acid fractions. The percentage of saturated C₂₀, C₂₂, C₂₄, and C₂₆ fatty acids increased in the series **N-A**, **T-A**, **B-A**, though the

Table 4: Composition of unsaponifiable neutral compound fractions from extracts of *P. thunbergii* needles (N-U), defoliated twigs (T-U), and outer bark (B-U) by GC/MS^a.

Compound	RI	N-U	T-U	B-U
<i>n</i> -Octadecane	1800			0.5
Cembrene (thunbergene)	1927			0.5
Manoyl oxide	1988	1.0	1.1	
<i>n</i> -Eicosane	2000			0.4
13- <i>epi</i> -Manoyl oxide	2009	2.4	1.2	
Dehydroabietane + manool (major)	2054		0.7	
Abieta-7,13-diene	2079		0.3	
Phytol	2121	2.6		
<i>cis</i> -Abienol (major) + unidentified [M ⁺ >277]	2149		1.0	
Pimarinal	2162		0.8	0.5
Manoyl oxide-18-al ^b	2195		0.8	
Unidentified [M ⁺ =286]	2220	1.3		
Unidentified [M ⁺ =286] + 13- <i>epi</i> -manoyl oxide-18-al ^b	2217		2.2	
Sandaracopimarinal	2252		1.1	
Sandaracopimaric acid methyl ester	2252	0.4		
Dehydroabietal	2269		1.0	
18-Hydroxymanoyl oxide ^b	2295		2.7	
Abietal	2305		1.1	
trans-Communal	2317	3.8	3.7	
18-Hydroxy-13-<i>epi</i>-manoyl oxide^b	2321	4.0	8.0	
Dehydroabietic acid methyl ester	2342	0.4	1.3	2.1
13-<i>epi</i>-Torulosol	2359		3.6	
Dehydroabietol	2368		0.6	
Neoabietal	2374		1.4	
Abietic acid methyl ester	2384	0.3		0.7
Abietol	2399		0.9	
<i>iso</i> -Agathal	2457		1.0	
Unidentified [M ⁺ =332] + neoabietol	2461		1.0	
Abieta-7,13,15-trienoic acid methyl ester	2497		0.4	
<i>n</i> -Pentacosane	2500		0.4	
15-Hydroxydehydroabietic acid methyl ester	2528		0.7	
<i>n</i> -Heptacosane + unidentified [M ⁺ =?]	2700	0.9	1.7	
1-Tetracosanol	2708			1.2
10-Nonacosanol	3089	3.1		
3,5-Stigmastadiene	3091			0.7
10-Nonacosanol	3110	38.0	3.0	
α -Tocopherol	3138	1.1		
Unidentified [M ⁺ =?] + [M ⁺ =434]	3210	5.1	0.8	
Campesterol	3229	1.0	1.2	0.7
β-Sitosterol	3323	15.8	19.8	14.3
Unidentified [M ⁺ =450]	3337	2.6		
5 α -Stigmastan-3-one	3372			1.2
Stigmast-4-en-3-one	3456			2.7
3 β -Methoxylanosta-9(11)-dienol	3486			0.8
Unidentified [M ⁺ =410]	3499			3.5
Unidentified [M ⁺ =440]	3553			4.5
Serratanoids [M ⁺ =454] + [M ⁺ =426]	3627			6.2
3 β ,21-Dimethoxyserrat-14-ene	3669		3.0	5.0
29-Nor-3 β -methoxyserrat-14-en-21-one	3677		0.9	5.8
3 α -Methoxyserrat-14-en-21-one	3687		0.3	1.4
3β-Methoxyserrat-14-en-21-one	3741		15.7	27.5
3 β -Methoxyserrat-14-en-21 β -ol			2.0	1.9
Serrat-14-en-3,21-dione			1.9	4.6
Unidentified [M ⁺ =440]			2.6	2.5
High-volatile compounds (not tabulated)		11.9	6.3	6.9
Total		95.4	96.2	95.9

^aContent of compounds is given in %; only major unidentified components are tabulated.^bCompounds with tentative configuration of stereogenic centers at atoms C-4 and C-13.

content of unsaturated fatty acids was insignificant in all acid fractions. The prevalent component of the acids from the pine needles, *trans*-communic acid, was reported among the dominant compounds in oleoresin and a cone extract of *Pinus luchuensis* (subsect. *Pinus*) [11,18], a species which is very close to *P. thunbergii* in terms of botanical classification. Predominance of levopimaric with palustric acids over abietic acid in the fraction **T-A** indicates additionally that the analyzed samples of plant material were taken from healthy trees, non-infected with the invasive pine wood nematode, *Bursaphelenchus xylophilus*, widespread in East Asia nowadays [19].

The fraction of unsaponifiable neutral compounds from the needle extract (**N-U**) contained *ca.* 12% of high-volatile compounds (mostly sesquiterpenoids), *ca.* 12% of diterpenoids, and *ca.* 17% of sterols. The major components in the fraction **N-U** were 10-nonacosanol (38.0%), β -sitosterol (15.8%), labdane type alcohols, 18-hydroxy-13-*epi*-manoyl oxide (4.0%) and *trans*-communal (3.8%). The total content of mono- and sesquiterpenoids

in the unsaponifiable neutral compounds from the twig extract (**T-U**) was only *ca.* 6%, whereas the percentages of diterpenoids and triterpenoids made up *ca.* 37% and *ca.* 21%, respectively. The main components were β -sitosterol (19.8%), the triterpenoid 3 β -methoxyserrat-14-en-21-one (15.7%), and the labdanoids (18-hydroxy-13-*epi*-manoyl oxide (8.0%), *trans*-communal (3.7%), and 13-*epi*-torulosol (3.6%)). The greatest variety of diterpene aldehydes and alcohols was found in fraction **T-U**. The fraction of bark unsaponifiable compounds (**B-U**) comprised high-volatile components (*ca.* 7%), diterpenoids (*ca.* 4%), sterols (*ca.* 20%), and a large amount of lanostane and serratane type triterpenoids (*ca.* 62%). The dominant compounds in the fraction were 3 β -methoxyserrat-14-en-21-one (27.5%) and β -sitosterol (14.3%). Compared with the parent **NC** fractions, the content of β -sitosterol in the fractions **N-U** and **T-U** was *ca.* 4 times higher, while in the fraction **B-U** only *ca.* 2 times. The fraction **T-U** contained maximal amounts of diterpenoids, while **B-U** was particularly rich in triterpenoids.

The content of fatty acids in the fraction of acids after saponification of the neutral compounds from needles (**N-S**) was *ca.* 34%, the major ones being lauric, oleic, linoleic, and palmitic acids (*ca.* 20% in total). In the analogues fraction obtained from the extract of defoliated twigs (**T-S**) the dominant acids, linoleic, oleic, behenic, and lignoceric acids, made up to 66%. At the same time, the fraction of acids after saponification of the neutral compounds from outer bark (**B-S**) contained mostly saturated C₂₀-C₂₆ even fatty acids (*ca.* 61% in sum), the major one being lignoceric acid (*ca.* 31%). The composition of the *P. thunbergii* extractive substances resembles that for *P. densiflora* [20], another two-needle pine from subsect. *Pinus*, growing in the same region. The principal differences in the composition of lipophilic extractives in the case of these two geographically and systematically related species enclose the following features. First, all the bicyclic labdane type acids (*trans*-communic, imbricatol, imbricatol, dihydroagathic acids) from the *P. thunbergii* extracts possessed the same (4*S*)-stereoconfiguration, whereas the analogous metabolites produced by *P. densiflora* were found to have the (4*R*)-configuration. Secondly, despite the fact that cembrene (thunbergene) was first found in roots of *P. thunbergii*, only minor amounts of it could be detected among the extractives of the outer bark. On the contrary, cembrene type diterpenoids are abundantly produced in the vegetative organs of *P. densiflora*.

The distribution was studied of low-volatile lipophilic compounds in needles, defoliated twigs, and outer bark of *P. thunbergii*. Acid compounds in the extract of needles comprised mainly labdane type diterpenoids (*trans*-communic acid), while in the extracts of defoliated twigs and outer bark the acids were represented predominantly by abietane type compounds (neoabietic, dehydroabietic, abietic, levopimaric and palustric acids). The major neutral components of the needle extract were 10-nonacosanol, labdanoids (18-hydroxy-13-*epi*-manoyl oxide, *trans*-communal), and β -sitosterol; in the case of the extract of defoliated twigs, they were labdanoids (18-hydroxy-13-*epi*-manoyl oxide, *trans*-communal, 13-*epi*-torulosol), serratane triterpenoids (3 β -methoxyserrat-14-en-21-one), and β -sitosterol; whereas serratane (3 β -methoxyserrat-14-en-21-one) alone dominated among the neutral compounds of the outer bark extract. Most of the neutral components and the labdane type acids were detected for the first time in organs and tissues of *P. thunbergii*. The low-volatile lipophilic compounds found in the studied parts of the pine extend the scope of its secondary metabolites detected earlier in essential oils.

Experimental

Plant material: Foliated twigs and outer bark of *P. thunbergii* Parl. were collected in September 2010 near Gangneung city, Republic of Korea. The specimens were identified by Dr E.A. Korolyuk (Central Siberian Botanical Garden (CSBG), Novosibirsk). Voucher specimens ("Korea, near Gangneung city, 09. 2010. Collect. Popov S.") are deposited with the Central Siberian Botanical Garden (Novosibirsk; NS).

Drying, extraction, and fractionation: Foliated twigs of *P. thunbergii* were separated to give needles and defoliated twigs. Needles, defoliated twigs, and outer bark were dried to constant weight (residual moisture content ~5%) in a drying cabinet at 30–50°C for 24 h. Dried plant material was cut into pieces of 10–15 mm and extracted with *n*-hexane in a Soxhlet apparatus for 24 h. The extracts were concentrated in vacuum, diluted with MeOBu-*t* and fractionated to give acids (A) and neutral compounds (NC) by treatment of ether solutions of the initial extracts with 1% NaOH aqueous solution. Portions of the NC fractions were saponified by boiling in 5% KOH aqueous ethanol solution and separated to give acids (S) and unsaponifiable neutral compounds (U). Portions of the acid fractions (A, S) were converted to methyl esters by treatment with a diethyl ether solution of diazomethane [21].

GC-MS analysis: GC-MS was performed using a Hewlett-Packard instrument with a HP 5890 Series II gas chromatograph with HP 5971 (EI, 70 eV) mass-selective detector and a HP-5ms capillary column (30 m × 0.25 mm internal diameter, film thickness 0.25 µm); carrier gas was He at a flow rate of 1 mL/min; programmed column temperature was at 50°C for 2 min, from 50°C to 300°C at 4°C/min, and 300°C for 30 min (or from 50°C to 320°C at 4°C/min, and 320°C for 30 min when an analyte contained triterpenoids); vaporizer temperature was 280°C; ion-source temperature was 175°C, and scan rate was 1.2 scans/s in the mass range 30–650 amu. Linear retention indices (*R*_Is) of the components were calculated as described in [22] by interpolation of *R*_Is of *n*-alkanes C₇–C₄₀. The components were identified: 1) by comparing their mass spectra with reference MS from commercial libraries (Wiley 7n and NIST 02), home-made library, literature data [22,23], and ones of authentic compounds; and 2) by comparing their calculated *R*_Is with those of authentic standards and the data published in the literature [22]. The contents (%) of compounds were determined from peak areas in chromatograms without using correction coefficients.

Acknowledgments - This research was financially supported by the Ministry of Education, Science and Technology (MEST), Gangwon Province, Gangneung City (Republic of Korea), Gangneung Science Industry Foundation (GSIF) as the R&D Project for Gangneung Science Park promoting program. The authors thank Dr Elena A. Korolyuk (CSBG) for identification of the species.

References

- (a) Kim KY, Chung HJ. (2000) Flavor compounds of pine sprout tea and pine needle tea. *Journal of Agricultural and Food Chemistry*, **48**, 1269–1272; (b) Lee GS. (2001) Preparation method of drink using black pine needles. R Korea patent: KR20010008094.
- Kim SM, Kang SW, Jeon JS, Um BH. (2012) A comparison of pycnogenol and bark extracts from *Pinus thunbergii* and *Pinus densiflora*: Extractability, antioxidant activity and proanthocyanidin composition. *Journal of Medicinal Plants Research*, **6**, 2839–2849.
- Park SM, Park JY. (2009) Cosmetic composition containing pine endodermis extract from *Pinus thunbergii* Parl. R Korea patent: KR20090123435.
- Kim YM, Wang MH, Rhee HI. (2004) A novel α-glucosidase inhibitor from pine bark. *Carbohydrate Research*, **339**, 715–717.
- Won JN, Lee SY, Song DS, Poo H. (2013) Antiviral activity of the plant extracts from *Thuja orientalis*, *Aster spathulifolius*, and *Pinus thunbergii* against influenza virus A/PR/8/34. *Journal of Microbiology and Biotechnology*, **23**, 125–130.
- Kim HS, Lee BS, Yun KW. (2013) Comparison of chemical composition and antimicrobial activity of essential oils from three *Pinus* species. *Industrial Crops and Products*, **44**, 323–329.
- Tsuruta K, Yoshida Y, Kusumoto N, Sekine N, Ashitani T, Takahashi K. (2011) Inhibition activity of essential oils obtained from Japanese trees against *Skeletonema costatum*. *Journal of Wood Science*, **57**, 520–525.
- (a) Idžojić M. (2000) Composition of volatiles from needles of *Pinus sylvestris*, *P. nigra*, *P. densiflora* and *P. thunbergiana*. *Šumarski List*, **124**, 263–270; (b) Pfeifhofer HW, Idžojić M, Zebec M. (2008) The needle volatile composition of *Pinus nigra* J.F. Arnold, *P. sylvestris* L., *P. densiflora* Siebold et Zucc. and *P. thunbergiana* Franco trispecies hybrids. *Silvae Genetica*, **57**, 221–226; (c) Sekine N, Shibutani S, Yatagai M. (2012) Water solubility of essential oils from coniferous needles with special reference to effect on germination and antitermite activity. *Mokuzai Gakkaishi*, **58**, 44–53.
- Park JS, Lee GH. (2011) Volatile compounds and antimicrobial and antioxidant activities of the essential oils of the needles of *Pinus densiflora* and *Pinus thunbergii*. *Journal of the Science of Food and Agriculture*, **91**, 703–709.
- Vasilyev SN, Roshchin VI, Yagodin VI. (1995) Extractive compounds of woody green of *Pinus sylvestris* L. *Rastitelnye Resursy*, **31**, 79–119.
- Song Z. (1998) *Characteristics of oleoresin and classification of Pinus in China*. China Academy of Forestry, Nanjing, 1–220.
- Yano S, Furuno T. (1994) Resin acids from extracts of pine cones of Kuromatsu (*Pinus thunbergii*). *Mokuzai Gakkaishi*, **40**, 72–77.
- Kariyone T, Isoi K, Yoshikura M. (1959) Chemical constituents of the plants of *Coniferae* and allied orders. XXVII. Plant waxes 14. Neutral estolide. *Yakugaku Zasshi*, **79**, 61–62.
- Wang C, Shen Z, Chen X. (1994) Structure of polyprenols from five conifer needles. *Linchan Huaxue Yu Gongye*, **14**, 1–8.
- Hata K, Sogo M. (1954) Detection of the phenolic substances of *Pinus densiflora*, *P. thunbergii*, *P. pentaphylla* var. *himekomatsu* and *P. armandii*. *Journal of the Japanese Forestry Society*, **36**, 8–11.
- Kobayashi H, Akiyoshi S. (1963) The structure of thunbergene. *Bulletin of the Chemical Society of Japan*, **36**, 823–826.
- Norin T, Winell B. (1972) Extractives from the bark of the Scots pine, *Pinus sylvestris*. *Acta Chemica Scandinavica*, **26**, 2297–2304.
- Minami T, Wada S, Tokuda H, Tanabe G, Muraoka O, Tanaka R. (2002) Potential antitumor-promoting diterpenes from the cones of *Pinus luchuensis*. *Journal of Natural Products*, **65**, 1921–1923.
- Xu G, Su Z. (1994) Study on the terpenoids in *Pinus thunbergii* Parl. infected with *Bursaphelenchus xylophilus*. *Linchan Huaxue Yu Gongye*, **14**, 49–54.
- Khohrina EA, Shpatov AV, Popov SA, Salnikova OI, Shmidt EN, Um BH. (2013) Non-volatile compounds of needles, trimmed saplings, and outer bark of *Pinus densiflora*. *Chemistry of Natural Compounds*, **49**, 561–565.
- Zinkel DF, Han JS. (1986) GLC determination of the resin acid composition in rosins and oleoresins: State of the art. *Naval Stores Review*, **96**, 14–19.
- Tkachev AV. (2008) *Study of volatile compounds of plants*. Ofset, Novosibirsk, 1–969.
- Zinkel DF, Zank LC, Wesolowski MF. (1971) *Diterpene resin acids: a compilation of infrared, mass, nuclear magnetic resonance, ultraviolet spectra and gas chromatographic retention data (of the methyl esters)*. US Forest Service Forest Products Laboratory, Madison, 1–191.

Lipid Constituents of the Edible Mushroom, *Pleurotus giganteus* Demonstrate Anti-Candida Activity

Chia-Wei Phan^{a,b}, Guan-Serm Lee^{a,b}, Ian G. Macreadie^c, Sri Nurestri Abd Malek^{a,b}, David Pamela^{a,d} and Vikineswary Sabaratnam^{a,b}

^aMushroom Research Centre, Institute of Biological Sciences, Faculty of Science, University of Malaya, 50603 Kuala Lumpur, Malaysia

^bInstitute of Biological Sciences, Faculty of Science, University of Malaya, 50603 Kuala Lumpur, Malaysia

^cSchool of Applied Sciences, RMIT University, Victoria 3083, Australia

^dDepartment of Anatomy, Faculty of Medicine, University of Malaya, 50603 Kuala Lumpur, Malaysia

wiki@um.edu.my

Received: May 10th, 2013; Accepted: October 10th, 2013

Different solvent extracts of *Pleurotus giganteus* fruiting bodies were tested for antifungal activities against *Candida* species responsible for human infections. The lipids extracted from the ethyl acetate fraction significantly inhibited the growth of all the *Candida* species tested. Analysis by GC/MS revealed lipid components such as fatty acids, fatty acid methyl esters, ergosterol, and ergosterol derivatives. The sample with high amounts of fatty acid methyl esters was the most effective antifungal agent. The samples were not cytotoxic to a mammalian cell line, mouse embryonic fibroblasts BALB/c 3T3 clone A31. To our knowledge, this is the first report of antifungal activity of the lipid components of *Pleurotus giganteus* against *Candida* species.

Keywords: *Pleurotus giganteus*, *Candida*, yeast, Antifungal, Medicinal mushroom, Fatty acid, Fatty ester methyl ester, Ergosterol.

Fungal infections are problematic for human health and are responsible for high rates of morbidity and mortality worldwide. Species of *Candida* are the dominant cause of opportunistic mycoses and among them, *C. albicans*, *C. glabrata*, *C. parapsilosis*, *C. tropicalis* and *C. krusei* account for 95–97% of all *Candida* infections [1,2]. *C. albicans* and *C. tropicalis* are susceptible to polyenes, flucytosine, azoles and echinocandins, while *C. glabrata* is either susceptible or resistant to fluconazole[3]. Furthermore, *C. krusei* displays decreased susceptibility to amphotericin B, as well as fluconazole. Considering the increasing incidence of drug-resistant *Candida* infections, the search for more effective anti-*Candida* agents as an alternative to synthetic ones is needed. The interest in natural products from medicinal plants as a source of anti-*Candida* agents has grown dramatically. A wide variety of plant extracts have been reported to have anti-*C. albicans* activity. Examples include propolis, *Punica granatum* (pomegranate), *Streblus asper* (Siamese rough bush), *Vitis vinifera* (common grape vine), and tea tree oil from *Melaleuca alternifolia* [4].

Medicinal mushrooms are relatively less researched for their antifungal properties. However, in the last five years, there has been a renewed interest in using mushrooms as antimicrobial agents. *Lentinula edodes* (shiitake), *Boletus edulis* (Penny bun), *Pleurotus ostreatus* (oyster mushroom), *Coprinus comatus* (shaggy mane), *Astraeus hygrometricus* (earthstar mushroom), and *Cordyceps militaris* were shown to exhibit antifungal activity against *C. albicans* [5–8]. *Pleurotus giganteus* (Zhudugu, Dabeijun, morning glory mushroom), a saprobic mushroom, is one of the largest mushroom which grows on soil either as solitary or gregarious fruiting bodies [9]. This mushroom has gained popularity in China for its culinary properties. The medicinal properties of this mushroom are less known. We have previously reported the hepatoprotective and neuronal stimulating effects of *P. giganteus* [10,11]. In this study the antifungal activities of different solvent extracts of this mushroom were evaluated. The extracts prepared with different solvents had different profiles of fatty acids, and fatty

Table 1: Activity of different extracts of *Pleurotus giganteus* against *Candida* species.

Candida strains	Untreated	Solvent extracts (µg/mL)								
		Methanol			Ethyl acetate			Aqueous		
		25	50	100	25	50	100	25	50	100
<i>Candida albicans</i> WM1172	++++	++++	+++	++	+	-	-	++++	++++	++++
<i>Candida albicans</i> ATCC90028	++++	+++	+++	+	+	-	-	++++	++++	+++
<i>Candida dubliniensis</i>	++++	++++	+++	-	+	-	-	++++	++++	+++
<i>Candida glabrata</i> CBS138	++++	++++	++	-	+	-	-	++++	++++	++++
<i>Candida glabrata</i> ATCC90030	++++	++++	++++	-	+	-	-	++	+++	+++
<i>Candida krusei</i> ATCC6258	++++	+++	+++	+	-	-	-	++++	+++	+++
<i>Candida pseudotropicalis</i>	++++	+++	+++	++	-	-	-	++++	++++	++++
<i>Candida tropicalis</i> WM30	++++	+++	++++	++	+	-	-	++++	++++	++++

Strains were grown with different mushroom extracts at the concentrations shown for two days on YEPD media. Growth was scored from “-” to “++++”, indicating no growth to strong growth.

acids have been shown to demonstrate antimicrobial activities [12]. The main metabolites / components in the extracts were analysed by GC-MS. As a preliminary *in vitro* toxicity assessment, the *P. giganteus* extracts were also investigated for cytotoxicity to mouse embryonic 3T3 fibroblast cells.

The anti-*Candida* activity of methanol, ethyl acetate and aqueous extracts of *P. giganteus* against all yeast species tested are summarised in Table 1. *Candida* species showed strong growth (denoted as “++++”) when extracts were not added to the medium. The aqueous extract had minimum or no inhibitory activity against all *Candida* spp. The ethyl acetate extract completely inhibited the growth of all *Candida* spp. when tested at 50 and 100 µg/mL. Thus, the ethyl acetate extract was fractionated to identify the active component/s responsible for the antifungal activity. Sub-fractions A to H were obtained and the minimum inhibitory concentration (MIC) values against all the tested yeasts are given in Table 2. The MIC values for all the *Candida* spp. tested, ranged from 2.0 ± 1.0 to 10.3 ± 2.5 µg/mL for sub-fraction A; and 9.3 ± 2.3 to 34.3 ± 10.8 µg/mL for sub-fraction B; respectively. The MIC values of

Table 2: Activity of the sub-fractions of ethyl acetate extracts against *Candida* species.

Candida strains	Sub-fractions from ethyl acetate extract (MIC ^a)							IC ₅₀ (mM)	
	A	B	C	D	E	F	G H	Fluconazole	Amphotericin B
<i>Candida albicans</i> WM1172	7.3 ± 1.5	16.0 ± 6.1	26.6 ± 11.6	*	*	*	*	1.0	0.6
<i>Candida albicans</i> ATCC90028	7.0 ± 1.0	22.6 ± 11.5	40.6 ± 16.6	*	*	*	*	2.0	0.7
<i>Candida dubliniensis</i>	8.2 ± 3.7	31.6 ± 7.8	37.0 ± 7.0	*	*	*	*	24.0	2.1
<i>Candida glabrata</i> CBS138	8.1 ± 1.5	12.5 ± 2.4	28.3 ± 6.6	*	*	*	*	>10.0	1.3
<i>Candida glabrata</i> ATCC90030	9.2 ± 1.6	29.4 ± 7.0	41.3 ± 7.6	*	*	*	*	>10.0	1.2
<i>Candida krusei</i> ATCC6258	10.3 ± 2.5	37.2 ± 5.0	*	*	*	*	*	8.0	0.4
<i>Candida pseudotropicalis</i>	3.8 ± 1.4	9.3 ± 2.3	23.0 ± 11.0	*	*	*	*	9.0	0.6
<i>Candida tropicalis</i> WM30	2.0 ± 1.0	34.3 ± 10.8	*	*	*	*	*	<1.0	1.2

Results were from three independent experiments performed in triplicate. ^aMIC is expressed in µg/mL. * : >50 µg/mL

sub-fraction C varied from 23.0 ± 11.0 to >50 µg/mL; whereas the MIC values for sub-fractions D-H were all >50 µg/mL. Overall, sub-fraction A showed the lowest MIC value for all *Candida* spp. tested.


Sub-fractions A and B were further analysed by GC-MS. Both samples were pale yellow-colored oils with a distinct odor. Constituents of sub-fractions A and B are listed in Table 3. Twelve compounds were identified in sub-fractions A and B. Sample A was characterized by high amounts of fatty acid methyl esters, namely: methyl palmitate, ethyl palmitate, methyl linoleate, methyl oleate, methyl stearate, and ethyl oleate. Sample B contained fatty acids (palmitic acid and oleic acid), fatty acid methyl esters (methyl linoleate and methyl oleate), ergosterol, ergosta-5,7,9 (11),22-tetraen-3β-ol, ergost-5,8(14)-dien-3-ol, and γ-ergosterol.

Table 3: Chemical composition of lipids in sub-fractions A and B of *P. giganteus*.

Constituents	RT (min)	Percentage (%)	Quality
Sub-fraction A			
Methyl palmitate	20.50	14.8	99
Ethyl palmitate	21.81	1.2	98
Methyl linoleate	23.70	19.8	99
Methyl oleate	23.80	39.3	99
Methyl stearate	24.26	3.3	99
Ethyl oleate	24.99	12.3	99
Sub-fraction B			
Methyl palmitate	20.49	0.2	95
Palmitic acid	21.28	14.4	99
Methyl linoleate	23.68	0.4	93
Methyl oleate	23.79	1.0	93
Oleic acid	24.61	31.7	99
Ergosta-5,7,9(11),22-tetraen-3β-ol	39.83	2.2	90
Ergosterol	40.33	24.4	98
Ergost-5,8(14)-dien-3-ol	40.51	10.2	87
γ-Ergosterol	41.32	3.7	94

The methanol, ethyl acetate, and aqueous extracts were not toxic to 3T3 fibroblasts cells and the IC₅₀ values were more than 2 mg/mL (Fig. 1). Meanwhile, cell viability (%) decreased steadily with increasing concentrations of sub-fractions A and B at levels up to 500 µg/mL. The IC₅₀ value of sub-fraction A was 352 µg/mL and the R² value was 0.9609. For sub-fraction B, the IC₅₀ was 362 µg/mL with the R² value recorded at 0.9552.

To our knowledge, this is the first report on the antifungal activity of the lipid components of *P. giganteus*. It has been reported that crude extracts of *P. ostreatus* and *C. comatus* inhibited the growth of *C. albicans* [6]. However, the MICs were much higher (up to 1 mg/mL) when compared with this study, which recorded an MIC of

**Figure 1:** Cell viability of embryonic fibroblast cells after treatment with various extracts of *P. giganteus*.

100 µg/mL. The sub-fractions A and B were shown to contain several bioactive components. Since they are blends of fatty acids and fatty acid methyl esters, they do not act on specific targets in the fungal cells, and fungal resistance may be unlikely to occur. Furthermore, fatty acids and their methyl esters were reported to have fungicidal activity to *C. albicans*, *C. krusei*, *C. tropicalis* and *C. parapsilosis* [13]. The entities might play crucial roles in lipophilic or hydrophilic effects on the cell wall and membrane, hence affecting the distribution of the lipids in the cells [14]. Moreover, ergosterol present in the sample could disrupt the ergosterol biosynthesis pathway in the yeast, causing growth inhibition or cell death. This was further supported by a study of Irshad *et al.* [15], who reported that ergosterol-rich *Cassia fistula* oil significantly decreased the *in vivo* ergosterol content in the *Candida* cell wall.

In this study, the sub-fractions A and B were not cytotoxic to mouse fibroblasts at the concentrations tested (Fig. 1). Animal testing is becoming less popular and is gradually being replaced by *in vitro* methods for toxicity assessment of pharmaceutical products. In conclusion, *P. giganteus* lipids are promising natural products to be further explored as antifungal agents against *Candida* species.

Experimental

Mushroom: The fruiting bodies of *Pleurotus giganteus* (Berk) Karunaratna & K.D. Hyde were obtained from Nas Agro Farm, Selangor, Malaysia. A voucher specimen (KLU-M 1227) was deposited in the Herbarium in the University of Malaya.

Chemicals: Fluconazole and amphotericin B were purchased from Sigma Co. (St. Louis, MO, USA). The stocks were prepared in dimethyl sulfoxide (DMSO) prior to bioassays. [3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] (MTT), was also obtained from Sigma. Methanol (MeOH), ethyl acetate (EtOAc), *n*-hexane and acetone were from Merck (Darmstadt, Germany).

Extracts preparation: The fresh fruiting bodies of *P. giganteus* were sliced, freeze-dried and ground to a fine powder (500 g). The mushroom powder was extracted with 80% MeOH to yield a MeOH extract (115 g, 23.0%). This (125 g) was further partitioned in EtOAc-H₂O (100 mL: 100 mL) to give an EtOAc-soluble extract (6.96 g, 6.05%) and a H₂O extract (74.2 g, 64.52%).

Fractionation of extract: The EtOAc extract (5.00 g) was further fractionated by CC over silica gel. The extract was eluted with *n*-hexane containing increasing concentrations of acetone to obtain 8 fractions (A to H) based on similarity of spots on TLC.

Cell culture: Mouse embryonic fibroblasts were cultured in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10%, v/v, heat-inactivated fetal bovine serum (PAA), 100 U/mL penicillin and 100 µg/mL streptomycin. The cells were routinely passaged every 2-3 days and incubated at 37°C and 5%, v/v, CO₂ in a humidified atmosphere.

Cytotoxicity: The crude MeOH and fractionated EtOAc extracts were dissolved in DMSO (10 mg/mL) as stock solutions. The H₂O extract (10 mg/mL) was stocked in sterilised distilled water. The cytotoxic effects of varying concentrations of MeOH, EtOAc and H₂O extracts, as well as the fractions A-H in DMSO to 3T3 fibroblast cells were tested by the established colorimetric MTT assay [16]. The absorbance was measured at 550 nm using a microplate reader. The IC₅₀ is the concentration of extract or fraction that reduced fibroblast cell growth by 50%.

Anti-yeast activity: *Candida albicans* WM1172, *C. albicans* ATCC90028, *C. dubliniensis*, *C. glabrata* CBS138, *C. glabrata* ATCC90030, *C. krusei* ATCC6258, *C. pseudotropicalis*, and *C. tropicalis* WM30 were used in this study. The yeast inhibition assay was performed according to the method of Macreadie *et al.* [17]. The yeast strains were grown in YEPD (1% yeast extract, 2% peptone, 2% glucose). If required, media were solidified by the addition of 1.5% agar. Yeast inocula (100 µL) with a starting optical density at A₅₉₅ of 0.02-0.04 were added to each well of a 96-well microplate (Orange Scientific, Braine-l'Alleud, Belgium). Mushroom extracts were then added as two-fold serial dilutions commencing with a 100 µg/mL concentration. Fluconazole (0.1

mM) and amphotericin B (1.0 mM) were used as positive controls. A growth control DMSO solvent alone was also included. The microplate was incubated in a microplate shaker at 35°C. After 2 h and 4 h incubation, the A₅₉₅ was recorded using a microplate reader (Sunrise™, Tecan, Austria). Each sample was assayed in triplicate. The lowest concentration of extracts that inhibited growth of *Candida* spp. is the minimum inhibitory concentration (MIC).

Gas chromatography-mass spectrometry (GCMS): GCMS analysis was performed on sub-fractions A and B using Network Gas Chromatography system (Agilent Technologies 6890N) equipped with an Inert Mass Selective Detector (Agilent Technologies 5975) (70eV direct inlet) on a HP-5ms (5% phenyl methyl siloxane) capillary column (30 m × 250 µm × 0.25 µm) initially set at 150°C, then increased at 5°C per min to 300°C and held for 10 min. Helium was used as carrier gas at a flow rate of 1 mL per min. The total ion chromatogram obtained was autointegrated by chemstation and the constituents were identified by comparison with the accompanying mass-spectra database (Wiley 9th edition with NIST 11 Mass Spectral Library, USA) wherever possible.

Acknowledgments – This research is supported by UM High Impact Research Grant UM-MOHE UM.C/625/1/HIR/MOHE/F00002-21001 from the Ministry of Higher Education Malaysia. The authors thank the University of Malaya for Postgraduate Research Grant (PV007/2012A) and MRC 66-02-03-0074. We thank Prof. Andrew T. Smith, Dean of School of Applied Science, RMIT University for partial funding for the joint research.

References

- [1] Pfaller MA, Pappas PG, Wingard JR. (2006) Invasive fungal pathogens: current epidemiological trends. *Clinical Infectious Diseases*, **43**, S3–14.
- [2] Barros FMC, Pippi B, Dresch RR, Dauber B, Luciano SC, Apel MA, Fuentefria AM, von Poser GL. (2013) Antifungal and antichemotactic activities and quantification of phenolic compounds in lipophilic extracts of *Hypericum* spp. native to South Brazil. *Industrial Crops and Products*, **44**, 294–299.
- [3] Shao P-L, Huang L-M, Hsueh P-R. (2007) Recent advances and challenges in the treatment of invasive fungal infections. *International Journal of Antimicrobial Agents*, **30**, 487–495.
- [4] Casaroto AR, Lara VS. (2010) Phytomedicines for *Candida*-associated denture stomatitis. *Fitoterapia*, **81**, 323–328.
- [5] Santoyo S, Ramirez-Anguiano AC, Reglero G, Soler-Rivas C. (2009) Improvement of the antimicrobial activity of edible mushroom extracts by inhibition of oxidative enzymes. *International Journal of Food Science & Technology*, **44**, 1057–1064.
- [6] Vamanu E. (2013) *In vitro* antioxidant and antimicrobial activities of two edible mushroom mycelia obtained in the presence of different nitrogen sources. *Journal of Medicinal Food*, **16**, 155–166.
- [7] Lai TK, Biswas G, Chatterjee S, Dutta A, Pal C, Banerji J, Bhuvanesh N, Reibenspies JH, Acharya K. (2012) Leishmanicidal and anticandidal activity of constituents of Indian edible mushroom *Astraeus hygrometricus*. *Chemistry & Biodiversity*, **9**, 1517–1524.
- [8] Wong JH, Ng TB, Wang H, Sze SCW, Zhang KY, Li Q, Lu X. (2011) Cordymin, an antifungal peptide from the medicinal fungus *Cordyceps militaris*. *Phytomedicine*, **18**, 387–92.
- [9] Mortimer PE, Karunarathna SC, Li Q, Gui H, Yang X, Yang X, He J, Ye L, Guo J, Li H, Sysouphanthong P, Zhou D, Xu J, Hyde KD. (2012) Prized edible Asian mushrooms: ecology, conservation and sustainability. *Fungal Diversity*, **56**, 31–47.
- [10] Wong W-L, Abdulla MA, Chua K-H, Kuppusamy UR, Tan Y-S, Sabaratnam V. (2012). Hepatoprotective effects of *Panus giganteus* (Berk.) Corner against thioacetamide-(TAA-) induced liver injury in rats. *Evidence-based Complementary and Alternative Medicine*, Article ID 170303.
- [11] Phan CW, Wong WL, David P, Naidu M, Sabaratnam V. (2012). *Pleurotus giganteus* (Berk.) Karunarathna & K. D. Hyde : Nutritional value and *in vitro* neurite outgrowth activity in rat pheochromocytoma cells. *BMC Complementary and Alternative Medicine*, **12**(102).
- [12] Hou CT. (2008). New bioactive fatty acids. *Asia Pacific Journal of Clinical Nutrition*, **17**, 192–195.
- [13] Agoramoorthy G, Chandrasekaran M, Venkatesalu V, Hsu MJ. (2007) Antibacterial and antifungal activities of fatty acid methyl esters of the blind-your-eye mangrove from India. *Brazilian Journy of Microbiology*, **38**, 739–742.
- [14] Bakkali F, Averbeck S, Averbeck D, Idaomar M. (2008) Biological effects of essential oils-A review. *Food and Chemical Toxicology*, **46**, 446–475.
- [15] Irshad MD, Ahmad A, Zafaryab MD, Ahmad F, Manzoor N, Singh M, Rizvi MM. (2013) Composition of *Cassia fistula* oil and its antifungal activity by disrupting ergosterol biosynthesis. *Natural Product Communications*, **8**, 261–264.
- [16] Mosmann T. (1983) Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunological Methods*, **65**, 55–63.
- [17] Macreadie IG, Avery TD, Robinson TV., Macreadie P, Barraclough M, Taylor DK, Tiekink ERT. (2008) Design of 1,2-dioxines with anti-*Candida* activity: aromatic substituted 1,2-dioxines. *Tetrahedron*, **64**, 1225–1232.

Effect of Trehalose Addition on Volatiles Responsible for Strawberry Aroma

Mirela Kopjar^{a*}, Janez Hribar^b, Marjan Simčič^b, Emil Zlatič^b, Tomaž Požrl^b and Vlasta Piližota^a

^aFaculty of Food Technology, F. Kuhača 20, 31000 Osijek, Croatia

^bBiotechnical Faculty, Jamnikarjeva 101, 1000 Ljubljana, Slovenia

mirela.kopjar@ptfos.hr

Received: July 4th, 2012; Accepted: October 2nd, 2013

Aroma is one of the most important quality properties of food products and has a great influence on quality and acceptability of foods. Since it is very difficult to control, in this study the effect of addition of trehalose (3, 5 and 10%) to freeze-dried strawberry cream fillings was investigated as a possible means for retention of some of the aroma compounds responsible for the strawberry aroma. In samples with added trehalose, higher amounts of fruity esters were determined. Increase of trehalose content did not cause a proportional increase in the amount of fruity esters. However, results of our research showed that trehalose addition did not have the same effect on both γ -decalactone and furaneol.

Keywords: Aroma compounds, Trehalose addition, Freeze-dried strawberry cream filling.












Next to color and texture, aroma is one of the most important quality properties, and thus its stability in different foods has been of increasing interest due to its strong relationship with consumer's acceptability of foods. However, it is very difficult to control. Different factors, like processing, storage, addition of ingredients and packaging materials, can cause modifications of overall aroma by either reducing aroma compound intensity or producing off-flavor components. Incorporation of small amounts of aroma compounds into foods can greatly influence the final product quality, cost and consumer satisfaction. The food industry is continuously developing ingredients, processing methods, and packaging materials to improve aroma preservation and delivery [1]. Strawberry is a very delicious fruit, growing in nearly all countries of the world. Due to its typical, very attractive aroma and color, strawberry has always been a favored raw material for production of different fruit product such as jams, strawberry cream fillings, jelly, and its aroma has often been analysed. Volatile components of strawberries have been extensively studied and more than 360 volatiles are assumed to be involved in strawberry aroma. A complex mixture of esters, aldehydes, alcohols and sulfur compounds mainly determines strawberry aroma, but esters are quantitatively and qualitatively the most important class of volatiles [2]. During preparation of fruit products, different additives are used influencing food product physicochemical properties, aroma, texture and color. Carbohydrates are known to enclose volatile compounds [2-6], thus in this study the influence of trehalose on selected aroma compounds was investigated. Trehalose (α , α -trehalose) is a disaccharide formed by an α -1,1 linkage of two d-glucose molecules. It is a non-reducing sugar that is not easily hydrolysed by acid, and the glycosidic bond is not cleaved by α -glucosidase. The molecular formula and weight are $C_{12}H_{22}O_{11}$ and 342.31, respectively [7,8]. Due to its natural functions and technical qualities, trehalose could be applied in the food, cosmetic and medical industries [9-11]. Influence of trehalose addition on overall aroma, color and texture in strawberry cream filling was investigated in our previous article [5].

The objective of this study was to give broader insight of the influence of trehalose addition and its amount on selected fruity

esters, γ -decalactone and furaneol (the compounds that are the most responsible for strawberry aroma) in strawberry cream fillings. Properties and structure of selected compounds are presented in Table 1.

In samples without aroma addition (Figure 1), ethyl acetate and ethyl butanoate were detected, while ethyl pentanoate and ethyl hexanoate were not. In samples with trehalose addition, higher amounts of ethyl acetate and ethyl butanoate were observed in comparison with samples without trehalose addition. The highest amounts of both esters were observed when 5% trehalose was added, while samples with 10% trehalose addition had the lowest amount of these esters. With 3% trehalose addition, a sharp increase in the amounts of methyl butanoate and methyl hexanoate was observed. The highest amount of these two esters was observed when 5% of trehalose was added. In samples with the highest amount of trehalose addition (10%), in both cases, a decrease in amount was observed, but still the amount of methyl esters was higher than in samples without trehalose addition. Ethyl 2-methyl butanoate was detected in samples without trehalose addition, while ethyl 3-methyl butanoate was detected only in the samples with trehalose addition. Addition of trehalose caused an increase in the ethyl 2-methyl butanoate amount. Samples with addition of 3 and 5% of trehalose had almost the same amount, while addition of 10% of trehalose did not cause retention of ethyl 2-methyl butanoate. Increase in the amount of γ -decalactone and furaneol was not observed with trehalose addition; actually, with trehalose addition, the amount of γ -decalactone was lower. In samples with aroma addition more aroma compounds were detected (Figure 2). In samples with 3% of trehalose addition, ethyl acetate was determined in the highest amount, but with further addition, a reverse effect occurred. In samples with addition of 5 and 10% of trehalose lower amounts of this ester were observed than in samples without trehalose addition. In the case of ethyl butanoate, the highest amount was observed in samples with 5% of trehalose addition, while for ethyl pentanoate and ethyl hexanoate the highest amount was observed with 3% of trehalose addition. Methyl hexanoate was not detected and methyl butanoate was recorded only in samples with 5 and 10% of trehalose addition.

Table 1: Investigated strawberry aroma compounds and their properties.

Aroma compound	CAS number	MW	Vapour pressure mm/Hg	logP (o/w)	Description	Structural formula
Ethyl acetate	141-78-6	88.11	112.0	0.71	Etherial, fruity, sweet, grape and rum-like	
Ethyl butanoate	105-54-4	116.16	12.8	1.85	Fruity, juicy, pineapple, cognac	
Ethyl pentanoate	539-82-2	130.18	4.75	2.3	Sweet, fruity, acidic, pineapple, apple, green, berry and tropical	
Ethyl hexanoate	123-66-0	144.21	1.66	2.83	Sweet, fruity, pineapple, waxy, fatty and estery with a green banana nuance	
Methyl butanoate	623-42-7	102.13	31.1	1.24	Pungent, etherial, fruity, perfumey and fusel with a fermented, cultured, creamy undertone of fruity apple sweet banana pineapple	
Methyl hexanoate	106-70-7	130.13	3.95	2.3	Fruity, pineapple, ether	
Ethyl 2-methyl butanoate	7452-79-1	130.18	7.85	2.12	Fruity, estery and berry with fresh tropical nuances	
Ethyl 3-methyl butanoate	108-64-5	130.18	7.85	2.12	Sweet, diffusive, estery, fruity, sharp, pineapple, apple, green and orange	
Ethyl 2-methyl pentanoate	39255-32-8	144.21	2.91	2.65	Fruity, green, melon and waxy with a fatty nuance	
γ -decalactone	706-14-9	170.25	0.00512	2.72	Fresh, oily, waxy, peach, coconut, buttery, sweet	
Furaneol	3658-77-3	128.13	0.032	0.34	Sweet, slightly burnt brown caramelised, cotton candy with a savoury nuance	

Data were obtained from <http://www.thegoodscentscompany.com> and <http://www.chemicalbook.co>

The ethyl 2-methyl butanoate amount increased with increase of trehalose amount and the highest effect was with the addition of 5% trehalose. The behavior of ethyl 3-methyl butanoate was very interesting and had a different tendency from ethyl 2-methyl-butanoate. Samples without and with 3 and 10% trehalose addition had the same amount of ethyl 3-methyl butanoate, while the sample with 5% trehalose addition had a higher amount of this aroma compound. The tendency of ethyl 2-methyl pentanoate was not observed for any other ester. With addition of trehalose there was a decrease in the ethyl 2-methyl pentanoate amount. Decrease of γ -decalactone and furaneol amounts occurred in samples with trehalose addition.

Comparison of samples with native strawberry aroma (without addition of aroma) with samples with addition of strawberry aroma showed that the initial concentration of aroma compounds was very important, but also a different trend was observed in samples with trehalose addition in these two types of samples. As a consequence of previous studies [12,13], a possible explanation for the different behavior of aroma compounds in samples without and with strawberry aroma addition could be competitive and cooperative effects of the aroma compounds. Results illustrated in Figure 3 were obtained by statistical analysis (using CA and MANOVA/ANOVA) and from them it is evident that the amount of aroma compounds in samples depended on their initial concentration, as well as on the amount of trehalose addition.

The real mechanism of trehalose action is still not known. Three theories have been put forward to explain the mechanism of the action of trehalose: 1) water replacement hypothesis, 2) glass transformation hypothesis, and 3) chemical stability hypothesis. The ability of sugar molecules to bind protectively onto the surface of

molecular structures has been ascribed also to their ability to form hydrogen bonds, the so called 'water replacement hypothesis'. Unlike most other disaccharides, trehalose has no direct internal hydrogen bonds. All four internal bonds are indirectly connected via the two water molecules that form part of the native dihydrate structure. This arrangement gives to the molecule an unusual flexibility around the disaccharide bond, which may allow trehalose to fit more closely to the irregular surface of macromolecules than other, more rigid disaccharides, in which the rings are directly hydrogen bonded to each other [9]. The greater bioprotective action of trehalose on biological structures in comparison with sucrose and maltose was connected with its greater destructuring effect on the tetrahedral H-bond network of water [14]. The trehalose system showed the smallest partial molar volume value, indicative of a more packed conformation, together with a greater partial volume increase with temperature, which indicates a greater structural sensitivity in contrast to sucrose and maltose solutions. From a biological point of view, these findings could imply a greater ability of trehalose to encapsulate biomolecules in more rigid and packed structures and hence a greater bioprotector effectiveness of trehalose in respect to sucrose [15].

Formulation of food products is very important since the composition of the matrix strongly affects the quality of the foods, and its stability. Retention of aroma compounds is a very complex phenomenon, as can be seen from our results. It depends on matrix composition and interactions between volatile and non-volatile constituents of the matrix, as well as on the structure of aroma compounds and their initial concentration. Small modifications (like replacement of certain amount of sucrose with trehalose and addition of strawberry aroma) of food matrix composition greatly affect the retention of the aroma compounds in strawberry cream




Figure 1: Total peak areas of aroma compounds in freeze-dried samples without aroma addition.




Figure 2: Total peak areas of aroma compounds in freeze-dried samples with aroma addition.




Figure 3: Dendrogram of cluster analysis of aroma compounds in samples without aroma addition (a) and with aroma addition (b).

Experimental

Sample preparation: Raw material for preparation of strawberry cream fillings without and with trehalose addition (3, 5 and 10% as sucrose replacement) were obtained from the food company Fructal d.d. (Ajdovščina, Slovenia), where ingredients (commercial frozen

strawberry puree, starch, vegetable fat, sucrose, glucose syrup, sorbitol) were mixed together according to an industrial recipe. The total solids of the mixture of ingredients were 40%. Samples of strawberry cream fillings were prepared by freeze-drying of mixture of ingredients with 40% of total solids until 76% of total solids were achieved. Freeze drying of samples was performed in a Christ

Freeze Dryer (Gamma 2–20, Germany). This required a drying time of 50 h. The following conditions were used: freezing temperature -20°C ; temperature of sublimation from -20 to 0°C , and vacuum of 0.630 mbar. The temperature of an isothermal desorption was from 0 to 20°C under the vacuum of 0.01 mbar. Strawberry cream fillings were prepared without and with addition of strawberry aroma (as prepared in industry). Aroma was added after strawberry cream filling preparation. Samples were homogenised on a metallic plate and, after that, cream fillings were left 10 days to stabilise at room temperature. Sampling for aroma compounds evaluation was conducted before addition of strawberry aroma and after 10 days of sample stabilisation. Samples were prepared in triplicates.

Determination of aroma compounds: GC-MS analyses were carried out for evaluation of the fruity esters, γ -decalactone and furaneol (Table 1). Extraction of volatiles was carried out with a microextraction needle 85 μm Carboxen/PDMS (Supelco) at 50°C for 40 min. For analyses, a 6890N instrument (Agilent, SAD) equipped with a MS 5971A detector (Hewlett Packard, SAD) was used. Compounds were analyzed on a ZB-WAX column (60 m x 0.32 mm x 0.5 μm , Phenomenex). The temperature programming was as follows: 5 min at 40°C , temperature gradient $4^{\circ}\text{C}/\text{min}$, and final temperature 230°C for 5 min. The carrier gas was helium with

a flow rate of 1 mL/min at 40°C . Desorption of adsorbed volatiles was carried out by exposing the fiber in the injector port of the GC at 270°C for 5 min. For thermal desorption, the splitless injection mode was used and the split valve was opened after 0.5 min. Mass spectra were obtained with 70 eV electron impact ionization, while the mass spectrometer was continuously scanning m/z 30–300. Determination of the analysed compounds was confirmed by retention times of single compounds and from bibliographic data. The results were expressed as total peak area.

Statistical analyses: For evaluation of the influence of strawberry aroma addition and trehalose amount, total peak area of GC determinations of all investigated aroma compounds were taken into account and data were evaluated using General ANOVA/MANOVA (one-way ANOVA) and Cluster Analysis (CA). Statistical analyses were carried out using software program STATISTICA 8 (StatSoft, Inc, USA).

Acknowledgments - This work was supported by research on Croatian-Slovenian joint project and National Scientific Project financed by the Croatian Ministry of Science, Education and Sports. We would like to thank Fructal d.d. (Ajdovščina, Slovenia) for providing raw material for strawberry paste preparation.

References

- [1] Madene A, Jacquot M, Scher J, Desobry S. (2006) Flavour encapsulation and controlled release. Review. *International Journal of Food Science and Technology*, **41**, 1–21.
- [2] Komes D, Lovrić T, Kovačević Ganić K, Gracin L. (2003) Study of trehalose addition on aroma retention in dehydrated strawberry puree. *Food Technology and Biotechnology*, **41**, 111–119.
- [3] Komes D, Lovrić T, Kovačević Ganić K, Gajdos Kljusurić J, Banović M. (2005) Trehalose improves flavour retention in dehydrated apricot puree. *International Journal of Food Science and Technology*, **40**, 425–435.
- [4] Komes D, Lovrić T, Kovačević Ganić K. (2007) Aroma of dehydrated pear products. *LWT-Food Science and Technology*, **40**, 1578–1586.
- [5] Kopjar M, Piližota V, Hribar J, Simčič M, Zlatič E, Nedić Tiban N. (2008) Influence of trehalose addition and storage conditions on the quality of strawberry cream filling. *Journal of Food Engineering*, **87**, 341–350.
- [6] Galmarini MV, Scheebor C, Zamora MC, Chirife J. (2009) The effect of trehalose, sucrose and maltodextrin addition on physicochemical and sensory aspects of freeze-dried strawberry puree. *International Journal of Food Science and Technology*, **43**, 510–516.
- [7] Birch GG. (1963) Trehalose. In *Advances in Carbohydrate Chemistry*, Vol. **18**. Academic Press, New York, 201–225.
- [8] Elbein AD. (1974) The metabolism of α,α -trehalose. In *Advances in Carbohydrate Chemistry and Biochemistry*, Vol. 30. Academic Press, New York, 227–256.
- [9] Colaço CALS, Roser B. (1995) Trehalose - a multifunctional additive for food preservation. In *Food Packaging and Preservation*. Blackie Professional, London, 123–140.
- [10] Ohtake, S, Wang YJ. (2011) Trehalose: Current use and future applications. *Journal of Pharmaceutical Science*, **100**, 2020–2053.
- [11] Magazu S, Migliardo F. (2010) Molecular mechanisms of bioprotection process by trehalose. *Current Nutrition and Food Science*, **6**, 157–160.
- [12] Van Ruth SM, King C (2003) Effect of starch and amylopectin concentrations on volatile flavour release from aqueous model food systems. *Flavour and Fragrance Journal*, **18**, 407–416.
- [13] Rutschmann MA, Solms J. (1990) Formation of inclusion complexes of starch with different organic compounds. III. Study of ligand binding in binary model systems with (-) limonene. *LWT-Food Science and Technology*, **23**, 80–83.
- [14] Branca C, Magazu S, Maisano G, Migliardo P. (1999) Anomalous cryoprotective effectiveness of trehalose: Raman scattering evidences. *The Journal of Chemical Physics*, **111**, 281–287.
- [15] Branca C, Magazu S, Maisano G, Migliardo P. (2000) Experimental study of the hydration properties of homologous disaccharides. *Journal of Biological Physics*, **26**, 295–306.

***Pogostemon hirsutus* Oil, Rich in Abietane Diterpenes**Ramar Murugan^{a*}, Gopal Rao Mallavarapu^b, Veerappan Sudha^a and Pemaiah Brindha^a^aCentre for Advanced Research in Indian System of Medicine, School of Chemical and Biotechnology, SASTRA University, Tirumalaisamudram, Thanjavur 613 401, India^b602, A-Block, Renaissance Temple Bells, Opp. ISKCON Temple, Yeshwantpur, Bangalore 560 022, India

ramarmurugan@yahoo.com

Received: August 22nd, 2013; Accepted: September 26th, 2013

Volatile oil extracted from the leaves of *Pogostemon hirsutus* Benth., a rare species from Anamalai hills of the Western Ghats of Tamil Nadu, India was analyzed by GC-FID and GC-MS. Fifty-four compounds amounting to 90.8% of the oil were identified. The oil was rich in monoterpenes and diterpenes. The major components of the oil were α -pinene (11.5%), sabinene (8.5%), β -pinene (5.2%), abietatriene (16.3%), dehydroabietal (3.5%) and dehydroabietol (21.0%). Abietane diterpenes are reported for the first time in the genus *Pogostemon* of Lamiaceae with six abietanes representing 42.6% of the oil.

Keywords: *Pogostemon hirsutus*, Lamiaceae, Essential oil, Diterpenes, Abietatriene, Dehydroabietol.

The genus *Pogostemon* Desf. of Lamiaceae is represented by 97 species globally and 54 species in India [1a]. *Pogostemon cablin* (Blanco) Benth. is the only species in this genus commercially cultivated for its essential oil, known as patchouli, which is extensively used in perfumery. As part of our search for indigenous wild *Pogostemon* species whose essential oils can be utilized in perfumery, we have studied and reported the chemical constituents of the essential oils of three indigenous and endemic *Pogostemon* species such as *P. heyneanus* Benth., *P. travancoricus* Bedd. var. *travancoricus* and *P. speciosus* Benth. [1b,c,d]. In continuation of our studies on the bioprospecting of *Pogostemon* species, we have now investigated the volatile constituents of the leaves of *P. hirsutus* collected from the Anamalai hills of the Western Ghats, India. The aim of this study was to explore the possibility of commercial utilization of the wild aromatic plants of the Western Ghats based on the chemical composition of the essential oils and olfactive characteristics.

Pogostemon hirsutus was previously considered to be endemic to Sri Lanka, but later it was discovered from the Anamalai hills of the Western Ghats and reported as a new distributional record for India [1e]. It is a small hirsute herb, growing about a feet high, in humid, rocky slopes in high altitude grassland above 2000 m altitude. Both fresh and dried leaves are slightly aromatic. The inflorescence is simple, terminal spike with white flowers. A detailed description along with a photograph is given in our earlier report [1e].

Hydro-distillation of the dried leaf of *P. hirsutus* yielded 0.4% of essential oil on dry weight basis. The oil was yellow with a strong earthy odor. GC analysis of the oil showed more than 100 peaks. GC-FID and GC-MS analyses of the oil enabled the identification of 54 compounds representing 90.8% of the oil. The identified compounds with their RIs and relative concentration in the oil are listed in Table 1. The oil was rich in monoterpenes and diterpenes and particularly dominated by abietanes. The concentration of sesquiterpenes was minimum, which constituted less than 8% of the oil. The major components of the oil were α -pinene (11.5%), sabinene (8.5%), β -pinene (5.2%), abietatriene [1,2,3,4,4a,9,10,10a-octahydro-1,1,4a-trimethyl-7-(1-methylethyl)-, (4aS-trans)-phenanthrene] (16.3%), dehydroabietal [1,2,3,4,4a,9,10,10a-octahydro-1,4a-dimethyl-7-(1-methylethyl)-, (1R-(1 α ,4 α ,10 α))-1-phenanth-

Table 1: Composition of the essential oil of *P. hirsutus*.

Compound	RI		Area %	Method of identification
	Observed	literature		
α -Thujene	930	930	1.3	GC, GC-MS
α -Pinene	940	939	11.5	GC, GC-MS
Camphene	955	954	0.5	GC, GC-MS, PE
Sabinene	980	975	8.5	GC, GC-MS, PE
β -Pinene	982	979	5.2	GC, GC-MS, PE
β -Myrcene	992	991	0.1	GC, GC-MS, PE
<i>p</i> -Cymene	1024	1024	1.0	GC, GC-MS, PE
Limonene	1030	1029	0.3	GC, GC-MS, PE
γ -Terpinene	1062	1059	0.1	GC, GC-MS
<i>n</i> -Octanol	1070	1068	0.1	GC, GC-MS, PE
Terpinolene	1088	1088	0.2	GC, GC-MS
Linalool	1097	1096	0.2	GC, GC-MS, PE
<i>cis</i> -Verbenol	1139	1141	0.1	GC, GC-MS
<i>trans</i> -Pinocarveol	1141	1139	0.3	GC, GC-MS
<i>trans</i> -Verbenol	1146	1144	0.2	GC, GC-MS
Pinocarvone	1162	1164	0.1	GC, GC-MS
Terpinen-4-ol	1179	1177	0.8	GC, GC-MS, PE
Napthalene	1182	1178	0.2	GC, GC-MS, PE
<i>cis</i> -Pinocarveol	1185	1184	0.1	GC, GC-MS
α -Terpineol	1190	1191	0.2	GC, GC-MS, PE
Myrtenal	1196	1195	0.1	GC, GC-MS
<i>p</i> -Cymen-7-ol	1290	1290	0.2	GC, GC-MS
<i>trans</i> -Pinocarvyl acetate	1299	1298	0.2	GC, GC-MS
<i>cis</i> -Pinocarvyl acetate	1310	1312	0.2	GC, GC-MS
α -Cubebene	1350	1348	0.1	GC, GC-MS
α -Ylangene	1373	1375	0.1	GC, GC-MS
α -Copaene	1378	1376	2.5	GC, GC-MS
β -Cubebene	1386	1388	0.1	GC, GC-MS
β -Elemene	1390	1390	0.1	GC, GC-MS
(<i>E</i>)-Caryophyllene	1418	1418	0.1	GC, GC-MS
α -Guaiane	1437	1439	0.1	GC, GC-MS
α -Selinene	1495	1498	0.1	GC, GC-MS
<i>trans</i> - β -Guaiane	1500	1502	0.3	GC, GC-MS
β -Sesquiphellandrene	1517	1522	0.1	GC, GC-MS
δ -Cadinene + <i>trans</i> -Calamenene	1523	1524	1.4	GC, GC-MS
α -Cadinene	1540	1538	0.2	GC, GC-MS
(<i>E</i>)-Nerolidol	1561	1563	0.3	GC, GC-MS
Caryophyllene oxide	1582	1583	0.4	GC, GC-MS
Cubanol	1646	1649	0.1	GC, GC-MS
α -Cadinol	1652	1652	0.1	GC, GC-MS
Pogostone	1702	1702	0.8	GC, GC-MS
Isopimara-9(11),15-diene	1903	1905	2.8	GC, GC-MS
Pimara-8(14),15-diene	1953	1949	1.1	GC, GC-MS
Sclarene	1971	1974	0.1	GC, GC-MS
Kaur-15-ene	2000	1997	1.0	GC, GC-MS
Abietatriene	2054	2056	16.3	GC, GC-MS
Abieta-8,13-diene	2090	2087	0.2	GC, GC-MS
<i>p</i> -(3,4-Dihydro-2,2,4-trimethyl-2H-1-benzopyran-4-yl)-phenol	2111	--	1.5	GC-MS
Abita-8(14),13(15)diene	2155	2154	0.7	GC, GC-MS
13 α -Methyl,13-vinylpodocarp-7-en-3 α -ol	2160	--	2.5	GC-MS
Sclareol	2222	2223	0.6	GC, GC-MS
Dehydroabietal	2280	2275	3.5	GC, GC-MS
Abietal	2311	2313	0.9	GC, GC-MS
Dehydroabietol	2368	2368	21.0	GC, GC-MS
Total			90.8	

RI - Retention Index; PE - peak enrichment on co-injection with standard.

rene carboxaldehyde] (3.5%) and dehydroabietol [1,2,3,4,4a,9,10,10a-octahydro-1,4a-dimethyl-7-(1-methylethyl)-, (1*R*-(1 α ,4 α ,10 α))-1-phenanthrenemethanol] (21.0%).

The composition of *P. hirsutus* oil is quite different from that of the oil of *P. cablin* (patchouli oil) which is widely used in perfumery and flavours. Patchouli oil contains mainly sesquiterpenes and oxygenated sesquiterpenes and the main component being patchouli alcohol. On the contrary, the oil of *P. hirsutus* contained mostly monoterpenes and diterpenes. Patchouli alcohol was not detected in the oil. However, pogostone which is one of the important constituents of patchouli oil was found in the essential oil of *P. hirsutus*. The main component of *P. hirsutus* oil was dehydroabietol, which is reported to possess antiplasmodial activity and found to inhibit the growth of chloroquine-sensitive as well as chloroquine-resistant strains of *Plasmodium falciparum* that causes malarial fever [2].

Diterpenes are widespread in conifers. However, Asteraceae and Lamiaceae are the two families in angiosperms rich in diterpenes. They have been used in chemotaxonomic studies in conifers and in some families in angiosperms especially in Lamiaceae [3-8]. Abietanes are a large class of tricyclic diterpenoids, commonly found in conifers and in some angiosperms. In Lamiaceae, they have been reported in genera such as *Coleus*, *Hyptis*, *Lepechinia*, *Nepeta*, *Plectranthus*, *Rabdosia*, *Salvia* and *Stachys* [3,5,8,9]. However, abietanes have not been reported so far in the genus *Pogostemon*. In the present study, six abietane compounds representing 42.6% of the essential oil of *P. hirsutus* were identified. Therefore, the high content of abietanes in the essential oil of this species is significant.

Experimental

Plant material: Leaf sample of *P. hirsutus* was collected from Akkamalai of Valparai region of Anamalai hills of the Western

Ghats, Tamil Nadu, India in the month of January. Herbarium specimens (*R. Murugan 21*) were prepared for identification and deposited in the Herbaria of Madras Christian College (MCCH), Chennai and Institute of Ayurveda and Integrative Medicine (FRLH), Bangalore, India. The species was identified and confirmed by matching with the authentic herbarium specimens including *Type specimens* deposited in the Herbaria of the Botanical Survey India (CAL), Kolkata and the Royal Botanic Gardens (K), London.

Extraction of oil: The collected leaf sample was dried in the shade for about three weeks and subjected to hydro-distillation in a Clevenger-type apparatus for about 8 hours. Yellow coloured essential oil was obtained and dried over anhydrous Na₂SO₄. The oil was stored in refrigerator at 5°C until analyzed.

Analyses: The GC-FID and GC-MS analyses of the essential oil were done by the method adopted in our previously published report [1d]. Component identification was done by comparison of the retention indices (RIs) of the GC peaks obtained using homologous series of *n*-alkanes (C₈-C₂₅) with those of compounds reported in literature [10] and also by comparing the mass spectra of the peaks with standards reported in literature and matching the mass spectra of the compounds available in NIST and Wiley libraries. Peak enrichment (PE) on co-injection with available authentic standards was also done to confirm the identification. Peak area percentages were calculated from FID response without the use of correction factors.

Acknowledgments - We are grateful to the Hon'ble Vice-chancellor and Dean, School of Chemical and Biotechnology, SASTRA University for facilities and encouragement. We thank the Principal Chief Conservator of Forests, Tamil Nadu for granting permission to survey and to collect leaf sample.

References

- (a) Murugan R, Livingstone C. (2010) *Pogostemon raghavendranii* (Lamiaceae), a new species from Anamalai hills, India. *Rheedea*, **20**, 21-24.(b) Murugan R, Mallavarapu GR, Padmashree, KV, Rao RR, Livingston C. (2010) Volatile oil composition of *Pogostemon heyneanus* Benth., and comparison of its composition with patchouli oil. *Natural Product Communications*, **5**, 1961-1964. (c) Murugan R, Mallavarapu GR. (2012) Composition of the essential oil of *Pogostemon travancoricus* var. *travancoricus*. *Natural Product Communications*, **7**, 87-88. (d) Murugan R, Mallavarapu GR. (2013) α -Bisabolol, the main constituent of the oil of *Pogostemon speciosus*. *Industrial Crops and Products*, **49**, 237-239. (e) Murugan R, Ravikuman K, Livingston C. (2010) Addition of a wild aromatic plant *Pogostemon hirsutus* Benth. to the flora of India. *Current Science*, **95**, 835.
- Ziegler HL, Jensen TH, Christensen J, Staerk D, Hagerstrand H, Sittie AA, Olsen CE, Staalso T, Ekpe P, Jeroszewski JW. (2002) Possible artifacts in the *in vitro* determination of antimalarial activity of natural products that incorporate into lipid bilayer: Apparent antimalarial activity of dehydroabietinol, a constituent of *Hyptis suaveolens*. *Planta Medica*, **68**, 547-549.
- Otto A, Wilde V. (2011) Sesqui, Di and Triterpenoids as chemosystematic markers in extant conifers-A review. *The Botanical Review*, **67**, 141-238.
- Menezes F de S, Kaplan MAC. (1993) Ocorrências de diterpenóides na família Lamiaceae. *Revista Brasileira de Ciências Farmacêuticas*, **74**, 45-46.
- Alvarenga SAV, Gastmans JP, Rodrigues G de V, Moreno PRH, Emerenciano V de P. (2001) A computer-assisted approach for chemotaxonomic studies – diterpenes in Lamiaceae. *Phytochemistry*, **56**, 583-595.
- Figueiredo MR, Kaplan MAC, Gottlieb OR. (1995) Diterpenes, taxonomic markers? *Plant Systematics and Evolution*, **195**, 149-158.
- Menezes F de S, Kaplan MAC, Gottlieb OR. (1995) Chemo-evolutionary polarization in the Lamiiflorae. *Anais da Academia Brasileira de Ciências*, **67**, 455-460.
- Ulubelen A, Topcu G. (1992) Abietane diterpenoids from *Salvia pomifera*. *Phytochemistry*, **31**, 3941-3951.
- Piozzi F, Bruno M. (2009) Diterpenoids in the essential oils from the genus *Stachys*. *Records of Natural Products*, **3**, 120-125.
- Adams RP. (1995) Identification of essential oil components by Gas Chromatography / Quadrupole Mass Spectroscopy, Allured Publishing Corporation, Carol Stream, IL, USA.

Combined Analysis of the Root Bark Oil of *Cleistopholis glauca* by Chromatographic and Spectroscopic Techniques

Zana A. Ouattara^{a,b}, Jean Brice Boti^b, Coffy Antoine Ahibo^b, Félix Tomi^a, Joseph Casanova^a and Ange Bighelli^{a*}

^aUniversité de Corse-CNRS, UMR 6134 SPE, Equipe Chimie et Biomasse, Route des Sanguinaires, 20000 Ajaccio, France

^bLaboratoire de Chimie Organique Biologique, UFR-SSMT, Université Félix Houphouët Boigny-Abidjan, BPV 34 Abidjan, Côte d'Ivoire

ange.bighelli@univ-corse.fr

Received: August 15th, 2013; Accepted: October 9th, 2013

The composition of root bark oil from *Cleistopholis glauca* Pierre ex Engler & Diels growing wild in Ivory Coast was investigated by GC (in combination with retention indices) and ¹³C NMR spectroscopy after partition of hydrocarbons and oxygenated compounds on silica gel. Thirty-one compounds have been identified. *C. glauca* produces a sesquiterpene-rich oil, patchoulone (33.5%), cyperene (9.5%) and germacrene D (6.6%) being the main components. Special attention was paid to the identification and quantification of germacrene C (a heat-sensitive compound) and δ -elemene, which were achieved by a combination of GC(FID) and ¹³C NMR spectroscopy. The composition of *C. glauca* root bark and leaf oils differed drastically.

Keywords: *Cleistopholis glauca*, Root bark oil, Essential oil composition, Germacrene C, Patchoulone.

Cleistopholis glauca Pierre ex Engler & Diels (*Annonaceae*) is native to central Africa, growing wild in rainforest and in secondary bush, as well as along waterways. It is widespread in Cameroon, Central African Republic, Gabon, Congo-Kinshasa (actually Democratic Republic of Congo) and Angola. It was introduced thirty years ago in the Petit Yapo forest (South-Eastern Ivory Coast). *C. glauca* is a tree up to 35m high and up to 80 cm in diameter, with a cylindrical bole straight trunk covered with a fibrous bark. The bark of *C. glauca* is used in the manufacture of African huts, and fibers make excellent strings. In decoction, it is used as an emetic. The macerated bark of stems and roots is used as a vermifuge in Central African Republic [1,2].

Structural elucidation has been reported of compounds isolated from solvent extracts of *C. glauca*: oligorhamnosides [3,4]; lipids [5]; terpenes [4,5]; a phenyl propanoid [6]; and a furanic derivative [6]. The chemical composition of *C. glauca* leaf oil has been recently investigated by our group [7]. Oil samples isolated from leaves collected from individual trees were analyzed by gas chromatography (GC) using a flame ionisation detector (FID), and the compounds were characterized by a combination of retention indices (RI) and ¹³C NMR spectroscopy. *C. glauca* produced a sesquiterpene-rich leaf oil, germacrene D (16.4-46.5%) and (*E*)- β -caryophyllene (8.0-26.2%) being the major components. In a few samples, monoterpene hydrocarbons were present in appreciable amounts: myrcene (up to 39.7%), β -pinene (up to 24.8%) and α -pinene (up to 16.4%).

In the present paper, we report on the chemical composition of a root bark oil sample from *C. glauca*, investigated by a combination of chromatographic and spectroscopic techniques and that differed drastically from the leaf oils of the same species. Identification and quantitative determination, by a combination of GC (FID) and ¹³C NMR spectroscopy, of a heat sensitive compound, germacrene C, was investigated in detail.


Water distillation of root bark from *C. glauca* collected in Petit Yapo forest produced an essential oil with a yield of 0.04%, w/w, vs fresh material, much lower than that obtained from leaves (0.16-0.19%) [7]. The root bark oil was first analyzed by GC, in combination with retention indices (RI) on two capillary columns of different polarity and by ¹³C NMR spectroscopy following a computerized method developed in our laboratory [8-10]. The experimental procedure may be summarized as follow: the ¹³C NMR chemical shift values of the experimental spectrum are matched against the spectra of reference compounds compiled into two computerized libraries, the first one built with spectra recorded in our laboratory, the second one built with ¹³C NMR data reported in the literature. Combined analysis by GC (RI) and ¹³C NMR of the essential oil (EO) allowed the identification of the major components and demonstrated that *C. glauca* root bark oil was a sesquiterpene-rich oil. However, various minor components remained unidentified and some discrepancies appeared between the GC (FID) and NMR analyses, concerning particularly the content of δ -elemene, a compound that may be produced by the plant or that may result from thermal rearrangement of germacrene C, a heat-sensitive compound. We demonstrated that such a rearrangement may occur during GC (FID) and GC-MS analyses [11]. Therefore, the EO was fractionated over a silica gel chromatography column (CC) into a hydrocarbon fraction (FH) and an oxygenated compounds-containing fraction (FO); both fractions were analyzed, once again, by GC (RI) and by ¹³C NMR (Table 1).

The major component was patchoulone (33.5%) (Figure 1), a tricyclic sesquiterpene ketone first isolated from the oil of *Cyperus rotundus* L. [12] and whose ¹³C NMR data have been reported later [13]. Sesquiterpene hydrocarbons were well represented by cyperene (9.5%) and germacrene D (6.6%). Camphene (5.2%) was the main monoterpene hydrocarbon and bornyl acetate (4.5%) the major oxygenated monoterpene. Other sesquiterpene hydrocarbons and oxygenated sesquiterpenes usually produced by aromatic plants, such as alismol (2.4%), δ -cadinene (1.7%), *epi*-cubenol (1.3%),

Table 1: Components of *Cleistopholis glauca* root oil.

Components	R _{1a}	R _{1p}	EO	FH	FO	Identification
1 α -Pinene	931	1024	0.9	0.8	-	R _{1a} , R _{1p} ¹³ C NMR
2 Camphene	944	1072	5.2	4.2	-	R _{1a} , R _{1p} ¹³ C NMR
3 Myrcene	980	1164	0.2	0.3	-	R _{1a} , R _{1p} ¹³ C NMR
4 Limonene*	1020	1204	0.3	0.5	-	R _{1a} , R _{1p} ¹³ C NMR
5 1,8-Cineole*	1020	1211	0.9	-	1.2	R _{1a} , R _{1p} ¹³ C NMR
6 (<i>E</i>)- β -Ocimene	1038	1252	0.2	0.3	-	R _{1a} , R _{1p} ¹³ C NMR
7 Bornyl acetate	1269	1580	4.5	-	7.7	R _{1a} , R _{1p} ¹³ C NMR
8 δ -Elemene*	1336	1472	2.3	5.4	-	R _{1a} , R _{1p} ¹³ C NMR
9 Germacrene C [#]	-	-	1.6	3.7	-	¹³ C NMR
10 α -Copaene	1376	1494	0.2	0.5	-	R _{1a} , R _{1p} ¹³ C NMR
11 β -Elemene	1388	1591	1.5	3.8	-	R _{1a} , R _{1p} ¹³ C NMR
12 Cyperene	1402	1530	9.5	23.9	-	R _{1a} , R _{1p} ¹³ C NMR
13 (<i>E</i>)- β -Caryophyllene	1412	1598	0.6	2.2	-	R _{1a} , R _{1p} ¹³ C NMR
14 β -Copaene	1428	1591	1.1	2.7	-	R _{1a} , R _{1p} ¹³ C NMR
15 <i>trans</i> - α -Bergamotene	1433	1586	0.2	0.5	-	R _{1a} , R _{1p} ¹³ C NMR
16 α -Humulene	1451	1669	0.2	0.5	-	R _{1a} , R _{1p} ¹³ C NMR
17 Rotundene	1459	1638	0.7	1.6	-	R _{1a} , R _{1p} ¹³ C NMR
18 1 β H,7 α H,10 β H-Guaia-4,11-diene*	1474	1663	1.8	3.5	-	R _{1a} , R _{1p} ¹³ C NMR
19 γ -Murolene*	1474	1688	0.5	1.2	-	R _{1a} , R _{1p} ¹³ C NMR
20 Germacrene D*	1477	1709	6.6	15.7	-	R _{1a} , R _{1p} ¹³ C NMR
21 α -Amorphene*	1477	1692	0.3	0.4	-	R _{1a} , R _{1p} ¹³ C NMR
22 β -Selinene	1482	1714	0.3	0.6	-	R _{1a} , R _{1p} ¹³ C NMR
23 Hinesene	1492	1725	0.9	2.3	-	R _{1a} , R _{1p} ¹³ C NMR
24 δ -Cadinene	1515	1757	1.7	4.3	-	R _{1a} , R _{1p} ¹³ C NMR
25 Germacrene B	1552	1827	1.0	2.4	-	R _{1a} , R _{1p} ¹³ C NMR
26 Patchouloune	1604	2098	33.5	-	58.8	R _{1a} , R _{1p} ¹³ C NMR
27 Alismol	1612	2270	2.4	-	4.2	R _{1a} , R _{1p} ¹³ C NMR
28 <i>epi</i> -Cubenol	1629	2061	1.3	-	2.3	R _{1a} , R _{1p} ¹³ C NMR
29 α -Cadinol	1639	2242	0.5	-	1.3	R _{1a} , R _{1p} ¹³ C NMR
30 Juvenile hormone III	1862	2495	1.8	-	3.2	R _{1a} , R _{1p} ¹³ C NMR
31 10-oxo-(2 <i>E</i> ,6 <i>E</i>)-Methylfarnesoate	1891	2579	0.9	-	1.5	R _{1a} , R _{1p} ¹³ C NMR

Order of elution and percentages are given on apolar column, those with an asterisk* excepted, percentage on polar column. R_{1a}, R_{1p} = Retention indices on apolar (BP-1) and polar (BP-20) column, respectively. FH and FO: fraction containing hydrocarbons and oxygenated compounds, respectively. #: Percentages calculated by combination of GC (FID) and ¹³C NMR. ¹³C NMR (*italic*): components identified by NMR in a CC fraction.

**Figure 1:** Structure of patchouloune (left) and cyperene (right).

β -copaene (1.1%), (*E*)- β -caryophyllene (0.6%) and α -cadinol (0.5%), were identified by comparison of their RIs on two columns of different polarity and their ¹³C NMR data with those of reference compounds compiled in our laboratory-built library.

Identification of some other components was achieved by comparison of the chemical shift values of the EO spectrum with literature data. For example, two compounds bearing the farnesane skeleton were identified as 10-oxo-(2*E*,6*E*)-methyl-farnesoate (**31**, 0.9%) and juvenile hormone III (**30**, 1.8%, 10,11-epoxy-3-methylenedodeca-1,6(*E*)-diene), an insect hormone rarely found in plants. The NMR data of the first one fit with those reported by Seidel *et al* [14], who isolated this compound from a stem bark extract of *C. glauca*. The carbon chemical shifts of juvenile hormone fit perfectly with those reported by Mori and Mori [15], who synthesized both enantiomers of that molecule. Recently, we reported the occurrence of this hormone in the leaf and trunk bark oils from *Cleistopholis patens* [11].

In parallel, it should be mentioned that 1 β H,7 α H,10 β H-guaia-4,11-diene **18** (1.8%) was identified by comparison of its ¹³C NMR chemical shifts (in the CC fraction FH and in the EO) with those reported by Blay *et al.* [16], who obtained the compound by synthesis. This compound has been simultaneously reported by our groups for the first time as a natural compound in the present work and in *Xylopia rubescens* leaf oil [17]. Similarly, hinesene (**23**) was identified by comparison of its NMR data and RI (apolar) with those reported by Joulain and König [18].

Special mention should be attributed to the identification and quantification of δ -elemene and germacrene C. The sesquiterpene hydrocarbon δ -elemene (**8**) was identified by GC (FID) and by ¹³C NMR spectroscopy. However, its relative percentage attributed by GC (FID) (3.9% in the EO, 9.1% in the CC fraction FH) was over-evaluated with respect to the intensities of the corresponding signals in the ¹³C NMR spectra. In agreement with our previous investigations on the composition of *C. patens* trunk bark oil [11], the ¹³C NMR spectra of the EO and the FH fraction of CC exhibited the 15 signals of δ -elemene [18] and the 15 signals of germacrene C (**9**). It should be mentioned that the ¹³C NMR data of germacrene C were reported in C₆D₆ [19]. Therefore, the spectrum of FH was recorded in C₆D₆ and in CDCl₃ and the chemical shift values of germacrene C in CDCl₃ are reported in the experimental part. As previously reported [20], germacrene C thermally rearranges to δ -elemene following a [3,3] sigmatropic reaction (Figure 2). Under our experimental chromatographic conditions, the content of δ -elemene evaluated by GC (3.9% in the EO, 9.1% in the CC fraction) corresponded in fact to the sum of δ -elemene and germacrene C. Combining GC percentages of δ -elemene with the ratio of both compounds, evaluated by comparing the mean intensities of their respective signals, allowed a better quantitative determination of δ -elemene and germacrene C: 5.4% and 3.7%, respectively in the CC fraction, 2.3% and 1.6%, respectively in the EO. In parallel we observed the presence of β -elemene (1.5%) and germacrene B (1.0%), while germacrene A and γ -elemene were not detected by NMR.

In conclusion, the root bark of *C. glauca* produced a sesquiterpene-rich oil, whose composition was dominated by patchouloune and cyperene. Neither compound has been identified as a component of *C. glauca* leaf oil that contained mainly germacrene D (16.4-46.5%) and (*E*)- β -caryophyllene (8.0-26.2%). Combined analysis by GC (RI) and by ¹³C NMR spectroscopy allowed the correct identification and quantification of germacrene C, a compound that thermally rearranges to δ -elemene during GC analysis.




Figure 2: Thermal rearrangement of germacrene C to δ-elemene.

Experimental

Plant material: Root bark of *C. glauca* was collected in Petit Yapo forest, near Adzopé (100 km north of Abidjan), in January 2011. Plant material was authenticated by Pr L. Aké Assi, from the Centre National Floristique (CNF, Abidjan, Ivory Coast).

Essential oil isolation and fractionation: The bark (1200 g) was submitted to hydrodistillation in a Clevenger-type apparatus for 3 h. The oil was decanted, weighed (0.530 g), and stored in a glass vial at 5°C prior to analysis. An aliquot of the EO (190 mg) was chromatographed on silica gel (ICN, 250-500 μm, 10 g). Two fractions were eluted with *n*-pentane (FH, 67 mg) and with diethyl oxide (FO, 97 mg). It should be pointed out that part of the monoterpene hydrocarbons was lost during the removal of solvent under reduced pressure.

GC analysis: GC analysis was carried out with a Perkin-Elmer Clarus 500 Autosystem apparatus equipped with 2 flame ionisation detectors (FID), and fused capillary columns (50 m x 0.22 mm i.d., film thickness 0.25 μm), BP-1 (polymethylsiloxane) and BP-20 (polyethylene glycol). Carrier gas, helium; linear velocity, 0.8 mL/min. The oven temperature was programmed from 60°C to 220°C at 2°C/min and then held isothermal (20 min). Injector temperature: 250°C (injection mode: split 1/60). Detector

temperature: 250°C. The relative proportions of the essential oil constituents were expressed as percentages obtained by peak area normalization, all relative response factors being taken as one.

¹³C NMR analysis: ¹³C NMR spectra were recorded on a Bruker AVANCE 400 Fourier Transform spectrometer operating at 100.63 MHz for ¹³C, equipped with a 5 mm probe, in deuterated chloroform (CDCl₃), with all shifts referred to internal tetramethylsilane (TMS). Spectra were recorded with the following parameters: pulse width (PW), 5 μs (flip angle 45°); acquisition time, 2.7 s for 128 K data table with a spectral width (SW) of 24000 Hz (240 ppm); digital resolution 0.183 Hz/pt. The number of accumulated scans was 3,000 for the oil sample and fractions of chromatography (about 50 mg of essential oil in 0.5 mL of CDCl₃).

Identification of individual components: The oil sample and both CC fractions were analyzed by GC (RI) and ¹³C NMR. Every component was identified by NMR, at least in one fraction of CC. Identification of most individual components was based: (i) on comparison of their GC retention indices (RI) on apolar and polar columns, determined relative to the retention times of a series of *n*-alkanes with linear interpolation ('Target Compounds' software of Perkin-Elmer), with those of either authentic compounds or literature data [18], (ii) by ¹³C NMR spectroscopy, following the methodology developed and computerized in our laboratories, using home-made software, (see text) [8-10]. A few compounds have been identified by comparison of their ¹³C NMR chemical shift values with those reported in the literature (see text).

¹³C NMR data of germacrene C (CDCl₃): δ 145.3 C, 141.4 C, 129.5 C, 127.1 C, 124.9 CH, 121.6 CH, 39.9 CH₂, 39.8 CH₂, 36.6 CH, 31.6 CH₂, 27.5 CH₂, 22.3 CH₃, 22.0 CH₃, 20.9 CH₃, 16.5 CH₃.

References

- [1] Le Thomas A. (1969) Flore du Gabon, volume 16. Annonacées, Muséum National d'Histoire Naturelle, Paris.
- [2] Ake Assi L, Abeye J, Guinko S, Giguët R, Bangavou Y. (1981) Contribution à l'identification et au recensement des plantes utilisées dans la médecine traditionnelle et la pharmacopée en République Centrafricaine. Agence de Coopération Culturelle et Technique, Paris.
- [3] Tane P, Ayafor JF, Sondengam BL. (1988) Partially-acetylated dodecanyl tri- and tetra-rhamnoside derivatives from *Cleistopholis glauca* (Annonaceae). *Tetrahedron Letters*, **29**, 1837-1840.
- [4] Seidel V, Bailleul F, Waterman PG. (2000) Novel oligorhamnosides from the stem bark of *Cleistopholis glauca*. *Journal of Natural Products*, **63**, 6-11.
- [5] Ngiefu K, Paquot C, Vieux A. (1976) Les plantes à huile du Zaïre II. *Oleagineux*, **31**, 545-547.
- [6] Etse JT, Gray AI, Waterman PG. (1988) Chemistry in the Annonaceae XXV: Sesquiterpenes from the stem bark of *Cleistopholis glauca*. *Journal of Natural Products*, **51**, 314-318.
- [7] Ouattara ZA, Boti JB, Ahibo CA, Tomi F, Casanova J, Bighelli A. (2012) Chemical composition of the leaf oil of *Cleistopholis glauca* Pierre ex Engler & Diels from Côte d'Ivoire. *Journal of Essential Oil Research*, **24**, 471-474.
- [8] Tomi F, Bradesi P, Bighelli A, Casanova J. (1995) Computer-aided identification on individual components of essential oils using carbon-13 NMR spectroscopy. *Journal of Magnetic Resonance Analysis*, **1**, 25-34.
- [9] Tomi F, Casanova J. (2006) ¹³C-NMR as a tool for identification of individual components of essential oils from Labiatae. A review. *Acta Horticulturae*, **723**, 185-192.
- [10] Bighelli A, Casanova J. (2009) Analytical tools for analysing *Cymbopogon* oils. In *Essential Oil Bearing Grasses - Cymbopogons*, Akhila A. (Ed.). CRC Press, London, 195-221.
- [11] Ouattara ZA, Boti JB, Ahibo AC, Casanova J, Tomi F, Bighelli A. (2013) Analysis of *Cleistopholis patens* leaf and trunk bark oils using combined GC-flame ionisation detection, GC-retention index, GC-MS and ¹³C-NMR. *Phytochemical Analysis*, in press. DOI: 10.1002/pca.2435
- [12] Trivedi B, Motl O, Herout V, Sorm F. (1964) On terpenes CLXIV. Composition of the oil from *Cyperus rotundus* L. Structure of patchoulone. *Collection of Czechoslovak Chemical Communications*, **29**, 1675-1688
- [13] Hisham A, Wray V, Pieters L, Claeys M, Dommissie R, Vlietinck A. (1992) Complete ¹H and ¹³C NMR spectral assignment of patchoulone, a tricyclic sesquiterpene ketone. *Magnetic Resonance in Chemistry*, **30**, 295-297.
- [14] Seidel V, Bailleul F, Waterman PG. (1999) Partially acetylated tri- and tetra-rhamnoside dodecanyl ether derivatives from *Cleistopholis patens*. *Phytochemistry*, **52**, 465-472.
- [15] Mori K, Mori H. (1987) Synthesis of both enantiomers of juvenile hormone III. *Tetrahedron*, **43**, 4097-4106
- [16] Blay G, Garcia B, Molina E, Pedro JR. (2007) Synthesis of (+)-pechueloic acid and (+)-aciphyllene. Revision of the structure of (+)-aciphyllene. *Tetrahedron*, **63**, 9621-9626.
- [17] Yapi TA, Boti JB, Ahibo AC, Bighelli A, Casanova J, Tomi F. (2013) Combined analysis of *Xylopi rubescens* Oliv. leaf oil by GC-FID, GC-MS and ¹³C NMR: Structure elucidation of new compounds. *Flavour Fragrance Journal*, **28**, 373-379.
- [18] Joulain D, König WA. (1998) *The Atlas of Spectral Data of Sesquiterpene Hydrocarbons*. E.B.-Verlag, Hamburg.

- [19] Colby SM, Crock J, Dowdle-Rizzo B, Lemaux PG, Croteau R. (1998) Germacrene C synthase from *Lycopersicon esculentum* cv. VFNT Cherry tomato: cDNA isolation, characterization, and bacterial expression of the multiple product sesquiterpene cyclase. *Biochemistry Proceedings of National Academy of Sciences USA*, **95**, 2216-2221.
- [20] Adio AM. (2009) Germacrenes A-E and related compounds: thermal, photochemical and acid induced transannular cyclizations. *Tetrahedron*, **65**, 1533-1552.

Bioactivities and Compositional Analyses of *Cinnamomum* Essential Oils from Nepal: *C. camphora*, *C. tamala*, and *C. glaucescens*

Prabodh Satyal^a, Prajwal Paudel^a, Ambika Poudel^b, Noura S. Dosoky^a, Kiran Kumar Pokharel^c and William N. Setzer^{a,*}

^aDepartment of Chemistry, University of Alabama in Huntsville, Huntsville, AL 35899, USA

^bDepartment of Chemistry, Tribhuvan University, MMAMC Campus, Biratnagar, Nepal

^cNational Herbarium and Plant Laboratories, Kathmandu, Nepal

wsetzer@chemistry.uah.edu

Received: September 9th, 2013; Accepted: September 16th, 2013

This work examines the biological activity of essential oils of *Cinnamomum camphora* leaves, *C. glaucescens* fruit, and *C. tamala* root from Nepal. The oils were screened for phytotoxic activity against lettuce and perennial ryegrass, brine shrimp lethality, and antibacterial, antifungal, cytotoxic, insecticidal, and nematocidal activities. *C. camphora* leaf essential oil was phytotoxic to lettuce, antifungal to *Aspergillus niger*, and insecticidal, particularly toward midge and butterfly larvae, fruit flies, and fire ants. *C. camphora* oil was also toxic to brine shrimp and human breast tumor cells. *C. glaucescens* fruit essential oil showed notable nematocidal activity, as well as termiticidal and mosquito larvicidal activity. The root essential oil of *C. tamala* was toxic to mosquito larvae and fire ants.

Keywords: Allelopathy, Insecticidal, Larvicidal, Nematocidal, Antimicrobial.

Cinnamomum, a genus of evergreen aromatic trees belonging to the family Lauraceae, is comprised of approximately 250 species [1]. In this study three species: *Cinnamomum camphora* (L.) J. Presl, *C. tamala* (Buch.-Ham.) T. Nees & Nees, and *C. glaucescens* Hand.-Mazz. from Nepal were studied for chemotypical variation in the volatile components, which were subsequently analyzed for possible biological activity.

C. camphora is primarily native to Japan and has been cultivated in the Himalayan region to produce camphor commercially, which is used in the pharmaceutical as well as the flavor industry [2,3]. Traditionally in the Ayurvedic system, *C. camphora* has been used to treat bronchitis, cold, congestion, diarrhea, dysentery, edema, flu, gas, metabolic and heart problems, as well as various gynecological problems [2b]. In addition, the Yunani, “Greek”, medicinal system also uses *C. camphora* for a cephalic tonic, cardiac treatment, and also as an expectorant [2b]. *C. camphora* extracts have exhibited cytotoxic [3], anti-inflammatory and anti-oxidant activities [4], and termiticidal [5], acaricidal [6], antifungal [7], and anthelmintic [8] activities.

C. tamala, commonly known as “tezpat”, is widely distributed throughout tropical and subtropical regions of Australia, the Pacific region, South America and the Himalayan region of Asia [9], and typically grows at altitudes between 900-2500 m [10]. Morphologically, *C. tamala* is a perennial or small evergreen tree, attaining 8-12 m in height and a girth of 150 cm with the leaves ranging from 12-20 cm long, 5-8 cm broad, and with white flowers [11]. Pharmacologically the plant has been reported to exhibit antihyperglycemic, antidiarrheal, antihyperlipidemic, anti-inflammatory, acaricidal, hepatoprotective, gastroprotective, antioxidant, antibacterial, α -amylase-inhibitory, and immunomodulatory activities [12]. The bark of *C. tamala* is used in Nepalese traditional medicine to treat intestinal disorders and control nausea, while the leaf is used as a spice and to control diarrhea [13]. Leaves of *C. tamala* are used in India to repel stored grain pests [14].

C. glaucescens, commonly known as “sugandh kokila”, is native to the Himalayan range in Nepal and India [15], although it is also found in the plains and hilly regions at altitudes below 2500 m. Morphologically, *C. glaucescens* buds are enclosed in overlapping scales with fruits about 3 cm long, the base enclosed in a cup 10-12 mm across, while the leaves are alternate, between 7-10 cm long with elliptical shape, a well-marked midrib, and 4-5 pairs of lateral veins. *C. glaucescens* is extensively used by local people as a spice, in medicine, and in consumption of tobacco [16]. Therapeutically, *C. glaucescens* has been traditionally used as a demulgent and stimulant and has shown analgesic, antiseptic, astringent, and carminative properties [16]. Furthermore, *C. glaucescens* seed is used for treatment of the common cold, cough, toothache and taenias [16]. Seed paste is applied to treat muscular swellings [16]. Seed oil has also been demonstrated to treat muscular spasm, joint pain and body aches [16]. *C. glaucescens* fruit essential oil has been assessed for insecticidal, oviposition deterrent and antifungal activities [17].

Essential oil compositions: The essential oils were obtained in 1.3%, 0.9%, 1.0%, and 1.5% yields for *C. camphora* (#15) leaf, *C. camphora* (#72) leaf, *C. tamala* root, and *C. glaucescens* fruits, respectively. The chemical compositions of the oils are summarized in Table 1. *C. camphora* (#15) leaf oil was dominated by camphor (36.5%), camphene (11.7%), and limonene (9.0%), with lesser amounts of sabinene (6.3%) and β -pinene (6.3%), while the *C. camphora* (#72) leaf oil contained almost exclusively camphor (98.0%). Currently, there are five different chemotypes observed worldwide for *C. camphora*: camphor, linalool, 1,8-cineole, nerolidol, and borneol [18]. Specifically, camphor content varied from a Japanese sample with 51.5% to a Chinese oil sample with 83.9% and an Ivory Coast leaf oil containing 74% camphor [19].

C. tamala root essential oil was also found to be rich in camphor (35.0%), followed by linalool (10.6%), *p*-cymene (8.5%), *o*-cymene (6.8%), and 1,8-cineole (6.1%). A survey of the literature reveals

Table 1: Chemical compositions of *Cinnamomum camphora* leaf essential oil, *C. glaucescens* fruit essential oil, and *C. tamala* root essential oil from Nepal.

RI ^a	Compound	Percent Composition ^b			
		<i>C. camphora</i>		<i>C. glaucescens</i>	<i>C. tamala</i>
		#15	#72		
922	Tricyclene	0.7	—	—	0.1
933	Methyl hexanoate	—	—	tr ^c	—
934	α -Thujene	7.7	—	0.4	0.7
941	α -Pinene	4.7	tr	2.6	4.7
953	Camphene	11.7	0.2	0.1	2.0
957	Thuja-2,4(10)-diene	—	—	tr	tr
962	Benzaldehyde	—	—	0.2	0.1
976	Sabinene	6.3	—	5.7	0.5
980	β -Pinene	6.3	0.1	2.8	4.1
993	Myrcene	1.2	—	0.1	1.1
1000	δ -2-Carene	tr	—	—	tr
1005	α -Phellandrene	—	—	tr	—
1007	δ -3-Carene	tr	—	—	0.3
1016	α -Terpinene	—	—	0.3	—
1025	<i>p</i> -Cymene	1.8	—	0.3	8.5
1028	<i>o</i> -Cymene	1.8	—	0.2	6.8
1034	Limonene	9.0	—	—	—
1035	1,8-Cineole	2.7	0.2	24.8	6.1
1043	Salicylaldehyde	—	—	—	0.3
1060	γ -Terpinene	tr	—	1.0	—
1063	<i>cis</i> -Sabinene hydrate	tr	—	tr	tr
1068	<i>cis</i> -Linalool oxide (furanoid)	tr	—	0.2	0.3
1076	<i>trans</i> -Linalool oxide (furanoid)	—	—	—	0.3
1076	<i>trans</i> -Dihydrosesquioxide	tr	—	—	—
1079	<i>p</i> -Cresol	—	—	tr	—
1087	Terpinolene	—	—	0.3	—
1087	Fenchone	0.1	—	—	—
1093	6,7-Epoxyterpinene	tr	—	—	—
1097	α -Pinene oxide	tr	—	—	—
1102	Perillene	0.2	—	—	—
1102	Linalool	0.6	—	3.7	10.6
1113	<i>endo</i> -Fenchol	—	—	tr	—
1121	<i>cis</i> - <i>p</i> -Menth-2-en-1-ol	—	—	0.1	—
1126	Methyl octanoate	—	—	tr	—
1138	<i>trans</i> -Pinocarveol	—	—	0.1	—
1139	<i>trans</i> -Sabinol	—	—	0.1	—
1144	<i>trans</i> -Verbenol	—	—	tr	—
1146	Camphor	36.5	98.0	—	35.0
1157	Sabina ketone	—	—	tr	—
1162	Pinocarvone	—	—	0.1	—
1162	Isoborneol	—	—	0.1	tr
1165	Borneol	0.2	1.4	—	0.8
1166	Pinocarvone	0.2	—	—	0.2
1168	δ -Terpineol	—	—	0.9	—
1170	<i>cis</i> -Linalool oxide (pyranoid)	—	—	tr	tr
1179	Terpinen-4-ol	1.4	tr	4.8	1.8
1182	<i>p</i> -Methylacetophenone	0.1	—	tr	tr
1184	Thuj-3-en-10-al	—	—	tr	—
1186	Cryptone	—	—	0.1	0.3
1188	<i>p</i> -Cymen-8-ol	0.2	—	—	—
1192	α -Terpineol	1.5	0.2	7.4	2.2
1196	Myrtenal	0.3	—	—	0.2
1197	Estragole (= Methyl chavicol)	—	—	—	0.1
1198	Myrtenol	0.3	—	0.3	0.1
1203	<i>cis</i> -Piperitol	—	—	tr	tr
1207	<i>trans</i> -Piperitol	tr	—	—	tr
1209	Verbenone	0.1	—	0.1	0.1
1217	<i>trans</i> -Carveol	0.2	—	tr	0.1
1226	<i>cis</i> -Carveol	0.1	—	—	—
1227	Nerol	tr	tr	0.2	0.1
1235	Ascaridole	—	—	tr	—
1236	Neral	tr	—	tr	—
1237	Cuminal	0.1	—	0.1	0.1
1239	<i>o</i> -Anisaldehyde	—	—	—	0.1
1242	Carvone	—	—	—	0.1
1242	Carvone	0.2	—	tr	—
1251	Piperitone	0.1	—	tr	0.1
1255	Geraniol	—	—	0.1	tr
1260	(2 <i>E</i>)-Decenal	—	—	—	tr
1269	(<i>E</i>)-Cinnamaldehyde	—	—	—	0.4
1270	Geranial	—	—	tr	tr
1281	Neryl formate	—	—	tr	—
1283	Safrole	tr	—	—	—
1285	<i>neois</i> -3-Thujanol acetate	—	—	tr	—
1286	Bornyl acetate	—	—	—	0.8
1290	<i>p</i> -Cymen-7-ol	0.1	—	0.1	0.1
1299	Terpinen-4-ol acetate	—	—	—	0.1
1301	Carvacrol	—	—	—	tr
1302	Methyl (<i>Z</i>)-cinnamate	—	—	0.5	—
1305	(<i>E</i>)-Cinnamyl alcohol	—	—	—	0.2
1321	Methyl geranate	—	—	tr	—
1332	8-Hydroxy-neo-menthol	—	—	tr	—
1337	δ -Elemene	—	—	—	tr

RI ^a	Compound	Percent Composition ^b			
		<i>C. camphora</i>		<i>C. glaucescens</i>	<i>C. tamala</i>
		#15	#72		
1349	α -Cubebene	tr	—	tr	0.1
1366	Neryl acetate	—	—	—	tr
1371	Hydrocinnamyl acetate	—	—	—	0.1
1376	α -Copaene	0.2	—	0.3	1.3
1381	<i>trans</i> -Soberol	0.3	—	—	—
1382	<i>trans</i> - <i>p</i> -Menth-6-ene-2,8-diol	—	—	—	tr
1387	Geranyl acetate	—	—	—	0.1
1388	Methyl (<i>E</i>)-cinnamate	—	—	40.5	tr
1391	β -Cubebene	0.2	—	—	0.8
1394	β -Elemene	0.1	—	—	0.2
1419	(<i>E</i>)-Caryophyllene	0.1	tr	tr	1.0
1428	Carvone hydrate	0.1	—	0.3	tr
1430	β -Copaene	—	—	—	tr
1438	Coumarin	—	—	tr	0.6
1439	Aromadendrene	0.2	—	—	tr
1451	(<i>E</i>)-Cinnamyl acetate	—	—	—	1.2
1454	α -Humulene	0.1	tr	tr	1.2
1461	Alloaromadendrene	0.1	—	—	—
1462	9- <i>epi</i> -(<i>E</i>)-Caryophyllene	—	—	—	tr
1478	γ -Murolene	tr	—	—	0.1
1483	Amorpha-4,7(11)-diene	—	—	—	0.1
1485	Widdra-2,4(14)-diene	—	—	—	tr
1487	β -Selinene	tr	—	—	tr
1493	<i>trans</i> -Murola-4(14),5-diene	—	—	—	tr
1496	<i>epi</i> -Cubebol	tr	—	0.1	0.2
1497	Bicyclogermacrene	—	tr	—	—
1502	α -Murolene	tr	—	—	0.1
1507	Germacrene A	—	—	—	tr
1515	γ -Cadinene	tr	—	—	tr
1518	Cubebol	0.1	—	0.1	0.1
1524	<i>trans</i> -Calamenene	tr	—	—	—
1525	δ -Cadinene	—	—	0.1	0.3
1529	(<i>E</i>)- <i>o</i> -Methoxycinnamaldehyde	—	—	—	tr
1538	α -Cadinene	—	—	—	tr
1543	α -Calacorene	—	—	—	tr
1550	Elemol	—	tr	—	—
1554	(<i>Z</i>)-Caryophyllene oxide	—	—	—	0.1
1558	Germacrene B	—	—	—	tr
1565	β -Calacorene	—	—	—	tr
1579	Spathulenol	—	tr	tr	0.2
1585	Caryophyllene oxide	0.4	tr	0.3	1.3
1592	Viridiflorol	tr	—	tr	—
1595	Cubeban-11-ol	tr	—	—	—
1604	Ledol	tr	—	tr	—
1609	Humulene epoxide II	0.3	—	0.1	0.4
1623	α -Corocalene	—	—	—	tr
1629	1- <i>epi</i> -Cubanol	tr	—	tr	0.1
1634	Caryophylla-4(12),8(13)-dien-5-ol	—	—	—	0.1
1643	τ -Murolol	tr	—	tr	0.1
1646	α -Murolol (= Torreyol)	—	—	—	tr
1650	β -Eudesmol	—	—	tr	—
1655	α -Cadinol	—	—	tr	—
1656	Selin-11-en-4a-ol	—	tr	—	0.1
1659	<i>cis</i> -Calamenen-10-ol	—	—	tr	0.1
1667	<i>trans</i> -Calamenen-10-ol	—	—	tr	tr
1670	14-Hydroxy-9- <i>epi</i> -(<i>E</i>)-caryophyllene	—	—	tr	0.1
1673	Cadalene	—	—	tr	tr
1678	Mustakone	—	—	tr	—
1722	(2 <i>Z</i> ,6 <i>E</i>)-Farnesol	—	—	tr	—
1741	(2 <i>E</i> ,6 <i>E</i>)-Farnesol	—	—	tr	—
2037	Kaur-16-ene	—	—	—	0.2
Total Identified		98.5	100	99.4	99.4

^a RI = "Retention Index" determined on an HP-5ms column with respect to a homologous series of *n*-alkanes.

^b Raw percentages based on total ion current without standardization.

^c tr = "trace" (< 0.05%).

that this is the first study of *C. tamala* root essential oil. However, previous leaf essential oil studies have shown eugenol and cinnamaldehyde to be major components, with Indian samples reporting 81.7% to 78% eugenol [20,21]. However, other studies showed linalool (50.3-33.7%, 19.7%, and 15.7-15.3%) and cinnamaldehyde (12.7-8.7%, 52.8%, and 41.2-55.2%) as the main components of the leaf oil of *C. tamala*, while eugenol was found in trace amounts [22-24].

Hydrodistilled *C. glaucescens* fruit (berries) essential oil was found to contain primarily methyl (*E*)-cinnamate (40.5%) and 1,8-cineole

(24.8%). Smaller quantities of α -terpineol (7.4%), sabinene (5.7%), terpinen-4-ol (4.8%), and linalool (3.7%) were also present. In another study from Nepal, the fruit oil of *C. glaucescens* showed a qualitatively similar composition with major components being 1,8 cineole (13%), methyl (*E*)-cinnamate (14%), and α -terpineol (7%) [25]. In contrast, a *C. glaucescens* fruit essential oil from India showed no methyl cinnamate, but a large percentage of 1,8-cineole (43.6%) [17]. In comparison, leaf oil from India of *C. glaucescens* contained elemicin (92.9%) and methyl eugenol (4.9%) as major components [15].

Allelopathic activity: The allelopathic activities of Nepalese *Cinnamomum* essential oils were analyzed in terms of inhibition of seed germination, as well as inhibition of seedling growth against a representative dicot (lettuce, *Lactuca sativa*) and a representative monocot (perennial ryegrass, *Lolium perenne*). The allelopathic activities of *C. camphora*, *C. tamala*, and *C. glaucescens* essential oils are summarized in Table 2. *Lactuca sativa* seed germination was notably inhibited by *C. camphora* (#15) essential oil (IC_{50} = 149 μ g/mL), as well as its major component, camphor (IC_{50} = 239 μ g/mL). Although neither *C. tamala* nor *C. glaucescens* appreciably affected germination of either *L. sativa* or *L. perenne*, the major component of *C. glaucescens* oil, methyl (*E*)-cinnamate did inhibit *L. perenne* germination (IC_{50} = 610 μ g/mL).

L. sativa seedling growth was more sensitive to the *Cinnamomum* essential oils than *L. perenne*. *C. camphora* essential oil significantly inhibited *L. sativa* radicle growth at 250 μ g/mL, which can be attributed to the major component, camphor. *C. glaucescens* essential oil and methyl (*E*)-cinnamate were inhibitory to both *L. sativa* and *L. perenne* at the concentrations tested. *C. tamala* oil inhibited seedling growth of *L. sativa* at 500 μ g/mL and *L. perenne* at 1000 μ g/mL, likely due to the high concentration of camphor in this essential oil.

Antimicrobial activity: Essential oils from the three species were analyzed for antibacterial and antifungal activities, while the major components were also tested to determine the compounds responsible for the results (Table 3). Of the three essential oils tested, only *C. camphora* showed significant antifungal activity against *A. niger* (MIC = 20 μ g/mL). The antifungal activity is hypothesized to be due to synergism among the essential oil

components rather than to any one of the constituents. Camphor (36.5%), limonene (9.0%), and β -pinene (6.3%) all exhibited MICs of 156 μ g/mL against *A. niger* (Table 3). *C. camphora* oil also showed marginal activity against *B. cereus* and *S. aureus*, with a MIC = 313 μ g/mL. *C. camphora* essential oils have been previously noted to have antifungal [26-28] and antibacterial activities [18,29], in agreement with the antimicrobial activities of Nepalese *C. camphora*. *C. tamala* essential oil was also somewhat active against *A. niger* (MIC = 156 μ g/mL).

Cytotoxicity and brine shrimp lethality: All three essential oils exhibited *in-vitro* cytotoxic activity against MCF-7 cells, and these activities also correlate with brine shrimp lethality (Table 3). Anderson and co-workers had previously reported a correlation between brine shrimp lethality and cytotoxic activity [30]. Most of the essential oil components tested also showed cytotoxicity to MCF-7 cells, as well as brine shrimp lethality.

Insecticidal activity: The *Cinnamomum* essential oils were screened for insecticidal activity against mosquito (*Culex pipiens*) and midge (*Chaoborus plumicornis*) larvae, cabbage white butterfly (*Pieris rapae*) larvae, termites (*Reticulitermes virginicus*), fruit flies (*Drosophila melanogaster*), and red imported fire ants (*Solenopsis invicta* \times *richteri*) (Table 3). All three essential oils showed mosquito larvicidal activities comparable with or better than previously reported essential oils, for example, common myrtle (*Myrtus communis*) [31], bay laurel (*Laurus nobilis*) [32], amyris (*Amyris balsamifera*) [33], and spearmint (*Mentha spicata*) [34], against *Culex pipiens*. Consistent with the mosquito larvicidal activity, the essential oils and components showed larvicidal activity against *Chaoborus* larvae as well. *C. camphora* leaf oil had previously shown such activity against *Culex quinquefasciatus* [35] and *Aedes aegypti* [36].

C. camphora leaf oil was the most active essential oil against cabbage butterfly larvae, fruit flies, and fire ants (LC_{50} = 186, 153, and 176 μ g/mL, respectively). *C. camphora* leaf oil had previously shown insecticidal activity against fire ants (*Solenopsis invicta*) [37], the rice weevil (*Sitophilus oryzae*), the broadbean weevil (*Bruchus rufimanus*) [38], *C. camphora* oil also showed insect repellent activity against these stored grain beetles [38] as well as adult *Aedes aegypti*, *Anopheles stephensi*, and *Culex*

Table 2: Allelopathic activity of *Cinnamomum camphora* leaf, *C. glaucescens* fruit, and *C. tamala* root essential oils and essential oil components on lettuce (*Lactuca sativa*) and perennial ryegrass (*Lolium perenne*).

Material (Concentration, μ g/mL)	Germination Inhibition (%)		Seedling Growth (% of Controls)			
	<i>Lactuca sativa</i>	<i>Lolium perenne</i>	<i>Lactuca sativa</i> radicle	<i>Lactuca sativa</i> hypocotyl	<i>Lolium perenne</i> radicle	<i>Lolium perenne</i> hypocotyl
<i>C. camphora</i> oil (1000)	98.3 ^a	31.7 ^b	69.4 ^m	37.9 ^m	>100 ^a	>100 ^a
<i>C. camphora</i> oil (500)	71.7	25.0	81.6 ⁱ	82.9 ^m	>100 ^a	>100 ^a
<i>C. camphora</i> oil (250)	66.7	18.3	80.2 ^p	>100 ^s	>100 ^a	>100 ^s
<i>C. glaucescens</i> oil (2000)	50.0 ^c	45.0 ^d	46.4 ^m	45.5 ^m	52.2 ^m	4.8 ^m
<i>C. glaucescens</i> oil (1000)	35.0	30.0	54.7 ^m	51.6 ^m	60.4 ^m	28.0 ^m
<i>C. glaucescens</i> oil (500)	30.0	18.3	82.0 ⁿ	66.2 ^m	83.0 ^p	72.6 ^p
<i>C. glaucescens</i> oil (250)	15.0	15.0	87.0 ^p	76.8 ^m	90.9 ^f	73.3 ^q
<i>C. tamala</i> oil (2000)	98.3 ^c	63.3 ^f	0.0 ^m	0.0 ^m	60.6 ^m	21.0 ^m
<i>C. tamala</i> oil (1000)	41.7	16.7	86.6 ^p	64.7 ^m	70.1 ^m	88.1 ⁱ
<i>C. tamala</i> oil (500)	23.3	10	96.0 ^f	77.8 ^m	93.0 ^f	99.1 ^s
<i>C. tamala</i> oil (250)	18.3	6.7	97.7 ^s	79.4 ^m	97.8 ^s	99.7 ^s
Methyl (<i>E</i>)-cinnamate (2000)	48.3 ^s	100 ^h	46.9 ^m	46.9 ^m	—	—
Methyl (<i>E</i>)-cinnamate (1000)	20.0	93.3	63.3 ^m	52.0 ^m	16.9 ^m	2.4 ^m
Methyl (<i>E</i>)-cinnamate (500)	1.7	30.0	89.8 ^p	75.8 ^m	42.2 ^m	10.6 ^m
Methyl (<i>E</i>)-cinnamate (250)	—	20.0	—	—	45.4 ^m	28.4 ^m
Camphor (1000)	91.7 ^j	50.0 ^j	19.9 ^m	4.8 ^m	15.8 ^m	3.5 ^m
Camphor (500)	91.7	43.3	21.1 ^m	9.7 ^m	17.3 ^m	5.3 ^m
Camphor (250)	58.3	35.0	51.6 ^m	19.3 ^m	41.5 ^m	7.8 ^m
α -Pinene (4000)	6.67	13.3	>100 ^s	99.6 ^s	90.0 ^s	>100 ^s
β -Pinene (4000)	13.3	18.3	84.1 ^q	58.8 ^m	77.0 ^q	11.5 ^m
β -Pinene (2000)	1.7	25.0	98.6 ^s	85.1 ^s	82.5 ^s	39.4 ^m

^a IC_{50} = 218 μ g/mL; ^b IC_{50} > 1000 μ g/mL; ^c IC_{50} = 1620 μ g/mL; ^d IC_{50} > 1773 μ g/mL; ^e IC_{50} = 1084 μ g/mL; ^f IC_{50} = 1630 μ g/mL; ^g IC_{50} = 1828 μ g/mL; ^h IC_{50} = 610 μ g/mL; ⁱ IC_{50} = 239 μ g/mL; ^j IC_{50} = 1000 μ g/mL.

Significance vs. controls: ^m P < 0.001, ⁿ P \approx 0.005 ^o P \approx 0.01, ^p P \approx 0.05, ^q P \approx 0.2, ^r P \approx 0.5, ^s not significant.

Table 3: Bioactivity screening of *Cinnamomum camphora* leaf, *C. glaucescens* fruit, and *C. tamala* root essential oils and major essential oil components.

Bioassay	<i>C. camphora</i> leaf oil	<i>C. glaucescens</i> fruit oil	<i>C. tamala</i> root oil	Camphor	1,8- Cineole	<i>p</i> - Cymene	Limonene	Linalool	Methyl cinnamate	α - Pinene	β - Pinene	α - Terpineol
<i>Artemia salina</i> lethality (LC ₅₀ , μ g/mL)	2.5 \pm 2.5	20.1 \pm 2.6	28.1 \pm 3.0	21.7 \pm 9.9	26.9 \pm 13.1	4.7 \pm 1.0	6.3 \pm 3.4	36.0 \pm 3.9	20.7 \pm 3.1	21.4 \pm 5.7	15.7 \pm 6.1	> 100
MCF-7 cytotoxicity (% kill at 100 μ g/mL)	71.2 \pm 26.8	55.1 \pm 4.8	41.2 \pm 13.3	100	100	100	90.78 \pm 9.6	18.0 \pm 9.8	0	68.3 \pm 4.9	78.0 \pm 4.2	100
Antimicrobial (MIC, μ g/mL)												
<i>Bacillus cereus</i>	313	625	625	313	313	1250	625	625	1250	1250	1250	625
<i>Staphylococcus aureus</i>	313	313	625	1250	625	1250	313	156	1250	1250	1250	1250
<i>Escherichia coli</i>	1250	625	625	1250	1250	625	625	625	625	1250	1250	1250
<i>Pseudomonas aeruginosa</i>	625	313	625	1250	1250	625	1250	1250	625	1250	1250	1250
<i>Aspergillus niger</i>	19.5	313	156	625	625	313	1250	625	625	156	156	313
<i>C. elegans</i> nematocidal (LC ₅₀ , μ g/mL)	574 \pm 118	151 \pm 51	1320 \pm 130	> 2500	> 2500	289 \pm 118	85.1	> 2500	138 \pm 28	> 2500	> 2500	> 2500
Insecticidal (LC ₅₀ , μ g/mL)												
<i>Chaoborus plumicornis</i> (larvae)	61.5 \pm 25.0	186 \pm 54	116 \pm 53	188 \pm 22	342 \pm 177	79.6	92.8	278	nt	97.4	97.4	140
<i>Culex pipiens</i> (larvae)	46.4	21.5	17.2	7.2	150	nt	nt	nt	nt	nt	nt	nt
<i>Pieris rapae</i> (larvae)	186	820	653	nt	nt	nt	nt	nt	nt	nt	nt	nt
<i>Reticulitermes virginicus</i>	340	34	1075	66	1080	nt	158	nt	nt	nt	nt	nt
<i>Drosophila melanogaster</i>	153 \pm 50	348 \pm 83	245 \pm 25	117 \pm 58	160 \pm 81	18.4 \pm 4.0	249 \pm 62	28.0 \pm 7.3	167 \pm 67	220 \pm 54	260 \pm 123	383 \pm 184
<i>Solenopsis invicta</i> \times <i>richieri</i>	176 \pm 53	325 \pm 90	191 \pm 59	286 \pm 23	375 \pm 74	518 \pm 56	1285 \pm 80	613 \pm 65	333 \pm 39	> 1000	> 1000	282 \pm 33

^a nt = not tested

quinquefasciatus mosquitoes [39], and the red bud borer, *Resseliella oculiperda* [40]. Camphor was shown to be the insecticidal component in this present work and in a previous study on stored grain insect pests [41].

C. glaucescens fruit oil was the most active *Cinnamomum* oil against termites with an LC₅₀ = 34 μ g/mL. Several essential oils have shown termiticidal activity [42-47], but differences in assay protocols preclude comparisons of bioactivities.

Nematicidal activity: *C. glaucescens* fruit essential oil exhibited the strongest nematocidal activity with an LC₅₀ of 151 μ g/mL, followed by *C. camphora* leaf oil (LC₅₀ = 574 μ g/mL) and *C. tamala* root oil (LC₅₀ = 1320 μ g/mL). The nematocidal activity of *C. glaucescens* oil on *C. elegans* can be attributed to its major component, methyl (*E*)-cinnamate (LC₅₀ = 138 μ g/mL), which had previously shown nematocidal activity against the pine wood nematode, *Bursaphelenchus xylophilus* [48,49]. The other major component in *C. glaucescens* oil, 1,8-cineole, was inactive against *C. elegans*, as well as the root knot nematode, *Meloidogyne incognita* [50]. Notably, camphor, the major component in both *C. camphora* and *C. tamala* oils, was also inactive against *C. elegans*. Camphor was reported by Ntalli and co-workers [51] to be inactive against the root knot nematode, *M. incognita*, in contrast to an earlier report by Al-Banna *et al.* [52].

Plant-derived natural products have been used for thousands of years, both for their medicinal properties [53], as well as pesticidal utilities [54-56]. Essential oils are attractive as pesticidal agents because they are readily available, renewable, readily degraded, and non-polluting. Essential oils have been utilized as fungicides [57], herbicides [58], insecticides [59], and nematocides [60,61]. In this work, we have demonstrated the pesticidal potential of essential oils derived from *Cinnamomum camphora*, *C. glaucescens*, and *C. tamala*. In particular, *C. camphora* leaf essential oil was phytotoxic to lettuce, antifungal to *Aspergillus niger*, and insecticidal, particularly toward midge and butterfly larvae, fruit flies, and fire ants. *C. camphora* oil was also toxic to brine shrimp and human breast tumor cells. *C. glaucescens* fruit essential oil showed notable nematocidal activity, as well as termiticidal and mosquito larvicidal activity. The root essential oil of *C. tamala* was toxic to mosquito larvae and fire ants.

Experimental

Plant material: *Cinnamomum camphora* (#15), *C. tamala*, and *C. glaucescens* were collected on the 18th May, 2011, from Hetauda,

Makwanpur (27.42° N, 85.03° E, 1550 m above sea level), Nepal. An additional sample of *C. camphora* (#72) was collected from Dhudikhel, Kavre (27.61°N 85.59°E, 1512 m above sea level) on 12th July, 2011. The plants were identified by Kiran Kumar Pokharel and voucher specimens (M001, M002, and M003, respectively) have been deposited in the Botany Department, Tribhuvan University, Post-Graduate Campus. Fresh leaves (100.0 g) of *C. camphora*, roots (100.0 g) of *C. tamala*, and berries (200.0 g) of *C. glaucescens*, were shredded and hydrodistilled for 4 h using a Clevenger apparatus to give essential oils: 1.3 g, 0.9 g, 1.0 g, and 1.5 g for *C. camphora* (#15), *C. camphora* (#72), *C. tamala*, and *C. glaucescens*, respectively.

Gas chromatographic-mass spectral analysis: The *Cinnamomum* essential oils were analyzed by GC-MS using an Agilent 6890 GC with an Agilent 5973 mass selective detector [MSD, operated in the EI mode (electron energy = 70 eV), scan range = 45-400 amu, and scan rate = 3.99 scans/sec), and an Agilent ChemStation data system. The GC column was an HP-5ms fused silica capillary with a (5% phenyl)-polymethylsiloxane stationary phase, film thickness 0.25 μ m, length 30 m, and an internal diameter of 0.25 mm. The carrier gas was helium with a column head pressure of 48.7 kPa and a flow rate of 1.0 mL/min. Injector temperature was 200°C and detector temperature 280°C. The GC oven temperature program was used as follows: 40°C initial temperature, hold for 10 min; increased at 3°C/min to 200°C; increased 2°/min to 220°C. A 1%, w/v, solution of the sample in CH₂Cl₂ was prepared and 1 μ L was injected using a splitless injection technique.

Identification of the oil components was based on their retention indices determined by reference to a homologous series of *n*-alkanes, and by comparison of their mass spectral fragmentation patterns with those reported in the literature [62] and stored on the MS library [NIST database (G1036A, revision D.01.00)/ChemStation data system (G1701CA, version C.00.01.080)]. The percentages of each component are reported as raw percentages based on total ion current without standardization.

Allelopathic activity assays: An allelopathic bioassay based on lettuce (*Lactuca sativa* L.) and perennial rye grass (*Lolium perenne* L.) germination and subsequent radicle and hypocotyl growth [63] was measured to study the effects of the *Cinnamomum* essential oils and major essential oil components. Stock solutions of each essential oil or compound (2.0 g/L essential oil and 1.0 g/L Tween-80 in water) were prepared and used for the assays. Two-fold serial dilutions of stock test solutions were prepared to give test

concentrations of 4000, 2000, 1000, 500, and 250 µg/mL with the control being 1.0 g/L aqueous Tween-80. Seeds were placed in 6-well test plates (10 seeds per well), each well lined with 2 layers of Whatman No. 1 filter paper moistened with test solution, and the test plates were sealed with Parafilm®. The test plates were incubated at room temperature in the dark for 5 days, after which the number of germinated seeds was determined and the root (radicle) and shoot (hypocotyl) lengths were measured. Student's *t*-test [64] was used to compare radicle and hypocotyl test means with controls. Seed germination IC₅₀ values were determined using the Reed-Muench method [65].

Antimicrobial screening: The essential oils and major components were screened for antimicrobial activity against Gram-positive bacteria, *Bacillus cereus* (ATCC No. 14579) and *Staphylococcus aureus* (ATCC No. 29213); Gram-negative bacteria, *Pseudomonas aeruginosa* (ATCC No. 27853) and *Escherichia coli* (ATCC No. 10798). Minimum inhibitory concentrations (MIC) were determined using the microbroth dilution technique [66]. Dilutions of the essential oil were prepared in cation-adjusted Mueller Hinton broth (CAMHB) beginning with 50 µL of 1%, w/w, solutions of essential oil in DMSO plus 50 µL CAMHB. The extract solutions were serially diluted (1:1) in CAMHB in 96-well plates. Organisms at a concentration of approximately 1.5×10^8 colony forming units (CFU)/mL were added to each well. Plates were incubated at 37°C for 24 h; the final minimum inhibitory concentration (MIC) was determined as the lowest concentration without turbidity. Geneticin was used as a positive antibiotic control; DMSO was used as a negative control. Antifungal activity against *Aspergillus niger* (ATCC No. 16888) was determined as above using potato dextrose broth inoculated with *A. niger* hyphal culture diluted to a McFarland turbidity of 1.0. Amphotericin B was the positive control.

Cytotoxicity screening: Human MCF-7 breast adenocarcinoma cells (ATCC No. HTB-22) [67] were grown in a 3% CO₂ environment at 37°C in RPMI-1640 medium, supplemented with 10% fetal bovine serum, 100,000 units penicillin and 10.0 mg streptomycin per L of medium, 15 mM of Hepes, and buffered with 26.7 mM NaHCO₃, pH 7.35. Cells were plated into 96-well cell culture plates at 2.5×10^4 cells per well. The volume in each well was 100 µL. After 48 h, supernatant fluid was removed by suction and replaced with 100 µL growth medium containing 1.0 µL of DMSO solution of the essential oil (1%, w/w, in DMSO), giving a final concentration of 100 µg/mL for each well. Solutions were added to wells in 4 replicates. Medium controls and DMSO controls (10 µL DMSO/mL) were used. Tingenone [68] was used as a positive control. After the addition of compounds, plates were incubated for 48 h at 37°C in 5% CO₂; medium was then removed by suction, and 100 µL of fresh medium was added to each well. In order to establish percent kill rates, the MTT assay for cell viability was carried out [69]. After colorimetric readings were recorded (using a Molecular Devices SpectraMAX Plus microplate reader, 570 nm), average absorbances, standard deviations, and percent kill ratios (%kill_{cmpd}/%kill_{DMSO}) were calculated.

Brine shrimp lethality assay: The brine shrimp, *Artemia salina* (L.), lethality test was carried out using a modification of the procedure by McLaughlin [70]. *A. salina* eggs were hatched in a sea salt solution (Instant Ocean®, 38 g/L) with an incandescent light bulb as the heat source. After 48 h, the newly hatched nauplii were counted using a micropipette and transferred to 20 mL vials. Nine vials were prepared, each containing 10 *A. salina* nauplii in 10 mL of sea salt solution (same as the hatching solution). Three vials were labeled as controls with the first containing no DMSO, another

with 10 µL, and the last with 100 µL DMSO. Three replicate vials contained 10 µL of 1% essential oil solution in DMSO, and the other 3 were prepared by adding 100 µL of 1% essential oil solution in DMSO. Surviving *A. salina* were counted after 24 h. LC₅₀ values (Table 3) were determined using the Reed-Muench method [65].

Butterfly larvicidal activity assay: Cabbage white butterfly, *Pieris rapae* (L.), eggs were purchased from Carolina Biological Supply (Item # 144100). The egg strip was cut into smaller sections and the eggs placed onto organically grown collared green (*Brassica oleracea* L.) leaves. The eggs hatched in 48 h, after which they were placed on collard green leaves in a Petri dish. Three different essential oil concentrations were made in 1% tween water solutions (1000 µg/mL, 500 µg/mL, and 250 µg/mL). Leaf disks (6.5 cm) were prepared from collar leaves and dipped into the different concentrations of sample solution separately for 30 sec. Solvents were evaporated in a fume hood for 30 min. The control was made from 1% tween water solution only. The leaf disks were each placed in a Petri dish and 10 *P. rapae* larvae were transferred into each disk. The number of dead larvae was counted after 48 h.

Termiticidal assay: Worker termites, *Reticulitermes virginicus* (Banks), were purchased from Carolina Biological Supply (Item # 143736). The termiticidal assay was carried out using a 6-well culture plate, each well fitted with a filter paper disc. Solutions of the essential oils and major components were prepared in 1% aqueous Tween® 80 solution at 2000, 1000, and 500 µg/mL. Sample solutions (200 µL) of each concentration were sprayed into 3 of the wells. Water and 1% aqueous Tween® 80 were used as controls in the remaining wells. Six termites were placed into each well and termiticidal activity was determined after 24 h.

Mosquito larvicidal assay: Larvae of *Culex pipiens* L. were obtained from Carolina Biological Supply (Item # 144478). For the bioassay, 10 mL of sterile water was placed in five 20 mL vials. Into each vial, 10 mosquito larvae were transferred using a soft brush. Three vials were labeled as control with the first one containing 10 µL DMSO, the second containing 100 µL DMSO and the third containing only sterile water. Into the remaining 2 vials were added 10 µL of 1% solution of essential oil in DMSO and 100 µL of 1% essential oil/DMSO solution (i.e., final concentrations of 10 and 100 µg/mL). Surviving mosquito larvae were counted after 24 h. The experiments were carried out at $23 \pm 2^\circ\text{C}$. An analogous assay was carried out using midge (*Chaoborus plumicornis* Fab.) larvae (glassworm), which were obtained from a local aquarium shop.

Fruit fly lethality assay: Wild type *Drosophila melanogaster* Meigen were obtained from Carolina Biological Supply and a breeding colony maintained using a *Drosophila* culture kit. *Drosophila* medium (2 mL) was placed into each of five 20 mL glass vials. Three vials were labeled as control, the first containing only *Drosophila* medium, the second with 20 µL DMSO, and the third with 150 µL of DMSO. Another 2 vials contained 20 µL of 1% essential oil solution in DMSO, and the other vial was prepared by adding 150 µL of 1% essential oil solution in DMSO. Ten individual fruit flies were transferred into each vial. The bioassay was carried out in triplicate. Surviving fruit flies were counted after 24 h.

Fire ant lethality assay: Worker red imported fire ants, probably *Solenopsis invicta* × *richteri* hybrid [71], were collected from the University of Alabama in Huntsville. Sample solutions of 1000 µg/mL, 500 µg/mL and 250 µg/mL were prepared in 1% aqueous Tween-80® solution. The control was 1% Tween solution. Each

assay was carried out using a 400 mL beaker, fitted with a filter paper disk on the bottom. The filter paper was sprayed with 600 µL of sample solution and 20 fire ant workers were transferred to the beaker, which was sealed with Parafilm®. The mortality of fire ants was recorded after 24 h. The bioassay was carried out at room temperature in triplicate. LC₅₀ values were calculated using the Reed Muench method [65].

Nematicidal assay: A nematicidal assay using *Caenorhabditis elegans* (Maupas) was carried out using a modification of the procedure of Park and co-workers [72]. A 1% solution of each essential oil and major component in DMSO was prepared. Dilutions of the sample solution were prepared in sterile water solution beginning with 50 µL of the 1% essential oil solution plus 50 µL sterile water. The sample solution was serially diluted (1:1) with sterile water in a 96-well plate. Into each well, 10-30 C.

elegans (mixtures of juvenile and adult nematodes, male: female: juvenile ~1:1:2) per 50 µL of sample solution. Sterile water and serially diluted DMSO were used as controls. The dead and living nematodes were counted after 24 h under a microscope. Dead nematodes were identified by their immobility, and straight body, even after transfer to clean water. LC₅₀ values (Table 3) were determined using the Reed-Muench method [65].

Acknowledgments – PS and PP are grateful for MMAMC (Mahendra Morang Adarsha Multiple Campus) for providing lab equipment for the essential oil extraction. We would also like to thank Professor Debra Moriarity for technical assistance with the cytotoxicity assays and Professor Bernhard Vogler for technical assistance with the GC-MS data collection. WNS would like to thank an anonymous donor for the gift of the GC-MS instrumentation.

References

- [1] Mabberley DJ. (2008) *Mabberley's Plant-Book*, 3rd Ed. Cambridge University Press, UK.
- [2] (a) Joshi S, Padalia R, Bisht D, Mathela C. (2009) Terpenoid diversity in the leaf essential oils of Himalayan Lauraceae species. *Chemistry and Biodiversity*, **6**, 1364-1373; (b) Singh R, Jawaid T. (2012) *Cinnamomum camphora* (Kapur): Review. *Pharmacognosy Journal*, **4**, 1-5.
- [3] Hsieh TJ, Chen CH, Lo WL, Chen CY. (2006) Lignans from the stem of *Cinnamomum camphora*. *Natural Product Communications*, **1**, 21-25.
- [4] Lee HJ, Hyun EA, Yoon WJ, Kim BH, Rhee MH, Kang HK, Cho JY, Yoo ES. (2006) *In vitro* anti-inflammatory and anti-oxidative effects of *Cinnamomum camphora* extracts. *Journal of Ethnopharmacology*, **103**, 208-216.
- [5] Hashimoto K, Ohtani Y, Sameshima K. (1997) The termiticidal activity and its transverse distribution in camphor (*Cinnamomum camphora*) wood. *Journal of the Japanese Wood Research Society*, **43**, 566-573.
- [6] Kim SI, Na YE, Yi JH, Kim BS, Ahn YJ. (2007) Contact and fumigant toxicity of oriental medicinal plant extracts against *Dermanyssus gallinae* (Acari: Dermanyssidae). *Veterinary Parasitology*, **145**, 377-382.
- [7] Chen Y, Dai G. (2012) Antifungal activity of plant extracts against *Colletotrichum lagenarium*, the causal agent of anthracnose in cucumber. *Journal of the Science of Food and Agriculture*, **92**, 1937-1943.
- [8] Haque R, Mondal S, Ghosh P. (2011) Investigation of *in vitro* anthelmintic activity of *Cinnamomum camphora* leaves. *International Journal of Drug Development & Research*, **3**, 301-306.
- [9] Gupta R, Bajpai KG, Johri S, Saxena AM. (2008) An overview of 253 Indian novel traditional medicinal plants with anti-diabetic potentials. *African Journal of Traditional, Complementary and Alternative Medicines*, **5**, 1-17.
- [10] Ahmed A, Choudhary MI, Farooq A, Demirci B, Demirci F, Başer KHC. (2000) Essential oil constituents of the spice *Cinnamomum tamala* (Ham.) Nees & Eberm. *Flavour and Fragrance Journal*, **15**, 388-390.
- [11] Sharma G, Nautiyal A. (2011) *Cinnamomum tamala*: A valuable tree from Himalayas. *International Journal of Medicinal and Aromatic Plants*, **1**, 1-4.
- [12] Kumar S, Vasudeva N, Sharma S. (2012) Pharmacological and pharmacognostical aspects of *Cinnamomum tamala* Nees and Eberm. *Journal of Pharmacy Research*, **5**, 480-484.
- [13] Kunwar RM, Adhikari N. (2005) Ethnomedicine of Dolpa district, Nepal: The plants, their vernacular names and uses. *Lyonia*, **8**, 43-49.
- [14] Phattacharjee PP, Ray DC. (2010) Pest management beliefs and practices of Manipuri rice farmers in Barak Valley, Assam. *Indian Journal of Traditional Knowledge*, **9**, 673-676.
- [15] Baruah A, Nath SC. (2006) Leaf essential oils of *Cinnamomum glanduliferum* (Wall) Meissn and *Cinnamomum glaucescens* (Nees) Meissn. *Journal of Essential Oil Research*, **18**, 200-202.
- [16] Ravindran PN, Babu K, Shylaja M (Eds). (2004) *Cinnamon and Cassia. The genus Cinnamomum*, CRC Press, Boca Raton, Florida, USA.
- [17] Prakash B, Singh P, Yadav S, Singh SC, Dubey NK. (2013) Safety profile assessment and efficacy of chemically characterized *Cinnamomum glaucescens* essential oil against storage fungi, insect, aflatoxin secretion and as antioxidant. *Food and Chemical Toxicology*, **53**, 160-167.
- [18] Ho C, Wang E, Su Y. (2009) [Essential oil compositions and bioactivities of the various parts of *Cinnamomum camphora* Sieb. var. *linaloolifera* Fujita]. *Forest Research Quarterly*, **31**, 77-96. [article in Japanese]
- [19] Senanayake U. (1977) *The Nature, Description and Biosynthesis of Volatiles in Cinnamomum ssp.*, Ph.D. thesis, University of New South Wales, Australia.
- [20] Chopra R, Nayar S, Chopra I. (1956) *Glossary of Indian Medicinal Plants*. CSIR, New Delhi, India, 65-66G.
- [21] Rana V, Blazquez M. (2005) GC-MS analysis of the essential oil of *Cinnamomum tamala* L. leaves. *Indian Perfumer*, **49**, 475-478.
- [22] Gulati B, Agarwal S, Thappa R, Dhar K. (1977) Essential oil of Tejpatri (Kumaon) from *Cinnamomum tamala*. *Indian Perfumer*, **21**, 15-20.
- [23] Sood R, Padha C, Talwar Y, Jamwal R, Chopra M, Rao P. (1979) Essential oils from the leaves of *Cinnamomum tamala* Nees and Eberm growing in Himachal Pradesh. *Indian Perfumer*, **23**, 75-78.
- [24] Nigam M, Ahmed A. (1990) The essential oil of *Cinnamomum tamala* II. *Papai Journal*, **12**, 24-25.
- [25] Adhikary SR, Tuladhar BS, Sheak A, van Beek TA, Posthumus MA, Lelyveld GP. (1992) Investigation of Nepalese essential oils. I. The oil of *Cinnamomum glaucescens* (sugandha kokila). *Journal of Essential Oil Research*, **4**, 151-159.
- [26] Mishra AK, Dwivedi SK, Kishore N, Dubey NK. (1991) Fungistatic properties of the essential oil of *Cinnamomum camphora*. *Pharmaceutical Biology*, **29**, 259-262.
- [27] Liu CH, Mishra AK, He B, Tan RX. (2001) Composition and antifungal activity of essential oils from *Artemisia princeps* and *Cinnamomum camphora*. *International Pest Control*, **43**, 72-74.
- [28] Singh P. (2008) Fungal contamination of raw material of some herbal drugs and recommendation of *Cinnamomum camphora* oil as herbal fungitoxicant. *Microbial Ecology*, **56**, 555-560.
- [29] Rangari P. (2011) Screening of antibacterial sensitivity of essential oils of camphor and cinnamon. *Shodh Anusandhan Samachar*, **1**, 22-26.
- [30] Anderson JE, Goetz CM, McLaughlin JL, Suffness M. (1991) A blind comparison of simple bench-top bioassays and human tumour cell cytotoxicities as antitumor prescreens. *Phytochemical Analysis*, **2**, 107-111.

- [31] Traboulsi AF, Taoubi K, El-Haj S, Bessiere JM, Rammal S. (2002) Insecticidal properties of essential plant oils against the mosquito *Culex pipiens molestus* (Diptera: Culicidae). *Pest Management Science*, **58**, 491-495.
- [32] Traboulsi AF, El-Haj S, Tueni M, Taoubi K, Nader NA, Mrad A. (2005) Repellency and toxicity of aromatic plant extracts against the mosquito *Culex pipiens molestus* (Diptera: Culicidae). *Pest Management Science*, **61**, 597-604.
- [33] Zhu J, Zeng X, Liu T, Qian K, Han Y, Xue S, Tucker B, Schultz G, Coats J, Rowley W, Zhang A. (2006) Adult repellency and larvicidal activity of five plant essential oils against mosquitoes. *Journal of the American Mosquito Control Association*, **22**, 515-522.
- [34] Koliopoulos G, Pitarokili D, Kioulos E, Michaelakis A, Tzakou O. (2010) Chemical composition and larvicidal evaluation of *Mentha*, *Salvia*, and *Melissa* essential oils against the West Nile virus mosquito *Culex pipiens*. *Parasitology Research*, **107**, 327-335.
- [35] Ranaweera SS. (1996) Mosquito-larvicidal activity of some Sri Lankan plants. *Journal of the National Science Foundation of Sri Lanka*, **24**, 63-70.
- [36] Amer A, Melhorn H. (2006) Larvicidal effects of various essential oils against *Aedes*, *Anopheles*, and *Culex* larvae (Diptera, Culicidae). *Parasitology Research*, **99**, 466-472.
- [37] Tang L, Sun YY, Zhang QP, Zhou Y, Zhang N, Zhang ZX. (2013) Fumigant activity of eight plant essential oils against workers of red imported fire ant, *Solenopsis invicta*. *Sociobiology*, **60**, 35-40.
- [38] Liu CH, Mishra AK, Tan RX, Tang C, Yang H, Shen YF. (2006) Repellent and insecticidal activities of essential oils from *Artemisia princeps* and *Cinnamomum camphora* and their effect on seed germination of wheat and broad bean. *Bioresource Technology*, **97**, 1969-1973.
- [39] Amer A, Melhorn H. (2006) Repellency effect of forty-one essential oils against *Aedes*, *Anopheles*, and *Culex* mosquitoes. *Parasitology Research*, **99**, 478-490.
- [40] van Tol RWHM, Swarts HJ, van der Linden A, Visser JH. (2007) Repellence of the red bud borer *Resseliella oculiperda* from grafted apple trees by impregnation of rubber budding strips with essential oils. *Pest Management Science*, **63**, 483-490.
- [41] Rozman V, Kalinovic I, Korunic Z. (2007) Toxicity of naturally occurring compounds of Lamiaceae and Lauraceae to three stored-product insects. *Journal of Stored Products Research*, **43**, 349-355.
- [42] Zhu BCR, Henderson G, Chen F, Fei H, Laine RA. (2001) Evaluation of vetiver oil and seven insect-active essential oils against the Formosan subterranean termite. *Journal of Chemical Ecology*, **27**, 1617-1625.
- [43] Chang ST, Cheng SS. (2001) Antitermitic activity of leaf essential oils and components from *Cinnamomum osmophloeum*. *Journal of Agricultural and Food Chemistry*, **50**, 1389-1392.
- [44] Cheng SS, Wu CL, Chang HT, Kao YT, Chang ST. (2004) Antitermitic and antifungal activities of essential oil of *Calocedrus formosana* leaf and its composition. *Journal of Chemical Ecology*, **30**, 1957-1967.
- [45] Park IK, Shin SC. (2005) Fumigant activity of plant essential oils and components from garlic (*Allium sativum*) and clove bud (*Eugenia caryophyllata*) oils against the Japanese termite (*Reticulitermes speratus* Kolbe). *Journal of Agricultural and Food Chemistry*, **53**, 4388-4392.
- [46] Clausen CA, Yang VW. (2008) Fumigant toxicity of essential oils to *Reticulitermes flavipes*. *Proceedings of the 104th Annual Meeting of the American Wood Protection Association*, **104**, 49-53.
- [47] Seo SM, Kim J, Lee SG, Shin CH, Shin SC, Park IK. (2009) Fumigant antitermitic activity of plant essential oils and components from ajowan (*Trachyspermum ammi*), allspice (*Pimenta dioica*), caraway (*Carum carvi*), dill (*Anethum graveolens*), geranium (*Pelargonium graveolens*), and litsea (*Litsea cubeba*) oils against the Japanese termite (*Reticulitermes speratus* Kolbe). *Journal of Agricultural and Food Chemistry*, **57**, 6596-6602.
- [48] Kong JO, Lee SM, Moon YS, Lee SG, Ahn YJ. (2007) Nematicidal activity of cassia and cinnamon oil compounds and related compounds toward *Bursaphelenchus xylophilus* (Nematoda: Parasitaphelenchidae). *Journal of Nematology*, **39**, 31-36.
- [49] Kim J, Seo SM, Park IK. (2011) Nematicidal activity of plant essential oils and components from *Gaultheria fragrantissima* and *Zanthoxylum alatum* against the pine wood nematode, *Bursaphelenchus xylophilus*. *Nematology*, **13**, 87-93.
- [50] Ntalli NG, Ferrari F, Giannakou I, Menkissoglu-Spiroudi U. (2010) Phytochemistry and nematicidal activity of the essential oils from 8 Greek Lamiaceae aromatic plants and 13 terpene components. *Journal of Agricultural and Food Chemistry*, **58**, 7856-7863.
- [51] Ntalli NG, Ferrari F, Giannakou I, Menkissoglu-Spiroudi U. (2011) Synergistic and antagonistic interactions of terpenes against *Meloidogyne incognita* and the nematicidal activity of essential oils from seven plants indigenous to Greece. *Pest Management Science*, **67**, 341-351.
- [52] Al-Banna L, Darwish RM, Aburjai T. (2003) Effect of plant extracts and essential oils on root-knot nematode. *Phytopathologia Mediterranea*, **42**, 123-128.
- [53] Kinghorn AD, Balandrin MF (Eds). (1993) *Human Medicinal Agents from Plants*. ACS Symposium Series 534. American Chemical Society, Washington DC.
- [54] Copping LG, Menn JJ. (2000) Biopesticides: A review of their action, applications and efficacy. *Pest Management Science*, **56**, 651-676.
- [55] Duke SO, Baerson SR, Dayan FE, Rimando AM, Scheffler BE, Tellez MR, Wedge DE, Schrader KK, Akey DH, Arthur FH, De Lucca AJ, Gibson DM, Harrison HF, Peterson JK, Gealy DR, Tworokoski T, Wilson CL, Morris JB. (2003) United States Department of Agriculture – Agriculture Research Service research on natural products for pest management. *Pest Management Science*, **59**, 708-717.
- [56] Copping LG, Duke SO. (2007) Natural products that have been used commercially as crop protection agents. *Pest Management Science*, **63**, 524-554.
- [57] Soylu EM, Kurt S, Soylu S. (2010) *In vitro* and *in vivo* antifungal activities of the essential oils of various plants against tomato grey mould disease agent *Botrytis cinerea*. *International Journal of Food Microbiology*, **143**, 183-189.
- [58] Macias FA, Molinillo JMG, Varela RM, Galindo JCG. (2007) Allelopathy – a natural alternative for weed control. *Pest Management Science*, **63**, 327-348.
- [59] Rajendra S, Srranjini V. (2008) Plant products and fumigants for stored-product insect control. *Journal of Stored Products Research*, **44**, 126-135.
- [60] Ntalli NG, Caboni P. (2012) Botanical nematocides: A review. *Journal of Agricultural and Food Chemistry*, **60**, 9929-9940.
- [61] Andrés MF, González-Coloma A, Sanz J, Burillo J, Sainz P. (2012) Nematicidal activity of essential oils: A review. *Phytochemistry Reviews*, **11**, 371-390.
- [62] Adams RP. (2007) *Identification of Essential Oil Components by Gas Chromatography / Mass Spectrometry*, 4th Ed. Allured Publishing, Carol Stream, Illinois, USA.
- [63] Kennedy JE, Davé PC, Harbin LN, Setzer WN. (2011) Allelopathic potential of *Sassafras albidum* and *Pinus taeda* essential oils. *Allelopathy Journal*, **27**, 111-122.
- [64] Zar JH. (1996) *Biostatistical Analysis*, 3rd Ed. Prentice Hall, Upper Saddle River, New Jersey, USA, 123-137.
- [65] Reed LZ, Muench H. (1938) A simple method of estimating fifty per cent endpoints. *American Journal of Hygiene*, **27**, 493-497.
- [66] Sahm DH, Washington JA (1991) Antibacterial susceptibility tests: Dilution methods. In *Manual of Clinical Microbiology*. Balows A, Hausler WJ, Herrmann KL, Isenberg HD, Shamody HJ (Eds). American Society for Microbiology, Washington DC, USA.
- [67] Soule HD, Vazquez J, Long A, Albert S, Brennan M. (1973) A human cell line from a pleural effusion derived from a breast carcinoma. *Journal of the National Cancer Institute*, **51**, 1409-1416.
- [68] Setzer WN, Setzer MC, Hopper AL, Moriarity DB, Lehrman GK, Niekamp NK, Morcomb SM, Bates RB, McClure KJ, Stessman CC, Haber WA. (1998) The cytotoxic activity of a *Salacia* liana species from Monteverde, Costa Rica, is due to a high concentration of tingenone. *Planta Medica*, **64**, 583.

- [69] Ferrari M, Fornasiero MC, Isetta AM. (1990) MTT colorimetric assay for testing macrophage cytotoxic activity in vitro. *Journal of Immunological Methods*, **131**, 165-172.
- [70] McLaughlin JL. (1990) Bench-top bioassays for the discovery of bioactive compounds in higher plants. *Brenesia*, **34**, 1-14.
- [71] Chen L, Lu YY, Hu QB, Fadamiro HY. (2012) Similarity in venom alkaloid chemistry of alate queens of imported fire ants: Implication for hybridization between *Solenopsis richteri* and *S. invicta* in the southern United States. *Chemistry and Biodiversity*, **9**, 702-713.
- [72] Park IK, Kim J, Lee SG, Shin SC. (2007) Nematicidal activity of plant essential oils and components from ajowan (*Trachyspermum ammi*), allspice (*Pimenta dioica*) and litsea (*Litsea cubeba*) essential oils against pine wood nematode (*Bursaphelenchus xylophilus*). *Journal of Nematology*, **39**, 257-279.

Essential Oil Characterization of Two Azorean *Cryptomeria japonica* Populations and Their Biological Evaluations

Cristina Moiteiro^{a,*}, Teresa Esteves^a, Luís Ramalho^b, Rosario Rojas^c, Sandra Alvarez^d, Susana Zacchino^d and Helena Bragança^e

^aUniversidade de Lisboa, Faculdade de Ciências, Departamento de Química e Bioquímica; Centro de Química e Bioquímica, Campo Grande, 1749-016 Lisboa, Portugal

^bLaboratório Nacional de Energia e Geologia, LBA-Laboratório de Biocombustíveis e Ambiente, Estrada do Paço do Lumiar, 22, 1648-039 Lisboa, Portugal

^cLaboratorios de Investigación y Desarrollo, Facultad de Ciencias y Filosofía, Universidad Peruana Cayetano Heredia, Av. Honorio Delgado 430 Urb. Ingeniería S.M.P. Lima 31, Perú

^dFacultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Área Farmacognosia, Suipacha 531, 200 Rosario, Argentina

^eInstituto Nacional de Investigação Agrária e Veterinária I.P., Quinta do Marquês, 2780-159 Oeiras, Portugal

cmmoiteiro@fc.ul.pt

Received: July 15th, 2013; Accepted: September 10th, 2013

Essential oils from foliage, bark and heartwood of *Cryptomeria japonica* D. Don from Azores Archipelago (Portugal) were analyzed by GC and GC-MS. Two populations, of black and reddish heartwood color, were studied. The main compounds found in the foliage of both populations were α -pinene (9.6-29.5%), (+)-phyllocladene (3.5-26.5%), *ent*-kaur-16-ene (0.2-20.6%), sabinene (0.5-19.9%) and limonene (1.4-11.5%), with a large variation in individual compounds from each population. Heartwood oils were characterized by a high content of cubebol (2.8-39.9%) and *epi*-cubebol (4.1-26.9%) isomers, which were absent in the foliage. Elemol and eudesmol isomers were found in the foliage and heartwood oils, while (+)-phyllocladene was absent in heartwood. Black and reddish bark oils were composed of the diterpenes dehydroferruginol (1.9-5.1%) and ferruginol (2.6-11.5%), along with the sesquiterpenes δ -cadinene (10.4-15.9%), α -muurolene (3.3-5.4%), *epi*-zonarene (4.0-5.0%), cubenol (9.3-14.0%), τ -muurolol (4.8-10.7%), β -eudesmol (3.0-9.9%), γ -eudesmol (1.9-7.0%) and hedyacariol (1.4-6.2%). Azorean *C. japonica* oils exhibited significant chemical differences compared with native plants from Asia.

The essential oils showed moderate antimicrobial activity against the pathogenic fungus *Cryptococcus neoformans* and human pathogenic bacteria (especially against multidrug-resistant *Mycobacterium tuberculosis*). The antimicrobial activity of the essential oils may be attributed to compounds such as *ent*-kaur-16-ene, (+)-phyllocladene, ferruginol and elemol, which are present in different proportions within the complex oil mixture. These results suggest a potential use for *C. japonica* oils obtained from wood industry leftovers.

Keywords: *Cryptomeria japonica*, Essential oils, GC-MS, Pathogenic fungi, *Mycobacterium tuberculosis*.

Cryptomeria japonica D. Don. (Cupressaceae), also known as Sugi or Japanese cedar, is an Asian conifer that was introduced into the Azores Islands in the mid-19th century. *C. japonica* progressively overcame endemic tree species, and nowadays is widely distributed in the islands, representing 57% of the total wood producing forest area, and with sawmills generating wastes having only minor and non-commercial applications [1]. Typically, *C. japonica* heartwood has a red color, but, in some cases, a black/dark color can also occur, reducing its commercial value [2,3]. During the last years, the incidence of reddish heartwood color of *C. japonica* trees in the Azores islands has been declining, leading to a rise of economically lower valued black heartwood specimens. The regional Azores government, in 1998, initiated a Forest Breeding Program in order to evaluate the genetic basis of the Azores sugi population. This program intended to improve the genetic species of *Cryptomeria* and to increase the productivity and quality of wood [4].

In recent years, particular interest has been placed on the study of *C. japonica* essential oils (EOs) from Japan, Korea, Taiwan and China, with some significant activities being reported, such as antiulcer [5], antifungal [6,7], antimicrobial [8-10], cancer chemopreventive [11], neuropharmacological [12], antimosquito [13-15], antitermite [16,17] and insecticidal [18,19].

In a strategy of valorization of an unrecognized natural resource in the Azorean islands, the main purpose of this study was to find a promising potential use for the unutilized bark, foliage and wood leftovers of *C. japonica*, by obtaining and chemically identifying their EOs, as well as evaluating their antimicrobial activities against wood deteriorating fungi, human pathogenic fungi and bacteria. This is the first study reporting the composition of the EOs of the two different *C. japonica* populations, black and reddish heartwood trees, of Azores, Portugal (Faial island).

For these purposes, two tree sampling sites were chosen, taking into account sugi heartwood color, a physical property that could influence the composition of the EOs. The black heartwood population was assigned as A, and the reddish heartwood population as B. Table 1 shows the yields of obtained EOs from the different tissues of *C. japonica* analyzed.

Comparing oil yields, taking into account the heartwood color of each *C. japonica* specimen, the variation in the foliage of black heartwood trees was higher than in the oils of reddish heartwood foliage, based on fresh weight (A- 1.1-1.9% and B- 0.5-0.8%). For heartwood oils of both populations, the yields obtained showed little differences (A- 0.6-2.0% and B- 0.5-2.1%), with the same being

Table 1: Percentage limits of the components detected in EOs from the foliage, woods and barks of *C. japonica* from two populations of Faial Island (Azores).

Compound	RI ^a	Relative content (%)					
		Foliage A*	Foliage B*	Heartwood A*	Heartwood B*	Bark A*	Bark B*
Tricyclene	830	0.3 - 0.4	t - 0.4	-	-	-	-
α -Thujene	915	t - 1.2	0.3 - 1.2	-	-	-	-
α-Pinene	931	9.6 - 29.5	13.5 - 23.8	-	-	0.8	2.1
Camphene	943	1.8 - 2.8	0.7 - 2.6	-	-	-	-
Sabinene	964	0.5 - 19.9	1.0 - 10.5	-	-	-	-
β -Pinene	970	0.5 - 1.7	0.7 - 1.2	-	-	-	-
Myrcene	978	2.5 - 3.9	2.4 - 3.1	-	-	-	-
Δ^3 -Carene	1007	0.5 - 1.9	t - 1.4	-	-	-	-
α -Terpinene	1011	t - 2.1	0.1 - 1.5	-	-	-	-
<i>p</i> -Cymene	1013	0.4 - 0.5	t - 0.4	-	-	-	-
Limone	1023	1.4 - 5.0	1.8 - 11.5	-	-	T	0.6
γ -Terpinene	1049	t - 3.5	0.2 - 2.5	-	-	T	-
Terpinolene	1079	0.2 - 1.3	0.3 - 1.1	-	-	-	-
2- <i>p</i> -Menthen-1-ol	1106	t - 0.3	t - 0.2	-	-	-	-
Terpinen-4-ol	1165	t - 8.9	0.4 - 4.6	-	-	-	-
α -Terpineol	1174	t - 0.4	t - 0.3	-	-	-	0.4
Bornyl acetate	1273	1.1 - 2.6	0.5 - 1.8	-	-	-	-
α -Cubebene	1354	-	-	0.1 - 0.4	t - 0.4	-	-
α -Copaene	1383	-	-	T	t - 0.2	-	-
β -Elemene	1394	-	-	0.2 - 0.8	0.4 - 1.0	-	-
β -Caryophyllene	1426	t - 0.2	t - 0.2	t - 0.7	-	-	-
<i>trans</i> - β -Farnesene	1454	-	-	t - 0.5	0.4 - 0.8	-	-
α -Humulene	1461	-	-	0.4 - 0.7	t - 0.8	0.4	0.7
<i>epi</i> -Bicyclosquiphellandrene	1469	-	-	-	-	1.3	3.4
γ -Murolene	1471	-	-	0.8 - 1.4	t - 1.5	0.4	-
Germacrene-D	1480	0.2 - 0.4	0.2 - 0.8	T	-	-	-
β -Cubebene	1499	-	-	t - 2.0	t - 1.9	-	-
<i>epi</i> -Cubebol	1500	-	-	4.1 - 26.9	12.3 - 21.2	-	0.6
α-Murolene	1503	0.1 - 1.5	0.3 - 0.5	2.0 - 3.3	1.5 - 2.9	3.3	5.4
γ -Cadinene	1516	0.2 - 1.2	0.2 - 0.3	t - 3.7	-	-	-
Cubebol	1520	-	-	2.7 - 39.9	14.6 - 33.8	-	0.5
<i>epi</i> -Zonarene	1522	-	-	-	-	4.0	5.0
δ-Cadinene	1524	0.8 - 6.5	0.8 - 1.4	6.2 - 10.8	6.4 - 10.9	10.4	15.9
Cadina-1,4-diene	1526	-	-	0.5 - 1.6	0.6 - 1.0	2.2	2.7
α -Calacorene	1532	-	-	0.43 - 0.6	0.4 - 0.9	1.1	1.9
Elemol	1542	0.2 - 12.7	4.1 - 8.1	1.7 - 14.1	1.8 - 9.2	2.6	1.1
Nerolidol	1551	-	-	t - 0.3	t - 0.3	0.5	1.4
Caryophyllene oxide	1580	0.2 - 3.0	0.3 - 0.7	t - 0.6	t - 0.5	-	-
<i>Allo</i> -Hedycariol	1580	-	-	0.7 - 2.9	0.7 - 2.5	2.3	1.7
Cubebol	1619	t - 0.7	0.3 - 1.2	5.5 - 17.2	4.0 - 18.4	14.0	9.3
γ-Eudesmol	1626	0.2 - 9.1	2.8 - 7.6	t - 2.9	t - 1.8	7.0	1.9
τ -Cadinol	1635	0.6 - 4.7	0.9 - 1.4	1.1 - 3.9	1.0 - 3.9	2.4	2.5
π-Muurolol	1631	-	-	1.3 - 2.8	t - 6.9	10.7	4.8
β-Eudesmol	1646	0.2 - 4.8	2.1 - 4.9	2.5 - 6.1	2.7 - 6.0	9.9	3.0
α -Cadinol	1647	-	2.3 - 3.2	t - 5.2	t - 4.0	t	1.8
α -Eudesmol	1652	4.4 - 7.1	1.6 - 5.9	0.6 - 10.7	2.2 - 5.8	3.4	1.3
Hedycariol	1661	-	-	t - 5.5	1.1 - 3.3	6.2	1.4
Caladene	1665	-	t - 0.1	-	t - 0.3	1.2	1.4
Tremetone	1704	-	-	0.6 - 1.5	0.2 - 3.1	1.4	-
Pimara-8,15-diene	1938	0.5 - 2.1	t - 2.4	t - 0.4	t	-	-
Cupressene	1954	0.6 - 2.8	0.2 - 0.9	-	t	-	-
Sandaracopimaradiene	1967	0.5 - 1.3	0.3 - 1.4	-	-	-	-
Pimara-8(14),15-diene	1986	0.2 - 0.7	0.3 - 0.7	t - 0.4	t - 0.4	t	0.8
Isopimaradiene	2018	t - 0.3	0.2 - 0.3	-	-	-	-
(+)-Phyllocladene	2053	7.9 - 24.1	3.5 - 26.5	-	-	-	0.2
ent-Kaur-16-ene	2078	0.2 - 0.4	0.4 - 20.6	t - 0.3	t - 0.2	-	0.3
8 β -Hydroxysandaracopimarene	2149	0.6 - 3.6	1.2 - 3.3	0.2 - 0.9	0.2 - 0.9	-	-
Sandaracopimarinal	2193	t - 0.2	t	0.2 - 0.5	0.2 - 1.6	1.6	0.9
Kauran-16-ol	2223	-	t	0.1 - 0.5	t - 0.8	1.3	1.6
Sandaracopimararinol	2271	-	t	t - 0.5	t - 2.0	2.2	0.5
Dehydroferruginol	2296	-	t	t - 0.1	t - 0.5	1.9	5.1
Ferruginol	2305	-	t	t - 0.5	0.1 - 1.2	2.6	11.5
Identified components (%)		94.6 - 99.2	97.8 - 99.3	97.9 - 99.5	95.8 - 97.5	95.1	92.1
N° Identified components		38	44	38	37	30	33
Grouped components variation (%)							
Monoterpene hydrocarbons		37.1 - 50.5	37.1 - 43.4	-	-	0.8	2.7
Oxygenated monoterpenes		2.0 - 10.7	1.4 - 5.6	-	-	-	0.4
Sesquiterpene hydrocarbons		1.4 - 9.7	1.5 - 3.2	12.3 - 21.7	12.0 - 32.7	24.3	36.5
Oxygenated sesquiterpenes		13.1 - 30.2	16.0 - 28.2	77.9 - 84.0	60.7 - 80.4	60.4	31.5
Diterpene hydrocarbons		12.5 - 28.2	19.1 - 32.9	t - 1.1	t - 0.4	-	1.3
Oxygenated diterpenes		0.6 - 3.6	1.2 - 3.3	0.8 - 2.2	0.5 - 6.4	9.6	19.6
Oil yields (% w/w)		1.1 - 1.9	0.5 - 0.8	0.6 - 2.0	0.5 - 2.1	0.2	0.1

^aRelative to C₈-C₂₄ n-alkanes on a OV-101 column; t- trace (<0.1%); * Population A- Black color of the heartwood; Population B- reddish color of the heartwood.

observed for the barks (A- 0.2% and B- 0.1%). It appears that no significant relationship exists between the color of heartwood and oil yields. The sapwoods EO yield was extremely low (0.1%).

The 22 oil samples analyzed by GC were complex mixtures and, depending on the part of the plant, a range of 30 to 44 compounds

were identified, totalizing a percentage variation of 94.6-99.5%. The compounds identified in the EOs from the different tissues analyzed, as well as their percentage limits, are shown in Table 1. It can be seen that, from each individual sample in the same population, a larger range of percentage limits of each component among the foliage and heartwoods were detected.

The foliage of both populations are characterized by high contents of monoterpene hydrocarbons (A- 37.1-50.5% and B- 37.1-43.4%), followed by oxygenated sesquiterpenes (A- 13.1-30.2% and B- 16.0-28.2%), diterpene hydrocarbons (A- 12.5-28.2% and B- 19.1-32.9%), and oxygenated monoterpenes (A- 2.0-10.7% and B- 1.4-5.6%), but a low content of sesquiterpene hydrocarbons (A- 1.4-9.7% and B- 1.5-3.2%) and oxygenated diterpenes (A- 0.7-3.6% and B- 1.2-3.3%). The main common constituents of the foliage of the black and reddish population were α -pinene (A- 9.6-29.5% and B- 13.5-23.8%), (+)-phyllocladene (A- 7.9-24.1% and B- 3.5-26.5%), *ent*-kaur-16-ene (A- 0.2-0.4% and B- 0.4-20.6%), sabinene (A- 0.5-19.9% and B- 1.0-10.5%), limonene (A- 1.4-5.0% and B- 1.8-11.5%), elemol (A- 0.2-12.7% and B- 4.1-8.1%), α -eudesmol (A- 4.4-7.1% and B- 1.6-5.9%), β -eudesmol (A- 0.1-4.8% and B- 2.1-4.9%) and γ -eudesmol (A- 0.2-9.1% and B- 2.8-7.6%). The isolation and characterization of the isomers (+)-phyllocladene and *ent*-kaur-16-ene allowed us to conclude that, among the EO components, *ent*-kaur-16-ene is clearly the unique compound that is present only in the reddish population, while its isomer, (+)-phyllocladene, exists in high amount in both populations. α -Pinene, *ent*-kaur-16-ene, sabinene, elemol and eudesmol isomers have also been identified in the leaves of *C. japonica* EOs from Taiwan [6,15,18,19] and Korea [8], although with different percentages than those found in Azorean oils. On the contrary, the main compounds identified in the EOs of the leaves of *C. japonica* from China were α -elemol (20.1%), kaur-16-ene (14.8%), β -phellandrene (6.0%) and β -elemene (5.9%) [20].

Heartwood oils were characterized by a high content of two isomers, cubebol (A- 2.7-39.9% and B- 14.6-33.8%) and *epi*-cubebol (A- 4.1-26.9% and B- 12.3-21.2%), which are absent in leaves. So, the oxygenated sesquiterpenes form, respectively, 77.9-84.0% and 60.7-80.4% of the black and reddish heartwood populations, followed by sesquiterpene hydrocarbons (A- 12.3-21.7% and B- 12.0-32.7%). These groups of terpenes are also representative of the foliage EOs, while the more volatile monoterpene hydrocarbons and oxygenated monoterpenes present in the leaves were lacking in the heartwood oils. Furthermore, cubenol (A- 5.5-17.2% and B- 4.0-18.4%), elemol (A- 1.7-14.1% and B- 1.8-9.2%), α -eudesmol (A- 0.6-10.7% and B- 2.2-5.8%), β -eudesmol (A- 2.5-6.1% and B- 2.7-6.0%), τ -cadinol (A- 1.1-4.0% and B- 1.0-3.9%), τ -muurulol (A- 1.3-2.8% and B- t-6.9%) and δ -cadinene (A- 6.2-10.8% and B- 6.4-10.9%) were also present in both heartwood populations. The main foliage constituent, (+)-phyllocladene, is not present in heartwood EOs and, consequently, diterpene fractions were largely decreased in heartwood EOs, for both populations.

Oxygenated diterpenoids, such as sandaracopimarinal, kauran-16-ol, sandaracopamarinol, dehydroferruginol and ferruginol, are characteristic of black and reddish bark oils, representing about 9.6% and 19.6%, respectively, of the total oil percentage. δ -Cadinene (A- 10.4% and B- 15.9%), α -muurulene (A- 3.3% and B- 5.4%), and *epi*-zonarene (A- 4.0% and B- 5.0%) are the main sesquiterpene hydrocarbons present in both bark populations, which represent 24.3% and 36.5% of the total bark EOs. Oxygenated sesquiterpenes (A- 60.4% and B- 31.5%) are the major group of constituents of black and reddish bark populations, due to a considerable presence of cubenol (A- 14.0% and B- 9.3%), τ -muurulol (A- 10.7% and B- 4.8%), β -eudesmol (A- 9.9% and B- 3.0%), γ -eudesmol (A- 7.0% and B- 1.9%) and hedycariol (A- 6.2% and B- 1.4%). So, sesquiterpenes, both oxygenated and hydrocarbons, are the main group constituents for both bark EOs, although black barks had higher amounts of oxygenated sesquiterpenes than reddish barks.

Comparing the results obtained from Azores *C. japonica* EOs with those reported for the same species from other countries, deep differences, not only in chemical constituents, but also in their percentages, were observed. According to the study of Cheng *et al.* [14], there is no apparent relationship between the age of *C. japonica* specimens and the chemical composition of the leaf EO from the same origin. Cheng also suggests that the discrepancy with other authors could be related with different seasons in which the samples were collected. Therefore, the differences observed in the Azorean EOs are probably due to specific microclimate and temperature of a Macaronesian region, in association with edaphic factors, and to the plant adaptation to this specific environment. Another important contribution to the differences observed in compound amounts among the several EOs analyzed could be the use of individual *C. japonica* specimens, instead of a homogenized mixture of foliage and woods. These results may provide useful information for further studies of EOs of *C. japonica* trees from other Archipelago islands.

C. japonica EOs from foliage, heartwoods and barks, together with the main compounds isolated from EOs, were tested for antimicrobial activity against human pathogenic bacteria and pathogenic and wood surface contaminant fungi. The antifungal evaluation is depicted in Table 2. The fungus *B. cinerea* was inhibited by all EOs at a MIC of 100 μ g/mL with fungistatic activity; inversely, the isolated compounds, (+)-phyllocladene, *ent*-kaur-16-ene and a mixture of elemol, cubebol and *epi*-cubebol did not display any antifungal activity (MIC > 200 μ g/mL). The fungi *F. circinatum* and *C. parasitica* showed resistance towards all the samples tested. The wood surface contaminants *A. niger* and *T. harzianum* displayed a different behavior towards the tested samples, with almost all of them showing a moderate activity against the last fungus and being, in general, inactive against *A. niger* (MIC \geq 200 μ g/mL). All the oil samples tested inhibited the growth of *C. cladosporioides* and *Cladosporium* sp, with MICs between 100-200 μ g/mL. Among the pure compounds, only a mixture of cubebol, *epi*-cubebol and elemol, from the heartwood EO, exhibited antifungal activity against *Cladosporium* spp. This result is significant, since sapstain fungi are responsible for wood deterioration processes.

Additionally, none of the EOs inhibited any of the human pathogenic fungi up to 250 μ g/mL. Regarding the activity of the main compounds isolated from EOs, only sandaracopimarinal and a fraction enriched in elemol showed moderate MICs of 62.5 and 125 μ g/mL, respectively, against *C. neoformans* (Table 2). By taking into account this last result, we deepened the knowledge of the inhibition capacity of the oils and components by analyzing the percentage inhibition of *C. neoformans* by each sample at 250 μ g/mL and by each two-fold dilution of this concentration up to 7.8 μ g/mL (Table 3). It was shown that the EOs do not possess significant antifungal properties against *C. neoformans*, since the foliage EO inhibited only 15.6% of growth at 250 μ g/mL (it is important to take into account that at least 50% inhibition is necessary to consider either an oil or a compound with antifungal properties). In contrast, some components of the oil showed interesting activities. Among them, (+)-phyllocladene and ferruginol, showed 50% inhibition at 250 μ g/mL. In addition, sandaracopimarinal, which had a MIC₁₀₀ = 62.5 μ g/mL, showed 50% of inhibition at 31.25 μ g/mL. Also, the enriched fraction in elemol, which had a MIC₁₀₀ = 125 μ g/mL, showed a MIC₅₀ = 62.5 μ g/mL.

Furthermore, EOs of foliage and heartwood of *C. japonica* populations were tested *in vitro* against the sensitive strain H₃₇Rv

Table 2: Minimum inhibitory concentration (MIC in µg/mL) of *Cryptomeria japonica* EOs and of the main compounds isolated from them against pathogenic fungi and yeast.

	<i>Ca</i>	<i>Ct</i>	<i>Sc</i>	<i>Cn</i>	<i>Afl</i>	<i>Afu</i>	<i>Ani</i>	<i>Mg</i>	<i>Tr</i>	<i>Tm</i>	<i>Th</i>	<i>Bc</i>	<i>Cc</i>	<i>Csp</i>
EO Foliage	i	i	i	i	i	i	i	i	i	i	150	100*	200	200
EO Heartwood	i	i	i	i	i	i	i	i	i	i	200	100*	200*	i
EO Bark	n.t.	n.t.	n.t.	n.t.	n.t.	n.t.	200	n.t.	n.t.	n.t.	200	100*	200*	i
Sandaracopimarinal	i	i	i	i	i	i	i	i	i	i	n.t.	n.t.	n.t.	n.t.
Sandaracopimarinal	i	i	i	62.5	i	i	i	i	i	i	n.t.	n.t.	n.t.	n.t.
β-Eudesmol	i	i	i	i	i	i	i	i	i	i	n.t.	n.t.	n.t.	n.t.
Ferruginol	i	i	i	i	i	i	i	i	i	i	n.t.	n.t.	n.t.	n.t.
(+)-Phyllocladene	i	i	n.t.	i	i	i	i	i	i	i	i	i	i	i
ent-Kaur-16-ene	i	i	n.t.	i	i	i	i	i	i	i	>200	>200	>200	>200
Pt-Cj-Ele+Eud ^a	i	i	i	i	i	i	i	i	i	i	n.t.	n.t.	n.t.	n.t.
Mix-Elemol ^a	i	i	i	125	i	i	i	i	i	i	n.t.	n.t.	n.t.	n.t.
Cubebol+epi-Cubebol+Elemol	n.t.	n.t.	n.t.	n.t.	n.t.	n.t.	i	n.t.	n.t.	n.t.	i	i	i	100*
Amphotericin B	0.98	1.95	0.49	0.25	0.49	0.49	0.49	0.12	0.06	0.06	n.t.	n.t.	n.t.	n.t.
Terbinafine	1.95	1.95	3.90	0.49	0.98	0.98	1.96	0.04	0.01	0.03	n.t.	n.t.	n.t.	n.t.
Ketoconazole	0.49	0.98	0.49	0.25	0.12	0.49	0.25	0.06	0.03	0.03	n.t.	n.t.	n.t.	n.t.
Carbendazim	n.t.	n.t.	n.t.	n.t.	n.t.	n.t.	0.4*	n.t.	n.t.	n.t.	200*	200*	0.2*	0.2*

^aThese fractions are enriched in compounds; i = >250 µg/mL; n.t. = not tested; * = bacteriostatic / fungistatic activities; *Ca*: *C. albicans*; *Ct*: *C. tropicalis*; *Sc*: *S. cerevisiae*; *Cn*: *C. neoformans*; *An*: *A. niger*; *Afu*: *A. fumigatus*; *Afl*: *A. flavus*; *Mg*: *M. gypseum*; *Tr*: *T. rubrum*; *Tm*: *T. mentagrophytes*; *Th*: *T. harzianum*; *Bc*: *B. cinerea*; *Cc*: *C. cladosporioides*; *Csp*: *Cladosporium* sp.

Table 3: Antifungal activity (% inhibition = mean ± SD) of EOs and pure compounds against *Cryptococcus neoformans*.

	MIC (µg/mL)					
	250	125	62.5	31.25	15.6	7.8
EO Foliage	15.6 ± 2.2	9.9 ± 2.7	4.6 ± 0.5	0	0	0
EO Heartwood	0	0	0	0	0	0
(+)-Phyllocladene	58.7 ± 0.6	38.2 ± 7.5	12.3 ± 8.2	0	0	0
ent-kaur-16-ene	26.0 ± 2.02	24.8 ± 7.9	17.7 ± 2.2	12.1 ± 2.9	0	0
Sandaracopimarinal	28.4 ± 12.5	8.7 ± 1.1	4.0 ± 1.8	0	0	0
Sandaracopimarinal	100	100	100	54.3 ± 5.2	0	0
β-Eudesmol	41.7 ± 9.1	39.3 ± 10.4	33.8 ± 3.7	33.6 ± 6.2	34.3 ± 3.7	32.1 ± 2.8
Ferruginol	51.5 ± 13.1	48.7 ± 8.5	46.5 ± 3.8	44.6 ± 7.3	35.8 ± 2.6	31.7 ± 3.5
Elemol+Eudesmol	4.6 ± 10.8	1.4 ± 3.6	0	0	0	0
Mix-Elemol	100	100	50.5 ± 4.9	30.53 ± 4.49	18.5 ± 4.9	0
Amphotericin B	100	100	100	100	100	100
Ketoconazole	100	100	100	100	100	100

and a multidrug-resistant (MDR) strain of *Mycobacterium tuberculosis* (Table 4). EOs of the foliage (MIC 25 µg/mL) proved to be more active than those of the heartwood (MIC 50 µg/mL), especially against the MDR strain. The main compounds present in these EOs were isolated and tested in the antimycobacterial assay. None of them alone was active against the sensitive H₃₇Rv and MDR strain (MIC ≥ 25 µg/mL), except for (+)-phyllocladene and ent-kaur-16-ene, which showed a good activity (MIC 12.5 µg/mL), against the latter strain. These results lead us to think that the antimicrobial activity of the EOs of foliage may be attributed to the presence of active compounds such as ent-kaur-16-ene and (+)-phyllocladene, and probably also to the synergetic effects between them and the proportions in which they are present within a complex mixture. This complexity makes it often difficult to explain the aforementioned activities, as well as relating them to specific compounds or mixtures present in oils.

Table 4: *In vitro* activity of EOs and their main constituents against *M. tuberculosis* H₃₇Rv and MDR strains.

Samples	MIC (µg/mL)	
	H ₃₇ Rv	MDR
EO Foliage A	50	25
EO Foliage B	50	25
EO Heartwood A	50	50
EO Heartwood B	200	50
(+)-Phyllocladene	>25	12.5
ent-Kaur-16-ene	>25	12.5
Elemol	>25	>25
Cubebol+epi-Cubebol	>25	25
Isoniazid	0.125	4
Rifampin	0.063	16

A- Black population; B- reddish population.

The use of the wastes produced by the *C. japonica* wood industry in Azores Archipelago, until now being only seen as disposable, can represent an extremely important commercial advantage for the islands economy. The amount of wood/foliage not used and lost in the fields is considerable, and the upscale of the oil production is relatively easy and inexpensive.

Experimental

Plant material: Foliage (needle-like juvenile foliage), bark, and heartwood from 5 individual plants of 2 *C. japonica* populations, black and reddish heartwood color specimens from Faial island, were collected in July 2007 and provided by the Azorean Direcção Regional dos Recursos Florestais (DRRF). The individual plant samples were collected from 2 locations in Faial: black populations, altitude 140 m; latitude N 38:32:52.883; longitude W 28:38:21.171; reddish populations, altitude 480 m; latitude N 38:34:25.88; longitude W 28:40:20.203. The foliage and woods were air dried during 4 days and 6 weeks, respectively, at room temperature.

Extraction procedure: The leaves were cut into small pieces and heartwoods were ground in a Fritsch-Germany mill through a 1-mm screen width. Leaves and woods were then hydrodistilled for 4 h using a Clevenger-type apparatus [21]. The yields of EOs obtained are given in Table 1.

Analysis of essential oils: The EOs were analyzed by gas chromatography with flame ionization detection (GC-FID) and gas chromatography-mass spectrometry (GC-MS). The GC-FID analysis was performed on a Hewlett Packard model 5890 series II gas chromatograph equipped with an Automatic sampler HP 6890 series injector and a flame ionization detector (FID), using 2 different fused-silica capillary columns: OV-101 (50 m x 0.25 mm, 0.25 µm film thickness) and Supelcowax 10 (30 m x 0.25 mm, i.d. 0.25 µm film thickness). The oven temperature was held at 70°C for 5 min and then programmed to 210°C at 2°C/min. Injector and detector temperatures were 260°C and 270°C, respectively. The carrier gas was helium at a constant pressure of 55 kPa for the OV-101 and 45 kPa for the Supelco wax10 columns. The intensity of each peak was integrated. Each sample was analyzed 3 times. The average peak areas of all GC signals were added together and the percentage of each component peak was calculated by comparing its average area to the total areas.

The GC-MS analysis was carried out on a Carlo Erba GC-MS system, model HRGC/MS, with a Kratos mass detector model MS25RF (sector instrument), fitted with a HP-5MS column (30 m x 0.25 mm, 0.25 μ m film thickness). The oven was programmed initially from 70°C with 2 min hold up time to the final temperature of 250°C with a 5°C/min ramp. The final temperature hold time was 5 min, injector temperature was 250°C and GC/MS interface temperature was 280°C. The carrier gas was helium at a constant pressure of 90 kPa, with a split ratio 1:20, ionization energy 70 eV source at 200°C with full scan (25–450 amu) and a scan time of 0.5 s. The GC identification of the EO components is given in Table 1 and was assigned by comparison of their retention indices (RI) relative to a C₈–C₂₄ *n*-alkane series, and by analysis of standards in the same gas chromatographic conditions. The mass spectra of EO components were identified by comparing the obtained mass spectra of the analytes with those of authentic standards from the mass spectra of Wiley 6.0 and Mass Spectra Library (NIST 98) and with corresponding RI and mass spectral data of components from reference oils analyzed in our laboratory.

General methods: Medium pressure column chromatography (CC) was carried out on silica gel 60 (12 x 75 mm, 40–63 μ m, Buchi, 044884) filled columns. Preparative TLC, silica gel 60 F₂₅₄ (20x20 cm 0.25 mm thickness, Merck, Germany) or RP-18 preparative TLC (20 x 20 cm 0.25 mm thickness, F254, Merck, 1.05559) plates were used for isolation of compounds. 1D and 2D NMR, Brüker AMX 300 MHz NMR spectrometer. Optical rotation, $[\alpha]_D$, Perkin-Elmer 241 Polarimeter.

Isolation procedures: Samples of the oils (*ca.* 100 mg) from the foliage of both populations with the higher percentage of kaurene and phyllocladene diterpenoids, selected by GC analysis, were subjected to silica gel preparative TLC and eluted with *n*-hexane and *n*-hexane/ethyl acetate (3:1), successively. The fractions obtained were analyzed, and the pure compounds characterized by GC-FID, GC-MS, ¹H NMR, ¹³C NMR, MS and $[\alpha]_D$ measurements, and by comparison with data reported in the literature [22–24]. In order to determine the absolute stereochemistry of phyllocladene and kaurene, both pure isolated compounds were compared with retention times obtained in a cyclodextrine chiral capillary GC column (program: 70°C for 2 min to 175°C at 5°C/min and 175°C for 35 min; ASTEC CHIRALDEXTM G-T 30m x 0.25mm, 0.12 μ m) for the authentic samples [22], as well as by comparing $[\alpha]_D$ values with those in the literature [22,23]. The two isomers were lastly assigned as (+)-phyllocladene and *ent*-kaur-16-ene.

A sample of the oil obtained from heartwood (1g) was fractionated by silica gel medium pressure CC, using *n*-hexane and *n*-hexane/diethyl ether mixtures as eluents. One of the fractions obtained (F54-66; 144.5 mg) was subjected to preparative RP-18 TLC and eluted successively with methanol/water (8:2), to obtain elemol and a mixture of the isomers *epi*-cubebol and cubebol, which were analyzed and characterized by ¹H NMR, ¹³C NMR and GC-MS. NMR and MS data of elemol are in agreement with those of the literature [25]. Cubebol and *epi*-cubebol were also identified by comparison of the mass fragmentation patterns and ¹³C NMR spectra described in the literature [26,27].

Bioassays: Fungal and bacterial strains. Pathogenic fungi used in this study: *Botrytis cinerea* CCM1 899, *Trichoderma harzianum* CCM1 711, *Cladosporium cladosporioides* CCM1 478, and *Cladosporium* sp. CCM1 700 were obtained from the Culture Collection of Industrial Microorganisms (CCMI) Laboratório de Microbiologia Industrial, ex-INETI, Lisbon, Portugal. *Fusarium circinatum* FC0004 and *Cryphonectria parasitica* CO722 were

obtained from the Culture Collection of Instituto Nacional de Investigação Agrária e Veterinária I.P., Oeiras, Portugal.

Candida albicans ATCC 10231, *C. tropicalis* CCC 131 2000, *Saccharomyces cerevisiae* ATCC 9763, *Cryptococcus neoformans* ATCC 32264, *Aspergillus flavus* ATCC 9170, *A. fumigatus* ATCC 26934, *A. niger* ATCC 9029, *Trichophyton rubrum* CCC 110, *T. mentagrophytes* ATCC 9972 and *Microsporium gypseum* CCC 115 were obtained from the American Type Culture Collection (ATCC), Rockville, MD, USA, and CEREMIC (CCC), Centro de Referencia en Micología, Facultad de Ciencias Bioquímicas y Farmacéuticas, Rosario, Argentina. Inocula of cell or spore suspensions were adjusted to 1–5 x 10³ cells/spores with colony forming units (CFU) /mL [28].

Antimicrobial activity screenings: The Minimum Inhibitory Concentration (MIC) yielding no visible growth of the active compounds, and of the oils, was determined by the broth dilution method, according to reported procedures [28,29], from 250 to 0.98 μ g/mL, with a final DMSO concentration \leq 1%. Minimal bactericidal/fungicidal activities were determined by sub-cultivation of the samples into normal culture media at appropriate temperature and incubation times. MICs were visually recorded at 48 h for yeasts, and, for the rest of the fungi, at a time according to the control fungal growth. Carbendazim®, Amphotericin B, Terbinafine and Ketoconazole were used as positive controls for fungi. Compounds with MICs > 250 μ g/mL were considered inactive.

Inhibition percentage determination: The test was performed in 96-well microplates according to previous reports [30]. Tests were performed in duplicate. Reduction of growth for each compound concentration was calculated as follows: % of inhibition = 100 - (OD₄₀₅ ETW - OD₄₀₅ SCW)/(OD₄₀₅ GCW - OD₄₀₅ SCW), where ETWs = EOs or compounds test wells; GCW = growth control well; SCW = sterility control well and OD = optical density.

MIC₅₀ was defined as the lowest concentration of a compound that showed 50% reduction of the growth control and was determined from the results obtained in the inhibition percentage determination.

Antimycobacterial activity in vitro: Bioassays were conducted on sensitive H₃₇Rv ATCC 27294 (American Type Culture Collection) and MDR (clinical isolate, strain 02TBDM039EP097) *Mycobacterium tuberculosis* strains. For the preparation of the inoculum, a suspension of *M. tuberculosis* was made by mixing growth from slants (20–30 days old) with 100 μ L of Tween 80 in 0.2% bovine serum albumin (Sigma Chemical Co., St. Louis, MO). Turbidity of the suspension was then adjusted to a McFarland standard No. 1 (3 x 10⁷ CFU/mL) by adding Tween 80 and bovine serum albumin. The bacterial suspension (300 μ L) was further transferred to 7.2 mL of 7H9GC broth (4.7 g of Middlebrook), 7H9 broth base (Difco, Detroit, MI), 20 mL of 10% glycerol, 1 g of Bacto Casitone (Difco), 880 mL of distilled water and 100 mL of OADC (oleic acid, albumin, dextrose, catalase) (Remel, Lenexa, KS). For the bioassay, the compounds were resuspended in DMSO at a concentration of 1 mg/mL (stock solution). These stock solutions were further diluted with appropriate volumes of 7H9GC broth to yield final concentrations of 0.1 to 50 μ g/mL. Final drug concentration ranges of standard antibiotics used as positive controls were 0.125 to 32 μ g/mL for isoniazid and 0.063 to 16 μ g/mL for rifampin. The standard drugs or compounds (100 μ L) were mixed in the wells with 100 μ L of bacterial inoculum, resulting in a final bacterial concentration of approximately 1.2 x 10⁶ CFU/mL. The wells in column 11 served as inoculum-

only controls. Solvent (DMSO) was included in every experiment as a negative control. The plates were sealed in plastic bags and then incubated at 37°C for 5 days. On day 5, 50 µL of the tetrazolium-Tween 80 mixture {1.5 mL of tetrazolium [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] (Aldrich Chemical Co., Milwaukee, WI) at a dilution of 1 mg/mL in absolute ethanol and 1.5 mL of 10% Tween 80} was added to the wells, and the plate was incubated at 37°C for 24 h. After this incubation period, the growth of the microorganism was visualized by the

change in color of the dye from yellow to purple. The tests were carried out in triplicate. MIC is defined as the lowest drug concentration that prevents the aforementioned change in color.

Acknowledgments - Collaborative work was performed under the auspices of the CYTED, project PIBATUB, ex-INETI and project Pest-OE/UI0612/2011. The authors are grateful to the Government of the Azores and Azorean DRRF for providing the *C. japonica* and the facilities for this study.

References

- Mendes JFP. (2008) A Estratégia para a Floresta na Região Autónoma dos Açores, DRRF. In *Seminário de Certificação Florestal na Região Autónoma dos Açores*, Ponta Delgada, 13-14 Março.
- Ishiguri F, Saitoh K, Andoh M, Abe Z, Yokota S, Yoshizawa N. (2000) Improvement of heartwood color of black-colored sugi (*Cryptomeria japonica* D. Don) by UV irradiation after smoke heating. *Holzforschung*, **54**, 294-300.
- Matsunaga H, Shiotari R, Matsumura J, Oda K, Utsumi Y, Yamamoto K. (2006) Potassium distribution in black heartwood of sugi (*Cryptomeria japonica*) I: localization in axial parenchyma cells. *Journal of Wood Science*, **52**, 95-100.
- Belierique J, Nóbrega C, Quintela M, Faria C, Almeida MH. (2012) In *VI Jornadas Florestais Insulares*, 21-24 May, Faial, Pico and São Jorge.
- Matsunaga T, Hasegawa C, Kawasuji T, Suzuki H, Saito H, Sagioka T, Takahashi R, Tsukamoto H, Morikawa T, Akiyama T. (2000) Isolation of the antitumor compound in essential oil from the leaves of *Cryptomeria japonica*. *Biological & Pharmaceutical Bulletin*, **23**, 595-598.
- Cheng SS, Lin HY, Chang ST. (2005) Chemical composition and antifungal activity of essential oils from different tissues of Japanese cedar (*Cryptomeria japonica*). *Journal of Agricultural and Food Chemistry*, **53**, 614-619.
- Takao Y, Kuriyama I, Yamada T, Mizoguchi H, Yoshida H, Mizushima Y. (2012) Antifungal properties of Japanese cedar essential oil from waste wood chips made from used sake barrels. *Molecular Medicine Reports*, **5**, 1163-1168.
- Cha JD, Jeong MR, Jeong SI, Moon SE, Kil BS, Yun SI, Lee KY, Song YH. (2007) Chemical composition and antimicrobial activity of the essential oil of *Cryptomeria japonica*. *Phytotherapy Research*, **21**, 295-299.
- Chang C, Chang W, Chang S, Cheng S. (2008) Antibacterial activities of plant essential oils against *Legionella pneumophila*. *Water Research*, **42**, 278-286.
- Geong-Ho L, Lee BK, Kim JH, Lee SH, Hong SK. (2009) Comparison of chemical compositions and antimicrobial activities of essential oils from three conifer trees; *Pinus densiflora*, *Cryptomeria japonica*, and *Chamaecyparis obtusa*. *Journal of Microbiology Biotechnology*, **19**, 391-396.
- Cha JD, Kim JY. (2012) Essential oil from *Cryptomeria japonica* induces apoptosis in human oral epidermoid carcinoma cells via mitochondrial stress and activation of caspases. *Molecules*, **17**, 3890-3901.
- Cheng WW, Lin CT, Chu FH, Chang ST, Wang SY. (2009) Neuropharmacological activities of phytoncide released from *Cryptomeria japonica*. *Journal of Wood Science*, **55**, 27-31.
- Cheng SS, Chang S, Chang S, Tsai K, Chen W. (2003) Bioactivity of selected plant essential oil against the yellow fever mosquito *Aedes aegypti* larvae. *Bioresource Technology*, **89**, 99-102.
- Cheng SS, Chua MT, Chang EH, Huang CG, Chen WJ, Chang ST. (2008) Larvicidal activity of tectoquinone isolated from red heartwood-type *Cryptomeria japonica* against two mosquito species. *Bioresource Technology*, **99**, 3617-3622.
- Gu HJ, Cheng SS, Lin CY, Huang CG, Chen WJ, Chang ST. (2009) Repellency of essential oils of *Cryptomeria japonica* (Pinaceae) against adults of the mosquitoes *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae). *Journal of Agricultural and Food Chemistry*, **57**, 11127-11133.
- Cheng SS, Chang ST. (2002) Antitermitic activity of essential oils from *Cryptomeria japonica*. *Quarterly Journal of Chinese Forestry*, **35**, 193-199.
- Cheng SS, Lin CY, Chung MJ, Chang ST. (2012) Chemical composition and antitermitic activity against *Coptotermes formosanus* Shiraki of *Cryptomeria japonica* leaf essential oil. *Chemistry & Biodiversity*, **9**, 352-358.
- Cheng SS, Chua MT, Chang EH, Huang CG, Chen WJ, Chang ST. (2009) Variations in insecticidal activity and chemical compositions of leaf essential oils from *Cryptomeria japonica* at different ages. *Bioresource Technology*, **100**, 465-470.
- Wang S, Lai W, Chu F, Lin C, Shen S, Chang ST. (2006) Essential oil from the leaves of *Cryptomeria japonica* acts as a silverfish (*Lepisma saccharina*) repellent and insecticide. *Journal of Wood Science*, **52**, 522-526.
- Xie Y, Wang J, Yang F, Lei C. (2011) Comparative analysis of essential oil components of two *Cryptomeria* species from China. *Industrial Crops and Products*, **34**, 1226-12230.
- European Pharmacopoeia by Council of Europe (2010) European Pharmacopoeia Commission; European Directorate for the Quality of Medicines & Healthcare. 7th Edition. Strasbourg.
- Kenmoku H, Tanaka M, Ogiyama K, Kato N, Sassa T. (2004) Identification of (+)-phylloladene, (-)-sandaracopimaradiene, and (+)-kaurene as new fungal metabolites from fusicoocin-producing *Phomopsis amygdali* F6. *Bioscience, Biotechnology, and Biochemistry*, **68**, 1574-1577.
- Yasue M, Ogiyama K, Saito M. (1976) The diterpene hydrocarbons in the leaves of *Cryptomeria japonica*. *Journal of the Japanese Forest Society*, **58**, 285-290.
- Hanson JR, Siversn M, Piozzi F, Savona G. (1976) The ¹³C nuclear magnetic resonance spectra of kauranoid diterpenes. *Journal of the Chemical Society, Perkin Transactions*, **1**, 114-117.
- Raharivelomanana P, Bianchini PJ, Cambon A, Azzaro M, Faure R. (1995) Two-dimensional NMR of sesquiterpenes. 8-Complete assignment of ¹H and ¹³C NMR spectra of seven sesquiterpene alcohols from *Neocallitropsis pancheri*. *Magnetic Resonance in Chemistry*, **33**, 233-235.
- Shibutani S, Takata K, Doi S. (2007) Quantitative comparisons of antitermite extractives in heartwood from the same clones of *Cryptomeria japonica* planted at two different sites. *Journal of Wood Science*, **53**, 285-290.
- Cornwell CP, Reddy N, Leach DN, Wyllie S. (2000) Origin of (+)-δ-cadinene and the cubenols in the essential oils of the Myrtaceae. *Flavour and Fragrance Journal*, **15**, 352-361.
- CLSI, Clinical and Laboratory Standards Institute (formerly NCCLS, National Committee for Clinical and Laboratory Standards) (2008). Method M27-A3, 3rd ed.; method M38-A2, 2nd ed.; Wayne PA. (Ed.). NCCLS, Pennsylvania, Vol. **28** (14), pp. 1-25 and Vol. **28** (16), pp. 1-35.
- Muroi H, Kubo I. (1996). Antibacterial activity of anacardic acids and totarol alone and in combination with methicillin, against methicillin-resistant *Staphylococcus aureus*. *Journal of Applied Bacteriology*, **80**, 387-394.
- Kouznetsov V, Meléndez Gómez C, Derita M, Svetaz L, del Olmo E, Zacchino S. (2012). Synthesis and antifungal activity of diverse C-2 pyridinyl and pyridinylvinyl substituted quinolines. *Bioorganic and Medicinal Chemistry*, **20**, 6506-6512.

Antioxidant, Antiproliferative and Antimicrobial Activities of the Volatile Oil from the Wild Pepper *Piper capense* Used in Cameroon as a Culinary Spice

Verlaine Woguem^{a,b}, Filippo Maggi^{c,*}, Hervet P. D. Fogang^{a,b}, Léon A. Tapondjou^a, Hilaire M. Womeni^b, Luana Quassinti^c, Massimo Bramucci^c, Luca A. Vitali^c, Dezemona Petrelli^d, Giulio Lupidi^e, Fabrizio Papa^c, Sauro Vittori^c and Luciano Barboni^c

^aLaboratory of Environmental and Applied Chemistry, Faculty of Science, University of Dschang, P.O. Box 183, Dschang, Cameroon

^bLaboratory of Biochemistry of Medicinal Plants, Food Science and Nutrition, Faculty of Science, University of Dschang, P.O. Box 67, Dschang, Cameroon

^cSchool of Pharmacy, University of Camerino, I-62032 Camerino, Italy

^dSchool of Biosciences and Biotechnology, University of Camerino, I-62032 Camerino, Italy

^eSchool of Science and Technology, Chemistry Division, University of Camerino, I-62032 Camerino, Italy

filippo.maggi@unicam.it

Received: August 25th, 2013; Accepted: October 1st, 2013

Wild pepper (*Piper capense* L.f., Piperaceae) is a spice traditionally used in western Cameroon to make soups called 'Nkui' and 'Nah poh'. In the present work, the essential oil hydrodistilled from fruits was analyzed by GC-FID and GC-MS, and for *in vitro* biological activities, namely cytotoxic, antioxidant and antimicrobial, by MTT, DPPH, ABTS and agar disc diffusion methods. The oil composition was dominated by monoterpene hydrocarbons (56.5%) responsible for the pepper odor, such as β -pinene (33.2%), sabinene (10.0%) and α -pinene (8.9%). The oil induced a concentration-dependent inhibitory effect on human tumor cells MDA-MB 231 (breast adenocarcinoma), A375 (malignant melanoma) and HCT116 (colon carcinoma), showing IC₅₀ values of 26.3, 76.0 and 22.7 μ g/ml, respectively. The oil showed total antioxidant activity with a Trolox equivalent antioxidant concentration (TEAC) value of 140 μ mol/g. The essential oil of *P. capense* proved to be an effective scavenger of the ABTS⁺ radical, with an activity only about 30 times lower than that of Trolox. Moderate activity was observed against the Gram-positive species *Staphylococcus aureus* and *Enterococcus faecalis*, and the yeast *Candida albicans*. The notable inhibition of some human tumor cells is worthy of further investigation to discover the possible mechanisms of action responsible for the observed cytotoxic effect of this essential oil.

Keywords: *Piper capense*, Essential oil, Cytotoxic activity, Antioxidant, Antimicrobial.

Species belonging to the genus *Piper* L. (Piperaceae) are distributed largely in tropical and sub-tropical regions of the world. They have high commercial, economic and medicinal value since they constitute an important source of secondary metabolites. Notably, they produce aromatic fruits which are mainly used as spices. Especially in Cameroon, two traditional soups made by people in the western region, called 'Nkui' and 'Nah poh', contain the fruits of wild pepper (*P. capense* L.f.), which is one of three *Piper* species recognized to be indigenous to Cameroon [1]. The plant is an under storey shrub common in gallery forest, with V-shaped branchless and pointed ovate leaves bearing seven basal nerves. The fruits ripen around January and March [1]. They are believed to be the cause of sweating and sleepiness [2]. The leaves are used in S. Tomé and Príncipe, on the western equatorial coast of Central Africa, as a stomachic and carminative in cases of indigestion, flatulence and colic. Furthermore, a handful of leaves is included in Cameroonian recipes to treat epilepsy in Fongo-Tongo village [3]. The roots are traditionally used in South Africa to treat wounds, vaginal discharge, infertility, sore throat and tongue sores. The same parts in Kenya are boiled and used against malaria [4].

Main secondary metabolites isolated from the plant are lignans [5,6], terpenoids [2,7], amides [8], amide alkaloids and flavones [9]. As far as we know, few investigations have been made on the biological activity of volatiles obtained from this spice. The essential oils obtained from leaves and fruits collected in the

western province of Cameroon were reported to show negligible antifungal activity [10]. Other papers described the toxicity of the essential oil against the insect infecting maize grains, *Sitophilus zeamais* [11], and the larvicidal activity against the malaria vector *Anopheles gambiae* [12].

As part of our continuous search for bioactivity of essential oils from Cameroonian spices [13,14], we evaluated the antioxidant, cytotoxic, and antimicrobial capacities of *P. capense* essential oil. Chemical investigation has been undertaken to establish the composition-activity relationship.

Sixty-two components were identified in the oil (Table 1), corresponding to 89.5% of the total composition. The oil was mainly constituted by monoterpenes (64.7%), among which the hydrocarbons were predominant (56.5%). Main representatives of this chemical class were β -pinene (33.2%), sabinene (10.0%) and α -pinene (8.9%), accounting together for 52.1% of the whole oil. This high concentration of hydrocarbons is essential to give the pepper odor [15]. The second fraction in terms of abundance was represented by sesquiterpenes (24.1%), among which hydrocarbons were again the most abundant (17.8%). Main representatives of this group were (*E*)-caryophyllene (6.3%) and germacrene D (3.8%). The chemical composition of the oil obtained from dry fruits was quite consistent with that reported in previous studies [10,11].

Table 1: Chemical composition of the essential oil hydrodistilled from fruits of *Piper capense*.

N.	Constituent ^a	RF ^b	Calc. LRI ^c	Lit. LRI ^d		Essential oil		ID ^f
				ADAMS	NIST08	% ^e	mg/g	
1	α -Thujene	1.1	927	930	927	0.6±0.0	14.5±0.3	RI,MS
2	α -Pinene	1.1	932	939	932	8.9±0.0	234.8±4.1	Std
3	Camphene	1.1	947	954	947	0.6±0.0	16.2±0.4	Std
4	Sabinene	1.1	973	975	973	10.0±0.8	264.9±24.5	RI,MS
5	β -pinene	1.1	975	979	965	33.2±0.7	876.9±5.9	Std
6	Myrcene	1.1	993	990	993	0.9±0.0	23.0±0.5	Std
7	α -Phellandrene	1.1	1005	1002	1005	Tr±0.0	0.7±0.0	RI,MS
8	δ -3-Carene	1.1	1011	1011	1011	0.5±0.0	12.3±0.3	Std
9	α -Terpinene	1.1	1019	1017	1019	Tr±0.0	0.5±0.1	RI,MS
10	<i>p</i> -Cymene	1.1	1028	1024	1028	0.4±0.0	11.4±0.3	Std
11	Limonene	1.1	1031	1029	1033	1.8±0.0	48.1±0.9	Std
12	1,8-Cineole	1.5	1034	1031	1034	0.4±0.0	10.2±0.7	Std
13	(<i>Z</i>)- β -Ocimene	1.1	1045	1034	1043	Tr±0.0	0.8±0.0	RI,MS
14	(<i>E</i>)- β -Ocimene	1.1	1055	1050	1052	Tr±0.0	0.9±0.0	RI,MS
15	γ -Terpinene	1.1	1063	1059	1063	Tr±0.0	1.1±0.2	Std
16	<i>cis</i> -Sabinene hydrate	1.5	1071	1070	1068	0.4±0.0	11.8±0.3	RI,MS
17	Terpinolene	1.1	1089	1088	1089	Tr±0.0	0.9±0.1	Std
18	<i>trans</i> -Sabinene hydrate	1.5	1098	1098	1098	Tr±0.0	0.8±0.0	RI,MS
19	Linalool	1.5	1103	1096	1103	1.3±0.0	34.2±0.9	Std
20	<i>cis</i> - <i>p</i> -Menth-2-en-1-ol	1.5	1125	1121	1126	0.1±0.0	3.1±0.5	RI,MS
21	<i>trans</i> -Pinocarveol	1.5	1140	1139	1141	0.6±0.0	15.7±0.7	Std
22	Camphor	1.5	1146	1146		0.2±0.0	4.5±0.8	Std
23	Pinocarvone	1.5	1165	1164	1165	0.1±0.0	3.2±0.3	RI,MS
24	Borneol	1.5	1167	1169	1167	0.6±0.0	14.9±0.5	Std
25	Terpinen-4-ol	1.5	1179	1177	1179	1.2±0.0	31.6±0.5	Std
26	α -Terpineol	1.5	1192	1188	1192	0.6±0.0	16.6±0.4	Std
27	Myrtenal	1.5	1195	1195	1195	0.3±0.0	8.3±0.3	Std
28	Myrtenol	1.5	1197	1195	1196	0.4±0.0	10.2±0.9	RI,MS
29	Isobornyl acetate	1.5	1286	1285		2.0±0.0	52.9±1.3	Std
30	α -Cubebene	1.1	1348	1348		0.4±0.1	10.3±2.1	RI,MS
31	Cyclosativene	1.1	1360	1371	1360	0.2±0.0	6.0±0.1	RI,MS
32	α -Copaene	1.1	1372	1376		0.6±0.0	16.0±0.2	RI,MS
33	β -Bourbonene	1.1	1380	1388		0.1±0.0	1.6±0.2	RI,MS
34	β -Cubebene	1.1	1387	1388	1387	1.2±0.0	32.7±0.3	RI,MS
35	β -Elemene	1.1	1389	1390		1.0±0.0	26.0±0.2	RI,MS
36	(<i>E</i>)-Caryophyllene	1.1	1413	1419		6.3±0.1	166.8±1.6	Std
37	β -Copaene	1.1	1424	1432		0.1±0.0	3.9±0.3	RI,MS
38	γ -Elemene	1.1	1431	1436		0.3±0.0	6.7±0.3	RI,MS
39	6,9-Guaiadiene	1.1	1440	1444		0.2±0.0	5.4±0.2	RI,MS
40	α -Humulene	1.1	1448	1454	1448	1.1±0.0	29.4±0.5	Std
41	(<i>E</i>)- β -Farnesene	1.1	1459	1456	1459	0.1±0.0	1.5±0.3	RI,MS
42	Germacrene D	1.1	1476	1485		3.8±0.0	99.7±0.6	RI,MS
43	β -selinene	1.1	1480	1490	1480	0.3±0.0	7.4±0.1	RI,MS
44	<i>trans</i> -Muurolo-4(14),5-diene	1.1	1485	1493		0.1±0.0	3.2±0.0	RI,MS
45	α -Muurolole	1.1	1496	1500		0.6±0.0	16.1±0.0	RI,MS
46	<i>n</i> -Pentadecane	1.4	1500	1500	1500	0.2±0.0	5.9±0.1	Std
47	β -Bisabolene	1.1	1506	1505		Tr	0.7±0.0	RI,MS
48	<i>trans</i> -Calamenene	1.1	1519	1522		Tr±0.0	0.5±0.0	RI,MS
49	δ -Cadinene	1.1	1523	1523		0.8±0.0	21.8±0.5	RI,MS
50	<i>trans</i> -Cadina-1,4-diene	1.1	1528	1534		Tr±0.0	0.4±0.0	RI,MS
51	Germacrene B	1.1	1551	1561	1552	0.6±0.0	16.7±0.4	RI,MS
52	Elemicin	1.4	1561	1557	1561	Tr±0.0	0.8±0.0	RI,MS
53	(<i>E</i>)-Nerolidol	1.3	1564	1563	1564	1.5±0.0	40.3±0.6	RI,MS
54	Spathulenol	1.3	1573	1578		Tr±0.0	0.8±0.0	RI,MS
55	Caryophyllene oxide	1.3	1576	1583	1576	2.8±0.0	74.2±0.8	Std
56	Salvial-4(14)-en-1-one	1.3	1588	1594		0.2±0.0	5.1±0.5	RI,MS
57	Humulene epoxide II	1.3	1602	1608	1602	0.3±0.0	8.2±0.3	RI,MS
58	1,10-di- <i>epi</i> -Cubenol	1.3	1626	1619	1623	0.4±0.0	11.6±0.6	RI,MS
59	Caryophylla-4(12),8(13)-dien-5-ol ^h	1.3	1632	1640	1635	0.1±0.1	2.9±0.6	RI,MS
60	Cubenol	1.3	1638	1646		0.3±0.0	7.4±0.4	RI,MS
61	α -Muurolol	1.3	1644	1646	1644	0.4±0.0	10.0±0.3	RI,MS
62	Eudesma-4(15),7-dien-1- β -ol	1.3	1682	1688		0.3±0.0	9.2±1.9	RI,MS
	Total identified (%)					89.5		
	Oil yield (%)					1.9		
	Grouped compounds							
	Monoterpene hydrocarbons					56.5	1495.5	
	Oxygenated monoterpenes					8.2	165.1	
	Sesquiterpene hydrocarbons					17.8	474.1	
	Oxygenated sesquiterpenes					6.3	169.7	
	Others					0.7	71.5	

^a Compounds are listed in order of their elution from a HP-5MS column. Their nomenclature was in accordance with Adams [30]. ^b Response factor (RF) of FID detector for the main chemical groups occurring in the essential oil. ^c Linear retention index on HP-5MS column experimentally determined using homologous series of C₈-C₃₀ alkanes. ^d Linear retention index taken from Adams [30] and/or NIST08 [31]. ^e Percentage values are means of three determinations. ^f Identification methods: Std, based on comparison with authentic compounds; MS, based on comparison with Wiley, ADAMS and NIST08 MS database; RI, based on comparison of RI with those reported in ADAMS and NIST08. ^g Tr, traces (mean value below 0.1%). ^h Correct isomer not identified.

In these studies the major volatiles of fruits from western Cameroon were again monoterpene hydrocarbons, namely β -pinene, sabinene and α -pinene (46.8, 17.4 and 14.4%, and 59.3, 14.7 and 10.5%, respectively), and sesquiterpene hydrocarbons, namely (*E*)-caryophyllene and germacrene D (4.0 and 5.2%, and 3.4 and 2.5%, respectively), but percentages were slightly different with respect to those detected by us. The aerial parts of *P. capense* growing in S. Tomé and Príncipe [2] were also characterized by monoterpenes such as β -pinene (32.5%) and α -pinene (8.6%), and sesquiterpenes such as (*E*)-caryophyllene (12.6%), along with a low content of sabinene (0.7%), and benzenepropanoic acid ethyl ester (7.8%), which was not detected in our study. Finally, the essential oil obtained from the whole plant collected in Kenya [12] showed a different composition consisting mainly of sesquiterpene hydrocarbons (43.9%) such as δ -cadinene (16.8%), β -bisabolene (5.6%) and bicyclogermacrene (3.3%), while monoterpene hydrocarbons, including β -pinene (7.2%) and α -phellandrene (4.8%), were less abundant (30.6%).

The antioxidant activity of the essential oil from *P. capense* was evaluated using DPPH and ABTS methods. From results reported in Table 2 we observed that the oil showed lower antioxidant activity in the DPPH assay, while worthy of mention was the activity against the ABTS⁺ radical. On the other hand, no reducing capacity power (FRAP assay) of the oil was pointed out. Proton radical scavenging activity is an important attribute of antioxidants and ABTS⁺, a protonated radical, has a characteristic maximum absorbance at 734 nm, which decreases with the scavenging of the proton radicals [16]. As shown, the essential oil from *P. capense* was an effective scavenger of the ABTS⁺ radical with an activity about 30 times lower than that of Trolox and the scavenging activity was much higher than that against the DPPH[•] radical. Factors like radical stereoselectivity or sample solubility in different testing systems have been reported to affect the capacity of samples to react and quench different radicals [17]. Wang and co-workers [18] found that some compounds having ABTS⁺ scavenging activity did not show inhibition against the DPPH radical. Low values of DPPH scavenging activity were also reported for different kinds of essential oils with a typical monoterpene hydrocarbon pattern [19-21]. In addition, the discrepancy observed in the results of ABTS⁺ and DPPH[•] scavenging ability might be due to their different mechanisms of reaction: ABTS⁺ assay is based on hydrogen transfer reaction, while DPPH assay is based on electron transfer [22]. Evaluation of the scavenging activity of pure compounds used as standards, conducted in our laboratories, showed no activity for α -pinene, β -pinene, and (*E*)-caryophyllene, and very low activity for limonene. Therefore, the antioxidant effectiveness of the oil, as already demonstrated [19], may be due to the synergism of different constituents.

Table 2: *In vitro* antioxidant activity of the essential oil from *Piper capense*.

Essential oil	(IC ₅₀ µg/ml) ^a		
	ABTS	DPPH	FRAP
<i>P. capense</i>	300 (± 6)	>>1000	N.A. ^b
Reference			
Trolox	10.1(±0.6)	3.9 (± 0.4)	

^a IC₅₀ = The concentration of compound that affords a 50% reduction). ^b N.A., not active.

To investigate the cytotoxic activity of *P. capense* essential oil, we evaluated its effect on the human tumor cell lines MDA-MB 231 (human breast adenocarcinoma), A375 (human malignant melanoma) and HCT116 (human colon carcinoma) by MTT assay. As shown in Table 3, the essential oil induced a concentration-dependent inhibitory effect on all cancer cell lines tested in the dilution range 0.78-200 µg/mL.

Table 3: *In vitro* cytotoxic activity of the essential oil from *Piper capense*.

Essential oil	Cell line (IC ₅₀ µg/mL)		
	MDA-MB 231 ^a	A375 ^b	HCT116 ^c
<i>P. capense</i>	26.3	76.0	22.7
95% C.I.	23.2 - 29.9	65.7 - 83.6	20.7 - 25.0
Major constituents			
β -Pinene	78.5	> 200	59.2
95% C.I.	73.2 - 84.4		54.5 - 64.0
(<i>E</i>)-Caryophyllene	45.3	63.3	55.7
95% C.I.	37.3 - 52.7	54.8 - 77.8	50.3 - 62.7
Reference			
Cisplatin	2.4	0.16	2.8
95% C.I.	2.0 - 2.7	0.12 - 0.22	2.6 - 3.0

IC₅₀ = Concentration of compound that affords a 50% reduction in cell growth (after 72 h of incubation). ^a Human breast adenocarcinoma cell line. ^b Human malignant melanoma cell line. ^c Human colon carcinoma cell line.

The IC₅₀ values of the oil were 26.3, 76.0 and 22.7 µg/mL on MDA-MB 231, A375 and HCT116 cell lines, respectively. The cytotoxic activity of *P. capense* essential oil may be attributed to specific components of the oil. We tested the cytotoxic activity of β -pinene and (*E*)-caryophyllene on the same cell lines used to analyze *P. capense* essential oil (Table 3). In a previous study, β -pinene proved to be less active than α -pinene against the same cell lines [13]. According to Bakkali and coworkers [23], sabinene has also a dose-dependent cytotoxic activity when tested either alone or added to the essential oil. The concentrations of β -pinene (33.2%), sabinene (10%) and α -pinene (8.9%) cannot fully justify the cytotoxic activity of *P. capense* essential oil, which means that some other compounds are active, probably sesquiterpenes such as (*E*)-caryophyllene (6.3%) and germacrene D (3.8%). (*E*)-caryophyllene showed cytotoxic activity against the tumor cell lines (Table 3), with IC₅₀ values in the range 45.3-63.3 µg/mL. To our knowledge, also germacrene D was active against tumor cells such as human breast adenocarcinoma (MDA-MB 231 and MCF-7), human ductal carcinoma (Hs 578T), and human hepatocellular carcinoma (Hep G2) [24]. In addition, minor components could also contribute to the cytotoxic activity of the oil or be responsible for synergism along with other compounds [25,26].

Table 4: Activity of *P. capense* essential oil and some major constituents on bacterial and yeast species. Values are expressed as inhibition zones (mm).^a

Essential oil	<i>S. aureus</i>	<i>E. faecalis</i>	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>C. albicans</i>
<i>P. capense</i>	10	10	6	6	8
Major constituents					
α -pinene	6	6	6	6	11
β -pinene	6	6	6	6	9
(<i>E</i>)-caryophyllene	8	10	8	6	6

^a Sterile filter paper discs (6 mm in diameter) were placed on the surface of inoculated plates and spotted with 10 µL of a 1:1 dilution of essential oil/reference compound in DMSO.

Results of antimicrobial activity tests are summarized in Table 4. The essential oil showed a measurable, albeit low, activity against the two Gram-positive bacterial species *S. aureus* and *E. faecalis* and the yeast *C. albicans*. Activity on Staphylococci has already been reported for the methanolic extract from barks [27], while no data were available for the antibacterial activity of the essential oil from barks of this species. The moderate activity against *C. albicans* has been previously observed, but was measured by a microdilution method [10]. Given the differences in the methods, the results may be considered comparable and the activity presented herein as a sort of general positive control of the antimicrobial activity test. A remarkably strong activity was observed in a previous study using an acetone extract from roots, whose composition was, however, not determined [28]. Comparison between activities of the essential oil and three major constituents of the oil, namely α -pinene, β -pinene, and (*E*)-caryophyllene, showed

that the last may be responsible for the activity observed against all the bacterial species, except for *P. aeruginosa*, against which neither the oil nor the pure compounds were effective. In the *E. coli* culture we observed an inhibition by (*E*)-caryophyllene (zone diameter of 8 mm), not accompanied by an appreciable activity of the whole essential oil. This result may be due to the relative low concentration of (*E*)-caryophyllene within the oil. On the contrary, *C. albicans* was only susceptible to both forms of pinene. This may account for the measurable antifungal activity of the essential oil.

In conclusion, the high content of monoterpene hydrocarbons found in *P. capense* essential oil furnishes a typical pepper note and supports the use of the plant as an odorous spice in African traditional cuisine. The oil was an effective scavenger of the ABTS⁺ radical with an activity only about 30 times lower than that of Trolox and so could be considered as a good food preservative. The essential oil showed notable inhibition activity against some human tumor cells, being worthy of further investigation to discover the possible mechanisms of action responsible for the observed cytotoxic effect. Finally, the results of antimicrobial activity against the bacterial Gram-positive and yeast species may sufficiently support the described use of some parts of this plant in the treatment of some infection related conditions (e.g. wounds, vaginal discharge, and sore throat).

Experimental

Plant material: Dry fruits of *P. capense* were bought in a market of Dschang (Cameroon, Menoua Division, Western Region). Identification was made by Mr Nana Victor, taxonomist at the Cameroon National Herbarium (Yaoundé), where a voucher specimen was deposited (N. 6018/HNC/SRF).

Extraction of the essential oil: The dry fruits (80 g) of *P. capense* were ground and subjected to hydrodistillation in a Clevenger-type apparatus using 750 mL of deionized water for 3 h yielding 1.98%, w/w, of a strong smelling, pale yellow oil. The oil collected was dried over anhydrous sodium sulfate, and then stored in the refrigerator (+4°C) until used.

GC-FID and GC-MS analyses: For gas chromatographic separations, an Agilent 4890D instrument coupled to an ionization flame detector (FID) was used. Volatile components were separated on a HP-5 capillary column (5% phenylmethylpolysiloxane, 25 m, 0.32 mm i.d.; 0.17 µm film thickness) (J and W Scientific, Folsom, CA), with the following temperature program: 5 min at 60°C, subsequently 4°C/min up to 220°C, then 11°C/min up to 280°C, held for 15 min, for a total run of 65 min. Injector and detector temperatures were 280°C. Helium was used as the carrier gas, at a flow rate of 1.4 mL/min; injection volume: 1 µL; split ratio, 1:34. A mixture of aliphatic hydrocarbons (C₈-C₃₀) (Sigma, Milan, Italy) in *n*-hexane was directly injected into the GC injector under the above temperature program, in order to calculate the retention index of each compound. Oil samples were diluted 1:100 in *n*-hexane and injected in a volume of 1 µL. Analysis was repeated 3 times. Data were collected by using HP3398A GC Chemstation software (Hewlett Packard, Rev. A.01.01). The relative amounts of essential oil components, expressed as percentages and mg/g e.o., were obtained according to a reported procedure [13].

GC-MS analysis was performed on an Agilent 6890N gas chromatograph coupled to a 5973N mass spectrometer using a HP-5MS (5% phenylmethylpolysiloxane, 30 m, 0.25 mm i.d., 0.1 µm film thickness) (J & W Scientific, Folsom) capillary column. The temperature programme was the same as above. Injector and detector temperatures were 280°C. Helium was used as carrier gas,

at a flow rate of 1 mL/min. Split ratio: 1:50; acquisition mass range: 29-400 *m/z*. All mass spectra were acquired in electron-impact (EI) mode with an ionization voltage of 70 eV. Oil samples were diluted 1:100 in *n*-hexane and the volume injected was 2 µL. Data were analyzed by using MSD ChemStation software (Agilent, Version G1701DA D.01.00). Whenever possible, volatile components were identified by co-injection with authentic standards and isolated compounds. Otherwise, the peak assignment was carried out by the interactive combination of chromatographic linear retention indices that were consistent with those reported in the literature [29,30], and MS data consisting of computer matching with WILEY275, NIST 08 ADAMS, and a home-made library (based on the analyses of reference oils and commercially available standards).

Evaluation of antioxidant activity

DPPH free-radical scavenging activity: The DPPH free-radical scavenging assay was carried out on a microplate according to a previously described procedure [31]. Extract concentration providing 50% inhibition (IC₅₀) was calculated from the linear regression algorithm of the graph plotting inhibition percentage against extract concentration. Experiments were conducted in triplicate.

ABTS assay: Radical scavenging capacity was measured by a modified method originally described by Re and coworkers [32] for application to a 96-well microplate assay [33]. Trolox was used as positive control. Each experiment was repeated 3 times.

Ferric reducing antioxidant power (FRAP) assay: Determination of antioxidant activity by ferric reducing antioxidant power (FRAP) assay was carried out according to Müller and coworkers [34]. The ability of the essential oil to scavenge the different radicals in the assays was compared with that of Trolox, used as positive control, and the activity of the oil was expressed as tocopherol equivalent antioxidant capacity (µmol TE/g product). Tests were repeated 3 times.

MTT cytotoxicity assay: Human colon carcinoma cell line HCT116 was cultured in RPMI1640 medium with 2 mM L-glutamine, 100 IU/mL penicillin, 100 µg/mL streptomycin, and supplemented with 10% heat-inactivated fetal bovine serum (HI-FBS) (PAA Laboratories GmbH, Austria). Human breast adenocarcinoma cell line MDA-MB 231, and human malignant melanoma cell line A375 were cultured in Dulbecco's Modified Eagle's Medium (DMEM) with 2 mM L-glutamine, 100 IU/mL penicillin, 100 µg/mL streptomycin, and supplemented with 10% HI-FBS. Cells were cultured in a humidified atmosphere at 37°C in the presence of 5% CO₂. The MTT assay was used as a relative measure of cell viability. Cell-viability assays were carried out according to a previously reported procedure [35]. Cytotoxicity was expressed as the concentration of compound inhibiting cell growth by 50% (IC₅₀). The IC₅₀ values were determined with the GraphPad Prism 4 computer program (GraphPad Software, S. Diego, CA, USA).

Antimicrobial activity: Microorganisms included in this study were: *Staphylococcus aureus* ATCC 25923, *Escherichia coli* ATCC 25922, *Pseudomonas aeruginosa* ATCC 27853, *Enterococcus faecalis* ATCC 29212, and *Candida albicans* ATCC 24433. Antimicrobial activity of the essential oil of *P. capense* was assessed by the disc diffusion test, as reported by the European Committee for Antimicrobial Susceptibility testing [36], with previously described minor modifications due to the nature of the substance tested [37]. Ten µL of each reference compound (α -pinene, β -pinene, (*E*)-caryophyllene) per paper disc was used in the control experiments; the known antimicrobials ciprofloxacin (5 µg

disc) and fluconazole (25 µg disc) were used as a reference against bacteria and fungi, respectively. Each test was repeated at least twice.

Acknowledgments – The authors are grateful to the Italian Ministry of Education (MIUR) for the financial support (COOPERLINK 2011 Prot. CII113PPUC).

References

- [1] Hutchinson J, Dalziel JM (1963) *Flora of West Tropical Africa*. 2nd ed., Vol. 2. Crown Agents, London.
- [2] Martins AP, Salgueiro L, Vila R, Tomi F, Cañigüeral S, Casanova J, Da Cunha AP, Adzet T. (1998) Essential oils from four *Piper* species. *Phytochemistry*, **49**, 2019-2023.
- [3] Noumi E, Fozi FL. (2003) Ethnomedical botany of epilepsy treatment in Fongo-Tongo village, western Province, Cameroon. *Pharmaceutical Biology*, **41**, 330-339.
- [4] Koch A, Tamez P, Pezzuto J, Soejarto D. (2005) Evaluation of plants used for antimalarial treatment by the Maasai of Kenya. *Journal of Ethnopharmacology*, **101**, 95-99.
- [5] Green TP, Wiemer DF. (1991) Four neolignan ketones from *Piper capense*. *Phytochemistry* **30**, 3759-3762.
- [6] Green TP, Galinis DL, Wiemer DF. (1991) Three neolignans from the roots of *Piper capense*. *Phytochemistry*, **30**, 1649-1652.
- [7] Chen TB, Green TP, Wiemer DF. (1992) Capentin: a novel sesquiterpene from the roots of *Piper capense*. *Tetrahedron Letters*, **33**, 5673-5676.
- [8] Pedersen ME, Metzler B, Stafford GI, van Staden J, Jäger AK, Rasmussen HB. (2009) Amides from *Piper capense* with CNS activity – A preliminary SAR analysis. *Molecules*, **14**, 3833-3843.
- [9] Kaou AM, Mahiou-Leddé V, Canlet C, Debrauwer L, Hutter S, Azas N, Ollivier E. (2010) New amide alkaloid from the aerial part of *Piper capense* L.f. (Piperaceae). *Fitoterapia*, **81**, 632-635.
- [10] Amvam Zollo PH, Biyiti L, Tchoumboungang F, Menut C, Lamaty G, Bouchet Ph. (1998) Aromatic plants of tropical central Africa. Part XXXII. Chemical composition and antifungal activity of thirteen essential oils from aromatic plants of Cameroon. *Flavour and Fragrance Journal*, **13**, 107-114.
- [11] Tchoumboungang F, Jazet DPM, Sameza ML, Fombotioh N, Wouatsa NAV, Amvam ZPH, Menut C. (2009) Comparative essential oils composition and insecticidal effect of different tissues of *Piper capense* L., *Piper guinense* Schum. et Thonn., *Piper nigrum* L., and *Piper umbellatum* L., grown in Cameroon. *African Journal of Biotechnology*, **8**, 424-431.
- [12] Matasyoh JC, Wathuta EM, Kariuki ST, Chepkorir R. (2011) Chemical composition and larvicidal activity of *Piper capense* essential oil against the malaria vector, *Anopheles gambiae*. *Journal of Asia-Pacific Entomology*, **14**, 26-28.
- [13] Fogang HPD, Taponjdjou LA, Womeni HM, Quassinti L, Bramucci M, Vitali LA, Petrelli D, Lupidi G, Maggi F, Papa F, Vittori S, Barboni L. (2012) Characterization and biological activity of essential oils from fruits of *Zanthoxylum xanthoxyloides* Lam. and *Zanthoxylum lepreurii* Guill. and Perr., two culinary plants from Cameroon. *Flavour and Fragrance Journal*, **27**, 271-279.
- [14] Fogang HPD, Womeni HM, Piombo G, Barouh N, Taponjdjou LA. (2012) Bioefficacy of essential and vegetable oils of *Zanthoxylum xanthoxyloides* seeds against *Acanthoscelides obtectus* (Say) (Coleoptera: Bruchidae). *Journal of Food Protection*, **75**, 547-555.
- [15] Jirovetz L, Buchbauer G, Ngassoum M. (1997) Investigation of the essential oils from the dried fruits of *Xylopi aethiopica* (West African "Peppertree") and *Xylopi a parviflora* from Cameroun. *Ernährung*, **21**, 324-325.
- [16] Mathew S, Abraham ET. (2006) *In vitro* antioxidant activity and scavenging effects of *Cinnamomum verum* leaf extract assayed by different methodologies. *Food and Chemical Toxicology*, **44**, 198-206.
- [17] Yu L, Haley S, Perret J, Harris M, Wilson J, Qian M. (2002) Free radical scavenging properties of wheat extracts. *Journal of Agricultural and Food Chemistry*, **50**, 1619-1624.
- [18] Wang M, Li J, Rangarajan M, Shao Y, La Voie EJ, Huang T, Ho C. (1998) Antioxidative phenolic compounds from sage (*Salvia officinalis*). *Journal of Agricultural and Food Chemistry*, **46**, 4869-4873.
- [19] Ruberto G, Baratta MT. (2000) Antioxidant activity of selected essential oil components in two lipid model systems. *Food Chemistry*, **69**, 167-174.
- [20] Sacchetti G, Maietti S, Muzzoli M, Scaglianti M, Manfredini S, Radice M, Bruni R. (2005) Comparative evaluation of 11 essential oils of different origin as functional antioxidants, antiradicals and antimicrobials in foods. *Food Chemistry*, **91**, 621-632.
- [21] Pednekar PP, Vakil BV, Sane RT, Datar AG. (2005) Phytochemical profile and antioxidant activity of the essential oil from *Blumea eriantha* DC. *International Journal of Pharmacy and Pharmaceutical Sciences*, **5**, 404-413.
- [22] Prior RL, Wu X, Schaich K. (2005) Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *Journal of Agricultural and Food Chemistry*, **53**, 4290-4302.
- [23] Bakkali F, Averbeck S, Averbeck D, Idaomar M. (2008) Biological effects of essential oils-a review. *Food and Chemical Toxicology*, **46**, 446-475.
- [24] Setzer WN, Schmidt JM, Noletto JA, Vogler B. (2006) Leaf oil compositions and bioactivities of Abaco bush medicine. *Pharmacologyonline*, **3**, 794-802.
- [25] Lattaoui N, Tantaoui-Elaraki A. (1994) Individual and combined oils. *Rivista Italiana EPPOS*, **13**, 13-19.
- [26] Yu JQ, Lei JQ, Yu HD, Cai X, Zou GL. (2004) Chemical composition and antimicrobial activity of essential oil of *Scutellaria barbata*. *Phytochemistry*, **65**, 881-884.
- [27] Steenkamp V, Fernandes AC, van Rensburg CEJ. (2007) Antibacterial activity of Venda medicinal plants. *Fitoterapia*, **78**, 561-564.
- [28] Samie A, Tambani T, Harshfield E, Green E, Ramalivhana E, Bessong PO. (2010) Antifungal activities of selected Venda medicinal plants against *Candida albicans*, *Candida krusei* and *Cryptococcus neoformans* isolated from South African AIDS patients. *African Journal of Biotechnology*, **9**, 2965-2976.
- [29] Adams RP. (2007) *Identification of essential oil components by gas chromatography/mass spectrometry*. Allured Publishing Corporation, Carol Stream, Illinois.
- [30] National Institute of Standards and Technology (2008) *NIST 08, Mass spectral library (NIST/EPA/NIH)*, Gaithersburg, USA.
- [31] Srinivasan R, Chandrasekar MJN, Nanjan MJ, Suresh B. (2007) Antioxidant activity of *Caesalpinia digyna* root. *Journal of Ethnopharmacology*, **113**, 284-291.
- [32] Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. (1999) Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology & Medicine*, **26**, 1231-1237.
- [33] Esparza-Rivera JR, Stone MB, Stuchnoff C, Pilon-Smiths E, Kendall PA. (2006) Effects of ascorbic acid applied by two hydrocooling methods on physical and chemical properties of green leaf stored at 5°C. *Journal of Food Science*, **71**, 270-276.
- [34] Müller L, Fröhlich K, Böhm V. (2011) Comparative antioxidant activities of carotenoids measured by ferric reducing antioxidant power (FRAP), ABTS bleaching assay (αTEAC), DPPH assay and peroxyl radical scavenging assay. *Food Chemistry*, **129**, 139-148.
- [35] Quassinti L, Lupidi G, Maggi F, Sagratini G, Papa F, Vittori S, Bianco A, Bramucci M. (2013) Antioxidant and antiproliferative activity of *Hypericum hircinum* L. subsp. *majus* (Aiton) N. Robson essential oil. *Natural Product Research*, **27**, 862-868.
- [36] Antimicrobial susceptibility testing EUCAST disk diffusion method, Version 2.1, (2012).

- [37] Quassinti L, Bramucci M, Lupidi G, Barboni L, Ricciutelli M, Sagratini G, Papa F, Caprioli G, Petrelli D, Vitali LA, Vittori S, Maggi F. (2013) *In vitro* biological activity of essential oils and isolated furanosesquiterpenes from the neglected vegetable *Smyrniololus atrum* L. (Apiaceae). *Food Chemistry*, **138**, 808-813.

Boldine and Related Aporphines: From Antioxidant to Antiproliferative Properties

Darina Muthna*, Jana Cmielova, Pavel Tomsik and Martina Rezacova

Department of Medical Biochemistry, Faculty of Medicine in Hradec Kralove, Charles University in Prague, Simkova 870, Hradec Kralove 500 38, Czech Republic

muthnad@lfhk.cuni.cz

Received: May 21st, 2013; Accepted: September 8th, 2013

Plant and folk medicine represent nowadays a source of either new therapeutic substances or substrates for drug synthesis. One such promising group for possible further exploitation is the family of aporphine alkaloids containing boldine and related compounds. In this mini-review we focus on boldine and its newly described effects, which predominantly arise from its antioxidant properties. Moreover, we try to compare its antiproliferative properties with other better known members of the aporphine group.

Keywords: Boldine, Aporphine alkaloids, Antioxidant, Antiproliferative properties.

A wide range of compounds of natural origin that have been exploited in folk medicine for a long time are currently being intensively studied. Because of their multiple benefits, these compounds represent a possible source of new therapeutic substances for the treatment of various diseases. One such promising group is the family of aporphine alkaloids including boldine, neolitsine, dicentrine, glaucine, cassythine, actinodaphnine, and nantenine [1-3]. Because of their chemical structures they are potent antioxidants. Oxidative stress results in direct or indirect reactive oxygen species (ROS) mediated damage of nucleic acids, proteins, and lipids, and has been implicated in carcinogenesis, neurodegeneration, atherosclerosis, diabetes and aging [4]. Thus the regulation of redox status remains a promising therapeutic approach [5]. In this review we focus on boldine and its newly described effects, which predominantly arise from its antioxidant properties. Moreover, we try to compare its described *in vitro* antiproliferative properties with other better known members of the aporphine group.




Figure 1: Boldine - (6a*S*)-1,10-Dimethoxy-6-methyl-5,6,6a,7-tetrahydro-4*H*-dibenzo[*de,g*]quinoline-2,9-diol

Antioxidant activities of boldine: Main sources of boldine are the leaves and bark of the Chilean tree *Peumus boldus* (common name, boldo) from the Monimiaceae family, but it can also be isolated from plants of the related Lauraceae family, among others [2,6,7]. Like other members of the aporphine group, boldine behaves as a potent antioxidant. Although boldine has various beneficial effects, the majority of them arise just from its antioxidant properties. Due to its free radical scavenging properties, boldine influences cell proliferation, survival, differentiation and metabolism [4]. Aporphines generally are easily oxidized to dehydro- and oxo-aporphines. Even in the absence of hydroxyl groups in non-phenolic aporphines, the antioxidant capacity is maintained by the stabilization of the benzylic C-6a radical with the nitrogen lone pair [8].

Generally, with respect to the phenolic structure, these phytochemicals can exert a biphasic effect. In cancer cells, they can scavenge the constitutively high amounts of H₂O₂ that stimulate proliferation. On the other hand, the same phytochemicals under certain experimental conditions can exhibit pro-oxidant activity and generate ROS in amounts beyond the threshold for its tolerance [9]. This is true also for boldine. At concentrations of 10 and 50 μM, it exhibits intense free radical scavenging properties; however, at a concentration of 100 μM, boldine potentiates lipoperoxidation induced by ischemia, indicating a pro-oxidant effect in rat hippocampal slices [10]. Similar concentration-dependent effects can be observed on regulators of adipogenesis and adiponectin levels. Boldine protects adiponectin from both oxidative stress and inflammatory cytokines at lower concentrations (10–25 μM), giving better protection than higher concentrations (more than 50 μM) [11]. In diabetic conditions, boldine restores NO bioavailability and thus attenuates oxidative stress and endothelial dysfunction [12]. Another property of boldine and related alkaloids which seems to be closely related to its free radical scavenging effects is its antinociceptive activity. Boldine inhibited the rapid pain response and exerted a central antinociceptive effect [2]. Moreover, boldine as an antioxidant, strongly inhibits the synthesis of prostaglandin, which results in its anti-inflammatory and antipyretic effects [13].

Enzyme inhibitory activities of boldine: Besides exhibiting significant antioxidant properties, boldine exerts inhibitory activity on some enzymes. Boldine inhibits acetylcholinesterase activity with an IC₅₀ value of 8.5±1.4 μM. For comparison, the IC₅₀ value of the standard acetylcholinesterase substrate inhibitor, huperzine A, is 1.8 μM [6]. The inhibition of acetylcholinesterase and the consequent increase in colon motility are the reasons why the infusion of *P. boldus* is used in folk medicine to treat digestive and hepatic problems [14]. However, boldine itself possess an ileum relaxant property [15]. In addition, boldine exhibits other properties with respect to which a connection with antioxidant capacity has not yet been proven. Boldine might be able to interact with the promoter of the peroxisome proliferator-activated receptor (PPAR) gene directly and/or modulate its responsive genes and thus have strong beneficial effects on obesity-related diseases. This effect seems to be independent of its antioxidant property [11].

Hoet *et al.* [16] focused on boldine as well as other aporphine alkaloids within the context of DNA. These alkaloids are non-planar molecules and thus lack the structure of conventional DNA intercalators. Nevertheless, boldine intercalates weakly into DNA and may also be an adaptive intercalator such as dicentrine, undergoing a conformational change to a planar conformation upon binding to DNA [17]. Despite this weak intercalation, no effect on topoisomerase I and II activity was observed after boldine treatment. Its cytotoxicity is, therefore, probably independent of interactions with DNA or topoisomerases.

Nevertheless, boldine is a popular compound in the synthesis of derivatives with improved inhibitory activities. Boldine methine can inhibit angiotensin-II-induced leukocyte-endothelial cell interactions *in vivo*. The effects are mediated partly through the inhibition of the generation of ROS, the down-regulation of P-selectin expression on the endothelial cell, the inhibition of CXC chemokine and PAF release, and the blockade of PAF receptor. In contrast, boldine did not markedly affect the inflammatory activity elicited by angiotensin-II [18]. Other activities of boldine are of an anti- α -glucosidase, anti-leishmanial and anti-fungal nature [6].

Antiproliferative activities of boldine: Whereas the antioxidant activities of boldine are well documented [19], relatively little is known about its antiproliferative properties. A considerable difference is observed in the effects of boldine alone compared with boldo extract, which is a mosaic of active components. In infusions of boldo leaves about 40 compounds were detected. Thus, the medicinal properties reported for such infusions should be attributed not only to the presence of catechin and boldine, but also to other compounds [20]. That is, that the various effects of the decoction or extract from boldo leaves were not confirmed or observed after simple treatment with boldine, catechin, quercetin or rutin, even at concentrations exceeding those in the extract [21].

The anti-proliferative effect of *P. boldus* water extract is strongly cell dependent. The extract at a concentration as high as 3.5 mg of *P. boldus* extract/mL did not cause any appreciable decrease in Caco-2 cell viability. For HeLa cells, a *P. boldus* water extract concentration of 0.66 mg/mL was able to reduce cell viability by 50% [14]. In addition, the methanolic extract (containing 1.05% of boldine) inhibited cell growth and induced apoptosis in the M14 cell line [21].

Despite its weak effect on cell viability, a strong modification in the protein profile of HeLa cells appears. Some of these proteins may correspond to the cadherins or to the connexins. This may help cell communication in the gap-junctions, as treatment with *P. boldus* results in the detachment of cells [14]. In the M14 human cancer cell line, *P. boldus* leaf extract at 5 and 10 mg/mL induced a reduction in Hsp70 protein expression. This decrease was in correlation with a high level of DNA fragmentation and a significant increase in caspase-3 enzyme activity. A parallel growth in ROS production could amplify the apoptosis cascades. *P. boldus* leaf extract at higher concentrations (20–40 mg/mL) causes necrosis associated with high lactate dehydrogenase release and a further increase in ROS production. These processes generate intolerable oxidative stress in cancer cells [21]. A very important fact is that the extract of leaves of *P. boldus* exhibited no effect on the viability of normal fibroblast cells [20]. Also Gerhardt *et al.* suggested that boldine is selectively toxic because it may not affect hippocampal cells to the extent that it affects tumor cells [22].

Boldine reduced cellular proliferation by inducing G2 arrest followed by a reduction in mitotic cells in U138-MG exposed to 80 μ M for 72 hours. Boldine did not induce apoptosis in these cells.

Neither activation of caspases 3 or 9 nor an increase in the cleavage of PARP or DNA fragmentation were observed after exposure for 24 hours. Only C6 cells were propidium iodide positive after 500 μ M boldine treatment for 72 hours [22].

Further, the antiproliferative properties of boldine are improved by the synthesis of different derivatives. Two new compounds – 3-thiocarbamateboldine and 2,9-*O,O*-diacetyl-3-thiocarbamateboldine – were synthesized and their cytotoxicity evaluated by Thomet *et al* [23]. The derivatization of boldine to a diphenylphosphinyl derivative results in its enhanced lipophilicity as well as an increase in its intercalating behavior and thus in significant cytotoxic activity in breast cancer cells lines (MDA-MB-231 and MCF-7). Boldine itself has no cytotoxic capacity at an indicated concentration of 100 μ M for 72 hours [24]. Finally, Thomet *et al.* used a boldine derivative as ligand to synthesized boldiplatin. This compound has a comparable inhibitory concentration to oxaliplatin and, moreover, exerts greater selectivity against cancer tumor cell lines [25].

Antiproliferative activities of related alkaloids: It was reported that aporphine alkaloids, especially those containing a 1,2-methylenedioxy group, are effective against cancer cell lines [26]. On the other hand, nantenine, with a 9,10-methylenedioxy, and apomorphine, with no methylenedioxy group, also have proven antiproliferative properties [27,28]. Thus which structural part is responsible for the antiproliferative activity remains an open question.

All the discussed alkaloids can be isolated from various plant species; only representatives of these plants are named. Further information about sources of aporphine alkaloids are reviewed by Guinaudeau, Leboeuf and Cave [29,30].

Nantenine




Figure 2: Nantenine- (6a*S*)-1,2-Dimethoxy-6-methyl-5,6,6a,7-tetrahydro-4*H*-benzo[*de*][1,3]benzodioxolo[5,6-*g*]quinoline

Nantenine can be isolated from *Nandina domestica* (Berberidaceae) and *Uvaria chamae* (Annonaceae). Nantenine exerts a cytotoxic effect on both HCT-116 and Caco-2 cell lines with a potency comparable with that of the clinically available anticancer therapeutic drug etoposide [27], and a cytotoxic effect in mouse L929 transformed cells [28]. Like boldine, nantenine also seems to be selectively cytotoxic. At a concentration corresponding to the IC₅₀ against L929 cells, it was nontoxic to mouse thymocytes [28]. The C1 alkoxy derivative and the C1 benzoate derivative of nantenine exhibit marginally higher selective cytotoxicity to normal cells. The C1 phenolic group on the aporphine scaffold is detrimental to the cytotoxic activity [27].

Dicentrine

Dicentrine, isolated from *Lindera megaphylla* (Lauraceae), is an adaptive intercalator, a molecule that can adopt a relatively flat conformation. It is a strong topoisomerase II inhibitor with an IC₅₀ of 27 μ M, but is not a topoisomerase II poison [17]. Dicentrine may be compared with classical anti-cancer drugs such as anthracyclines. Its DNA intercalation, together with its inhibition of topoisomerases I and II, results in the formation of DNA strand breaks, which is followed by the activation of multiple signaling routes such as BRCA1-mediated DNA damage response, p53 signaling, G1/S and G2/M cell cycle regulation, and




Figure 3: Dicentrine - (7aS)-10,11-Dimethoxy-7-methyl-6,7,7a,8-tetrahydro-5H-[1,3]benzodioxolo[6,5,4-de]benzo[g]quinoline

aryl hydrocarbon receptor pathways. CDKN1A appeared as one of the differentially expressed genes upon dicentrine treatment. Moreover, dicentrine is selectively cytotoxic to EGFR-expressing cells; the overexpression of EGFR is associated with different types of cancer [3]. Dicentrine significantly inhibited cell growth and biosynthesis of the macromolecules DNA and RNA, and decreased the colony formation. Its cytotoxicity was confirmed in multiple human, as well as mouse tumor cell lines by an MTT and *in vitro* tumor growing assay [31,32]. Also, its derivative, (–)-nordicentrine, showed cytotoxicity against various human cancer cell lines [33].

Anonaine




Figure 4: Anonaine - (7aR)-6,7,7a,8-Tetrahydro-5H-[1,3]benzodioxolo[6,5,4-de]benzo[g]quinoline

Anonaine from *Michelia alba* (Magnoliaceae) induces dose-dependent DNA damage that is correlated with increased intracellular nitric oxide, ROS, glutathione depletion, disruptive mitochondrial transmembrane potential, activation of caspase 3, 7, 8 and 9, and PARP cleavage. (–)-Anonaine up-regulated the expression of Bax and p53 proteins in HeLa cancer cells. It is worth noting that the DNA damage did not occur in (–)-anonaine-treated non-cancer cell lines, either MDCK or Vero cells [34].

Apomorphine




Figure 5: Apomorphine - (6aR)-6-Methyl-5,6,6a,7-tetrahydro-4H-dibenzo[de,g]quinoline-10,11-diol

Apomorphine displays scavenger properties at low concentrations [35]. After treatment with low concentrations, an antiapoptotic response (increased bcl-xL, bcl-2; decreased bax) and increased cell viability were manifested [36]. At high concentrations of apomorphine a proapoptotic response, including the up-regulation of caspases-3 and 10, fas and fas-ligand, NF-κB p105 subunit, and tumor suppressor protein p53 mRNAs was observed [36]. It causes cell death at higher concentrations. Typical necrotic DNA degradation was found in C6 cells treated with 400 μM of apomorphine for 6 hours (loss of membrane integrity; degeneration of cytoplasmic organelles, especially mitochondria; and DNA fragmentation) [37].

References

- [1] Stévinny C, Bailly C, Quetin-Leclercq J. (2005) Cytotoxic and antitumor potentialities of aporphinoid alkaloids. *Current Medicinal Chemistry - Anti-Cancer Agents*, **5**, 173-82.
- [2] Zhao Q, Zhao Y, Wang K. (2006) Antinociceptive and free radical scavenging activities of alkaloids isolated from *Lindera angustifolia* Chen. *Journal of Ethnopharmacology*, **106**, 408-13.
- [3] Konkimalla VB, Efferth T. (2010) Inhibition of epidermal growth factor receptor over-expressing cancer cells by the aporphine-type isoquinoline alkaloid, dicentrine. *Biochemical Pharmacology*, **79**, 1092-1099.
- [4] Ray PD, Huang BW, Tsuji Y. (2012) Reactive oxygen species (ROS) homeostasis and redox regulation in cellular signaling. *Cellular Signalling*, **24**, 981-90.
- [5] Matés JM, Segura JA, Alonso FJ, Márquez J. (2012) Oxidative stress in apoptosis and cancer: an update. *Archives of Toxicology*, **86**, 1649-1665.

Roemerine




Figure 6: Roemerine - (7aR)-7-Methyl-6,7,7a,8-tetrahydro-5H-[1,3]benzodioxolo[6,5,4-de]benzo[g]quinoline

Roemerine, isolated from *Annona senegalensis* (Annonaceae), has no cytotoxic properties alone, but is able to enhance the cytotoxic response mediated by vinblastine with multidrug-resistant KB-V1 cells. This effect seems to be achieved via its interaction with P-glycoprotein [38]. Some cytotoxicity of roemerine was proven against K562S cells [39].

Duguetine




Figure 7: Duguetine - (7aS)-10,11-Dimethoxy-7-methyl-6,7,7a,8-tetrahydro-5H-[1,3]benzodioxolo[6,5,4-de]benzo[g]quinoline-8-ol

Duguetine, isolated from *Duguetia furfuracea* (Annonaceae), proved to have *in vitro* antitumor activity comparable with that of doxorubicin [26].

Conclusion: After the ascendancy of synthetic chemistry over natural product drug discovery and development during the latter part of the 20th century [40], nowadays the interest returns to the plant and folk medicine as a source of either new therapeutic substances or chemical scaffolds for drug synthesis. One such promising group is the family of aporphine alkaloids comprising boldine and its derivatives that is used in folk medicine to treat headache, earache, rheumatism, dyspepsia, menstrual pain, urinary tract inflammation and others [19]. Ample evidence demonstrates the ability of boldine to act as a potent free radical-scavenger and antioxidant molecule. However, recent studies show that boldine and other aporphine alkaloids also exhibit antitumor properties which are not related to oxidative stress management. Boldine can be isolated quite easily from various plants, and, due to its beneficial properties, could represent another new phytotherapeutic drug arising from folk medicine.

Abbreviations: ROS reactive oxygen species, PPAR peroxisome proliferator-activated receptor, PAF platelet-activating factor, PARP poly ADP ribose polymerase

Acknowledgments - This work was supported by Charles University project Prvok P37/1. J. Cmielova is co-financed by the European Social Fund and the state budget of the Czech Republic. Project no. CZ.1.07/2.3.00/30.0022.

- [6] Mollataghi A, Coudiere E, Hadi AH, Mukhtar MR, Awang K, Litaudon M, Ata A. (2012) Anti-acetylcholinesterase, anti- α -glucosidase, anti-leishmanial and anti-fungal activities of chemical constituents of *Beilschmiedia* species. *Fitoterapia*, **83**, 298-302.
- [7] Han Z, Zheng Y, Chen N, Luan L, Zhou C, Gan L, Wu Y. (2008) Simultaneous determination of four alkaloids in *Lindera aggregata* by ultra-high-pressure liquid chromatography-tandem mass spectrometry. *Journal of Chromatography A*, **1212**, 76-81.
- [8] Cassels BK, Asencio M, Conget P, Speisky H, Videla LA, Lissi EA. (1995) Structure-antioxidative activity relationships in benzyloisoquinoline alkaloids. *Pharmacological Research*, **31**, 103-107.
- [9] Loo G. (2003) Redox-sensitive mechanisms of phytochemical-mediated inhibition of cancer cell proliferation. *The Journal of Nutritional Biochemistry*, **14**, 64-73.
- [10] Konrath EL, Santin K, Nassif M, Latini A, Henriques A, Salbego C. (2008) Antioxidant and pro-oxidant properties of boldine on hippocampal slices exposed to oxygen-glucose deprivation *in vitro*. *NeuroToxicology*, **29**, 1136-1140.
- [11] Yu B, Cook C, Santanam N. (2009) The aporphine alkaloid boldine induces adiponectin expression and regulation in 3T3-L1 cells. *Journal of Medicinal Food*, **12**, 1074-1083.
- [12] Lau YS, Tian XY, Huang Y, Murugan D, Achike FI, Mustafa MR. (2013) Boldine protects endothelial function in hyperglycemia-induced oxidative stress through an antioxidant mechanism. *Biochemical Pharmacology*, **85**, 367-375.
- [13] Backhouse N, Delporte C, Givernau M, Cassels BK, Valenzuela A, Speisky H. (1994) Anti-inflammatory and antipyretic effects of boldine. *Agents and Actions*, **42**, 114-117.
- [14] Falé PL, Amaral F, Amorim Madeira PJ, Sousa Silva M, Florêncio MH, Frazão FN, Serralheiro ML. (2012) Acetylcholinesterase inhibition, antioxidant activity and toxicity of *Peumus boldus* water extracts on HeLa and Caco-2 cell lines. *Food and Chemical Toxicology*, **50**, 2656-2662.
- [15] Speisky H, Squella JA, Núñez-Vergara LJ. (1991) Activity of boldine on rat ileum. *Planta Medica*, **57**, 519-522.
- [16] Hoet S, Stévinny C, Block S, Oppendoes F, Colson P, Baldeyrou B, Lansiaux A, Bailly C, Quetin-Leclercq J. (2004) Alkaloids from *Cassytha filiformis* and related aporphines: antitrypanosomal activity, cytotoxicity, and interaction with DNA and topoisomerases. *Planta Medica*, **70**, 407-413.
- [17] Woo SH, Sun NJ, Cassady JM, Snapka RM. (1999) Topoisomerase II inhibition by aporphine alkaloids. *Biochemical Pharmacology*, **57**, 1141-1145.
- [18] Estellés R, Milian L, Nabah YN, Mateo T, Cerdá-Nicolás M, Losada M, Ivorra MD, Issekutz AC, Cortijo J, Morcillo EJ, Blázquez MA, Sanz MJ. (2005) Effect of boldine, secoboldine, and boldine methine on angiotensin II-induced neutrophil recruitment *in vivo*. *Journal of Leukocyte Biology*, **78**, 696-704.
- [19] O'Brien P, Carrasco-Pozo C, Speisky H. (2006) Boldine and its antioxidant or health-promoting properties. *Chemico-Biological Interactions*, **159**, 1-17.
- [20] Simirgiotis MJ, Schmeda-Hirschmann G. (2010) Direct identification of phenolic constituents in Boldo Folium (*Peumus boldus* Mol.) infusions by high-performance liquid chromatography with diode array detection and electrospray ionization tandem mass spectrometry. *Journal of Chromatography A*, **1217**, 443-449.
- [21] Russo A, Cardile V, Caggia S, Gunther G, Troncoso N, Garbarino J. (2011) Boldo prevents UV light and nitric oxide-mediated plasmid DNA damage and reduces the expression of Hsp70 protein in melanoma cancer cells. *Journal of Pharmacy and Pharmacology*, **63**, 1219-29.
- [22] Gerhardt D, Horn AP, Gaelzer MM, Frozza RL, Delgado-Cañedo A, Pelegrini AL, Henriques AT, Lenz G, Salbego C. (2009) Boldine: a potential new antiproliferative drug against glioma cell lines. *Investigational New Drugs*, **27**, 517-525.
- [23] Thomet FA, Pinyol P, Villena J, Espinoza LJ, Reveco PG. (2010) Cytotoxic thiocarbamate derivatives of boldine. *Natural Product Communications*, **5**, 1587-90.
- [24] Thomet FA, Piñol P, Villena J, Reveco PG. (2011) *In vitro* cytotoxic evaluation of a novel phosphinyl derivative of boldine. *Molecules*, **16**, 2253-2258.
- [25] Thomet FA, Pinyol P, Villena J, Reveco PG. (2012) Towards a more selective analogue of oxaliplatin: Synthesis of [Pt((1*R*,2*R*)-diaminocyclohexane)(3-carboxypredicentrinato)]. *Inorganica Chimica Acta*, **384**, 255-259.
- [26] da Silva DB, Tulli EC, Militão GC, Costa-Lotufo LV, Pessoa C, de Moraes MO, Albuquerque S, de Siqueira JM. (2009) The antitumoral, trypanocidal and antileishmanial activities of extract and alkaloids isolated from *Duguetia furfuracea*. *Phytomedicine*, **16**, 1059-1063.
- [27] Ponnala S, Chaudhary S, González-Sarrias A, Seeram NP, Harding WW. (2011) Cytotoxicity of aporphines in human colon cancer cell lines HCT-116 and Caco-2: an SAR study. *Bioorganic & Medicinal Chemistry Letters*, **21**, 4462-4464.
- [28] Philipov S, Ivanovska N, Istatkova R, Velikova M, Tuleva P. (2000) Phytochemical study and cytotoxic activity of alkaloids from *Uvaria chamae* P. Beauv. *Die Pharmazie*, **55**, 688-689.
- [29] Shamma M, Guinaudeau H. (1986) Aporphinoid alkaloids. *Natural Product Reports*, **3**, 345-351.
- [30] Guinaudeau H, Leboeuf M, Cave A. (1975) Aporphine alkaloids. *Lloydia*, **38**, 275-338.
- [31] Huang RL, Chen CC, Huang YL, Ou JC, Hu CP, Chen CF, Chang C. (1998) Anti-tumor effects of d-dicentrine from the root of *Lindera megaphylla*. *Planta Medica*, **64**, 212-215.
- [32] Kondo Y, Imai Y, Hojo H, Endo T, Nozoe S. (1990) Suppression of tumor cell growth and mitogen response by aporphine alkaloids, dicentrine, glaucine, corydine, and apomorphine. *Journal of Pharmacobio-dynamics*, **13**, 426-431.
- [33] Lekphrom R, Kanokmedhakul S, Kanokmedhakul K. (2009) Bioactive styryllactones and alkaloid from flowers of *Goniotalamus laoticus*. *Journal of Ethnopharmacology*, **125**, 47-50.
- [34] Chen CY, Liu TZ, Tseng WC, Lu FJ, Hung RP, Chen CH, Chen CH. (2008) (-)-Anonaine induces apoptosis through Bax- and caspase-dependent pathways in human cervical cancer (HeLa) cells. *Food and Chemical Toxicology*, **46**, 2694-2702.
- [35] El-Bachá RS, Netter P, Minn A. (1999) Mechanisms of apomorphine cytotoxicity towards rat glioma C6 cells: protection by bovine serum albumin and formation of apomorphine-protein conjugates. *Neuroscience Letters*, **263**, 25-28.
- [36] Weinreb O, Mandel S, Youdim MB. (2003) cDNA gene expression profile homology of antioxidants and their antiapoptotic and proapoptotic activities in human neuroblastoma cells. *The FASEB Journal*, **17**, 935-937.
- [37] dos Santos El-Bachá R, Daval J, Koziel V, Netter P, Minn A. (2001) Toxic effects of apomorphine on rat cultured neurons and glial C6 cells, and protection with antioxidants. *Biochemical Pharmacology*, **61**, 73-85.
- [38] You M, Wickramaratne DB, Silva GL, Chai H, Chagwedera TE, Farnsworth NR, Cordell GA, Kinghorn AD, Pezzuto JM. (1995) (-)-Roemerine, an aporphine alkaloid from *Annona senegalensis* that reverses the multidrug-resistance phenotype with cultured cells. *Journal of Natural Products*, **58**, 598-604.
- [39] Baghdikian B, Mahiou-Leddert V, Bory S, Bun SS, Dumetre A, Mabrouki F, Hutter S, Azas N and Ollivier E. (2013) New antiplasmodial alkaloids from *Stephania rotunda*. *Journal of Ethnopharmacology*, **145**, 381-385.
- [40] Fowler MW. (2006) Plants, medicines and man. *Journal of the Science of Food and Agriculture*, **86**, 1797-1804.
- [41] Chemical structures and nomenclature: www.chemspider.com

New Therapeutic Potentials of Milk Thistle (*Silybum marianum*)Nataša Milić^a, Nataša Milošević^{a*}, Ljiljana Suvajdžić^a, Marija Žarkov^b and Ludovico Abenavoli^c^aDepartment of Pharmacy, Faculty of Medicine, University of Novi Sad, Novi Sad, Serbia^bDepartment of Neurology, Faculty of Medicine, University of Novi Sad, Novi Sad, Serbia^cDepartment of Health Sciences, University Magna Graecia, Catanzaro, Italy

milosevic_natasa@gmail.com

Received: August 26th, 2013; Accepted: September 18th, 2013

Silymarin is a bioflavonoid complex extract derived from dry seeds of Milk thistle [*Silybum marianum*(L.) Gaertn. (Fam. Asteraceae/Compositaceae)] whose hepatoprotective effect has clinically been proved. Low toxicity, favorable pharmacokinetics, powerful antioxidant, detoxifying, preventive, protective and regenerative effects and side effects similar to placebo make silymarin extremely attractive and safe for therapeutic use. The medicinal properties of silymarin and its main component silibinin have been studied in the treatment of Alzheimer's disease, Parkinson's disease, sepsis, burns, osteoporosis, diabetes, cholestasis and hypercholesterolemia. Owing to its apoptotic effect, without cytotoxic effects, silymarin possesses potential applications in the treatment of various cancers. Silymarin is being examined as a neuro-, nephro- and cardio-protective in the damage of different etiologies due to its strong antioxidant potentials. Furthermore, it has fetoprotective (against the influence of alcohol) and prolactin effects and is safe to be used during pregnancy and lactation. Finally, the cosmetics industry is examining the antioxidant and UV-protective effects of silymarin. Further clinical studies and scientific evidence that silymarin and silibinin are effective in the therapy of various pathologies are indispensable in order to confirm their different flavonolignan pharmacological effects.

Keywords: Milk thistle, Silymarin, Silibinin, Therapeutic, Protective.

Milk thistle [*Silybum marianum* (L.) Gaertn., family Asteraceae/Compositaceae] is probably the oldest and the best-studied plant in the treatment of liver diseases. The first records of milk thistle healing effects are recorded in the Old Testament [1]. Silymarin standardized dry extract is a bioflavonoid complex consisting of at least seven flavonolignans (Figure 1). Silibinin is the major component of this complex extract (about 60-70%) and is biologically the most active constituent of silymarin, followed by silicristin (20%), silidianin (10%), and isosilybin (5%). Silibinin is a mixture of two stereoisomers, silibinin A and silibinin B in the equimolar ratio 1:1 [2].

Modern Western medicine has confirmed hepatoprotective properties of silymarin by conducting animal studies and clinical trials [1]. Silymarin has antioxidant, anti-inflammatory, antifibrotic, detoxifying and regenerative properties. It stimulates protein synthesis and liver regeneration. Silymarin has numerous hepatoprotective effects: it prevents lipid peroxidation by scavenging free radicals and increases the levels of reduced glutathione (GSH). It regulates membrane permeability and membrane stability in damage caused by xenobiotics, regulates nuclear expression and inhibits the transformation of hepatocytes in myofibroblasts (with cirrhosis) [3, 4]. New *in vivo* studies have proved silymarin to be a strong hepatoprotective in hepatic fibrosis induced by carbon tetrachloride (CCl₄). After silymarin had been administered to the rats treated previously with CCl₄, the levels of glutamic oxaloacetic transaminase and glutamic pyruvic transaminases were normalized and the secretion of the connective tissue growth factor was decreased, indicating that silymarin ameliorates the destructive changes in the liver after hepatic fibrosis was provoked [5]. An insight has recently been published into the possible mechanism of silymarin defense capacity, as an antioxidant against various hepatotoxicants. Orally administered silymarin to the experimental mice in only three doses of 100 or 200 mg/kg increased the amount of metabolites generated from homocysteine

in the transsulfuration pathway (cystathionine, cysteine, and glutathione), elevated the activity of cystathionine β -synthase, while down-regulated cysteine dioxygenase. This resulted in the augmented oxygen radical scavenging capacity of the liver cytosol and reduced lipid peroxidation. For the first time, it was demonstrated that the antioxidant capacity of silymarin was connected to the hepatic glutathione production in the liver by cysteine synthesis increment and inhibited degradation to taurin [6]. Silymarin inhibits the absorption of α -falloidine and α -amanitin toxins (from the mushroom *Amanita phalloides*) as it prevents their binding to the cell surface and limits the transportation systems on the cell membranes. Silymarin/silibinin (SIL) stimulates the regeneration of hepatocytes by nuclear A polymerase activation, increasing protein synthesis and inhibiting the expression of the adhesion molecules (E-selection) [7]. The anti-inflammatory effect of silymarin prevents the activation of the intrahepatic nuclear factor kappa B (NF- κ B), reducing the levels of tumor necrosis factor alpha (TNF- α), interleukin-2 (IL-2), interferon gamma (IFN- γ) and the inducible nitric oxide synthase (iNOS) [8]. It was shown that silymarin was hepatoprotective [9], and it was used in the treatment of nonalcoholic fatty liver [4], alcoholic liver cirrhosis [10], and acute and chronic hepatitis [11].

SIL possesses some other potential remedying properties and the new therapeutic effects of milk thistle extract have been considered and proved in various preclinical and clinical studies which are presented in this paper.

Pharmacokinetics of SIL: Silymarin/silibinin is rapidly, but poorly absorbed due to its high liposolubility and poor water solubility. After an oral administration of silymarin (equivalent to 120 mg silibinin), the peak plasma concentration was reached within 1-2 h, but the maximum concentration was only 1.1 to 1.3 mg/mL [12]. The increased bioavailability of silymarin was achieved by complexation with phosphatidylcholine, β -dextran or a suitable




Figure 1. Flavonolignans of Milk thistle extracts.

material for capsulation. Silibinin in plasma is rapidly distributed in liver, lungs, skin, prostate, and the pancreas, and the maximum level is achieved within just one hour after oral administration. About 70% of silibinin binds to plasma proteins, and the concentration in the bile is 100 times higher than in plasma. In humans, higher doses of silibinin are subjected to I and II metabolism phases, whereby the liver microsomes produce one dominant demethylated metabolite in the first phase, while three mono-hydroxyl metabolites and one dihydroxyl metabolite are generated in small amounts. Silibinin monoglucuronide, silibinin diglucuronide, silibinin monosulfate and silibinin glucuronide sulfate are produced in the second phase, and then *O*-demethyl silibinin glucuronide and silibinin triglucuronide. The half-time elimination of silymarin (silibinin equivalent to 25%) is 6.32 h and it is mainly excreted via bile. Only 1 to 7% of silymarin is excreted into the urine [13-15]. However, the urinary excretion of 31 detected metabolites was followed by clearance for 24 h while substantial amounts of glucuronides, sulfo-glucuronides and diglucuronides were excreted within 48 h. A non-marginal entero-hepatic recirculation, which preferentially occurs for silymarin sulfate-glucuronides and diglucuronides, may be the explanation for a delayed clearance of the conjugated metabolites. Although, polyphenols are drastically modified by the colon microbiota, no microbially derived compounds were detected after the oral administration of silymarin [16].

Silymarin side effects: German Commission E approved silymarin for various liver conditions and stated that it had no side effects at the recommended doses. It is safe for use in pregnancy (560 mg/day), in children (20-50 mg intravenously) and in elders over 75 years (420 mg/day). The toxic effects of the milk thistle extracts have not been recorded at a dose of 1200 mg/day, although it was administered at doses of 160-600 mg/day divided in three equal doses in most of the clinical studies. However, mild allergies were

recorded at doses greater than 1500 mg a day. A few patients reported heartburn, diarrhea, bloating, flatulence and dyspepsia. It is proved that silymarin is well-tolerated and safe for use up to 41 months [17].

Silymarin interaction with drugs: Silymarin inhibited the enzymes of phase I metabolism: CYP2E1, CYP2D6, CYP2C19, CYP1A2 and CYP2A6 and the enzymes of the phase II metabolism such as uridine diphosphoglucuronyltransferase (UGT) 1A1 isozymes, and due to this ability it could cause interactions with the drugs that are metabolized with the same enzymes. It was also detected that silymarin, in a dose-dependent manner, inhibited P-glycoprotein, which is responsible for the excretion of the xenobiotics from the liver into the hepatobiliary duct and could cause the drug accumulation that is excreted via those routes. However, only the interactions of silymarin with metronidazole (increases the clearance of metronidazole by about 30%), pyrazinamide (prevents its excretion) [13] and warfarin (increases the risk of bleeding) have been detected so far [18].

Effect on Alzheimer's disease and dementia: Alzheimer's disease (AD) is characterized by progressive cognitive impairment and protein plaque deposits. The extracellular aggregation of amyloid β ($A\beta$ -peptide), a protein plaque component, found in the brain cells of the patients with AD, presents the major histopathological change in this disease. It is assumed that $A\beta$ -peptide generates H_2O_2 , which is instantly converted into a highly reactive hydroxyl radical which causes an oxidative stress. In the *in vitro* study on human neuroblastoma SH-SZ5Y, it was proved that silibinin, in a dose-dependent manner, inhibited the aggregation of $A\beta$ -peptide, decreased $A\beta$ -peptide cytotoxicity, reduced hydrogen peroxide production and cell injury. Silibinin is a powerful anti-oxidative agent which increases GSH and superoxide dismutase (SOD) levels

and inhibits lipid peroxidation. The study confirmed the anti-oxidative effects of silibinin and assumed that the decrease of H_2O_2 production, induced by A β aggregation, was the main route for its neuroprotective activity [19].

Pretreatment with silibinin in a dose-dependent manner improved the spatial memory and energy metabolism in the brain, reduced cholinergic dysfunction and prevented the accumulation of lipid peroxides in mice after streptozotocin (STZ) injection had been administered. Several studies reported that the impaired brain energy metabolism was the key event in AD pathophysiology. It was demonstrated that dysregulated Ca^{2+} intracellular homeostasis and increased free radical formation disturbed the mitochondrial function and caused the changes in brain energy metabolism. The pretreatment with silibinin eliminated the disruption of the metabolism energy (the defects in the electron-transport chain of mitochondria and the reduced production of ATP in the mice brains that STZ caused), reduced the levels of reactive oxygen species (ROS) and malonyldialdehyde (MDA), and increased GSH and Ca^{2+} levels in the synaptic space. Loss or down-regulation of the neuronal nicotinic acetylcholine receptors (nAChRs) was also observed in AD. Silibinin reduced the anti-acetylcholinesterase action, restored the function of acetylcholinesterase and mRNA expression in a dose-dependent manner in the brain of the mice that were treated with STZ. In addition, it prevented the decreased expression of the nAChRs (particularly the subtypes which are predominant in the brain and cause the impairment of the cognitive functions) [20].

The cognitive deficit was observed in patients with diabetes and this disease was often cited as a risk factor for the development of vascular dementia and Alzheimer's disease. It is supposed that the stimulation of ubiquitous and inducible heme oxygenase (HO)-1, caused by silibinin, was accountable for the silibinin antioxidant and anti-inflammatory activity as the HO isoforms catalyzed the conversion of heme to carbon monoxide and bilirubin, and its increased expression was the response to the oxidative stress in the CNS. The level of HO-1 was reduced in the forebrain and cerebellum of the diabetic mice, while it was unexpectedly increased in the brainstem; treatment with silibinin caused a considerable induction of HO-1 in the forebrain and cerebellum, but it provoked inhibition in the brainstem. Silibinin decreased isoprostanes and 8-OH deoxyguanosine (markers of lipid peroxidation and DNA damage, respectively) in the hippocampus and hypothalamus of the treated diabetic animals. Silibinin protected DNA from the oxidative stress and therefore, it could be considered as an important phytotherapeutic agent in the treatment strategy of the accompanying diabetic side effects on the CNS [21].

Silibinin's favorable effects on the cognitive function was confirmed by *in vivo* studies on mice whose memory loss was caused by methamphetamine (METH), which reduced the ability to recognize a new object and the five-fold choice. METH provokes hallucinations and delusions in humans, and METH chronic use causes cognitive deficits after withdrawal. Silibinin neutralized the METH effects and reduced the time necessary to recognize the new items (impairment in the novel object recognition test). Silibinin administered to the healthy mice did not cause any changes in the serotonin and dopamine levels, but annulled the effects that METH had on these systems if it had been given as a pretreatment. It is believed that silibinin inhibited monoamine oxygenase (MAO) and reduced the dopamine and serotonin metabolisms. The silibinin antioxidant effect reduced the peroxynitrite concentration which caused the dopamine and serotonin concentrations to be maintained in the brain. Silibinin anti-inflammatory effects protected microglial

activation in the midbrain, striatum, thalamus, orbitofrontal and insular cortices provoked by METH. It additionally gave protection from the neurotoxicity caused by lipopolysaccharides that initiated a pro-inflammatory cascade and released the cytokines. Therefore, silibinin can have a potentially important role in the therapy of drug addicts [22].

Effect on Parkinson's disease: The possible causes of the degeneration of the dopaminergic neurons in Parkinson's disease (PD) are: reduced GSH levels, DNA damage, iron deposition and primarily, the oxidative stress that impairs the oxidative phosphorylation and energy metabolism that lead to dopaminergic neuron death. Silymarin postponed the oxidative damage of the neurons (reducing the synthesis of ROS and maintaining SOD levels) that had been developed in many neurodegenerative diseases by inhibiting MAO in glial cells, by stimulating peroxidation product secretion and, probably, by stimulating the synthesis of ribosomal RNA as it bonds via the estradiol receptors [23].

Anticarcinogenic effects: The anti-tumor effect of silymarin was detected in tumors induced in the epidermis of mice in 1994 [24]; this led to a number of studies. Silymarin is classified in the group of effective chemoprotective and chemopreventive agents in the treatment of various cancer conditions. In 1999, it was reported that silibinin was probably responsible for its anti-proliferative effect since it inhibited DNA synthesis in prostate carcinoma LNCaP and DU145 cells, breast carcinoma MCF-7 cells and cervical carcinoma A431 cells [25].

Effect on skin cancer: Silymarin demonstrated the capacity to reduce the expression of the TNF- α endogenous promoter in SENCAR mice with chemically induced skin cancer. Orally administered silymarin not only inhibited the tumor growth, but it further decreased the volume of the existing tumors (80-97%) with no signs of toxicity to the animals. Silymarin inhibited all the proteins of the mitogen-activated protein kinase (MAPK) family: ERK1/2, JNK and p38 and induced apoptosis; it was irrelevant whether silymarin had been administered orally or topically [26-28]. It can be applied in a benign condition such as premalignant keratosis. It was found that silymarin possessed a protective effect in SKH-1 mice with UVB radiation-induced skin tumor initiation, tumor growth and complete carcinogenesis. Silymarin significantly affected the inhibition of UVB-induced sunburns and the formation of apoptotic cells, skin oedema, catalase activity reduction, induction of expression and activity of cyclooxygenase (COX) and ornithine decarboxylase (ODC), when applied either topically or orally before or after exposure to UVB. Regulating the p53-p21 cascade that protected the DNA from damage, silymarin reduced a number of the thymidine dimer-positive cells and slowed the kinetics of the DNA damage by stimulating its reparation after the UV exposure. Silymarin induced the level of Cip1/p21 protein in the epidermis of the mice during the acute radiation with UVB rays, thus inhibiting the cell proliferation and expression of the genes responsible for the cell growth in a cascade way [29-31]. In addition, in novel studies, it was found that the main constituent of silymarin, silibinin, saved the cells from apoptosis through inhibition of IGF-1R activation followed by repression of ERK1/2 and JNK phosphorylation [32] and induced autophagy protection of human carcinoma A431 cells from UVB-induced apoptosis [33] in UVB-irradiated A431 cells. Furthermore, silibinin contributed to the reduction of A, E and D1 cyclins, increased the protein levels of CDKIs, Cip1/p21 and Kip1/p27 in the tumor cells, inhibited the activity of cyclin-dependent kinase (CDK) and thus decreased the proliferation and growth of the tumor cells. Therefore, it is expected that the silibinin non-toxic photo-protective effects will be

considered as a potentially useful chemo-preventive agent for the treatment of skin cancer [34].

Effect on hepatocellular carcinoma: The positive effects of silymarin in the treatment of hepatocellular carcinoma were reported in several *in vivo* and *in vitro* studies. In the study conducted on Wistar rats with induced hepatocellular carcinoma, treatment with silymarin (1000 ppm orally, 16 weeks before or 5 weeks after tumor induction) regulated the increased levels of aminotransferases (AST and ALT), phosphatases (ACP and ALP), lactate dehydrogenase, gamma-glutamyl transferase and 5'-nucleotidase, tumor marker alpha-fetoprotein (AFP) and carcinoembryonic antigen (CEA). The reduction in MDA-DNA formation was recorded in the animals with tumors that had received silymarin compared with the control group [35]. The research on H4IIE hepatocytes, where the tumor had been induced with a high ethanol concentration, confirmed the hepatoprotective effects of SIL: silibinin inhibited CYP2E1 (but not alcohol dehydrogenase) and ethanol metabolism, reduced the ROS level, while it also reduced the proliferation and progression of the cell cultures in the rats suffering from hepatocellular carcinoma at a dose of 10 μ M [36], which was consistent with the findings that silibinin inhibited cell proliferation, matrix metalloproteinase 2 enzymatic activity, NO production and ERK 1/2 phosphorylation in a dose-dependent manner in the HepG-2 cell line [37]. Furthermore, recent studies confirmed that pretreatment with silymarin down-regulated the cytokines-gene expression in the human hepatocellular carcinoma cell lines susceptible to amphotericin B-induced oxidative stress [38].

Effect on breast cancer: Silymarin inhibited both anchorage-dependent and anchorage-independent cell growth in dose- and time-dependent manners in the treatment of human breast carcinoma. Its effects on the cell growth and proliferation were probably connected with an induction of up to 19-fold in the protein expression of CDK inhibitor Cip1/p21G1 which caused the arrest in the cell cycle progression [39]. In addition, silibinin had a synergistic effect with anticancer chemotherapeutics such as doxorubicin, cisplatin and carboplatin in the human breast cancer cells. Silibinin, in a dose-dependent manner, inhibited the expression of matrix metalloproteinase 9 (MMP-9), preventing the transcription of this protein by blocking the pathway that went through MAPK and by suppressing the AP-1 proteins with oligonucleotides containing the sequence for the AP-1 DNA at the binding site of the MMP-9, preventing the expression of MMP-9 in the *in vitro* cultures of the human breast cancer cells [40, 41]. The results were even more impressive with the use of silibinin-loaded lipid nanoparticles containing D- α -tocopheryl-polyethyleneglycol 1000 succinate (TPGS) [42]. It was recently found that silibinin was a Wnt/ β -catenin signaling inhibitor by suppressing Wnt co-receptor LRP6 expression at the transcription level, and that the anti-cancer activity of silibinin was associated with its inhibitory effect on Wnt/LRP6 signaling [43].

Effect on ovarian cancer: Silipide (IDB 1016 - silibinin complex with phosphatidylcholine), administered orally to hairless mice which had been embedded with the xenograft of human ovarian cancer cells, contributed to a significant reduction in the tumor mass by 78% [44]. It was found that silibinin enabled the overcoming of resistance to paclitaxel. It reduced the invasiveness of the paclitaxel-resistant tumor cells in another *in vitro* study on paclitaxel-sensitive and paclitaxel-resistant cells [45]. If these effects are proved in *in vivo* conditions, silibinin in combination with paclitaxel could become a part of the strategy in the treatment of tumors in patients resistant to paclitaxel.

Effect on cervical cancer: Silymarin has a possible application in chemo-prevention and in the treatment of cervical carcinoma. Silymarin caused the apoptosis of HeLa cells of human cervical cancer at low doses (80 μ mol/L) and cell necrosis at high doses (160 μ mol/L). The apoptosis caused by silymarin via MAPK activation was due to chromatin condensation and nuclear fragmentation of the apoptotic cells [46]. It was found that silibinin inhibited hypoxia-induced HIF-1 α accumulation and HIF-1 transcriptional activity in human cervical (HeLa) and hepatoma (Hep3B) cells, activated Akt, and reduced hypoxia-induced vascular endothelial growth factor (VEGF) released by HeLa and Hep3B cells, and finally, inhibited cell proliferation [47].

Effect on bladder cancer: Silibinin inhibited cell growth and caused cell cycle arrest in the G1 and G2/M phases of the highly invasive TCC-SUP cells (apoptosis), and the G1 phase of the cell cycle in T-24 of human bladder carcinoma cells (without apoptosis), respectively. Silibinin acted in the T-24 cells through erbB1-mediated mitogenic signalization, inhibiting the cell growth without causing apoptosis [48]. It was found that orally administered silymarin, at a dose of 1000 ppm for 8 weeks during the initiation of cancer, or 24 weeks after the initiation, caused statistically significant reduction in the number of pre-neoplastic lesions and the rate of the disease progression of bladder cancer in the *in vivo* study on ICR mice with induced bladder cancer [49]. Furthermore, silibinin inhibited the tumor RT4 xenograft growth by decreasing the tumor volume and weight, reduced cell proliferation and increased apoptosis in the tumors in the *in vivo* studies. The assumed mechanism of action was decrement of the surviving protein expression and its nuclear localization [50].

Effect on lung cancer: Silibinin slowed the growth of A549 human lung cancer cells and enhanced the effect of doxorubicin modulation of the NF- κ B route thus inhibiting the resistance to doxorubicin, significantly reduced the levels of iCOX-2 and diminished the adverse effects of the chemotherapeutics [51]. In addition, silibinin reversed the resistance to etoposide and to doxorubicin in human small-cell lung carcinoma and acted synergistically with the chemotherapeutics [52]. Oral pretreatment with silibinin statistically reduced the number of tumors and large tumors (> 1.5 mm) in relation to the control group in the study on A/J mice, where lung cancer had been provoked by urethane. This study evaluated the density of blood vessels where it was found that silibinin decreased the number of highly vascularized tumors and the density of the blood vessels in the vascularized tumors. Finally, silibinin decreased the levels of iNOS and COX-2 enzymes whose expression increased in the process of angiogenesis responsible for the tumor development [53]. The pretreatment with silibinin reduced the phosphorylation of STAT1 and STAT3 induced by cytokines responsible for the proliferation of the A549 human lung cancer cells under the *in vitro* conditions. Silibinin inhibited the AP-1 transcription factor of DNA, blocked the MAPK cascade and inhibited iNOS at the concentration of 200 μ mol/L, indicating that it could be a potential chemo-preventive agent for lung cancer treatment [54]. Silibinin-meglumine, a water-soluble form, inhibited the growth of non-small-cell lung carcinoma mouse xenografts as efficiently as the "gold standard" gefitinib and reversed the resistance of gefitinib-unresponsive tumors [55].

Effect on prostate cancer: SIL possessed anti-carcinogenic activity both with hormone-dependent and hormone-independent prostate cancer and its component isosilybin B was demonstrated as the most effective suppressor of the topo IIa gene promoter activity. The inhibiting effect on the prostate cancer cells in the LNCaP, PC3 and DU145 cultures that SIL exhibited was not registered in the

normal prostate epithelial cells [56]. Silibinin induced increased expression of cytokeratins and chromogranin A and modulated the phosphorylation status of retinoblastoma (Rb) and Rb-related proteins in human prostate carcinoma LNCaP cells. Furthermore, it caused a considerable reduction in the prostate specific antigen (PSA) in the LNCaP cells, which was important for cell growth inhibition. SIL down-regulated epidermal growth factor receptor (EGFR) signaling in PCA, which consequently led to the cascade that inhibited CDK and cancer cell cycle arrest. Silibinin/silymarin inhibited tumor growth factor (TGF- α) mRNA expression and decreased secreted and cellular levels of TGF- α in both LNCaP and DU145 cells. It not only reduced NF- κ B signaling activity in DU145 prostate cancer cells, but also increased the sensitivity of the cells to TNF α -induced apoptosis that was commonly inhibited [57]. Additionally, the silipide/silibinin effect in prostate cancer was accomplished by reducing the level of insulin-like growth factor-1 (IGF-1) and/or elevating the level of insulin-like growth factor-binding protein-3 (IGFBP-3) in TRAMP mice [58]. A dose-dependent decrease was also observed in cyclin B1, cyclin E, and cyclin A protein levels by silibinin, which suggested that oral silibinin blocked PCA growth and progression via modulation of the tumor IGF-IGFBP-3 axis and cell cycle regulation [59].

Effect on oral cancer: Silymarin was proved to decrease the viability of human pharynx squamous carcinoma (FaDu) cells in an *in vitro* study. Apoptosis of the FaDu cells was preceded by 10-fold diminished Akt phosphorylation and five to six-fold upregulated expression of the phosphatase and tensin homolog. This caused the inhibition of Bcl-2 expression and elevation of caspases activity which led to the apoptosis of the FaDu cells [60].

Effect on gastric cancer: Silibinin dose-dependently inhibited the growth of human gastric carcinoma SGC-7901 cells. Namely, it provoked G2 phase arrest in the cell-cycle progression of the SGC-7901 cell line. SGC-7901 cells growth inhibition was observed after silibinin significantly decreased the expression of p34cdc2 levels and induced p53 and p21 expression. A final common pathway, involving the activation of caspase, is usual for most chemotherapeutic agents. Surprisingly, silibinin caused apoptotic death of SGC-7901 cells in a caspase-independent manner [61].

Effect on colon cancer: Silibinin, in a dose-dependent manner, induced cell cycle arrest in the G2/M phase of the FET and GEO cell lines of human colon cancer and in the poorly differentiated HCT116 cells in the G1 phase. As with prostate cancer, silibinin inhibited CDK and the activity of D1, and E cyclins in colon carcinoma, but did not reduce COX-2 expression and activity. It also inhibited autocrine TGF- α secretion, and its binding to the EGFR and the EGFR expression [62]. Significant reduction in the frequency of colonic ACF (aberrant crypt foci) in a dose-dependent manner in F344 rats was achieved after dietary administration of silymarin (100, 500 and 1,000 ppm), either during or after 4 weeks of exposure to the carcinogens [63]. Orally administered silibinin, before or during the tumor initiation, caused statistically significant reduction in the number of formed tumors, while it prevented the development of macroscopically visible tumors (only polyps were developed) during the entire period of the study on Wistar rats with induced colon cancer, using 1,2-dimethylhydrazine (DMH). Silibinin modulated the enzyme activity of an intestinal micro flora, reduced oxidative stress in the colon and prevented the retoxification of DMH (it inhibited cytochrome P450 activity, the creation of the DMH carcinogenic metabolites, and also induced the enzyme induction of second phase metabolism which facilitates DMH excretion). Furthermore, silibinin facilitated the establishment of the antioxidant enzyme functions [64]. Another *in vivo* study

reported that silibinin had an anti-proliferative effect against human colorectal carcinoma HT29 xenograft and an antiangiogenic effect, which is very important in preventing metastasis. The silibinin antiproliferative activity was achieved via down-regulation of ERK1/2 and Akt phosphorylation, as well as the cyclin D1 expression. The silibinin antiangiogenic activity *in vivo* was connected with the inhibition of iNOS, COX-1, COX-2, hypoxia-inducing factor-1 α (HIF-1 α) and VEGF [65]. Therefore, silibinin could be considered as a promising chemo-preventive agent for the treatment of colon cancer.

Effect on pancreatic cancer: Dietary silibinin suppressed the growth of human pancreatic carcinoma BxPC-3 and PANC-1 tumor xenografts (representing early and late clinical stages of the disease, respectively), inhibited the cell proliferation and induced the apoptosis in BxPC-3 tumors in *in vitro* and *in vivo* studies. It caused a strong G1 arrest in BxPC-3 cells in a dose-and/or time-dependent manner and a moderate arrest in PANC-1 cells. The authors assumed that silibinin inhibited the G1/S cell cycle progression via down-regulation of cyclin D1 and CDK4/2 or by up-regulation of CDKIs [66].

Effect on leukemia: Silibinin, in a dose-dependent manner, inhibited the proliferation of the human promyelocytic leukemia HL-60 cell line and induced cell differentiation, which was even more pronounced when combined with 1,25-(OH)₂D₃ vitamin via both protein kinase C α and β level elevation and enhanced the activity [67]. Silymarin strongly inhibited Akt protein levels in K562 leukemia cells, caused caspases activation and PARP cleavage, which inhibited the growth of the human leukemia cells and induced their apoptosis [68]. This was confirmed for both silibinin A and silibinin B on the same cell culture, which showed that the diastereoisomers were more potent than the silibinin racemic mixture [69].

Nephroprotective effect: Silibinin-hemisuccinate (also silicristin and silidianin) stimulated the proliferation of Vero cells similar to fibroblasts (African green monkey, *C. aethiops*) by about 23% more than the control group, increased the synthesis of DNA molecules in the cells, protein synthesis, and reduced the toxic effects of paracetamol. In addition, silibinin had a positive effect on cells chemically damaged by cisplatin and vincristine, but only when it had been administered before these drugs [70]. Silymarin treatment of diabetic nephropathy patients at the final phase caused the normalization of immunoregulatory defects in *in vitro* studies [71]. Silymarin, in a dose-dependent manner, acted preventively in ischemia and reperfusion after renal injury and decreased the morphological changes that included the dilation and vacuolization of tubules, pelvic inflammation, interstitial inflammation, perirenal fat infiltration, and glomerular and tubular necrosis caused by the nephrotoxic drugs: cisplatin, doxorubicin, aminoglycosides, vincristine, cyclosporine and acetaminophen in *in vitro* studies [72].

Neuroprotective effect: Silymarin had a neuroprotective effect in rats with brain damage caused by acetaminophen in an *in vivo* study. In the group of rats treated only with silymarin, increased GSH and ascorbic acid levels and slightly increased SOD levels were observed in relation to the control, while the MDA levels were significantly reduced in the group that had received silymarin three days before paracetamol was administered. It is supposed that silymarin binds nonenzymatic ROS into CNS, or it increases the protein synthesis involved in the antioxidant activity, and, therefore, the authors recommended the use of silymarin in the treatment of neurodegenerative and neurotoxic disorders [73]. Silibinin, administered intragastrically 30 min before a permanent middle

cerebral artery occlusion, significantly alleviated neurological deficit, reduced an infarct volume and suppressed a brain oedema, which was assumed to have been accomplished by up-regulation of pAkt, pmTOR, HIF-1 α , Bcl-2 and by down-regulation of Bax, and NF- κ B in the ischemic brain tissue after a stroke [74]. The pretreatment with silymarin had a statistically significant neuroprotective effect in the *in vivo* studies on the rats with the neurotoxicity caused by Na-fluoride since it increased the activity of the antioxidant systems (SOD, CAT and GSH) [75].

Cardioprotective effect: Fat infiltration in the myocardium of obese db/db mice was induced to correspond to myocardium steatosis of obese and diabetic patients or to patients with non-alcoholic steatosis. Silibinin decreased the ALT levels in the serum three times in comparison with the untreated group and reduced the insulin and glucose levels in the serum during the fasting period causing the elimination of the generated insulin resistance. The treatment with silibinin significantly decreased myocardial injury and mitigated morphological abnormalities of the majority of the myocardiocytes, reduced the damage caused by oxidative stress and maintained the GSH level. Silibinin completely eliminated the increased TNF- α expression and decreased the expression of genes for IL-6, which was observed in the obese mice [76]. Silibinin attenuated the phenylephrine-induced hypertrophic response in H9c2 cardiac cells, up-regulation of atrial natriuretic peptide and the increase of cellular protein levels by blocking ERK1/2 and Akt signaling pathways. Pretreatment of the H9c2 cells with silibinin also protected them from H₂O₂-induced cellular stress [77].

Hypocholesterolemic and hypolipidemic effects: Silymarin had the same effect on serum cholesterol as probucol (an antioxidant hypocholesterolemic drug) in rats on a high-cholesterol diet. Silymarin increased the HDL level mildly, reduced the cholesterol content in the liver and stimulated LDL cholesterol excretion from the liver, which contributed to the anti-atherogenic effect [78]. The results were confirmed in a study in which silymarin dose-dependently reduced total cholesterol, LDL and triglyceride levels in rabbits on a high cholesterol diet and thus inhibited the formation of atherosclerotic plaques [79]. In a four-month randomized double blind study silymarin, administered three times a day (200 mg tablet) with a standard therapy in non-insulin dependent hyperlipidemic patients caused a statistically significant reduction in total cholesterol, LDL and triglyceride levels compared with the control group which received only a standard therapy with placebo [80]. Silibinin-cyclodextrin significantly decreased triglyceride levels in relation to the placebo group in a double-blind six-month-study with patients with poorly controlled non-insulin dependent diabetes [81]. The treatment with silymarin, during one month, significantly reduced total cholesterol, LDL and VLDL levels, and increased HDL levels slightly (silymarin showed the same effect as lovastatin) in the randomized clinical study with 57 patients with dyslipidemia [82].

Effect on diabetes: In a study on Wistar rats, silibinin showed a significant hypoglycemic effect in the liver, inhibiting gluconeogenesis during fasting and glycolysis in satiety, respectively, opening the possibility of the clinical application of silibinin in the treatment of diabetes [8]. The blood glucose level of patients with diabetes mellitus II, in the group who received silymarin, was statistically significantly lower after four months of administration in relation to the placebo group, where the increased glucose level was recorded in a double-blind placebo-controlled clinical study. The patients treated with silymarin, compared with the control group, had a slight, but not statistically significant weight loss and reduction in blood pressure [80].

A long-term treatment with silymarin reduced lipoperoxidation and insulin resistance in the patients with alcohol liver cirrhosis who had similar disease histories and pathological characteristics, like patients with alcoholic liver disease and NASH. Namely, the significant decrease in fasting blood glucose levels, insulin level decrement (40%), reduction of exogenous insulin needs, decrease of mean daily blood glucose levels, reducing daily glycosuria and HbA1c levels, were recorded in the group that received silymarin together with the standard therapy after only 4 months of administration [83]. In addition, the rise in glucose and fatty acid levels in the diabetics led to ROS increment accompanied by insulin resistance, dysfunction of the pancreatic β -cells and insulin secretion – the effects that may be significantly mitigated due to the silymarin antioxidant action [84].

Effect on sepsis: The endotoxins (lipopolysaccharides) that activate macrophages and produce cytokines and lead to oxidative stress are considered to have the key role in sepsis development and multiple body function disorders. Pretreatment with silymarin had the same effect as pretreatment with antioxidant *N*-acetylcysteine (NAC) in a study on rats with induced septic lung and brain damage. The survival rate in the group treated with NAC was the same as in the group treated with silymarin after 72 h. Both silymarin and NAC effectively eliminated the increase of TNF- α , IL-1 and IL-6, LDH and MDA caused by sepsis. Silymarin inhibited the oxidative injury caused by the lungs and brain sepsis owing to its ability to balance the antioxidative status and to regulate the inflammatory mediators [85].

Effects on osteoporosis and osteoarthritis: Seidlova-Wuttke *et al.* were the first to observe that silymarin had a raloxiphen-like effect on bone structure [86]. In another study it was demonstrated that silymarin did not affect the luteinizing hormone (LH) level, or cholesterol, LDL and HDL levels in rats after ovariectomy, but it increased the uterine weight, height and hypertrophy of endometrial luminal epithelium compared with the control group, probably due to partial binding to ER α receptors in the uterus. However, silymarin augmented the density of the trabecular part of the long bones, did not affect the osteocalcin levels, but reduced the alkaline phosphatase levels and increased the calcium and phosphorus levels in the serum and stimulated the secretion of parathyroid hormone 5.5 times more than that with ethinyl estradiol. The potential use of silymarin against osteoporosis development was confirmed in this study, assuming that its effect could not be achieved only through the estrogen receptor, but some other mechanisms of action should be included [87].

The destructive changes of moniodoacetate-provoked osteoarthritis in rats were ameliorated with both celecoxib and silymarin. Celecoxib (100 mg/kg) and silymarin (50 mg/kg) applied separately or in parallel (celecoxib in a dose of 100 mg/kg and silymarin in a dose of 25 mg/kg) lowered the osteoarthritis MDA and NO levels. A combination therapy, containing both celecoxib and silymarin, resulted in IL-1 β reduction in the serum more significantly than the application of each drug separately. Silymarin potentiated the effect of celecoxib, and applied together, they also amended the histopathological findings (fibrillated surface, presence of osteoclasts and connective tissue) in the rats with induced osteoarthritis. The authors proposed further testing of celecoxib and silymarin as a combined osteoarthritis therapy in humans [88].

Preventive effect for fetal alcohol syndrome: Research on the influence of silymarin on fetal alcohol syndrome was made on Fisher female rats with confirmed pregnancies. The hatchlings from the ethanolic group had significantly less mass, less total brain mass

and less mass of the corpus callosum than the control and in the group treated with silymarin, while the female group, along with ethanol was fed with silymarin during the pregnancy, gave birth to offspring of a nearly identical body mass as the control group [89]. In another study, exposure to alcohol during fetal development influenced the spatial orientation of the adult rats, and the spatial memory impairment was more pronounced in the female rats, but was significantly lower in the group pretreated with silymarin. It is assumed that silymarin scavenged the free radicals produced by ethanol oxidation thus preventing the potential negative effects on the migration of the cells, then, it acted on the hippocampus stimulating the DNA and proteins synthesis, maintaining the number of cerebral Purkinje cells, or it acted through the NMDA receptors [90].

Prolactive effect: Female rats were treated either with a standardized extract (silymarin BIO-C® = Pilütatte® - Sil®) or with metoclopramide intraperitoneally (the control). Sil® caused a statistically significant increase in body weight and serum prolactin levels in the female rats compared with the control. After interruption of the treatment with Sil® after 66 days, the prolactin level was still elevated. Bromocriptine, an agonist of D2-dopamine receptors, dose-dependently, significantly reduced the prolactin levels in the serum induced by Sil® treatment. It is clear that silymarin increased the circulating prolactin levels in the female rats and that the dopamine D2 receptors were probably at least partially involved in this effect [91].

Immunomodulatory effect: Silymarin modulated CD4⁺ splenocytes proliferation in mice activated α CD3 mAb (anti-mouse CD3 monoclonal antibody) in the *in vitro* study, but did not have a direct mitogenic activity and cytotoxic effects. Silymarin, at the concentration of 50 μ M, significantly inhibited the CD3-induced NF- κ B nuclear translocation and production of IL-2 and IFN- γ in the activated splenocytes after 72 h in relation to the untreated cells and showed immunomodulatory potential [92]. Silibinin exerted anti-inflammatory and anti-fibrotic effects on the CD14⁺ cells by NF- κ B mediated inhibition of TNF- α , IL-10, TNF- β 1, PGE₂ and NO production [93]. The down-regulation of NF- κ B, induced by pretreatment with silibinin, prevented mouse-ovalbumin induced allergic airway inflammation in the *in vivo* study [94]. Silibinin inhibited the leukotriene formation significantly by Kupffer cells *in vitro*, inhibiting the 5-lipo-oxygenase pathway [95], which is

consistent with the studies indicating that silymarin and its constituent silibinin inhibited arachidonic acid metabolism *in vitro* [96, 97].

Effect on burns: The burns caused by hot water in Wistar rats caused severe oxidative stress and tissue damage. The dermal or dermal/oral application of silymarin restored the increased LDH and MDA levels and decreased the GSH levels to the values of the control group. When the burns were treated with silymarin, the higher levels of TNF- α were reduced compared with the group without the silymarin treatment. Myeloperoxidase activity, which indicated the increased infiltration of neutrophils in the damaged tissue, was significantly increased 48 h after the burns had been caused, but was fully restored to the level of the control group after the silymarin treatment. Silymarin eliminated the increased thromboplastin activity observed in the burns and reduced the oxidative damage of the epidermis and dermis caused by heat [98].

Effect on cholestasis: It was shown that silibinin had a therapeutic application in cholestasis because it affected the transport of tauro- and glyco-conjugated bile salts through the canalicular membrane in isolated hepatocytes of Wistar rats. Silibinin possessed an anti-cholelithic effect on estradiol-17 β -D-glucuronide and taurochenodeoxycholate, and induced cholestasis by increasing the level of cytosolic Ca²⁺ through cAMP, which opened the possibility for further research of the silibinin activity in the prevention of gallstone formation [99].

SIL effect in cosmetics: Cosmetic products containing silymarin to treat rosacea and to maintain the skin moist and lips smooth are already on the market. As the aging processes are accompanied by oxidative stress, SIL is expected to be used in cosmetic preparations against wrinkles and aging skin. The ability of silymarin to protect the skin from epidermal hyperplasia and DNA epidermal cell damage caused by UVB radiation makes SIL appropriate and benefitting to be used in cosmetics preparations with a UV protection factor in sunscreens [100].

Acknowledgments - This article has been prepared within the project TR31029 "Functional products based on cereals for persons with metabolic disorders", funded by the Ministry of Education, Science and Technological Development of the Republic of Serbia.

References

- [1] Kren V, Walterová D. (2005) Silybin and silymarin-new effects and applications. *Biomedical Papers of the Medical Faculty of the University Palacký, Olomouc, Czechoslovakia*, **149**, 29-41.
- [2] Kroll DJ, Shaw HS, Oberlies NH. (2007) Milk thistle nomenclature: Why it matters in cancer research and pharmacokinetic studies. *Integrative Cancer Therapies*, **6**, 110-119.
- [3] Abenavoli L, Capasso R, Milic N, Capasso F. (2010) Milk thistle in liver diseases: past, present and future. *Phytotherapy Research*, **24**, 1423-1432.
- [4] Abenavoli L, Aviello G, Capasso R, Milic N, Capasso F. (2011) Milk thistle for treatment of nonalcoholic fatty liver disease. *Hepatitis Monthly*, **11**, 173-177.
- [5] Tzeng JI, Chen MF, Chung HH, Cheng JT. (2013) Silymarin decreases connective tissue growth factor to improve liver fibrosis in rats treated with carbon tetrachloride. *Phytotherapy Research*, **27**, 1023-1028.
- [6] Kwon do Y, Jung YS, Kim SJ, Kim YS, Choi DW, Kim YC. (2013) Alterations in sulfur amino acid metabolism in mice treated with silymarin: a novel mechanism of its action involved in enhancement of the antioxidant defense in liver. *Planta Medica*, **79**, 997-1002.
- [7] Pradhan SC, Girish C. (2006) Hepatoprotective herbal drug, silymarin from experimental pharmacology to clinical medicine. *The Indian Journal of Medical Research*, **124**, 491-504.
- [8] Colturato CP, Constantin RP, Maeda AS Jr, Constantin RP, Yamamoto NS, Bracht A, Ishii-Iwamoto EL, Constantin J. (2012) Metabolic effects of silibinin in the rat liver. *Chemico-biological Interactions*, **195**, 119-132.
- [9] Shaker E, Mahmoud H, Mnaa S. (2010) Silymarin, the antioxidant component and *Silybum marianum* extracts prevent liver damage. *Food and chemical toxicology: an international journal published for the British Industrial Biological Research Association*, **48**, 803-806.
- [10] Parés A, Planas R, Torres M, Caballería J, Viver JM, Acero D, Panés J, Rigau J, Santos J, Rodés J. (1998) Effects of silymarin in alcoholic patients with cirrhosis of the liver: results of a controlled, double-blind, randomized and multicenter trial. *Journal of Hepatology*, **28**, 615-621.
- [11] Mayer KE, Myers RP, Lee SS. (2005) Silymarin treatment of viral hepatitis: a systematic review. *Journal of Viral Hepatitis*, **12**, 559-567.

- [12] Kim YC, Kim EJ, Lee ED, Kim JH, Jang SW, Kim YG, Kwon JW, Kim WB, Lee MG. (2003) Comparative bioavailability of silibinin in healthy male volunteers. *International Journal of Clinical Pharmacology and Therapeutics*, **41**, 593–596.
- [13] Wu JW, Lin LC, Tsai TH. (2009) Drug–drug interactions of silymarin on the perspective of pharmacokinetics. *Journal of Ethnopharmacology*, **121**, 185–193.
- [14] Ghosh A, Ghosh T, Jain S. (2010) Silymarin- a review on the pharmacodynamics and bioavailability enhancement approaches. *Journal of Pharmaceutical Science and Technology*, **2**, 348–355.
- [15] Wen Z, Dumas TE, Schrieber SJ, Hawke RL, Fried MW, Smith PC. (2008) Pharmacokinetics and metabolic profile of free, conjugated, and total silymarin flavonolignans in human plasma after oral administration of milk thistle extract. *Drug Metabolism and Disposition: The Biological Fate of Chemicals*, **36**, 65–72.
- [16] Calani L, Brighenti F, Bruni R, Del Rio D. (2012) Absorption and metabolism of milk thistle flavanolignans in humans. *Phytomedicine*, **20**, 40–46.
- [17] Post-White J, Ladas EJ, Kelly KM. (2007) Advances in the Use of Milk Thistle (*Silybum marianum*). *Integrative Cancer Therapies*, **6**, 104–109.
- [18] Brantley SJ, Oberlies NH, Kroll DJ, Paine MF. (2010) Two flavonolignans from milk thistle (*Silybum marianum*) inhibit CYP2C9-mediated warfarin metabolism at clinically achievable concentrations. *The Journal of Pharmacology and Experimental Therapeutics*, **332**, 1081–1087.
- [19] Yin F, Liu J, Ji X, Wang Y, Zidichouski J, Zhang J. (2011) Silibinin: a novel inhibitor of A β aggregation. *Neurochemistry International*, **58**, 399–403.
- [20] Tota S, Kamat PK, Shukla R, Nath C. (2011) Improvement of brain energy metabolism and cholinergic functions contributes to the beneficial effects of silibinin against streptozotocin induced memory impairment. *Behavioural Brain Research*, **221**, 207–215.
- [21] Marrazzo G, Bosco P, La Delia F, Scapagnini G, Di Giacomo C, Malaguarnera M, Galvano F, Nicolosi A, Li Volti G. (2011) Neuroprotective effect of silibinin in diabetic mice. *Neuroscience Letters*, **504**, 252–256.
- [22] Lu P, Mamiya T, Lu L, Mouri A, Niwa M, Kim HC, Zou LB, Nagai T, Yamada K, Ikejima T, Nabeshima T. (2010) Silibinin attenuates cognitive deficits and decreases of dopamine and serotonin induced by repeated methamphetamine treatment. *Behavioural Brain Research*, **207**, 387–393.
- [23] Baluchnejadmojarad T, Roghani M, Mafakheri M. (2010) Neuroprotective effect of silymarin in 6-hydroxydopamine hemi-Parkinsonian rat: Involvement of estrogen receptors and oxidative stress. *Neuroscience Letters*, **480**, 206–210.
- [24] Kaur M, Agarwal R. (2007) Silymarin and epithelial cancer chemoprevention: how close we are to bedside? *Toxicology and Applied Pharmacology*, **224**, 350–359.
- [25] Bhatia N, Zhao J, Wolf DM, Agarwal R. (1999) Inhibition of human carcinoma cell growth and DNA synthesis by silibinin, an active constituent of milk thistle: comparison with silymarin. *Cancer Letters*, **147**, 77–84.
- [26] Zho J, Sharma Y, Agarwal R. (1999) Significant inhibition by the flavonoid antioxidant silymarin against 12-O-tetradecanoylphorbol 13-acetate caused modulation of antioxidant and inflammatory enzymes, and cyclooxygenase 2 and interleukin-1 alpha expression in SENCAR mouse epidermis: Implications in the prevention of stage I tumor promotion. *Molecular Carcinogenesis*, **26**, 321–333.
- [27] Zi X, Agarwal R. (1999) Modulation of mitogen-activated protein kinase activation and cell cycle regulators by the potent skin cancer preventive agent silymarin. *Biochemical and Biophysical Research Communications*, **263**, 528–536.
- [28] Singh RP, Tyagi AK, Zhao J, Agarwal R. (2002) Silymarin inhibits growth and causes regression of established skin tumors in SENCAR mice via modulation of mitogen-activated protein kinases and induction of apoptosis. *Carcinogenesis*, **23**, 499–510.
- [29] Katiyar SK, Korman NJ, Mukhtar H, Agarwal R. (1997) Protective effects of silymarin against photocarcinogenesis in a mouse skin model. *Journal of the National Cancer Institute*, **89**, 556–566.
- [30] Katiyar-Santosh K. (2002) Treatment of silymarin, a plant flavonoid, prevents ultraviolet light-induced immune suppression and oxidative stress in mouse skin. *International Journal of Oncology*, **21**, 1213–1222.
- [31] Pinnell SR. (2003) Cutaneous photodamage, oxidate stress, and topical antioxidant protection. *Journal of the American Academy of Dermatology*, **48**, 1–19.
- [32] Liu W, Otkur W, Li L, Wang Q, He H, Zang L, Hayashi T, Tashiro S, Onodera S, Xia M, Ikejima T. (2013) Interference of silibinin with IGF-1R signalling pathways protects human epidermoid carcinoma A431 cells from UVB-induced apoptosis. *Biochemical and Biophysical Research Communications*, **432**, 314–319.
- [33] Liu W, Otkur W, Li L, Wang Q, He H, Ye Y, Zhang Y, Hayashi T, Tashiro S, Onodera S, Ikejima T. (2013) Autophagy induced by silibinin protects human epidermoid carcinoma A431 cells from UVB-induced apoptosis. *Journal of Photochemistry and Photobiology. B: Biology*, **123**, 23–31.
- [34] Singh RP, Agarwal R. (2005) Mechanisms and preclinical efficacy of silibinin in preventing skin cancer. *European Journal of Cancer*, **41**, 1969–1979.
- [35] Ramakrishnan G, Augustine TA, Jagan S, Vinodhkumar R, Devaki T. (2007) Effect of silymarin on *N*-nitrosodiethylamine induced hepatocarcinogenesis in rats. *Experimental Oncology*, **29**, 39–44.
- [36] Brandon-Warner E, Sugg JA, Schrum LW, McKillop IH. (2010) Silibinin inhibits ethanol metabolism and ethanol-dependent cell proliferation in an *in vitro* model of hepatocellular carcinoma. *Cancer Letters*, **291**, 120–129.
- [37] Momeny M, Khorramizadeh MR, Ghaffari SH, Yousefi M, Yekaninejad MS, Esmaili R, Jahanshahi Z, Nooridaloui MR. (2008) Effects of silibinin on cell growth and invasive properties of a human hepatocellular carcinoma cell line, HepG-2, through inhibition of extracellular signal-regulated kinase 1/2 phosphorylation. *European Journal of Pharmacology*, **591**, 13–20.
- [38] Zi X, Feyes DK, Agarwal R. (1998) Anticarcinogenic effect of a flavonoid antioxidant, silymarin, in human breast cancer cells MDA-MB 468: induction of G1 arrest through an increase in Cip1/p21 concomitant with a decrease in kinase activity of cyclin-dependent kinases and associated cyclins. *Clinical Cancer Research: an official journal of the American Association for Cancer Research*, **4**, 1055–1064.
- [39] Lin ZY, Chuang WL. (2011) Influence of silibinin on differential expressions of total cytokine genes in human hepatocellular carcinoma cell lines. *Biomedicine and Preventive Nutrition*, **1**, 91–94.
- [40] Lee SO, Jeong YJ, Im HG, Kim CH, Chang YC, Lee IS. (2007) Silibinin suppresses PMA-induced MMP-9 expression by blocking the AP-1 activation via MAPK signaling pathways in MCF-7 human breast carcinoma cells. *Biochemical and Biophysical Research Communications*, **354**, 165–171.
- [41] Kim S, Choi JH, Lim HI, Lee SK, Kim WW, Kim JS, Kim JH, Choe JH, Yang JH, Nam SJ, Lee JE. (2009) Silibinin prevents TPA-induced MMP-9 expression and VEGF secretion by inactivation of the Raf/MEK/ERK pathway in MCF-7 human breast cancer cells. *Phytomedicine*, **16**, 573–580.
- [42] Xu P, Yin Q, Shen J, Chen L, Yu H, Zhang Z, Li Y. (2013) Synergistic inhibition of breast cancer metastasis by silibinin-loaded lipid nanoparticles containing TPGS. *International Journal of Pharmaceutics*, **454**, 21–30.
- [43] Lu W, Lin C, King TD, Chen H, Reynolds RC, Li Y. (2012) Silibinin inhibits Wnt/ β -catenin signaling by suppressing Wnt co-receptor LRP6 expression in human prostate and breast cancer cells. *Cellular Signalling*, **24**, 2291–2296.
- [44] Gallo D, Giacomelli S, Ferlini C, Raspaglio G, Apollonio P, Prislei S, Riva A, Morazzoni P, Bombardelli E, Scambia G. (2003) Antitumour activity of the silybin-phosphatidylcholine complex, IdB 1016, against human ovarian cancer. *European Journal of Cancer*, **39**, 2403–2410.
- [45] Zhou L, Liu P, Chen B, Wang Y, Wang X, Chiriva Internati M, Wachtel MS, Frezza EE. (2008) Silibinin restores paclitaxel sensitivity to paclitaxel-resistant human ovarian carcinoma cells. *Anticancer Research*, **28**, 1119–1127.

- [46] Huang Q, Wu LJ, Tashiro S, Onodera S, Li LH, Ikejima T. (2005) Silymarin augments human cervical cancer HeLa cell apoptosis via P38/JNK-MAPK pathways in serum-free medium. *Journal of Asian Natural Products Research*, **7**, 701-709.
- [47] Garcia-Maceira P, Mateo J. (2009) Silibinin inhibits hypoxia-inducible factor-1 α and mTOR/p70S6K/4E-BP1 signalling pathway in human cervical and hepatoma cancer cells: implications for anticancer therapy. *Oncogene*, **28**, 313-324.
- [48] Tyagi A, Agarwal C, Harrison G, Glode LM, Agarwal R. (2004) Silibinin causes cell cycle arrest and apoptosis in human bladder transitional cell carcinoma cells by regulating CDKI-CDK-cyclin cascade, and caspase 3 and PARP cleavages. *Carcinogenesis*, **25**, 1711-1720.
- [49] Vinh PQ, Sugie S, Tanaka T, Hara A, Yamada Y, Katayama M, Deguchi T, Mori H. (2002) Chemopreventive effects of a flavonoid antioxidant silymarin on N-butyl-N-(4-hydroxybutyl)nitrosamine-induced urinary bladder carcinogenesis in male ICR mice. *Japanese Journal of Cancer Research: Gann*, **93**, 42-49.
- [50] Singh RP, Tyagi A, Sharma G, Mohan S, Agarwal R. (2008) Oral silibinin inhibits *in vivo* human bladder tumor xenograft growth involving down-regulation of survivin. *Clinical Cancer Research: an official journal of the American Association for Cancer Research*, **14**, 300-308.
- [51] Singh RP, Mallikarjuna GU, Sharma G, Dhanalakshmi S, Tyagi AK, Chan DC, Agarwal C, Agarwal R. (2004) Oral silibinin inhibits lung tumor growth in athymic nude mice and forms a novel chemocombination with doxorubicin targeting NF-kB-mediated inducible chemoresistance. *Clinical Cancer Research: an official journal of the American Association for Cancer Research*, **10**, 8641-8647.
- [52] Sadava D, Kane SE. (2013) Silibinin reverses drug resistance in human small-cell lung carcinoma cells. *Cancer Letters*, **339**, 102-106.
- [53] Singh RP, Deep G, Chittezhath M, Kaur M, Dwyer-Nield LD, Malkinson AM, Agarwal R. (2006) Effect of silibinin on the growth and progression of primary lung tumors in mice. *Journal of the National Cancer Institute*, **98**, 846-855.
- [54] Chittezhath M, Deep G, Singh RP, Agarwal C, Agarwal R. (2008) Silibinin inhibits cytokine-induced signaling cascades and down-regulates inducible NO synthase in human lung carcinoma A549 cells. *Molecular Cancer Therapeutics*, **7**, 1817-1826.
- [55] Cufi S, Bonavia R, Vazquez-Martin A, Corominas-Faja B, Oliveras-Ferreros C, Cuyàs E, Martín-Castillo B, Barrajón-Catalán E, Visa J, Segura-Carretero A, Bosch-Barrera J, Joven J, Micol V, Menendez JA. (2013) Silibinin meglumine, a water-soluble form of milk thistle silymarin, is an orally active anti-cancer agent that impedes the epithelial-to-mesenchymal transition (EMT) in EGFR-mutant non-small-cell lung carcinoma cells. *Food and Chemical Toxicology: an international journal published for the British Industrial Biological Research Association*, **60**, 360-368.
- [56] Davis-Searles PR, Nakanishi Y, Kim NC, Graf TN, Oberlies NH, Wani MC, Wall ME, Agarwal R, Kroll DJ. (2005) Milk thistle and prostate cancer: differential effects of pure flavonolignans from *Silybum marianum* on antiproliferative end points in human prostate carcinoma cells. *Cancer Research*, **65**, 4448-4457.
- [57] Singh RP, Agarwal R. (2004) A cancer chemopreventive agent silibinin, targets mitogenic and survival signaling in prostate cancer. *Mutation Research*, **555**, 21-32.
- [58] Verschoyle RD, Greaves P, Patel K, Marsden DA, Brown K, Steward WP, Gescher AJ. (2008) Evaluation of the cancer chemopreventive efficacy of silibinin in genetic mouse models of prostate and intestinal carcinogenesis: Relationship with silibinin levels. *European Journal of Cancer*, **44**, 898-906.
- [59] Raina K, Blouin MJ, Singh RP, Majeed N, Deep G, Varghese L, Glodé LM, Greenberg NM, Hwang D, Cohen P, Pollak MN, Agarwal R. (2007) Dietary feeding of silibinin inhibits prostate tumor growth and progression in transgenic adenocarcinoma of the mouse prostate model. *Cancer Research*, **67**, 11083-11091.
- [60] Su CH, Chen LJ, Liao JF, Cheng JT. (2013) Increase of phosphatase and tensin homolog by silymarin to inhibit human pharynx squamous cancer. *Journal of Medicinal Food*, **16**, 778-784.
- [61] Zhang Y, Li Q, Ge Y, Chen Y, Chen J, Dong Y, Shi W. (2013) Silibinin triggers apoptosis and cell-cycle arrest of SGC7901 cells. *Phytotherapy Research*, **27**, 397-403.
- [62] Hogan FS, Krishnegowda NK, Mikhailova M, Kahlenberg MS. (2007) Flavonoid, silibinin, inhibits proliferation and promotes cell-cycle arrest of human colon cancer. *The Journal of Surgical Research*, **143**, 58-65.
- [63] Kohno H, Tanaka T, Kawabata K, Hirose Y, Sugie S, Tsuda H, Mori H. (2002) Silymarin, a naturally occurring polyphenolic antioxidant flavonoid, inhibits azoxymethane-induced colon carcinogenesis in male F344 rats. *International Journal of Cancer*, **101**, 461-468.
- [64] Sangeetha N, Viswanathan P, Balasubramanian T, Nalini N. (2012) Colon cancer chemopreventive efficacy of silibinin through perturbation of xenobiotic metabolizing enzymes in experimental rats. *European Journal of Pharmacology*, **674**, 430-438.
- [65] Singh RP, Gu M, Agarwal R. (2008) Silibinin inhibits colorectal cancer growth by inhibiting tumor cell proliferation and angiogenesis. *Cancer Research*, **68**, 2043-2050.
- [66] Nambiar D, Prajapati V, Agarwal R, Singh RP. (2012) *In vitro* and *in vivo* anticancer efficacy of silibinin against human pancreatic cancer BxPC-3 and PANC-1 cells. *Cancer Letters*, pii: S0304-3835(12)00539-3. doi: 10.1016/j.canlet.2012.09.004.
- [67] Kang SN, Lee MH, Kim KM, Cho D, Kim TS. (2001) Induction of human promyelocytic leukemia HL-60 cell differentiation into monocytes by silibinin: involvement of protein kinase C. *Biochemical Pharmacology*, **61**, 1487-1495.
- [68] Zhong X, Zhu Y, Lu Q, Zhang J, Ge Z, Zheng S. (2006) Silymarin causes caspases activation and apoptosis in K562 leukemia cells through inactivation of Akt pathway. *Toxicology*, **227**, 211-216.
- [69] Zhang J, Luan Q, Liu Y, Lee DY, Wang Z. (2011) A comparison of the diastereoisomers, silybin A and silybin B, on the induction of apoptosis in K562 cells. *Natural Product Communications*, **6**, 1653-1656.
- [70] Sonnenbichler J, Scalera F, Sonnenbichler I, Weyhenmeyer R. (1999) Stimulatory effects of silibinin and silicristin from the milk thistle *Silybum marianum* on kidney cells. *The Journal of Pharmacology and Experimental Therapeutics*, **290**, 1375-1383.
- [71] Dietzmann J, Thiel U, Ansorge S, Neumann KH, Tager M. (2002) Thiol-inducing and immunoregulatory effects of flavonoids in peripheral blood mononuclear cells from patients with end-stage diabetic nephropathy. *Free Radical Biology and Medicine*, **33**, 1347-1354.
- [72] Dashti-Khavidaki S, Shahbazi F, Khalili H, Lessan-Pezeshki M. (2012) Potential renoprotective effects of silymarin against nephrotoxic drugs: a review of literature. *Journal of Pharmaceutical Sciences*, **15**, 112-123.
- [73] Nencini C, Giorgi G, Micheli L. (2007) Protective effect of silymarin on oxidative stress in rat brain. *Phytomedicine*, **14**, 129-135.
- [74] Wang C, Wang Z, Zhang X, Zhang X, Dong L, Xing Y, Li Y, Liu Z, Chen L, Qiao H, Wang L, Zhu C. (2012) Protection by silibinin against experimental ischemic stroke: up-regulated pAkt, pmTOR, HIF-1 α and Bcl-2, down-regulated Bax, NF-kB expression. *Neuroscience Letters*, **529**, 45-50.
- [75] Nabavi SM, Sureda A, Nabavi SF, Latifi AM, Moghaddam AH, Hellio C. (2012) Neuroprotective effects of silymarin on sodium fluoride-induced oxidative stress. *Journal of Fluorine Chemistry*, **142**, 79-82.
- [76] Salamone F, Galvano F, Marino A, Paternostro C, Tibullo D, Buccieri F, Mangiameli A, Parola M, Bugianesi E, Li Volti G. (2012) Silibinin improves hepatic and myocardial injury in mice with nonalcoholic steatohepatitis. *Digestive and Liver Disease: Official Journal of the Italian Society of Gastroenterology and the Italian Association for the Study of the Liver*, **44**, 334-342.
- [77] Anestopoulos I, Kavo A, Tentes I, Kortsaris A, Panayiotidis M, Lazou A, Pappa A. (2013) Silibinin protects H9c2 cardiac cells from oxidative stress and inhibits phenylephrine-induced hypertrophy: potential mechanisms. *The Journal of Nutritional Biochemistry*, **24**, 586-594.

- [78] Krecman V, Skottová N, Walterová D, Ulrichová J, Simánek V. (1998) Silymarin inhibits the development of diet-induced hypercholesterolemia in rats. *Planta Medica*, **64**, 138-142.
- [79] Radjabian T, Huseini HF. (2010) Anti-hyperlipidemic and anti-atherosclerotic activities of silymarins from cultivated and wild plants of *Silybum marianum* L. with different contents of flavonolignans. *Iranian Journal of Pharmacology and Therapeutics*, **9**, 63-67.
- [80] Huseini HF, Larijani B, Heshmat R, Fakhrzadeh H, Radjabipour B, Toliat T, Raza M. (2006) The efficacy of *Silybum marianum* (L.) Gaertn. (silymarin) in the treatment of type II diabetes: a randomized, double-blind, placebo-controlled, clinical trial. *Phytotherapy Research*, **20**, 1036-1039.
- [81] Lirussi F, Beccarello A, Zanette G, De Monte A, Donadon V, Velussi M, Crepaldi G. (2002) Silybin-beta-cyclodextrin in the treatment of patients with diabetes mellitus and alcoholic liver disease. Efficacy study of a new preparation of an anti-oxidant agent. *Diabetes, Nutrition and Metabolism*, **15**, 222-231.
- [82] Mesheimish BAR, Hussain SAR, Ismail SH, Hussein KI, Sulaiman AA. (2007) Hypolipidemic effect of silymarin in dyslipidaemia of different etiologies. *Journal of the Faculty of Medicine, Baghdad*, **49**, 449-456.
- [83] Velussi M, Cernigoi AM, De Monte A, Dapas F, Caffau C, Zilli M. (1997) Long-term (12 months) treatment with an anti-oxidant drug (silymarin) is effective on hyperinsulinemia, exogenous insulin need and malondialdehyde levels in cirrhotic diabetic patients. *Journal of Hepatology*, **26**, 871-879.
- [84] Abenavoli L, Milic N, Capasso F. (2012) Anti-oxidant therapy in non-alcoholic fatty liver disease: the role of silymarin. *Endocrine*, **42**, 754-755.
- [85] Toklu HZ, Tunali Akbay T, Velioglu-Ogunc A, Ercan F, Gedik N, Keyer-Uysal M, Sener G. (2008) Silymarin, the antioxidant component of *Silybum marianum*, prevents sepsis-induced acute lung and brain injury. *The Journal of Surgical Research*, **145**, 214-222.
- [86] Seidlová-Wuttke T, Becker V, Christoffel H, Jarry W, Wuttke D. (2003) Silymarin is a selective estrogen receptor β (ER β) agonist and has estrogenic effects in the metaphysis of the femur but no or antiestrogenic effects in the uterus of ovariectomized (ovx) rats. *Journal of Steroid Biochemistry & Molecular Biology*, **86**, 179-188.
- [87] El-Shitany NA, Hegazy S, El-Desoky K. (2010) Evidences for antiosteoporotic and selective estrogen receptor modulator activity of silymarin compared with ethinylestradiol in ovariectomized rats. *Phytomedicine*, **17**, 116-125.
- [88] Ashkavand Z, Malekinejad H, Amniatlab A, Rezaei-Golmisheh A, Vishwanath BS. (2012) Silymarin potentiates the anti-inflammatory effects of Celecoxib on chemically induced osteoarthritis in rats. *Phytomedicine*, **19**, 1200-1205.
- [89] Moreland N, La Grange L, Montoya R. (2002) Impact of in utero exposure to EtOH on corpus callosum development and paw preference in rats: protective effects of silymarin. *BMC Complementary and Alternative Medicine*, **2**, 10.
- [90] Neese S, La Grange L, Trujillo E, Romero D. (2004) The effects of ethanol and silymarin treatment during gestation on spatial working memory. *BMC Complementary and Alternative Medicine*, **4**, 4.
- [91] Capasso R, Aviello G, Capasso F, Savino F, Izzo AA, Lembo F, Borrelli F. (2004) Silymarin BIO-C[®], an extract from *Silybum marianum* fruits, induces hyperprolactinemia in intact female rats. *Phytomedicine*, **16**, 839-844.
- [92] Gharagozloo M, Velardi E, Bruscoli S, Agostini M, Di Sante M, Donato V, Amirghofran Z, Riccardi C. (2010) Silymarin suppress CD4⁺ T cell activation and proliferation: Effects on NF-kB activity and IL-2 production. *Pharmacological Research: the Official Journal of the Italian Pharmacological Society*, **61**, 405-409.
- [93] Bannwart CF, Nakaira-Takahagi E, Golim MA, Medeiros LTL, Romão M, Weel IC, Peraçoli MT. (2010) Downregulation of nuclear factor kappa B (NF-kB) pathway by silibinin in human monocytes challenged with *Paracoccidioides brasiliensis*. *Life Sciences*, **86**, 880-886.
- [94] Choi YH, Jin GY, Guo HS, Piao HM, Li Lc, Li GZ, Lin ZH, Yan GH. (2012) Silibinin attenuates allergic airway inflammation in mice. *Biochemical and Biophysical Research Communications*, **427**, 450-455.
- [95] Dehmlow C, Erhard J, De Groot H. (1996) Inhibition of Kupffer cell functions as an explanation for the hepatoprotective properties of silybinin. *Hepatology*, **23**, 749-754.
- [96] Dehmlow C, Murawski N, de Groot H. (1996) Scavenging of reactive oxygen species and inhibition of arachidonic acid metabolism by silibinin in human cells. *Life Sciences*, **58**, 1591-1600.
- [97] Kang JS, Jeon YJ, Park SK, Yang KH, Kim HM. (2004) Protection against lipopolysaccharide-induced sepsis and inhibition of interleukin-1 β and prostaglandin E2 synthesis by silymarin. *Biochemical Pharmacology*, **67**, 175-181.
- [98] Toklu HZ, Tunali-Akbay T, Erkanli G, Yüksel M, Ercan F, Sener G. (2007) Silymarin, the antioxidant component of *Silybum marianum*, protects against burn-induced oxidative skin injury. *Burns*, **33**, 908-916.
- [99] Crocenzi FA, Basiglio CL, Pérez LM, Portesio MS, Pozzi EJ, Roma MG. (2005) Silibinin prevents cholestasis-associated retrieval of the bile salt export pump, Bsep, in isolated rat hepatocyte couplets: Possible involvement of cAMP. *Biochemical Pharmacology*, **69**, 1113-1120.
- [100] Singh RP, Agarwal R. (2009) Cosmeceuticals and silibinin. *Clinics in Dermatology*, **27**, 479-484.

Biomedical Properties of Edible Seaweed in Cancer Therapy and Chemoprevention Trials: A Review

Farideh Namvar^{a,*}, Paridah Md. Tahir^a, Rosfarizan Mohamad^{a,b}, Mahnaz Mahdavi^c, Parvin Abedi^d, Tahereh Fathi Najafi^e, Heshu Sulaiman Rahman^f and Mohammad Jawaaid^a

^aInstitute of Tropical Forestry and Forest Products (INTROP), Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia

^bFaculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia

^cShiraz Branch, Islamic Azad University, Shiraz, Iran

^dReproductive Health Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

^eDepartment of Medicine, Applied Biology Research Center, Mashhad Branch, Islamic Azad University, Mashhad Iran

^fDepartment of Microbiology and Pathology, Faculty of Veterinary Medicine, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia

farideh.namvar@gmail.com

Received: May 15th, 2013; Accepted: September 3rd, 2013

This review article summarizes *in vitro* and *in vivo* experiments on seaweed anticancer activity and seaweed chemical components. Seaweed use in cancer therapy, chemopreventive randomized control trials (RCTs) and quasi-experiments are discussed. The literature reviewed in this article was obtained from various scientific sources and encompasses publications from 2000-2012. Seaweed therapeutic effects were deemed scientifically plausible and may be partially explained by the *in vivo* and *in vitro* pharmacological studies described. Although the mechanisms of action remain unclear, seaweed's anticancer properties may be attributable to its major biologically active metabolites. Much of the seaweed research outlined in this paper can serve as a foundation for explaining seaweed anticancer bioactivity. This review will open doors for developing strategies to treat malignancies using seaweed natural products.

Keywords: Anticancer, Apoptosis, Cancer Therapy, Chemoprevention, Cytotoxicity, Macro algae, Seaweeds.

The number of patients diagnosed with cancer has been increasing each year, and cancer remains a major cause of death. According to the WHO, cancer accounts for approximately 13.3% of deaths worldwide. Rates of occurrence are expected to rise by 50% worldwide over the next 20 years [1]. Individuals diagnosed with cancer struggle with managing symptoms and coping with treatment side effects. Many clinicians and patients turn to alternative or traditional medicine to avoid the unpleasant or dangerous side effects inherent to modern cancer therapies. Recently, chemopreventive strategies using natural products and dietary supplements have become attractive options as potential means of controlling cancer incidence [2]. Gathering information on and analyzing purported traditional medicine practices is an important element of anticancer drug development involving natural products. For thousands of years, humans have utilized certain marine algae for multiple purposes [3]. Seaweed medical use dates to at least 5,000 years ago in ancient China, where the emperor ShenNung listed seaweed as a medicine in his classic pharmacopeia, the Pen Ts'ao [4]. Seaweeds have been used in traditional Chinese medicine to treat goiter, scrofula, swelling and pain of testes, and edema [5]. Additionally, some seaweed species have also been used in modern Chinese medicine to treat arteriosclerosis, skin diseases, high blood pressure, hepatosplenomegaly, neurosis, angina pectoris, acute esophagitis, and chronic bronchitis [6]. Marine algae belong to one of two groups: microalgae or macroalgae (seaweeds). Macroalgae are plant-like organisms. They are classified according to their pigmentation into green (chlorophytes), red (rhodophytes) and brown (phaeophytes) macroalgae.

Known for their richness in polysaccharides, minerals and certain vitamins, seaweeds also contain other bioactive substances, such as proteins, lipids and polyphenols. These chemicals possess potential for use in treating cancer [7], oxidative stress [3,8], inflammation [9], allergy [10], diabetes [11], thrombosis [12], obesity [13], lipidemia [14], hypertensive [15,16] and other degenerative diseases. This review provides an overview of seaweed *in vitro* and *in vivo* investigations, and it summarizes our own research on seaweed components and their anticancer activity. Additionally, seaweed RCTs and quasi-experiments directed toward cancer therapy and prevention are discussed.

The chemical and nutritional contents of seaweeds vary with age, organism, species, and habitat [17]. Seasonal, environmental, and physiological variations also affect the contents [18]. Cultivated seaweed composition is affected by the time of harvest and the processing methods (e.g., drying, cutting, and milling) [19,20]. Typical air-dried seaweed contains 3–47% carbohydrate, 33–75% protein, 1.5–4% lipid, and 10–35% ash [21,22].

Seaweed contains considerably more protein than high-protein legumes, such as soybeans, and the protein content is generally higher in red and green algae (10–47% DM) than in brown algae (3–16%DM) [23]. The quality of a food protein depends on the variety and amounts of essential amino acids that that particular protein contains. Macroalgal proteins contain all essential amino acids; however, variations in specific concentrations occur between phyla [24]. The organoleptic characteristics of algae arise primarily from their free amino acid profile, which in turn depends on the

environment where the algae grow [25]. Some contain important light-harvesting, protein-pigment complexes called phycobiliproteins. When present, these proteins generally constitute 1–10% of the dry algal biomass. Phycobiliproteins are oligomeric proteins containing two chromophore-bearing polypeptides. This broad class of proteins encompasses three protein subclasses that absorb various red and blue wavelengths – phycoerythrins (PEs), phycocyanins and allophycocyanins. PEs are contained in certain red algae, and phycocyanins can be found in certain red and blue-green algae [8]. Phycobiliproteins also fluoresce *in vitro*. This property has enabled them to be used in biotechnological applications. For example, the biomedical diagnostic community employs phycobiliproteins, primarily in immunochemical methods. Moreover, in some countries, phycobiliproteins are used as natural colorants for food and cosmetics (e.g., chewing gum, dairy products, jellies, ice sherbets, lipsticks and eyeliners) [26]. The most commonly used phycobiliprotein is phycoerythrin, which is present in fluorescent immunoassays, fluorescent immunohistochemistry and other methodologies [8].

Recent studies have shown that the antioxidative capacity of phycobiliproteins can be exploited to prevent or treat several chronic diseases [27]. Phycobiliproteins also exhibit anti-inflammatory, neuroprotective, hypocholesterolemic, hepatoprotective, antiviral, antitumor, antiarteriosclerotic, serum lipid-reducing and lipase inhibition activities. Lectins are glycoproteins that have health applications and uses in clinical diagnosis. Within organisms that possess them, lectins function in aggregation and recognition roles [28]. They have been isolated from primitive marine organisms, including fishes, sponges and sea invertebrates [28]. Recently, a new lectin structural family was identified in *Bryothamnion triquetrum*, a red marine alga. The lectin contained 91 amino acids and two disulfide bonds. The amino acid sequence of the *B. triquetrum* lectin was not similar to any known animal or plant lectin structure [3]. Finally, mitogenic and antineoplastic isoagglutinin glycoproteins have been discovered in the red alga *Solieria robusta* [8].

A normal total lipid content in a macroalgae is between 1.5 and 5%; therefore, its contribution as an energy source appears to be low [29]. Generally, seaweed fatty acids are even-numbered, linear chains that possess one or more double bonds [30]. Algae contain higher proportions of saturated and unsaturated fatty acids than terrestrial plants [18]. PUFAs account for almost half of the seaweed lipids. Much of the seaweed PUFA occurs in the form of omega-3 (n-3) and omega-6 (n-6) fatty acids, such as eicosapentaenoic acid (EPA) and arachidonic acid (AA) [31]. In addition to fatty acids, the unsaponifiable fraction of macroalgal lipids contains carotenoids (e.g., fucoxanthin in brown seaweeds; carotene, lutein and violaxanthin in red and green seaweeds), tocopherols, sterols (e.g., fucosterol in brown seaweeds) and terpenoids [32].

Algal polysaccharides differ from terrestrial plant polysaccharides, and polysaccharide type varies greatly between algal species (Table 1). Although their carbohydrate content is relatively high (33–75% DM), algae are not energy-rich foods because the digestibility of these carbohydrates is low [33]. Most algal polysaccharides simply pass through the human gastrointestinal tract as dietary fiber [34]. “Dietary fibers” are a wide range of complex materials, and numerous definitions have been proposed for this term. Dietary fibers are classified as soluble (SDF) or insoluble (IDF) according to their water solubility. The total dietary fiber content of seaweeds ranges from 29.3–62.3 g/100 g [35], which is higher than the fiber content of most fruits and vegetables. The typical Asian individual’s algal intake of 8 g/day satisfies up to 12.5% of the daily recommended dietary fiber requirement [31].

Table 1: Carbohydrate content of seaweeds

Division	IDF	SDF
Phaeophyta (Brown A.)	-Structural cellulose -Insoluble alginates	-Alginate (10–40%) -Fucans (3.6 to 20%)/ -Fucoidans -Ascophyllans -Glycuronogalactofucans -Neutral glucans (up to 30%) -Laminarans -Mannitol
Rhodophyta (Red A.)	-Cellulose -Residual polysaccharides	-Soluble xylans -Sulfated galactans -Porphyrans -Agar (mixture of sulfated galactans, D-galactose and 3,6-anhydro-L-lactose) -Carrageenan (linear sulfated galactans)
Chlorophyta (Green A.)	-Glucosylan -Glucuronomannan, Amorphous cellulose uronic acids. Rhamnose, xylose, galactose, arabinose	-Ulvan (approximately 8–29% DM)

IDF: insoluble dietary fiber; SDF: soluble dietary fiber

The health benefits of algal fiber consumption are well documented [2]. Polysaccharide composition is a principal characteristic used in classifying algae. Algae contain three types of polysaccharides: structural cell wall polysaccharides, intercellular mucilage polysaccharides, and storage polysaccharides. Storage polysaccharides, such as carrageenans and alginates, are common to seaweeds and are the most commercially exploited seaweed components. Storage polysaccharides exhibit textural and stabilizing properties that are useful in several food applications [31].

Found extensively in the cell walls of macroalgae, fucoidans are a class of sulfated, fucose-rich polymers. Fucoidans are a major therapeutic component of brown algae. Depending on the specific seaweed species, these polymers constitute up to 25–30% of algal dry weight [36]. Fucoidans display numerous bioactive properties, including anticoagulant, antiviral, antibacterial [37], antithrombotic, anti-inflammatory, antitumor and antioxidant activities [38]. Fucoidans can be easily extracted using either hot water or acidic solutions. In addition, the therapeutic potential of fucoidans increases with the degree of sulfation [39].

Fucoidans vary extensively in their polysaccharide structures. Every fucoidan-producing species possesses multiple types of fucoidan that differ in their monosaccharide proportions. In addition, brown algae synthesize species-specific fucoidan sugar compositions. Fucoidan structure and biological significance has been studied [38,40,41]. Fucoidans of various molecular weights and of different origins (and also other seaweed contents) have exhibited a marked anticancer effect *in vitro* and *in vivo*, whether they are administered orally or injected directly into the blood or peritoneal cavity (Table 2).

A significant reduction in the cancer development was observed in mice and rats after cancer cells had been implanted [41,49]. Similar effects were observed in several other cancer animal models, including leukemia and breast cancer [50]. The phenomenon was attributed to the stimulation of both innate and specific immunity. Isolated fucoidans from brown algae have also been shown to possess broad immunomodulating effects. In human studies, fucoidan has been shown to stimulate the enzyme granzyme A, which is believed to induce cytokine production in a variety of cells [51]. This observation may partially explain the mechanism of

fucoidan immunomodulation [49,52]. In another human study involving fucoidan ingestion, low serum levels of fucoidans coincided with the presence of stromal cell-derived factor 1 and increased chemokine receptor CXCR4-expressing hematopoietic stem cells and chemokine IFN- γ [52].

Table 2: Bioactivity of seaweed components.

Characteristic	Bioactivity	Study design	Ref.
Total polysaccharide	Antitumor	<i>In vitro</i> (HepG2, A549 and MGC-803 cells)	[42]
Carrageenan	Antitumor immunomodulation	<i>In vivo</i> (Mice)	[43]
Agar	Antitumor	<i>In vivo</i> (inhibited the transplantation of Ehrlich ascites carcinoma in mice)	[43]
Laminaran	Stimulates immune systems; B and helper T cell	<i>In vivo</i>	[44]
Porphyran	Potentially apoptotic	<i>In vitro</i>	[45]
Carotenoids	Cancer prevention	<i>In vitro</i> - <i>In vivo</i>	[46]
Fucoanthin	Apoptotic	<i>In vitro</i> (human promyelocytic leukemia cells)	[47]
Peptides	Antitumor	<i>In vitro</i> - <i>In vivo</i>	[48]




Figure 1: Flowchart representing the mechanism of action of fucoxanthin antitumor initiation, promotion and progression. VEGF: vascular endothelial growth factor; FGF: fibroblast growth factor; HPA: hypothalamic-pituitary-adrenal; NK: natural killer cell; COX-2: cyclooxygenase-2; iNOS: inducible nitric oxide synthase; NF κ B: nuclear factor kappa B; cFLIP: cellular form of FLICE inhibitory protein; Bcl2: B-cell lymphoma; ROS: reactive oxygen species; and RNS: reactive nitrogen species.

Large, controlled studies have been proposed to evaluate fucoidans therapeutic effects for inoperable, advanced or recurrent colorectal cancer [49]. Strong archival support documents the promising bioactivities of fucoidan. Researchers hope to use fucoidans for treating thrombosis and some forms of cancer [37]. Wijesekara *et al.* [53] reported that algal sulfated polysaccharides possess potential for new pharmaceutical and food anticancer products.

Fucoxanthin is an orange carotenoid found in edible brown seaweeds. This carotenoid exhibits anticarcinogenic effects, apoptotic induction of cancer cells, anti-inflammatory effects, and radical-scavenging activity [23,54] (Figure 1). Additionally, fucoxanthin significantly inhibits numerous enzymes, e.g., matrix metalloproteinases, hyaluronidases and elastases [55]. This inhibitory activity limits tissue breakdown in inflammatory settings caused by injury and disease; moreover, the activity even inhibits metastasis [56].

Laminaran (or laminarin) is a water-soluble, approximately 5000 Da polysaccharide containing 20-25 glucose units [57] and is the second most numerous storage glucan in brown algae. Most

laminarans are dietary fibers because they form complex structures that are stabilized by inter-chain hydrogen bonds and are resistant to hydrolysis in the upper gastrointestinal tract (GIT). Laminaran modulates intestinal metabolism by affecting mucus composition, intestinal pH and short-chain fatty acid production. In addition to their role as dietary fibers, laminarans function as prebiotics and possess antibacterial and antitumor activities [58]. Environmental factors such as water temperature, salinity, waves, sea currents and depth affect the structure and biological activities of laminaran [57]. The antimetastatic effects of laminaran sulfate have been investigated [57,59]. Laminaran sulfate inhibited heparanase activity in mouse B16-BL6 melanoma cells and rat 13762 MAT mammary adenocarcinoma cells. Consequently, the tumor cells' ability to degrade heparan sulfate in their extracellular matrixes was reduced, and an antimetastatic effect was, therefore, produced [59].

At the effective concentrations, laminaran sulfate exerted a small effect on tumor cell proliferation and primary tumor growth *in vivo*. These results further illustrate heparanase involvement in tumor metastasis, and the study supports potential clinical applications of laminaran sulfate in cancer treatment [59].

Alginic acid, or alginate, is the common name given to a family of linear polysaccharides produced by brown seaweed [60]. Alginates are available commercially in both acid and salt (sodium and calcium) forms. The acid form is a linear polyuronic acid referred to as alginic acid. The salt form is an important cell wall component in all brown seaweed, constituting up to 40 - 47 % of dry algal biomass. Alginate is widely used in the food and pharmaceutical industries because of its ability to chelate metal ions and to form highly viscous solutions [57]. Alginate possesses diverse bioactivities: anticoagulative [61], antitumor [62], antiviral, anti-hypertension and antioxidant [53]. Alginate protects against carcinogens by clearing them from the digestive system and protecting stomach and intestinal surface membranes from the effects of carcinogens [50]. Intestinal enzymes cannot digest alginic acid. Consequently, heavy metals are gelated or rendered insoluble by alginic acid in the intestines and cannot be absorbed into the body [58]. In several countries, the use of alginic acid and its derivatives for gastritis, gastro duodenal ulcer and anti-ulcer treatment is patent protected. Clinical trials have shown that sodium alginate promotes the regeneration of stomach mucous membranes, suppresses inflammation, and eradicates mucous membrane colonies of *Helicobacter pylori* [63]. Additionally, alginate promotes the restoration of intestinal biogenesis [8]. Other studies show effects on fecal microbial fauna, changes in compound and acid concentrations, and prebiotic properties that can promote health, particularly in regard to cancer prevention [64,65].

Carrageenans are a family of linear, hydrophilic, sulfated galactans that are obtained from different Rhodophyta species [13]. Carrageenans are traditionally split into six categories based on their chemical composition and commercial use [71]. Because of their sulfate group concentration, all carrageenan fractions are soluble in water and insoluble in organic solvents, oil and fats [13]. Carrageenans are classic agents for inducing inflammation and inflammatory pain in experiments [72]. From a human health perspective, carrageenans exhibit several promising pharmaceutical properties [62], including antitumor [73], immunomodulatory [74], antihyperlipidemic [13,75,76], anticoagulative [77] and antiviral activities [78]. Recently, researchers confirmed that carrageenan is an extremely potent infection inhibitor of a broad range of human genital papillomaviruses (HPVs); therefore, carrageenan-based sexual lubricant gels may offer protection against HPV transmission [13,52,58,79].

At concentrations approximately 1000 times higher than those required to inhibit papilloma viruses, carrageenans inhibited HIV [78,80]. Carrageenans are approved and widely used as a food additive [71,81]. In the food industry, they are employed as a thickener, emulsifier, and stabilizer [10,82]. The food products that contain carrageenan include chocolate milk, ice cream, whipped cream, dietetic foods, low-fat processed meats, infant formula, beer, and salad dressings. Carrageenans' unique ability to combine with milk proteins improves the texture of many food products [6,13,76]. The safety of carrageenans as food additives have been investigated by many researchers. In a review of 45 studies using animal models (rats, mice, rabbits and Guinea pigs), low molecular weight (degraded) carrageenans were associated with the induction and proliferation of intestinal neoplasms and ulcerations [76]. These results warn of potential problems associated with human carrageenan consumption, such as inflammatory bowel disease and gastrointestinal malignancies [13,36,83]. One research group performed a time-trend analysis to see if carrageenan use correlated with the incidence of mammary carcinoma in twentieth century United States. This analysis revealed that the increasing consumption of several alginate gums correlated positively with increased incidences of breast carcinoma [81].

Mannitol is an important sugar alcohol produced by photosynthesis [84]. This sugar alcohol occurs naturally in many brown algal species, accounting for 20–30% of the dry weight of some *Laminaria* species [60,85]. Mannitol is only 70% as sweet as sugar, but has a low glycemic index rating because it cannot be metabolized by intestinal cells [86]. Mannitol is commonly used in chewing gum and mints. The mannitol content of a seaweed varies widely with season and, particularly, environment. The applications of this sugar alcohol are extremely diverse. Because of its hydrating and antioxidant properties, mannitol is used in numerous cosmetic and pharmaceutical products [60,70,86]. In addition, mannitol is used in the paint and varnish industry, leather and paper manufacture, the plastics industry and explosives. A variety of foods contain mannitol. It can replace sucrose in candy and chocolate-flavored coatings. Sugar-free chocolates are popular among people with diabetes, a growing problem in modern society. Mannitol is used as a flavor enhancer because of its sweet taste and pleasantly cool feel in the mouth. Mannitol is non-hygroscopic and chemically inert; therefore, it can be used to maintain the proper moisture level in foods and increase shelf-life and stability [86]. Mannitol is non-carcinogenic. Finally, it can be used in pediatric and geriatric food products because it does not contribute to tooth decay [58,87].

Ulvans are highly charged, sulfated polyelectrolytes with average molecular weights from 189 to 8,200 kDa [88]. These cell wall polysaccharides comprise 8 to 29% of dry algal matter [58,66]. Ulvans' unique physicochemical properties make them attractive candidates for novel food, pharmaceutical, chemical aqua cultural, and agricultural bioactive polymers. Two major types have been identified: water soluble ulvans and insoluble, cellulose-like ulvans. Numerous researchers have demonstrated ulvans' antitumor, immunomodulatory [89], antiviral [90], anticoagulative [91] and antioxidant activities [92,93]. Ulvans were able to affect the adhesion and proliferation of normal and tumor human colonic cells. Ulvans were also able to affect the expression of transforming growth factors (TGF) and surface glycosyl markers related to cellular differentiation [89,94]. Moreover, ulvans induced mucin secretion in rat colon and, therefore, increased protection of the colonic mucosa. Because of their ion-exchange properties, ulvans are used as bioindicators for monitoring heavy metal pollution in coastal waters [95]. Lahaye and Robic extensively reviewed the available ulvan literature, emphasizing its structure and properties [94].

Species of *Porphyra* contain a sulfated, complex galactan called porphyran [66,67]. This has been used as a gelling agent, a nutritional supplement and an antioxidant [66,68]. Porphyran is a quality dietary fiber that chemically resembles agar. A 20% nori (*Porphyra*) powder mixed with a basic diet and given orally to rats prevented, 1,2-dimethylhydrazine-induced intestinal carcinogenesis. Porphyran has exhibited appreciable antitumor activity against Meth-A fibrosarcoma [69,70]. In addition, porphyran was demonstrated to significantly lower an artificially enhanced level of hypertension and bloodcholesterol in rats [77]. An agar-type polysaccharide obtained from a cold water extract of a *Gracilaria* species displayed antioxidant and antitumor activities [43]. Agar-oligosaccharides have also been shown to suppress the formation of a pro-inflammatory cytokine and an enzyme associated with nitric oxide production [96]. Agarose can be separated from agar in 42% yield; agar content varies seasonally from 26 to 42% in *Gelidium* species.

Algal oligosaccharides are carbohydrate molecules with low degrees of polymerization [77]. They are produced naturally and by chemical, physical or biochemical degradations of algal polysaccharides. Numerous oligosaccharides have been shown to possess antioxidant, immune stimulative and antitumor properties [22]. Additionally, a diet containing oligosaccharides benefits health because oligosaccharides are prebiotics. These digestion-resistant oligomers function as food sources for and promote the growth of beneficial bacteria [22]. For example, xylo- and fructo-oligosaccharides remain intact until the large intestine, where aerobic bacteria, such as bifidobacteria and lactobacilli, preferentially consume these sugars [97]. Moreover, non-digestible oligosaccharides have exhibited other biological activity, such as immunostimulative, antioxidative, antiangiogenic and antithrombotic properties [22]. In their review [77], Mussatto and Mancil showed that transgalactosylated disaccharide intake reduces fecal pH and ammonia, *p*-cresol and in dole concentrations. They also assert an increase in bifidobacteria and lactobacilli populations and a decrease in Bacteroidaceae populations. Because these changes in fecal physiological parameters are believed to reduce the risk of cancer, macroalgal non-digestible oligosaccharides are considered potential anticarcinogenic food ingredients [66].

Generally, algae contain a variety and high concentrations of minerals (8–40%) [21]. Their high mineral contents result from the ability of their cell surface polysaccharides to retain various inorganic marine substances [31]. However, actual mineral content varies between species and because of oceanic residence time, the geographic harvesting site, wave exposure, seasonal changes, environmental factors, physiological factors, processing, and the mineralization methods used [21,23]. Physiologically important minerals accumulate in seaweeds at considerably higher levels than in edible terrestrial plants and animals [31]. In fact, edible seaweed value in human nutrition is based on its high content of several essential minerals – As, Na, Mg, P, K, I, Fe, and Zn. Algal Na and K levels are particularly high, although their Na/K ratios are usually below 1:5 [98]. Because seaweeds also contain large amounts of trace elements that are limited in terrestrial vegetables, algal-based supplements could provide adults with the daily requirements of these minerals [21]. All seaweeds contain relatively large amounts of sulfates (1.3–5.9%), which are typical components of their polysaccharides. Fucans in brown algae and galactans in red algae bind sulfates; such sulfated mucilages are not found in terrestrial plants [99]. Sulfate levels relate to specific aspects of ionic regulation and salt concentrations in the algal aqueous environments [100]. Most algae contain higher Mg levels (500–1,000 mg/100 g) than terrestrial plants and animals [21]. Algae also

contain higher Fe levels than terrestrial organisms, but the concentration varies considerably depending on the species and harvesting season. A study on iron bioavailability revealed that iron absorption and retention was lower in seaweed-fed animals than a control. The researchers attributed the cause to the polyphenolic compounds or peptides from the partially digested algal protein in the diet [101]. Calcium is a major element in algae and is present at concentrations of 470–1,400 mg/100 g of DM [21]. P is found at levels of 200–800 mg/100 g of DM. Because of its high Ca/P ratio (3:5), algae could compensate for the calcium scarcity in several foods, such as cereals and meats [16,21]. Moreover, seaweed consumption may prove useful to individuals at risk of calcium deficiency, namely expectant mothers, adolescents and the elderly. Algal iodine concentrations, which vary greatly among species [101,102], also differ according to growth conditions, habitat, and pre-harvest conditions [103]. Brown seaweeds accumulate many elements. They are a good source of magnesium, copper, iron, iodine and other nutritionally rare elements. In the developed world, Japan has one of the lowest age-adjusted breast cancer mortality rates [1,104]. The incidence of breast cancer in Japanese immigrants to the United States and in their successive generations, has gradually reached the rates of white United States women, which suggests a dietary link [105]. High iodine intake may be a key protective factor against breast cancer development in Japanese women. One correlation study in Spain found a significant positive association between regions where iodine intake was low and breast cancer mortality rates high [106]. The use of iodine in treating breast cancer warrants study. Traditional eastern Asian medicine has long used iodine-rich seaweeds as a cancer treatment to “soften” tumors and “reduce” nodulation [107]. Recent *in vivo* animal studies suggest iodine possesses an antitumor effect [108]. In dimethylbenz [a] anthracene-induced mammary carcinoma in rats, iodine supplementation suppressed the disease’s development [106]. This suppressive activity was enhanced when the iodine treatment was combined with progesterone treatment. The suppressed tumors possessed significantly higher mean iodine content than the nonsuppressed tumors. Progesterone enhanced the iodine uptake [109]. An anticarcinogenic role for iodine in experimental animals was suggested by the work of Funahashi and co-workers. They observed that Lugol’s iodine or iodine-rich Wakame seaweed administration to rats treated with the carcinogen 7,12-dimethylbenzanthracene suppressed mammary tumor development [35]. Those researchers additionally demonstrated that seaweed induced a greater degree of apoptosis in human breast cancer cells than fluorouracil, a chemotherapeutic agent used to treat breast cancer. This finding led the authors to speculate that ‘seaweed may be applicable for prevention of breast cancer’ [109].

A major nutritive characteristic of edible algae is their high vitamin content, primarily with regard to water-soluble vitamins C and B and fat-soluble vitamins A and E. The vitamin C levels in green and brown algae range between 50 and 300 mg/100 g of DM, whereas the values for red algae range from 10 to 80 mg/100 g of DM [21]. Certain dried green and red algae, e.g. *Spirulina*, contain extensive amounts of vitamin B12. However, other edible algae contain little or none [110]. Daily ingestion of only 1 g of *Spirulina* supplies the daily requirements of B12. Because of their vitamin B12 content, algae have been recommended as dietary supplements for vegetarians, among whom megaloblastic anemia is common. Seaweed folate content (as folic acid or vitamin B9) ranges from 61.4 to 161.6 µg/100 g of DM [21]. Red and brown algae contain high levels of folic acid and folate derivatives, including 5-methyltetrahydrofolate, 5-formyltetrahydrofolate and tetrahydrofolate. On a DM basis, thiamine (vitamin B1) content is

high compared with that of terrestrial vegetables [111]. Brown seaweeds contain higher vitamin E levels (2.3–41.2 mg/100 g of DM) than green and red seaweeds (0.8 mg/100 g of DM) [112]. Brown algae contain α , β , and γ -tocopherols; green and red algae contain only α -tocopherol [113].

Phenolic compounds in seaweed possess a wide range of bioactivity: antimicrobial, antioxidant, antiviral and antitumor. These molecules are assumed to function in algae as chemical defenses against grazers, pathogens and epiphytes [114,115]; phenolic compounds in algae are also assumed to provide photoprotection [116]. Algal polyphenols, also called phlorotannins, are composed of phloroglucinol units (1,3,5-trihydroxybenzene). They constitute a heterogeneous group of molecules that vary greatly in their structure and degree of polymerization. Phlorotannins at high levels are found primarily in Fucales (20–30% dw). However, their extractable polyphenol levels (less than 0.4% algal DM) are lower than other phytochemicals. Phlorotannin concentrations are genetically variable and depend on the season of harvesting, habitat, and other extrinsic environmental factors such as salinity, nutrient and light availability, ultraviolet radiation, and herbivore grazing. Furthermore, phenolic content varies within an individual alga. Brown seaweeds display the highest phlorotannin concentrations [117]. Although phlorotannins have different structures and molecular weights, their chemical properties are similar [118]. Phlorotannins are ecologically important. They protect brown algae from ultraviolet irradiation and may be a chemical defense against herbivores. Like gallotannins or condensed tannins in terrestrial plants, phlorotannins are capable of precipitating proteins or metal ions. Polymeric phlorotannins inhibit enzymes such as phospholipase A, lipoxygenase, cyclooxygenase-1, hyaluronidase [118], and tyrosinase. Several polyphenolic compounds and complex mixtures have been isolated from brown algal species to explore the potential phlorotannin role in preventing degenerative diseases linked to oxidative stress [119]. Nutritionists agree that cancer incidence and neurodegenerative disease probability can be diminished by diets rich in fruits, grains and vegetables. The damage of DNA by reactive oxygen species is widely believed to be a significant contributor to age-related cancer development [63]. Until recently, algae were largely ignored as sources of antioxidant. Studies over the last several years have revealed that several seaweed components have appreciable antioxidant capability [20,120]. Tierney *et al.* thoroughly reviewed the available literature on macroalgal antioxidant activities [121]. Numerous researchers have studied seaweed health benefits *in vitro* [122] and *in vivo*.

A few human studies have assessed the impact of seaweed consumption on the risk for future disease (Table 3). Such data should be interpreted with caution because they indicate associative relationships but not necessarily causal relationships between seaweed intake and health outcomes. Different seaweeds appear to have different effects on disease risk. Yang reported that premenopausal women who ate the greatest amounts of seaweed were 56% less likely to be diagnosed with breast cancer than premenopausal women who ate the least amounts of seaweed [123]. An analysis of postmenopausal women revealed an even more dramatic risk reduction (68%) when comparing the highest and lowest quintiles of seaweed intake [123]. Preliminary research has demonstrated that selected seaweed phytochemicals (i.e., alginic acid, fucoidan, fucoxanthin and laminarin) interfere with cancer cell proliferation *in vitro* [124]. Human studies offer additional clues regarding possible mechanisms. Insulin-like growth factor 1 (IGF-1) is an anabolic hormone important for growth and development.

Table 3: Summary of recent observational studies on seaweed intake and health.

Disease/ Health condition	Type of study	Study design	Ref.
Breast cancer	Case-control study	362 cases	[123]
Cardiovascular disease	Prospective study	40547 Japanese men and women	[135]
Allergic rhinitis	Cross-sectional study	1002 Pregnant Japanese women	[136]
Serum IGF-1 Concentrations	Double-blinded, randomized, placebo-controlled crossover clinical trial.	30 Healthy Postmenopausal American Women	[125]
Obesity	3760 Japanese women aged 18-20 yr.	Cross-sectional study	[137]
Diabetes, Type II	Retrospective study	3,405 Korean individuals, aged 20 -65 yr.	[138]
post-load plasma glucose and insulin concentrations	Double-blind, randomized, placebo-controlled crossover study	Twenty-three participants (11 men, 12 women) aged 19-59 yr.	[129]
Serum total cholesterol	Retrospective study	> 7000 people	[139]
estrogen and phytoestrogen metabolism	Double-blind trial	15 Healthy postmenopausal women	[126]
Metabolic syndrome	Randomized double-blinded placebo-controlled trial	13 Men (mean age 47.4 ± 9.9 yr) and 14 women (average age 45.6 ± 12.2 yr) with at least one symptom of the metabolic syndrome	[140]
Immunomodulatory effects	open-label combined Phase I and II study	10 Healthy individuals aged between 18 and 65 yr	[52]

High-circulating serum concentrations of IGF-1 in adulthood are associated with an increased risk of postmenopausal breast cancer. Breast cancer incidence is typically lower in Asian countries where soy is commonly consumed. Notably, in trials with American women, soy supplements significantly increased IGF-1. However, seaweed is also consumed regularly in Asian countries with low breast cancer risk. Thirty healthy postmenopausal women participated in a 14-week double-blinded, randomized, placebo-controlled, crossover clinical trial. Participants consumed 5 g/day of either placebo or seaweed in capsules for 7 weeks. During the 7th week, a high-soy protein isolate powder was added. Soy significantly increased serum IGF-1 concentrations compared with the placebo. Seaweed reduced this soy increase by approximately 40%. The researchers concluded that concurrent seaweed and soy consumption may be important for moderating the effects of soy on IGF-1 serum concentrations, hence providing a possible mechanism for seaweed anticancer effects [125].

In another study, fifteen healthy postmenopausal women participated in a double-blind trial of seaweed supplementation with a soy challenge. Seaweed positively altered estrogen and phytoestrogen metabolism [126]. These observations may explain why the rates of estrogen-dependent cancers are higher in Western and lower in Eastern countries [127]. Results from our lab have shown that polyphenol-rich red seaweed is tumor-suppressive. The mechanism occurred via apoptosis induction, down regulating the endogenous estrogen biosynthesis, and improvement of the antioxidative status of the rats [128]. Because cholesterol is a precursor to sex hormone biosynthesis, kelp consumption may alter circulating sex hormone levels and menstrual cycling patterns. In particular, dietary kelp may be beneficial to women at high risk for estrogen-dependent diseases.

Additionally, brown seaweed was administered to three premenopausal women with either abnormal menstrual cycling patterns or menstruation-related disease histories. Seaweed intake was associated with significant increases in menstrual cycle lengths. Moreover, hormone measurements for one woman revealed significant anti-estrogenic and progestagenic effects following kelp administration. These pilot data suggest that dietary kelp may lengthen the menstrual cycle and exert anti-estrogenic effects in premenopausal women. Furthermore, these studies suggest that, in addition to soy, seaweed consumption may be responsible for the reduced risk of estrogen-related cancers observed in Japanese populations [127]. In another study, seaweed consumption was associated with a 12.1% reduction in the insulin incremental area. Glucose and insulin responses were similar between men and women. No adverse effects were associated with seaweed capsule consumption. These data suggest that brown seaweed may alter the insulin homeostasis in response to carbohydrate ingestion [129].

High levels of circulating insulin and an inadequate intake of iodine, a trace mineral found in seaweeds, are thought to contribute to breast cancer incidence. However, not every type of seaweed afforded the same level of protection. These results highlight the variability in the bioactive components of seaweed. Even within a single species, research has suggested that significant seasonal variations in nutritional content exist, which is likely to impact the biological effects of edible components outlined above.

Historically, seaweed has been consumed around the world. However, seaweed has been consumed in appreciable amounts only in certain areas. Although chemical analyses suggest numerous nutritional benefits to consuming seaweed, more evidence is needed linking seaweed dietary intake to health. Epidemiological studies suggest dietary factors could be critical to understanding cancer rates. Seaweed whole extracts or purified components have exhibited high antitumor activity *in vitro*. *In vivo* research with different animal models have demonstrated that fresh seaweed or seaweed extracts (as part of a regular diet, drinking water and/or as an injection) both inhibit and treat cancer.

Although little is known about relative cancer risk and seaweed intake in humans, a small body of *in vivo* and *in vitro* research suggests seaweed may be useful in cancer prevention and treatment. Seaweeds have been used to treat tumors in traditional Chinese and Japanese folk medicine. Populations that consume seaweed regularly, as in Japan, have dramatically lower rates of hormone-sensitive cancers, such as breast and prostate cancers [130]. Epidemiological studies performed in Japan before Westernized diets were common reported that Japanese women had lower rates of premenopausal and postmenopausal breast cancer [131]. In addition, after developing breast cancer, Japanese women were more likely to survive five years or more than United States women with breast cancer [132].

No clinical studies of cancer and seaweed have been reported. However, in a large prospective dietary study in Japan, investigators reported that, after 9 years of follow-up, miso soup (made from a concentrated hot water extract of seaweed plus a tablespoon or less of soybean paste and usually a few vegetables) was the food most closely associated with low breast cancer risk out of the foods investigated [133]. Women who ate three or more bowls of miso soup daily had approximately half the rate of breast cancer as those who ate two or fewer bowls. In two other epidemiological studies of diet and breast cancer in Japan, daily miso soup consumption was associated with 15% lower rate of breast cancer; women who drank miso soup at least five times /week had 13% lower rates [134].

References

- [1] Siegel R. (2012) Cancer statistics, 2012. *Cancer Journal for Clinicians*, **62**, 10–29.
- [2] Gerhauser C. (2013) Cancer chemoprevention and nutri-epigenetics: State of the art and future challenges. *Topics in Current Chemistry*, **329**, 73–132.
- [3] Blunden G. (2001) Biologically active compounds from marine organisms. *Phytotherapy Research*, **15**, 89–94.
- [4] Liu L, Heinrich M, Myers S, Dworjany SA. (2012) Towards a better understanding of medicinal uses of the brown seaweed *Sargassum* in Traditional Chinese Medicine: A phytochemical and pharmacological review. *Journal of Ethnopharmacology*, **142**, 591–619.
- [5] El Gamal AA. (2010) Biological importance of marine algae. *Saudi Pharmaceutical Journal*, **18**, 1–25.
- [6] Laurienzo P. (2010) Marine polysaccharides in pharmaceutical applications: an overview. *Marine Drugs*, **8**, 2435–2465.
- [7] Namvar F, Suhaila M, Gasemi Fard S, Behravan J. (2012) Polyphenol-rich seaweed (*Eucheuma cottonii*) extract suppresses breast tumour via hormone modulation and apoptosis induction. *Food Chemistry*, **130**, 376–382.
- [8] Aneiros A, Garateix A. (2004) Bioactive peptides from marine sources: pharmacological properties and isolation procedures. *Journal of Chromatography. B, Analytical technologies in the biomedical and life sciences*, **803**, 41–53.
- [9] Khan M, Choi J, Lee M, Kim E, Nam T. (2008) Anti-inflammatory activities of methanol extracts from various seaweed species. *Journal of Environmental Biology*, **29**, 465–469.
- [10] Zuercher AW, Fritsché R, Corthésy B, Mercenier A. (2006) Food products and allergy development, prevention and treatment. *Current Opinion in Biotechnology*, **17**, 198–203.
- [11] Perez GRM, Zavala SMA, Perez GS, Perez GC. (1998) Antidiabetic effect of compounds isolated from plants. *Phytomedicine*, **5**, 55–75.
- [12] Nishino T, Fukuda A, Nagumo T, Fujihara M, Kaji E. (1999) Inhibition of the generation of thrombin and factor Xa by a fucoidan from the brown seaweed *Ecklonia kurome*. *Thrombosis Research*, **96**, 37–49.
- [13] Campo VL, Kawano DF, Silva DB, Carvalho I. (2009) Carrageenans: Biological properties, chemical modifications and structural analysis – A review. *Carbohydrate Polymers*, **77**, 167–180.
- [14] Miyashita K. (2009) The carotenoid fucoxanthin from brown seaweed affects obesity. *Lipid Technology*, **21**, 186–190.
- [15] Wada K, Nakamura K, Tamai Y, Tsuji M, Sahashi Y, Watanabe K, Ohtsuchi S, Yamamoto K, Ando K, Nagata C. (2011) Seaweed intake and blood pressure levels in healthy pre-school Japanese children. *Nutrition Journal*, **10**, 83, 1–7.
- [16] Mohamed S, Hashim SN, Rahman HA. (2012) Seaweeds: A sustainable functional food for complementary and alternative therapy. *Trends in Food Science & Technology*, **23**, 83–96.
- [17] Ortiz J, Romero N, Robert P. (2006) Dietary fiber, amino acid, fatty acid and tocopherol contents of the edible seaweeds *Ulva lactuca* and *Durvillaea antarctica*. *Food Chemistry*, **99**, 98–104.
- [18] Colombo ML, Risé P, Giavarini F, De Angelis L, Galli C, Bolis CL. (2006) Marine macroalgae as sources of polyunsaturated fatty acids. *Plant Foods for Human Nutrition*, **61**, 67–72.
- [19] BobinDubigeon C, Lahaye M. (1997) Factors limiting the biodegradation of *Ulva* sp cell-wall polysaccharides. *Journal of the Science of Food and Agriculture*, **75**, 341–351.
- [20] Costa LS, Fidelis GP, Cordeiro SL, Oliveira RM, Sabry DA, Câmara RBG, Nobre LTDB, Costa MSSP, Almeida-Lima J, Farias EHC, Leite EL, Rocha HAO. (2010) Biological activities of sulfated polysaccharides from tropical seaweeds. *Biomedicine & Pharmacotherapy*, **64**, 21–28.
- [21] Bocanegra A, Bastida S, Benedí J, Ródenas S, Sánchez-Muniz FJ. (2009) Characteristics and nutritional and cardiovascular-health properties of seaweeds. *Journal of Medicinal Food*, **12**, 236–258.
- [22] Courtois J. (2009) Oligosaccharides from land plants and algae: production and applications in therapeutics and biotechnology. *Current Opinion in Microbiology*, **12**, 261–273.
- [23] Fleurence J. (1999) Seaweed proteins: biochemical, nutritional aspects and potential uses. *Trends in Food Science & Technology*, **10**, 26–29.
- [24] Galland-Irmouli A, Fleurence J. (2012) Nutritional value of proteins from edible seaweed *Palmaria palmata*. *Journal of Nutritional Biochemistry*, **10**, 353–359.
- [25] Ramos M, Cristina A, Moreira R., Carvalho AD. (2000) Amino acid composition of some Brazilian seaweed species. *Journal of Food Biochemistry*, **24**, 33–39.
- [26] Sekar S, Chandramohan M. (2007) Phycobiliproteins as a commodity: trends in applied research, patents and commercialization. *Journal of Applied Phycology*, **20**, 113–136.
- [27] Smit AJ. (2004) Medicinal and pharmaceutical uses of seaweed natural products: A review. *Journal of Applied Phycology*, **16**, 245–262.
- [28] Ogawa T, Watanabe M, Naganuma T, Muramoto K. (2011) Diversified carbohydrate-binding lectins from marine resources. *Journal of Amino Acids*. ID: 838914
- [29] Mabeau S, Fleurence J. (1993) Seaweed in food products: biochemical and nutritional aspects. *Trends in Food Science & Technology*, **4**, 927–929.
- [30] Dembitsky V. (1991) Glycolipids and fatty acids of some seaweeds and marine grasses from the Black Sea. *Phytochemistry*, **30**, 6–8.
- [31] MacArtain P, Gill C, Brooks M, Campbell R, Rowland L. (2007) Nutritional value of edible seaweeds. *Nutrition Reviews*, **65**, 535–543.
- [32] Pioveti L, Deffo P, Valls R, Peiffer G. (1991) Determination of sterols and diterpenoids from brown algae (Cystoseiraceae). *Journal of Chromatography*, **588**, 99–105.
- [33] Lahaye M, Kafeer B. (1997) Seaweed dietary fibres: structure, physico-chemical and biological properties relevant to intestinal physiology. *Sciences Des Aliments. Lavoisier*, **17**, 563–584.
- [34] Dawczynski C, Schubert R, Jahreis G. (2007) Amino acids, fatty acids, and dietary fibre in edible seaweed products. *Food Chemistry*, **103**, 891–899.
- [35] Funahashi H, Imai T, Tanaka Y, Tsukamura K, Hayakawa Y, Kikumori T, Mase T, Itoh T, Nishikawa M, Hayashi H, Shibata A, Hibi Y, Takahashi M, Narita T. (1999) Wakame seaweed suppresses the proliferation of 7,12-dimethylbenz(a)-anthracene-induced mammary tumors in rats. *Japanese Journal of Cancer Research: Gann*, **90**, 922–927.
- [36] Fitton J. (2005) Fucoidans: healthful saccharides from the sea. *Gynecoscience & Nutrition*, **6**, 1–6.
- [37] Morya VK, Kim J, Kim E-K. (2012) Algal fucoidan: structural and size-dependent bioactivities and their perspectives. *Applied Microbiology and Biotechnology*, **93**, 71–82.
- [38] Pomin VH, Mourão P.S. (2008) Structure, biology, evolution, and medical importance of sulfated fucans and galactans. *Glycobiology*, **18**, 1016–1027.
- [39] De Paula SM, Silveira M. (2009) Studies on molluscan shells: contributions from microscopic and analytical methods. *Micron*, **40**, 669–690.
- [40] Fitton JH. (2011) Therapies from fucoidan; multifunctional marine polymers. *Marine Drugs*, **9**, 1731–1760.
- [41] Berteau O, Mulloy B. (2003) Sulfated fucans, fresh perspectives: structures, functions, and biological properties of sulfated fucans and an overview of enzymes active toward this class of polysaccharide. *Glycobiology*, **13**, 29–40.
- [42] Ye H, Wang K, Zhou C, Liu J, Zeng X. (2008) Purification, antitumor and antioxidant activities *in vitro* of polysaccharides from the brown seaweed *Sargassum pallidum*. *Food Chemistry*, **111**, 428–432.

- [43] Fernández LE, Valiente OG, Mainardi V, Bello JL, Vélez H RA. (1989) Isolation and characterization of an antitumor active agar-type polysaccharide of *Gracilaria domingensis*. *Carbohydrate Research*, **190**, 77–83.
- [44] Hoffman R, Paper DH, Donaldson J, Alban S, Franz G. (1995) Characterisation of a laminarin sulphate which inhibits basic fibroblast growth factor binding and endothelial cell proliferation. *Journal of Cell Science*, **108**, 3591–3598.
- [45] Plaza M, Cifuentes A, Ibanez E. (2008) In the search of new functional food ingredients from algae. *Trends in Food Science & Technology*, **19**, 31–39.
- [46] Astorg P. (1997) Food carotenoids and an overview of current cancer prevention: properties. *Trends in Food Science & Technology*, **8**, 406–413.
- [47] Kotake-Nara E, Terasaki M, Nagao A. (2005) Characterization of apoptosis induced by fucoxanthin in human promyelocytic leukemia cells. *Bioscience, Biotechnology, and Biochemistry*, **69**, 224–227.
- [48] Pereira RC, Costa-lotufo LV. (2012) Bioprospecting for bioactives from seaweeds: potential, obstacles and alternatives. *Revista Brasileira de Farmacognosia*, **22**, 894–905.
- [49] Ikeguchi M, Yamamoto M, Arai Y, Maeta Y, Ashida K, Katano K, Miki Y, Kimura T. (2011) Fucoidan reduces the toxicities of chemotherapy for patients with unresectable advanced or recurrent colorectal cancer. *Oncology Letters*, **2**, 319–322.
- [50] Sithranga Boopathy N, Kathiresan K. (2010) Anticancer drugs from marine flora: an overview. *Journal of Oncology*, **ID**, 214186.
- [51] Hirayasu H. (2005) Sulfated the polysaccharides esterase activity derived from dietary seaweeds increase of a lymphocyte granzyme. *Journal of Nutritional Science and Vitaminology*, **51**, 475–477.
- [52] Myers SP, O'Connor J, Fitton JH, Brooks L, Rolfe M, Connellan P, Wohlmuth H, Cheras PA, Morris C. (2011) A combined Phase I and II open-label study on the immunomodulatory effects of seaweed extract nutrient complex. *Biologics: Targets & Therapy*, **5**, 45–60.
- [53] Wijesekara I, Pangestuti R, Kim S-K. (2011) Biological activities and potential health benefits of sulfated polysaccharides derived from marine algae. *Carbohydrate Polymers*, **84**, 14–21.
- [54] D'Orazio N, Gemello E, Gammone MA, De Girolamo M, Ficoneri C, Riccioni G. (2012) Fucoxanthin: a treasure from the sea. *Marine Drugs*, **10**, 604–616.
- [55] Thring TSA, Hili P, Naughton DP. (2009) Anti-collagenase, anti-elastase and anti-oxidant activities of extracts from 21 plants. *BMC Complementary and Alternative Medicine*, **9**, 27.
- [56] Senni K, Gueniche F, Foucault-Bertaud A, Igondjo-Tchen S, Fioretti F, Collicie-Jouault S, Durand P, Guezennec J, Godeau G, Letourneur D. (2006) Fucoidan a sulfated polysaccharide from brown algae is a potent modulator of connective tissue proteolysis. *Archives of Biochemistry and Biophysics*, **445**, 56–64.
- [57] Gupta S, Abu-Ghannam N. (2011) Bioactive potential and possible health effects of edible brown seaweeds. *Trends in Food Science & Technology*, **22**, 315–326.
- [58] Holdt SL, Kraan S. (2011) Bioactive compounds in seaweed: functional food applications and legislation. *Journal of Applied Phycology*, **23**, 543–597.
- [59] Miao HQ, Elkin M, Aingorn E, Ishai-Michaeli R, Stein CA, Vlodavsky I. (1999) Inhibition of heparanase activity and tumor metastasis by laminarin sulfate and synthetic phosphorothioate oligodeoxynucleotides. *International Journal of Cancer. Journal International du Cancer*, **83**, 424–431.
- [60] Zubia M, Payri C, Deslandes E. (2008) Alginate, mannitol, phenolic compounds and biological activities of two range-extending brown algae, *Sargassum mangroveense* and *Turbinaria ornata* (Phaeophyta: Fucales), from Tahiti (French Polynesia). *Journal Applied Phycology*, **20**, 1033–1043.
- [61] Yang E-J, Moon J-Y, Kim M-J, Kim DS, Kim C-S, Lee WJ, Lee NH, Hyun C-G. (2010) Inhibitory effect of Jeju endemic seaweeds on the production of pro-inflammatory mediators in mouse macrophage cell line RAW 264.7. *Journal of Zhejiang University. Science, B*, **11**, 315–322.
- [62] Murata M, Nakazoe J. (2001) Production and use of marine algae in Japan. *Japan Agricultural Research Quarterly*, **35**, 281–290.
- [63] Reddy L, Odhav B, Bhoola KD. (2003) Natural products for cancer prevention: a global perspective. *Pharmacology & Therapeutics*, **99**, 1–13.
- [64] Brownlee IA, Allen A, Pearson JP, Dettmar PW, Havler ME, Atherton MR, Onsøyen E. (2005) Alginate as a source of dietary fiber. *Critical Reviews in Food Science and Nutrition*, **45**, 497–510.
- [65] Wang Y, Han F, Hu B, Li J, Yu W. (2006) *In vivo* prebiotic properties of alginate oligosaccharides prepared through enzymatic hydrolysis of alginate. *Nutrition Research*, **26**, 597–603.
- [66] O'Sullivan L, Murphy B, McLoughlin P, Duggan P, Lawlor PG, Hughes H, Gardiner GE. (2011) Prebiotics from marine macroalgae for human and animal health applications. *Marine Drugs*, **8**, 2038–2064.
- [67] Jimenez-Escrig FJS-M. (2000) Dietary fibre from edible seaweeds: Chemical structure, physicochemical properties and effects on cholesterol metabolism. *Nutrition Research*, **20**, 585–598.
- [68] Cornish ML, Garbary DJ. (2010) Antioxidants from macroalgae: potential applications in human health and nutrition. *Algae*, **25**, 155–171.
- [69] Noda H, Amano H, Arashima K, Nisizawa K. (1990) Antitumor activity of marine algae. *Hydrobiologia*, **204-205**, 577–584.
- [70] Noda H, Amano H, Arashima K. (1989) Antitumor activity of polysaccharides and lipids from marine algae. *Nippon Suisan Gakkaishi*, **55**, 1265–1271.
- [71] Di Rosa M. (1972) Biological properties of carrageenan. *Journal of Pharmacy and Pharmacology*, **24**, 89–102.
- [72] Morris CJ. (2003) Carrageenan-induced paw edema in the rat and mouse. *Methods in Molecular Biology*, **225**, 115–121.
- [73] Dias PF, Siqueira JM, Vendruscolo LF, De Jesus Neiva T, Gagliardi AR, Maraschin M, Ribeiro-do-Valle RM. (2005) Antiangiogenic and antitumoral properties of a polysaccharide isolated from the seaweed *Sargassum stenophyllum*. *Cancer Chemotherapy and Pharmacology*, **56**, 436–446.
- [74] Zhou G, Sun Y, Xin H, Zhang Y, Li Z, Xu Z. (2004) *In vivo* antitumor and immunomodulation activities of different molecular weight lambda-carrageenans from *Chondrus ocellatus*. *Pharmacological Research*, **50**, 47–53.
- [75] Cáceres PJ, Carlucci MJ, Damonte EB, Matsuhira B, Zúñiga EA. (2000) Carrageenans from Chilean samples of *Stenogramme interrupta* (Phyllophoraceae): structural analysis and biological activity. *Phytochemistry*, **53**, 81–86.
- [76] Shah ZC, Huffman FG. (2003) Current availability and consumption of carrageenan-containing foods. *Ecology of Food and Nutrition*, **42**, 357–371.
- [77] Mussatto SI, Mancilha IM. (2007) Non-digestible oligosaccharides: A review. *Carbohydrate Polymers*, **68**, 587–597.
- [78] Schaeffer DJ, Krylov VS. (2000) Anti-HIV activity of extracts and compounds from algae and cyanobacteria. *Ecotoxicology and Environmental Safety*, **45**, 208–227.
- [79] Buck CB, Thompson CD, Roberts JN, Müller M, Lowy DR, Schiller JT. (2006) Carrageenan is a potent inhibitor of papillomavirus infection. *PLoS Pathogens*, **2**, e69.
- [80] Marais D, Gawarecki D, Allan B, Ahmed K, Altini L, Cassim N, Gopolang F, Hoffman M, Ramjee GWA. (2011) The effectiveness of Carraguard, a vaginal microbicide, in protecting women against high-risk human papillomavirus infection. *Antiviral Therapy*, **16**, 1219–1226.
- [81] Tobacman JK, Wallace RB, Zimmerman MB. (2001) Consumption of carrageenan and other water-soluble polymers used as food additives and incidence of mammary carcinoma. *Medical Hypotheses*, **56**, 589–598.

- [82] Gupta S, Abu-Ghannam N. (2011) Recent developments in the application of seaweeds or seaweed extracts as a means for enhancing the safety and quality attributes of foods. *Innovative Food Science & Emerging Technologies*, **12**, 600–609.
- [83] Lordan S, Ross RP, Stanton C. (2011) Marine bioactives as functional food ingredients: potential to reduce the incidence of chronic diseases. *Marine Drugs*, **9**, 1056–1100.
- [84] Iwamoto K, Shiraiwa Y. (2005) Mini-review: Salt-regulated mannitol metabolism in algae. *Marine Biotechnology*, **7**, 407–415.
- [85] Noda H. (1993) Health benefits and nutritional properties of nori. *Journal of Applied Phycology*, **5**, 255–258.
- [86] Le AS, Mulderrig KB. (2001) Sorbitol and mannitol. *Food Science and Technology*. Marcel Dekker, New York, 317–334.
- [87] Nabors L. (2004) Alternative sweeteners. *Agro Food Industry Hi-Tech*, **15**, 39–41.
- [88] Lahaye MKB. (1997) Seaweed dietary fibres: structure, physico-chemical and biological properties relevant to intestinal physiology. *Sciences des Aliments*, **17**, 563–584.
- [89] York GTVS-N, Škerget M. (1999) Biological properties of ulvan, a new source of green seaweed sulfated polysaccharides, on cultured normal and cancerous colonic epithelial cells. *Journal of Medicinal Plant and Natural Product Research*, **65**, 527–231.
- [90] Wang W, Wang S-X, Guan H-S. (2012) The antiviral activities and mechanisms of marine polysaccharides: an overview. *Marine Drugs*, **10**, 2795–2816.
- [91] Mao W, Zang X, Li Y, Zhang H. (2006) Sulfated polysaccharides from marine green algae *Ulva conglobata* and their anticoagulant activity. *Journal of Applied Phycology*, **18**, 9–14.
- [92] Alves A, Sousa RA, Reis RL. (2012) A practical perspective on ulvan extracted from green algae. *Journal of Applied Phycology*. (DOI 10.1007/s10811-012-9875-4).
- [93] Qi H, Zhang Q, Zhao T, Hu R, Zhang K, Li Z. (2006) *In vitro* antioxidant activity of acetylated and benzoyleated derivatives of polysaccharide extracted from *Ulva pertusa* (Chlorophyta). *Bioorganic & Medicinal Chemistry Letters*, **16**, 2441–2445.
- [94] Lahaye M, Robic A. (2007) Structure and functional properties of ulvan, a polysaccharide from green seaweeds. *Biomacromolecules*, **8**, 24–30.
- [95] Nagaoka M, Shibata H, Kimura-Takagi I, Hashimoto S, Kimura K, Makino T, Aiyama R, Ueyama S, Yokokura T. (1999) Structural study of fucoidan from *Cladosiphon okamuranus* Tokida. *Glycoconjugate Journal*, **16**, 19–26.
- [96] Wang J, Jiang X, Mou H, Guan H. (2004) Anti-oxidation of agar oligosaccharides produced by agarase from a marine bacterium. *Journal of Applied Phycology*, **16**, 333–340.
- [97] Wang Y. (2009) Prebiotics: Present and future in food science and technology. *Food Research International*, **42**, 8–12.
- [98] Rupe P. (2002) Mineral content of edible marine seaweeds. *Food Chemistry*, **79**, 23–26.
- [99] Karthikaidevi G. (2010) Mineral composition of marine macroalgae from Mandapam coastal regions; Southeast coast of India. *American-Eurasian Journal of Botany*, **1**, 58–67.
- [100] Kloareg B, Quatrano R.S. (1988) Structure of the cell walls of marine algae and ecophysiological functions of the matrix polysaccharides. *Oceanography and Marine Biology Annual Review*, **26**, 259–315.
- [101] Shaw NS, Liu YH. (2000) Bioavailability of iron from purple laver (*Porphyra* spp.) estimated in a rat hemoglobin regeneration bioassay. *Journal of Agricultural and Food Chemistry*, **48**, 1734–1737.
- [102] Hou X, Chai C, Qian Q, Yan X, Fan X. (1997) Determination of chemical species of iodine in some seaweeds (I). *Science of the Total Environment*, **204**, 215–221.
- [103] Wap B. (1950) The seasonal variation in the cellulose content of the common Scottish Laminariaceae and Fucaceae. *Journal of the Marine Biological Association of the United Kingdom*, **29**, 379–387.
- [104] Parker SL, Davis KJ, Wingo PA, Ries LA, Heath CW. (1998) Cancer statistics by race and ethnicity. *CA: a Cancer Journal for Clinicians*, **48**, 31–48.
- [105] Ziegler RG, Hoover RN, Pike MC, Hildesheim A, Nomura AM, West DW, Wu-Williams AH, Kolonel LN, Horn-Ross PL, Rosenthal JF, Hyer MB. (1993) Migration patterns and breast cancer risk in Asian-American women. *Journal of the National Cancer Institute*, **85**, 1819–1827.
- [106] Cann SA, Van Netten JP, Van Netten C. (2000) Hypothesis: iodine, selenium and the development of breast cancer. *Cancer Causes & Control:CCC*, **11**, 121–127.
- [107] Lin H, Liu J ZY. (2011) Developments in cancer prevention and treatment using traditional Chinese medicine. *Frontiers of Medicine*, **5**, 127–133.
- [108] Venturi S. (2001) Is there a role for iodine in breast diseases? *Breast (Edinburgh, Scotland)*, **10**, 379–382.
- [109] Funahashi H, Imai T, Mase T, Sekiya M, Yokoi K, Hayashi H, Shibata A, Hayashi T, Nishikawa M, Suda N, Hibi Y, Mizuno Y, Tsukamura K, Hayakawa A, Tanuma S. (2001) Seaweed prevents breast cancer? *Japanese Journal of Cancer Research:Gann*, **92**, 483–487.
- [110] Sánchez-Machado DI, López-Cervantes J, López-Hernández J, Paseiro-Losada P. (2004) Simultaneous determination of thiamine and riboflavin in edible marine seaweeds by high-performance liquid chromatography. *Journal of Chromatographic Science*, **42**, 117–120.
- [111] Fleurence J, Morancès M, Dumay J, Decottignies P, Turpin V, Munier M, Garcia-Bueno N, Jaouen P. (2012) What are the prospects for using seaweed in human nutrition and for marine animals raised through aquaculture? *Trends in Food Science & Technology*, **27**, 57–61.
- [112] Kumari P, Kumar M, Gupta V, Reddy CRK, Jha B. (2010) Tropical marine macroalgae as potential sources of nutritionally important PUFAs. *Food Chemistry*, **120**, 749–757.
- [113] Matanjun P, Mohamed S, Mustapha NM, Muhammad K. (2008) Nutrient content of tropical edible seaweeds, *Eucheuma cottonii*, *Caulerpa lentillifera* and *Sargassum polycystum*. *Journal of Applied Phycology*, **21**, 75–80.
- [114] Dohrn SZA, Marine T. (2006) New trends in marine chemical ecology. *Estuaries and Coasts*, **29**, 531–551.
- [115] Amsler CD FV. (2006) Defensive and sensory chemical ecology of brown algae. *Advances in Botanical Research*, **43**, 1–91.
- [116] Pavia H, Cervin G, Lindgren A, Åberg P. (1997) Effects of UV-B radiation and simulated herbivory on phlorotannins in the brown alga *Ascophyllum nodosum*. *Marine Ecology Progress*, **157**, 139–146.
- [117] Wang T, Jónsdóttir R, Ólafsdóttir G. (2009) Total phenolic compounds, radical scavenging and metal chelation of extracts from Icelandic seaweeds. *Food Chemistry*, **116**, 240–248.
- [118] Shibata T, Fujimoto K. (2002) Inhibitory activity of brown algal phlorotannins against hyaluronidase. *International Journal of Food Science and Technology*, **37**, 703–709.
- [119] Kang HS, Chung HY, Kim JY, Son BW, Jung HA, Choi JS. (2004) Inhibitory phlorotannins from the edible brown alga *Ecklonia stolonifera* on total reactive oxygen species (ROS) generation. *Archives of Pharmacology Research*, **27**, 194–198.
- [120] Vinayak RC, Sabu A S, Chatterji A. (2011) Bio-prospecting of a few brown seaweeds for their cytotoxic and antioxidant activities. *Evidence-based Complementary and Alternative Medicine: eCAM*. Article ID:673083.
- [121] Tierney MS, Croft AK, Hayes M. (2010) A review of antihypertensive and antioxidant activities in macroalgae. *Botanica Marina*, **53**, 387–408.
- [122] Moo-Puc R, Robledo D, Freile-Pelegrin Y. (2011) Enhanced antitumoral activity of extracts derived from cultured *Udotea flabellum* (Chlorophyta). *Evidence-based Complementary and Alternative Medicine: eCAM*. Article ID: 969275.
- [123] Yang YJ, Nam S-J, Kong G, Kim MK. (2010) A case-control study on seaweed consumption and the risk of breast cancer. *The British Journal of Nutrition*, **103**, 1345–1353.

- [124] Namvar F, Baharara J, Mahdi A. (2013) Antioxidant and anticancer activities of selected Persian Gulf algae. *Indian Journal of Clinical Biochemistry*, DOI: 10.1007/s12291-013-0313-4
- [125] Teas J, Irhimeh MR, Druker S, Hurley TG, Hébert JR, Savarese TM, Kurzer MS. (2011) Serum IGF-I concentrations change with soy and seaweed supplements in healthy postmenopausal American women. *Nutrition and Cancer*, **63**, 743–748.
- [126] Hurley TG, Hebert JR, Franke AA, Sepkovic DW, Kurzer MS. (2009) Dietary seaweed modifies estrogen and phytoestrogen metabolism in healthy postmenopausal women. *Nutrition Journal*, **139**, 1779.
- [127] Skibola CF. (2004) The effect of *Fucus vesiculosus*, an edible brown seaweed, upon menstrual cycle length and hormonal status in three premenopausal women: a case report. *BMC Complementary and Alternative Medicine*, **4**, 4–10.
- [128] Namvar F, Mohamed S, Fard SG, Behravan J, Mustapha NM, Alitheen NBM, Othman F. (2012) Polyphenol-rich seaweed (*Eucheuma cottonii*) extract suppresses breast tumour via hormone modulation and apoptosis induction. *Food Chemistry*, **130**, 376–382.
- [129] Paradis M-E, Couture PBL. (2011) A randomised crossover placebo-controlled trial investigating the effect of brown seaweed (*Ascophyllum nodosum* and *Fucus vesiculosus*) on postchallenge plasma glucose and insulin levels in men and women. *Applied Physiology, Nutrition, and Metabolism*, **36**, 913–919.
- [130] Hebert JR, Hurley TG, Olendzki BC, Teas J, Ma Y, Hampl JS. (1998) Nutritional and socioeconomic factors in relation to prostate cancer mortality: a cross-national study. *Journal of the National Cancer Institute*, **90**, 1637–1647.
- [131] Kodama M, Kodama T, Miura SYM. (1991) Nutrition and breast cancer risk in Japan. *Anticancer Research*, **11**, 745–754.
- [132] Morrison AS, Black MM, Lowe CR, MacMahon BYS. (1973) Some international differences in histology and survival in breast cancer. *International Journal of Cancer*, **11**, 261–267.
- [133] Yamamoto S, Sobue T, Kobayashi M, Sasaki S, Tsugane S. (2003) Soy, isoflavones, and breast cancer risk in Japan. *Journal of the National Cancer Institute*, **95**, 906–913.
- [134] Key TJ, Sharp GB, Appleby PN, Beral VGM. (1999) Soya foods and breast cancer risk: a prospective study in Hiroshima and Nagasaki, Japan. *British Journal of Cancer*, **81**, 1248–1256.
- [135] Shimazu T, Kuriyama S, Hozawa A, Ohmori K, Sato Y, Nakaya N, Nishino Y, Tsubono Y, Tsuji I. (2007) Dietary patterns and cardiovascular disease mortality in Japan: a prospective cohort study. *International Journal of Epidemiology*, **36**, 600–609.
- [136] Miyake Y, Sasaki S, Ohya Y, Miyamoto S, Matsunaga I, Yoshida T, Hirota Y, Oda H. (2006) Dietary intake of seaweed and minerals and prevalence of allergic rhinitis in Japanese pregnant females: baseline data from the Osaka Maternal and Child Health Study. *Annals of Epidemiology*, **16**, 614–621.
- [137] Okubo H, Sasaki S, Murakami K, Kim MK, Takahashi Y, Hosoi Y, Itabashi M. (2008) Three major dietary patterns are all independently related to the risk of obesity among 3760 Japanese women aged 18–20 years. *International Journal of Obesity*, **32**, 541–549.
- [138] Lee HJ, Kim HC, Vitek L, Nam CM. (2010) Algae consumption and risk of type 2 diabetes: Korean National Health and Nutrition Examination Survey in 2005. *Journal of Nutritional Science and Vitaminology*, **56**, 13–18.
- [139] Kondo I, Funahashi K, Nakamura M, Ojima T, Yoshita K, Nakamura Y. (2010) Association between food group intake and serum total cholesterol in the Japanese population. *Journal of Epidemiology*, **20**, S576–S581.
- [140] Teas J, Baldeón ME, Chiriboga DE, Davis JR, Sarriés AJ, Braverman LE. (2009) Could dietary seaweed reverse the metabolic syndrome? *AsiaPacific Journal of Clinical Nutrition*, **18**, 145–154.

Methods for Extraction and Determination of Phenolic Acids in Medicinal Plants: A Review

Agnieszka Arceusz, Marek Wesolowski* and Pawel Konieczynski

Department of Analytical Chemistry, Medical University of Gdansk, 80-416 Gdansk, Poland

marwes@gumed.edu.pl

Received: July 18th, 2013; Accepted: October 8th, 2013

Phenolic acids constitute a group of potentially immunostimulating compounds. They occur in all medicinal plants and are widely used in phytotherapy and foods of plant origin. In recent years, phenolic acids have attracted much interest owing to their biological functions. This paper reviews the extraction and determination methods of phenolic acids in medicinal plants over the last 10 years. Although Soxhlet extraction and ultrasonic assisted extraction (UAE) are commonly used for the extraction of phenolic acids from plant materials, alternative techniques such as supercritical fluid extraction (SFE), and accelerated solvent extraction (ASE) can also be used. After extraction, phenolic acids are determined usually by liquid chromatography (LC) owing to the recent developments in this technique, especially when it is coupled with mass spectrometry (MS). Also detection systems are discussed, including UV-Vis, diode array, electrochemical and fluorimetric. Other popular techniques for the analysis of this group of secondary metabolites are gas chromatography coupled with mass spectrometry (GC-MS) and capillary electrophoresis (CE).

Keywords: Phenolic acids, Medicinal plants, Extraction, Chromatographic techniques, Capillary electrophoresis, Secondary metabolites.

Introduction

Vascular plants synthesise many organic compounds, often called secondary metabolites [1]. These low-molecular-weight compounds determine the basic life processes of plants. It has been estimated that about 8000 compounds naturally occurring in plants are phenols. Their characteristic structural feature is an aromatic ring with at least one hydroxyl substituent. Many phenolic compounds occur constitutively, but some stress factors contribute to increases in or *de novo* synthesis of phenolics, such as infection, plant tissue damage, UV radiation and elevated temperature [2].

The popularity and consumption of medicinal plants have grown significantly in recent years. This phenomenon has led to a faster and better evaluation of the quality of plant products. One of the tools playing a crucial role as an element controlling the quality of plant material used in medicine is the so-called fingerprint analysis [3-5].

Determination of phenolic compounds can be very helpful in estimation of pharmacological activity of medicinal plants [6]. Within this huge group of compounds, a significant role is played by phenolic acids, aromatic secondary metabolites widespread in the plant kingdom. These compounds contain both hydroxyl and carboxyl groups. Phenolic acids include hydroxyl derivatives of benzoic and cinnamic acids (Table 1). Much attention has been focused on gallic, vanillic, salicylic, caffeic and *p*-coumaric acids, which are active constituents of many plants [7,8]. Examples include thyme, whose phenolic acids comprise gallic, caffeic and rosmarinic acids, also sage containing ferulic, gallic, rosmarinic, vanillic, and caffeic acids, as well as rosemary, characterised by a high content of vanillic, caffeic and rosmarinic acids [7].

Source and roles played by phenolic acids in plants

Phenolic acids are secondary metabolites and constitute an important group of hydrophilic compounds in plant tissues [9,10]. They seldom occur in the free form and, therefore, they appear in low concentrations. In plants, phenolic compounds can appear mainly in their bound forms, for example as glycosides (phenolic glycosides)

and in esters as depsides and depsidones [11]. In addition, in plant tissues, compounds of phenolic acids with other natural constituents have been identified, for instance, with flavonoids, fatty acids, sterols and cell wall polymers.



The contents of phenolic acids in plant raw materials, as well as of other phenolic compounds, depend on many factors. The biggest impacts come from climatic conditions, cultivation, fertilization and time of harvest [2,12,13]. Also important are storage conditions of the plant materials and the methods of preparation [10]. It is known that plants from the *Lamiaceae* family contain especially large amounts of phenolic acids [14]. These compounds can also be found in large amounts in fruits and vegetables. The main source of phenolic acids is herbal infusions; they also occur in black and green tea, and coffee [15]. In food products, phenolic acids may influence the color, flavor, fragrance and oxidation stability of food. In this case, their level depends, among other factors, on the type of technological process applied to a given plant material [10].

Source and roles of phenolic acids in human

Phenolic acids constitute an important group of compounds, characterised by a wide spectrum of pharmacological activity. They are responsible for free radical scavenging, metallic ions chelation, and changing enzymatic activity. Moreover, these compounds exhibit antiviral activity, for instance rosmarinic acid, anti-inflammatory activity, for example a mixture of esters of benzoic and cinnamic acids; diuretic, anti-allergic, as well as chologogic and cholcretic activities [15-17]. Phenolic acids participate in regeneration and adaptation processes in humans and are used in prevention against many diseases [18,19]. It was proved that they prevent coronary disease, inflammation, type-2 diabetes, as well as aiding in the treatment of cancer. Several of these compounds, for instance ferulic and caffeic acids, are called cancer growth inhibitors. Furthermore, recent studies have also shown pro-coagulant activity of phenolic acids isolated from *Blumea riparia* DC [20] and possible degradation pathway of salvianolic acid B in water solution and simulated gastric and intestinal fluids [21].

Table 1: Structural formulas of phenolic acids.

Hydroxybenzoic acids	R ₁	R ₂	R ₃	R ₄
Benzoic acid	H	H	H	H
<i>p</i> -Hydroxybenzoic acid	H	H	OH	H
Vanillic acid	H	OCH ₃	OH	H
Gallic acid	H	OH	OH	OH
Syringic acid	H	OCH ₃	H	OCH ₃
Protocatechic acid	H	OH	OH	H
Gentisic acid	OH	H	H	OH
Veratric acid	H	OCH ₃	OCH ₃	H
Salicylic acid	OH	H	H	H
Hydroxycinnamic acids	R ₁	R ₂	R ₃	R ₄
Cinnamic acid	H	H	H	H
<i>o</i> -Coumaric acid	OH	H	H	H
<i>m</i> -Coumaric acid	H	OH	H	H
<i>p</i> -Coumaric acid	H	H	OH	H
Ferulic acid	H	H	OH	OCH ₃
Caffeic acid	H	OH	OH	H



The availability of phenolic acids in humans is mostly influenced by their chemical form, as well as the morphological part of the plant [22]. More hydrophilic compounds are characterised by a higher bioavailability and are more readily absorbed in the upper part of the digestive system. On the other hand, substances in their bound form are absorbed after enzymatic hydrolysis, which is mediated by intestinal microflora.

The biological significance of phenolic acids calls for the need of elaboration of appropriate analytical methods enabling their monitoring in drugs and foods of plant origin, as well as in plant raw materials used for their production. Bearing all this in mind, the aim of this work is to present an up-to-date review of the most often used methods for extraction and determination of phenolic acids in plant materials.

Extraction methods

One of the main sources of errors committed during analytical procedure is the stage of sample preparation. As this stage determines the time of analysis, it is crucial to shorten the procedure and to enhance its accuracy and selectivity while identifying the majority of chemical compounds [23]. In order to separate the analytes from solid samples, the most often used solvents are those with low evaporation heats and low boiling points. The solvents also have to be non-toxic, non-flammable and chemically neutral, and they should not influence negatively the instrumentation and stability of the analysed substances. Solvents such as methanol, ethanol, acetone, ethyl acetate and their mixtures with water are commonly used [6,10,24,25].

Phenols and phenolic acids also exist as insoluble bound complexes, which are coupled to cell wall polymers through ester and glycosidic links and they are not extractable by organic solvents [11,26]. Bound phenolic acids are typically liberated using base hydrolysis, acid hydrolysis, or both before extraction [27-29]. Many extraction procedures incorporate the use of an antioxidant as stabilizer; compounds that have been used for this purpose include butylated hydroxyanisole, tert-butylhydroquinone and ascorbic acid.

Soxhlet extraction

Extraction is a first stage of the process leading to isolation of secondary metabolites from plant materials. A classical method used for this purpose is Soxhlet extraction. This method ensures multiple digestion of a plant material and continuous extraction, while a fresh portion of a solvent is delivered. Thanks to this, the largest difference

between the concentrations of analyte in the cell solution and in the solvent is maintained throughout, which leads to total extraction of a plant material.

Multiple extraction with the use of a Soxhlet apparatus was applied for extraction of phenolic acids from plant material (*Sambucus nigra* L., *Polygonum aviculare*). Methanol was a solvent and the digestion process lasted 15 h. Application of this modification of extraction in the case of wild common lilac allowed a high extraction yield of *p*-hydroxybenzoic, vanillic and ferulic acids [30]. Soxhlet extraction was also used for qualitative and quantitative analysis of phenolic compounds occurring in *Salvia halophila* and *S. virgata* [31]. To optimize the extraction conditions, *n*-hexane, ethyl acetate, methanol and 50% aqueous methanol solution were used during 8 h. It was found that the highest extraction yield of total phenolics was obtained when 50% methanol was used, and the lowest by using *n*-hexane. Kořar et al. [32] tested the same solvents when they investigated extraction of *S. halophila*.

Soxhlet extraction was also used by Karasová and Lehotay [33], who reported on the isolation of benzoic acid derivatives from *Melissa officinalis*. Different extraction times were tested (1 h, 4 h, 8 h), as well as the solvent, which was a mixture of methanol with water at different volume ratios (60:40 and 80:20). Eventually, for separation of the derivatives, the 80:20 mixture of methanol and water was applied, and the extraction was performed during 1 h. It has been concluded that the time of extraction had no influence on the results. The only exception was gallic acid, because, after extending the extraction time, the lowest concentrations of this compound were obtained.

The main disadvantages of Soxhlet extraction are that it is a time-consuming process and uses costly solvents, which must be of appropriate quality. For these reasons, in recent years other methods have been preferred that are less time-consuming and require smaller volumes of solvents [34]. Among these methods are ultrasound-assisted extraction (UAE), supercritical fluid extraction (SFE), and accelerated solvent extraction (ASE). These methods can be successfully used for extraction of phenolic compounds from plant materials. A comparison of these methods is presented in Table 2.

Ultrasound-assisted extraction (UAE)

Ultrasound-assisted extraction (UAE) is based on the action of ultrasonic vibrations directed toward an extracted sample, which enhance efficacy of penetration of a sample by solvent. This method

Table 2: Comparison of extraction techniques.

Parameter	Soxhlet	UAE	MAE	SFE	ASE
Costs	low	low	average	high	high
Time of extraction	6-48 h	< 30 min	< 30 min	< 60 min	< 30 min
Solvent volume to be used [mL]	200-600	< 50	< 40	< 10	< 100

is characterized by high speed, simplicity, and usually takes several minutes. Apart from the type of a solvent, sample size, pH of extract, temperature and pressure, factors such as particle size, duration of sonification and its amplitude, also have an impact on the efficacy of the extraction process [35,36]. Regardless of these factors, the method is considered as the simplest one possible to perform in a laboratory [37,38]. One of its advantages is the possibility to carry out the extraction of several samples simultaneously within a relatively short time. However, it is necessary to decant the extract or to filter it through appropriate paper filters. On the other hand, an opaque solution after extraction could get stuck on an HPLC column. In practice, during the UAE process disruption of plant cell walls takes place, this enabling the effective extraction of metabolites contained in the cells. This method was applied by Pawar *et al.* [39], who analysed different species of ginger. The UAE process was conducted for 30 min, using methanol as a solvent.

Ultrasound-assisted extraction is not the sole process used for extraction. Sometimes it is preceded by another procedure in order to prepare adequately the sample, for example to moisten it with a solvent. Such a procedure has been proposed by You *et al.* [29], who extracted the fruits of blackberry with an 80:20 (v/v) methanol/water mixture, with addition of acetic acid, during 16 h in a dark place. Then, sonication was performed and the analysed samples were centrifuged twice. Benetis *et al.* [40] have also used ultrasound for the analysis of phenolic compounds in *Achillea millefolium* L. Preliminary studies conducted with the use of ethanol/water solutions of concentrations from 40% to 96% (v/v), and applying different extraction times of 5, 10, 20, 30 and 60 min, have shown that the best results were obtained with the 70% ethanol solution. An extraction profile in relation to ultrasonication time has shown that by extending the time of extraction the yield increased, but after 60 min a decrease in yield was noticed. Finally, for further experiments, a 70% ethanolic solution was chosen as solvent and a 30-min ultrasonication used, with a 0.25g sample and 25 mL of solvent.

Among numerous references describing application of UAE to sample preparation, there is one, which compares the efficacy of this method with the classical liquid-liquid extraction [41]. For both processes, the following solvents were used: a 60% methanolic solution, a 60% acetone solution, water, and a 60:30 (v/v) mixture of water and ethyl acetate. The analysed material was aromatic plants, including *Rosmarinus officinalis* and *Origanum majorana*. Classical digestion was performed in a water bath at 90°C for 2 h, whereas ultrasonication was performed for 1 h at 25° and 60°C. Both processes were carried out in triplicate using each of the above mentioned solvents. The methanolic solvent appeared to be the best, and the temperature of 60°C. Under these conditions, the highest yield of extraction was achieved.

Supercritical fluid extraction (SFE)

This type of extraction often precedes analysis by HPLC [42]. Supercritical fluid extraction is a modern technique, which has many advantages over the classical extraction methods [38,43]. Among these are low temperatures, which is a positive feature in the case of analysis of thermally labile compounds. Other advantages are high selectivity, significant reduction in solvent volumes used for extraction, low mass of sample for extraction (around several mg),

short extraction time, possibility of automation, as well as off-line and on-line coupling with a majority of chromatographic techniques (GC, HPLC). The solvent used in SFE is a fluid in a supercritical state, usually carbon dioxide, owing to its low price and low toxicity. However, in the case of phenolic compounds, it is not the best solvent, since its polarity is low in comparison with that of phenolics [44]. In spite of this, Castro-Vargas *et al.* [45] used carbon dioxide for isolation of phenolic fractions from the seeds of *Psidium guajava* L., family *Myrtaceae*.

Many parameters influence the efficacy of extraction by supercritical fluids. They can be divided into two groups [46]. The first encompasses parameters related to conducting the extraction, such as pressure, temperature, time, sample weight, and flow intensity. The other group involves parameters connected with the matrix of a sample, such as its form, homogeneity, solubility and desorption ability of analytes. Apart from this, the SFE process depends on the pH. Its variations can influence the extraction yield and the speed of extraction of the analyte from an aqueous phase.

SFE is commonly applied in the pharmaceutical industry, as well as in the food and cosmetic areas [38]. Supercritical fluid extraction serves for the isolation of biologically active compounds from plant materials, mainly for those which cannot be separated by the use of simple solvent extraction.

Accelerated solvent extraction (ASE)

The accelerated solvent extraction method is a relatively new technique developed and distributed mainly by Dionex (Dionex Corporation, Sunnyvale, CA). In the literature, there are alternative names for this technique, such as pressurized fluid extraction (PFE) and pressurized liquid extraction (PLE).

In the ASE technique the same solvents are used as those in classical methods, but a higher pressure (about 3.3-20.3 MPa) and elevated temperature, about 40-200°C, are applied [38]. A sample extracted by this technique is placed in an extraction vessel made of stainless steel. The time of analysis is short, within the range of 5-15 min [47].

Basic advantages of ASE are: the possibility to extract samples with high humidity, short extraction time, better penetration of a sample by solvent, good extraction kinetics, automation of the process and relatively easy use of instrumentation [38]. Also, lesser volumes of solvents are needed. For example, during ASE extraction lasting about 20 min, only 10-50 mL of solvent were used, whereas in classical methods, such as Soxhlet extraction, extraction time is usually from 10 to 48 h, and the volume of solvent exceeds 400 mL [47].

Shake extraction

Beside the above-mentioned methods of sample extraction used for the analysis of plant material, extraction with various shaking devices has been performed. The process of shaking is aimed at enlarging the surface area where the solvent interacts with plant material, and in this way enhancing efficacy of the whole process and shortening its time. In this method, a sample is suspended in a certain volume of solvent and then shaken at a set stirrer speed. The method was applied by Li *et al.* [48] for the analysis of phenolics in *Phalaris canariensis* L. The

procedure was as follows. To the analysed sample, an 85:15 (v/v) mixture of 95% ethanol and 1 M hydrochloric acid was added, and the suspension was shaken on an orbital (rotary) shaker for 1 h at 30°C. After centrifugation of the extract, the plant material was re-extracted under the same conditions, and finally the combined filtrates were evaporated under reduced pressure. For further analysis, the dry residue was used after dissolving it in a 50% methanolic solution.

Fernand *et al.* [49] have also applied the shaking process for the extraction of a plant material. An ethanolic solution was used as the extractant, whereas the dry residue, after evaporation of the solvent, was dissolved in a 1:1 (v/v) mixture of ethanol and water. In the next step, solid-phase extraction was used to remove from the extract constituents interfering with phenolic components.

Santos-Gomes *et al.* [50] conducted their investigation in a similar way. However, instead of an ethanolic solution they used acetone and a shortened extraction time. A triple extraction was applied and simultaneous reduction of the solvent's volume: 50 mL over 15 min, 50 mL over 10 min and 25 mL over 5 min. Then, the residue left after concentration of the extracts under reduced pressure was dissolved in a methanolic solution, which was analysed by HPLC. Extraction by shaking was also applied for the analysis of rosmarinic acid in the leaves of lemon balm [51]. Aqueous methanol solutions were used as solvents, with methanol/water ratios of 40, 60 and 80% (v/v). Also, the impact of extraction time (30, 60 and 90 min) and temperature (25, 40 and 55°C) on the process was studied. The optimum extraction conditions were obtained when the process was conducted during 60 min with use of 60:40 (v/v) methanol/water solution.

Solid-liquid extraction carried out in a separation funnel is also considered as one of the shaking methods, but extraction yields are more variable because shaking speed and strength cannot be controlled with great accuracy. In the case of analysis using gas chromatography (GC), the most often applied extraction processes are: solid-phase micro-extraction (SPME), headspace single-drop micro-extraction (HSDME) and microwave-assisted extraction (MAE) [52,53]. The last is commonly used for isolation of active substances from plant material [54]. The parameters most affecting its efficacy, are the following: the type and volume of solvent, radiation potential, and extraction time and temperature of the process [53,55-58]. Usually organic solvents are used, and their volume depends on the kind and mass of a sample [53,58]. In comparison with Soxhlet extraction (Table 2), lesser amounts of solvent are needed, and the extraction time is shorter. The root of *Salvia miltiorrhizae* was extracted by the MAE technique in order to determine selected phenolic acids [59], as well as dry roots of *Eucommia ulmoides*, in which chlorogenic acid, among others, was determined by using a methanol and water mixture and conducting the extraction at 40°C [60]. A slightly lower temperature, 30°C, was used for determining phenolic compounds in the herbs of *Hypericum perforatum* and *Thymus vulgaris*. HCl solution was used as the extractant [61].

Methods for the determination of phenolic acids

For separation, purification and identification of phenolic compounds in plant materials numerous chromatographic methods have been applied [10]. Moreover, chromatographic techniques are also used for investigation of interactions of phenolic compounds with other food constituents.

High-performance liquid chromatography (HPLC)

Quantitative analysis of phenolic compounds in plant material depends on the chemical nature of the constituents, the method of extraction, particle size, time and conditions of storage of the plant

material prior to analysis, as well as on the determination method and the presence of interfering agents, such as fats, terpenes and chlorophyll [10]. Usually, chromatographic methods are applied, of which high-performance liquid chromatography [62,63] is the most common. In this method, different types of columns, mobile phases, column temperature and, to a smaller extent, the flow rate of the mobile phase, have been tested. Water, methanol and acetonitrile are the most common constituents of the mobile phase. Sometimes it is necessary to add modifiers, which facilitate the resolution of the components. The most popular modifiers are formic acid, ammonium acetate and acetic acid, the presence of which prevents tailing in the chromatograms. As far as the time of analysis is concerned, it is not a fixed parameter, because by modification of the flow rate of the mobile phase it is possible to extend or to shorten it. In the HPLC technique it is possible to use detection systems such as a UV-Vis spectrometer with either single wavelength or diode-array capability, chemiluminescence detector (CL), coulometric electrode array system (CEAD) and mass spectrometer (MS).

Reversed-phase, high-performance, liquid chromatography with UV-Vis detection was used by Waksmundzka-Hajnos *et al.* [30] for the analysis of the inflorescence of *Sambucus nigra* L. and the foliage of *Polygonum aviculare* L. Isocratic elution was applied with the use of two mobile phases. The first, used for *S. nigra*, was a 22:78 (v/v) mixture of methanol and orthophosphoric acid, and the other, applied for *P. aviculare* L., was a methanol/water (25:75 v/v) mixture, with addition of 1% of acetic acid. Phenolic compounds were detected at 520 nm.

Santos-Gomes *et al.* [50] applied HPLC for the analysis of phenolics in several plant species of *Lamiaceae*. The mobile phase was a mixture of acetonitrile, water and acetic acid (15:84:0.85 v/v) as solvent A and methanol as solvent B. Gradient elution was applied and the flow rate was set at 0.8 mL/min. The analytical wavelength was 280 nm, and identification of particular phenolic compounds was performed based on comparison of the retention times of the compounds with those of the standards.

Benetis *et al.* [40] have used HPLC with UV-Vis detection for the analysis of phenolic compounds in *Achillea millefolium* L. Gradient elution was applied with a mobile phase of acetonitrile/water, and trifluoroacetic acid (TFA) as a modifier to prevent ionization of phenolic groups. Before performing the analysis, two types of columns were tested: Xterra RP18 (Waters) and Ascentis RP-Amide Supelco; the latter column was chosen. The phenolic compounds were analysed at 25°C, and a 10 µL sample was injected into the column with a flow rate of 1.5 mL/min and detection at 360 nm.

Liquid chromatography with UV detection was also applied for the determination of phenolic acids in *Echinacea purpurea* [64], the leaves of lemon balm [65], aqueous extracts of *Hypericum perforatum* [66], and in 32 medicinal plants growing in Poland [67]. In the first case, the derivatives of caffeic acid were determined at 330 nm. In the second case, caffeic, ferulic and *p*-coumaric acids were determined at 325 nm, gallic, vanillic and syringic acids at 280 nm, and in the third case, all phenolic acids at 210 nm.

Liquid chromatography with diode array detection (DAD), in comparison with single wavelength UV/Vis detection, enables registration of the absorption spectrum of a compound over a wide range of wavelengths and determination of the absorbance maxima. This type of detection was applied for establishing the chemical composition of extracts of *Salvia halophila* [32], for identification and quantification of *p*-hydroxybenzoic acid derivatives in lemon balm (*Melissa officinalis*) [33], phenolic acids in the roots of *Salvia*

Table 3: Comparison of LC-MS with GC-MS for the analysis of phenolic compounds in medicinal plants [77].

Parameters	LC-MS	GC-MS
Time of sample preparation	20 min	180 min
Time of analysis	60 min	50 min
Range of linearity	limited	good
Selectivity	good	high
Limit of detection	5-15 ng/mL	10-80 ng/mL
Identification	possibility of calculation of empiric equation	mass spectral library for large group of compounds; fragmentation enables evaluation of molecular structure
Ruggedness of the system	satisfactory	very good

miltiorrhiza [68], in preparations of Chinese plants [69], and in plant-derived foods [27]. Another example of UV detection of polyphenolic compounds in HPLC was the use of a photodiode array detector by scanning between 200 and 400 nm, with a resolution of 1.2 nm [28].

In recent years, a growing interest has been gained in HPLC with chemiluminescent detection, the advantage of which is high sensitivity and selectivity. However, this method also has some drawbacks. First is its limited application, due to the small number of HPLC-CL chemiluminescent reactions. There are several chemical agents enabling derivatization, and in spite of the fact that they broaden the identification range for constituents not connected with chemiluminescent reactions, their excess or lack of integration can interfere with identification of analytes. Another disadvantage appears when the sample constituents are detected at a slow flow rate, and the mobile phase is incompatible with the chemiluminescent reactions; this precluding identification of these constituents. Another drawback of the method is that the intensity of chemiluminescence depends on such environmental factors as the kind of solvent, pH and ionic strength [70]. Despite those disadvantages, Cui *et al.* [70] decided to use this method in their research. They performed both isocratic and gradient elution at 25°C and a flow rate of 1.0 mL/min. The mobile phase was chosen out of three mixtures: acetonitrile-orthophosphoric acid-water, methanol-acetic acid-water, and phosphate-acetonitrile-methanol. It was found that when the concentration of acetic acid was 1.5%, the intensity of chemiluminescence was highest. Similar effect was obtained with a 35% methanolic solution and, accordingly, the 35:65 (v/v) of methanol-1.5% acetic acid mixture was selected as the mobile phase for the analysis. After completion of these measurements, it was found that peaks in the chromatograms registered using CL detection were higher than those obtained by the DAD detection system.

A less commonly used type of detection system is coulometric electrode array detector (CEAD). This detector was applied for determining ferulic and *p*-coumaric acids in the seeds of common flax and in the roots of nettle [71]. HPLC-CEAD is characterised by high sensitivity and precision, based on redox activity of the analyte [72].

In some laboratories, HPLC coupled with mass spectrometry (MS) is also used [73]. Guo *et al.* [74] have applied the LC-MS technique for the simultaneous determination of six phenolic acids in rat plasma after intravenous administration of a traditional Chinese medicinal preparation *Guanxinning* in a form of a lyophilised powder containing *Salvia miltiorrhiza* Bge. The same type of detection was used for determining caffeic acid in products from *Echinacea* sp. [45], chlorogenic acid in the leaves of lemon balm [65], and caffeic and protocatechuic acids in the roots of sage [75]. Furthermore, LC-MS and NMR techniques were also used for identification and structure establishment of eighteen phenolic acids and mono- and diglycosidic flavonoids [76].

Gas chromatography (GC)

Gas chromatography is one of the less common techniques used for the determination of phenolic acids. Using this technique, it is

possible to analyse only several small molecule phenolic acids (below 600 D) [66,77]. Moreover, GC requires high temperatures, which can lead to sample decomposition [78].

For the analysis of phenolic acids in plant materials, GC coupled with mass spectrometry (GC-MS) is very often used. In comparison with LC-MS, this method allows better selectivity, precision and accuracy, especially when small amounts of compounds have to be analysed in plant material [77].

GC-MS was applied by Fiamegos *et al.* [79], who determined the contents of phenolic acids in medicinal plant raw materials and their infusions. They used GC-MS instrumentation with SIM (*selective ion monitoring*). Owing to this modification of the method, the currents were registered only for selected ions with characteristic masses, typical of the studied analyte. The method with SIM is useful for determining constituents occurring at low concentrations in complex mixtures.

Proestos *et al.* [41] have also applied GC-MS for the determination of phenolic acids in aromatic plants, such as nettle and common rue. They used capillary gas chromatography, silylation as a derivatization procedure, and *N,O*-bis(trimethylsilyl)-trifluoro-acetamide and trimethylchlorosilane as silylating agents. Along with capillary gas chromatography (CGC-MS), they also used HPLC with UV detection at 280 nm.

A comparison of GC-MS with LC-MS (Table 3) has shown that the first method requires a time-consuming sample preparation step prior to analysis, but the time of analysis is shorter, and the linearity and selectivity of the method is much better than that in the case of LC/MS [77]. The sample preparation step includes dynamic sonication-assisted extraction of herbal samples, followed by the extracts being further treated by liquid-liquid extraction and derivatization. In GC-MS the compounds are identified by comparison of their spectra with those available in the mass spectra library.

It is also worth mentioning that, among the chromatographic methods, high-speed, counter-current chromatography (HSCCC) [80] was used by Yang *et al.* [81] for isolation and purification of phenolic acids originating from a Chinese medicinal plant, *Smilax china*.

Capillary electrophoresis (CE)

Recently, capillary electrophoresis (CE) has achieved growing significance in the analysis of phenolic compounds [82]. This method allows the separation and determination of polar substances of both ionic and non-ionic character, as well as non-polar, non-ionic substances. Advantages of CE are a small volume of electrolytes, short analysis time, high resolution potential and small sample weight [83-86]. In order to achieve proper resolution by this method, it is necessary to optimize such parameters as the type of buffer, its pH and concentration, the type of capillary and its volume, temperature of electrophoresis, voltage and the way of injection of a sample [87].

Each of these parameters will depend on CE methodology and on the chemical properties of the studied phenolic acid and its matrix.

Before analysis by capillary electrophoresis, the active compounds are extracted into solution, usually by supported liquid extraction (SLE) [37,44,88-92], supercritical fluid extraction (SFE) [93,94], and liquid-liquid extraction (LLE) [9,25]. The less common methods are microwave-assisted solvent extraction (MAE), taking only several minutes [84], and Soxhlet extraction [95].

Apart from CE, another popular technique applied for the determination of phenolic compounds, is *capillary zone electrophoresis* (CZE) [96,97]. This method was used, among others, for the determination of rosmarinic acid in commercial sage tea-bags [85]. Of the tested solvents, such as methanol, acetone and acetonitrile, the best was methanol. As the extraction technique, ultrasonication was used, and for detection, UV spectrometry at 210 nm. Resolution of the compounds was performed in a quartz capillary, 50 μm in diameter, and a solution of borate as a separating buffer was used. Beside CE and CZE, sometimes micellar electrokinetic chromatography (MEKC) is also used. By using this technique, phenolic acids were determined in the roots and pods of *Echinacea purpurea* [9], as well as in the herb of *Artemisia capillaris* [98]. *Capillary electrochromatography* (CEC) was applied for the analysis of extracts obtained from the flowers of chamomile [82].

As detection methods in capillary electrophoresis and its modifications, optical methods are used (fluorimetric, phosphorimetric, chemiluminescent, UV-Vis, IR and Raman spectrometries, and refractometry), as well as electrochemical methods (conductometric, potentiometric, amperometric and voltamperometric), and other, such as mass spectrometry and radiometry. The most frequently used detectors are those based on UV, MS and amperometric methods. They assure high sensitivity and selectivity.

CE with UV detection is commonly used for the analysis of phenolic acids in plant materials, and the most often applied electrolyte is a borate buffer of pH 9.2 [99]. This buffer was used for the determination of phenolic acids in *Strobili lupuli* [91], *Cortex fraxini* [92], and the fruits of sea buckthorn berries [100]. Less frequently, a MOPSO buffer solution is used; this was applied together with Tris solution and boric acid, pH 8.3, for the determination of chlorogenic acid in *Hypericum perforatum* [101]. Also a phosphate buffer with acetonitrile, pH 2.8, was used for the determination of caffeic and chlorogenic acids in *Matricaria chamomilla* L. [82].

MS detection is useful for identification of chemical compounds in mixtures as it can elucidate chemical and structural information about molecules from their molecular weight and distinctive fragmentation patterns [102]. In the CE method coupled with MS, resolution of the constituents can be regulated by changing the pH. This technique is

useful when other analytical techniques do not provide reliable results. Moreover, the CE-MS combination opens more analytical options, such as increase in sensitivity and specificity of the method. Application of this hyphenated method does not require using complicated sample preparation procedures and in comparison with the chromatographic methods, it is a good alternative for simultaneous analysis of phenolic compounds, because it provides rapid and efficient separations and uses reduced sample and solvent consumption [103].

Concluding remarks

Phenolic acids are secondary metabolites widespread in the plant kingdom. Because of their wide spectrum of biological activity, isolation, quantification and structure establishing of these compounds is still a crucial matter. On the one hand, there are methods of isolation of phenolic acids from medicinal and dietary plants, on the other, analytical techniques for quantitation of these secondary metabolites. Literature screening has shown that although Soxhlet and ultrasound-assisted (UAE) extractions are commonly used for isolation of polyphenols from plant matrices, alternative techniques of extraction, such as supercritical fluid (SFE) and accelerated solvent (ASE) are more and more often used. Non-questionable advantages of Soxhlet extraction are its low cost and good recovery, but there are some disadvantages, such as large volume of solvent, long time of extraction and tedious handling procedure. Recently introduced advanced techniques of extraction, such as SFE and ASE, are not as profitable as initially expected. Supercritical fluid extraction takes place under subcritical conditions, and this time-consuming and expensive procedure is limited to compounds of low or medium polarity. On the other hand, accelerated solvent extraction is performed with high extraction temperatures that may lead to degradation of thermolabile compounds. Furthermore, this technique is considered as a potential alternative to SFE for extraction of polar compounds.

Quantitation of phenolic acids is usually performed by liquid chromatography (LC) owing to the recent developments in this technique, especially when it is coupled with mass spectrometry (MS). Other advanced techniques used for quantitation of the secondary metabolites are gas chromatography coupled with mass spectrometry (GC-MS) and capillary electrophoresis (CE). All the above techniques have some limitations. For example, LC with DAD and MS provided short time analysis with less sample preparation, while GC method, which is used very often with MS detection, required a relatively high temperature, which can lead to sample decomposition. Moreover, GC can be used only for several small molecule phenolic acids (below 600 D). Recently, CE has achieved growing significance in quantitation of phenolic acids. This technique provides short analysis time and small volume of electrolytes, and the most common detectors are UV, MS and amperometric.

References

- [1] Lattanzio V, Lattanzio VMT, Cardinali A. (2006) Role of phenolics in the resistance mechanisms of plant against fungal pathogens and insects. *Phytochemistry: Advances in Research*, 23-67. Research Signpost, Trivandrum, Kerala, India.
- [2] Li Z-H, Wang Q, Ruan X, Pan C-D, Jiang D-A. (2010) Phenolics and plant allelopathy. *Molecules*, **15**, 8933-8952.
- [3] Alaerts G, Van Erps J, Pieters S, Dumarey M, van Nederkassel AM, Goodarzi M, Smeyers-Verbeke J, Van der Heyden Y. (2012) Similarity analyses of chromatographic fingerprints as tools for identification and quality control of green tea. *Journal of Chromatography B*, **910**, 61-70.
- [4] Ma T, Huang C, Meng X, Zhang Q, Zhang L, Lv X, Jin Y, Xie J, Li J. (2011) Fingerprint analysis of Hawk-tea by high-performance liquid chromatography. *Food Chemistry*, **129**, 551-556.
- [5] Xie B, Gong T, Tang M, Mi D, Zhang X, Liu J, Zhang Z. (2008) An approach based on HPLC/fingerprint and chemometrics to quality consistency evaluation of *Liuwei Dihuang* pills produced by different manufactures. *Journal of Pharmaceutical and Biomedical Analysis*, **48**, 1261-1266.
- [6] Şahin S, Demir C, Malyer H. (2011) Determination of phenolic compounds in *Prunella* L. by liquid chromatography-diode array detection. *Journal of Pharmaceutical and Biomedical Analysis*, **55**, 1227-1230.
- [7] Kaefer CM, Milner JA. (2008) The role of herbs and spices in cancer prevention. *Journal of Nutritional Biochemistry*, **19**, 347-361.

- [8] Inbaraj BS, Lu H, Kao TH, Chen BH. (2010) Simultaneous determination of phenolic acids and flavonoids in *Lycium barbarum* Linnaeus by HPLC-DAD-ESI-MS. *Journal of Pharmaceutical and Biomedical Analysis*, **51**, 549–556.
- [9] Pomponio R, Gotti R, Hudaib M, Cavrini V. (2002) Analysis of phenolic acids by micellar electrokinetic chromatography: application to *Echinacea purpurea* plant extracts. *Journal of Chromatography A*, **945**, 239–247.
- [10] Naczki M, Shahidi F. (2006) Phenolics in cereals, fruits and vegetables: Occurrence, extraction and analysis. *Journal of Pharmaceutical and Biomedical Analysis*, **41**, 1523–1542.
- [11] Mattila P, Hellstrom J. (2007) Phenolic acids in potatoes, vegetables, and some of their products. *Journal of Food Composition and Analysis*, **20**, 152–160.
- [12] Robbins RJ. (2003) Phenolic acids in foods: An overview of analytical methodology. *Journal of Agricultural and Food Chemistry*, **51**, 2866–2887.
- [13] Carlsen SC, Kudsk P, Laursen B, Mathiasen SK, Mortensen AG, Fomsgaard IS. (2009) Allelochemicals in rye (*Secale cereale* L.): cultivar and tissue differences in the production of benzoxazinoids and phenolic acids. *Natural Product Communications*, **4**, 199–208.
- [14] Zgórk G, Kawka S. (2001) Application of conventional UV, photodiode array (PDA) and fluorescence (FL) detection to analysis of phenolic acids in plant material and pharmaceutical preparations. *Journal of Pharmaceutical and Biomedical Analysis*, **24**, 1065–1072.
- [15] Atoui AK, Mansouri A, Boskou G, Kefalas P. (2005) Tea and herbal infusions: Their antioxidant activity and phenolic profile. *Food Chemistry*, **89**, 27–36.
- [16] Robbins R, Bean SR. (2004) Development of a quantitative high-performance liquid chromatography-photodiode array detection measurement system for phenolic acids. *Journal of Chromatography A*, **1038**, 97–105.
- [17] Zheng W, Wang S. (2001) Antioxidant activity and phenolic compounds in selected herbs. *Journal of Agricultural and Food Chemistry*, **49**, 5165–5170.
- [18] Slavin J, Marquart L, Jacobs D. (2000) Consumption of whole-grain food and decreased risk of cancer: proposed mechanisms. *Cereal Foods Worlds*, **45**, 54–58.
- [19] Weng CJ, Yen GC. (2012) Chemopreventive effects of dietary phytochemicals against cancer invasion and metastasis: phenolic acids, monophenol, polyphenol, and their derivatives. *Cancer Treatment Reviews*, **38**, 76–87.
- [20] Huang L, Lin C, Li A, Wei B, Teng J, Li L. (2010) Pro-coagulant activity of phenolic acids isolated from *Blumea riparia*. *Natural Product Communications*, **5**, 1263–1266.
- [21] Xu M, Han J, Li HF, Fan L, Liu AH, Guo DA. (2008) Analysis on the stability of total phenolic acids and salvianolic acid B from *Salvia miltiorrhiza* by HPLC and HPLC-MSn. *Natural Product Communications*, **3**, 669–676.
- [22] Lafay S, Gil-Izquierdo A. (2008) Bioavailability of phenolic acids. *Phytochemistry Reviews*, **7**, 301–311.
- [23] Rostagno MA, D'Arrigo M, Martinez JA. (2010) Combinatory and hyphenated sample preparation for the determination of bioactive compounds in foods. *Trends in Analytical Chemistry*, **29**, 553–561.
- [24] Kivilompolo M, Hyötyläinen T. (2009) On-line coupled dynamic sonication-assisted extraction-liquid chromatography for the determination of phenolic acids in *Lamiaceae* herbs. *Journal of Chromatography A*, **1216**, 892–896.
- [25] Hurtado-Fernández E, Gómez-Romero M, Carrasco-Pancorbo A, Fernández-Gutiérrez A. (2010) Application and potential of capillary electrophoresis methods to determine antioxidant phenolic compounds from plant food material. *Journal of Pharmaceutical and Biomedical Analysis*, **53**, 1130–1160.
- [26] Plazonić A, Bucar F, Maleš Ž, Mornar A, Nigović B, Kujundžić N. (2009) Identification and quantification of flavonoids and phenolic acids in burr parsley (*Caucalis platycarpus* L.), using high-performance liquid chromatography with diode array detection and electrospray ionization mass spectrometry. *Molecules*, **14**, 2466–2490.
- [27] Mattila P, Kumpulainen J. (2002) Determination of free and total phenolic acids in plant-derived foods by HPLC and diode-array detection. *Journal of Agricultural and Food Chemistry*, **50**, 3660–3667.
- [28] Belščak-Cvitanović A, Stojanović R, Manojlović V, Komes D, Juranović Cindrić I, Nedović V, Bugarski B. (2011) Encapsulation of polyphenolic antioxidants from medicinal plant extracts in alginate-chitosan system enhanced with ascorbic acid by electrostatic extrusion. *Food Research International*, **44**, 1094–1101.
- [29] You Q, Wang B, Feng Ch, Huang Z, Wang X, Luo P. (2011) Comparison of anthocyanins and phenolics in organically and conventionally grown blueberries in selected cultivars. *Food Chemistry*, **125**, 201–208.
- [30] Waksmundzka-Hajnos M, Oniszczyk A, Szewczyk K, Wianowska D. (2007) Effect of sample preparation methods on the HPLC quantitation of some phenolic acids in plant materials. *Acta Chromatographica*, **19**, 227–237.
- [31] Akkol EK, Göger F, Kosar M, Can Baser KH. (2008) Phenolic composition and biological activities of *Salvia halophila* and *Salvia virgata* from Turkey. *Food Chemistry*, **108**, 942–949.
- [32] Koşar M, Göger F, Hüsni Can Başer K. (2011) In vitro antioxidant properties and phenolic composition of *Salvia halophila* Hedge from Turkey. *Food Chemistry*, **129**, 374–379.
- [33] Karasová G, Lehotay J. (2005) Chromatographic determination of derivatives of *p*-hydroxybenzoic acid in *Melissa officinalis* by HPLC. *Journal of Liquid Chromatography & Related Technologies*, **28**, 2421–2431.
- [34] Rajaei A, Barzegar M, Hamidi Z, Sahari MA. (2010) Optimization of extraction conditions of phenolic compounds from pistachio (*Pistachia vera*) green hull through response surface method. *Journal of Agricultural Science and Technology*, **12**, 605–615.
- [35] Khan MK, Abert-Vian M, Fabiano-Tixier AS, Dangles O, Chemat F. (2010) Ultrasound-assisted extraction of polyphenols (flavanone glycosides) from orange (*Citrus sinensis* L.) peel. *Food Chemistry*, **119**, 851–858.
- [36] Kagan IA. (2011) Effects of pH, sample size, and solvent partitioning on recovery of soluble phenolic acids and isoflavonoids in leaves and stems of red clover (*Trifolium pratense* cv. Kenland). *Natural Product Communications*, **6**, 1657–1660.
- [37] Casazza AA, Alikbarian B, Mantegna S, Cravotto G, Perego P. (2010) Extraction of phenolics from *Vitis vinifera* wastes using non-conventional techniques. *Journal of Food Engineering*, **100**, 50–55.
- [38] Garcia-Salas P, Morales-Soto A, Segura-Carretero A, Fernandez-Gutierrez A. (2010) Phenolic-compound-extraction systems for fruit and vegetable samples. *Molecules*, **15**, 8813–8826.
- [39] Pawar N, Pai S, Nimbalkar M, Dixit G. (2011) RP-HPLC analysis of phenolic antioxidant compound 6-gingerol from different ginger cultivars. *Food Chemistry*, **126**, 1330–1336.
- [40] Benetis R, Radusiene J, Jakstas V, Janulis V, Malinauskas F. (2008) Development of a RP-HPLC method for the analysis of phenolic compounds in *Achillea millefolium* L. *Journal of Liquid Chromatography and Related Technologies*, **31**, 596–610.
- [41] Proestos C, Komaitis M. (2006) Ultrasonically assisted extraction of phenolic compounds from aromatic plants: Comparison with conventional extraction techniques. *Journal of Food Quality*, **29**, 567–582.
- [42] Žiaková A, Brandšteterová E, Blahová E. (2003) Matrix solid-phase dispersion for the liquid chromatographic determination of phenolic acids in *Melissa officinalis*. *Journal of Chromatography A*, **983**, 271–275.
- [43] Teixeira DM, Patão RF, Coelho AV, Teixeira da Costa C. (2006) Comparison between sample disruption methods and solid-liquid extraction (SLE) to extract phenolic compounds from *Ficus carica* leaves. *Journal of Chromatography A*, **1103**, 22–28.

- [44] Pinelo M, Ruiz-Rodríguez A, Sineiro J, Señoráns FJ, Reglero G, Núñez MJ. (2007) Supercritical fluid and solid-liquid extraction of phenolic antioxidants from grape pomace: a comparative study. *European Food Research and Technology*, **226**, 199-205.
- [45] Castro-Vargas HI, Rodríguez-Varela LI, Ferreira SRS, Parada-Alfonso F. (2010) Extraction of phenolic fraction from guava seeds (*Psidium guajava* L.) using supercritical carbon dioxide and co-solvents. *Journal of Supercritical Fluids*, **51**, 319-324.
- [46] Kazazi H, Rezaei K. (2009) Effect of various parameters on the selective extraction of main components from Hyssop using supercritical fluid extraction (SFE). *Food Science and Technology Research*, **15**, 645-652.
- [47] Dai J, Mumper RJ. (2010) Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules*, **15**, 7313-7352.
- [48] Li W, Qui Y, Patterson CA, Beta T. (2011) The analysis of phenolic constituents in glabrous canaryseed groats. *Food Chemistry*, **127**, 10-20.
- [49] Fernand VE, Dinh DT, Washington SJ, Fakayode SO, Losso JN, Van Ravenswaay RO, Warner IM. (2008) Determination of pharmacologically active compounds in root extracts of *Cassia alata* L. by use of high performance liquid chromatography. *Talanta*, **74**, 896-902.
- [50] Santos-Gomes PC, Seabra RM, Andrade PB, Fernandes-Ferreira M. (2003) Determination of phenolic antioxidant compounds produced by calli and cell suspensions of sage (*Salvia officinalis* L.). *Journal of Plant Physiology*, **160**, 1025-1032.
- [51] Kim S, Yun EJ, Bak JS, Lee H, Lee SJ, Kim CT, Lee J-H, Kim KH. (2010) Response surface optimized extraction and chromatographic purification of rosmarinic acid from *Melissa officinalis* leaves. *Food Chemistry*, **121**, 521-526.
- [52] Jiang Y, David B, Tu P, Barbain Y. (2010) Recent analytical approaches in quality control of traditional Chinese medicines: A review. *Analytica Chimica Acta*, **675**, 9-18.
- [53] Madej K. (2009) Microwave-assisted and cloud-point extraction in determination of drugs and other bioactive compounds. *Trends in Analytical Chemistry*, **28**, 436-446.
- [54] Deng C, Liu N, Gao M, Zhang X. (2007) Recent developments in sample preparation techniques for chromatography analysis of traditional Chinese medicines. *Journal of Chromatography A*, **1153**, 90-96.
- [55] Eskilsson CS, Björklund E. (2000) Analytical-scale microwave-assisted extraction. *Journal of Chromatography A*, **902**, 227-250.
- [56] Chan C, Yusoff R, Ngoh GC, Kung FW. (2011) Microwave-assisted extraction of active ingredients from plants. *Journal of Chromatography A*, **1218**, 6213-6225.
- [57] Mandal V, Mohan Y, Hemalatha S. (2007) Microwave assisted extraction-an innovative and promising extraction tool for medicinal plant research. *Pharmacognosy Reviews*, **1**, 7-18.
- [58] Tatke P, Jaiswal Y. (2011) An overview of microwave assisted extraction and its applications in herbal drug research. *Research Journal of Medicinal Plant*, **5**, 21-31.
- [59] Fang X, Wang J, Zhou H, Jiang X, Zhu L, Gao X. (2009) Microwave-assisted extraction with water for fast extraction and simultaneous RP-HPLC determination of phenolic acids in radix *Salvia miltiorrhizae*. *Journal of Separation Science*, **32**, 2455-2466.
- [60] Li H, Chen B, Zhang Z, Yao S. (2004) Focused microwave assisted extraction and HPLC determination of effective constituents in *Eucommia ulmoides* Oliv. *Talanta*, **63**, 659-665.
- [61] Sterbova D, Matejicek D, Vleck J, Kuban V. (2004) Combined microwave assisted isolation and solid-phase purification procedures prior to the chromatographic determination of phenolic compounds in plant materials. *Analytica Chimica Acta*, **513**, 435-444.
- [62] Yin L, Lu B, Qi Y, Xu L, Han X, Xu Y, Peng J, Sun C. (2009) Simultaneous determination of 11 active components in two well-known traditional Chinese medicines by HPLC coupled with diode array detection for quality control. *Journal of Pharmaceutical and Biomedical Analysis*, **49**, 1101-1108.
- [63] Robards K. (2003) Strategies for the determination of bioactive phenols in plants, fruits and vegetables. *Journal of Chromatography A*, **1000**, 657-691.
- [64] Iranshahi M, Amanzadeh Y. (2008) Rapid isocratic HPLC analysis of caffeic acid derivatives from *Echinacea purpurea* cultivated in Iran. *Chemistry of Natural Compounds*, **44**, 190-193.
- [65] Marques V, Farah A. (2009) Chlorogenic acids related compounds in medicinal plants and infusions. *Food Chemistry*, **113**, 1370-1376.
- [66] Temerdashev ZA, Frolova NA, Kolychev IA. (2011) Determination of phenolic compounds in medicinal herbs by reserved-phase HPLC. *Journal of Analytical Chemistry*, **66**, 407-414.
- [67] Wojdyło A, Oszmiański J, Czemerys R. (2007) Antioxidant activity and phenolic compounds in 32 selected herbs. *Food Chemistry*, **105**, 940-949.
- [68] Liu A-H, Li L, Xu M, Lin Y-H, Guo H-Z, Guo D-A. (2006) Simultaneous quantification of six major phenolic acids in the roots of *Salvia miltiorrhiza* and four related traditional Chinese medicinal preparations by HPLC-DAD method. *Journal of Pharmaceutical and Biomedical Analysis*, **41**, 48-56.
- [69] Li Y, Zhu J, Wan L, Yu Q, Li X, Cheng Z, Guo C. (2010) Combinative method using HPLC fingerprint and quantitative analyses for quality consistency evaluation of an herbal medicinal preparation produced by different manufactures. *Journal of Pharmaceutical and Biomedical Analysis*, **52**, 597-602.
- [70] Cui H, Zhou J, Xu F, Lai Ch-Z, Wan G-H. (2004) Determination of phenolic compounds using high-performance liquid chromatography with Ce^{4+} -Tween 20 chemiluminescence detection. *Analytica Chimica Acta*, **511**, 273-279.
- [71] Kraushofer T, Sontag G. (2002) Determination of some phenolic compounds in flax seed and nettle roots by HPLC with coulometric electrode array detection. *European Food Research and Technology*, **215**, 529-533.
- [72] Ma L, Zhang X, Zhang H, Gan Y. (2007) Development of a fingerprint of *Salvia miltiorrhiza* Bunge by high-performance liquid chromatography with a coulometric electrode array system. *Journal of Chromatography B*, **846**, 137-146.
- [73] Steinmann D, Ganzera M. (2011) Recent advances on HPLC/MS in medicinal plant analysis. *Journal of Pharmaceutical and Biomedical Analysis*, **55**, 744-757.
- [74] Guo XR, Chen XH, Li L, Shen ZD, Wang XL, Zheng P, Duan FX, Ma YF, Bi KS. (2008) LC-MS determination and pharmacokinetic study of six phenolic components in rat plasma after taking traditional Chinese medicinal-preparation: Guanxinning lyophilized powder of injection. *Journal of Chromatography B*, **873**, 51-58.
- [75] Wu J-L, Yee LP, Jiang Z-H, Cai Z. (2007) One single LC-MS/MS analysis for both phenolic components and tanshinones in Radix *Salviae Miltiorrhizae* and its medicinal products. *Talanta*, **73**, 656-661.
- [76] Williams JD, Wojcińska M, Calabria LM, Linse K, Clevinger JA, Mabry TJ. (2009) The flavonoids and phenolic acids of the genus *Silphium* and their chemosystematic value. *Natural Product Communications*, **4**, 435-446.
- [77] Kivilompolo M, Obürka V, Hyötyläinen T. (2007) Comparison of GC-MS and LC-MS methods for the analysis of antioxidant phenolic acids in herbs. *Analytical and Bioanalytical Chemistry*, **388**, 881-887.
- [78] Lin J-T, Liu S-H, Tsay GJ, Yang D-J. (2010) Composition of flavonoids and phenolic acids in *Glycin tomentella* Hayata cultivated in various soils. *Food Chemistry*, **121**, 659-665.
- [79] Fiamegos YC, Nanos CG, Vervoot J, Stalikas CD. (2004) Analytical procedure for the in-vial derivatization-extraction of phenolic acids and flavonoids in methanolic and aqueous plant extracts followed by gas chromatography with mass-selective detection. *Journal of Chromatography A*, **1041**, 11-18.
- [80] Ito Y. (2005) Golden rules and pitfalls in selecting optimum conditions for high-speed counter-current chromatography. *Journal of Chromatography A*, **1065**, 145-168.
- [81] Yang C, Tang Q, Liu J, Zhang Z, Liu W. (2008) Preparative isolation and purification of phenolic acids from *Smilax china* by high-speed counter-current chromatography. *Separation and Purification Technology*, **61**, 474-478.

- [82] Fonseca F, Tavares MFM, Horváth C. (2007) Capillary electrochromatography of selected phenolic compounds of *Chamomilla recutita*. *Journal of Chromatography A*, **1154**, 390-399.
- [83] Gan Z, Chen Q, Fu Y, Chen G. (2012) Determination of bioactive constituents in *Flos Sophorae Immaturus* and *Cortex Fraxini* by capillary electrophoresis in combination with far infrared-assisted solvent extraction. *Food Chemistry*, **130**, 1122-1126.
- [84] Duan H, Chen Y, Chen G. (2010) Far infrared-assisted extraction followed by capillary electrophoresis for the determination of bioactive constituents in the leaves by *Lycium barbarum* Linn. *Journal of Chromatography A*, **1217**, 4511-4516.
- [85] Başkan S, Öztekin N, Erim FB. (2007) Determination of carnosic acid and rosmarinic acid in sage by capillary electrophoresis. *Food Chemistry*, **101**, 1748-1752.
- [86] Crego AL, Ibáñez E, Garcia E, Rodriguez de Pablos R, Señarás FJ, Reglero G, Cifuentes A. (2004) Capillary electrophoresis separation of rosemary antioxidants from subcritical water extracts. *European Food Research and Technology*, **219**, 549-555.
- [87] Ballus CA, Meinhardt AD, Bruns RE, Godoy H. (2011) Use of multivariate statistical techniques to optimize the simultaneous separation of 13 phenolic compounds from extra-virgin olive oil by capillary electrophoresis. *Talanta*, **83**, 1181-1187.
- [88] Teixeira DM, Teixeira de Costa C. (2005) Novel methods to extract flavonones and xanthenes from the root bark of *Maclura pomifera*. *Journal of Chromatography A*, **1062**, 175-181.
- [89] Peng Y, Yuan J, Liu F, Ye J. (2005) Determination of active components in rosemary by capillary electrophoresis with electrochemical detection. *Journal of Pharmaceutical and Biomedical Analysis*, **39**, 431-437.
- [90] Šafra J, Pospíšilova M, Honegr J, Spilkova J. (2007) Determination of selected antioxidants in *Melissae herba* by isotachopheresis and capillary zone electrophoresis in the column-coupling configuration. *Journal of Chromatography A*, **1171**, 124-132.
- [91] Helmja K, Vaher M, Pussa T, Kamsol K, Orav A, Kaljurand M. (2007) Bioactive components of the hop strobilus: comparison of different extraction methods by capillary electrophoretic and chromatographic methods. *Journal of Chromatography A*, **1155**, 222-229.
- [92] Wang XK, He YZ, Qian LL. (2007) Determination of polyphenol components in herbal medicines by micellar electrokinetic capillary chromatography with Tween 20. *Talanta*, **74**, 1-6.
- [93] Ibáñez E, Cifuentes A, Crego AL, Señarás FJ, Cavero S, Reglero G. (2000) Combined use of supercritical fluid extraction, micellar electrokinetic chromatography, and reverse phase high performance liquid chromatography for the analysis of antioxidants from rosemary (*Rosmarinus officinalis* L.). *Journal of Agricultural and Food Chemistry*, **48**, 4060-4065.
- [94] Vaher M, Koel M. (2003) Separation of polyphenolic compounds extracted from plant matrices using capillary electrophoresis. *Journal of Chromatography A*, **990**, 225-230.
- [95] Yue ME, Jiang TF, Shi YP. (2004) Fast determination of flavonoids in *Hippophae rhamnoides* and its medicinal preparation by capillary zone electrophoresis using dimethyl- β -cyclodextrin as modifier. *Talanta*, **62**, 695-699.
- [96] Lee ISL, Boyce MC, Breadmore MC. (2011) A rapid quantitative determination of phenolic acids in *Brassica oleracea* by capillary zone electrophoresis. *Food Chemistry*, **127**, 797-801.
- [97] Bonoli M, Pelillo M, Lercker G. (2003) Fast separation and determination of carnosic acid and rosmarinic acid in different rosemary (*Rosmarinus officinalis*) extracts by capillary zone electrophoresis with ultra violet-diode array detection. *Chromatographia*, **57**, 505-512.
- [98] Sheu SJ, Chieh CL, Weng WC. (2001) Capillary electrophoretic determination of the constituents of *Artemisiae Capillaris herba*. *Journal of Chromatography A*, **911**, 285-293.
- [99] Gotti R. (2011) Capillary electrophoresis of phytochemical substances in herbal drugs and medicinal plants. *Journal of Pharmaceutical and Biomedical Analysis*, **55**, 775-801.
- [100] Gorbatsová J, Lougas T, Vokk R, Kaljurand M. (2007) Comparison of the contents of various antioxidants of sea buckthorn berries using CE. *Electrophoresis*, **28**, 4136-4142.
- [101] Hamoudova R, Pospíšilova M, Spilkova J. (2006) Analysis of selected constituents in methanolic extracts of *Hypericum perforatum* collected in different localities by capillary ITP-CZE. *Electrophoresis*, **27**, 4820-4826.
- [102] McMahon G. (2007) *Analytical Instrumentation. A Guide to Laboratory, Portable and Miniaturized Instruments*. Wiley, Chichester, 36-49.
- [103] Arráez-Román D, Gómez-Caravaca AM, Gómez-Romero M, Segura-Carretero A, Fernández-Gutiérrez A. (2006) Identification of phenolic compounds in rosemary honey using solid-phase extraction by capillary electrophoresis-electrospray ionization-mass spectrometry. *Journal of Pharmaceutical and Biomedical Analysis*, **41**, 1648-1656.

Natural Product Communications

2013

Volume 8

Natural Product Communications 8 (1-12) 1-1830 (2013)

ISSN 1934-578X (print)

ISSN 1555-9475 (online)

Natural Product Communications

Contents of Volume 8 2013

Number 1

Argan Tree Metabolites and Argan Oil (Guest Editor: Dom Guillaume)

The Social and Environmental Context of Argan Oil Production Yann le Polain de Waroux	1
Climatic Conditions and Herbivory Effects on Morphological Plasticity of <i>Argania spinosa</i> Fatima Ain-Lhout, Maria Zunzunegui, Mari Cruz Díaz Barradas, Juan Jáuregui, Tarik Tagma and Said Boutaleb	5
Some Secrets of <i>Argania spinosa</i> Water Economy in a Semiarid Climate Mari Cruz Díaz Barradas, Maria Zunzunegui, Mari Paz Esquivias, Said Boutaleb, Javier Valera-Burgos, Tarek Tagma and Fátima Ain-Lhout	11
Germination Success and Seedling Development of <i>Argania spinosa</i> under Different Climatic Conditions and Browsing Intensity Maria Zunzunegui, Juan Jáuregui, Fatima Ain-Lhout, Said Boutaleb, Leonor Álvarez-Cansino and MariPaz Esquivias	15
Modeling of the Distribution of Heavy Metals and Trace Elements in Argan Forest Soil and Parts of Argan Tree Faez A. E. Mohammed, Rahma Bchitou, Mohamed Boulmane, Ahmed Bouhaouss and Dominique Guillaume	21
Can Fruit-form be a Marker for Argan Oil Production? Saïd Gharby, Hicham Harhar, Badr Eddine Kartah, Hanae El Monfalouti, Clément Denhez, Miloudi Hilali, Dom Guillaume and Zoubida Charrouf	25
Chemical Changes in Extra Virgin Argan Oil after Thermal Treatment Saïd Gharby, Hicham Harhar, Badr Eddine Kartah, Dom Guillaume and Zoubida Charrouf	29
Volatile Compound Formation During Argan Kernel Roasting Hanae El Monfalouti, Zoubida Charrouf, Manuela Giordano, Dominique Guillaume, Badreddine Kartah, Hicham Harhar, Saïd Gharby, Clément Denhez and Giuseppe Zeppa	33
Quality Parameters for Cold Pressed Edible Argan Oils Bertrand Matthäus	37
Triterpenoids from <i>Argania spinosa</i>: 20 Years of Research Imane Chafchaoui-Moussaoui, Zoubida Charrouf and Dom Guillaume	43
Argan Oil-contained Antioxidants for Human Mitochondria Luis C. López, Carmen Cabrera-Vique, Carmen Venegas, Laura García-Corzo, Marta Luna-Sánchez, Dario Acuña-Castroviejo and Germaine Escames	47
Effect of Argan and Olive Oil Consumption on the Hormonal Profile of Androgens Among Healthy Adult Moroccan Men Abdelfettah Derouiche, Ali Jafri, Issam Driouch, Mohammed El Khasmi, Ahmed Adlouni, Nada Benajiba, Youssef Bamou, Rachid Saïle and Mohammed Benouhoud	51
Argan Oil and Postmenopausal Moroccan Women: Impact on the Vitamin E Profile Hanae El Monfalouti, Zoubida Charrouf, Asma El Hamdouchi, Hanane Labraimi, Imane Chafchaoui-Moussaoui, Badreddine Kartah, Khalid El Kari, Yahya Bensouda, Abdelfettah Derouich, Sylvie Dodin, Clément Denhez, Dom Guillaume and Hassan Agnaou	55

Microbial Transformations of Isophorone by <i>Alternaria alternata</i> and <i>Neurospora crassa</i> Ismail Kiran, Özge Özşen, Turgay Çelik, Semra İlhan, Bükay Yenice Gürsu and Fatih Demirci	59
Headspace, Solid-phase Micro-extraction, Gas Chromatographic-Mass Spectrometric Analysis of Terpenoids in the Latex of <i>Euphorbia</i> Species Asmita V. Patel, Stephen Sumner, H. Leslie Thompson, Gerald Blunden, David Wright, Liu Jun-feng and Zan Jun-feng	63
Trichodermaerin, a New Diterpenoid Lactone from the Marine Fungus <i>Trichoderma erinaceum</i> Associated with the Sea Star <i>Acanthaster planci</i> Zhong-Liang Xie, Hou-Jin Li, Lai-You Wang, Wan-Ling Liang, Wei Liu and Wen-Jian Lan	67
Chemical Composition and Biological Activities of Soldiers of the Brazilian Termite Species, <i>Nasutitermes macrocephalus</i> (Isoptera: Natutitermitinae) Márcia N. S. de la Cruz, Helvécio M. S. Júnior, Denilson F. Oliveira, Leticia V. Costa-Lotufo, Antonio G. Ferreira, Daniela S. Alviano and Claudia M. Rezende	69
Evaluation of the Quality of Chinese and Vietnamese Cassia Using LC-MS and Multivariate Analysis Ken Tanaka, Feng Li, Yasuhiro Tezuka, Shiro Watanabe, Nobuo Kawahara and Hiroaki Kida	75
Norcucurbitane Triterpenoids from the Fruits of <i>Momordica charantia</i> var. <i>abbreviata</i> Yun-Wen Liao, Chiy-Rong Chen, Jue-Liang Hsu, Yun-Sheng Lin, Hsueh-Ling Cheng, Wen-Ling Shih, Yueh-Hsiung Kuo and Chi-I Chang	79

Fusaodavinin, a Novel Metabolite Containing Sulfur from the Endophytic Fungus <i>Fusarium</i> sp. (CTGU-ZL-34) Liang Zhang, Yanhui Liu, Zhangshuang Deng, Zhiyong Guo, Jianfeng Chen, Xuan Tu and Kun Zou	83
Summation Solute Hydrogen Bonding Acidity Values for Hydroxyl Substituted Flavones Determined by NMR Spectroscopy William L. Whaley, Ekua M. Okoso-amaa, Cody L. Womack, Anna Vladimirova, Laura B. Rogers, Margaret J. Risher and Michael H. Abraham	85
Phenolic Content and DPPH Radical Scavenging Activity of the Flowers and Leaves of <i>Trifolium repens</i> Agnieszka Kicel and Maria Wolbiś	99
Kenganthranol F, a new Anthranol from <i>Psorospermum aurantiacum</i> Gesquiere M. Tiani, Ishtiaq Ahmed, Karsten Krohn, Ivan R. Green and Augustin E. Nkengfack	103
Inhibition of <i>In Vitro</i> Leukotriene B₄ Biosynthesis in Human Neutrophil Granulocytes and Docking Studies of Natural Quinones Premysl Landa, Zsotia Kutil, Veronika Temml, Jan Malik, Ladislav Kokoska, Ute Widowitz, Marie Pribylova, Marcela Dvorakova, Petr Marsik, Daniela Schuster, Rudolf Bauer and Tomas Vanek	105
Effect of <i>Hypogymnia physodes</i> Extracts and their Depsidones on Micronucleus Distribution in Human Lymphocytes Igor Ž. Stojanović, Miroslava Stanković, Olga Jovanović, Goran Petrović, Andrija Šmelcerović and Gordana S. Stojanović	109
A New Lyoniresinol Derivative from <i>Smilax microphylla</i> Li-Sha Liu, Hui-Lian Huang, Rong-Hua Liu, Gang Ren, Feng Shao, Yao-Hui Ye and Tao Lin	113
Anti-metastatic Activities of Bibenzyls from <i>Dendrobium pulchellum</i> Pithi Chanvorachote, Akkarawut Kowitdamrong, Thidarat Ruanghirun, Boonchoo Sritularak, Chutichot Mungmee and Kittisak Likhitwitayawuid	115
Synthesis of 3,5,3',4'-Tetrahydroxy-trans-stilbene-4'-O-β-D-glucopyranoside by Glucosyltransferases from <i>Phytolacca americana</i> Tomoya Iwakiri, Hiroya Imai, Hiroki Hamada, Toru Nakayama and Shin-ichi Ozaki	119
Challenges of Curcumin Bioavailability: Novel Aerosol Remedies Parasuraman Aiya Subramani and Venkata R. Narala	121
Molecular Analysis of <i>Vitex</i> Species Using Candidate DNA Barcoding and PCR-RFLP of the <i>matK</i> Gene for Authentication of <i>Vitex glabrata</i> Waranyoo Phoolcharoen and Suchada Sukrong	125
Volatiles from <i>Syzygium paniculatum</i> Fruit Clara E. Quijano-Célis, Daniel Echeverri-Gil, Yinet Ruiz and Jorge A. Pino	129
Seasonal Variations in the Composition of the Essential Oils of Rosemary (<i>Rosmarinus officinalis</i>, Lamiaceae) Dmitar Lakušić, Mihailo Ristić, Violeta Slavkovska and Branislava Lakušić	131
Chemical Compositions and Antimicrobial and Antioxidant Activities of the Essential Oils from <i>Magnolia grandiflora</i>, <i>Chrysactinia mexicana</i>, and <i>Schinus molle</i> Found in Northeast Mexico Laura Guerra-Boone, Rocío Álvarez-Román, Ricardo Salazar-Aranda, Anabel Torres-Cirio, Verónica Mayela Rivas-Galindo, Noemí Waksman de Torres, Gloria María González González and Luis Alejandro Pérez-López	135
Chemical Composition and Biological Activities of the Essential Oil from <i>Calamintha nepeta</i> Plants from the Wild in Southern Italy Emilia Mancini, Laura De Martino, Hanna Malova and Vincenzo De Feo	139
Composition and <i>in-vitro</i> Cytotoxic Activities of the Leaf Essential Oil of <i>Beilschmiedia erythrophloia</i> from Taiwan Yu-Chang Su and Chen-Lung Ho	143

Number 2

Drimendiol, A Drimane Sesquiterpene with Quorum Sensing Inhibition Activity Cristian Paz, Gerardo Cárcamo, Mario Silva, José Becerra, Homero Urrutia and Katherine Sossa	147
Microbial Transformation of Curcumin by <i>Aspergillus niger</i> Chen Li-Xia, Zhang Hui, Zhao Qian, Yin Shi-Yu, Zhang Zhong, Li Tian-Xian and Qiu Feng	149
Chemopreventive Effect of Sarcophine-diol on NOR-1-Induced TPA-Promoted Skin Carcinogenesis in Female HOS:HR-1 Mice Pawel T. Szymanski, Safwat A. Ahmed, Sherief Khalifa, Harukuni Tokuda, Eiichiro Ichiishi, Akira Iida, Nobutaka Suzuki and Hesham Fahmy	153
Carmichaeline A: A New C₂₀-diterpenoid Alkaloid from <i>Aconitum carmichaeli</i> Shu-hua Li, Jun-ru Xiong, Yuan-qin Zhang, Qing-xiang Xiang and Feng-zheng Chen	155
Steroidal Saponins from <i>Dracaena marginata</i> Abdelmalek Rezgui, Anne-Claire Mitaine-Offer, David Pertuit, Tomofumi Miyamoto, Chiaki Tanaka, Stéphanie Delemasure, Patrick Dutartre and Marie-Aleth Lacaille-Dubois	157
Zephyrabetaine: A New Betaine-type Amaryllidaceae Alkaloid from <i>Zephyranthes grandiflora</i> Deepali Katoch, Dharmesh Kumar, Upendra Sharma, Neeraj Kumar, Yogendra S. Padwad, Brij Lal and Bikram Singh	161
Antioxidant and Anti-inflammatory Compounds in the Popular Landscape Plant <i>Berberis thunbergii</i> var. <i>atropurpurea</i> Chuan-Rui Zhang, Robert E. Schutzki and Muraleedharan G. Nair	165
Two New Amides from <i>Streptomyces michiganensis</i> Jinghua Xue, Liangxiong Xu, Zi-Hua Jiang and Xiaoyi Wei	169
Determination of Bioactive Compounds in the Juice of Pummelo (<i>Citrus grandis</i> Osbeck) Marina Russo, Ivana Bonaccorsi, Germana Torre, Antonella Cotroneo, Paola Dugo and Luigi Mondello	171
Antiplasmodial Activity of Compounds from the Surface Exudates of <i>Senecio roseiflorus</i> Leonidah Omosa Kerubo, Jacob Ogweni Midiwo, Solomon Derese, Moses K. Langat, Hosea M. Akala, Norman C. Waters, Martin Peter and Matthias Heydenreich	175

Anti-inflammatory, Antioxidant and Cytotoxicity Activities of Methanolic Extract and Prenylated Flavanones Isolated from Leaves of <i>Eysehhardtia platycarpa</i> Valeri Domínguez-Villegas, Vanessa Domínguez-Villegas, María Luisa García, Ana Calpena, Beatriz Clares-Naveros and María Luisa Garduño-Ramírez	177
Phenolic Glycosides from <i>Lindera obtusiloba</i> and their Anti-allergic Inflammatory Activities Hyun Gyu Choi, Hwa Dong Lee, Sang Hyun Kim, MinKyun Na, Jeong Ah Kim and Seung Ho Lee	181
Antiproliferative Effects of an Analog of Curcumin in Hep-2 cells: A Comparative Study with Curcumin Mohankumar Kumaravel, Pajaniradje Sankar, Periyasamy Latha, Chellakan S Benson and Rajagopalan Rukkumani	183
Antiproliferative Activity of epi-Cercosporin in Human Solid Tumor Cell Lines Ángel Trigos, César Espinoza, Maricela Martínez, Olivia Márquez, Leticia G. León, José M. Padrón, Manuel Norte and José J. Fernández	187
New Anthraquinone Derivatives from <i>Geosmithia lavendula</i> Lourin G. Malak, Daoud W. Bishay, Afaf M. Abdel-Baky, Ahmed M. Moharram, Stephen J. Cutler and Samir A. Ross	191
Pancreatic Lipase Inhibitory Activity of Cassiamin A, a Bianthraquinone from <i>Cassia siamea</i> Dilip Kumar, Aniket Karmase, Sneha Jagtap, Ruchi Shekhar and Kamlesh K. Bhutani	195
Antifeedant Activity of Spin-Labeled Derivatives of Deoxypodophyllotoxin against <i>Brontispa longissima</i> Larvae Gang Feng, Jing Zhang, Liu Yang, Ying-Qian Liu, Zhi-Wei Zhang, Xuan Tian, Qi-An Jin and Zheng-Qiang Peng	199
Determination of Organic Acids in <i>Salicornia herbacea</i> by Solid-phase Extraction Combined with Liquid Chromatography Dandan Han, Minglei Tian, Dong Wha Park and Kyung Ho Row	203
Hypoglycemic Effect of <i>Bumelia sartorum</i> Polyphenolic Rich Extracts Halliny S. Ruela, Katia C. C. Sabino, Ivana C. R. Leal, Ana M. Landeira-Fernandez, Michelle R. A. de Almeida, Talita S. M. Rocha and Ricardo M. Kuster	207
Protoanemonin Content Variation between <i>Clematis</i> spp.: Leaf, Stem and Root Fangming Jin, Christian Narkowicz and Glenn A Jacobson	211
Methanolic Extract of <i>Nigella sativa</i> Seed Inhibits SiHa Human Cervical Cancer Cell Proliferation Through Apoptosis Tarique N. Hasan, Gowhar Shafi, Naveed A Syed, Muhammad A Alfawaz, Mohammed A. Alsaif, Anjana Munshi, Kai Y. Lei and Ali A. Alshatwi	213
Glucosinolate Biosynthesis in Hairy Root Cultures of Broccoli (<i>Brassica oleracea</i> var. <i>italica</i>) Sun-Ju Kim, Woo Tae Park, Md. Romij Uddin, Yeon Bok Kim, Sang-Yong Nam, Kwang Hyun Jho and Sang Un Park	217
Characterization of Volatile Components of <i>Zingiber roseum</i> Essential Oil Using Capillary GC on Modified Cyclodextrins VPPalayam S Pragadheesh, Anju Yadav, Manju Singh and Chandan S Chanotiya	221
Essential Oil of Flowers of <i>Anaphalis contorta</i>, an Aromatic and Medicinal Plant from India Rajesh K. Joshi	225
Composition of Essential Oils from Seeds of <i>Abies koreana</i> Anna Wajs-Bonikowska, Karol Olejnik, Radosław Bonikowski and Piotr Banaszczyk	227
Chemical Composition of <i>Hypericum richeri</i> subsp. <i>grisebachii</i> Essential Oil from Croatia Igor Jerković, Maja Marasović, Zvonimir Marijanović, Kroat Hazler Pilepić, Željko Maleš and Mladen Miloš	231
Chemical Composition of the Essential Oils from Two Vietnamese <i>Asarum</i> Species: <i>A. glabrum</i> and <i>A. cordifolium</i> Tran Huy Thai, Ophélie Bazzali, Tran Minh Hoi, Nguyen Anh Tuan, Félix Tomi, Joseph Casanova and Ange Bighelli	235
Essential Oils from Five Species of Annonaceae from Vietnam Tran D. Thang, Do N. Dai, Tran M. Hoi and Isiaka A. Ogunwande	239
Essential Oils from the Leaves of Six Medicinal Plants of Nigeria Isiaka A. Ogunwande, Nudewheni O. Avoseh, Guido Flamini, Alimot-Sadiat O. Hassan, AbdulRazaq O. Ogunmoye, Akindele O. Ogunsanwo, Kamorudeen O. Yusuf, Atuonwu O. Kelechi, Zainab A. Tiamiyu and Godgift O. Tabowei	243
Comparative Study on <i>In Vitro</i> Activities of Citral, Limonene and Essential Oils from <i>Lippia citriodora</i> and <i>L. alba</i> on Yellow Fever Virus Luz Angela Gómez, Elena Stashenko and Raquel Elvira Ocazonez	249
TLC-Bioautographic Evaluation of <i>In Vitro</i> Anti-tyrosinase and Anti-cholinesterase Potentials of Sandalwood Oil Biswapriya B. Misra and Satyahari Dey	253
Composition, Mosquito Larvicidal, Biting Deterrent and Antifungal Activity of Essential Oils of Different Plant Parts of <i>Cupressus arizonica</i> var. <i>glabra</i> ('Carolina Sapphire') Abbas Ali, Nurhayat Tabanca, Betül Demirci, K. Husnu Can Baser, Jane Ellis, Sandra Gray, Brett R. Lackey, Christine Murphy, Ikhlas A. Khan and David E. Wedge	257
Composition of <i>Cassia fistula</i> Oil and its Antifungal Activity by Disrupting Ergosterol Biosynthesis Md. Irshad, Aijaz Ahmad, Md. Zafaryab, Farah Ahmad, Nikhat Manzoor, Man Singh and M. Moshahid A. Rizvi	261
Chemical Composition and Biological Activity of the Essential Oil of <i>Amomum biflorum</i> Chakkrapat Singtothong, Michel J. Gagnon and Jean Legault	265
Chemical Composition and Antibacterial Activity of Essential Oils from <i>Myrcia alagoensis</i> (Myrtaceae) Aline do N. Silva, Ana Paula T. Uetanabaro and Angélica M. Lucchese	269
Composition, <i>in-vitro</i> Anticancer, and Antimicrobial Activities of the Leaf Essential Oil of <i>Machilus mushaensis</i> from Taiwan Yu-Chang Su, and Chen-Lung Ho	273
Chemical Constituents and Cytotoxic Evaluation of Essential Oils from Leaves of <i>Porcelia macrocarpa</i> (Annonaceae) Erica Biolcati P. da Silva, Alisson L. Matsuo, Carlos R. Figueiredo, Mariana H. Chaves, Patricia Sartorelli and João Henrique G. Lago	277

Book Review

Medicinal Plants of China, Korea, and Japan: Bioresources for Tomorrow's Drugs and Cosmetics by Chritophe Wiart, PharmD, PhD Stephen Pyne	281
---	-----

Number 3

Metabolites from the Aerial Parts of the Sicilian Population of <i>Artemisia alba</i> Antonella Maggio, Sergio Rosselli, Celeste Laura Brancazio, Vivienne Spadaro, Francesco Maria Raimondo and Maurizio Bruno	283
New Bioactive Secondary Metabolites from Bornean Red Alga, <i>Laurencia similis</i> (Ceramiales) Takashi Kamada and Charles Santhanaraju Vairappan	287
A New Bioactive Sesquiterpenoid Quinone from the Mediterranean Sea Marine Sponge <i>Dysidea avara</i> Ashraf Nageeb El-Sayed Hamed, Wim Wätjen, Roland Schmitz, Yvonne Chovolou, RuAngelie Edrada-Ebel, Dina T. A. Youssef, Mohammed Salah Kamel and Peter Proksch	289
An Efficient Oxyfunctionalization of Quinopimaric Acid Derivatives with Ozone Oxana B. Kazakova, Elena V. Tretyakova, Irina E. Smirnova, Timur I. Nazyrov, Ol'ga S. Kukovinets, Genrikh A. Tolstikov and Kiryll Yu. Suponitskii	293
Plants from Northeast Mexico with Anti-HSV Activity David Silva-Mares, Ernesto Torres-López, Ana M Rivas-Estilla, Paula Cordero-Pérez, Noemí Waksman-Minsky and Verónica M Rivas-Galindo	297
The True Identity of the Triterpene Component of <i>Wyethia mollis</i>, a Lanosta-diene Containing a Tetrahydropyran E-ring Cole T. Smith, Bruce Noll, Thomas G. Waddell and Kyle S. Knight	299
Structures and Biological Activities of Typicosides A₁, A₂, B₁, C₁ and C₂, Triterpene Glycosides from the Sea Cucumber <i>Actinocucumis typica</i> Alexandra S. Silchenko, Anatoly I. Kalinovskiy, Sergey A. Avilov, Pelageya V. Andryaschenko, Pavel S. Dmitrenok, Ekaterina A. Martynyas, Vladimir I. Kalinin, P. Jayasandhya, Gigi C. Rajan and Krishna P. Padmakumar	301
Preparation and Application of Reversed Phase Chromatorotor™ for the Isolation of Natural Products by Centrifugal Preparative Chromatography Ilias Muhammad, Volodymyr Samoylenko, Francis Machumi, Mohamed Ahmed Zaki, Rabab Mohammed, Mona H. Hetta and Van Gillum	311
Chemical Constituents of the Leaves of <i>Dracaena thalioides</i> Akihito Yokosuka, Atsushi Sekiguchi and Yoshihiro Mimaki	315
7-Oxodioscin, a New Spirostan Steroid Glycoside from the Rhizomes of <i>Dioscorea nipponica</i> Zulfiqar Ali, Troy J. Smillie and Ikhlās A. Khan	319
Two Spirostan Steroid Glycoside Fatty Esters from <i>Dioscorea cayenensis</i> Zulfiqar Ali, Troy J. Smillie and Ikhlās A. Khan	323
GC-MS Investigation of Amaryllidaceae Alkaloids in <i>Galanthus xvalentinei</i> nothosubsp. <i>subplicatus</i> Buket Bozkurt Sarikaya, Strahil Berkov, Jaime Bastida, Gulen Irem Kaya, Mustafa Ali Onur and Nehir Unver Somer	327
Antifouling Indole Alkaloids from Two Marine Derived Fungi Fei He, Zhuang Han, Jiang Peng, Pei-Yuan Qian and Shu-Hua Qi	329
Andrographidine G, a New Flavone Glucoside from <i>Andrographis paniculata</i> Swarna D. Hapuarachchi, Zulfiqar Ali, Naohito Abe, Suresh T. Sugandhika, Senerath T. P. Sandun and Ikhlās A. Khan	333
Geranyl Flavonoids from <i>Robinia pseudoacacia</i> Lai-Bin Zhang, Jie-Li Lv and Hui Zhang	335
Antibacterial Activity Against <i>Ralstonia solanacearum</i> of Phenolic Constituents Isolated from Dragon's Blood Hui Wang, Ying Luo, Haofu Dai and Wenli Mei	337
Interaction of Dihydromyricetin and α-Amylase Lei Chen, Chao Wang, Qingyi Wei, Zhengxiang Ning and Erdong Yuan	339
Effect of the Isoflavone Genistein on Tumor Size, Metastasis Potential and Melanization in a B16 Mouse Model of Murine Melanoma Corina Danciu, Florina Borcan, Florina Bojin, Istvan Zupko and Cristina Dehelean	343
Quantitative Analysis of Total Flavonoids and Total Phenolic Acids in Thirty <i>Hypericum</i> Taxa Kroata Hazler Pilepić, Željko Maleš and Maja Crkvenčić	347
New Diarylheptanoids and a Hydroxylated Otelione from <i>Ottelia alismoides</i> Thomas R. Hoye, Seif-Eldin N. Ayyad, Hollie J. Beckord and Susan G. Brown	351
Evaluation of the <i>in vivo</i> Anti-inflammatory and Analgesic and <i>in vitro</i> Anti-cancer Activities of Curcumin and its Derivatives James N. Jacob, Dinesh K. Badyal, Suman Bala and Masoud Toloue	359
Chemical Constituents from <i>Dendropanax dentiger</i> Yi-Chun Lai and Shuei-Sheng Lee	363
Six New Phenolic Glycosides and a New Ceramide from the Flowers of <i>Wedelia biflora</i> and Their Cytotoxicity Against Some Cancer Cell Lines Nguyen T. H. Thu, Le T. Ha, Vo T. Nga, Pham N. K. Tuyen, Ton T. Quang, Forde-Riddick Danielle, Pratt Lawrence M. and Nguyen K. P. Phung	367
Antimicrobials from the Marine Algal Endophyte <i>Penicillium</i> sp. Andrew J. Flewelling, John A. Johnson and Christopher A. Gray	373
Antioxidant and Cytoprotective Effects of an Ethanol Extract of <i>Acalypha wilkesiana</i> var. <i>macafeana</i> from Malaysia Wardah M. Din, Jessica Chu, Garry Clarke, Khoo T. Jin, Tracey D. Bradshaw, Jeff R. Fry and Christophe Wiart	375
A New Antimicrobial and Anticancer Peptide Producing by the Marine Deep Sediment Strain "<i>Paenibacillus profundus</i>" sp. nov. SI 79 Nataliya I. Kalinovskaya, Lyudmila A. Romanenko, Anatoly I. Kalinovskiy, Pavel S. Dmitrenok and Sergey A. Dyshlovoy	381

Potential Applications for <i>Annona squamosa</i> Leaf Extract in the Treatment and Prevention of Foodborne Bacterial Disease Achara Dholvitayakhun, Nathanon Trachoo, Uthai Sakee and T.P. Tim Cushnie	385
Genoprotective Effect of the Chinese Herbal Decoction Xiao Jian Zhong Tang Yim-Tong Szeto, Ngok-Fung Cheng, Sok-Cheon Pak and Wouter Kalle	389
Effect of Nine Plant Volatiles in the Field on the Sex Pheromones of <i>Leguminivora glycinivorella</i> Daihua Hu, Juntao Feng, Zhihui Wang, Hua Wu and Xing Zhang	393
Induction of Defensive Response in <i>Eucalyptus globulus</i> Plants and its Persistence in Vegetative Propagation Christian Troncoso, Claudia Perez, Victor Hernandez, Manuel Sanchez-Olate, Darcy Rios, Aurelio San Martin and José Becerra	397
Chemical Composition of the Essential Oil of <i>Baccharoides lilacina</i> from India Rajesh K. Joshi	401
Chemical Composition and Anti-<i>Trypanosoma cruzi</i> Activity of Essential Oils Obtained from Leaves of <i>Xylopia frutescens</i> and <i>X. laevigata</i> (Annonaceae) Thanany Brasil da Silva, Leociley Rocha Alencar Menezes, Marília Fernanda Chaves Sampaio, Cássio Santana Meira, Elisalva Teixeira Guimarães, Milena Botelho Pereira Soares, Ana Paula do Nascimento Prata, Paulo Cesar de Lima Nogueira and Emmanouel Vilaça Costa	403
Chemical Composition and Biological Activities of the Essential Oil from <i>Artemisia herba-alba</i> Growing Wild in Tunisia Ismail Amri, Laura De Martino, Aurelio Marandino, Hamrouni Lamia, Hanana Mohsen, Elia Scandolera, Vincenzo De Feo and Emilia Mancini	407
<i>Aloe ferox</i> Seed: A Potential Source of Oil for Cosmetic and Pharmaceutical Use Rachael Dangarembizi, Eliton Chivandi and Kennedy Erlwanger	411
<u>Review/Account</u>	
Therapeutic Effects of Glycyrrhizic Acid Lee Jia Ming and Adeline Chia Yoke Yin	415
Natural Products of Mineral Origin Massimiliano Laudato, Luigi Pescitelli and Raffaele Capasso	419

Addition/Correction

Chemical Composition and Biological Activities of Soldiers of the Brazilian Termite Species, <i>Nasutitermes macrocephalus</i> (Isoptera: Natutitermitinae) Márcia N. S. de la Cruz, Helvécio M. S. Júnior, Denilson F. Oliveira, Leticia V. Costa-Lotufo, Antonio G. Ferreira, Daniela S. Alviano and Claudia M. Rezende	424
---	-----

Number 4

Anti-melanogenesis Constituents from the Seaweed <i>Dictyota coriacea</i> Ryeo Kyeong Ko, Min-Chul Kang, Sang Suk Kim, Tae Heon Oh, Gi-Ok Kim, Chang-Gu Hyun, Jin Won Hyun and Nam Ho Lee	427
Methyl Carnosate, an Antibacterial Diterpene Isolated from <i>Salvia officinalis</i> Leaves Elisa Climati, Fabio Mastrogiovanni, Maria Valeri, Laura Salvini, Claudia Bonechi, Nilufar Zokirzhonovna Mamadalieva, Dilfuza Egamberdieva, Anna Rita Taddei and Antonio Tiezzi	429
Cytotoxicity of Meroterpenoids from <i>Sargassum siliquastrum</i> against Human Cancer Cells Jung Im Lee, Myoung K. Kwak, Hee Y. Park and Youngwan Seo	431
Isolation of Methyl 27-caffeoyloxyoleanolate – A New Oleanane Triterpenoid from the Roots of <i>Hibiscus vitifolius</i> Duraisamy Ramasamy and Ariamuthu Saraswathy	433
Synthesis and Cytotoxic Activity of New Betulin and Betulinic Acid Esters with Conjugated Linoleic Acid (CLA) Barbara Tubek, Paweł Mituła, Natalia Niezgoda, Katarzyna Kempnińska, Joanna Wietrzyk and Czesław Wawrzęńczyk	435
Analysis of Pyrrolizidine Alkaloids and Evaluation of Some Biological Activities of Algerian <i>Senecio delphinifolius</i> (Asteraceae) Soukaina Tidjani, Philippe N. Okusa, Amar Zellagui, Laetitia Moreno Y Banuls, Caroline Stévigny, Pierre Duez and Salah Rhouati	439
Berbanine: a New Isoquinoline-isoquinolone Alkaloid from <i>Berberis vulgaris</i> (Berberidaceae) Anna Hošťálková, Zdeněk Novák, Milan Pour, Anna Jirošová, Lubomír Opletal, Jiří Kuneš and Lucie Cahliková	441
Dicentrine Production in Callus and Cell Suspension Cultures of <i>Stephania venosa</i> Tharita Kitisripanya, Jukrapun Komaikul, Nirachara Tawinkan, Chuennapha Atsawinkowit and Waraporn Putalun	443
New Flavan and Alkyl α,β-Lactones from the Stem Bark of <i>Horsfieldia superba</i> Nabil Ali Al-Mekhlafi, Khozirah Shaari, Faridah Abas, Ethyl Jeyaseela Jeyaraj, Johnson Stanslas, Shaik Ibrahim Khalivulla and Nordin H. Lajis	447
New Flavonol Triglycosides from the Leaves of Soybean Cultivars Yoshinori Murai, Ryoji Takahashi, Felipe Rojas Rodas, Junichi Kitajima and Tsukasa Iwashina	453
Melitidin: A Flavanone Glycoside from <i>Citrus grandis</i> 'Tomentosa' Wei Zou, Yonggang Wang, Haibin Liu, Yulong Luo, Si Chen and Weiwei Su	457
Two New Chalones from the Flowers of <i>Clerodendrum inerme</i> Shaik Khadar Shahabuddin, Rachakunta Munikishore, Golakoti Trimurtulu, Duvvuru Gunasekar, Alexandre Deville and Bernard Bodo	459
A Novel Phenolic Compound from <i>Phyllanthus emblica</i> Gaimei She, Ruiyang Cheng, Lei Sha, Yixia Xu, Renbin Shi, Lanzhen Zhang and Yajian Guo	461

Anti-austeric Activity of Phenolic Constituents of Seeds of <i>Arctium lappa</i> Yasuhiro Tezuka, Keiichi Yamamoto, Suresh Awale, Feng Li, Satoshi Yomoda and Shigetoshi Kadota	463
Bioactive Lignans from the Leaves and Stems of <i>Schisandra wilsoniana</i> Guang-Yu Yang, Rui-Rui Wang, Zhong-Hua Gao, Yin-Ke Li, Liu-Meng Yang, Xiao-Nian Li, Shan-Zhai Shang, Yong-Tang Zheng, Wei-Lie Xiao and Han-Dong Sun	467
Antioxidative / Acetylcholinesterase Inhibitory Activity of Some Asteraceae Plants Ivana Generalić Mekinić, Franko Burčul, Ivica Blažević, Danijela Skroza, Daniela Kerum and Višnja Katalinić	471
Antioxidant and Antimicrobial Activities, and Phenolic Compounds of Selected <i>Inula</i> species from Turkey Alper Gökbulut, Onural Özhan, Basri Satılmış, Kadir Batçoğlu, Selami Günel and Engin Şarer	475
Two New Dihydrostilbenoid Glycosides Isolated from the Leaves of <i>Litsea coreana</i> and their Anti-inflammatory Activity Wenjian Tang, Weili Lu, Xiaoqing Cao, Yilong Zhang, Hong Zhang, Xiongwen Lv and Jun Li	479
Inhibitory Activity of Benzophenones from <i>Anemarrhena asphodeloides</i> on Pancreatic Lipase Yang Hee Jo, Seon Beom Kim, Jong Hoon Ahn, Qing Liu, Bang Yeon Hwang and Mi Kyeong Lee	481
Identification and Quantification of Furanocoumarins in Stem Bark and Wood of Eight Algerian Varieties of <i>Ficus carica</i> by RP-HPLC-DAD and RP-HPLC-DAD-MS Samia Rouaiguia-Bouakkaz, Habiba Amira-Guebailia, Céline Rivière, Jean-Claude Delaunay, Pierre Waffo-Tégouo and Jean-Michel Mérillon	485
UPLC-Q-TOF/MS Coupled with Multivariate Statistical Analysis as a Powerful Technique for Rapidly Exploring Potential Chemical Markers to Differentiate Between <i>Radix Paeoniae Alba</i> and <i>Radix Paeoniae Rubra</i> Nian-cui Luo, Wen Ding, Jing Wu, Da-wei Qian, Zhen-hao Li, Ye-fei Qian, Jian-ming Guo and Jin-ao Duan	487
Antimicrobial Activity of Crude Methanolic Extract from <i>Phyllanthus niruri</i> Darah Ibrahim, Lim Sheh Hong and Nanthianantham Kuppan	493
Cellulose Contents of Some Abundant Indian Seaweed Species Arup K. Siddhanta, Sanjay Kumar, Gaurav K. Mehta, Mahesh U. Chhatbar, Mihir D. Oza, Naresh D. Sanandya, Dharmesh R. Chejara, Chirag B. Godiya and Stalin Kondaveeti	497
Anti-inflammatory Potential of Silk Sericin Pornanong Aramwit, Pasarapa Towiwat and Teerapol Srichana	501
Composition of Essential Oil from Aerial and Underground Parts of <i>Geum rivale</i> and <i>G. urbanum</i> Growing in Poland Aleksandra Owczarek, Jan Gudej and Agnieszka Kice	505
Volatile Composition of Six Horsetails: Prospects and Perspectives Françoise Fons, Didier Froissard, Jean-Marie Bessière, Alain Fruchier, Bruno Buatois and Sylvie Rapior	509
Chemical Compositions of the Rhizome, Leaf and Stem Oils from Malaysian <i>Hornstedtia leonurus</i> Nor Akmalazura Jani, Hasnah Mohd. Sirat, NorAzah Mohamad Ali and Azrina Aziz	513
Effect on Emotional Behavior and Stress by Inhalation of the Essential oil from <i>Chamaecyparis obtusa</i> Hikaru Kasuya, Erika Hata, Tadaaki Satou, Masaki Yoshikawa, Shinichiro Hayashi, Yoshinori Masuo and Kazuo Koike	515
Chemical Composition and Antibacterial Activity of Rhizome Oils from Five <i>Hedychium</i> Species Ratchuporn Suksathan, Siriwoot Sookkhee, Somboon Anuntalabhochai and Sunee Chansakaow	519
Chemical Composition and Antimicrobial Activity of Three Essential Oils from <i>Curcuma wenyujin</i> Jingjing Zhu, Agnieszka D. Lower-Nedza, Meng Hong, Song Jie, Zhimin Wang, Dong Yingmao, Christine Tschiggerl, Franz Bucar and Adelheid H. Brantner	523
Essential Oil Composition and Antimicrobial Activity of Aerial Parts and Ripe Fruits of <i>Echinophora spinosa</i> (Apiaceae) from Italy Daniele Fraternali, Salvatore Genovese and Donata Ricci	527
Composition and <i>in vitro</i> Anticancer Activities of the Leaf Essential Oil of <i>Neolitsea variabilissima</i> from Taiwan Yu-Chang Su, Kuan-Ping Hsu, Eugene I-Chen Wang and Chen-Lung Ho	531

Review/Account

Natural Products from Marine Algae of the Genus <i>Osmundaria</i> (Rhodophyceae, Ceramiales) Kelvin Osako and Valéria Laneuville Teixeira	533
Phenols, Alkaloids and Terpenes from Medicinal Plants with Antihypertensive and Vasorelaxant Activities. A Review of Natural Products as Leads to Potential Therapeutic Agents Francesco Maione, Carla Cicala, Giulia Musciacco, Vincenzo De Feo, Anibal G. Amat, Armando Ialenti and Nicola Mascolo	539
Diosmin – Isolation Techniques, Determination in Plant Material and Pharmaceutical Formulations, and Clinical Use Anna Bogucka – Kocka, Michał Woźniak, Marcin Feldo, Janusz Kocki and Katarzyna Szewczyk	545

Number 5

Gerald Blunden Award (2012)

Cytotoxic Agents of the Crinane Series of Amaryllidaceae Alkaloids Jerald J. Nair, Jaime Bastida, Francesc Viladomat and Johannes van Staden	553
--	-----

Original Paper

Chemosystematics of the Thai Liverwort <i>Cheilolejeunea</i> (Marchantiophyta, Lejeuneaceae) Phiangphak Sukkharak and Yoshinori Asakawa	565
---	-----

Cytotoxic Properties of <i>Marrubium globosum</i> ssp. <i>libanoticum</i> and its Bioactive Components Mariangela Marrelli, Filomena Conforti, Daniela Rigano, Carmen Formisano, Maurizio Bruno, Felice Senatore and Francesco Menichini	567
Cytotoxic Scalarane Sesterterpenoids from a Marine Sponge <i>Hippospongia</i> sp. Yuh-Ming Fuh, Mei-Chin Lu, Chia-Hung Lee and Jui-Hsin Su	571
Ursane-Type Saponins from <i>Zygophyllum cornutum</i> Soumeiya Bencharif-Betina, Tomofumi Miyamoto, Chiaki Tanaka, Zahia Kabouche, Anne-Claire Mitaine-Offer and Marie-Aleth Lacaille-Dubois	573
Vasorelaxant Alkaloids from <i>Spirospermum penduliflorum</i> (Menispermaceae), a Plant Used to Treat Hypertension in Malagasy Traditional Medicine Guy E. Raoelison, Mamy H. Rafamantanana, René Razafindrazaka, Adolphe Randrianntsoa, Suzanne Urverg-Ratsimamanga, Nicole Morel and Joëlle Quetin-Leclercq	575
PPARα Signaling is Activated by Cocoa in Mouse Liver Marco Fidaleo and Claudia Sartori	579
Chemical Analysis of Flowers of <i>Bombax ceiba</i> from Nepal Khem Raj Joshi, Hari Prasad Devkota and Shoji Yahara	583
Chemical Investigation of <i>Caragana arborescens</i> Shoots Daniil N. Olennikov, Larisa M. Tankhaeva and Vyacheslav V. Partilkhaev	585
A New Metabolite from the Endophytic Fungus <i>Penicillium citrinum</i> Xinlan Li, Liang Zhang, Yanhui Liu, Zhiyong Guo, Zhangshuang Deng, Jianfeng Chen, XuanTu and Kun Zou	587
Antioxidant Activity of the Isoflavonoids from the Roots of <i>Maackia amurensis</i> Nadezda I. Kulesh, Sergey A. Fedoreyev, Marina V. Veselova, Natalia P. Mischenko, Vladimir A. Denisenko, Pavel S. Dmitrenok, Yakov F. Zverev and Svetlana V. Zamyatina	589
The Effect of Pyridinecarbothioamides on Isoflavonoid Production in <i>Genista tinctoria</i> Cultures in Vitro Lenka Tůmová, Věra Klimešová and Anna Vildová	593
A New Homoisoflavanone from the Rhizomes of <i>Polygonatum cyrtonema</i> Li-She Gan, Jin-Jie Chen, Man-Fei Shi, and Chang-Xin Zhou	597
Two 2-Phenylbenzofuran Derivatives from <i>Morus atropurpurea</i> Wen-Jing Wang, Dong-Ling Wu, Sen-Tai Liao, Chun-Lin Fan, Guo-Qiang Li, Xian-Tao Zhang, Ying Wang, Xiao-Qi Zhang and Wen-Cai Ye	599
10'-Deoxy-10' α-hydroxyasochlorin, a New Cell Migration Inhibitor and Other Metabolites from <i>Acremonium</i> sp., a Fungal Endophyte in <i>Ephedra trifurca</i> W. M. Anoja P. Wanigesekara, E. M. Kithsiri Wijeratne, A. Elizabeth Arnold and A. A. Leslie Gunatilaka	601
Polyphenols in the Aqueous Extracts of Garden Thyme (<i>Thymus vulgaris</i>) Chemotypes Cultivated in Hungary Blanka Szilvássy, Gábor Rak, Szilvia Sárosi, Ildikó Novák, Zsuzsanna Pluhár and László Abrankó	605
Resveratrol Production from Hairy Root Cultures of <i>Scutellaria baicalensis</i> Sang-Won Lee, Young Seon Kim, Md. Romij Uddin, Do Yeon Kwon, Yeon Bok Kim, Mi Young Lee, Sun-Ju Kim and Sang Un Park	609
Anti-periodontal Pathogen and Anti-inflammatory Activities of Oxyresveratrol Waranyoo Phoolcharoen, Sireerat Soampon, Boonchoo Sritularak, Kittisak Likhitwitayawuid, Jintakorn Kuvatanasuchati and Prasit Pavasant	613
The Triple Botanical Origin of Russian Propolis from the Perm Region, Its Phenolic Content and Antimicrobial Activity Milena Popova, Boryana Trusheva, Rail Khismatullin, Natalja Gavrilova, Galina Legotkina, Jaroslav Lyapunov and Vassya Bankova	617
Comparing Different Solvent Extracts of <i>Rhus semialata</i> var. <i>roxburghiana</i> Stem against Ferrous Ion-Induced Lipid Peroxidation in Mice Liver Mitochondria Pei-Chin Lin, Wei-Fung Bi, Che-Hsuan Lin, Fei-Peng Lee and Ling-Ling Yang	621
Antioxidant Action of Solid Preparation of Xingnaojing in SHRSP Yang Liu, Naomi Yasui, Aya Kishimoto, Jian-ning Sun and Katsumi Ikeda	627
Antioxidant and Antityrosinase Activity of <i>Cissus quadrangularis</i> Extract Ikuko Suzu, Hiroki Goto, Nami Hiwatashi, Shinichiro Hattori, Kanjana Rotjanapan, Wilairat Leeanansaksiri and Seiji Okada	629
Eight New Alkyne and Alkene Derivatives from Four <i>Saussurea</i> Species Collected in China Yoshinori Saito, Yuko Iwamoto, Yasuko Okamoto, Takayuki Kawahara, Xun Gong, Chiaki Kuroda and Motoo Tori	631
The Apoptotic Activity of one VLC Fraction of the Sponge <i>Petrosia tuberosa</i> on Human Cervical Cells and the Subsequent Isolation of a Bioactive Polyacetylene Avin Ramanjooloo, Girish Beedessee, Deepak Arya, Rob WM. vanSoest, Thierry Cresteil and Daniel E.P. Marie	635
Chemical Characterization, Mineral Content and Radical Scavenging Activity of <i>Sideritis scardica</i> and <i>S. raeseri</i> from R. Macedonia and R. Albania Marija Karapandzova, Bujar Qazimi, Gjoshe Stefkov, Katerina Bačeva, Trajče Stafilov, Tatjana Kadifkova Panovska and Svetlana Kulevanova	639
Phytochemical and Micromorphological Traits of <i>Geranium dalmaticum</i> and <i>G. macrorrhizum</i> (Geraniaceae) Dario Kremer, Dubravka Vitali Čepo, Valerija Dunkić, Ivna Dragojević Müller, Ivan Kosalec, Nada Bezić and Edith Stabenheiner	645
GC-MS Fingerprints and Other Physico-chemical Characteristics of Rare Unifloral <i>Prunus cerasus</i> L. Honey Piotr Marek Kuś, Igor Jerković, Carlo Ignazio Giovanni Tuberose, Zvonimir Marijanović and Mladenka Šarolić	651
Volatile Fraction Composition and Total Phenolic and Flavonoid Contents of <i>Elionurus hensii</i>—Antioxidant Activities of Essential Oils and Solvent Extracts Yin Yang, Marie-Cécile De Cian, Samuel Nsikabaka, Pierre Tomi, Thomas Silou, Jean Costa and Julien Paolini	655
Leaf Essential Oils of Six Vietnamese Species of <i>Fissistigma</i> (Annonaceae) Martina Höferl, Do Ngoc Dai, Tran Dinh Thang, Leopold Jirovetz and Erich Schmidt	663

Studies on the Antimicrobial and Antioxidant Activity and Chemical Composition of the Essential Oils of <i>Kitaibelia vitifolia</i> Pavle Mašković, Marija Radojković, Mihailo Ristić and Slavica Solujić	667
Angiotensin Converting Enzyme Inhibition Activity of Fennel and Coriander Oils from India Sushil Kumar Chaudhary, Niladri Maity, Neelesh Kumar Nema, Santanu Bhadra, Bishnu Pada Saha and Pulok Kumar Mukherjee	671
Effect of Coriander Oil (<i>Coriandrum sativum</i>) on Planktonic and Biofilm Cells of <i>Acinetobacter baumannii</i> Andreia F. Duarte, Susana Ferreira, Rosário Oliveira and Fernanda C. Domingues	673
Essential Oil from <i>Caesalpinia peltophoroides</i> Flowers – Chemical Composition and <i>in vitro</i> Cytotoxic Evaluation Bianca A. de Carvalho, Olivia S. Domingos, Murilo Massoni, Marcelo H. dos Santos, Marisa Ionta, João Henrique G. Lago, Carlos R. Figueiredo, Alisson L. Matsuo and Marisi G. Soares	679
Antimicrobial, Antioxidant, and Cytotoxic Activities of the Essential Oil of <i>Tarchonanthus camphoratus</i> Nasser A. Awadh Ali, Mohamed A. Al-Fatimi, Rebecca A. Crouch, Annika Denkert, William N. Setzer and Ludger Wessjohann	683

Number 6

Australasian Natural Product Studies (Guest Editor: Sylvia Urban)

Epimanol and a New 2,6-Dideoxy-hexopyran-3-ulose Derivative from <i>Celmisia viscosa</i> Catherine E. Sansom, Lesley Larsen, Alison C. Evans and Nigel B. Perry	689
7-Bromo-1-ethyl-β-carboline, an Alkaloid from the New Zealand Marine Bryozoan <i>Pterocella vesiculosa</i> Michèle R. Prinsep and Méline Dumté	693
Alkaloids from the Roots and Leaves of <i>Stichoneuron halabalensis</i> and their Acetylcholinesterase Inhibitory Activities Rosdayati Alino Ramli, Wilford Lie and Stephen G. Pyne	695
Establishment of a Phenotypic-based Sand Dollar <i>Fellaster zelandiae</i> Embryo Development Assay and its Application in Defining the Structure-Activity Relationship of Discorhabdin Alkaloids Tanja Grkovic and Brent R. Copp	699
Sharing of Pyrazine Semiochemicals Between Genera of Sexually Deceptive Orchids Bjorn Bohman, Ryan D. Phillips, Gavin Flematti, Rod Peakall and Russell A. Barrow	701
Constituents of the Indonesian Epiphytic Medicinal Plant <i>Drynaria rigidula</i> Ari S. Nugraha, Rachada Haritakun and Paul A. Keller	703
Variability of the Polyphenolic Content and Antioxidant Capacity of Methanolic Extracts of Pomegranate Peel Antony Kam, Kong M Li, Valentina Razmovski-Naumovski, Srinivas Nammi, Kelvin Chan and George Q Li	707
Combination Effects of Curcumin and Aqueous Extract of <i>Lignosus rhinocerotis</i> Mycelium on Neurite Outgrowth Stimulation Activity in PC-12 Cells Priscilla A. John, Kah - Hui Wong, Murali Naidu, Vikineswary Sabaratnam and Pamela David	711
HPLC-NMR Chemical Profiling and Dereplication Studies of the Marine Brown Alga, <i>Cystophora torulosa</i> Sylvia Urban and Michael Timmers	715
Two Polyketides from a Co-culture of Two Marine-derived Fungal Strains Miriam H. Kossuga, Antonio G. Ferreira, Lara D. Sette and Roberto G. S. Berlinck	721
Cyclic Peroxides from a Two-Sponge Association of <i>Plakortis communis</i>-<i>Agelas mauritiana</i> Pinus Jumaryatno, Lynette K. Lambert, John N. A. Hooper, Joanne T. Blanchfield and Mary J. Garson	725
Relative Configuration of the Marine Natural Product Elatenyne using NMR Spectroscopic and Chemical Derivatization Methodologies Robert Brkljača and Sylvia Urban	729
Antimicrobial, Antimalarial and Cytotoxicity Activities of Constituents of a Bhutanese Variety of <i>Ajania nubigena</i> Phurpa Wangchuk, Paul A. Keller, Stephen G. Pyne, John Korth, Samten, Malai Taweechotipatr, Roonglawan Rattanajak and Sumalee Kamchonwongpaisan	733
Dihydrotagetone, an Unusual Fruity Ketone, is Found in Enantiopure and Enantioenriched Forms in Additional Australian Native Taxa of <i>Phebalium</i> (Rutaceae: Boronieae) Nicholas J. Sadgrove, Ian R. H. Telford, Ben W. Greatrex, Ashley Dowell and Graham L. Jones	737
Antimicrobial Activity of Essential Oils and Solvent Extracts from <i>Zieria</i> species (Rutaceae) Nicholas J. Sadgrove and Graham L. Jones	741
Characterization and Bioactivity of Essential Oils from <i>Geijera parviflora</i> (Rutaceae): A Native Bush Medicine from Australia Nicholas J. Sadgrove and Graham L. Jones	747

Synthesis and Insecticidal Activities of Novel Derivatives of 1β,4α,6α,9α-Tetrahydroxy-2β,12-epoxymethano-β-dihydroagarofuran Jiwen Zhang, Zhan Hu, Shengkun Li, Shuding Yang and Wenjun Wu	753
Novel α-Glucosidase Activator from <i>Pulicaria undulata</i> Nasir Rasool, Muhammad Abid Rashid, Saleha Suleman Khan, Zulfiqar Ali, Muhammd Zubair, Viqar Uddin Ahmad, Shamsun Nahar Khan, M. Iqbal Choudhary and Rasool Bakhsh Tareen	757
Phytochemical, Antimicrobial and Antiplasmodial Investigations of <i>Terminalia brownii</i> Francis Machumi, Jacob O. Midiwo, Melissa R. Jacob, Shabana I. Khan, Babu L. Tekwani, Jin Zhang, Larry A. Walker and Ilias Muhammad	761

A New Sulfated Triterpene Saponin from <i>Gypsophila trichotoma</i> Iliana Krasteva, Maya Yotova, Kristina Jenett-Siems, Petranka Zdraveva and Stefan Nikolov	765
Bioactive Triterpene Saponins from the Leaves of <i>Schefflera elegantissima</i> Amira S. Wanas, Mostafa A. Fouad, Mohamed S. Kamel, Katsuyoshi Matsunami and Hideaki Otsuka	767
Synthesis and Mass Spectral Fragmentation Patterns of Brassinolide Early Biosynthetic Precursors Labeled at C-26 Vladimir A. Khrupach, Danuše Tarkowská, Vladimir N. Zhabinskii, Olga V. Gulyakevich, Yurii V. Ermolovich, Pavel Drašar and Miroslav Strnad	771
Severibuxine, Isolated from <i>Severinia buxifolia</i>, Induces Apoptosis in HL-60 Leukemia Cells Chihiro Ito, Tomiyasu Murata, Midori Kato, Natsu Suzuki, Tian-Shung Wu, Norio Kaneda, Hiroshi Furukawa and Masataka Itoigawa	775
Two New Morphinandienone Alkaloids from <i>Croton micradenus</i> Armando L. Payo-Hill, Daysi Sandoval-López, Hermán T. Vélez-Castro, Iraida Spengler-Salabarría and Luca Rastrelli	779
Isolation and Cholinesterase Inhibitory Activity of <i>Narcissus</i> Extracts and Amaryllidaceae Alkaloid Lucie Čahlíková, Miroslav Ločárek, Nina Benešová, Radim Kučera, Jakub Chlebek, Zdeněk Novák and Lubomír Opletal	781
Antioxidant Activity and Mechanism of Tetrahydroamentoflavone <i>in vitro</i> Xican Li, Weijuan Han, Wenqiong Mai and Li Wang	787
Blue Mood for Superfood Kari Taulavuori, Riitta Julkunen-Tiitto, Valtteri Hyöky and Erja Taulavuori	791
Acetylcholinesterase Inhibitors from <i>Croton sylvaticus</i> Ethyl Acetate Leaf Extract and their Mutagenic Effects Mutalib A. Aderogba, Ashwell R. Ndhlala and Johannes Van Staden	795
Coumarin Precursor from <i>Micromelum integerrimum</i> Leaves Wong Phakhodee, Preetiya Pongparn, Angkana Saovapakhiran and Surat Laphookhieo	799
Antifilarial Activity of Constituents of <i>Calophyllum inophyllum</i> and their Derivatives Jankiprasad, Gunaganti Naresh, Jyoti Gupta, Shailja M. Bhattacharya, Siron M. Rajendran, Shailendra K. Awasthi and Tadigoppula Narendar	803
Two New Lignans from the Wood of <i>Cunninghamia konishii</i> Chi-I Chang, Yen-Cheng Li, Ching-Chuan Kuo, Che-Yi Chao, Hsun-Shuo Chang, Jyh-Horng Wu, Sheng-Yang Wang and Yueh-Hsiung Kuo	805
Biological Studies of Turmeric Oil, Part 1: Selective <i>in vitro</i> Anticancer Activity of Turmeric Oil (TO) and TO-Paclitaxel Combination James N. Jacob and Masoud Toloue	807
Biological Studies of Turmeric Oil, Part 2: Isolation and Characterization of Turmeric Oil Components; Cytotoxic Activity Against Pancreatic Cancer Cells Wudan Yan, Wayne D. Bowen, Russell Hopson, Abraham E. Mathew and James N. Jacob	811
Ushikulide C, a New Immunosuppressant from <i>Streptomyces</i> sp. IUK-102 Kosaku Takahashi, Eri Fukushima, Jun Kawabata, Hideyuki Matsuura and Kazuhiko Kurosawa	815
Influence of the Bioclimatic Area on the Polyphenolic Composition, and Antioxidant and Bacteriostatic Activities of <i>Rosmarinus officinalis</i> María J. Jordán, Vanesa Lax, María C. Rota, Susana Lorán and José A. Sotomayor	817
Antibacterial and Antioxidant Activities of Various Medicinal Plants Used in Oriental Medicine Junho Seo, Jiyeon Kim, Geon Go, Jung-Suk Sung and Kwang-Geun Lee	823
Aconitamide, a Novel Alkaloid from the Roots of <i>Aconitum carmichaeli</i> Tzong-Huei Lee, Yu-Chang Chen, Ching-Kuo Lee, Hsun-Shuo Chang, Ching-Chuan Kuo, Che-Yi Chao, Jing-Jer Lin, Lee-Chiang Lo and Yueh-Hsiung Kuo	827
The Chemical Composition and Antibacterial Activities of the Essential Oils from Three <i>Aframomum</i> Species from Cameroon, and Their Potential as Sources of (<i>E</i>)-(<i>R</i>)-Nerolidol Sylvie Kwanga Nguikwie, Maximilienne A. Nyegue, Florentine Ndoye-Foe Belinga, Rosalie A. Ngono Ngane, Bernard Romestand, Achraf Kouzayha, Hervé Casabianca, Paul H. Amvam Zollo and Chantal Menut	829
Chemical Composition and Antimicrobial Activity of <i>Daucus aureus</i> Essential Oils from Algeria Nawel Meliani, Mohammed El Amine Dib, Nassim Djabou, Jean Costa, Hocine Allali, Boufeldja Tabti and Alain Muselli	835
Chemical Composition of the Essential Oils of Three Species of Apiaceae Growing Wild in Sicily: <i>Bonannia graeca</i>, <i>Eryngium maritimum</i> and <i>Opopanax chironium</i> Antonella Maggio, Maurizio Bruno, Carmen Formisano, Daniela Rigano and Felice Senatore	841

Number 7

New Aspects in Natural Product Synthesis: Methodology and Strategy (Guest Editors: Hisahiro Hagiwara and Toshio Suzuki)

Regioselective Hydroxylation and Glucosylation of α- and β-Pinenes with Cultured Cells of <i>Eucalyptus perriniana</i> Kei Shimoda, Naoji Kubota, Manabu Hamada, Ken Suwada, Hatsuyuki Hamada and Hiroki Hamada	847
Synthesis of Sterically-Crowded Olefins, <i>gem</i>-Dihaloalkenes, Butatrienes, and Thioketenes through the Reaction of Substituted Bornane-2-thiones or Bornane-2-selones with Conventional Carbenes or Metal Carbenoids Kazuaki Shimada, Jun Sasaki, Anna Kishi, Shigenobu Aoyagi and Yuji Takikawa	851
Modification of Valencene by Bio- and Chemical Transformation Yoshinori Asakawa, Toshihiro Hashimoto, Yoshiaki Noma and Mai Furusawa	859

Enantioselective Synthesis of Natural Trinorsesquiterpene Tetralones by Chemo-enzymatic Approaches Stefano Serra	863
Structural Factors in the Odor of α-Santalol Derivatives Toshio Hasegawa, Hiroaki Izumi and Hideo Yamada	869
Second Generation Synthesis of the Neo-Clerodane Diterpenoid Methyl Barbascoate Hisahiro Hagiwara, Naomi Honma, Kimihiko Kinugawa, Shota Sato, Takashi Hoshi and Toshio Suzuki	873
Synthesis of a Key Intermediate, 10-Acetyl-7-(<i>t</i>-butyldiphenylsilyloxymethyl)-4-methylbicyclo[5.3.0]dec-4-en-1-ol, in the Synthesis of Pseudolaric Acid A Katsuyuki Nakashima, Naoki Kikuchi, Tasuku Takehara, Takayuki Shiozawa, Shigeru Takaoka and Motoo Tori	877
Use of RCM Reactions for Construction of Eight-Membered Carbocycles and Introduction of a Hydroxy Group at the Junction Between Five- and Eight-Membered Carbocycles Takahiro Morimitsu, Reiko Mizutani, Katsuyuki Nakashima, Yoshinori Saito and Motoo Tori	883
Synthetic Approach Toward α-Aminomethyl-γ-butyrolactones from β-Lactam Synthons Elaborated by SmI₂-mediated Reductive Coupling Reactions Masaki Takahashi, Takahiro Sudo, Yusuke Murata, Tetsuya Sengoku and Hidemi Yoda	889
Formal Synthesis of (+)-Madindoline A, a Potent IL-6 Inhibitor, Utilizing Enzymatic Discrimination of Quaternary Carbon Ken-ichi Shimizu, Mina Tomita, Ken-ichi Fuhshuku, Takeshi Sugai and Mitsuru Shoji	897
Glucosylation of Taxifolin with Cultured Plant Cells Kei Shimoda, Naoji Kubota, Manabu Hamada, Masahiro Sugamoto, Kohji Ishihara, Hatsuyuki Hamada and Hiroki Hamada	903
Regioselective Hydroxylation and Glucosylation of Flavanones with Cultured Plant Cells of <i>Eucalyptus perriniana</i> Ryusuke Hosoda, Yoshiyuki Horio, Kei Shimoda, Manabu Hamada, Hatsuyuki Hamada and Hiroki Hamada	905
Synthesis of Resveratrol Glycosides by Cultured Plant Cells Kei Shimoda, Manabu Hamada, Hatsuyuki Hamada, Mai Takemoto and Hiroki Hamada	907
Synthesis of 3-Farnesyl Salicylic Acid, a Novel Antimicrobial from <i>Piper multiplinervium</i> George A. Kraus, Divya Chaudhary, Sean Riley, Feng Liu, Allison Schlapkohl, Megan Weems and Gregory J. Phillips	911
Total Synthesis of Bisbibenzyl Dibenzofuran Asterelin A via Intramolecular Oxidative Coupling Kosho Makino, Kenichi Harada, Miwa Kubo, Hideaki Hioki and Yoshiyasu Fukuyama	915
Cyclopentanoids from Cyclopentadiene: Synthesis of (–)-Methyl jasmonate and (+)-12-Oxophytodienoic acid Junzo Nokami, Kazuhiko Fujii, Yusuke Mizutani, Rikiya Omatsu, Kiyoshi Watanabe, Hiroshi Yasuda and Tsutomu Inokuchi	919
Synthesis of Both Enantiomers of 12-Methyl-13-tridecanolide and 14-Methyl-15-pentadecanolide (Muscolide) Yoshihiro Noda, Natsuki Mamiya and Hitoshi Kashin	925
A New Variant of Fused Cyclic Ether Synthesis Based on Ireland-Claisen Rearrangement and RCM Daisuke Doman, Kenshu Fujiwara, Natsumi Kawamura, Ryo Katoono, Hidetoshi Kawai and Takanori Suzuki	929
 <u>Review/Account</u>	
Carvone as a Versatile Chiral Building Block for Total Syntheses of Heterocyclic Sesquiterpenoids Hisahiro Hagiwara	935
Synthesis of a Hydrindenone in Rings C and D of YW3699 Reiko Mizutani, Takahiro Morimitsu, Katsuyuki Nakashima and Motoo Tori	949
Collective Total Synthesis of PPAPs: Total Synthesis of Clusianone via Intramolecular Cyclopropanation Masahiro Uwamori and Masahisa Nakada	955
Synthetic Approaches to Tetracyclic Pyrrole Imidazole Marine Alkaloids Takuya Imaoka, Makoto Iwata, Takafumi Akimoto and Kazuo Nagasawa	961
Lynbouilloside and Related Macrolides from Marine Cyanobacteria Abdelatif ElMarrouni, Amandine Kolleth, Raphael Lebeuf, Julian Gebauer, Sébastien Prevost, Montserrat Heras, Stellios Arseniyadis and Janine Cossy	965
Enantioselective Total Synthesis of Otteliones A and B, Novel and Powerful Antitumor Agents from the Freshwater Plant <i>Ottelia alismoides</i> Tadashi Katoh	973
A New Approach to the Synthesis of Chiral Blocks for Cyclopentanoids Airat M. Gimazetdinov, Nadezhda A. Ivanova and Mansur S. Miftakhov	981
Stereocontrolled Total Synthesis of Tetrodotoxin from <i>myo</i>-Inositol and D-Glucose by Three Routes: Aspects for Constructing Complex Multi-Functionalized Cyclitols with Branched-Chain Structures Ken-ichi Sato, Shoji Akai and Juji Yoshimura	987
Development of a Second Generation Palladium-Catalyzed Cycloalkenylation and its Application to Bioactive Natural Product Synthesis Masahiro Toyota	999
Stereoselective Aminopalladation and Oxypalladation and Their Application to the Synthesis of Natural Products Hidefumi Makabe	1005
Evolution of the Total Syntheses of Batzellasides, the First Marine Piperidine Iminosugar Tetsuya Sengoku, Jolanta Wierzejska, Masaki Takahashi and Hidemi Yoda	1011
Recent Applications and Developments of Organic Azides in Total Synthesis of Natural Products Hiroki Tanimoto and Kiyomi Kakiuchi	1021

Number 8

Natural Clovanes from the Gorgonian Coral <i>Rumphella antipathies</i> Hsu-Ming Chung, Wei-Hsien Wang, Tsong-Long Hwang, Yang-Chang Wu and Ping-Jyun Sung	1037
Microbial Biotransformation of 16α,17-Epoxy-ent-kaurane-19-oic acid by <i>Beauveria sulfurescens</i> ATCC 7159-F Ricardo A. Furtado, G. M. Kamal B. Gunaherath, Jairo K. Bastos and A. A. Leslie Gunatilaka	1041
Establishment of <i>In vitro</i> Adventitious Root Cultures and Analysis of Andrographolide in <i>Andrographis paniculata</i> Shiv Narayan Sharma, Zenu Jha and Rakesh Kumar Sinha	1045
New Cycloartane-type Triterpenes from <i>Marcetia latifolia</i> (Melastomataceae) and <i>in silico</i> Studies on <i>Candida parapsilosis</i> Protease Tonny C. C. Leite, Franco H. A. Leite, Ivo J. C. Vieira, Raimundo Braz Filho and Alessandro Branco	1049
Triterpene Glycosides from the Sea Cucumber <i>Eupentacta fraudatrix</i>. Structure and Biological Action of Cucumariosides I₁, I₃, I₄, Three New Minor Disulfated Pentaosides Alexandra S. Silchenko, Anatoly I. Kalinovskiy, Sergey A. Avilov, Pelageya V. Andryjaschenko, Pavel S. Dmitrenok, Ekaterina A. Martyyas and Vladimir I. Kalinin	1053
Improved Extraction and Complete Mass Spectral Characterization of Steroidal Alkaloids from <i>Veratrum californicum</i> Christopher M. Chandler, Jeffrey W. Habig, Ashley A. Fisher, Katherine V. Ambrose, Susana T. Jiménez and Owen M. McDougal	1059
TLC-Image Analysis of Non-Chromophoric Tuberostemonine Alkaloid Derivatives in <i>Stemona</i> Species Sumet Kongkiatpaiboon, Vichien Keeratinjakal and Wandee Gritsanapan	1065
Two New Compounds from Gorgonian-associated Fungus <i>Aspergillus</i> sp. Xin-Ya Xu, Xiao-Yong Zhang, Fei He, Jiang Peng, Xu-Hua Nong and Shu-Hua Qi	1069
New Metabolites from the Algal Associated Marine-derived Fungus <i>Aspergillus carneus</i> Olesya I. Zhuravleva, Shamil Sh. Afyatullof, Ekaterina A. Yurchenko, Vladimir A. Denisenko, Natalya N. Kirichuk and Pavel S. Dmitrenok	1071
Chiroptical Studies of Flavanone Marcelo A. Muñoz, María A. Bucio and Pedro Joseph-Nathan	1075
Flavonoids with Anti-HSV Activity from the Root Bark of <i>Artocarpus lakoocha</i> Boonchoo Sritularak, Kullasap Tantrakarnsakul, Vimolmas Lipipun and Kittisak Likhitwitayawuid	1079
Orphan Flavonoids and Dihydrochalcones from <i>Primula</i> Exudates Tshering Doma Bhutia, Karin M. Valant-Vetschera and Lothar Brecker	1081
First Identification of α-Glucosidase Inhibitors from Okra (<i>Abelmoschus esculentus</i>) Seeds Wannisa Thanakosai and Preecha Phuwapraisirisan	1085
<i>In vitro</i> Anti-proliferative Effect of Naturally Occurring Oxyprenylated Chalcones Serena Fiorito, Francesco Epifano, Celine Bruyère, Robert Kiss and Salvatore Genovese	1089
Kaempferol 3,7,4'-glycosides from the Flowers of <i>Clematis</i> Cultivars Keisuke Sakaguchi, Junichi Kitajima and Tsukasa Iwashina	1093
7-O-Methylpelargonidin Glycosides from the Pale Red Flowers of <i>Catharanthus roseus</i> Fumi Tatsuzawa	1095
Pyranocoumarin and Triterpene from <i>Millettia richardiana</i> Manitriniaina Rajemiarimiraho, Jean Théophile Banzouzi, Stéphane Richard Rakotonandrasana, Pierre Chalard, Françoise Benoit-Vical, Léa Herilala Rasoaiva, Amélie Raharisololalao and Roger Randrianja	1099
A Concise and Efficient Total Synthesis of α-Mangostin and β-Mangostin from <i>Garcinia mangostana</i> Dandan Xu, Ying Nie, Xizhou Liang, Ling Ji, Songyuan Hu, Qidong You, Fan Wang, Hongchun Ye and Jinxin Wang	1101
Variation in the Contents of Neochlorogenic Acid, Chlorogenic Acid and Three Quercetin Glycosides in Leaves and Fruits of Rowan (<i>Sorbus</i>) Species and Varieties from Collections in Lithuania Kristina Gaivelyte, Valdas Jakstas, Almantas Razukas and Valdimaras Janulis	1105
A New Cytotoxic Phenolic Derivative from the Roots of <i>Antidesma acidum</i> Sutin Kaennakam, Jirapast Sichaem, Pongpun Siripong and Santi Tip-pyang	1111
A New Nervogenic Acid Glycoside with Pro-coagulant Activity from <i>Liparis nervosa</i> Qin Song, Qingyao Shou, Xiaojun Gou, Fengzhen Chen, Jing Leng and Weifeng Yang	1115
Biologically Active Secondary Metabolites from <i>Asphodelus microcarpus</i> Mohammed M. Ghoneim, Guoyi Ma, Atef A. El-Hela, Abd-Elsalam I. Mohammad, Saeid Kottob, Sayed El-Ghaly, Stephen J. Cutler and Samir A. Ross	1117
Dibenzylbutane Lignans from the Stems of <i>Schisandra bicolor</i> Yinning Chen, Na Li, Yuehui Zhu, Cuilan Zhang, Xiaofei Jiang, Jianxiang Yang, Zhifang Xu, Samuel X. Qiu and Riming Huang	1121
Absolute Configuration of Falcarinol (9Z-heptadeca-1,9-diene-4,6-diyn-3-ol) from <i>Pastinaca sativa</i> Mireia Corell, Emile Sheehy, Paul Evans, Nigel Brunton and Juan Valverde	1123
A New Macrolide from a Marine-derived Fungus <i>Aspergillus</i> sp. Jie Bao, Xin-Ya Xu, Xiao-Yong Zhang and Shu-Hua Qi	1127
HPLC-MS and GC-MS Analyses Combined with Orthogonal Partial Least Squares to Identify Cytotoxic Constituents from Turmeric (<i>Curcuma longa</i> L.) Jianlan Jiang, Huan Zhang, Zidan Li, Xiaohang Zhang, Xin Su, Yan Li, Bin Qiao and Yingjin Yuan	1129
Determination of Antiplasmodial Activity and Binding Affinity of Selected Natural Products towards <i>Pf</i>TrxR and <i>Pf</i>GR Ranjith Muniguntti, Katja Becker, Reto Brun and Angela I. Calderón	1135
<i>In vitro</i> Anti-inflammatory and Antioxidant Potential of Si-Miao-San, its Modifications and Pure Compounds Agneszka D. Lower-Nedza, Carmen Kuess, Haiyu Zhao, Baolin Bian and Adelheid H. Brantner	1137

Smooth Muscle Relaxation Activity of an Aqueous Extract of Dried Immature Fruit of <i>Poncirus trifoliata</i> (PF-W) on an Isolated Strip of Rat Ileum	
Kyu-Sang Kim, Won-Sik Shim, Ike Campomayor dela Peña, Eun-Kyung Seo, Woo-Young Kim, Hyo-Eon Jin, Dae-Duk Kim, Suk-Jae Chung, Jae-Hoon Cheong and Chang-Koo Shim	1143
Invertebrate Water Extracts as Biocompatible Reducing Agents for the Green Synthesis of Gold and Silver Nanoparticles	
Lina Han, Yeong Shik Kim, Seonho Cho, and Youmie Park	1149
Elemental Analysis of <i>Ginkgo biloba</i> Leaf Samples Collected during One Vegetation Period	
Szilvia Czige, Erzsébet Háznagy-Radnai, Klára Pintye-Hódi, Jaroslav Tóth, Daniela Tekeľová and Imre Máthé	1153
Wild <i>Thymra capitata</i> from Western Rif (Morocco): Essential Oil Composition, Chemical Homogeneity and Yield Variability	
Khadija Bakhy, Ouafae Benhabib, Chaouki Al Faiz, Ange Bighelli, Joseph Casanova and Felix Tomi	1155
Essential Oils of <i>Chiliadenus lopadusanus</i> (Asteraceae)	
Pietro Zito, Maurizio Sajeve, Elena Scirica, Maurizio Bruno, Sergio Rosselli, Antonella Maggio and Felice Senatore	1159
Chemical Composition of the Essential Oil of <i>Lepidagathis fasciculata</i> from Bondla Forest of Goa, India	
Rajesh K. Joshi	1163
Leaf Oil from <i>Vepris madagascariensis</i> (Rutaceae), Source of (<i>E</i>)-Anethole	
Delphin J. R. Rabehaja, Harilala Ihandriharison, Panja A. R. Ramanoelina, Suzanne Ratsimamanga-Urverg, Ange Bighelli, Joseph Casanova and Félix Tomi	1165
Cardiovascular Effects of the Essential Oil of <i>Croton zehntneri</i> Leaves in DOCA-salt Hypertensive, Conscious Rats	
Rodrigo José Bezerra de Siqueira, Gloria Pinto Duarte, Pedro Jorge Caldas Magalhães and Saad Lahlou	1167
Chemical Composition and Antioxidant Activity of the Essential Oil and Fatty Acids of the Flowers of <i>Rhanterium adpressum</i>	
Chahrazed Hamia, Nadhir Gourine, Hadjer Boussoussa, Mokhtar Saidi, Emile M. Gaydou and Mohamed Yousfi	1171
Chemical Composition and Biological Activity of <i>Conyza bonariensis</i> Essential Oil Collected in Mérida, Venezuela	
Liliana Araujo, Laila M. Moujir, Janne Rojas, Luis Rojas, Juan Carmona and María Rondón	1175
Chemical Compositions, Phytotoxicity, and Biological Activities of <i>Acorus calamus</i> Essential Oils from Nepal	
Prabodh Satyal, Prajwal Paudel, Ambika Poudel, Noura S. Dosoky, Debra M. Moriarity, Bernhard Vogler and William N. Setzer	1179
 <u>Review/Account</u>	
Review on Research of Suppression Male Fertility and Male Contraceptive Drug Development by Natural Products	
Vijay Kumar Bajaj and Radhey S Gupta	1183

Number 9

Alternate Biosynthesis of Valerenadiene and Related Sesquiterpenes	
Shashikumar K. Paknikar, Shahuraj H. Kadam, April L. Ehrlich and Robert B. Bates	1195
A Facile Synthesis of (±)-Heliannuol-D	
Tao Zhang, Liang-Zhu Huang, You-Qiang Li, Yimg-Meng Xu and Zhen-Ting Du	1197
A New Bioactive Diterpene Glycoside from <i>Molinia retusa</i> from the Madagascar Dry Forest	
Alexander L. Eaton, Liva Harinantenaina, Peggy J. Brodie, Maria B. Cassera, Jessica D. Bowman, Martin W. Callmänder, Richard Randrianaivo, Roland Rakotondrajaona, Etienne Rakotobe, Vincent E. Rasamison and David G. I. Kingston	1201
Nitric Oxide and Tumor Necrosis factor-α Inhibitory Substances from the Rhizomes of <i>Kaempferia marginata</i>	
Kanidta Kaewkroek, Chatchai Wattanapiromsakul, Palangpon Kongsaree and Supinya Tewtrakul	1205
Biscembranoids from the Marine Sponge <i>Petrosia nigricans</i>	
Nguyen Xuan Nhiem, Ngo Van Quang, Chau Van Minh, Dan Thi Thuy Hang, Hoang Le Tuan Anh, Bui Huu Tai, Pham Hai Yen, Nguyen Thi Hoai, Do Cong Thung and Phan Van Kiem	1209
Isolation of Cycloecalenol from <i>Boophone disticha</i> and Evaluation of its Cytotoxicity	
Emmanuel Adekanmi Adewusi, Paul Steenkamp, Gerda Fouche and Vanessa Steenkamp	1213
Chemical Constituents from an Endophytic Fungus <i>Chaetomium globosum</i> Z1	
Chun-Yan Zhang, Xiao Ji, Xuan Gui and Bao-Kang Huang	1217
Determination of C-23 Configuration in (20<i>R</i>)-23-Hydroxycholestane Side Chain of Steroid Compounds by ^1H and ^{13}C NMR Spectroscopy	
Alla A. Kicha, Anatoly I. Kalinovsky, Alexander S. Antonov, Oleg S. Radchenko, Natalia V. Ivanchina, Timofey V. Malyarenko, Alexander M. Savchenko and Valentin A. Stonik	1219
Oxasetin from <i>Lophiostoma</i> sp. of the Baltic Sea: Identification, <i>in silico</i> Binding Mode Prediction and Antibacterial Evaluation against Fish Pathogenic Bacteria	
Muftah Ali M. Shushni, Faizul Azam and Ulrike Lindequist	1223
Chemical Constituents from the Fruit Body of <i>Chlorophyllum molybdites</i>	
Zushang Su, Ping Wang, Wei Yuan, and Shiyong Li	1227
Pulchranins B and C, New Acyclic Guanidine Alkaloids from the Far-Eastern Marine Sponge <i>Monanchora pulchra</i>	
Tatyana N. Makarieva, Ekaterina K. Ogurtsova, Yuliya V. Korolkova, Yaroslav A. Andreev, Irina V. Mosharova, Ksenya M. Tabakmakher, Alla G. Guzii, Vladimir A. Denisenko, Pavel S. Dmitrenok, Hyi-Seung Lee, Eugene V. Grishin and Valentin A. Stonik	1229
Cloning and Characterization of a cDNA Encoding Calcium/Calmodulin-dependent Glutamate Decarboxylase from <i>Scutellaria baicalensis</i>	
Yeon Bok Kim, Md Romij Uddin, Do Yeon Kwon, Min-Ki Lee, Sun-Ju Kim, Chanhui Lee and Sang Un Park	1233

Biflavonoids, Main Constituents from <i>Garcinia bakeriana</i> Leaves Ahmed Al-Shagdari, Adonis Bello Alarcón, Osmany Cuesta-Rubio, Anna Lisa Piccinelli and Luca Rastrelli	1237
Analysis of Flavonoids and Iridoids in <i>Vitex negundo</i> by HPLC-PDA and Method Validation Somendu K. Roy, Khemraj Bairwa, Jagdeep Grover, Amit Srivastava and Sanjay M. Jachak	1241
Chemical Constituents of the Leaves of <i>Triumfetta semitriloba</i> Alejandra Barraza-Morales, Deisy Medrano-Nahuat, Sergio R. Peraza-Sánchez	1245
Phytochemical Evaluation of <i>Lythrum salicaria</i> Extracts and Their Effects on Guinea-pig Ileum Tímea Bencsik, Loránd Barthó, Viktor Sándor, Nóra Papp, Rita Benkó, Attila Felinger, Ferenc Kilár and Györgyi Horváth	1247
New Flavonol Glycosides from the Leaves of <i>Triantha japonica</i> and <i>Tofieldia nuda</i> Tsukasa Iwashina, Minoru N. Tamura, Yoshinori Murai and Junichi Kitajima	1251
Cytotoxic Activity of Dihydrochalcones Isolated from <i>Corema album</i> Leaves against HT-29 Colon Cancer Cells Antonio J. León-González, Miguel López-Lázaro, José L. Espartero and Carmen Martín-Cordero	1255
Immunomodulatory Activities of α-Mangostin on Peripheral Blood Mononuclear Cells Pimolkan Kasemwattanaoj, Primchanien Moongkarndi, Kovit Pattanapanyasat, Supachoke Mangmool, Ekkarat Rodpai, Jutima Samer, Julaporn Konlata and Kasama Sukapirom	1257
Antiplasmodial Quinones from the Rhizomes of <i>Kniphofia foliosa</i> Martha Induli, Meron Gebru, Negera Abdissa, Hosea Akala, Ingrid Wekesa, Robert Byamukama, Matthias Heydenreich, Sylvia Murunga, Ermias Dagne and Abiy Yenesew	1261
Biphenyl Derivatives from <i>Garcinia schomburgkiana</i> and the Cytotoxicity of the Isolated Compounds Chihiro Ito, Takuya Matsui, Eri Noda, Nijisri Ruangrunsi and Masataka Itoigawa	1265
Anticarcinogenic Effect and Carcinogenic Potential of the Dietary Phenolic Acid: <i>o</i>-Coumaric Acid Alaattin Sen, Pelin Atmaca, Gulsum Terzioğlu and Sevkî Arslan	1269
Bioproduction and Optimization of Rosmarinic Acid Production in <i>Solenostemon scutellarioides</i> through Media Manipulation and Conservation of High Yielding Clone via Encapsulation Ranabir Sahu, Saikat Dewanjee and Moumita Gangopadhyay	1275
<i>In vitro</i> Anti-diabetic Activity of <i>Sclerocarya birrea</i> and <i>Ziziphus mucronata</i> Nuno M.H. Da Costa Mousinho, Jacob J. van Tonder and Vanessa Steenkamp	1279
Secondary Metabolites from the Fungus <i>Emericella nidulans</i> Amer H. Tarawneh, Francisco León, Mohamed M. Radwan, Luiz H. Rosa and Stephen J. Cutler	1285
A New Glucuronolactone Glycoside Phoenixoside B from the Seeds of <i>Phoenix dactylifera</i> Sumbul Azmat, Rehana Ifzal, Faryal Vali Mohammad, Viqar Uddin Ahmad and Aqib Zahoor	1289
Cancer-Suppressive Potential of Extracts of Endemic Plant <i>Helichrysum zivojinii</i>: Effects on Cell Migration, Invasion and Angiogenesis Ivana Z. Matić, Ivana Aljančić, Vlatka Vajs, Milka Jadranin, Nevenka Gligorićević, Slobodan Milosavljević and Zorica D. Jurančić	1291
Analysis of Volatile Components, Fatty Acids, and Phytosterols of <i>Abies koreana</i> growing in Poland Anna Wajs-Bonikowska, Karol Olejnik, Radosław Bonikowski and Piotr Banaszczyk	1297
Cytotoxic Effects of Air Freshener Biocides in Lung Epithelial Cells Jung-Taek Kwon, Mimi Lee, Gun-Baek Seo, Hyun-Mi Kim, Ilseob Shim, Doo-Hee Lee, Taksoo Kim, Jung Kwan Seo, Pilje Kim and Kyunghee Choi	1301
GC/GC-MS Analysis, Isolation and Identification of Bioactive Essential Oil Components from the Bhutanese Medicinal Plant, <i>Pleurospermum amabile</i> Phurpa Wangchuk, Paul A. Keller, Stephen G. Pyne, Malai Taweechotipatr and Sumalee Kamchonwongpaisan	1305
Antibacterial Activity of the Essential Oil of <i>Heracleum sibiricum</i> Dragoljub L. Miladinović, Budimir S. Ilić, Tatjana M. Mihajilov-Krsteć, Dejan M. Nikolić, Olga G. Cvetković, Marija S. Marković and Ljiljana C. Miladinović	1309
Assessment of the Chemical Composition and <i>in vitro</i> Antimicrobial Potential of Extracts of the Liverwort <i>Scapania aspera</i> Danka R. Bukvicki, Amit K. Tyagi, Davide G. Gottardi, Milan M. Veljic, Snezana M. Jankovic, Maria E. Guerzoni and Petar D. Marin	1313
Essential Oils of <i>Alpinia rafflesiana</i> and Their Antimicrobial Activities Shariha Jusoh, Hasnah Mohd. Sirat and Farediah Ahmad	1317
Chemical Composition and Synergistic Antioxidant Activities of Essential Oils from <i>Atractylodes macrocephala</i> and <i>Astragalus membranaceus</i> Jinkui Li, Feng Li, Yan Xu, Wenjian Yang, Lili Qu, Qian Xiang, Cong Liu and Dapeng Li	1321
Chemical Analysis and Antioxidant Activity of the Essential Oils of Three Piperaceae Species Growing in the Central Region of Cuba Elisa Jorge Rodríguez, Yanelis Saucedo-Hernández, Yvan Vander Heyden, Ernesto F. Simó-Alfonso, Guillermo Ramis-Ramos, Maria Jesús Lerma-García, Urbano Monteagudo, Luis Bravo, Mildred Medina, Yuriam de Armas and José Manuel Herrero-Martínez	1325
The Composition, Anti-mildew and Anti-wood-decay Fungal Activities of the Leaf and Fruit Oils of <i>Juniperus formosana</i> from Taiwan Yu-Chang Su, Kuan-Ping Hsu, Eugene I-Chen Wang and Chen-Lung Ho	1329

Meeting/Report

Meeting Report: First National Meeting on Aloe, April 20-21, 2013, Isernia, Italy New Perspectives in Aloe Research: from Basic Science to Clinical Application Raffaele Capasso, Massimiliano Laudato and Francesca Borrelli	1333
---	------

Review/Account**Alkaloids of the South African Amaryllidaceae: a Review**

Jerald J. Nair, Jaime Bastida, Carles Codina, Francesc Viladomat and Johannes van Staden 1335

Number 10**Guaiool - A Naturally Occurring Insecticidal Sesquiterpene**

Tao Liu, Chun-Juan Wang, Hui-Qin Xie and Qing Mu 1353

3-Oxoabolene and 1-Oxocurcuphenol, Aromatic Bisabolanes from the Sponge *Myrmekioderma* sp.

Afsaneh Yegdaneh, Sumaitt Putchakarn, Supreeya Yuenyongsawad, Alireza Ghannadi and Anuchit Plubrukarn 1355

Acanthoic Acid Inhibits Melanogenesis through Tyrosinase Down-regulation and Melanogenic Gene Expression in B16 Melanoma Cells

Weon-Jong Yoon, Young-Min Ham, Hun Seok Yoon, Wook-Jae Lee, Nam Ho Lee and Chang-Gu Hyun 1359

Cembranoids from the Cultured Soft Coral *Sinularia gibberosa*

Hsiu-Fen Lin, Huey-Jen Su, Nai-Lun Lee and Jui-Hsin Su 1363

Major Constituents of *Boswellia carteri* Resin Exhibit Cyclooxygenase Enzyme Inhibition and Antiproliferative Activity

Sami I. Ali, Chuan-Rui Zhang, Amal A. Mohamed, Farouk K. EL-Baz, Ahmad K. Hegazy, Maimona A. Kord and Muraledharan G. Nair 1365

Cucurbitane-type Triterpenes from *Citrullus lanatus* (Watermelon) Seeds

Takashi Kikuchi, Rina Okada, Yu Harada, Kenji Ikushima, Takahiro Yamakawa, Takeshi Yamada and Reiko Tanaka 1367

A New Taraxerol Derivative from the Roots of *Microcos tomentosa*

Sutin Kaennakam, Jirapast Sichae, Suttira Khumkratok, Pongpun Siripong and Santi Tip-pyang 1371

Two New Triterpenoids from *Gelsemium elegans* and *Aglaia odorata*

Bing Liu, Lin Yang, You-Kai Xu, Shang-Gao Liao, Huai-Rong Luo and Zhi Na 1373

Inhibition of Tumor Cells Multidrug Resistance by Cucumarioside A₂-2, Frondoside A and their Complexes with Cholesterol

Ekaterina S. Menchinskaya, Dmitry L. Aminin, Sergey A. Avilov, Aleksandra S. Silchenko, Pelageya V. Andryjashchenko, Vladimir I. Kalinin and Valentin A. Stonik 1377

Anticholinesterase and Antioxidant Activities of Fucoxanthin Purified from the Microalga *Phaeodactylum tricornutum*

Arthitaya Kawee-ai, Ampin Kuntiya and Sang Moo Kim 1381

Impact of Ploidy Change on Secondary Metabolites and Photochemical Efficiency in *Solanum bulbocastanum*

Immacolata Caruso, Fabrizio Dal Piaz, Nicola Malafronte, Nunziatina De Tommasi, Riccardo Aversano, Cristian Wulff Zottele, Maria-Teresa Scarano and Domenico Carputo 1387

Alkaloids from an Endophytic *Streptomyces* sp. YIM66017

Hao Zhou, Yabin Yang, Jucheng Zhang, Tianfeng Peng, Lixing Zhao, Lihua Xu and Zhongtao Ding 1393

Novel Decaturin Alkaloids from the Marine-Derived Fungus *Penicillium oxalicum*

Pei-le Wang, Dan-yi Li, Lei-rui Xie, Xin Wu, Hui-ming Hua and Zhan-lin Li 1397

Monanchomycalin C, a New Pentacyclic Guanidine Alkaloid from the Far-Eastern Marine Sponge *Monanchora pulchra*

Ksenya M. Tabakmakher, Vladimir A. Denisenko, Alla G. Guzii, Pavel S. Dmitrenok, Sergey A. Dyshlovoy, Hyi-Seung Lee and Tatyana N. Makarieva 1399

Antiinflammatory and Antioxidant Flavonoids from *Helichrysum kraussii* and *H. odoratissimum* Flowers

Percival B. Legoale, Mahlori J. Mashimbye and Teunis van Ree 1403

Antiproliferative Effect of Flavonoids from the Halophyte *Vitex rotundifolia* on Human Cancer Cells

You Ah Kim, Hojun Kim and Youngwan Seo 1405

Identification of a Xanthine Oxidase-inhibitory Component from *Sophora flavescens* using NMR-based Metabolomics

Ryuichiro Suzuki, Yuka Hasuike, Moeka Hirabayashi, Tatsuo Fukuda, Yoshihito Okada and Yoshiaki Shirataki 1409

Flavone C-Glycosides from *Lychnis senno* and their Antioxidative Activity

Hari Prasad Devkota, Kumiko Fukusako, Koji Ishiguro and Shoji Yahara 1413

Anti-oxidative and DNA Protecting Effects of Flavonoids-rich *Scutellaria lateriflora*

Madhukar Lohani, Manuj Ahuja, Manal A Buabeid, Dean Schwartz, Dennis Shannon, Vishnu Suppiramaniam, Barbara Kempainen and Muralikrishnan Dhanasekaran 1415

Polyphenolic Compounds from Callus Cultures of *Iris pseudacorus*

Darya V. Tarbeeva, Sergey A. Fedoreyev, Marina V. Veselova, Anatoliy I. Kalinovskiy, Ludmila D. Seletskaya, Tamara I. Mazurok and Victor P. Bulgakov 1419

Pterocarpanes from the Root Bark of *Aeschynomene fascicularis*

Edgar Caamal-Fuentes, Rosa Moo-Puc, Luis W. Torres-Tapia and Sergio R. Peraza-Sanchez 1421

Cytotoxic Constituents of *Pachyrhizus tuberosus* from Peruvian Amazon

Olga Leuner, Jaroslav Havlik, Milos Budesinsky, Vladimir Vrkoslav, Jessica Chu, Tracey D. Bradshaw, Jana Hummelova, Petra Miksatkova, Oldrich Lapcik, Irena Valterova and Ladislav Kokoska 1423

(+)-Rumphiin and Polyalthurea, New Compounds from the Stems of *Polyalthia rumphi*

Tian-Shan Wang, You-Ping Luo, Jing Wang, Meng-Xiong He, Ming-Guo Zhong, Ying Li and Xiao-Ping Song 1427

Citriquinones A and B, New Benzoquinones from *Penicillium citrinum*

P. K. Vinitha Ranji, S. Chandrani Wijeyaratne, K. Hector Jayawardana and G. M. Kamal B. Gunaherath 1431

Potent Microbial and Tyrosinase Inhibitors from Stem Bark of *Bauhinia rufescens* (Fabaceae)

Aminu Muhammad and Hasnah Mohd Sirat 1435

Inhibitory Effects of Stilbene Derivatives from <i>Parthenocissus tricuspidata</i> on Adipocyte Differentiation and Pancreatic Lipase Sang Hoon Lee, Qing Liu, Bang Yeon Hwang and Mi Kyeong Lee	1439
HPLC-PDA Analysis and Anti-inflammatory Effects of Mori Cortex Radicis Chang-Seob Seo, Hye-Sun Lim, Soo-Jin Jeong, HyeKyung Ha and Hyeun-Kyoo Shin	1443
Development of HPLC Fingerprint for Quality Assessment of Bulbus Lili Kunming Qin, Hao Cai, Lijuan Zheng, Miao Zhang, Xinghai Zhang, Jie Gu and Baochang Cai	1447
Antifungal and Antioxidant Pyrrole Derivative from <i>Piper pedicellatum</i> Chandan Tamuly, Partha P. Dutta, Manobjyoti Bordoloi and Jayanta Bora	1451
Quorum Sensing: A Non-conventional Target for Antibiotic Discovery Varsha Naik and Girish Mahajan	1455
An Alkenylresorcinol Derivative from <i>Hakea sericea</i> Fruits and their Antimicrobial Activity Ángelo Luís, Carla Cruz, Ana Paula Duarte and Fernanda Domingues	1459
Glucosinolates in Two Endemic Plants of the <i>Aurinia</i> Genus and their Chemotaxonomic Significance Ivica Blažević, Gina Rosalinda De Nicola, Sabine Montaut and Patrick Rollin	1463
Proteinaceous Protease Inhibitor from <i>Lawsonia inermis</i>: Purification, Characterization and Antibacterial Activity Arvind Dabhade, Priti Patel and Ulhas Patil	1467
Essential Oils Composition of <i>Croton</i> Species from the Amazon Nathalie A. Turiel, Alcy F. Ribeiro, Elisangela Elena N. Carvalho, Vanessa D. Domingos, Flávia Cristina A. Lucas, Léa Maria M. Carreira, Eloisa Helena A. Andrade and José Guilherme S. Maia	1471
Comparative Analysis of the Essential Oil Composition of <i>Murraya paniculata</i> and <i>M. exotica</i> Hai-Ning Lv, Xiao-Yu Guo, Peng-Fei Tu and Yong Jiang	1473
Chemical Composition of the Essential Oil of <i>Erechtites valerianaeifolia</i> from Mérida, Venezuela Johanna Hernández, Ismer Bracho, Luis B. Rojas-Fermin, Alfredo Usubillaga and Juan Carmona	1477
Effect of Different Drying Methods on the Essential Oils of Mint (<i>Mentha haplocalyx</i>) Yachun Shu, Yajun Chen, Kunming Qin, Hao Cai, Li Wu, Huan Li, Lijuan Xu and Baochang Cai	1479
<i>Juglans regia</i> and <i>J. nigra</i>, Two Trees Important in Traditional Medicine: A Comparison of Leaf Essential Oil Compositions and Biological Activities Prajwal Paudel, Prabodh Satyal, Noura S. Dosoky, Samjhana Maharjan and William N. Setzer	1481

Review/Account

Natural Compounds against Flaviviral Infections Md Abubakr, Subhash C Mandal and Sugato Banerjee	1487
Microwave-Assisted Techniques (MATs); a Quick Way to Extract a Fragrance: A Review Antonios K. Kokolakis and Spyridon K. Golfopoulos	1493

Number 11

Three New Iridoid Glucoside Salts from <i>Hedyotis tenelliflora</i> Growing in Vietnam Luu H. Van Long, Vo Thi Nga, Nguyen Phuc Dam, Mai Anh Hung, Tu Duc Dung, Ton That Quang and Nguyen K. Phi Phung	1507
<i>In vitro</i> Anti-inflammatory Activity of <i>Ligustrum vulgare</i> Extracts and Their Analytical Characterization Anna Macková, Pavel Mučaji, Ute Widowitz and Rudolf Bauer	1509
Revised Structures of Nardal and Nardin: Identity with Valerenal and Valerenic Acid Shahuraj H. Kadam, Shashikumar K. Paknikar and G. Venkateswara Rao	1513
Chemical Relationships between Liverworts of the Family Lejeuneaceae (Porellales, Jungermanniopsida) Agnieszka Ludwiczuk, Phiangphak Sukkharak, Robert Gradstein, Yoshinori Asakawa and Kazimierz Glowinski	1515
Minor Ent-abietane Diterpenoids from <i>Euphorbia guyoniana</i> Hamada Haba, Laurence Marcourt, Mohammed Benkhaled and Christophe Long	1519
Isolation, Characterization and Sensory Evaluation of a Hexa β-D-Glucopyranosyl Diterpene from <i>Stevia rebaudiana</i> Indra Prakash, Venkata Sai Prakash Chaturvedula and Avetik Markosyan	1523
Structure and Biological Action of Cladolosides B₁, B₂, C, C₁, C₂ and D, Six New Triterpene Glycosides from the Sea Cucumber <i>Cladolabes schmeltzii</i> Alexandra S. Silchenko, Anatoly I. Kalinovskiy, Sergey A. Avilov, Pelageya V. Andryjaschenko, Pavel S. Dmitrenok, Ekaterina A. Yurchenko, Igor Yu. Dolmatov, Vladimir I. Kalinin and Valentin A. Stonik	1527
5α,8α-Epidioxysterols from a Formosan Sponge, <i>Axinyssa</i> sp. Tzu-Rong Su, Kai-Ju Liang, Michael Y. Chiang, Mei-Chin Lu, Yu-Jen Wu and Jui-Hsin Su	1535
Ecology- and Bioassay-Guided Drug Discovery for Treatments of Tropical Parasitic Disease: 5α,8α-Epidioxysterols-6-en-3β-ol Isolated from the Mollusk <i>Dolabrifera dolabrifera</i> Shows Significant Activity against <i>Leishmania donovani</i> Kathryn E. Clark, Angela Capper, Gina Della Togna, Valerie J. Paul, Luz I. Romero, Timothy Johns, Luis Cubilla-Rios and Todd L. Capson	1537
Alkaloids from <i>Chlidanthus fragrans</i> and their Acetylcholinesterase, Butyrylcholinesterase and Prolyl Oligopeptidase Activities Lucie Cahliková, Martina Hrabínová, Andrea Kulhánková, Nina Benešová, Jakub Chlebek, Daniel Jun, Zdeněk Novák, Kateřina Macáková, Jiří Kuneš, Kamil Kuča and Lubomír Opletal	1541

Further Evidence on the Intramolecular Lactonization in Rat Liver Microsomal Metabolism of 12-O-acetylated Retronecine-type Pyrrolizidine Alkaloids Jun Tang, Zhengtao Wang, Teruaki Akao and Masao Hattori	1545
<i>Uncaria tomentosa</i> Alkaloidal Fraction Reduces Paracellular Permeability, IL-8 and NS1 Production on Human Microvascular Endothelial Cells Infected with Dengue Virus Raimundo Sousa Lima-Junior, Cintia da Silva Mello, Antonio Carlos Siani, Ligia M. Marino Valente and Claire Fernandes Kubelka	1547
Antioxidant Activity of Zyzzyanones and Makaluvamines from the Marine Sponge <i>Zyzzya fuliginosa</i> Natalia K. Utkina	1551
Quantitative Analysis of Catechin, Flavonoid, and Saponin Constituents in “Tea Flower”, the Flower Buds of <i>Camellia sinensis</i>, from Different Regions in Taiwan Toshio Morikawa, I-Jung Lee, Shuhei Okugawa, Sohachiro Miyake, Yoshinobu Miki, Kiyofumi Ninomiya, Niichiro Kitagawa, Masayuki Yoshikawa and Osamu Muraoka	1553
<i>In Vitro</i> Inhibitory Effects of <i>Moringa oleifera</i> Leaf Extract and Its Major Components on Chemiluminescence and Chemotactic Activity of Phagocytes Boonyadist Vongsak, Wandee Gritsanapan, Yuvadee Wongkrajang and Ibrahim Jantan	1559
Acyated Delphinidin Glycosides from Violet and Violet-Blue Flowers of <i>Clematis</i> Cultivars and their Coloration Keisuke Sakaguchi, Junichi Kitajima and Tsukasa Iwashina	1563
Antimicrobial Activity of Extracts and Fractions from Aerial Parts of Selected Plants (<i>Garcinia achachairu</i>, <i>Macrosiphonia velame</i>, <i>Rubus niveus</i> and <i>Pilea microphylla</i>) Against Some Pathogenic Microorganisms Carla Melim, Karoliny Guimarães, Zhelmy Martin-Quintal, Aurea Damaceno Alves, Domingos Tabajara de Oliveira Martins, Franco Delle Monache, Valdir Cechinel Filho, Alexandre Bella Cruz and Rivaldo Niero	1567
Characterization of cDNA Encoding Resveratrol Synthase and Accumulation of Resveratrol in Tartary Buckwheat Yeon Bok Kim, Aye Aye Thwe, YeJi Kim, Sun Kyung Yeo, Chanhui Lee and Sang Un Park	1571
Norbergenin Derivatives from <i>Diospyros crassiflora</i> (Ebenaceae) Carine Mvot Akak, Augustin Ephrem Nkengfack and Peng-Fei Tu	1575
Inhibitory Effects of Ethyl Acetate-Soluble Fraction from <i>Morus alba</i> on Lipid Accumulation in 3T3-L1 Cells Hee-Sook Park, Soon-Mi Shim and Gun-Hee Kim	1579
Neuritogenic and Neuroprotective Activities of Fruit Residues Sarin Tadtong, Mayuree Kanlayavattanukul and Nattaya Lourith	1583
Total Polyphenol Content, <i>in vitro</i> Antifungal and Antioxidant Activities of Callus Cultures from <i>Inula crithmoides</i> Anahi Buccini, Laura Giamperi and Donata Ricci	1587
Novel Antiplasmodial Agents from <i>Christia vespertilionis</i> Harish C. Upadhyay, Brijesh S. Sisodia, Harveer S. Cheema, Jyoti Agrawal, Anirban Pal, Mahendra P. Darokar and Santosh K. Srivastava	1591
Synthesis and Biological Evaluation of Nonaprenylsulfates Suryachandra R. Ravada, Lakshmareddy Emani, Ganeswara R. Mallula and Trimurtulu Golakoti	1595
Antifungal Activity of Tetrasulfanes against <i>Botrytis cinerea</i> Brigitte Czepukojc, Michaela Leroch, Franz Salm, Uma M. Viswanathan, Torsten Burkholz, Matthias Hahn and Claus Jacob	1599
Identification of Sake Extract as a New Anti-melanogenic Ingredient by <i>in vitro</i> and Clinical Trials Jeong-Ah Hwang, Myeong-Jin Goh, Eun-Joo Kim, Myong-Ryul Lee, Nok-Hyun Park, Yong-Joo Na, Jun-Cheol Cho and Hae-Kwang Lee	1605
Therapeutic Effect of <i>Broussonetia papyrifera</i> and <i>Lonicera japonica</i> in Ovalbumin-induced Murine Asthma Model Seong-Ho Hong, Jung-Taek Kwon, Ji-Young Shin, Ji-Eun Kim, Arash Minai-Tehrani, Kyeong-Nam Yu, Somin Lee, Sung-Jin Park, Seung-Hee Chang, Hu-Lin Jiang, M Vibin, Kiwon Han, Kun-Ho Son, Wie-Jong Kwak, Chanhee Chae, Sung-Hye Bang and Myung-Haing Cho	1609
<i>Crambe tataria</i>: The Glucosinolate/Myrosinase System <i>in vivo</i> and <i>in vitro</i> Anna Piovan, Raffaella Filippini and Rosy Caniato	1615
Antimicrobial and Biofilm Effects of Herbs Used in Traditional Chinese Medicine Denny Lau and Balbina J. Plotkin	1617
Homologous <i>Canavalia</i> Lectins Elicit Different Patterns of Antinociceptive Responses Nilson Vieira Pinto, Cláudia Ferreira Santos, Benildo Sousa Cavada, Kyria Santiago do Nascimento, Francisco Nascimento Pereira Junior, Alana de Freitas Pires and Ana Maria Sampaio Assreuy	1621
Volatile Constituents of <i>Trifolium pratense</i> spp. <i>nivale</i> Quantified at Different Growth Stages, and Evaluation of their Antimicrobial Activity Roberto Cecotti, Emanuele Carpana, Patrizia Bergomi and Aldo Tava	1625
Chemical Composition and Free Radical Scavenging Activity of the Essential Oil of <i>Achillea ligustica</i> Growing Wild in Lipari (Aeolian Islands, Sicily) Mariem Ben Jemia, Zyed Rouis, Antonella Maggio, Alessandro Venditti, Maurizio Bruno and Felice Senatore	1629
Chemical Analysis and Calcium Channel Blocking Activity of the Essential Oil of <i>Perovskia abrotanoides</i> Abdul Jabbar Shah, Munawwer Rasheed, Qaiser Jabeen, Amir Ahmed, Rasool Bakhsh Tareen, Anwarul Hassan Gilani, Muhammad Nadir and Viqar Uddin Ahmad	1633
Chemical Composition and <i>In vitro</i> Cytotoxic Activity of the Essential Oils of <i>Stachys rupestris</i> and <i>Salvia heldreichiana</i>, Two Endemic Plants of Turkey Elif Ayşe Erdogan, Ayşe Everest, Laura De Martino, Emilia Mancini, Michela Festa and Vincenzo De Feo	1637
Chemical Composition of the Essential Oil from the Leaves of <i>Carapa guianensis</i> Collected from Venezuelan Guayana and the Antimicrobial Activity of the Oil and Crude Extracts Gina Meccia, Patricia Quintero, Luis B. Rojas, Alfredo Usabillaga, Judith Velasco, Tulia Diaz, Clara Diaz, Jesús Velásquez and Maria Toro	1641

Chemical Composition of the Essential Oil of the Flowering Aerial Parts of <i>Pimpinella monoica</i> Rajesh K. Joshi	1643
Cytotoxic Activity of Essential Oils of Aerial Parts and Ripe Fruits of <i>Echinophora spinosa</i> (Apiaceae) Daniele Fraternale, Donata Ricci, Cinzia Calcabrini, Michele Guescini, Chiara Martinelli and Piero Sestili	1645
<u>Review/Account</u>	
Human Body Scents: Do they Influence our Behavior? Sophie Mildner and Gerhard Buchbauer	1651

Number 12

New Humulenes from <i>Hyptis incana</i> (Labiatae) Mitsuru Satoh, Yoshio Satoh, Yasuhiro Anzai, Daisuke Ajisawa, Keiichi Matsuzaki, Mitsuko Makino and Yasuo Fujimoto	1665
Inhibitory Effects against Pasture Weeds in Brazilian Amazonia of Natural Products from the Marine Brown Alga <i>Dictyota menstrualis</i> Rainiomar Raimundo Fonseca, Antonio Pedro Silva Souza Filho, Roberto Campos Villaça and Valéria Laneuville Teixeira	1669
Isolation of the Plant Hormone (+)-Absciscic acid as an Antimycobacterial Constituent of the Medicinal Plant Endophyte <i>Nigrospora</i> sp. Trevor N. Clark, Katelyn Ellsworth, Haoxin Li, John A. Johnson and Christopher A. Gray	1673
New Cembranoid Diterpene from the South China Sea Soft Coral <i>Sarcophyton</i> sp. Fei Cao, Jing Zhou, Kai-Xia Xu, Meng-Qi Zhang and Chang-Yun Wang	1675
Crotofolane Diterpenoids from <i>Croton caracasanus</i> Katuska Chávez, Reinaldo S. Compagnone, Ricarda Riina, Alexander Briceño, Teresa González, Emilio Squitieri, Carlos Landaeta, Humberto Soscún and Alírica I. Suárez	1679
Development and Validation of a Modified Ultrasound-Assisted Extraction Method and a HPLC Method for the Quantitative Determination of Two Triterpenic Acids in <i>Hedyotis diffusa</i> Yu-Chiao Yang, Ming-Chi Wei, Hui-Fen Chiu and Ting-Chia Huang	1683
New Triterpenoid Saponins from the Roots of <i>Saponaria officinalis</i> Barbara Moniuszko-Szajwaj, Łukasz Pecio, Mariusz Kowalczyk, Ana M. Simonet, Francisco A. Macias, Małgorzata Szumacher-Strabel, Adam Cieślak, Wiesław Oleszek and Anna Stochmal	1687
Minor Triterpene Saponins from Underground Parts of <i>Lysimachia thyrsiflora</i>: Structure elucidation, LC-ESI-MS/MS Quantification, and Biological Activity Irma Podolak, Paweł Żmudzki, Paulina Koczurkiewicz, Marta Michalik, Paweł Zajdel and Agnieszka Galanty	1691
Variation of Saponin Contents and Physiological Status in <i>Quillaja saponaria</i> Under Different Environmental Conditions Angélica Grandón S, Miguel Espinosa B, Darcy Ríos L, Manuel Sánchez O, Katia Sáez C, Víctor Hernández S. and José Becerra A	1697
New Access to 7,17-seco C₁₉-Diterpenoid Alkaloids via Vacuum Pyrolysis of <i>N</i>-Deethyl-8-acetyl Derivatives Ling Wang, Qi-Feng Chen and Feng-Peng Wang	1701
Alkaloids from <i>Boophone haemanthoides</i> (Amaryllidaceae) Jerald J. Nair, Lucie Rárová, Miroslav Strnad, Jaume Bastida and Johannes van Staden	1705
Supinidine Viridiflorates from the Roots of <i>Chromolaena pulchella</i> Mario A. Gómez-Hurtado, J. Martín Torres-Valencia, Rosa E. del Río, Gabriela Rodríguez-García, Virginia Motilva, Sofía García-Mauriño, Carlos M. Cerda-García-Rojas and Pedro Joseph-Nathan	1711
<i>N</i>-Containing Metabolites from the Marine Sponge <i>Agelas clathrodes</i> Fan Yang, Rui-Hua Ji, Jiang Li, Jian-Hong Gan and Hou-Wen Lin	1713
Two New Compounds and Anti-complementary Constituents from <i>Amomum tsao-ko</i> Jiahong Jin, Zhihong Cheng and Daofeng Chen	1715
Antiangiogenic Activity of Flavonoids from <i>Melia azedarach</i> Shigenori Kumazawa, Satomi Kubota, Haruna Yamamoto, Naoki Okamura, Yasumasa Sugiyama, Hirokazu Kobayashi, Motoyasu Nakanishi and Toshiro Ohta	1719
Application of Mixture Analysis to Crude Materials from Natural Resources (IV)^[1(a-c)]: Identification of <i>Glycyrrhiza</i> Species by Direct Analysis in Real Time Mass Spectrometry (II) Eriko Fukuda, Yoshihiro Uesawa, Masaki Baba and Yoshihito Okada	1721
Comparison of Total Phenolic Content, Scavenging Activity and HPLC-ESI-MS/MS Profiles of Both Young and Mature Leaves and Stems of <i>Andrographis paniculata</i> Lee Suan Chua, Ken Choy Yap and Indu Bala Jaganath	1725
Xanthones from aerial parts of <i>Hypericum laricifolium</i> Juss. Irama Ramírez-González, Juan Manuel Amaro-Luis and Ali Bahsas	1731
A New Xanthone from the Pericarp of <i>Garcinia mangostana</i> Manqin Fu, Samuel X. Qiu, Yujuan Xu, Jijun Wu, Yulong Chen, Yuanshan Yu and Gengsheng Xiao	1733
Isolation of a Phomoxanthone A Derivative, a New Metabolite of Tetrahydroxanthone, from a <i>Phomopsis</i> sp. Isolated from the Mangrove, <i>Rhizophora mucronata</i> Yoshihito Shiono, Takehiro Sasaki, Fumiaki Shibuya, Yukito Yasuda, Takuya Koseki and Unang Supratman	1735
Anti-allergic Inflammatory Effects of Cyanogenic and Phenolic Glycosides from the Seed of <i>Prunus persica</i> Geum Jin Kim, Hyun Gyu Choi, Ji Hyang Kim, Sang Hyun Kim, Jeong Ah Kim and Seung Ho Lee	1739

Isolation, Synthesis and Biological Evaluation of Phenylpropanoids from the Rhizomes of <i>Alpania galanga</i> Sumit S Chourasiya, Eppakayala Sreedhar, K. Suresh Babu, Nagula Shankaraiah, V. Lakshma Nayak, S. Ramakrishna, S. Sravani and M.V. Basaveswara Rao	1741
Rosmarinic Acid Interaction with Planktonic and Biofilm <i>Staphylococcus aureus</i> Livia Slobodníková, Silvia Fialová, Helena Hupková and Daniel Grančai	1747
New Butenolide and Pentenolide from <i>Dysidea cinerea</i> Phan Van Kiem, Nguyen Xuan Nhiem, Ngo Van Quang, Chau Van Minh, Nguyen Hoai Nam, Nguyen Thi Cuc, Hoang Le Tuan Anh, Bui Huu Tai, Pham Hai Yen, Nguyen Xuan Cuong, Nguyen Phuong Thao, Nguyen Thi Hoai, Nan Young Kim, Seon Ju Park and Kim Seung Hyun	1751
A New Cyclopeptide from Endophytic <i>Streptomyces</i> sp. YIM 64018 Xueqiong Yang, Yabin Yang, Tianfeng Peng, Fangfang Yang, Hao Zhou, Lixing Zhao, Lihua Xu and Zhongtao Ding	1753
Involvement of Trypsin-Digested Silk Peptides in the Induction of RAW264.7 Macrophage Activation Kyoung-Ho Pyo, Min-Ki Kim, Kwang-Soon Shin, Hyang Sook Chun and Eun-Hee Shin	1755
Low-Volatile Lipophilic Compounds in Needles, Defoliated Twigs, and Outer Bark of <i>Pinus thunbergii</i> Alexander V. Shpatov, Sergey A. Popov, Olga I. Salnikova, Ekaterina A. Khokhrina, Emma N. Shmidt and Byung Hun Um	1759
Lipid Constituents of the Edible Mushroom, <i>Pleurotus giganteus</i> Demonstrate Anti-Candida Activity Chia-Wei Phan, Guan-Serm Lee, Ian G. Macreadie, Sri Nurestri Abd Malek, David Pamela and Vikineswary Sabaratnam	1763
Effect of Trehalose Addition on Volatiles Responsible for Strawberry Aroma Mirela Kopjar, Janez Hribar, Marjan Simčič, Emil Zlatić, Tomaž Požrl and Vlasta Piližota	1767
<i>Pogostemon hirsutus</i> Oil, Rich in Abietane Diterpenes Ramar Murugan, Gopal Rao Mallavarapu, Veerappan Sudha and Pemaiah Brindha	1771
Combined Analysis of the Root Bark Oil of <i>Cleistopholis glauca</i> by Chromatographic and Spectroscopic Techniques Zana A. Ouattara, Jean Brice Boti, Coffy Antoine Ahibo, Félix Tomi, Joseph Casanova and Ange Bighelli	1773
Bioactivities and Compositional Analyses of <i>Cinnamomum</i> Essential Oils from Nepal: <i>C. camphora</i>, <i>C. tamala</i>, and <i>C. glaucescens</i> Prabodh Satyal, Prajwal Paudel, Ambika Poudel, Noura S. Dosoky, Kiran Kumar Pokharel, and William N. Setzer	1777
Essential Oil Characterization of Two Azorean <i>Cryptomeria japonica</i> Populations and Their Biological Evaluations Cristina Moiteiro, Teresa Esteves, Luís Ramalho, Rosario Rojas, Sandra Alvarez, Susana Zacchino and Helena Bragança	1785
Antioxidant, Antiproliferative and Antimicrobial Activities of the Volatile Oil from the Wild Pepper <i>Piper capense</i> Used in Cameroon as a Culinary Spice Verlaine Woguem, Filippo Maggi, Hervet P. D. Fogang, Léon A. Tapondjou, Hilaire M. Womeni, Luana Quassinti, Massimo Bramucci, Luca A. Vitali, Dezemona Petrelli, Giulio Lupidi, Fabrizio Papa, Sauro Vittori and Luciano Barboni	1791
<u>Review/Account</u>	
Boldine and Related Aporphines: From Antioxidant to Antiproliferative Properties Darina Muthna, Jana Cmielova, Pavel Tomsik and Martina Rezacova	1797
New Therapeutic Potentials of Milk Thistle (<i>Silybum marianum</i>) Nataša Milić, Nataša Milošević, Ljiljana Suvajdžić, Marija Žarkov and Ludovico Abenavoli	1801
Biomedical Properties of Edible Seaweed in Cancer Therapy and Chemoprevention Trials: A Review Farideh Namvar, Paridah Md. Tahir, Rosfarizan Mohamad, Mahnaz Mahdavi, Parvin Abedi, Tahereh Fathi Najafi, Heshu Sulaiman Rahman and Mohammad Jawaaid	1811
Methods for Extraction and Determination of Phenolic Acids in Medicinal Plants: A Review Agneszka Arceusz, Marek Wesolowski and Pawel Konieczynski	1821

Natural Product Communications

Volume 8 (1-12)

2013

Author Index

- | | | | |
|-----------------------------------|-------------------------------------|--------------------------------------|---------------------------------------|
| Abas, F 447 | Andryjashchenko, PV 1377,1527 | Belinga, FNF 829 | Brun, R 1135 |
| Abdel-Baky, AM 191 | Anh, HLT 1209,1751 | Benajiba, N 51 | Bruno, M283,567,841,1159,1629 |
| Abdissa, N 1261 | Antonov, AS 1219 | Bencharif-Betina, S 573 | Brunton, N 1123 |
| Abe, N 333 | Anuntalabhochai, S 519 | Bencsik, T 1247 | Bruyère, C 1089 |
| Abedi, P 1811 | Anzai, Y 1665 | Benešová, N 781,1541 | Buabeid, MA 1415 |
| Abenavoli, L 1801 | Aoyagi, S 851 | Benkhalel, M 1519 | Buatois, B 509 |
| Abraham, MH 85 | Aramwit, P 501 | Benkó, R 1247 | Bucar, F 523 |
| Abbrankó, L 605 | Araujo, L 1175 | Benlhabib, O 1155 | Bucchini, A 1587 |
| Abubakr, M 1487 | Arceusz, A 1821 | Benoit-Vical, F 1099 | Buchbauer, G 1651 |
| Acuña-Castroviejo, D 47 | Arnold, AE 601 | Benouhoud, M 51 | Bucio, MA 1075 |
| Aderogba, MA 795 | Arseniyadis, S 965 | Benson, CS 183 | Budesinsky, M 1423 |
| Adewusi, EA 1213 | Arsilan, S 1269 | Bensouda, Y 55 | Bukvicki, DR 1313 |
| Adlouni, A 51 | Arya, D 635 | Bergomi, P 1625 | Bulgakov, VP 1419 |
| Afiyatullof, SS 1071 | Asakawa, Y 565,859,1515 | Berkov, S 327 | Burcul, F 471 |
| Agnau, H 55 | Assreuy, AMS 1621 | Berlinck, RGS 721 | Burkholz, T 1599 |
| Agrawal, J 1591 | Atmaca, P 1269 | Bessière, JM 509 | Byamukama, R 1261 |
| Ahibo, CA 1773 | Atsawinkowit, C 443 | Bezić, N 645 | |
| Ahmad, A 261 | Aversano, R 1387 | Bhadra, S 671 | Caamal-Fuentes, E 1421 |
| Ahmad, F 261,1317 | Avilov, SA 301,1053,1377,1527 | Bhattacharya, SM 803 | Cabrera-Vique, CC 47 |
| Ahmad, VU 757,1289,1633 | Avoseh, NO 243 | Bhutani, KK 195 | Cahliková, L 441,781,1541 |
| Ahmed, A 1633 | Awale, S 463 | Bhutia, TD 1081 | Cai, B 1447,1479 |
| Ahmed, I 103 | Awasthi, SK 803 | Bi, WF 621 | Cai, H 1447,1479 |
| Ahmed, SA 153 | Ayyad, SEN 351 | Bian, B 1137 | Calcebrini, C 1645 |
| Ahn, JH 481 | Azam, F 1223 | Bighelli, A 235,1155,1165,1773 | Calderón, AI 1135 |
| Ahuja, M 1415 | Aziz, A 513 | Bishay, DW 191 | Callmander, MW 1201 |
| Ain-Lhout, F 5,11,15 | Azmat, S 1289 | Blanchfield, JT 725 | Calpena, A 177 |
| Ajisawa, D 1665 | | Blažević, I 471,1463 | Caniato, R 1615 |
| Akai, S 987 | Baba, M 1721 | Blunden, G 63 | Cao, F 1675 |
| Akak, CM 1575 | Bačeva, K 639 | Bodo, B 459 | Cao, X 479 |
| Akala, H 1261 | Badyal, DK 359 | Bogucka – Kocka, A 545 | Capasso, R 419,1333 |
| Akala, HH 175 | Bahsas, A 1731 | Bohman, B 701 | Capper, A 1537 |
| Akao, T 1545 | Bairwa, K 1241 | Bojin, F 343 | Capson, TL 1537 |
| Akimoto, T 961 | Bajaj, VK 1183 | Bonaccorsi, I 171 | Cárcamo, G 147 |
| Alarcón, AB 1237 | Bakhy, K 1155 | Bonechi, C 429 | Carmona, J 1175,1477 |
| Al-Fatimi, MA 683 | Bala, S 359 | Bonikowski, R 1297 | Carpana, E 1625 |
| Alfawaz, MA 213 | Bamou, Y 51 | Bonikowski, R 227 | Carpito, D 1387 |
| Ali, A 257 | Banaszczak, P 227,1297 | Bora, J 1451 | Carreira, LMM 1471 |
| Ali, NAA 683 | Banerjee, S 1487 | Borcan, F 343 | Caruso, I 1387 |
| Ali, NM 513 | Bang, SH 1609 | Bordoloi, M 1451 | Carvalho, EEN 1471 |
| Ali, SI 1365 | Bankova, V 617 | Borrelli, F 1333 | Casabianca, H 829 |
| Ali, Z 319,323,333,757 | Banuls, LMY 439 | Boti, JB 1773 | Casanova, J 235,1155,1165,1773 |
| Aljančić, I 1291 | Banzouzi, JT 1099 | Bouhaouss, A 21 | Cassera, MB 1201 |
| Allali, H 835 | Bao, J 1127 | Boulmane, M 21 | Cavada, BS 1621 |
| Al-Mekhlafi, NA 447 | Barboni, L 1791 | Boussoussa, H 1171 | Cecotti, R 1625 |
| Alsaif, MA 213 | Barradas, MCD 5,11 | Boutaleb, S 5,11,15 | Çelik, T 59 |
| Al-Shagdari, A 1237 | Barraza-Morales, A 1245 | Bowen, WD 811 | Čepo, DV 645 |
| Alshatwi, AA 213 | Barrow, RA 701 | Bowman, JD 1201 | Cerda-García-Rojas, CM 1711 |
| Alvarez, S 1785 | Barthó, L 1247 | Bracho, I 1477 | Chae, C 1609 |
| Álvarez-Cansino, L 15 | Basaveswara Rao, MV 1741 | Bradshaw, TD 375,1423 | Chafchaouani-Moussaoui, I 43,48 |
| Álvarez-Román, R 135 | Baser, KHC 257 | Bragança, H 1785 | Chalard, P 1099 |
| Alves, AD 1567 | Bastida, J 327,553,1335,1705 | Bramucci, M 1791 | Chan, K 707 |
| Alviano, DS 69,424 | Bastos, JK 1041 | Brancazio, CL 283 | Chandler, CM 1059 |
| Amaro-Luis, JM 1731 | Batçıoğlu, K 475 | Branco, A 1049 | Chang, CI 79,805 |
| Amat, AG 539 | Bates, RB 1195 | Brantner, AH 523,1137 | Chang, HS 805,827 |
| Ambrose, KV 1059 | Bauer, R 105,1509 | Bravo, L 1325 | Chang, SH 1609 |
| Aminin, DL 1377 | Bazzali, O 235 | Brecker, L 1081 | Chanotiya, CS 221 |
| Amira-Guebailia, H 485 | Bchitou, R 21 | Briceño, A 1679 | Chansakaow, S 519 |
| Amri, I 407 | Becerra, J 147,397,1697 | Brindha, P 1771 | Chanvorachote, P 115 |
| Andrade, EHA 1471 | Becker, K 1135 | Brkljača, R 729 | Chao, CY 805,827 |
| Andreev, YA 1229 | Beckord, HJ 351 | Brodie, PJ 1201 | Charrouf, Z 25,29,33,43,55 |
| Andryjaschenko, PV 301,1053 | Beedessee, G 635 | Brown, SG 351 | Chaturvedula, VSP 1523 |

- Chaudhary, D 911
 Chaudhary, SK 671
 Chaves, MH 277
 Chávez, K 1679
 Cheema, HS 1591
 Chejara, DR 497
 Chen, CR 79
 Chen, D 1715
 Chen, F 155,1115
 Chen, J 83,587
 Chen, JJ 597
 Chen, L 339
 Chen, QF 1701
 Chen, S 457
 Chen, Y 1121,1479,1733
 Chen, YC 827
 Cheng, HL 79
 Cheng, NF 389
 Cheng, R 461
 Cheng, Z 1715
 Cheong, JH 1143
 Chhatbar, MU 497
 Chiang, MY 1535
 Chiu, HF 1683
 Chivandi, E 411
 Chlebek, J 1541
 Chlebek, J 781
 Cho, JC 1605
 Cho, MH 1609
 Cho, S 1149
 Choi, HG 181,1739
 Choi, K 1301
 Choudhary, MI 757
 Chourasiya, SS 1741
 Chovolou, Y 289
 Chu, J 375,1423
 Chua, LS 1725
 Chun, HS 1755
 Chung, HM 1037
 Chung, SJ 1143
 Cian, MCD 655
 Cicala, C 539
 Cieślak, A 1687
 Clares-Naveros, B 177
 Clark, KE 1537
 Clark, TN 1673
 Clarke, G 375
 Clmati, E 429
 Cmielova, J 1797
 Codina, C 1335
 Compagnone, R S 1679
 Conforti, F 567
 Copp, BR 699
 Cordero-Pérez, P 297
 Corell, M 1123
 Cossy, J 965
 Costa, EV 403
 Costa, J 655,835
 Costa-Lotufo, LV 69,424
 Cotroneo, A 171
 Cresteil, T 635
 Crkvenčić, M 347
 Crouch, RA 683
 Cruz, AB 1567
 Cruz, C 1459
 Cruz, MNS 69,424
 Cubilla-Rios, L 1537
 Cuc, NT 1751
 Cuesta-Rubio, O 1237
 Cuong, NX 1751
 Cushnie, TPT 385
 Cutler, SJ 191,1117,1285
 Cvetković, OG 1309
 Czepukojc, B 1599
 Czigle, S 1153
 Da Costa Mousinho, NMH 1279
 da Silva, EBP 277
 Dabhade, A 1467
 Dagne, E 1261
 Dai, DN 239,663
 Dai, H 337
 Dam, NP 1507
 Danciu, C 343
 Dangarembizi, R 411
 Danielle, FR 367
 Darokar, MP 1591
 David, P 711
 de Almeida, MRA 207
 de Armas, Y 1325
 de Carvalho, BA 679
 De Feo, V 539
 Dehelean, C 343
 Delaunay, JC 485
 Delemasure, S 157
 Demirci, B 257
 Demirci, F 59
 Deng, Z 83,587
 Denhez, C 25,33,55
 Denisenko, VA 1399
 Denisenko, VA 587,1071,1229
 Denkert, A 683
 Derese, S 175
 Derouiche, A 51,55
 Deville, A 459
 Devkota, HP 583,1413
 Dewanjee, S 1275
 Dey, S 253
 Dhanasekaran, M 1415
 Dholvitayakhun, A 385
 Diaz, C 1641
 Diaz, T 1641
 Dib, MEA 835
 Din, WM 375
 Ding, W 487
 Ding, Z 1393,1753
 Djabou, N 835
 Dmitrenok, PS 301,381,587,1053
 Dmitrenok, PS 1071,1229,1399
 Dmitrenok, PS 1527
 Dodin, S 55
 Dolmatov, IY 1527
 Domingos, OS 679
 Domingos, VD 1471
 Domingues, F 1459
 Domingues, FC 673
 Domínguez-Villegas, V 177
 Domon, D 929
 dos Santos, MH 679
 Dosoky, NS 1179,1481,1777
 Dowell, A 737
 Drašar, P 771
 Driouch, I 51
 Du, ZT 1197
 Duan, JA 487
 Duarte, AF 673
 Duarte, AP 1459
 Duarte, GP 1167
 Duez, P 439
 Dugo, P 171
 Dumté, M 693
 Dung, TD 1507
 Dunkić, V 645
 Dutartre, P 157
 Dutta, PP 1451
 Dvorakova, M 105
 Dyshlovoy, SA 1399
 Dyshlovoy, SA 381
 Eaton, AL 1201
 Echeverri-Gil, D 129
 Edrada-Ebel, R 289
 Egamberdieva, D 429
 Ehrlich, AL 1195
 EL-Baz, FK 1365
 El-Ghaly, S 1117
 El-Hela, AA 1117
 Ellis, J 257
 Ellsworth, K 1673
 ElMarrouni, A 965
 Emani, L 1595
 Epifano, F 1089
 Erdogan, EA 1637
 Erlwanger, K 411
 Ermolovich, YV 771
 Escames, G 47
 Espartero, JL 1255
 Espinosa B, M 1697
 Espinoza, C 187
 Esquivias, MP 11,15
 Esteves, T 1785
 Evans, AC 689
 Evans, P 1123
 Everest, A 1637
 Fahmy, H 153
 Faiz, CA 1155
 Fan, CL 599
 Fedoreyev, SA 1419
 Fedoreyev, SA 587
 Feldo, M 545
 Felinger, A 1247
 Feng, G 199
 Feng, J 393
 Feng, Q 149
 Feo, VD 139,407,1637
 Fernández, JJ 187
 Ferreira, AG 69,424,721
 Ferreira, S 673
 Festa, M 1637
 Fialová, S 1747
 Fidaleo, M 579
 Figueiredo, CR 277,679
 Filho, APSS 1669
 Filho, RB 1049
 Filho, VC 1567
 Filippini, R 1615
 Fiorito, S 1089
 Fisher, AA 1059
 Flamini, G 243
 Flematti, G 701
 Flewelling, AJ 373
 Fogang, HPD 1791
 Fons, F 509
 Fonseca, RR 1669
 Formisano, C 567,841
 Fouad, MA 767
 Fouche, G 1213
 Froissard, D 509,527,1645
 Fruchier, A 509
 Fry, JR 375
 Fu, M 1733
 Fuh, YM 571
 Fuhshuku, K 897
 Fujii, K 919
 Fujimoto, Y 1665
 Fujiwara, K 929
 Fukuda, E 1721
 Fukuda, T 1409
 Fukusako, K 1413
 Fukushi, E 815
 Fukuyama, Y 915
 Furtado, RA 1041
 Furukawa, H 775
 Furusawa, M 859
 Gagnon, MJ 265
 Gaivelyte, K 1105
 Galanty, A 1691
 Gan, JH 1713
 Gan, LS 597
 Gangopadhyay, M 1275
 Gao, ZH 467
 García, ML 177
 García-Corzo, L 47
 García-Mauriño, S 1711
 Garduño-Ramírez, ML 177
 Garson, MJ 725
 Gavrilova, N 617
 Gaydou, EM 1171
 Gebauer, J 965
 Gebru, M 1261
 Genovese, S 527,1089
 Ghannadi, A 1355
 Gharby, S 25,29,33
 Ghoneim, MM 1117
 Giamperi, L 1587
 Gilani, AH 1633
 Gillum, V 311
 Gimazetdinov, AM 981
 Giordano, M 33
 Giovanni, CI 651
 Gligorijević, N 1291
 Glowniak, K 1515
 Go, G 823
 Godiya, CD 497
 Goh, MJ 1605
 Gökbulut, A 475
 Golakoti, T 1595
 Golfnopoulos, SK 1493
 Gómez, LA 249
 Gómez-Hurtado, MA 1711
 Gong, X 631
 González, GMG 135
 González, T 1679
 Goto, H 629
 Gottardi, DG 1313
 Gou, X 1115
 Gourine, N 1171
 Gradstein, R 1515
 Grančai, D 1747
 Grandón S, A 1697
 Gray, CA 373,1673
 Gray, S 257
 Greatrex, BW 737
 Green, IR 103
 Grishin, EV 1229
 Gritsanapan, W 1065,1559
 Grkovic, T 699
 Grover, J 1241
 Gu, J 1447
 Gudej, J 505
 Guerra-Boone, L 135
 Guerzoni, ME 1313
 Guescini, M 1645
 Gui, X 1217
 Guillaume, D 21,25,29,33,43,55
 Guimarães, ET 403
 Guimarães, K 1567
 Gulyakevich, OV 771
 Gunaherath, GMKB 1041,1431
 Günal, S 475
 Gunasekar, D 459
 Gunatilaka, AAL 601,1041
 Guo, JM 487
 Guo, XY 1473

- Guo, Y 461
Guo, Z 83,587
Gupta, J 803
Gupta, RS 1183
Gürsu, BY 59
Guzii, AG 1229,1399
- Ha, H 1443
Ha, LT 367
Haba, H 1519
Habig, JW 1059
Hagiwara, H 873,935
Hahn, M 1599
Ham, YM 1359
Hamada, H. 119,847,903,905,907
Hamada, M 847,903,905,907
Hamdouchi, AE 55
Hamed, AN 289
Hamia, C 1171
Han, D 203
Han, K 1609
Han, L 1149
Han, W 787
Han, Z 329
Hang, DTT 1209
Hapuarachchi, SD 333
Harada, K 915
Harada, Y 1367
Harhar, H 25,29,33
Harinantenaina, L 1201
Haritakun, R 703
Hasan, TN 213
Hasegawa, T 869
Hashimoto, T 859
Hassan, AO 243
Hasuie, Y 1409
Hata, E 515
Hattori, M 1545
Hattori, S 629
Havlik, J 1423
Hayashi, S 515
Háznagy-Radnai, E 1153
He, F 329,1069
He, MX 1427
Hegazy, AK 1365
Heras, M 965
Hernández S., V 1697
Hernández, J 1477
Hernandez, V 397
Herrero-Martínez, JM 1325
Hetta, MH 311
Heyden, YV 1325
Heydenreich, M 175,1261
Hilali, M 25
Hioki, H 915
Hirabayashi, M 1409
Hiwatashi, N 629
Ho, CL 143,273,531,1329
Hoai, NT 1209,1751
Höferl, M 663
Hoi, TM 235,239
Hong, LS 493
Hong, M 523
Hong, SH 1609
Honma, N 873
Hooper, JNA 725
Hopson, R 811
Horio, Y 905
Horváth, G 1247
Hoshi, T 873
Hosoda, R 905
Hošťálková, AH 441
Hoye, TR 351
Hrabínová, M 1541
- Hribar, J 1767
Hsu, JL 79
Hsu, KP 531,1329
Hu, A 1101
Hu, D 393
Hu, Z 753
Hua, H 1397
Huang, BK 1217
Huang, HL 113
Huang, LZ 1197
Huang, R 1121
Huang, TC 1683
Hui, Z 149
Hummelova, J 1423
Hung, MA 1507
Hupková, H 1747
Hwang, BY 481,1439
Hwang, JA 1605
Hwang, TL 1037
Hyöky, V 791
Hyun, CG 427,1359
Hyun, JW 427
Hyun, KS 1751
- Ialenti, A 539
Ibrahim, D 493
Ichiishi, E 153
Ifzal, R 1289
Ihandriharison, H 1165
Iida, A 153
Ikeda, K 627
Ikushima, K 1367
Ilhan, S 59
Ilić, BS 1309
Imai, H 119
Imaoka, T 961
Induli, M 1261
Inokuchi, T 919
Ionta, M 679
Irshad, M 261
Ishiguro, K 1413
Ishihara, K 903
Ito, C 775,1265
Itoigawa, M 775,1265
Ivanchina, NV 1219
Ivanova, NA 981
Iwakiri, T 119
Iwamoto, Y 631
Iwashina, T. 453,1093,1251,1563
Iwata, M 961
Izumi, H 869
- Jabeen, Q 1633
Jachak, SM 1241
Jacob, C 1599
Jacob, JN 359,807,811
Jacob, MR 761
Jacobson, GA 211
Jadrantin, M 1291
Jafri, A 51
Jaganath, IB 1725
Jagtap, S 195
Jakstas, V 1105
Jani, NA 513
Jankiprasad 803
Jankovic, SM 1313
Jantan, I 1559
Janulis, V 1105
Jáuregui, J 5,15
Jawaid, M 1811
Jayasandhya, P 301
Jayawardana, KH 1431
Jemia, MB 1629
Jenett-Siems, K 765
- Jeong, SJ 1443
Jerković, I 231,651
Jeyaraj, EJ 447
Jha, Z 1045
Jho, KH 217
Ji, L 1101
Ji, RH 1713
Ji, X 1217
Jiang, HL 1609
Jiang, J 1129
Jiang, X 1121
Jiang, Y 1473
Jiang, ZH 169
Jiec, S 523
Jiménez, ST 1059
Jin, F 211
Jin, HE 1143
Jin, J 1715
Jin, KT 375
Jin, QA 199
Jirošová, A 441
Jirovetz, L 663
Jo, YH 481
John, PA 711
Johns, T 1537
Johnson, JA 373,1673
Jones, GL 737,741,747
Jordán, MJ 817
Joseph-Nathan, P 1075,1711
Joshi, KR 583
Joshi, RK 225,401,1163,1643
Jovanović, O 109
Julkunen-Tiitto, RJ 791
Jumaryatno, P 725
Jun, D 1541
Jun-feng, L 63
Jun-feng, Z 63
Junior, FNP 1621
Junior, HMS 69,424
Juranic, ZD 1291
Jusoh, S 1317
- Kabouche, Z 573
Kadam, SH 1195,1513
Kadota, S 463
Kaennakam, S 1111,1371
Kaewkroek, K 1205
Kakiuchi, K 1021
Kalinin, VI. 301,1053,1377,1527
Kalinovskaya, NI 381
Kalinovsky, AI. 1219,1419,1527
Kalinovsky, AI 301,381,1053
Kalle, W 389
Kam, A 707
Kamada, T 287
Kamchonwongpaisan, S 1305
Kamchonwongpaisan, S 733
Kamel, MS 289,767
Kaneda, N 775
Kang, MC 427
Kanlayavattanukul, M 1583
Karapandzova, M 639
Kari, KE 55
Karmase, A 195
Kartah, B 33,55
Kartah, BE 25,29
Kasemwattanaaroj, P 1257
Kashin, H 925
Kasuya, H 515
Katalinić, V 471
Kato, M 775
Katoch, D 161
Katoh, T 973
Katoono, R 929
- Kawabata, J 815
Kawahara, N 75
Kawahara, T 631
Kawai, H 929
Kawamura, N 929
Kawee-ai, A 1381
Kaya, GI 327
Kazakova, OB 293
Keeratinijakal, V 1065
Kelechi, AO 243
Keller, PA 703,733,1305
Kempińska, K 435
Kemppainen, B 1415
Kerubo, LO 175
Kerum, D 471
Khalifa, S 153
Khalivulla, SI 447
Khan, IA 257,319,323,333
Khan, SI 761
Khan, SN 757
Khan, SS 757
Khasmi, ME 51
Khisमतullin, R 617
Khokhrina, EA 1759
Khrpach, VA 771
Khumkratok, S 1371
Kicel, A 99,505
Kicha, AA 1219
Kida, H 75
Kiem, PV 1209,1751
Kikuchi, N 877
Kikuchi, T 1367
Kilár, F 1247
Kim, DD 1143
Kim, EJ 1605
Kim, GH 1579
Kim, GJ 1739
Kim, GO 427
Kim, H 1405
Kim, HM 1301
Kim, J 823
Kim, JA 181,1739
Kim, JE 1609
Kim, JH 1739
Kim, KS 1143
Kim, MK 1755
Kim, NY 1751
Kim, P 1301
Kim, SB 481
Kim, SH 181,1739
Kim, SJ 217,609,1233
Kim, SM 1381
Kim, SS 427
Kim, T 1301
Kim, WY 1143
Kim, YA 1405
Kim, YB 217,609,1233,1571
Kim, YJ 1571
Kim, YS 609,1149
Kingston, DGI 1201
Kinugawa, K 873
Kiran, I 59
Kirichuk, NN 1071
Kishi, A 851
Kishimoto, A 627
Kiss, R 1089
Kitagawa, N 1553
Kitajima, J. 453,1093,1251,1563
Kitisripanya, T 443
Klimešová, V 593
Knight, KS 299
Ko, RK 427
Kobayashi, H 1719
Kocki, J 545

- Koczurkiewicz, P 1691
 Koike, K 515
 Kokolakis, AK 1493
 Kokoska, L 105,1423
 Kolleth, A 965
 Komaikul, J 443
 Kondaveeti, S 497
 Kongkiatpaiboon, S 1065
 Kongsaree, P 1205
 Konieczynski, P 1821
 Konlata, J 1257
 Kopjar, M 1767
 Kord, MA 1365
 Korolkova, YV 1229
 Korth, J 733
 Kosalec, I 645
 Koseki, T 1735
 Kossuga, MH 721
 Kottob, S 1117
 Kouzayha, A 829
 Kowalczyk, M 1687
 Kowitdamrong, A 115
 Krasteva, I 765
 Kraus, GA 911
 Kremer, D 645
 Krohn, K 103
 Kubelka, CF 1547
 Kubo, M 915
 Kubota, N 847,903
 Kubota, S 1719
 Kuča, K 1541
 Kučera, R 781
 Kuess, C 1137
 Kukovinets, OS 293
 Kulesh, NI 587
 Kulevanova, S 639
 Kulhánková, A 1541
 Kumar, D 161,195
 Kumar, N 161
 Kumar, S 497
 Kumaravel, M 183
 Kumazawa, S 1719
 Kuneš, J 441,1541
 Kuntiya, A 1381
 Kuo, CC 805,827
 Kuo, YH 79,805,827
 Kuppan, N 493
 Kuroda, C 631
 Kurosawa, K 815
 Kuš, PM 651
 Kuster, RM 207
 Kutil, Z 105
 Kuvatanasuchati, J 613
 Kwak, MK 431
 Kwak, WJ 1609
 Kwon, DY 609,1233
 Kwon, JT 1301,1609
- Labraimi, H 55
 Lacaille-Dubois, MA 157,573
 Lackey, BR 257
 Lago, JHG 277,679
 Lahlou, S 1167
 Lai, YC 363
 Lajis, NH 447
 Lakshma Nayak, V 1741
 Lakušić, B 131
 Lakušić, D 131
 Lal, B 161
 Lambert, LK 725
 Lamia, H 407
 Lan, WJ 67
 Landa, P 105
 Landaeta, C 1679
- Landeira-Fernandez, AM 207
 Langat, MK 175
 Lapcik, O 1423
 Laphookhieo, S 799
 Larsen, L 689
 Latha, P 183
 Lau, D 1617
 Laudato, M 1333
 Laudato, M 419
 Lawrence MP 367
 Lax, V 817
 Leal, ICR 207
 Lebeuf, R 965
 Lee, C 1233,1571
 Lee, CH 571
 Lee, CK 827
 Lee, DH 1301
 Lee, FP 621
 Lee, GS 1763
 Lee, HD 181
 Lee, HK 1605
 Lee, HS 1229,1399
 Lee, JJ 1553
 Lee, JJ 431
 Lee, KG 823
 Lee, M 1301
 Lee, MK 481,1233,1439
 Lee, MR 1605
 Lee, MY 609
 Lee, NH 427,1359
 Lee, NL 1363
 Lee, S 1609
 Lee, SH 181,1439,1739
 Lee, SS 363
 Lee, SW 609
 Lee, TH 827
 Lee, WJ 1359
 Leeanansaksiri, W 629
 Legault, J 265
 Legoale, PB 1403
 Legotkina, G 617
 Lei, KY 213
 Leite, FHA 1049
 Leite, TCC 1049
 Leng, J 1115
 León, LG 187
 León-González, AJ 1255
 Lerma-García, MJ 1325
 Leroch, M 1599
 Leuner, O 1423
 León, F 1285
 Li, D 1321,1397
 Li, F 75,463,1321
 Li, GQ 599,707
 Li, H 1479,1673
 Li, HJ 67
 Li, J 479,1321,1713
 Li, KM 707
 Li, N 1121
 Li, S 155,753,1227
 Li, X 587,787
 Li, XN 467
 Li, Y 1129,1427
 Li, YC 805
 Li, YK 467
 Li, YQ 1197
 Li, Z 1129,1397
 Li, ZH 487
 Liang, KJ 1535
 Liang, WL 67
 Liang, X 1101
 Liao, SG 1373
 Liao, ST 599
 Liao, YW 79
- Lie, W 695
 Likhitwitayawuid, K 115,613
 Likhitwitayawuid, K 1079
 Lim, HS 1443
 Lima-Junior, RS 1547
 Lin, CH 621
 Lin, HF 1363
 Lin, HW 1713
 Lin, JJ 827
 Lin, PC 621
 Lin, T 113
 Lin, YS 79
 Lindequist, U 1223
 Lipipun, V 1079
 Liu, B 1373
 Liu, C 1321
 Liu, F 911
 Liu, H 457
 Liu, LS 113
 Liu, Q 481,1439
 Liu, RH 113
 Liu, T 1353
 Liu, W 67
 Liu, Y 83,587,627
 Liu, YQ 199
 Li-Xia, C 149
 Lo, LC 827
 Ločárek, M 781
 Lohani, M 1415
 Long, C 1519
 Long, LHV 1507
 López, LC 47
 López-Lázaro, M 1255
 Lorán, S 817
 Lourith, N 1583
 Lower-Nedza, AD 523,1137
 Lu, MC 571,1535
 Lu, W 479
 Lucas, FCA 1471
 Lucchese, AM 269
 Ludwiczuk, A 1515
 Luis, Á 1459
 Luna-Sánchez, M 47
 Luo, HR 1373
 Luo, NC 487
 Luo, Y 337,457
 Luo, YP 1427
 Lupidi, G 1791
 Lv, HN 1473
 Lv, JL 335
 Lv, X 479
 Lyapunov, J 617
- Ma, G 1117
 Macáková, K 1541
 Machumi, F 311,761
 Macias, FA 1687
 Macková, A 1509
 Macreadie, IG 1763
 Magalhães, PJC 1167
 Maggi, F 1791
 Maggio, A 283,841,1159,1629
 Mahajan, G 1455
 Maharjan, S 1481
 Mahdavi, M 1811
 Mai, W 787
 Maia, JGS 1471
 Maione, F 539
 Maity, N 671
 Makabe, H 1005
 Makarieva, TN 1229,1399
 Makino, K 915
 Makino, M 1665
 Malafronte, N 1387
- Malak, LG 191
 Malek, SNA 1763
 Maleš, Z 231,347
 Malik, J 105
 Mallavarapu, GR 1771
 Mallula, GR 1595
 Malova, H 139
 Malyarenko, TV 1219
 Mamadaliyeva, NZ 429
 Mamiya, N 925
 Mancini, E 139,407,1637
 Mandal, SC 1487
 Mangmool, S 1257
 Manzoor, N 261
 Marandino, A 407
 Marasović, M 231
 Marcourt, L 1519
 Marie, DEP 635
 Marijanović, Z 231,651
 Marin, PD 1313
 Markosyan, A 1523
 Marković, MS 1309
 Márquez, O 187
 Marrelli, M 567
 Marsik, P 105
 Martin, AS 397
 Martín-Cordero, C 1255
 Martinelli, C 1645
 Martínez, M 187
 Martino, LD 139,407,1637
 Martin-Quintal, Z 1567
 Martins, DTO 1567
 Martyyas, EA 301,1053
 Mascolo, N 539
 Mashimbye, MJ 1403
 Mašković, P 667
 Massoni, M 679
 Mastrogiovanni, F 429
 Masuo, Y 515
 Máthé, I 1153
 Mathew, AE 811
 Matić, IZ 1291
 Matsui, T 1265
 Matsunami, K 767
 Matsuo, AL 277,679
 Matsuura, H 815
 Matsuzaki, K 1665
 Mazurok, TI 1419
 McDougal, OM 1059
 Meccia, G 1641
 Medinilla, M 1325
 Medrano-Nahuat, D 1245
 Mehta, GK 497
 Mei, W 337
 Meira, CS 403
 Mekinić, IG 471
 Meliani, N 835
 Melim, C 1567
 Mello, CS 1547
 Menchinskaya, ES 1377
 Menezes, LRA 403
 Menichini, F 567
 Menut, C 829
 Méryllon, JM 485
 Michalik, M 1691
 Midiwo, JO 175,761
 Miftakhov, MS 981
 Mihajilov-Krstev, TM 1309
 Miki, Y 1553
 Miksatkova, P 1423
 Miladinović, DL 1309
 Miladinović, LC 1309
 Mildner, S 1651
 Milić, N 1801

- Miloš, M..... 231
 Milosavljević, S..... 1291
 Milošević, N..... 1801
 Mimaki, Y..... 315
 Minai-Tehrani, A..... 1609
 Ming, LJ..... 415
 Minh, CV..... 1209
 Minh, CV..... 1751
 Mischenko, NP..... 587
 Misra, BB..... 253
 Mitaine-Offer, AC..... 157,573
 Mitula, P..... 435
 Miyake, S..... 1553
 Miyamoto, T..... 157
 Miyamoto, T..... 573
 Mizutani, R..... 883,949
 Mizutani, Y..... 919
 Mohamad, R..... 1811
 Mohamed, AA..... 1365
 Mohammad, FV..... 1289
 Mohammad, AI..... 1117
 Mohammed, FAE..... 21
 Mohammed, R..... 311
 Moharram, AM..... 191
 Mohsen, H..... 407
 Moiteiro, C..... 1785
 Monache, FD..... 1567
 Mondello, L..... 171
 Monfalouti, HE..... 25,33,55
 Moniuszko-Szajwaj, B..... 1687
 Montaut, S..... 1463
 Monteagudo, U..... 1325
 Moongkarndi, P..... 1257
 Moo-Puc, R..... 1421
 Morel, N..... 575
 Moriarity, DM..... 1179
 Morikawa, T..... 1553
 Morimitsu, T..... 883,949
 Mosharova, IV..... 1229
 Motilva, V..... 1711
 Moujir, LM..... 1175
 Mu, Q..... 1353
 Mučaji, P..... 1509
 Muhammad, A..... 1435
 Muhammad, I..... 311,761
 Mukherjee, PK..... 671
 Müller, ID..... 645
 Mungmee, C..... 115
 Munigunti, R..... 1135
 Munikishore, R..... 459
 Muñoz, MA..... 1075
 Munshi, A..... 213
 Murai, Y..... 453,1251
 Muraoka, O..... 1553
 Murata, T..... 775
 Murata, Y..... 889
 Murphy, C..... 257
 Murugan, R..... 1771
 Murunga, S..... 1261
 Musciacchio, G..... 539
 Muselli, A..... 835
 Muthna, D..... 1797

 Na, MK..... 181
 Na, YJ..... 1605
 Na, Z..... 1373
 Nadir, M..... 1633
 Nagasawa, K..... 961
 Naidu, M..... 711
 Naik, V..... 1455
 Nair, JJ..... 553,1335,1705
 Nair, MG..... 165,1365
 Najafi, TF..... 1811
 Nakada, M..... 955
 Nakanishi, M..... 1719
 Nakashima, K..... 877,883,949
 Nakayama, T..... 119
 Nam, NH..... 1751
 Nam, SY..... 217
 Nammi, S..... 707
 Namvar, F..... 1811
 Narala, VR..... 121
 Narender, T..... 803
 Naresh, G..... 803
 Narkowicz, C..... 211
 Nascimento, KS..... 1621
 Nazyrov, TI..... 293
 Ndhlala, AR..... 795
 Nema, NK..... 671
 Nga, VT..... 367,1507
 Ngane, RAN..... 829
 Nguikwie, SK..... 829
 Nhiem, NX..... 1209,1751
 Nicola, GRD..... 1463
 Nie, Y..... 1101
 Niero, R..... 1567
 Niezgodna, N..... 435
 Nikolić, DM..... 1309
 Nikolov, S..... 765
 Ning, Z..... 339
 Ninomiya, K..... 1553
 Nkengfack, AE..... 103,1575
 Noda, E..... 1265
 Noda, Y..... 925
 Nogueira, PCL..... 403
 Nokami, J..... 919
 Noll, B..... 299
 Noma, Y..... 859
 Nong, XH..... 1069
 Norte, M..... 187
 Novák, I..... 605
 Novák, Z..... 441,781,1541
 Nsikabaka, S..... 655
 Nugraha, AS..... 703
 Nyegue, MA..... 829

 Ocazionez, RE..... 249
 Ogunmoye, AO..... 243
 Ogunsanwo, AO..... 243
 Ogunwande, IA..... 239,243
 Ogurtsova, EK..... 1229
 Oh, TH..... 427
 Ohta, T..... 1719
 Okada, R..... 1367
 Okada, S..... 629
 Okada, Y..... 1409,1721
 Okamoto, Y..... 631
 Okamura, N..... 1719
 Okoso-amaa, EM..... 85
 Okugawa, S..... 1553
 Okusa, PN..... 439
 Olejnik, K..... 227,1297
 Olennikov, DN..... 585
 Oleszek, W..... 1687
 Oliveira, DF..... 69,424
 Oliveira, R..... 673
 Omatsu, R..... 919
 Onur, MA..... 327
 Opletal, L..... 441,781,1541
 Osako, K..... 533
 Otsuka, H..... 767
 Ouattara, ZA..... 1773
 Owczarek, A..... 505
 Oza, MD..... 497
 Ozaki, S..... 119
 Özhan, O..... 475
 Özşen, Ö..... 59

 Padmakumar, KP..... 301
 Padrón, JM..... 187
 Padwad, YS..... 161
 Pak, SC..... 389
 Paknikar, SK..... 1195,1513
 Pal, A..... 1591
 Pamela, D..... 1763
 Panovska, TK..... 639
 Paolini, J..... 655
 Papa, F..... 1791
 Papp, N..... 1247
 Park, DW..... 203
 Park, HS..... 1579
 Park, HY..... 431
 Park, NH..... 1605
 Park, SJ..... 1609,1751
 Park, SU..... 217,609,1233,1571
 Park, WT..... 217
 Park, Y..... 1149
 Partilkhaev, VV..... 585
 Patel, AV..... 63
 Patel, P..... 1467
 Patil, U..... 1467
 Pattanapanyasat, K..... 1257
 Paudel, P..... 1179,1481,1777
 Paul, VJ..... 1537
 Pavasant, P..... 613
 Payo-Hill, AL..... 779
 Paz, C..... 147
 Peakall, R..... 701
 Pecio, L..... 1687
 Peña, IC..... 1143
 Peng, J..... 329,1069
 Peng, T..... 1393,1753
 Peng, ZQ..... 199
 Peraza-Sánchez, SR..... 1245,1421
 Perez, C..... 397
 Pérez-López, LA..... 135
 Perry, NB..... 689
 Pertuit, D..... 157
 Pescitelli, L..... 419
 Peter, M..... 175
 Petrelli, D..... 1791
 Petrović, G..... 109
 Phakhodee, W..... 799
 Phan, CW..... 1763
 Phillips, GJ..... 911
 Phillips, RD..... 701
 Phoolcharoen, W..... 125,613
 Phung, NKP..... 367,1507
 Phuwapraisirisan, P..... 1085
 Piaz, FD..... 1387
 Piccinelli, AL..... 1237
 Pilepić, KH..... 231,347
 Piližota, V..... 1767
 Pino, JA..... 129
 Pinto, NV..... 1621
 Pintye-Hódi, K..... 1153
 Piovan, A..... 1615
 Pires, AF..... 1621
 Plotkin, BJ..... 1617
 Plubrukarn, A..... 1355
 Pluhár, Z..... 605
 Podolak, I..... 1691
 Pokharel, KK..... 1777
 Polain de Waroux, Y..... 1
 Pongparn, P..... 799
 Popov, SA..... 1759
 Popova, M..... 617
 Poudel, A..... 1179,1777
 Pour, M..... 441
 Požrl, T..... 1767
 Pragadheesh, VS..... 221
 Prakash, I..... 1523

 Prata, APN..... 403
 Prevost, S..... 965
 Pribylova, M..... 105
 Prinsep, MR..... 693
 Proksch, P..... 289
 Putalun, W..... 443
 Putschakarn, S..... 1355
 Pyne, S..... 281
 Pyne, SG..... 695,733,1305
 Pyo, KH..... 1755

 Qazimi, B..... 639
 Qi, SH..... 329,1069,1127
 Qian, DW..... 487
 Qian, PY..... 329
 Qian, YF..... 487
 Qian, Z..... 149
 Qiao, B..... 1129
 Qin, K..... 1447,1479
 Qiu, SX..... 1121,1733
 Qu, L..... 1321
 Quang, NV..... 1209,1751
 Quang, TT..... 1507
 Quang, TT..... 367
 Quassinti, L..... 1791
 Quetin-Leclercq, J..... 575
 Quijano-Célis, CE..... 129
 Quintero, P..... 1641

 Rabehaja, DJR..... 1165
 Radchenko, OS..... 1219
 Radojković, M..... 667
 Radwan, MM..... 1285
 Rafamantanana, MH..... 575
 Raharisololalao, A..... 1099
 Rahman, HS..... 1811
 Raimondo, FM..... 283
 Rajan, GC..... 301
 Rajemiarimiraho, M..... 1099
 Rajendran, SM..... 803
 Rak, G..... 605
 Rakotobe, E..... 1201
 Rakotonandrasana, SR..... 1099
 Rakotondrajaona, R..... 1201
 Ramakrishna, S..... 1741
 Ramalho, L..... 1785
 Ramanjooloo, A..... 635
 Ramanoelina, PAR..... 1165
 Ramasamy, D..... 433
 Ramírez-González, I..... 1731
 Ramis-Ramos, G..... 1325
 Ramli, RA..... 695
 Randrianaivo, R..... 1201
 Randrianja, R..... 1099
 Randriantsoa, A..... 575
 Ranji, PKV..... 1431
 Rao, GV..... 1513
 Raelison, GE..... 575
 Rapior, S..... 509
 Rárová, L..... 1705
 Rasamison, VE..... 1201
 Rasheed, M..... 1633
 Rashid, MA..... 757
 Rasoanaivo, LH..... 1099
 Rasool, N..... 757
 Rastrelli, L..... 779,1237
 Ratsimamanga-Urverg, S..... 1165
 Rattanajak, R..... 733
 Ravada, SR..... 1595
 Razafindrazaka, R..... 575
 Razmovski-Naumovski, V..... 707
 Razukas, A..... 1105
 Ree, TV..... 1403
 Ren, G..... 113

- Rezacova, M..... 1797
 Rezende, CM..... 424
 Rezende, CM..... 69
 Rezgui, A..... 157
 Rhouati, S..... 439
 Ribeiro, AF..... 1471
 Ricci, D..... 527,1587,1645
 Rigano, D..... 567,841
 Riina, R..... 1679
 Riley, S..... 911
 Río, RE..... 1711
 Ríos LD..... 1697
 Rios, D..... 397
 Risher, MJ..... 85
 Ristić, M..... 131,667
 Rivas-Estilla, AM..... 297
 Rivas-Galindo, VM..... 135,297
 Rivière, C..... 485
 Rizvi, MMA..... 261
 Rocha, TSM..... 207
 Rodas, FR..... 453
 Rodpai, E..... 1257
 Rodríguez, EJ..... 1325
 Rodríguez-García, G..... 1711
 Rogers, LB..... 85
 Rojas, J..... 1175
 Rojas, L..... 1175
 Rojas, LB..... 1641
 Rojas, R..... 1785
 Rojas-Fermin, LB..... 1477
 Rollin, P..... 1463
 Romanenko, LA..... 381
 Romero, LI..... 1537
 Romestand, B..... 829
 Rondón, M..... 1175
 Rosa, LH..... 1285
 Ross, SA..... 191,1117
 Rosselli, S..... 283,1159
 Rota, MC..... 817
 Rotjanapan, K..... 629
 Rouaiguia-Bouakkaz, S..... 485
 Rouis, Z..... 1629
 Row, KH..... 203
 Roy, SK..... 1241
 Ruanghirun, T..... 115
 Ruangrunsi, N..... 1265
 Ruela, HS..... 207
 Ruiz, Y..... 129
 Rukkumani, R..... 183
 Russo, M..... 171

 Sabaratnam, V..... 711,1763
 Sabino, KCC..... 207
 Sadgrove, NJ..... 737,741,747
 Sáez C, K..... 1697
 Saha, BP..... 671
 Sahu, R..... 1275
 Saidi, M..... 1171
 Saile, R..... 51
 Saito, Y..... 631
 Saito, Y..... 883
 Sajeva, M..... 1159
 Sakaguchi, K..... 1093,1563
 Sakee, U..... 385
 Salazar-Aranda, R..... 135
 Salm, F..... 1599
 Salnikova, OI..... 1759
 Salvini, L..... 429
 Samer, J..... 1257
 Samoylenko, V..... 311
 Sampaio, MFC..... 403
 Samten, J..... 733
 Sanandiyi, ND..... 497
 Sánchez O, M..... 1697
 Sanchez-Olate, M..... 397
 Sándor, V..... 1247
 Sandoval-López, DS..... 779
 Sandun, STP..... 333
 Sankar, P..... 183
 Sansom, CE..... 689
 Santos, CF..... 1621
 Saovapakhiran, A..... 799
 Saraswathy, A..... 433
 Şarer, E..... 475
 Sarikaya, BB..... 327
 Šarolić, M..... 651
 Šárosi, S..... 605
 Sartorelli, P..... 277
 Sartori, C..... 579
 Sasaki, J..... 851
 Sasaki, T..... 1735
 Satılmış, B..... 475
 Sato, K..... 987
 Sato, S..... 873
 Satoh, M..... 1665
 Satoh, Y..... 1665
 Satou, T..... 515
 Satyal, P..... 1179,1481,1777
 Saucedo-Hernández, Y..... 1325
 Savchenko, AM..... 1219
 Scandolera, E..... 407
 Scarano, MT..... 1387
 Schlapkohl, A..... 911
 Schmidt, E..... 663
 Schmitz, R..... 289
 Schuster, D..... 105
 Schutzki, RE..... 165
 Schwartz, D..... 1415
 Scirica, E..... 1159
 Sekiguchi, A..... 315
 Seletskaia, LD..... 1419
 Sen, A..... 1269
 Senatore, F..... 567,841,1159,1629
 Sengoku, T..... 889,1011
 Seo, CS..... 1443
 Seo, EK..... 1143
 Seo, GB..... 1301
 Seo, J..... 823
 Seo, JK..... 1301
 Seo, Y..... 431,1405
 Serra, S..... 863
 Sestili, P..... 1645
 Sette, LD..... 721
 Setzer, WN..... 683,1179,1481,1777
 Sha, L..... 461
 Shaari, K..... 447
 Shafi, G..... 213
 Shah, AJ..... 1633
 Shahabuddin, SK..... 459
 Shang, SZ..... 467
 Shankaraiah, N..... 1741
 Shannon, SD..... 1415
 Shao, F..... 113
 Sharma, SN..... 1045
 Sharma, U..... 161
 She, G..... 461
 Sheehy, E..... 1123
 Shekhar, R..... 195
 Shi, MF..... 597
 Shi, R..... 461
 Shibuya, F..... 1735
 Shih, WL..... 79
 Shim, I..... 1301
 Shim, SM..... 1579
 Shim, WS..... 1143
 Shim, CK..... 1143
 Shimada, K..... 851
 Shimizu, K..... 897
 Shimoda, K..... 847,903,905,907
 Shin, EH..... 1755
 Shin, HK..... 1443
 Shin, JY..... 1609
 Shin, KS..... 1755
 Shiono, Y..... 1735
 Shiozawa, T..... 877
 Shirataki, Y..... 1409
 Shi-Yu, Y..... 149
 Shmidt, EN..... 1759
 Shoji, M..... 897
 Shou, Q..... 1115
 Shpatov, AV..... 1759
 Shu, Y..... 1479
 Shushni, MAM..... 1223
 Siani, AC..... 1547
 Sichaem, J..... 1111
 Sichaem, J..... 1371
 Siddhanta, AK..... 497
 Silchenko, AS..... 301,1053,1377
 Silchenko, AS..... 1527
 Silou, T..... 655
 Silva, AN..... 269
 Silva, M..... 147
 Silva, TB..... 403
 Silva-Mares, D..... 297
 Simčić, M..... 1767
 Simó-Alfonso, EF..... 1325
 Simonet, AM..... 1687
 Singh, B..... 161
 Singh, M..... 221,261
 Singtothong, C..... 265
 Sinha, RK..... 1045
 Siqueira, RJB..... 1167
 Sirat, HM..... 513,1317,1435
 Siripong, P..... 1111,1371
 Sisodia, BS..... 1591
 Skroza, D..... 471
 Slavkovska, V..... 131
 Slobodniková, L..... 1747
 Šmelcerović, A..... 109
 Smillie, TJ..... 319,323
 Smirnova, IE..... 293
 Smith, CT..... 299
 Soares, MBP..... 403
 Soares, MG..... 679
 Solujić, S..... 667
 Somer, NU..... 327
 Son, KH..... 1609
 Song, Q..... 1115
 Song, XP..... 1427
 Soompon, S..... 613
 Sookkhee, S..... 519
 Soscún, H..... 1679
 Sossa, K..... 147
 Sotomayor, JA..... 817
 Spadaro, V..... 283
 Spengler-Salabarría, IS..... 779
 Squitieri, E..... 1679
 Sravani, S..... 1741
 Sreedhar, E..... 1741
 Srichana, T..... 501
 Sritularak, B..... 115,613,1079
 Srivastava, A..... 1241
 Srivastava, SK..... 1591
 Stabentheiner, E..... 645
 Staden, JV..... 553,795
 Stafilov, T..... 639
 Stanković, M..... 109
 Stanslas, J..... 447
 Stashenko, E..... 249
 Steenkamp, P..... 1213
 Steenkamp, V..... 1213,1279
 Stefkov, G..... 639
 Stévigny, C..... 439
 Stochmal, A..... 1687
 Stojanović, GS..... 109
 Stojanović, IŽ..... 109
 Stonik, VA..... 1219,1229,1377
 Stonik, VA..... 1527
 Strnad, M..... 1705
 Strnad, M..... 771
 Su, X..... 1129
 Su, HJ..... 1363
 Su, JH..... 571,1363,1535
 Su, TR..... 1535
 Su, W..... 457
 Su, YC..... 143,273,531,1329
 Su, Z..... 1227
 Suárez, AI..... 1679
 Subramani, PA..... 121
 Sudha, V..... 1771
 Sudo, T..... 889
 Sugai, T..... 897
 Sugamoto, M..... 903
 Sugandhika, ST..... 333
 Sugiyama, Y..... 1719
 Sukapirom, K..... 1257
 Sukkharak, P..... 565,1515
 Sukrong, S..... 125
 Suksathan, R..... 519
 Sumner, S..... 63
 Sun, HD..... 467
 Sun, J..... 627
 Sung, JS..... 823
 Sung, PJ..... 1037
 Suponitskii, KY..... 293
 Suppiramaniam, V..... 1415
 Supratman, U..... 1735
 Suresh Babu, K..... 1741
 Suvajdić, L..... 1801
 Suwada, K..... 847
 Suzu, I..... 629
 Suzuki, N..... 153,775
 Suzuki, R..... 1409
 Suzuki, T..... 873,929
 Syed, N..... 213
 Szeto, YT..... 389
 Szewczyk, K..... 545
 Szilvássy, B..... 605
 Szumacher-Strabel, M..... 1687
 Szymanski, PT..... 153

 Tabakmakher, KM..... 1229,1399
 Tabanca, N..... 257
 Tabowei, GO..... 243
 Tabti, B..... 835
 Taddei, AR..... 429
 Tadtong, S..... 1583
 Tagma, T..... 5,11
 Tahir, PM..... 1811
 Tai, BH..... 1209,1751
 Takahashi, K..... 815
 Takahashi, M..... 889,1011
 Takahashi, R..... 453
 Takaoka, S..... 877
 Takehara, T..... 877
 Takemoto, M..... 907
 Takikawa, Y..... 851
 Tamuly, C..... 1451
 Tamura, MN..... 1251
 Tanaka, C..... 157,573
 Tanaka, K..... 75
 Tanaka, R..... 1367
 Tang, J..... 1545
 Tang, W..... 479
 Tanimoto, H..... 1021
 Tankhaeva, LM..... 585

- Tantrakarnsakul, K. 1079
 Tapondjou, LA 1791
 Tarawneh, AH 1285
 Tarbeeva, DV 1419
 Tareen, RB 757,1633
 Tarkovská, D 771
 Tatsuzawa, F 1095
 Taulavuori, E 791
 Taulavuori, K 791
 Tava, A 1625
 Taweechoitipatr, M 733,1395
 Tawinkan, N 443
 Teixeira, VL 533,1669
 Tekeľová, D 1153
 Tekwani, BL 761
 Telford, IRH 737
 Temml, V 105
 Terzioglu, G 1269
 Tewtrakul, S 1205
 Tezuka, Y 75,463
 Thai, TH 235
 Thanakosai, W 1085
 Thang, TD 239,663
 Thao, NP 1751
 Thompson, HL 63
 Thu, NTH 367
 Thung, DC 1209
 Thwe, AA 1571
 Tiamiyu, ZA 243
 Tian, M 203
 Tian, X 199
 Tiani, GM 103
 Tian-Xian, L 149
 Tidjani, S 439
 Tiezzi, A 429
 Timmers, M 715
 Tip-pyang, S 1111,1371
 Togna, GD 1537
 Tokuda, H 153
 Toloue, M 359,807
 Tolstikov, GA 293
 Tomi, F 235,1155,1165,1773
 Tomi, P 655
 Tomita, M 897
 Tommasi, ND 1387
 Tomsik, P 1797
 Tori, M 631,877,883,949
 Toro, M 1641
 Torre, G 171
 Torres, NW 135
 Torres-Cirio, A 135
 Torres-López, E 297
 Torres-Tapia, LW 1421
 Torres-Valencia, JM 1711
 Tóth, J 1153
 Towiwat, P 501
 Toyota, M 999
 Trachoo, N 385
 Tretyakova, EV 293
 Trigos, Á 187
 Trimurtulu, G 459
 Troncoso, C 397
 Trusheva, B 617
 Tschiggerl, C 523
 Tu, PF 1473,1575
 Tu, X 83,587
 Tuan, NA 235
 Tubek, B 435
 Tuberoso, CIG 651
 Tůmová, L 593
 Turiel, NA 1471
 Tuyen, PNK 367
 Tyagi, AK 1313
 Uddin, MR 217,609,1233
 Uesawa, Y 1721
 Uetanabaro, APT 269
 Um, BH 1759
 Upadhyay, HC 1591
 Urban, S 715,729
 Urrutia, H 147
 Urverg-Ratsimamanga, S 575
 Usubillaga, A 1477,1641
 Utkina, NK 1551
 Uwamori, M 955
 Vairappan, CS 287
 Vajs, V 1291
 Valant-Vetschera, KM 1081
 Valente, LMM 1547
 Valera-Burgos, J 11
 Valeri, M 429
 Valterova, I 1423
 Valverde, J 1123
 van Staden, J 1335,1705
 van Tonder, JJ 1279
 Vanek, T 105
 vanSoest, RWM 635
 Velasco, J 1641
 Velásquez, J 1641
 Vélez-Castro, HT 779
 Veljic, MM 1313
 Venditti, A 1629
 Venegas, C 47
 Veselova, MV 587,1419
 Vibin, M 1609
 Vieira, IJC 1049
 Viladomat, F 553,1335
 Vildová A 593
 Villaça, RC 1669
 Viswanathan, UM 1599
 Vitali, LA 1791
 Vittori, S 1791
 Vladimirova, A 85
 Vogler, B 1179
 Vongsak, B 1559
 Vrkoslav, V 1423
 Waddell, TG 299
 Waffo-Téguo, P 485
 Wajs-Bonikowska, A 227,1297
 Waksman-Minsky, N 297
 Walker, LA 761
 Wanas, AS 767
 Wang, EIC 1329
 Wang, L 1701
 Wang, WH 1037
 Wang, C 339
 Wang, CJ 1353
 Wang, CY 1675
 Wang, EIC 531
 Wang, F 1101
 Wang, FP 1701
 Wang, H 337
 Wang, J 1101,1427
 Wang, L 787
 Wang, LY 67
 Wang, P 1227,1397
 Wang, RR 467
 Wang, SY 805
 Wang, TS 1427
 Wang, WJ 599
 Wang, Y 457,599
 Wang, Z 393,523,1545
 Wangchuk, P 733,1305
 Wanigsekara, WMAP 601
 Watanabe, K 919
 Watanabe, S 75
 Waters, NC 175
 Wätjen, W 289
 Wattanapiromsakul, C 1205
 Wawrzeńczyk, C 435
 Wedge, DE 257
 Weems, M 911
 Wei, MC 1683
 Wei, Q 339
 Wei, X 169
 Wekesa, I 1261
 Wesolowski, M 1821
 Wessjohann, L 683
 Whaley, WL 85
 Wiart, C 375
 Widowitz, U 105,1509
 Wierzejska, J 1011
 Wietrzyk, J 435
 Wijeratne, EMK 601
 Wijeyaratne, SC 1431
 Woguem, V 1791
 Wolbiś, M 99
 Womack, CL 85
 Womeni, HM 1791
 Wong, KH 711
 Wongkrajang, Y 1559
 Woźniak, M 545
 Wright, D 63
 Wu, DL 599
 Wu, H 393
 Wu, J 487,1733
 Wu, JH 805
 Wu, L 1479
 Wu, TS 775
 Wu, W 753
 Wu, X 1397
 Wu, YC 1037
 Wu, YJ 1535
 Xiang, Q 155,1321
 Xiao, G 1733
 Xiao, WL 467
 Xie, HQ 1353
 Xie, L 1397
 Xie, ZL 67
 Xiong, J 155
 Xu, D 1101
 Xu, KX 1675
 Xu, L 169,1393,1479,1753
 Xu, XY 1069,1127
 Xu, Y 461,1321,1733
 Xu, YK 1373
 Xu, YM 1197
 Xu, Z 1121
 Xue, J 169
 Yadav, A 221
 Yahara, S 583,1413
 Yamada, H 869
 Yamada, T 1367
 Yamakawa, T 1367
 Yamamoto, H 1719
 Yamamoto, K 463
 Yan, W 811
 Yang, F 1713,1753
 Yang, GY 467
 Yang, J 1121
 Yang, L 199,1373
 Yang, LL 621
 Yang, LM 467
 Yang, S 753
 Yang, W 1115,1321
 Yang, X 1753
 Yang, Y 655,1393,1753
 Yang, YC 1683
 Yap, KC 1725
 Yasuda, H 919
 Yasuda, Y 1735
 Yasui, N 627
 Ye, H 1101
 Ye, WC 599
 Ye, YH 113
 Yegdaneh, A 1355
 Yen, PH 1209,1751
 Yenesew, A 1261
 Yeo, SK 1571
 Yin, ACY 415
 Yingmao, D 523
 Yoda, H 889,1011
 Yokosuka, A 315
 Yomoda, S 463
 Yoon, HS 1359
 Yoon, WJ 1359
 Yoshikawa, M 515,1553
 Yoshimura, J 987
 Yotova, M 765
 You, Q 1101
 Yousfi, M 1171
 Youssef, DTA 289
 Yu, KN 1609
 Yu, Y 1733
 Yuan, E 339
 Yuan, W 1227
 Yuan, Y 1129
 Yuenyongsawad, S 1355
 Yurchenko, EA 1071
 Yurchenko, EA 1527
 Yusuf, KO 243
 Zacchino, S 1785
 Zafaryab, M 261
 Zahoor, A 1289
 Zajdel, P 1691
 Zaki, MA 311
 Zamyatina, SV 587
 Žarkov, M 1801
 Zdraveva, P 765
 Zellagui, A 439
 Zeppa, G 33
 Zhabinskii, VN 771
 Zhang, X 1447
 Zhang, C 1121
 Zhang, CR 165,1365
 Zhang, CY 1217
 Zhang, H 335,479,1129
 Zhang, J 199,753,761,1393
 Zhang, L 83,461,587
 Zhang, LB 335
 Zhang, M 1447
 Zhang, MQ 1675
 Zhang, T 1197
 Zhang, X 393,1129
 Zhang, XQ 599
 Zhang, XT 599
 Zhang, XY 1069,1127
 Zhang, Y 155,479
 Zhang, ZW 199
 Zhao, H 1137
 Zhao, L 1393,1753
 Zheng, L 1447
 Zheng, YT 467
 Zhong, MG 1427
 Zhong, Z 149
 Zhou, CX 597
 Zhou, H 1393,1753
 Zhou, J 1675
 Zhu, J 523
 Zhu, Y 1121
 Zhuravleva, OI 1071

Zito, P 1159
Zlatić, E 1767
Żmudzki, P 1691

Zollo, PHA 829
Zottele, CW 1387
Zou, K 83,587

Zou, W 457
Zubair, M 757
Zunzunegui, M 5,11,15

Zupko, I 343
Zverev, YF 587

Natural Product Communications

Volume 8 (1-12)

2013

Keywords Index

- Abelmoschus esculentus* 1085
Abies koreana 227,1297
 Abietane diterpenoids 293
 Abietatriene 1771
 Abscissic acid 1673
 Absolute configuration 1123,1675, 1711
 ABTS 475,1413,1551
Acalypha wilkesiana var. *macafeana* hort. 375
 Acanthaceae 333,1163
 Acanthoic acid 1359
Acanthopanax koreanum 1359
 ACE inhibitor 671
 Acetonylnipholone
 cyclooxanthrone 1261
 Acetylcholinesterase 695,781,795, 1541
 Acetylcholinesterase inhibition 471
 Acetylcholinesterase inhibitor 447
 Acetylcholineformine 1545
 Acetylthiamopyranosylisovitexin 1413
 AChE/BchE 1373
Achillea ligustica 1629
 Aciduminate 1111
Acinetobacter baumannii 673
 Aconitamide 827
Aconitum carmichaeli 155,827
Acorus calamus 1179
Acremonium sp. 601
Actinococcus typical 301
 Acylated anthocyanins 1563
 Adipocyte differentiation 1439,1579
 Adventitious roots 1045
Aedes aegypti 257
 Aeschynocarpin 1421
Aeschynomene fascicularis 1421
Aframomum dalzielii 829
Aframomum lesteuianum 829
Aframomum pruinosum 829
 Agaricaceae 1227
 Agavaceae 315
Agelas clathrodes 1713
Aglaia odorata 1373
Agrobacterium rhizogenes 217
Ajania nubigena 733
 Aldehyde 393
 Aldol reaction 877
 Alkaloids 1021,1335,1393,1397, 1427,1705
 Alkaloids 165,385,441,539,553,693,
 Alkene 631
 Alkyl α,β -lactones 447
 Allelochemical 1197
 Allelopathy 1179,1481,1777
 Aloe extract 1333
Aloe ferox 411
 Aloe gel 1333
 Aloe species 1333
Alpania galangal 1741
Alpinia rafflesiana 1317
Alternaria alternata 59
 Alum 419
 Alzheimer's disease 781
 Amaryllidaceae 327,553,1231, 1335,1541,1705
 Amaryllidaceae alkaloids 161
 Amazonian yam bean 1423
 Amide glycoside 169
 Aminomethyl- γ -butyrolactone 889
 Aminopalladation 1005
Amomum biflorum 265
Amomum tsao-ko 1715
 Amylase 339,1279
Anaphalis contorta 225
 Androgens 51
 Andrographidine G 333
Andrographis paniculata 333,1045, 1725
 Andrographolide 1045
Anemarrhena asphodeloides 481
 Anethole 671,1165
 Angiogenesis 1719
 Angiogenesis *in vitro* 1291
Annona squamosa 385
 Annonaceae 663
 Anoikis 115
Anthocleista djalensis 243
 Anthraquinones 191,1117
 Anti-allergic inflammatory 181
 Anti-allergic inflammatory activities 1739
 Anti-bacteria 613
 Antibacterial 265,385,823,1175, 1223,1305
 Antibacterial activity 269,287,337, 519,829, 1309,1363,1459,1567, 1747
 Antibacterial property 429
 Anti-biofilm activity 673
 Antibiotic 373,493
 Antibiotics 911
 Anticancer 213,553,703,1811
 Anticancer activity 83,273,287, 359,381,531,807,1427
 Anticandidal 261
 Anticholinesterase 253,327,1381
Antidesma acidum 1111
 Anti-diabetic 415
 Antifertility 1183
 Antifilarial activity 803
 Antifouling 329
 Antifungal 257,741,1179,1451,1763
 Antifungal activity 1587,1599,1691
 Anti-HIV-1 activities 467
 Anti-HSV 703
 Anti-HSV activity 1979
 Antihypertensive activity 575
 Anti-inflammatory 415,479,613, 1037,1443
 Anti-inflammatory activity 105,177, 1403
 Anti-inflammatory effects 1509
 Antileishmanial 1117,1537
 Antileukemia 1117
 Anti-malaria 703
 Antimalarial 733,1117,1305,1591
 Anti-metastasis 115
 Antimicrobial 191,415,617,741,761, 1117,1317,1777
 Antimicrobial activities 667,835
 Antimicrobial activity 273,407,493, 523,527,817,1313,1617,1641,1735
 Antimicrobial evaluation 59
 Antimildew activity 1329
 Antimicrobial 733
 Antimycobacterial activity 1673
 Antioxidant 55,265,339,375,389, 415,439,621,627,707,787,823, 1381,1451,1791,1797
 Antioxidant activities 667
 Antioxidant activity 99,139,177,335, 471,655,817,1121,1171,1325, 1403,1575,1587
 Antioxidants 47,1279,1415
 Antioxidation 629
 Antiplasmodial 761,1201,1591
 Antiplasmodial activity 175
 Antiproliferation 1355
 Antiproliferative 1201
 Antiproliferative activity 187,1089
 Antiproliferative properties 1797
 Antispermatic 1183
 Anti-TB 703
 Anti-telomerase 827
 Anti-tumor 415
 Antitumor activities 467
 Antitumor agents 973
 Antitussive 457
 Anti-tyrosinase 253
 Antiviral 249,415,1547
 Antiviral activity 297,1127
 Antiviral effect 149
 Antivirus 43
 Anti-wood-decay fungal activity 1329
 Apiaceae 841
 Apiofuransyl dendranthemoside A 363
 Apoptosis 775,1269,1645,1811
 Aporphine alkaloids 575,1787
 Apoptosis 213
 Aqueous extract 711
 Arabinose 1011
 Araliaceae 363,767,1289
Arctium lappa 463
 Argan oil 1,25,29,33,37,51,55,
 Argan tree 21
 Arginine 43
 Aroma compounds 1767
 Aroma variation 33
Artabotrys hongkongensis 239
Artemia salina 177,1179
Artemisia alba Turra 283
Artemisia herba-alba Asso 407
 Arthropod 1487
Artocarpus lakoocha 1079
 Asarone 1179
Asarum cordifolium 235
Asarum glabrum 235
 Ascomycota 191
 ascorbic acid 627
 Asparagaceae 157
Aspergillus 859
Aspergillus carneus 1071
Aspergillus niger 149,1179
Aspergillus sp. 1069,1127
Aspergillus sydowii 329
Aspergillus microcarpus 1117
 Aspirin 359
 Asteraceae 175,283,439,471,631, 683,689,1175,1477,1523,1711
 Asterelin A 915
 Asthma, Mouse asthma model 1609
Astragalus membranaceus 1321
 Asymmetric dihydroxylation 1011
Atractylodes macrocephala 1321
AtrB induction 1599
 Atrofurin A 599
 Atrofurin B 599
Aurinia leucadea 1463
Aurinia sinuate 1463
 Auxin 609
Axinyssa 1535
 B16A5 343
Baccharoides lilacina 401
Bacillus cereus 429
 Bacteria 493,1435
 Bactericidal 385
 Bactericidal activity 673
 Baker's yeast 863
 Batzellaside B 1011
Bauhinia rufescens 1435
Beauveria sulfurescens 1041
 Behavior 1651
Beilschmiedia erythrophloia 143
 Benzenediols 1715
 Benzisothiazolinone 1301
 Benzophenone 481
 Benzoquinones 1431

- Berbanine 441
 Berberidaceae 441
 Berberine 165
Berberis vulgaris 441
 Bergamotene 663
 Betaine 161
Betula pendula 617
 Betulin 435
 Betulinic acid 435,1099
 Bhutanese traditional medicine 733
 Bibenzyls 115
 Bicyclogermacrene 663,1195,1641
 Bifidobacteria 527
 Biflavonoid 787,1237
 Binder-free RP rotors 311
 Bioactivity 265,747,1675
 Bioassay-guided fractionation 195, 1459
 Bioavailability 121
 Biocides 1301
 Bioclimatic area 817
 Biodiversity 1629
 Biofilm 147,1617,1747
 Biological activities 69,533
 Biomarker 515
 Biorational pesticides 753
 Biosynthesis 771,1195
 Biotransformation 59,859,1041
 Biphenyl derivatives 1265
 Bisabolane 1355
 Bisbibenzyl 915
 Biscembranoid 1209
 Bismuth 419
 Bisuracilyl-substituted 1713
 Biting deterrent 257
 Blue light 791
 Body scents 1651
 Boldine 1797
 Bombacaceae 583
Bombax ceiba 583
Bonannia graeca 841
 Bondowoso 703
Boophone disticha 1213
Boophone haemanthoides 1705
 Bornyl acetate 1629,1643
Boswellia carteri resin 1365
Botryodiplodia 859
Botryosphaeria 859
Botrytis cinerea 1599
 Branched-chain sugar 987
Brassica oleracea var. *italica* 217
 Brassicaceae 1615
 Brassinolide 771
 Brassinosteroids 771
 Brine shrimp lethality 1179
 Bromoindoles 287
 Bromomethylpropanol 925
Brontispa longissima 199
Broussonetia papyrifera 1609
 Bryozoan 693
Bulbus Lili 1447
Bumelia sartorum 207
 Burdock Fruit 463
 Butatriene 851
 Butyldimethylsilyl-2,3-diethyl- β -cyclodextrin 221
 Butyrylcholinesterase 1541
 Butyrylcholinesterase 781
 C57BL/6J 343
 Cadinene 225,1163
 Cadinol 273
 Cadinol 531
 Cadinol 531,1329
Caesalpinia peltophoroides 679
 Caffeic acid 203
Caladenia 701
Calamintha nepeta 139
 Calanolides 803
 Calcium 419
 Calcium antagonist 1633
 Callus 593
 Callus cultures 1587
Calophyllum inophyllum 803
Camellia sinensis 1553
 Camphor 1159
 Canadian medicinal plant 1673
Canavalia 1621
 Cancer 183
 Cancer cells 1053,1377
 Cancer chemopreventive agents 1365
 Cancer Therapy 1811
Candida, yeast 1763
 Canonical correlation analysis 1129
 Capillary electrophoresis 1821
Caragana arborescens Lam. 585
Carapa guianensis 1641
 Carboline 693
 Carbon isotope discrimination 11
 Carbon-13 NMR 235,1633
 Carcinogen activation 1269
 Cardiovascular diseases 539
 Cardiovascular effects 1167
 Carmichaeline A 155
 Carnosic acid 429
 Carolina Sapphire 257
 Carotenoids 171
 Carrageenan 501
 Carvacrol-rich oil 1155
 Carvone 935
 Caryophyllaceae 1687
 Caryophyllene 143,225,401,531,663 1481,1643
 Caspase 775
 Caspases-3 213
 Caspases-8 213
 Caspases-9 213
Cassia fistula 261
Cassia occidentalis 243
Cassia siamea 195,243
 Cassiamin A 195
 Catalytic activity 339
 Catalytic cyclization 999
 Catechins 579
Catharanthus roseus 903,1095
 Catheter-related infections 1747
 CD11b 1755
 CD40 1755
Celastraceae 753
 Cell culture 1419
 Cell cycle arrests 1269
 Cell growth 1301
 Cell migration inhibition 601
 Cell suspension culture 443
 Cellulose 497
Celmisia viscosa 689
 Cembrane diterpenoid 1675
 Cembrane-based 1363
 Cembrene 841
 Centrifugal chromatography 311
 Ceramide 367
Cercospora piaropi 187
 Cerebroside 1285
Chaetomium globosum 1217
 Chakasaponin 1553
 Chalcogenirane 851
 Chalcogenocarbonyl ylide 851
 Chalcones 459,1089
Chamaecyparis obtuse 515
Cheilelejeunea 565
 Chelating 787
 Chemical composition 235,265,523,655,1171,1313,1321,16 37,
 Chemical constituents 1227
 Chemical defenses 397
 Chemical ecology 701,1537
 Chemical homogeneity 1155
 Chemical markers 487
 Chemical studies 1523
 Chemical variability 835
 Chemiluminescence 1559
 Chemoprevention 153,1811
 Chemosignals 1651
 Chemosystematics 565,1515
 Chemotaxis 1559
 Chemotaxonomy 737,1463
 Chemotype 605
 Chemotypes 565,747
Chiliadenus 1159
 Chinese cassia 75
 Chinese herb 827
 Chinese medicine 621
 Chiral 737
 Chiral chromatography 1123
 Chiral pool 1011
 Chiral selector 221
 Chiral synthon 935
Chlidanthus fragrans 1541
Chlorella 859
 Chlorogenic acid 1105
 Chlorophyceae 497
Chlorophyllum molybdites 1227
 Cholinergic 1247
Christia vespertilionis 1591
 Chromanol 431
 Chromatographic techniques 1821
 ChromatorotorsTM 311
Chromolaena pulchella 1711
Chrysactinia Mexicana 135
 Chrysanthenyl acetate 1629
 Chrysandicin 10-methyl ether 1261
 Cineole 1629
Cinnamomum 1777
Cinnamomum cassia 75
 Ciprofibrate 579
 Circular dichroism 1751
 Circular dichroism detection 1075
Cissus quadrangularis 629
 Citral 249,1301
 Citriquinone A 1431
 Citriquinone B 1431
Citrullus lanatus seeds 1367
Citrus grandis 'Tomentosa' 457
Citrus grandis Osbeck 171
 CLA 435
Cladolabes schmeltzii 1527
 Cladolides 1527
 Claisen rearrangement 1101
Cleistopholis glauca 1773
Clematis 211,1093,1563
 Clerodane 873
Clerodendrum inerme 459
 Click chemistry 729
 Clinical trial 1605
 Clonogenic assay 635
 Clovane, *Rumphella antipathies* 1037
 Clusiaceae 1237
 Clusianone 955
 Cluster analysis 131
Cnestis ferruginea 243
 Cocoa 579
 Co-culture 721
 Coenzyme Q10 47
 Coenzyme Q9 47
 Colic 1633
 Colon adenocarcinoma 1255
 Combination effect 711
 Comet assay 389
 Complement inhibitor 1715
 Compositae 225
 Conjugated linoleic acid 435
 Constituents 1217
 Contraception 1183
Conyza bonariensis 1175
Coptis chinensis 823
 CoQ 47
Corema album 1255
 Coriander essential oil 673
 Coriander oil 671
 Coumarin 799
 Coumarin precursor 799
 Coumarins 283,583,803
Crambe tataria 1615
 Crimson Pigmy 165
 Crinane 553,1335
 Crotofolanes 1679
Croton caracasanus 1679
Croton draconoides 1471
Croton micradenus 779
Croton sylvaticus 795
Croton urucurana 1471
Croton zehntneri 1167
 Crown structure 5
 Crypto-chlorogenic acid 1559
Cryptomeria japonica 1785
 Crystal structure 1451
 Cucumariosides 1053
 Cucurbitaceae 79
 Cucurbitane-type triterpene 1367
 Cultured plant cells 847,903,905,907,
Cunninghamia konishii 805
Cupressus arizonica var. *glabra* 257
Curcuma wenyujin 523
 Curcumene 1163
 Curcumene 225,1163
 Curcumin 121,183,359
 Curcumin analog 183
 Curcumol 149
 Curcuphenol 1355
 Curtius rearrangement 1021
 Cyanobacteria 965
 Cyanogenic glycosides 1739
 Cyclic peroxides 725
 Cyclin D1 183
 Cycloadducts 981
 Cycloalkenylation 999
 Cycloartanes 1049
 Cyclodipeptide 169
 Cycloeucaleanol 1213
 Cyclooxygenase 1509
 Cyclooxygenase inhibitory 1365
 Cyclooxygenase-2 501
 cyclopamine 1059
 Cyclopentadiene 919,981
 Cyclopentanediene 897
 Cyclopentanoid 919
 Cyclopeptide 1753
 Cyclopropanation 955

- Cyclopropylcarbinylcation rearrangement 1195
 Cymene 1645
Cystophora torulosa 715
 Cytoprotection 375
 Cytotoxic 439,553,635,683
 Cytotoxic action 1291
 Cytotoxic activities 1053,1175,1527
 Cytotoxic activity 143,161,177,315, 679,1209,1255,1265,1367,1399, 1637,1741,1791
 Cytotoxicity 289,431,435,447,567, 571,601,733,811,1111,1179,1213, 1227,1301,1305,1371,1393,1405, 1535,1645,1665,1691,1733,1811
- Dark 1571
 DART-MS 1721
 Date palm 1289
Daucus aureus 835
 Davanone derivatives 283
 Deacetylasperulosidic acid sodium salt 1507
 Decaturin E 1397
 Decaturin F 1397
 Dehydro-6-deoxoteasterone 771
 Dehydroabietol 1771
 Delphinidin glycosides 1563
Dendrobium pulchellum 115
Dendropanax dentiger 363
 Dengue 1547
 Density functional theory calculations 1075
 Deoxoteasterone 771
 Deoxy-hydroxyascocochlorin 601
 Deoxypretazettine 1541
 Depsidones 109
 Dereplication 715
 Dermatophytes 135
 Development 1
 DHEAS 51
 Diabetes mellitus 1085
 Diacetylcurcumin 359
 Diarylheptanoids 1715
 Diarylheptanoids 351,1715
 Dibenzylbutane lignan 1121
 Dicentrine 443
 Dichloroketene 981
Dictyota coriacea 427
Dictyota menstrualis 1669
 Diglutarylcurcumin 359
 Dihydroagarofuran 753
 Dihydrobenzofuran 1071
 Dihydrochalcones 1081,1255
 Dihydromyricetin 339
 Dihydrostilbenoid 479
 Dihydrotagetone 737
 Dihydroxy-4-methylphenyl-nona-3(Z)-enoic acid 1459
 Dihydroxycrenulide 427
 Dihydroxy-methoxy-xanthone 1731
 Dihydroxyolean-12-en-28-oic acid 761
 Dihydroxy-trimethoxylignan-9',9'-olide 805
Dioscorea cayenensis 323
Dioscorea nipponica 319
Dioscoreaceae 319,323
 Diosmin 545
Diospyros crassiflora 1575
 Diphenyl-propan-1-one 1081
 Dipolar cycloaddition 1021
 Discorhabdin alkaloids 699
- Distichaminol 1705
 Diterpene 689,1201
 Diterpene lactones 1725
 Diterpenes 75,1771,1755
 Diterpenoid 67,873
 Diterpenoid alkaloid 155
 Diterpenoid alkaloid 1701
 Diterpenoids 1679
 Diynes 631
 DNA barcodes 125
 DNA damage 389
 DNA fragmentation 635
 DOCA-salt hypertensive rats 1167
 Docking 1223
 DPPH 475,639,1137
Dracaena marginata 157
Dracaena thalioides 315
 Dragon's blood 337
 Drimane sesquiterpenoids 147
 Drought adaptation 11
 Drug delivery 121
 Drug discovery 539
 Drug interaction 1269
 Drying 1479
Drynaria rigidula 703
 Dual COX/LOX inhibition 105
 Duciformine 1545
Dysidea avara 289
Dysidea cinerea 1751
 Dysideidae 1751
 Dysideolide A 1751
 Dysideolide B 1751
- E/Z-isomers 869
 EA.hy926 1291
 Earthworms 1149
 Ebenaceae 1575
 Echinoderm embryo development assay 699
Echinophora spinosa 527,1645
 Edible oils 47
 Eight-membered carbocycle 883
 Elastase 1037
 Elatinyne 729
 Eleganoid A 1373
 Elemicin 235
 Elemol 1329
 Elephantiasis 803
 Elevated-plus maze test 515
 Elicitor 443
 Elicitor 593
Elionurus hensii 655
Emericella nidulans 1285
 Enantiomer 221,737
 Enantiomeric bicyclic γ -lactones 981
 Enantioselective synthesis 925
 Endo-fenchol 683
 Endoperoxide 1537
 Endophyte 373,1673,1735
 Endophytic fungus 83,587,601
 Endothelial cells 1547
 Ent-abietane diterpenoids 1519
 Ent-kaurane diterpene glycosides 1523
 Ent-kauranes 1041
 Enzymatic reaction 919
 Enzymatic reduction 897
 Enzyme-mediated resolution 863
Ephedra trifurca 601
 epi-cercosporin 187
 Epidioxysterol 1535
 Epilolide 427
 Epimanool 689
- Epiphytic medicinal plant 703
Epi- α -cadinol 401
 Equator Apricot with Red Eye 1095
Equisetum 509
Erechtites valerianaefolia 1477
 Ergosterol 1217,1763
 Ergosterol inhibition 261
 Ericaceae 1255
Eryngium maritimum 841
 Eschenmoser rearrangement 949
 Essential oil 135,143,225,227,231, 265,273,403,407,505,515,523,531, 645,655,667,683,733,737,841, 1167,1171,1297,1305,1309,1325, 1329,1477,1493,1633,1637,1643, 1645,1771,1791
 Essential oil composition 139,243, 277,679,1163,1175,1179,1471, 1481,1625,1773
 Essential oils 131,269,519,527,741, 747,829,835,1317,1321,1479,1777, 1785
 Esters 435
 Ethyl acetate-soluble fractions 1579
Eucalyptus 397
Eucalyptus perriniana 847,903,905,907
 Eugenol 505,1481
Eupentacta fraudatrix 1053
Euphorbia guyoniana 1519
Euphorbia latex 63
 Euphorbiaceae 779,795,1111,1471,1519,1679
 Extraction 1821
 Extractive compounds 1755
Eysenhardtia platycarpa 177
- Fabaceae 585,1421,1591,1625
 Falcarinol 1123
 Fatty acid 1297,1763
 Fatty acid composition 411
 Fatty acids 1171
 Fatty ester methyl ester 1763
 Fennel oil 671
 Ferrier(II) reaction 987
Ferula ferulaeoides 1353
 Ferulic acid 203
 Ferulic acid isomers 509
 Fibroin 1755
Ficus carica 485
 Fir 227,1297
 Fish pathogenic bacteria 1223
Fissistigma bicolor 663
Fissistigma chloroneurum 663
Fissistigma cupreonitens 663
Fissistigma maclurei 239
Fissistigma pallens 663
Fissistigma petelotii 663
Fissistigma rufinerve 239
Fissistigma shangtzeense 663
 Flavan 447
 Flavanone 1075
 Flavanone 905,1409
 Flavanones 177
 Flavivirus 249,1487
 Flavone C-glycoside 1413
 Flavones 85
 Flavonoid 605,733,791,1241,1719
 Flavonoid glucoside 333
 Flavonoids 171,283,583,585,703, 1079,1081,1403,1405,1415,1731
 Flavonol glycosides 1093,1251
 Flavonols 99
- Flavor variation 33
 Floratheasaponin 1553
 Flow Cytometry 183
 Flower colors 1563
 Flowers 459,1171
 Fluorescence 339
 Forest degradation 1
 Formal synthesis 897
Fragaria virginiana 1673
 Free radical scavenging activity 1629
 Free-radical oxidation 589
 Free-radical scavenging 621
 Freeze-dried strawberry cream filling 1767
 Fruit 79
 Fruit shape 25
 Fruit volatiles 129
 Fucoxanthin 1381
Fucus spiralis 373
 Functionalized cyclitol 987
 Fungi 1435
 Furanocoumarins 485
 Furocoumarins 171
 Furostanol bisdesmoside 315
 Fusaodavinvin 83
Fusarium sp. 83
 Fused ring compound 929
- Galactolipids 1285
 Galanthamine 1335
Galanthus xvalentini nothosubsp.
Subplicatus 327
 Gallic acid 621
 Gamma-aminobutyric acid 1233
Garcinia 1237
Garcinia mangostana 1733
Garcinia schomburgkiana 1265
 Gas chromatography-mass spectrometry 63
 Gas exchanges 1387
 Gastrointestinal motility 1143
 GC 401
 GC/FID 231
 GC/MS 33,225,231,401,407,639, 781,1163,1643,1755
 GC-FID 651
 GC-MS 129,135,139,211,327,651, 683,1129,1321,1325,1463,1479, 1633,1785
 GC-MS 139
Geijera parviflora 747
Gelsemium elegans 1373
 Genista 593
 Genistein 343
Geosmithia lavendula 191
 Geranium 645
 Geranyl flavonoid 335
 Geranylated compounds 103
 Germacrene C 1773
 Germacrene D 663,841,1481
 Germination treatments 15
Geum rivale 505
Geum urbanum 505
Ginkgo biloba 1153
Ginkgonis folium 1153
 Globalization 1
 Glucose uptake 1279
 Glucosidase 1085,1279
 Glucosidase activator 757
 Glucosidase inhibition 757
 Glucoside 119,847
 Glucosinolates 217,1463,1615

- Glucosylation 847,903,905,907
 Glucosyltransferase 119
 Glucuronolactone glycoside 1289
 Glutamate decarboxylase 1233
 Glutathione reductase 1135
 Glyceroglycolipid related compound 315
 glyceryl-leu-ala-leu-leu-val-leu-ala 381
Glycine max 453,907
 Glycoalkaloids 1387
 Glycolipids 165
 Glycosides 689,703,1697
 Glycoxy uracil 1227
Glycyrrhiza 1617
Glycyrrhiza glabra 1721
Glycyrrhiza spp. 1089
Glycyrrhiza uralensis 823
Glycyrrhiza uralensis 1721
 Glycyrrhizic acid 415
 Gold nanoparticles 1149
 Gorgonian-associated fungus 1069
Gossypium hirsutum 907
 Green Chemistry 1493
 Green leaf volatiles 393
 Green synthesis 1149
 Guaiol 1353
 Guanidine alkaloids 1229
 Guinea-pig ileum 1247
 Guttiferaceae 1265,1731
Gypsophila trichotoma 765
- Hairy root 217
 Hairy root culture 609
Hakea sericea 1459
 Harziandione 67
 Headspace 63
 Heat-labile 385
 Heavy metals 21
Hedychium coronarium 519
Hedychium flavescens 519
Hedychium neocarneum 519
Hedychium speciosum 519
Hedychium stenopetalum 519
Hedyotis corymbosa 1683
Hedyotis diffusa 1683
Hedyotis tenelliflora 1507
 Heliannuol-D 1197
Helichrysum kraussii 1403
Helichrysum odoratissimum 1403
Helichrysum zivojinii 1291
 Hemorrhagic disorders 545
 Henry reaction 987
 Hep-2 183,1431
 HepG2 cell 375
 Heptenal 509
Heracleum sibiricum 1309
 Herbal tea 479
 Herbivory 15
 Herbs 1617
 Herpes simplex 297
 Heterocycle 935
 Hexenol 505
Hibiscus vitifolius 433
 High-performance liquid chromatographic fingerprint 1447
Hippospongia 571
 HL-60 cells 315,775
 HLA 1651
 Homoisoflavanone 597
 Homolycorine 781
Hornstedtia leonurus 513
 Horsetails 509
- Horsfieldia superba* 447
 HPLC 171,471,1045,1075,1105, 1241,1463
 HPLC-DAD 485
 HPLC-ESI-MS/MS 1725
 HPLC-MS-MS 1129
 HPLC-NMR, 1443
 HPLC-PDA 1443
 HSQC-ROESY 815
 HT-29 1255
 Human cancer cells 431,1405
 Human lymphocytes 109
 Human pheromones 1651
 Humulene 143
 Humulene-type sesquiterpene 1665
 Hydrodistillation 1493
 Hydrogen-bonding acidity 85
 Hydroxycholestane side chain 1219
 Hydroxylation 905
 Hydroxyphenethyl-methoxy-methylphenol 1435
 Hydroxythymol 367
 Hypericaceae 103
Hypericum 347
Hypericum laricifolium 1731
Hypericum richeri Vill. subsp. *grisebachii* (Boiss.) Nyman. 231
 Hypertension 539
 Hypoglycemic 1085
 Hypoglycemic activity 207
Hypogymnia physodes 109
Hyptis incana 1665
- IBX oxidation 883
 ICP-AES 21,639
 IL-6 181
 Image analysis 1065
 Iminosugar 1011
 Immune response 1755
 Immunomodulatory activities 1257
 Immunosuppressant 815
In silico 1049
in vitro cytotoxic potential 277
 Indol-3-carboxylic acid 1245
 Indole alkaloid 329
 Indonesia 703
 Inducer 1455
 Inflammation 501,1609
 Inhalation 515
 Inhibition of cancer cell lines 367
 Inhibitor 393
 Inhibitory effects 1669
 Insect antifeedants 199
 Insecticidal 1777
 Insecticidal activity 289,1353
 Insulin 1279
 Integerriminol 799
 Intestinal dysbiosis 527
 Intachub, 1567
 Intramolecular copigmentation 1563
 Intramolecular lactonization 1545
Inula crithmoides 1587
Inula sp. 475
 Invasiveness 1291
 Invertebrate extracts 1149
 Iridoid 1241,1507
Iris pseudacorus 1419
 Irregular sesquiterpenes 283
 Isocarvone 949
 Isocercosporin (Helix P) 187
 isocoumarin 381,1575
 Isoflavones 99
 Isoflavonoids 589,593,1419,1423
- Isolated thoracic aorta 1167
 Isomyristicin 1305
 Isophorone 59
 Isoquinoline-isoquinolone dimer 441
 Isorhamnetin 1251
 Isotope labeling 771
- Jatropha dioica* 297
J_{H-H} values 725
 Joziknipholones A and B 1261
Juglans nigra 1481
Juglans regia 1481
 Juglone 1481
Julocroton triqueter 1471
Juniperus formosana 1329
- Kaempferia marginata* 1205
 Kaempferol 1251,1719
 Kaempferol 3,7,4'-glycosides 1093
 Kaempferol 3-O-glucoside 1559
 Kaempferol 3-O-triglycosides 453
 Kenganthranol F 103
Kitaibelia vitifolia 667
Kniphofia foliosa 1261
 Knipholone anthrone 1261
 Koninginin 1285
 Kynurenic acid 651
- Labdane diterpenoids 567
 Labdanoids 1755
 Labiatae 1665
 Lactam 1069
 Lactam Synthon 889
 Lactobacilli 527
 Lactone 1069
 LAI 5
 Lamiaceae 125,1771
 Lankongensisine A 1545
 Lanostadiene 299
 Lanostane 299
 Larvicidal 257,1777
 Laser polarimetric detection 1075
Laurencia elata 729
Laurencia similis Aristolan 287
 Leaf essential oil 1165,1641
 Leaves 269,453
 Lectins 1621
Leguminivora glycinivorella 393
 Leguminosae 335
 Lejeuneaceae 1515
 Lejeuneoideae 1515
Lepidagathis fasciculata 1163
 Leukemia cell 1665
 Levopimaric acid 293
 LH 51
 Light 1571
 Lignan 113,805,1427
 Lignans 463,467
Lignosus rhinocerotis 711
Ligustrum vulgare 1509
 Lilac aldehydes 651
 Limonene 129,221,227,249
 Linalool 671
Lindera obtusiloba 181
 Linoleic acid 1551
 Linoleic acid rich 25
Liparis nervosa 1115
 Lipid Peroxidation 1137
 Lipoxygenase 105,1509,1595
Lippia essential oil 249
 Litchi 1583
Litsea coreana 479
- Liverworts 565,1313,1515
 LMA 5
 Lonchocarpenin 1099
Lonocera japonica 1609
Lophiostoma 1223
 L-Pyrogutamic acid 1011
 Lung cancer 115
Lychnis senno 1413
 Lycorine 1335
 Lyngbouilloside 965
 Lyngbyaliosides 965
 Lyoniresinol derivative 113
Lysimachia thyriflora 1691
Lythrum salicaria 1247
- Maackia amurensis* 589
Machilus mushaensis 273
 Macroalgae 1811
 Macroelements 645
 Macrolactones 965
 Macrolide 815,1127
 Macrophage 1755
 Madagascar 1165
 Madindoline 897
Magnolia grandiflora 135
 Makaluvamines 1551
 Malaria 1135,1261
 MALDI MS 301
 Mangostin 1101,1257
 Mangrove 1735
Marcetia latifolia 1049
 Marine fungi 721,1223
 Marine fungus 67
 Marine invertebrate 1537
 Marine natural product 1669
 Marine red algae 533
 Marine sponge 1209,1399,1713, 1751
 Marine sponges 1551
 Marine-derived fungus 1071,1127
Marrubium globosum libanoticum 567
 Mass spectrometry 485,605,771,1059
 Mate choice 1651
matK 125
 MDA 627
 MDA-MB-231 1291
 MDR1 resistant strain 1599
 Medicinal mushroom 711,1763
 Medicinal plants 385,823,1305,1821
 Medium 609
 Medium ring ethers 929
 Medium strength 1045
 Megastigmane glycoside 363
 Melanin 343,629,1359
 Melanin synthesis inhibition 427
 Melanogenesis 1359,1605
 Melanogenesis inhibitory activity 1367
 Melanoma 567
 Melastomataceae 1049
 Melatonin 47
Melia azedarach 1719
 Meliaceae 1641
Melissococcus plutonius 1625
 Melitidin 457
Melodorum fruticosum 239
 Menispermaceae 575
Mentha haplocalyx 1479
 Metabolic diseases 1439
 Metall nanoparticles 1149
 Metallo-protease 1467

- Metastasis 343
Methanolic extracts 1567,1641
Methoxy-D-mannono-1,4-lactone 363
Methoxymedicarpin 1421
Methyl 27-caffeoyloxyoleanolate 433
Methyl barbascoate 873
Methyl carnosate 429
Methyl salicylate 1481
Methyl-13-tridecanolide 925
Methylanthocyanidin 1095
Methylbenzylamine 981
Methyleugenol 513
Methylpelargonidin 1095
Methylpelargonidin 3-galactoside 1095
Methylpelargonidin 3-robinobioside 1095
Methyl-pentadecanolide 925
Methyl- β -cyclodextrin 221
MHC 1651
MIC 59,385
Microbial transformation product 149
Microcin 1371
Microcos tomentosa 1371
Micromelum integerrimum 799
Micronucleus test 109
Microwave accelerated distillation 1493
Milk thistle 1801
Milletia richardiana 1099
Mimosa pudica 1669
Minerals 639
Minimum inhibitory concentration 493
Mitochondria 47
Mitochondrial 635
Mixed fermentation 721
Mixture 1301
Mixture analysis 1721
Modified Si-Miao-San 1137
Modified ultrasound-assisted extraction 1683
Molecular authentication 125
Molecular docking 105
Molinaea retusa 1201
Momordica charantia var. *abbreviate* 79
Monanchomycalin 1399
Monanchora pulchra 1229,1399
Monoterpene 519
Mood 1651
Moraceae 599
Mori Cortex Radicis 1443
Moringa oleifera 1559
Moroccan Western Rif 1155
Morphinandienone alkaloids 779
Morus alba 1579
Morus atropurpurea 599
Mosquito 257
Mountain tea 639
Mouse liver 579
mRNA expression 1205
MRSA 1617
MS 1229
MTT 635
Mucic acid 3-O-gallate 461
Mucor 859
Multidrug resistance 1377
Multidrug resistant strain 1305
Multivariant analyses 707
Multivariate statistical analysis 487
Muscolide 925
Mutagenicity 795
Mycobacterium tuberculosis 1785
Myrcia 269
Myristicaceae 447
Myrmekioderma 1355
Myrosinase 1615
Myrtanal 505
N,N-aminal 961
Nanoparticles 121
Narcissus 781
Nardal 1513
Nardin 1513
Nasutitermes macrocephalus 69
Natural compounds 105,1487
Natural preservative 829
Natural product 955
Natural Product Synthesis 929
Natural products 293,533,715,729, 965,973,1135
N-containing 1713
Nematicidal 1777
Neochlorogenic acid 1105
Neolitsea variabilis 531
Nepal 1179
Nerolidol 829
Nervogenic acid glycoside 1115
Neurite outgrowth stimulation activity 711
Neuritogenicity 1583
Neurodegenerative 795
Neuroprotection 1415
Neuroprotective 1583
Neurospora crassa 59
New anthocyanidin 1095
NF- κ B 289
Nicotiana tabacum 903
Nigella sativa seeds 213
Nigropsora sp. 1673
Nitric oxide 1205
NMR 157,779,301,597,779,1049, 1053,1093,1229,1237,1289,1399, 1519,1527,1687
NMR spectra 1219
NMR spectroscopy 1711
NMR spectroscopy 85
NMR-based metabolomics 1409
Nociception 1621
nOe NMR enhancements 729
Nonaprenylsulfate 1595
Non-targeted screening 605
Nootkatone 859
Norbergenin 1575
Norcucurbitane 79
Obesity 195,481
Ocimene 129
o-Coumaric acid 1269
Oct-1-en-3-ol 1625
Octen-3-ol 505
Odoratanone A 1373
Okra 1085
Olean-12-ene saponins 1691
Oleanane 767
Oleanolic acid 1683
Oleic acid rich 25
Olive oil 51
Opopanax chironium 841
Optical resolution 981
Orchid 701
Organic azides 1021
Oroidin 961
Orthogonal partial least squares 1129
Osmundaria 533
Ottelia alismoides 351
Otteliones 351,973
Oxacycles 929
Oxasetin 1223
Oxepin-containing alkaloids 1071
Oxidation 293
Oxidative alkylation 999
Oxidative coupling 915
oxidative damages 375
Oxidative stability 25,37
Oxidative state 37
Oxidative stress 1301,1415
Oxidative stress damage protection 621
Oxindole alkaloid 1547
Oxo- α -ionol 509
Oxodioscin 319
Oxyphenylated secondary metabolites 1089
Oxyresveratrol 613
Ozone 293
P19-derived neurons 1583
p53 213
Pachyrrhizus sp. 1423
Paclitaxel 807
Paenibacillus 381
Paenibacillus larvae 1625
Pale red flower color 1095
Palladium (II) 999
PANC-1 463
Pancreatic cancer 463,811
Pancreatic lipase 195,481,1439
Paniculosis 757
Papain 1467
Parmeliaceae 109
Parthenocissus tricuspidata 1439
Pastinaca sativa 1123
Patchoulene 1773
Pathogenic fungi 1785
Paw edema 359
PBMcs 1257
PCR-RFLP 125
Pedicellamide 1451
Penicillium 373
Penicillium citrinum 587,1431
Penicillium oxalicum 1397
Penicillium sp. 329
Penicillocitricin A 587
Pentacyclic guanidine alkaloids 1399
Pepsin 1467
Peptide antibiotic 381
Periodontal ligament (PDL) 613
Permeability 1547
Perovskia abrotanoides 1633
Peroxisome proliferator activated receptor 579
Persistence in propagules 397
Petronigrone 1209
Petrosia nigricans 1209
Phaeodactylum tricornutum 1381
Phaeophyceae 497
Phakellin 961
Phakellstatin 961
Pharmaceutical formulations 545
Phebalium 737
Phenol glycoside 479
Phenolic acids 605,1725,1821
Phenolic compounds 207,475,567,
Phenolic constituents 337
Phenolic glucoside 181,367
Phenolic glycosides 1739
Phenolics 165
Phenols 539,1715
Phenyl-4*H*-oxepino[2,3-*b*]pyran-4-one 1081
Phenylacetaldehyde 1625
Phenylbenzofuran 599
Phenylethanol 1625
Phenylpropanoids 513,585,1089,1387,1781
Phoenix dactylifera 1289
Phoenixoside B 1289
Phomopsis sp. 1735
Phomoxanthone 1735
Phyllanthus emblica 461
Phyllanthus niruri 493
Physicochemical properties 411
Phytoalexins 485
Phytochemical analysis 621
Phytochemical study 1755
Phytochemicals 791,1435
Phytochemistry 1081
Phytohormones 1275
Phytolacca Americana 119
Phytosterol 1297
Phytotoxic activity 139
Phytotoxicity 407
Piceatannol 119
Pimpinella monoica 1643
Pinene 129,841,847
Pinus thunbergii 1755
Piper aduncum 1325
Piper auritum 1325
Piper capense 1791
Piper pedicellatum 1451
Piper umbellatum 1325
Piperaceae 1451
Piperidine alkaloids 1005
Plakortis 725
Plant material 545
Plant volatiles 393
Plasmodium falciparum 1135
Plasticity index 5
Pleurosporum amabile 1305
Pleurotus giganteus 1763
pMDI 121
Podophyllotoxin 199
Pogostemon hirsutus 1771
Pollen 645
Pollinator 701
Pollinators 701,1159
Polyalthia longifolia var. *pendula* 239
Polyalthia rumphii 1427
Polyalthurea 1427
Polyamines 1275
Polycyclic ethers 929
Polygonatum cyrtoneura 597
Polyketides 373,587,721,1005,1089
Polyphenol 1553
Polyphenolic compounds 1419
Polyphenolic profile 817
Polyphenols 707,1247,1725
Polyploidy 1387
Pomegranate peel 707
Poncirus trifoliata 1143
Populus nigra 617
Populus tremula 617
Porcelia macrocarpa 277
Postmenopausal women 55

- Potato biodiversity 1387
 Preferential cytotoxicity 463
 Prenylated compounds 103
 Preservation 29
Primula exudates 1081
 Principal component analysis 21,523
 Procatechuic acid 203
 Pro-coagulant activity 1115
 Procyanidin 75
 Progenin III linoleate 323
 Progenin III palmitate 323
 Prolyl oligopeptidase 1541
 Propolis 617
 Protective 1801
 Protein kinase inhibition 289
 Protoanemonin 211
 Protocatechuic acid 1579
Prunus cerasus L. honey 651
Prunus persica 1739
psbA-trnH intergenic spacer 125
 Pseudoionone 509
 Pseudolaric acid A 877
Pseudomonas aeruginosa 1467
 Psoralen 485
Psorospermum aurantiacum 103
 Pterocarpan 589,1421
Pterocella vesiculosa 693
 Ptychanthoideae 1515
Pulicaria undulate 757
 Pulicarside 757
 Pummelo juice 171
 Pyocyanin 1455
 Pyranocoumarin 1099
 Pyrazine 701
 Pyridinecarbothioamide 593
 Pyrolysis 1701
 Pyrrole derivative 1451
 Pyrrole-imidazole alkaloid 961
 Pyrrolizidine alkaloids 439,1545, 1711
- Quality 37
 Quality assessment 1447
 Quality evaluation 1553
 Quantitative analysis 1553
 Quantitative determination 99
 Quercetin 1085,1251,1719
 Quercetin 3-*O*-glucoside 1559
 Quercetin 3-*O*-triglycoside 453
 Quercetin glycosides 1105
Quillaja saponaria 1697
 Quinolin-2,4-dione alkaloid 775
 Quinopimaric acid 293
 Quinovic acid 573
 Quorum sensing 147,1455
- Radical scavenging activity 1551
 Radical-scavenging 787
 Radix *Paconiae* Alba 487
 Radix *Paconiae* Rubra 487
Ralstonia solanacearum 337
 Ranunculaceae 211,827
 Rat ileum 1143
 RCM reaction 883,949
 Reactive oxygen species 1255
 Rearrangement 293
 Reduction 905
 Regio-isomer 1213
 Relative configuration 725,729
 Resveratrol 609,907
 Resveratrol synthase 1571
 Reversed-phase chromatography 311
- Rhantherium adpressum* 1171
 Rhizomes 1261
 Rhodophyceae 497,533
Rhus semialata var. *roxburghiana* extracts 621
 Ring closing metathesis 877
 Rippertane 69
Robinia pseudoacacia 335
 Root bark oil 1773
 Root elongation 15
 Root/stem ratio 15
 Roots 433,827
 Rosaceae 1105
 Rosemary 817
 Rosmarinic acid 471,1275,1747
Rosmarinus officinalis 131
 Rotation Planar Chromatography 311
 Rotenoid 1423
 RP-HPLC 475
 Rumphiin 1427
 Rutaceae 775,799
- Safrole 235
 Sake 1605
 Salacca 1583
Salicornia herbacea 203
 Salicylic acid 911
Salvia heldreichiana 1637
Salvia officinalis,
 Sand dollar 699
 Sandalwood oil 253
Sandaracopimarinal 1163
 Santalol 253,869
 Santalol derivatives 869
 Sapodilla 1583
Saponaria officinalis 1687
 Saponins 43,1201,1697
 Sapotaceae 207
 Sarcophine 153
 Sarcophine-diol 153
Sarcophyton sp. 1675
Sarcotragus sp. 1595
Sargassum siliquastrum 431
Saussurea graminea 631
Saussurea hieracioides 631
Saussurea stella 631
Saussurea tatsienensis 631
 Sclerane 571
Scapania aspera 1313
Schefflera elegatissima 767
 Schibicolignan A 1121
Schinu molle 135
Schisandra bicolor 1121
Schisandra wilsoniana 467
 Schmidt reaction 1021
Sclerocarya birrea 1279
Scutellaria 1617
Scutellaria baicalensis 1233
Scutellaria baicalensis 609
Scutellaria lateriflora 1415
 Sea cucumbers 301,1053,1377,1527
 Seasonal variations 131
 Seaweeds 497,533,1811
 Seco-diterpenoid alkaloid 1701
 Secondary metabolites 1071,1697,1821
 Sedative 1195
 Seed 227
 Seed oil 411
Senecio delphinifolius 439
Senecio roseiflorus 175
 Senecioneae tribe 1477
- Senna obtusifolia* 1669
 Sensory evaluation 1523
 Sensory evaluation 37
 SERCA 207
 Sericin 501
 Serratanoids 1755
 Sesquiterpenes 269, 289,1195,1665
 Sesquiterpenic fraction 139
 Sesquiterpenoids 565,571,935
 Seven-membered carbocycle 877
 Severibuxine 775
Severinia buxifolia 775
 Sex 1651
 Sex pheromone 393
 Sexual deception 701
 SHRSF 627
 SH-SY5Y cells 1213
 SiHa cells 213
 Silibinin 1801
 Silicates 419
 Silk peptide 1755
 Silver nanoparticles 1149
 Silymarin 1801
 Si-Miao-San 1137
 Simultaneous analysis 1443
Sinularia gibberosa 1363
 Sitosterol glycoside 1245
 Skin cancer 153
 Skin whitening 1359
 SmI₂-mediated Reductive Coupling Reaction 889
 Smilacaceae 113
Smilax microphylla 113
 Smooth muscle relaxation 1143
 Snails 1149
 Soft coral 1675
Solanum erianthum 243
Solanum torvum 243
Solenostemon scutellarioides 1275
 Solid-phase extraction 203
 Solid-phase micro-extraction 63
 Solvent extracts 655,741
Sophora flavescens 1409
 Sorbus 1105
 Soybean 453
 Spasmogenic effect 1247
 Spectral data 1523
 Spectrophotometry 347
 Spectroscopic and RP-HPLC analyses 99
 Spin-labeled 199
Spirospermum penduliflorum 575
 Spirostan glycoside-fatty acid conjugates 323
 Spirostan glycosides 319
 SPME GC-MS 1313
 Sponge 1229,1535
 Squalene 1245
 Stability 29,385,1381
Stachys rupestris 1637
Staphylococcus aureus 135,1467,1747
 Staudinger-aza-Wittig reaction 1021
 Steam Distillation 1493
Stemona alkaloids 695
Stemona phyllantha 1065
Stemona tuberosa 1065
Stephania venosa 443
 Stereochemistry 1219
 Stereoselective synthesis 863,1005,1741
 Stereoselectivity 955
 Steroid 1219
- steroidal alkaloids 1059
 Steroidal saponins 157
 Steroids 165,1227
 Sterol 191
 Sterols 765
Stevia rebaudiana 1523
Stichoneuron halabalensis 695
 Stilbene derivatives 1439
Streptococcus pyogenes 135
Streptomyces sp. 1393,1753
Streptomyces 169
Streptomyces michiganensis 169
 Stress 515
 Striatane 565
 Structural elucidation 83,587,1041,1431
 Structure analysis 1393
 Structure characterization 1523
 Structure-activity relationships 1551
 structure-odor relationship 869
 Subcritical extraction 1719
 Sub-inhibitory 1617
 Successive Ring-opening and Lactonization 889
 Sucrose 1275
 Sulfated saponin 765
 Superfood 791
 Superoxide anion 1037
 Surface exudates 175
 Suspension 593
 Suzuki-Miyaura coupling 915
 Synergism 1567
 Synergistic 393
 Synergistic antioxidant 1321
 Synthesis 911
 Synthetic seed 1275
Syzygium paniculatum 129
- Tail flick 359
 Taiwan 1553
 Tamarind 1583
 Tannic acid 1455
Tapiscia sinensis 587
Taraxacum 1617
Tarchonanthes 683
 Tartary buckwheat 1571
 TAS 627
 Taxifolin 3',7-*O*-diglucoside 903
 Taxifolin 3'-*O*-glucoside 903
 Taxifolin 7-*O*-glucoside 903
 Taxifolin β-D-glucoside 903
 Taxodiaceae 805
 TCM 1447
 TD DFT 597
 Tea flower 1553
 Teneoside D 1507
 Teneoside E 1507
Terminalia brownie 761
 Terpenes 69,227,539,703,1591
 Terpenoids 63,1159,1515
 Terpeneol 129
 Testosterone 51
 Tetrahydroamentoflavone 787
 Tetralones 863
 Tetrasulfanes 1599
 Tetradotoxin 987
 Therapeutic 1801
 Thermal treatment 29
 Thin-layer chromatography 1065
 Thioketene 851
 Thioredoxin reductase 1135
Thymra capitata 1155
Thymus vulgaris 605

- Tiliaceae 1371
 Time-kill 385
 Timing of emergence 15
 Tirucallic acid 1365
 TLC-bioautography 253
 TNF- α 181
 Tocopherol 25
Tofieldia nuda 1251
 Tofieldiaceae 1251
 Total flavonoids 347
 Total phenolic acids 347
 Total phenols 639
 Total polyphenol content 1587
 Total synthesis 873,919,935,955,
 961,973,1011,1021,1101,1197
 Trace elements 21,645,1153
 Traditional Chinese medicine 1447
 Transcript level 1233,1571
 Trehalose addition 1767
Triantha japonica 1251
Trichoderma erinaceum 67
 Trichodermaerin 67
 Trichomes 645
Trichophyton 741
 Triclosan 1301
 Trifarane 565
Trifolium pratense L. ssp. *Nivale*
 1625
Trifolium repens 99
 Trihydroxy-3'-methoxy-3,4-
 methylenedioxygnan-9',9-olide
 805
 Trinervitane 69
 Trinorsesquiterpenes 863
 Triterpene 43,299,1099
 Triterpene glycosides 301,1053,
 1377,1527
 Triterpene saponins 573,767
 Triterpenes 1049,1697,1755
 Triterpenoid 79
 Triterpenoid saponins 1687
Triumfetta semitriloba 1245
 Triumfettalarein 1245
 TRPV1 1229
 TRPV3 1229
 Trunk bark oil 1165
 Trypanocidal activity 403
 Trypsin 1467,1755
 Tuberostemonine 1065
 Tubulin polymerization inhibitors
 973
 Tumor 343
 Tumor necrosis factor- α 1205
 Tumor suppressors 1269
 Turmeric oil 807,811
 Turmeric total extracts 1129
Tussilago 1617
 Type 2 diabetes mellitus 1279]
 Typicosides 301
 Tyrosinase 629,1137,1605
 Tyrosinase enzyme 1435
 U937 cells 1645
 Ubiquinone 47
 UHPLC-MS 1247
 Unactivated olefins 999
Uncaria tomentosa 1547
 UPLC-ESI-MS/MS quantification
 1691
 UPLC-Q-TOF/MS 487
 Urea 1427
 Ursolic acid 1683
 Ushikulide 815
 Vago-vagal reflex 1167
 Valencene 859
 Valerenal 1513
 Valerenic acid 1513
 Valerian 1195
 Valerianaceae 1513
 Vanilloyl-taraxerol 1371
 Vein wall remodeling 545
 Venous stasis 545
Vepris madagascarica 1165
Veratrum californicum 1059
 Verbenaceae 459
 Vibrational circular dichroism 1075
 Vietnam 235
 Vietnamese cassia 75
 Vinaceuline 1753
 Vinylguaiaicol 509
 Violaceine 147
 Viridiflorol 1629
 Vitamin E 55
Vitex glabrata 125
Vitex negundo 1241
Vitex rotundifolia 1405
 VLC 635
 Volatile Constituents 1493
 Volatile Organic Compounds 509
 Volatiles 227,1159,1297
 Volcano plot 1721
 Vomifoliol 651
 Water extract 1143
 Water tea-infusion 639
 Water use efficiency 11
Wedelia biflora 367
 Wittig reaction 1101
 WUE 11
Wyethia mollis 299
 Xanthine oxidase 1409
 Xanthone 1237,1733
 Xanthones 583,1731
 Xiao Jian Zhong Tang 389
 Xingnaojing 627
 X-ray 1353
 X-ray analysis 293
 X-ray-fluorescence analysis 1153
 XRD 497
Xylopia frutescens 403
Xylopia laevigata 403
 Yellow-Fever Virus 249
 YW3699 883,949
Zephyranthes grandiflora 161
Zieria 741
 Zimoside A 481
 Zinc 419
Zingiber roseum 221
 Zingiberace 1205
 Zingiberaceae 829,1317,1715
Ziziphus mucronata 1279
 Zygophyllaceae 573
Zygophyllum cornutum Coss. 573
 Zyzzyanones 1551

Natural Product Communications

Manuscripts in Press Volume 8, Number 12 (2013)

Differences in the Chemical Composition of *Arnica montana* Flowers from Wild Populations of North Italy

Maria Clauser, Nicola Aiello, Fabrizio Scartezzini, Gabbriella Innocenti and Stefano Dall'Acqua

Essential Oil Compositions of Two Populations of *Salvia samuelssonii* Growing in Different Biogeographical Regions of Jordan

Ammar Bader, Pier Luigi Cioni, Nunziatina De Tommasi and Guido Flamini

In Vivo Anti-inflammatory Activity of Some Naturally Occurring *O*- and *N*-Prenyl Secondary Metabolites

Francesco Epifano, Salvatore Genovese, Serena Fiorito, Roberto della Loggia, Aurelia Tubaro and Silvio Sosa

New Diterpenes from *Azorella spinosa*

Luis Astudillo, Margarita Gutiérrez, Luisa Quesada, Aurelio San-Martín, Luis Espinoza and Patricio Peñailillo

Isolation of a New Anti-inflammatory 20, 21, 22, 23, 24, 25, 26, 27-Octanorcurbitacin-type Triterpene from *Ibervillea sonora*

Angel Jardón-Delgado, Gil Alfonso Magos-Guerrero and Mariano Martínez-Vázquez

Anti-trypanosomal Phenolic Derivatives from *Baccharis uncinella*

Simone dos S. Grecco, Maria Júlia P. Félix, João Henrique G. Lago, Érika G. Pinto, André G. Tempone, Paulete Romoff, Marcelo José P. Ferreira and Patricia Sartorelli

Volatile Organic Compounds of six French *Dryopteris* Species: Natural Odorous and Bioactive Resources

Didier Froissard, Sylvie Rapior, Jean-Marie Bessi  re, Alain Fruchier, Bruno Buatois and Fran  oise Fons

Intraspecific Variability of the Essential Oil of *Cladanthus mixtus* from Morocco

Anass Elouaddari, Abdelaziz El Amrani, Jamal JamalEddine, Jos   G. Barroso, Luis G. Pedro and Ana Cristina Figueiredo

Antiplatelet Aggregation Effects of Phenanthrenes from *Calanthe arisanensis*

Chia-Lin Lee, Ming-Hon Yen, Fang-Rong Chang, Chin-Chung Wu, and Yang-Chang Wu

Chiral Resolution and Absolute Configuration of 3 α ,6 β -Dicinnamoyloxytropene and 3 α ,6 β -Di(1-ethyl-1*H*-pyrrol-2-ylcarbonyloxy)tropene, Constituents of *Erythroxylum* Species

Marcelo A. Mu  oz, Solange Arriagada and Pedro Joseph-Nathan

Chromatographic Analysis and Antioxidant Capacity of *Tabernaemontana catharinensis*

Aline A. Boligon, Mariana Piana, Thiago G. Schawnz, Romaiana P. Pereira, Jos   B. T. Rocha and Margaret L. Athayde

Enzyme-treated *Asparagus officinalis* Extract Shows Neuroprotective Effects and Attenuates Cognitive Impairment in Senescence-accelerated Mice

Takuya Sakurai, Tomohiro Ito, Koji Wakame, Kentaro Kitadate, Takashi Arai, Junetsu Ogasawara, Takako Kizaki, Shogo Sato, Yoshinaga Ishibashi, Tomonori Fujiwara, Kimio Akagawa, Hitoshi Ishida and Hideki Ohno

Antifungal Activity of Metabolites from the Marine Sponges *Amphimedon* sp. and *Monanchora arbuscula* against *Aspergillus flavus* Strains Isolated from Peanuts (*Arachis hypogaea*)

Cynthia Arevabini, Yasmin D. Criveleni, Mariana H. de Abreu, Tamires A. Bitencourt, M  rio F. C. Santos, Roberto G. S. Berinck, Eduardo Hajdu, Ren   de O. Belebony, Ana L. Fachin and Mozart Marins

New Guaian-type Sesquiterpene from *Wikstroemia indica*

Mamoru Kato, Yu-Min He, Dya Fita Dibwe, Feng Li, Suresh Awale, Shigetoshi Kadota, and Yasuhiro Tezuka

A New Diterpenoid from the Aerial Parts of *Andrographis paniculata*

Chun-Hua Wang, Wen Li, Rui-Xia Qiu, Miao-Miao Jiang and Guo-Qiang Li

Chemical Composition of the Essential Oil from *Croton kimosorum*, an Endemic Species to Madagascar

Delphin J. R. Rabehaja, Harilala Ihandriharison, Panja A. R. Ramanoelina, Rakotonirina Benja, Suzanne Ratsimamanga-Urverg, Ange Bighelli, Joseph Casanova and F  lix Tomi

Antinociceptive and Anti-edematous Activities of the Essential Oils of Two Balkan Endemic *Laserpitium* Species

Vi  nja Popovi  , Silvana Petrovi  , Maja Tomi  , Radica Stepanovi  -Petrovi  , Ana Micov, Milica Pavlovi  -Drobac, Maria Couladis and Marjan Niketi  

Flavonoids Produced by Tissue Culture of *Dracaena cambodiana*

Hui Wang, Guanyong Luo, Jiayuan Wang, Haiyan Shen, Ying Luo, Haofu Dai and Wenli Mei

A New Dolabellane Diterpenoid and a Sesquiglian from *Aglaia odorata* var. *microphyllina*

Shuai Liu, Yang Wei, Shou-Bai Liu, Hui Wang, Zhi-Kai Guo, Yan-Bo Zeng, Wen-Hua Dong, Wen-Li Mei and Hao-Fu Dai

8-Hydroxycudraxanthone G Suppresses IL-8 Production in SP-C1 Tongue Cancer Cells

Arlette S. Setiawan, Roosje R. Oewen, Supriatno, Willyanti Soewondo, Sidik and Unang Supratman

Determination of Triterpenic Acids and Screening for Valuable Secondary Metabolites in *Salvia* sp. Suspension Cultures

Sibylle K  mmritz, Christiane Haas, Atanas I. Pavlov, Doris Geib, Roland Ulber, Thomas Bley and Juliane Steingroewer

How to Deal with Nomenclatorial Ambiguities of Trivial Names for Natural Products? – A Clarifying Case Study Exemplified for "Corymbosin"

Vatsavaya Ramabharathi and Wolfgang Schuehly

Aporphine Alkaloids of *Cinnamomum molliissimum* and their Bioactivities

Fatin Fasihah Masnon, Najmah PS Hassan and Farediah Ahmad

Modernization of Ayurveda: A Brief Overview of Indian Initiatives

Ambarish Mukherjee, Mousumi Banerjee, Vivekananda Mandal, Amrithesh C Shukla and Subhash C. Mandal

Chemical Compositions and Antimicrobial Activity of the Essential Oils of *Hornstedtia havilandii* (Zingiberaceae)

Ernieyanti Hashim, Hasnah Mohd Sirat and Khong Heng Yen

Anticancer Activity of Binary Toxins from *Lysinibacillus sphaericus* IAB872 against Human Lung Cancer Cell Line A549

Wenjuan Luo, Cuicui Liu, Ruijuan Zhang, Jianwei He and Bei Han

A New Eudesmane Sesquiterpene from *Dichrocephala integrifolia*

Fang Qin, Yi-Bing Wu, Rui-xia Guo, Mei Dong, Fran  oise Sauriol, Qing-Wen Shi, Yu-Cheng Gu and Hiromasa Kiyota

Simultaneous Determination of 13 Chemical Marker Compounds in Gwakhyangjeonggi-san, a Herbal Formula, with Validated Analytical Methods

Jung-Hoon Kim, Hyeun-Kyoo Shin and Chang-Seob Seo

PSY-1, a *Taxus chinensis* var. *mairei* Extract, Inhibits Cancer Cell Metastasis by Interfering with MMPs

Zao-qian Zheng, Ying-Ying Fu, Bo-Heng Li, Mei-Ling Zhang, Xiu-Li Yang, Chuan-wei Xin, Jia-na Shi, Yin Ying and Ping Huang

5-(Hydroxymethyl)-2-furaldehyde Inhibits Adipogenic and Enhances Osteogenic Differentiation of Rat Bone Mesenchymal Stem Cells

Xiang-ling Tan, Yan-Hong Zhang, Jian-Ping Cai, Li-Hua Zhu, Wen-Jie Ge and Xian Zhang

Anti-*L. donovani* Activity in Macrophage/Amastigote Model of Palmarumycin CP₁₈ and its Large Scale Production

Humberto E. Ortega, Eliane de Moraes Teixeira, Ana Rabello, Sarah Higginbotham and Luis Cubilla-R  os

Rosmarinic Acid Interaction with Planktonic and Biofilm <i>Staphylococcus aureus</i> Livia Slobodníková, Silvia Fialová, Helena Hupková and Daniel Grančai	1747
New Butenolide and Pentenolide from <i>Dysidea cinerea</i> Phan Van Kiem, Nguyen Xuan Nhiem, Ngo Van Quang, Chau Van Minh, Nguyen Hoai Nam, Nguyen Thi Cuc, Hoang Le Tuan Anh, Bui HuuTai, Pham Hai Yen, Nguyen Xuan Cuong, Nguyen Phuong Thao, Nguyen Thi Hoai, Nan Young Kim, Seon Ju Park and Kim Seung Hyun	1751
A New Cyclopeptide from Endophytic <i>Streptomyces</i> sp. YIM 64018 Xueqiong Yang, Yabin Yang, Tianfeng Peng, Fangfang Yang, Hao Zhou, Lixing Zhao, Lihua Xu and Zhongtao Ding	1753
Involvement of Trypsin-Digested Silk Peptides in the Induction of RAW264.7 Macrophage Activation Kyoung-Ho Pyo, Min-Ki Kim, Kwang-Soon Shin, Hyang Sook Chun and Eun-Hee Shin	1755
Low-Volatile Lipophilic Compounds in Needles, Defoliated Twigs, and Outer Bark of <i>Pinus thunbergii</i> Alexander V. Shpatov, Sergey A. Popov, Olga I. Salnikova, Ekaterina A. Khokhrina, Emma N. Shmidt and Byung Hun Um	1759
Lipid Constituents of the Edible Mushroom, <i>Pleurotus giganteus</i> Demonstrate Anti-Candida Activity Chia-Wei Phan, Guan-Sern Lee, Ian G. Macreadie, Sri Nurestri Abd Malek, David Pamela and Vikineswary Sabaratnam	1763
Effect of Trehalose Addition on Volatiles Responsible for Strawberry Aroma Mirela Kopjar, Janez Hribar, Marjan Simčič, Emil Zlatič, Tomaž Požrl and Vlasta Piližota	1767
<i>Pogostemon hirsutus</i> Oil, Rich in Abietane Diterpenes Ramar Murugan, Gopal Rao Mallavarapu, Veerappan Sudha and Pemaiah Brindha	1771
Combined Analysis of the Root Bark Oil of <i>Cleistopholis glauca</i> by Chromatographic and Spectroscopic Techniques Zana A. Ouattara, Jean Brice Boti, Coffy Antoine Ahibo, Félix Tomi, Joseph Casanova and Ange Bighelli	1773
Bioactivities and Compositional Analyses of <i>Cinnamomum</i> Essential Oils from Nepal: <i>C. camphora</i>, <i>C. tamala</i>, and <i>C. glaucescens</i> Prabodh Satyal, Prajwal Paudel, Ambika Poudel, Noura S. Dosoky, Kiran Kumar Pokharel, and William N. Setzer	1777
Essential Oil Characterization of Two Azorean <i>Cryptomeria japonica</i> Populations and Their Biological Evaluations Cristina Moiteiro, Teresa Esteves, Luís Ramalho, Rosario Rojas, Sandra Alvarez, Susana Zacchino and Helena Bragança	1785
Antioxidant, Antiproliferative and Antimicrobial Activities of the Volatile Oil from the Wild Pepper <i>Piper capense</i> Used in Cameroon as a Culinary Spice Verlaine Woguem, Filippo Maggi, Hervet P. D. Fogang, Léon A. Tapondjou, Hilaire M. Womeni, Luana Quassinti, Massimo Bramucci, Luca A. Vitali, Dezemona Petrelli, Giulio Lupidi, Fabrizio Papa, Sauro Vittori and Luciano Barboni	1791
<u>Review/Account</u>	
Boldine and Related Aporphines: From Antioxidant to Antiproliferative Properties Darina Muthna, Jana Cmielova, Pavel Tomsik and Martina Rezacova	1797
New Therapeutic Potentials of Milk Thistle (<i>Silybum marianum</i>) Nataša Milić, Nataša Milošević, Ljiljana Suvajdžić, Marija Žarkov and Ludovico Abenavoli	1801
Biomedical Properties of Edible Seaweed in Cancer Therapy and Chemoprevention Trials: A Review Farideh Namvar, Paridah Md. Tahir, Rosfarizan Mohamad, Mahnaz Mahdavi, Parvin Abedi, Tahereh Fathi Najafi, Heshu Sulaiman Rahman and Mohammad Jawaid	1811
Methods for Extraction and Determination of Phenolic Acids in Medicinal Plants: A Review Agnieszka Arceusz, Marek Wesolowski and Pawel Konieczynski	1821

Natural Product Communications

2013

Volume 8, Number 12

Contents

<u>Original Paper</u>	<u>Page</u>
New Humulenes from <i>Hyptis incana</i> (Labiatae) Mitsuru Satoh, Yoshio Satoh, Yasuhiro Anzai, Daisuke Ajisawa, Keiichi Matsuzaki, Mitsuko Makino and Yasuo Fujimoto	1665
Inhibitory Effects against Pasture Weeds in Brazilian Amazonia of Natural Products from the Marine Brown Alga <i>Dictyota menstrualis</i> Rainiomar Raimundo Fonseca, Antonio Pedro Silva Souza Filho, Roberto Campos Villaca and Valéria Laneuville Teixeira	1669
Isolation of the Plant Hormone (+)-Absciscic acid as an Antimycobacterial Constituent of the Medicinal Plant Endophyte <i>Nigrospora</i> sp. Trevor N. Clark, Katelyn Ellsworth, Haoxin Li, John A. Johnson and Christopher A. Gray	1673
New Cembranoid Diterpene from the South China Sea Soft Coral <i>Sarcophyton</i> sp. Fei Cao, Jing Zhou, Kai-Xia Xu, Meng-Qi Zhang and Chang-Yun Wang	1675
Crotofolane Diterpenoids from <i>Croton caracasanus</i> Katuska Chávez, Reinaldo S. Compagnone, Ricarda Riina, Alexander Briceño, Teresa González, Emilio Squitieri, Carlos Landaeta, Humberto Soscún and Alírica I. Suárez	1679
Development and Validation of a Modified Ultrasound-Assisted Extraction Method and a HPLC Method for the Quantitative Determination of Two Triterpenic Acids in <i>Hedyotis diffusa</i> Yu-Chiao Yang, Ming-Chi Wei, Hui-Fen Chiu and Ting-Chia Huang	1683
New Triterpenoid Saponins from the Roots of <i>Saponaria officinalis</i> Barbara Moniuszko-Szajwaj, Łukasz Pecio, Mariusz Kowalczyk, Ana M. Simonet, Francisco A. Macias, Małgorzata Szumacher-Strabel, Adam Cieślak, Wiesław Oleszek and Anna Stochmal	1687
Minor Triterpene Saponins from Underground Parts of <i>Lysimachia thyriflora</i>: Structure elucidation, LC-ESI-MS/MS Quantification, and Biological Activity Irma Podolak, Paweł Zmudzki, Paulina Koczurkiewicz, Marta Michalik, Paweł Zajdel and Agnieszka Galanty	1691
Variation of Saponin Contents and Physiological Status in <i>Quillaja saponaria</i> Under Different Environmental Conditions Angélica Grandón S., Miguel Espinosa B., Darcy Ríos L., Manuel Sánchez O., Katia Sáez C., Víctor Hernández S. and José Becerra A	1697
New Access to 7,17-seco C₁₉-Diterpenoid Alkaloids via Vacuum Pyrolysis of <i>N</i>-Deethyl-8-acetyl Derivatives Ling Wang, Qi-Feng Chen and Feng-Peng Wang	1701
Alkaloids from <i>Boophone haemanthoides</i> (Amaryllidaceae) Jerald J. Nair, Lucie Rárová, Miroslav Strnad, Jaume Bastida and Johannes van Staden	1705
Supinidine Viridiflorates from the Roots of <i>Chromolaena pulchella</i> Mario A. Gómez-Hurtado, J. Martín Torres-Valencia, Rosa E. del Río, Gabriela Rodríguez-García, Virginia Motilva, Sofia García-Mauriño, Carlos M. Cerda-García-Rojas and Pedro Joseph-Nathan	1711
<i>N</i>-Containing Metabolites from the Marine Sponge <i>Agelas clathrodes</i> Fan Yang, Rui-Hua Ji, Jiang Li, Jian-Hong Gan and Hou-Wen Lin	1713
Two New Compounds and Anti-complementary Constituents from <i>Amomum tsao-ko</i> Jiahong Jin, Zhihong Cheng and Daofeng Chen	1715
Antiangiogenic Activity of Flavonoids from <i>Melia azedarach</i> Shigenori Kumazawa, Satomi Kubota, Haruna Yamamoto, Naoki Okamura, Yasumasa Sugiyama, Hirokazu Kobayashi, Motoyasu Nakanishi and Toshiro Ohta	1719
Application of Mixture Analysis to Crude Materials from Natural Resources (IV)^[1(a-c)]: Identification of <i>Glycyrrhiza</i> Species by Direct Analysis in Real Time Mass Spectrometry (II) Eriko Fukuda, Yoshihiro Uesawa, Masaki Baba and Yoshihito Okada	1721
Comparison of Total Phenolic Content, Scavenging Activity and HPLC-ESI-MS/MS Profiles of Both Young and Mature Leaves and Stems of <i>Andrographis paniculata</i> Lee Suan Chua, Ken Choy Yap and Indu Bala Jaganath	1725
Xanthenes from aerial parts of <i>Hypericum laricifolium</i> Juss. Irama Ramírez-González, Juan Manuel Amaro-Luis and Ali Bahsas	1731
A New Xanthone from the Pericarp of <i>Garcinia mangostana</i> Manqin Fu, Samuel X. Qiu, Yujuan Xu, Jijun Wu, Yulong Chen, Yuanshan Yu and Gengsheng Xiao	1733
Isolation of a Phomoxanthone A Derivative, a New Metabolite of Tetrahydroxanthone, from a <i>Phomopsis</i> sp. Isolated from the Mangrove, <i>Rhizophora mucronata</i> Yoshihito Shiono, Takehiro Sasaki, Fumiaki Shibuya, Yukito Yasuda, Takuya Koseki and Unang Supratman	1735
Anti-allergic Inflammatory Effects of Cyanogenic and Phenolic Glycosides from the Seed of <i>Prunus persica</i> Geum Jin Kim, Hyun Gyu Choi, Ji Hyang Kim, Sang Hyun Kim, Jeong Ah Kim and Seung Ho Lee	1739
Isolation, Synthesis and Biological Evaluation of Phenylpropanoids from the Rhizomes of <i>Alpania galanga</i> Sumit S Chourasiya, Eppakayala Sreedhar, K. Suresh Babu, Nagula Shankaraiah, V. Lakshma Nayak, S. Ramakrishna, S. Sravani and M.V. Basaveswara Rao	1741

Continued inside backcover