


NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering all
Aspects of Natural Products Research



**This Issue is Dedicated to
Professor Alejandro F. Barrero
On the Occasion of his 68th Birthday**

Volume 12. Issue 5. Pages 641-802. 2017
ISSN 1934-578X (printed); ISSN 1555-9475 (online)
www.naturalproduct.us

EDITOR-IN-CHIEF

DR. PAWAN K AGRAWAL

Natural Product Inc.
7963, Anderson Park Lane,
Westerville, Ohio 43081, USA
agrawal@naturalproduct.us

EDITORS

PROFESSOR ALEJANDRO F. BARRERO

Department of Organic Chemistry, University of Granada,
Campus de Fuente Nueva, s/n, 18071, Granada, Spain
afbarre@ugr.es

PROFESSOR MAURIZIO BRUNO

Department STEBICEF,
University of Palermo, Viale delle Scienze,
Parco d'Orleans II - 90128 Palermo, Italy
maurizio.bruno@unipa.it

PROFESSOR VLADIMIR I. KALININ

G.B. Elyakov Pacific Institute of Bioorganic Chemistry,
Far Eastern Branch, Russian Academy of Sciences,
Pr. 100-letya Vladivostoka 159, 690022,
Vladivostok, Russian Federation
kalinin@piboc.dvo.ru

PROFESSOR YOSHIHIRO MIMAKI

School of Pharmacy,
Tokyo University of Pharmacy and Life Sciences,
Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan
mimaki@ps.toyaku.ac.jp

PROFESSOR STEPHEN G. PYNE

Department of Chemistry, University of Wollongong,
Wollongong, New South Wales, 2522, Australia
spyne@uow.edu.au

PROFESSOR MANFRED G. REINECKE

Department of Chemistry, Texas Christian University,
Forts Worth, TX 76129, USA
m.reinecke@tcu.edu

PROFESSOR WILLIAM N. SETZER

Department of Chemistry, The University of Alabama in Huntsville,
Huntsville, AL 35809, USA
wsetzer@chemistry.uah.edu

PROFESSOR PING-JYUN SUNG

National Museum of Marine Biology and Aquarium
Checheng, Pingtung 944
Taiwan
pjsung@nmmba.gov.tw

PROFESSOR YASUHIRO TEZUKA

Faculty of Pharmaceutical Sciences, Hokuriku University,
Ho-3 Kanagawa-machi, Kanazawa 920-1181, Japan
y-tezuka@hokuriku-u.ac.jp

PROFESSOR DAVID E. THURSTON

Institute of Pharmaceutical Science
Faculty of Life Sciences & Medicine
King's College London, Britannia House
7 Trinity Street, London SE1 1DB, UK
david.thurston@kcl.ac.uk

HONORARY EDITOR

PROFESSOR GERALD BLUNDEN

The School of Pharmacy & Biomedical Sciences,
University of Portsmouth,
Portsmouth, PO1 2DT U.K.
axuf64@dsl.pipex.com

ADVISORY BOARD

Prof. Giovanni Appendino
Novara, Italy

Prof. Norbert Arnold
Halle, Germany

Prof. Yoshinori Asakawa
Tokushima, Japan

Prof. Vassaya Bankova
Sofia, Bulgaria

Prof. Roberto G. S. Berlinck
São Carlos, Brazil

Prof. Anna R. Bilia
Florence, Italy

Prof. Geoffrey Cordell
Chicago, IL, USA

Prof. Fatih Demirci
Eskişehir, Turkey

Prof. Francesco Epifano
Chieti Scalo, Italy

Prof. Ana Cristina Figueiredo
Lisbon, Portugal

Prof. Cristina Gracia-Viguera
Murcia, Spain

Dr. Christopher Gray
Saint John, NB, Canada

Prof. Dominique Guillaume
Reims, France

Prof. Duvvuru Gunasekar
Tirupati, India

Prof. Hisahiro Hagiwara
Niigata, Japan

Prof. Judith Hohmann
Szeged, Hungary

Prof. Tsukasa Iwashina
Tsukuba, Japan

Prof. Leopold Jirovetz
Vienna, Austria

Prof. Phan Van Kiem
Hanoi, Vietnam

Prof. Niel A. Koorbanally
Durban, South Africa

Prof. Chiaki Kuroda
Tokyo, Japan

Prof. Hartmut Laatsch
Gottingen, Germany

Prof. Marie Lacaille-Dubois
Dijon, France

Prof. Shoei-Sheng Lee
Taipei, Taiwan

Prof. M. Soledade C. Pedras
Saskatoon, Canada

Prof. Luc Pieters
Antwerp, Belgium

Prof. Peter Proksch
Düsseldorf, Germany

Prof. Phila Raharivelomanana
Tahiti, French Polynesia

Prof. Stefano Serra
Milano, Italy

Dr. Bikram Singh
Palampur, India

Prof. Leandros A. Skaltsounis
Zografou, Greece

Prof. John L. Sorensen
Manitoba, Canada

Prof. Johannes van Staden
Scottsville, South Africa

Prof. Valentin Stonik
Vladivostok, Russia

Prof. Winston F. Tinto
Barbados, West Indies

Prof. Sylvia Urban
Melbourne, Australia

Prof. Karen Valant-Vetschera
Vienna, Austria

INFORMATION FOR AUTHORS

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site <http://www.naturalproduct.us>.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the Natural Product Inc. (NPI). A photocopy license is available from the NPI for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

To Subscribe: Natural Product Communications is a journal published monthly. 2017 subscription price: US\$2,595 (Print, ISSN# 1934-578X); US\$2,595 (Web edition, ISSN# 1555-9475); US\$2,995 (Print + single site online); US\$595 (Personal online). Orders should be addressed to Subscription Department, Natural Product Communications, Natural Product Inc., 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

Vibrational Circular Dichroism: Recent Advances for the Assignment of the Absolute Configuration of Natural Products

Eleuterio Burgueño-Tapia^a and Pedro Joseph-Nathan^{b,*}

^a*Departamento de Química Orgánica, Escuela Nacional de Ciencias Biológicas, Instituto Politécnico Nacional, Prolongación de Carpio y Plan de Ayala, Col. Santo Tomás, México D.F., 11340 Mexico*

^b*Departamento de Química, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Apartado 14-740, México D.F., 07000 Mexico*

pjoseph@nathan.cinvestav.mx

Received: May 14th, 2015; Accepted: August 4th, 2015

Reprinted with permission of the Natural Product Inc. Reference: *Natural Product Communications*, 10, 1785-1795 (2015)

Vibrational circular dichroism (VCD) emerged during the last decade as a reliable tool for the absolute configuration (AC) determination of organic compounds. The principles, instrumentation, and methodology applied prior to early 2013 were recently reviewed by us. Since VCD is a very dynamic field, the aim of this review is to update VCD advances for the AC assignment of terpenoids, aromatic compounds, alkaloids, and other natural products for the 2013-2014 period, when VCD was applied to the AC assignment of some 70 natural products. In addition, although discovered in 2012, a brief introduction to the VCD exciton coupling approach and its applications in natural products AC assignment is presented.

Keywords: Natural products, Absolute configuration, Vibrational circular dichroism, Vibrational circular dichroism exciton coupling.

Chirality is the property of handedness, which is probably the most intriguing aspect found in Nature, that goes from the shape of galaxies to the shape of millions of molecules. It allows the existence of life since amino acids that constitute proteins have an *S* absolute configuration (AC), and in our organism these proteins interact differently with the two enantiomers of a given molecule. From here it follows that knowing the AC of organic molecules is fundamental to understanding many aspects of life. Of course, secondary metabolites, biosynthesized by living organism, have their own handedness [1,2].

Although the determination of the AC of natural products has been dominated during the 20th century by circular dichroism of electronic transitions, which was simply known as CD, and is currently known as electronic circular dichroism (ECD), the situation changed approximately at entry into the 21st century when circular dichroism of vibrational transitions gained relevance to perform this task [3]. One of the advantages of vibrational circular dichroism (VCD) over ECD is that electronic transitions that can be measured in the ultraviolet and visible regions are limited to the presence of chromophores, while transitions associated with vibrational modes of molecules, which are measured in the infrared region of the electromagnetic spectrum, are present in any chiral natural product regardless of its simplicity or complexity. An additional advantage of VCD over ECD is that the number of vibrations detected in the infrared region is significantly larger than the number of electronic transitions that can be detected for a given molecule, thus providing much more detailed information. A third advantage of VCD over ECD is that conformational details of molecules are more sensitive to vibrational transitions than to electronic transitions.

Of relevance is also to note that most AC determinations made during the 20th century were based on comparisons with model compounds, which in some cases led to the wrong AC assignment, which of course can be established correctly by VCD, as is the case of esquelande derivatives, which are constituents of a commercial

fragrance [4]. A similar situation was observed for chromane derivatives, for which the relationship between the stereostructure and their ECD was determined almost exclusively by the empirical chromane helicity rule, although failures of the rule have been detected in some cases [5].

VCD has allowed the re-examination of the AC of some natural products like (–)-brevianamide [6] and (+)-schizandrin [7], and it has eventually aided the re-examination and reassignment of the molecular structure of klaivanolide [8]. The VCD superiority over other spectroscopic methods has allowed the structural analysis of complex molecules, like the atropisomer distribution of bridled chiorporphyrins in solution [9], or the planar configuration of nona- and dodecamethoxycryptophanes [10]. It also has been used to determine the secondary structure of NH-indazoles such as (4*S*,7*R*)-campho[2,3-*c*]pyrazole [11]. In addition, VCD has been used for reaction monitoring, and to show solvent induced conformational changes in cyclic peptides [12]. VCD spectra of aminoacids (aa_s), peptides and proteins are difficult to obtain under biological conditions, making it necessary to work with highly concentrated samples to reach acceptable signal intensity, although eventually this is not possible due to low solubility and aggregation phenomena. The VCD signal intensity of aa_s and oligopeptides was enhanced by up to 2 orders of magnitude by coupling them to a paramagnetic metal ion [13].

Since VCD acquired relevance in AC assignments [14-17], efforts have been made in generating virtual spectrometers [18,19], and groups with large resources have systematized conformational analyses, VCD calculation curves, and comparison between experimental and calculated spectra [20].

Misassignments would be rare in VCD since the method is mainly based on the comparison of an experimental spectrum with one based on good level quantum chemical calculations. The determination of the experimental spectrum is a task that requires some 5-10 mg of the natural product, and it consumes typically

5–6 h instrument time, and an additional equal time to measure the solvent spectrum, which has to be subtracted from the spectrum containing the sample. In contrast, the quantum chemical calculation is a quite laborious procedure which can consume from many hours to a few weeks, depending on the number of electrons the sample possesses and the number of conformers which have to be considered. Thus, to illustrate this point, a critical example was a phloroglucinol derivative which shows its sole stereogenic center on a chain appending from an aromatic ring, and which required over 600 h to complete calculations when using a desktop computer operating at 3 GHz with 8 Gb RAM [21]. From here it follows that the limiting time, by far, is the calculation time and not the experimental instrument time.

Fortunately, not all are bad regarding quite long calculation times. It was shown a few years ago that exciton coupling, based on the interaction of two IR chromophores, is a convenient and versatile method to determine AC [22]. Similar to the ECD exciton coupling [23,24], vibrational circular dichroism exciton coupling (VCDEC) [25] is based on the through-space interaction of two or more chromophores which yield a pair of VCD signals with opposite signs around the absorption region of the chromophores. The sign of this split VCD signal reflects the AC of the molecule. Thus, when the interacting chromophores are counterclockwise oriented, the long wave number component of the associated exciton couplet can be expected to exhibit a negative Cotton effect, while when they are clockwise oriented, the long wavelength Cotton effect is positive (Figure 1). For this purpose carbonyl groups have been selected because of their well localized stretching vibration mode that gives rise to electronic transition moments whose direction is virtually parallel to the carbonyl bond. In addition, carbonyl groups can be routinely installed in a desired part of the molecule. Thus, for a chiral molecule containing two chromophores, the AC can be determined without the need of time expensive theoretical calculations.




Figure 1: Relative orientation of two carbonyl groups and amplitude of the VCDEC signals.

The VCDEC approach has been tested for AC assignment of a small number of natural compounds including spiroinducimides A and B [26], a 3-substituted isoindolinone [27], berkeleyamide D

[28], a germacranolide [29], and a series of angular 3'-acyloxypyranocumarins [25] and 3',4'-diacyloxypyranocumarins [30]. The method has also been used for monitoring the enolization of camphor diketones [31].

In a recent review [14], an introduction to the principles, instrumentation, and theoretical quantum chemistry methods for VCD spectra calculation, along with VCD methodology applied to chiral mono-, sesqui-, di-, mero- and triterpenoids, as well as other natural products prior to mid-2013 was provided. Since VCD is currently a very dynamic field, the aim of this review is to update VCD advances for the AC assignment of terpenoids, aromatic compounds, alkaloids, and other natural products for the 2013–2014 period.

Terpenoids

A few mono- and sesquiterpenes have been studied by VCD in the period covered by this review, and of course the most abundant group of studied terpenes is that comprised of diterpenes. Regarding triterpenes, the only studied molecule, already reviewed [14] is a lupeol derivative.

Monoterpenes


VCD is quite sensitive to slight differences in the conformation of chiral molecules and their local environment. A study about the effect of a hydroxy group was addressed using menthol (**1**) as a representative chiral molecule. Comparison of the VCD spectrum with calculations for various rotational conformations of substituents revealed that the 1064–1254 cm^{-1} region sensitively reflects conformational changes. The use of a Boltzmann-weighted analysis to account for the various conformers improved the agreement between the calculated and experimental results, although some discrepancies remained. This sensibility shows the potential application for conformational selective analysis of the local environment of the hydroxy group in chiral molecules and its interaction with neighboring solvent molecules [32].

Although over 150 naturally-occurring thymol derivatives containing a stereogenic center are known [33], (+)-(8*S*)-8,9-epoxy-6-hydroxy-10-benzoyloxy-7-oxothymol isobutyrate (**2**), isolated from *Ageratina cylindrica*, is the only compound whose AC has been assigned. After a Monte Carlo protocol at the MMFF level of theory, followed by complete optimization and harmonic vibrational frequencies calculation using DFT with the B3LYP/DGDZVP basic set and functional, IR and VCD Boltzmann-weighted spectra matched modestly (Table 1). B3PW91 and PBEPBE functionals with the same basis set were therefore used. The enantiomer similarity index increased significantly when using B3PW91. Although the similarity index of IR is smaller, the VCD spectrum is better, and the computer time per conformer is almost one half of that when using the PBEPBE functional [34].

Table 1: Confidence level data for the IR and VCD spectra of **2** calculated using DFT at different levels of theory.

Method	anH ^a	S _{IR} ^b	S _E ^c	S _E ^d	ESI ^e	C' (%)	CPU ^g (min)
B3LYP/DGDZVP	0.972	86.5	66.9	26.0	40.9	79	215
B3PW91/DGDZVP	0.962	92.8	71.6	21.4	20.0	100	266
PBEPBE/DGDZVP	1.002	70.3	72.5	13.4	59.1	100	154


^aAnharmonicity factor. ^bIR spectra similarity. ^cVCD spectra similarity for the correct enantiomer. ^dVCD spectra similarity for the incorrect enantiomer. ^eEnantiomer similarity index calculated as $S_E - S_{E'}$. ^fConfidence level for the stereochemical assignments. ^gJob CPU average time per conformer when using a node with eight processors at 2.93 GHz and 8 Gb of RAM.



Sesquiterpenes

The AC of (+)-3-ishwarone (**3**), isolated from *Peperomia scandens*, was assigned using three chiroptical methods. Based on the reported C1,C2,C9,C11 relative configuration, experimental and calculated ECD, ORD, and VCD spectra of the four possible diastereomers 1*R*,2*S*,4*S*,5*R*,9*R*,11*R*, 1*R*,2*S*,4*R*,5*R*,9*R*,11*R*, 1*R*,2*S*,4*S*,5*S*,9*R*,11*R*, and 1*R*,2*S*,4*R*,5*S*,9*R*,11*R* were evaluated. Comparison of ECD data failed to establish undoubtedly the AC of **3**. The electronic dissymmetry factor (EDF) (ratio of CD to absorption spectra) suggested that the AC of **3** may be 1*R*,2*S*,4*S*,5*R*,9*R*,11*R*, although the conclusion needed support. In turn, ORD data suggested the 1*R*,2*S*,4*S*,5*R*,9*R*,11*R* and 1*R*,2*S*,4*S*,5*R*,9*R*,11*R* diastereomers were feasible. Finally, VCD results provided better similarity for the 1*R*,2*S*,4*S*,5*R*,9*R*,11*R* diastereomer [35].

Addition of diazomethane has been used for many years to prove the presence of an exocyclic methylene group in an α,β -unsaturated lactone. The stereochemistry of this addition was confirmed recently using zaluzanin A (**4**) as a model. The AC of **4** was assured by VCD spectroscopy of its diacetyl derivative and confirmed by X-ray diffraction analysis, while the AC of the diazomethane addition product followed after ^1H NMR chemical shifts evaluation with respect to **4**, and confirmed by X-ray diffraction analysis, including Flack and Hooft parameters AC determination [36].





Diterpenes

The four spiro fused diterpene-indole alkaloids **5-8** were isolated from *Aspergillus terreus*. The relative configuration followed from ROESY experiments. Consequently, the 5*R*,7*S*,8*R*,9*R*,10*R*,13*S*,4*S*,16*S*,22*S*,23*S* AC for **5** and **6**, and the 5*R*,7*S*,8*R*,9*R*,10*R*,3*S*,14*S*,16*S*,22*S* AC for **7** and **8**, was established using a combination of Marfey's method [37] and by comparisons of the DFT B3LYP/6-31G(d,p) calculated and experimental ECD and VCD spectra [38].

The isoneoamphilectane diterpenes **9-11**, strong growth inhibitors of *Mycobacterium tuberculosis*, were isolated from the sponge *Svenzea flava*. Since the 7-formaldehyde group in **11** undergoes *cis/trans* isomerization, it was transformed into its amino derivative **12**. After geometry optimization of **12**, frequency, IR, and VCD calculations, followed by comparison with its experimental VCD spectrum, the 3*S*,4*R*,7*S*,8*S*,11*R*,12*S*,13*R* AC was established [39].

From *Acacia schaffneri*, a plant used to alleviate stomach pain and toothache, the macrocyclic dimeric diterpene **13**, containing a C_2 symmetry axis, was isolated. The structure was established by NMR data, with the HMBC correlation of H-7 with C-17' a key to




assume the dimeric structure. Comparison of its VCD spectrum with that calculated using DFT with the B3LYP/DGDZVP basis set and functional, assigned its preferred conformation and its 5*S*,7*S*,8*R*,9*R*,10*S*,17*S*,5'*S*,7'*S*,8'*R*,9'*R*,10'*S*,17'*S* AC, which was confirmed by evaluation of the Flack and Hooft parameters obtained after single crystal X-ray diffraction analysis [40].

The diterpene content of Chilean *Calceolaria* species up to now comprises 11 abietanes, 6 labdane, 34 pimarane, 11 stemarane, 13 scopadulane, and 6 metabolites having two diterpene units linked as a malonate diester. The AC of these compounds is quite confusing, since abietanes and scopadulanes have been depicted as *normal* diterpenes, while pimaranes, labdanes, and stemaranes have been depicted as *ent* diterpenes. The depicted AC of 13-acetoxyscopadulane (**14**), isolated from several *Calceolaria* species, was established by VCD in combination with DFT calculations, as well as by evaluation of Flack and Hooft single crystal X-ray parameters. It follows that this compound belongs to the normal enantiomeric series of diterpenes, and that at least 11 out of 13 scopadulanes, for which chemical or biogenetical relationships with **14** are established, also belong to the same stereochemical series [41].


Diterpenes **15-17** were isolated from *Solidago shortii*, an endangered species. Its relative 3*R**,4*S**,5*R**,7*S**,8*S**,9*S**,10*S** configuration was assigned from NMR data and analysis of coupling constant values using an energy-minimized model and the Tori equation. In turn, the AC was assigned by comparison of the calculated and experimental VCD spectra. The flexibility of the side chain of **15** resulted in a large number of low-energy conformers, giving a poor confidence level for the calculated spectra. ROESY correlations were therefore used to limit the number of possible conformers and to improve the quality of the calculated spectra. Evaluation of the VCD data using the Compare/OA software supported the 3*R*,4*S*,5*R*,7*S*,8*S*,9*S*,10*S* AC. Furthermore, the calculated OR ($[\alpha]_D +121$) compared well with the experimental ($[\alpha]_D +76$) value and further supports the proposed AC. [42].

A chemical study of the flowers and leaves of *Ageratine jocosotepecana* afforded diterpenoids **18-20**. After evaluation of the



14


15




16

17


reported controversial arguments for the assignment of the C-13 configuration, the coexistence of 13*R* and 13*S*-labdanes, belonging to the *normal* 5*S*,10*S* diterpene series, in the same species, was unexpected and demonstrated by VCD measurements of **19** and **21** in comparison with the DFT B3LYP/DGDZVP calculated spectra. To gain chemical evidence for the configuration at C-13 in **18** and **20**, these natural products were transformed into their respective 8-en methyl ester derivatives. Interestingly, the ^{13}C NMR chemical shifts of **19** and **21** were very similar, revealing the inability of ^{13}C NMR spectroscopy to distinguish these C-13 epimers [43].




18 R
19 H
19 CH₃



20 R
21 H
21 CH₃



22



23

Hyptisolide A (**22**), isolated from *Hyptis crenata* Pohl ex Benth, was the first diterpenoid that had the 7,8;11,12-*bis*-secoabietane skeleton. Its structure and relative stereochemistry were elucidated by HRESIMS, NMR, and X-ray diffraction analyses. The 5*S*,6*S*,10*R* AC of **22** was determined by comparing the experimental VCD spectrum with the calculated spectrum. Although an intense bisignate signal in the stretching carbonyl groups region was observed, no further evaluation was made [44] in terms of VCDEC.


(-)-(1*S*,3*E*,7*R*,8*R*,11*E*)-7,8-Epoxyembrene-3,11-dien-1-ol (**23**), its derived acetate, cembrene A, nephtenol, and cembrenol were isolated from *Burcera multijuga*. The conformational preference of **23** followed from molecular modeling, and its AC from comparison

of DFT B3LYP/DGDZVP calculated and experimental VCD curves [45].


Aromatic compounds

DFT calculations of two conformers of planchol E (**24a** and **24b**) at the DFT B3LYP/6-31+G(d) level of theory showed that the experimental ^1H NMR spectra display the features of **24a** without an intramolecular H-bond, while the measured IR spectrum originated from **24b** has an intramolecular H-bond. Calculated ECD and VCD at the same level of theory showed significant differences between **24a** and **24b**, although no comparison with experimental data was made [46].


The *S*-bridged dimeric pyronaphthoquinone hypogeamicin A (**25**), along with hypogeamicin B (**26**), and other monomeric precursors were isolated from a cave-derived actinomycete *Nonomuraea specus*. The AC of **26** was assigned using ECD, VCD, and ORD spectra and *ab initio* quantum chemical calculations. The four stereoisomers 1*R*,3*R*,4*aS*,10*aR*, 1*S*,3*R*,4*aS*,10*aR*, 1*R*,3*S*,4*aS*,10*aR*, and 1*S*,3*S*,4*aS*,10*aR* were considered. The similarity between the experimental and calculated spectra was determined via overlap. The consensus of quantum mechanical calculations, suggested the AC of (-)-**26** to be 1*R*,3*S*,4*aS*,10*aR*. On the basis of the likely conservation of the epoxide ring opening stereo- and regiochemistry and stereochemical assignment of **26**, the 1*R*,3*R*,4*aR*,10*aS*,10*R*, 3'*R*,4*a'**R*,10*a'**S* AC was proposed for **25** [47].




24a



24b




25



26

(+)-Schizandrin (**27**) is an antihepatotoxic, anti-HIV, and anti-tumor natural product, whose originally described 7*S*,8*S* AC was reassigned as 7*S*,8*R* after comparison of the DFT B3LYP/6-311+G(d) spectra of 7*S*,8*S* and 7*S*,8*R* diastereoisomers with the experimental spectrum. Both experimental and B3LYP/6-31G(d) calculated VCD spectra for (7*S*,8*S*)-**27** look similar in some regions. The major difference between the predicted and experimental VCD spectra is a strong negative signal near 1400 cm⁻¹ in the calculated spectrum, which was assumed to be caused by the use of low level computation sets. Therefore, the B3LYP/6-311+G(d) higher level of theory was used. Unfortunately, the predicted VCD and IR spectra look similar to those obtained at the B3LYP/6-31G(d) level. The B3LYP/6-311+G(d) predicted VCD and IR spectra for (7*S*,8*R*)-**27** look similar to the experimental results. It clearly exhibited that the AC is 7*S*,8*R*. ECD was also used to reach the same conclusion [7].



The first racemic total synthesis, chiral resolution, and AC elucidation of oxalicumone C (**28**) was described [48]. After enantiomeric resolution, determination of the AC was achieved by comparison of measured VCD and ECD spectra with those obtained from *ab initio* calculations. The VCD spectrum of **28** does not match well with the calculated one because of intermolecular hydrogen bonding of the two hydroxy groups. Only the 1050–1225 cm⁻¹ region showed reasonable agreement. To avoid hydrogen-bonding interactions, the hydroxy groups of both enantiomers were silylated to obtain their respective bis(trimethylsilyl) ethers. In the absence of hydrogen bonding, the experimental and calculated VCD spectra showed good agreement over the measured range (1050–1800 cm⁻¹). Nevertheless, the signals between 1600 and 1800 cm⁻¹ in the VCD spectra of **28** and its silylated compound were handled with caution because of birefringence artifacts, and were, therefore, not used to determine the AC. Solvation was treated with the integral equation formalism polarizable continuum model for CH₃CN during the second optimization and subsequent frequency analysis to obtain the vibrational spectra and time-dependent DFT calculations [TD-B3LYP/6-311G(d,p)] for the electronic spectra. Finally, the stereochemistry of the natural product was determined by comparing the optical rotation values from both synthesized enantiomers of **28** {(*S*)-**28** [α]_D = +57.6 (CH₃CN) and (*R*)-**28** [α]_D = -69.7 (CH₃CN)} with the reported data of naturally-occurring **28** {[α]_D = +11.25 (CHCl₃)} [48].

A complex mixture of lomatins (**29**) C-3' esters and (-)-*O*-angeloyllomatins (**30**) were isolated from the seeds of *Prinosciadium thapsoides*. A literature search revealed that some lomatins C-3' monoesters have positive specific rotations, while others had negative values. The mixture was hydrolyzed and resulting **29** was re-esterified to afford (-)-*O*-angeloyl- (**30**), (-)-*O*-acetyl- (**31**), (-)-*O*-isobutyryl- (**32**), (+)-*O*-isovaleroyl- (**33**), (+)-*O*-senecioid- (**34**), (+)-*O*-tigloyl- (**35**), and (+)-*O*-dodecanoyllomatins (**36**). The AC of all molecules was determined as (*R*) from the strong bisignate VCD couplet in the stretching carbonyl group, and by applying the VCDEC approach. In addition to the observed couplet, DFT calculations at the B3LYP/DGDZVP level allowed identification of some vibrational modes of (*R*)-acetylomatins, which show good similarity to all other esterified derivatives, thus validating the VCDEC conclusion [25].

To investigate the scope and limitations of the empirical chromane elicity rule [49], a combination of ECD spectroscopy, especially the


temperature dependence, single crystal X-ray diffraction analysis, and DFT calculations were used in a set of ten (*S*)-trolox (**37**) [(*S*)-6-hydroxy-2,3,7,8-tetramethyl]chromanes. VCD was applied to selected diacetyl derivative **38** because this compound does not show hydrogen bonding and, consequently, the interpretation of the VCD spectrum is facilitated. The spectra were calculated using the B3LYP functional and the 6-31G(d), 6-311++G(d,p) and TZVP basis sets. The best fit of the experimental and calculated spectra was obtained for the TZVP basis. The overall agreement of the predicted and experimental VCD spectra independently supported the configurational and conformational assignment of **38** obtained from ECD studies. The results demonstrated that the chromane helicity rule should always be applied cautiously [5].

Alkaloids


Mycosphaerella sp. fungus, isolated from *Aloe arborescens*, was cultivated in the presence of nicotinamide to afford mycosphines A-D (**39-42**). Although the ¹³C NMR spectrum of **39** resembled that of similin B, the chemical shifts of C-6 differed by 4.3 ppm. The relative configuration of similin B was determined as 2*S**,11*R** by X-ray crystallography, while that of **39** corresponded to the 2*S**,11*S** configuration, indicating that **39** was either the C-2 or the C-11 epimer of similin B. The AC at C-2 and C-11 followed after applying the VCD method using DFT calculations for the acetyl derivative of **39**. While the AC of **40** and **41** was established using biogenetic considerations, the 1*S*,2*S* AC of **42** was determined by the Mosher method [50].

The AC of (+)-caripyrin (**43**) was assigned as 2'*R*,3'*R* by VCD in combination with DFT calculations at the B3LYP/6-311G(d,p) level of theory using the integral equation formalism polarizable continuum model IEFPCM for CCl₄ solvation. IR and VCD for the relevant conformers were presented and discussed [51].

(-)-Berkeleyamide D (**44**), described as a matrix metalloproteinase-3 caspase-1 inhibitor, was synthesized as the racemate. After chiral separation of (+)-**44** the AC was determined using the VCDEC approach. The VCD spectrum of (-)-**44** showed a positive-negative couplet, going from lower to higher wave number, indicating a clockwise orientation of the two adjacent carbonyl groups, and thereby assigning the 5*S*,8*R*,9*R* AC for the natural product [28].




Alkaloids from plants of the genus *Stemona* (family Stemonaceae) typically are characterized by the incorporation of a pyrrolo[1,2-*a*]azepine core. *Stemona* lactam S (**45**) and tuberostemospiroline



(46) were isolated from *S. tuberosa* Lour, a plant used in Chinese traditional medicine [52]. Its structure and AC were established by X-ray and VCD spectroscopy. Three optimized conformers with essentially the same conformation, except for the side ethyl group of 45, and only one conformer for 46, were observed. After geometrical optimization, IR and VCD data were calculated using the DFT B3PW91/DGDZVP2 level of theory. The calculated VCD spectra of (8*R*,9*S*,10*S*,11*R*)-45 and (9*S*,9*aS*,11*R*)-46 were in good agreement with the respective experimental ones. In addition, stemona-amides C-D (47-48), alkaloids having a novel skeleton, were isolated from the same plant [53]. The relative configuration was established by X-ray crystallography and by NOESY experiments, while the AC for 47 and 48 followed after comparison of the experimental VCD spectra with those calculated for the 3*S*,8*R*,9*R*,9*aS*,10*S*,11*R*,12*S*,13*R*,18*S*,20*S* and 3*S*,8*R*,9*R*,9*aS*,10*S*,11*R*,12*S*,13*S*,18*S*,20*S* diastereoisomers, respectively, using DFT at the B3PW91/DGDZVP2 level of theory.

Tadalafil (49), which contains two stereogenic centers, is an approved drug for the treatment of erectile dysfunction, pulmonary arterial hypertension, and benign prostate hyperplasia. It was used to search if VCD allows the determination of the AC without prior knowledge of the relative stereochemistry. Besides VCD, IR and NMR spectra were also used to determine the relative stereochemistry. ECD and ORD spectroscopy, as possible alternative chiral spectroscopic methods to VCD, were used to investigate the complementarity of the three chiroptical techniques. ECD and ORD methods identified the 6*S*,12*aR*/6*R*,12*aS* enantiomeric pair, but for the 6*R*,12*aR*/6*S*,12*aS* pair the AC was uncertain since the lowest-energy conformers have opposite ORD values, and the specific rotation is small. As a result, VCD was clearly the superior discriminatory method for diastereoisomers, and the VCD and NMR combination was optimal [54].

Colchicine (50) is the main agent for the treatment of acute cases of gout, and although it was first isolated in 1820 [55] several unknown aspects remain to be explored. DFT studies of geometry, energy, and ^1H , ^{13}C , and ^{15}N NMR studies were described. In addition, the assignment of both enantiomers by using a chiral solvating agent, and IR and VCD studies of natural anhydrous (-)-50, (-)-50·2H₂O, anhydrous (±)-50, and (±)-50·2H₂O samples were reported. The two achiral samples do not display chiroptical response in CHCl₃ solution, as expected. However, they present a weak chiroptical response in the solid phase, which could be explained by a small enantiomeric excess. After VCD measurement of these four samples, and comparison with the DFT B3LYP/6-311++G(d,p) calculated spectra, natural (-)-colchicine was established as the *P*,7*S* diastereoisomer [56].




Brucine (51) and strychnine (52) are found in several species of the *Strychnos* genus. These alkaloids can be absorbed by the human body to produce stimulating central nervous system activity along with other effects. However, several negative effects have been described, indicating that the demarcation line between these activities is very narrow. DFT calculations at the B3LYP/6-311G++(d,p) level of theory were used to study the equilibrium geometry, vibrational spectra, thermodynamics, and non lineal optical (NLO) properties of 51 and 52. Calculation of the VCD spectra show good agreement with those previously described and indicate that the VCD peaks due to out-of phase stretching modes of aromatic rings and carbonyl stretching modes, in combination with CH stretching modes, are good configuration markers and can be used for AC identification [57].


The AC of azaheterocycles 53-55, resulting from the cascade rearrangement of enediynes, were assigned by the joint use of VCD and Cu X-ray diffraction using the Flack and Hooft parameters, thus demonstrating that the rearrangement of enediynes proceeded with retention of configuration. IR and VCD were calculated using DFT with the B3LYP and 6311+G (d,p) functional and basic set. Solvent effects (CH₂Cl₂) were introduced using the polarizable continuum model IEF-PCM combined with SMD [58] quantum mechanical continuum solvation parameters [59].

Diagnostic VCD bands of tropane alkaloids from *Erythroxylaceae* species can be related to the AC without the need to perform DFT calculations. Specifically, the *-/-/+/-* pattern has been observed in the VCD 1100-950 cm⁻¹ region of compounds which possess the 3*R*,6*R* AC, while the antipodal *+/+/-/-* pattern has been observed in compounds which possess the 3*S*,6*S* AC, independently of the ester identity on each hydroxy group. Racemic samples of 56 and 57 were resolved by chiral HPLC. The first eluate of 57 shows [α] = -36.2 (EtOH), in agreement with that reported for natural catuabine E ([α] = -35.4). For (-)-56 and (-)-57, four main bands with the *-/-/+/-* pattern, going from low to high wave numbers, were observed, while isomers (+)-56 and (+)-57 showed the opposite *+/+/-/-* pattern. Following this empirical rule, the AC of each isolated enantiomer was established as (-)-(3*R*,6*R*)-56, (+)-(3*S*,6*S*)-56, (-)-(3*R*,6*R*)-57, and (+)-(3*S*,6*S*)-57 [60].

Cycloaspeptide G (58) is a pentapeptide isolated from the fungus *Isaria farinosa*, which displays an excellent cytotoxic effect against HeLa and MCF7 cell lines. Although its AC was determined by spectroscopic techniques, its conformational characteristics were explored using DFT. Four conformers were considered, and after




DFT B3LYP/6-31G geometry optimization the low energy conformer was considered to study the ECD and VCD spectra. The computations revealed the B3LYP/6-31G(d) level is preferred over the B3LYP/6-31G level. Some NH, and carbonyl stretching modes, and CH bending modes were assigned and discussed at the DFT B3LYP/6-31G and B3LYP/6-31G(d) levels of theory [61].



The epipolythiodiketopiperazine (ETP) alkaloids are a large family of fungal secondary metabolites with very interesting cytotoxic activities. The isolation, structural determination, AC assignments, and conformational analyses of two ETPs, preussiadsins A (**59**) and B (**60**) from *Preussia tyoharum* were described. Assignment of the AC of **59** was initially hindered when all attempts to crystallize the compound and its derivatives failed. Based on the proposed 3*S**,5*aR**,10*bS**,11*S**,12*S**,3'*R**,5'*aS**,10'*bR**,11'*S**,12'*R** AC, quantum chemical computational calculations at the DFT B3LYP/6-31+G(d,p) level were carried out to generate VCD and ECD spectra, as well as a specific rotation value for **59**. In both cases, the calculated VCD and ECD spectra matched well with the experimental data. The calculated specific rotation value of **59** ($[\alpha]_D^{+53}$) was in agreement with the experimental observations ($[\alpha]_D^{+66}$). Therefore, the AC of **59** followed as 3*S**,5*aR**,10*bS**,11*S**,12*S**,3'*R**,5'*aS**,10'*bR**,11'*S**,12'*R*. The proposed structure and AC assignment for **60** was determined by *S*-methylation. This provided three products that were confirmed as being identical to the products of the cleavage/methylation reaction of **59**. Thus **59** and **60** share the same AC [62].

Other natural products


Cultivation of the filamentous fungus *Chaetomium indicem* yield spiroindocumines A (**61**), and B (**62**) as minor constituents. Their AC could not be deduced by Mosher derivatization, and was



therefore assigned using the VCDEC approach. Although the VCD spectra of **61** and **62**, at a 0.05 M concentration, showed small signals with no significant peaks above 1700 cm⁻¹, in their respective diacetates, **63** and **64**, a strong bisignate VCD signal with a positive-negative couplet, from lower to higher wave number, that suggested a clockwise orientation between the adjacent carbonyl groups at C1' and AcO-2', was observed using a 0.01 M solution, allowing it to be concluded that C2' is *R*, and the entire AC are (2'*R*,6*S*,7*S*)-**61** and (2'*R*,6*S*,7*R*)-**62**. These AC assignments represent the first application of the VCDEC approach to novel natural products [26].

Typically ECD provides dependable data for the AC assignment and is more sensitive than VCD. Comparison of simulated and measured VCD and ECD spectra of natural dioxolanes **65**–**70** from *Guignardia bedwelli*, and semisynthetic derivatives shows the superiority of VCD over ECD for the AC determination of these compounds. After conformation analysis, UV and ECD spectra were obtained as Boltzmann-weighted averages. The calculated ECD spectrum of **65** did not match the experimental data, although the calculated UV spectrum showed a reasonable agreement with the recorded absorption bands. The VCD spectrum of **65** reasonably compared with a B3LYP/6-311G(d,p) calculated spectrum considering CDCl₃ solvation. However, rotational strengths at 1251 and 1207 cm⁻¹, along with frequencies of the C=O stretching of the COOH group are not correctly predicted. This was attributed to the known aggregation effect of carboxylic acids in nonpolar solvents, which is difficult to represent *in silico*, while measurements in DMSO-*d*₆ did not improve this situation. Compound **65** was therefore transformed into its methyl ester, whereby the agreement between the predicted VCD spectrum and the experimental data was much better. A similar situation was observed for **66**, which was also transformed into its methyl ester. The naturally occurring esters **67**–**70** showed highly similar VCD spectra, each with a characteristic triplet consisting of a positive band around 1282 cm⁻¹ [$\nu_{as}(\text{C2}–\text{C6}–\text{O})$] and two negative bands around 1247 cm⁻¹ [$\nu(\text{C4}–\text{C5}) + \nu(\text{C1}'–\text{C8})$] and 1170 cm⁻¹ [$\nu(\text{C4}–\text{C5}) + \nu(\text{C1}'–\text{C8})$]. This constant pattern is predicted with high accuracy, excepting the rotational strengths of the lowest-frequency bands in **67** and **68**, which are overestimated by DFT. These results indicate all investigated compounds have the *S* AC [63].

From *Curcuma longa* L., a popular plant of Chinese herbal medicine, fourteen cytotoxic terpene-conjugated curcuminoids J–W were isolated. After relative configuration NMR assignments, the AC was established by ECD and confirmed for terpecurcumin **S** (**71**) by VCD using the DFT B3LYP/6-31G(d) level of theory [64].



The construction of well-arranged π -conjugated chromophores has attracted much attention since the relative orientations and distances of the chromophores are crucial in determining their photophysical properties. The tubular compound **72**, in which three π -conjugated chromophores are connected, was synthesized as a diastereomeric mixture with planar chirality, a pair of enantiomers and a *meso* compound. Each compound was separated by chiral HPLC. The inherent chirality of *m*-calex[3]amide (**72**) induced a predominantly one-handed helicity to the arrangement of π -conjugated chromophores, without the help of a chiral guest or a stereogenic center. The AC of *m*-calex[3]amide, and the preferred helicity of the bithiophene unit were determined by combined experimental and theoretical studies. The ECD and VCD analyses were mutually complementary in adding the interpretation of the chiral structure in the present system; ECD spectroscopy was sensitive to the helicity of the bithiophene chromophores, whereas VCD spectroscopy was sensitive to the planar chirality of *m*-calex[3]amide [65].


The dihydroxylated macrolide **73**, along with ten known compounds were isolated from a solid culture of the endophytic fungus *Pestalotiopsis manguiifera*. Structural elucidation and relative configuration followed after spectroscopic data including coupling constant values and NOESY evaluation. The 4*R*,7*R*,8*R*,9*S* AC of **73** was determined by VCD at the DFT B3LYP/6-31G(d) level [66].

Methyl mandelate (**74**) is a relatively small chiral α -hydroxyester which undergoes intra- and intermolecular hydrogen bonding interactions, and possesses a phenyl ring that influences the closeness and orientation of solvent molecules. It was selected to establish a general strategy to account for solvent effects. A comparative VCD study of **74** in methanol, dimethyl sulfoxide, and chloroform, using explicit and implicit solvent models, was performed. The results showed that the simultaneous inclusion of explicitly and implicitly solvent has a significant impact on the appearance of the vibrational absorption and VCD spectra, and is crucial for reliable spectral assignments when solvents are capable of hydrogen-bonding with solutes. When no strong solute solvent hydrogen-bonding interactions exist, like in chloroform, the gas phase monomer model is adequate for spectra interpretation, while inclusion of implicit solvation improves the frequency agreement with the experimental data [67].


Phyllostin (**75**), seytolide (**76**), and oxysporone (**77**) are promising natural herbicides for the weed biocontrol of some *Orabanche* species. Supported by the knowledge of the relative configuration, the combined application of ORD, ECD, and VCD approaches permitted the assignment of the 3*S*,4*aR*,8*S*,8*aR* AC to (–)-**75**. In **76**, theoretical analysis of both electronic and vibrational CD spectra provided consistent results, while ORD was found unsuitable. Therefore, in the case of (–)-**76**, only two of the used chiroptical methods allowed the 4*aR*,8*S*,8*aR* AC assignment. The good agreement between the experimental and calculated ORD and ECD spectra of **77** led to the AC assignment. Taking into account solvent effects, all calculated VCD bands gave a satisfactory agreement with the experimental spectrum. Consequently, all three chiroptical methods support the AC assignment of (+)-**77** as 4*S*,5*R*,6*R*. This study suggests that for flexible molecules, a concerted application of more than one chiroptical methodology should be considered [68].

The configuration complexity of oligostilbenoids increases with their degree of polymerization. Four dimeric stilbene glucosides, two diastereoisomers of (–)-gnemonoside A (**78** and **79**), (–)-gnemonoside C (**80**), and (–)-gnemonoside D (**81**), as well as a mixture of the enantiomers of gnetin C (**82**, **83**) were isolated from the rhizomes of *Gnetum africanum*. ¹H and ¹³C NMR spectra of **78** and **79** were superimposable, and similar to that reported for gnemonoside A. VCD spectra of **78** and **79** are not perfectly opposite, revealing that they are not enantiomers, but could be diastereomers. After enzymatic hydrolysis, **78** and **79** gave aglycones (–)-**83** and (+)-**82**, respectively. Their AC was assigned by VCD experiments in combination with calculations and ¹H NMR experiments. The VCD data indicated that the 7*aS*,8*aS* enantiomer of gnetin C is slightly predominant in natural gnetin C. Interestingly, all the bands observed in the VCD spectra of **80** and **81** had the same sign as those of compound **78**, but their intensities were slightly lower. The 7*aS*,8*aS* configuration was preserved for the stilbene dimers **78**, **80**, and **81**. [69].

Klaivanolide, originally assumed as **84**, was presented as a promising antileishmanial agent from *Uvaria klaineana*, whose AC determination was made by VCD spectroscopy. The difficulties experienced in the synthesis led to revise and re-assign its structure as the butyrolactone acetylmelodorinol **85**. The calculated VCD spectra of the originally assigned structure for klaivanolide and the revised structure known as acetylmelodorinol are similar. This observation was rationalized by the authors considering that the VCD phenomenon ensues from IR absorption properties of functional groups that are identical in both compounds, and are strongly correlated with their relative arrangement around the stereogenic center, which remains as *S* in both structures [8].



Hemicalide (**86**) is a highly bioactive marine natural product for which the relative configuration of the C8–C13 fragment and of the C18–C24 α,β -dihydroxy- δ -lactone subunit have been described. The relative configuration of the C36–C42 subunit was assigned by combining stereocontrolled synthesis with NMR, IR, and VCD analysis. Diastereomeric model compounds corresponding to the C36–C46 subunit were synthesized for ^1H and ^{13}C NMR data comparison with those of **86**. An attempt to determine the relative configuration at C42 by a statistical analysis of calculated and measured NMR chemical shifts was made. However, despite the seemingly high probabilities calculated, different conclusions were reached based on the ^{13}C and ^1H spectra. To make an unambiguous assignment of the relative configuration, a VCD analysis was performed for **87** and **88**. The AC at C42 in these two key epimeric compounds was particularly challenging for VCD analysis due to the large number of conformations and the presence of six stereocenters. The commonly used 6-31G(d) basis set overestimated the abundance of conformers showing intramolecular H-bonding. The use of the aug-cc-p-VDZ calculation level with a polarizable



continuum model gave a more realistic Boltzmann distribution. Although the use of VCD spectroscopy can be sufficient to assign the AC of epimers containing multiple stereocenters, spectra subtraction of the epimers, for the calculations and the experiments, greatly enhanced the power for epimer distinction. The AC determination at C42 in **87** and **88** allowed the assignment of the relative configuration of the C36–C42 subunit [70].

In summary, several reviews [3,14-17], and a just published one [71], show VCD is currently a superior and preferred technique for the AC determination of natural products. It accounts for some 70 additional natural products reviewed herein for less than the last two years period. The easy empirical interpretation of the VCDEC bisignated couplet will certainly attract more users to this methodology since thereby no long and tedious DFT calculations are required.

Acknowledgments - Partial financial support from CONACYT-Mexico (Grants 168066 and 152994), and SIP-IPN (Grant 20151556) is acknowledged.

References

- Peterson D, Schnell M, Doyle JM. (2013) Enantiomer-specific detection of chiral molecules via microwave spectroscopy. *Nature*, **497**, 475-478.
- Nafie LA. (2013) Handedness detected by microwaves. *Nature*, **497**, 446-448.
- Polavarapu PL. (2012) Determination of the structure of chiral natural products using vibrational circular dichroism. In *Comprehensive Chiroptical Spectroscopy. Applications in Stereochemical Analysis of Synthetic Compounds, Natural Products, and Biomolecules*; Berova N, Polavarapu PL, Nakanishi K, Woody RW. (Eds). Wiley, Hoboken. Vol. 2, Chapter 11, 387-420.
- Cerda-García-Rojas CM, Bucio MA, González SB, García-Gutiérrez HA, Joseph-Nathan P. (2015) Absolute configuration of esquelane derivatives from *Adesmia boronioides* by vibrational circular dichroism. *Tetrahedron: Asymmetry*, **26**, 136-140.
- Gorecki M, Suszczynska A, Woznica M, Baj A, Wolniak M, Cyranski MK, Witkowski S, Frelek J. (2014) Chromane helicity rule - scope and challenges based on an ECD study of various trolox derivatives. *Organic & Biomolecular Chemistry*, **12**, 2235-2254.
- Ren J, Li GY, Shen L, Zhang GL, Nafie LA, Zhu HJ. (2013) Challenges in the assignment of relative and absolute configurations of complex molecules: computation can resolve conflicts between theory and experiment. *Tetrahedron*, **69**, 10351-10356.
- He P, Wang X, Guo X, Ji Y, Zhou C, Shen S, Hu D, Yang X, Luo D, Dukor R, Zhu H. (2014) Vibrational circular dichroism study for natural bioactive schizandrin and reassignment of its absolute configuration. *Tetrahedron Letters*, **55**, 2965-2968.
- Ferrie L, Ferhi S, Bernadat G, Figadere B. (2014) Toward the total synthesis of klaivanolide: Complete reinterpretation of its originally assigned structure. *European Journal of Organic Chemistry*, **2014**, 6183-6189.
- Castaigns A, Marchon JC, Cavagnat D, Buffeteau T. (2013) Conformational equilibria of bridled chirophorophyrins in solution investigated by vibrational circular dichroism. *Chirality*, **25**, 480-486.
- Brotin T, Vanthuyne N, Cavagnat D, Ducasse L, Buffeteau T. (2014) Chiroptical properties of nona- and dodecamethoxy cryptophanes. *Journal of Organic Chemistry*, **79**, 6028-6036.
- Quesada-Moreno MM, Avilés-Moreno JR, López-González JJ, Claramunt RM, López C, Alkorta I, Elguero J. (2014) Chiral self-assembly of enantiomerically pure (4*S*,7*R*)-campho[2,3-*c*]pyrazole in the solid state: a vibrational circular dichroism (VCD) and computational study. *Tetrahedron: Asymmetry*, **25**, 507-515.
- Merten C, Li F, Bravo-Rodríguez K, Sánchez-García E, Xu Y, Sander W. (2014) Solvent-induced conformational changes in cyclic peptides: a vibrational circular dichroism study. *Physical Chemistry Chemical Physics*, **16**, 5627-5633.

- [13] Domingos SR, Huerta-Viga A, Baij L, Amirjalayer S, Dunnebie DAE, Walters AJC, Finger M, Nafie LA, De Bruin B, Buma WJan, Woutersen S. (2014) Amplified vibrational circular dichroism as a probe of local biomolecular structure. *Journal of the American Chemical Society*, **136**, 3530-3535.
- [14] Joseph-Nathan P, Gordillo-Román B. (2015) Vibrational circular dichroism absolute configuration determination of natural products. In *Progress in the Chemistry of Organic Natural Products*. Vol. 100, Kinghorn AD, Falk H, Kobayashi J. (Eds.). Springer International Publishing, Switzerland, 311–451.
- [15] Batista Jr. JM. (2013) Determination of absolute configuration using chiroptical methods. In *Stereoselective Synthesis of Drugs and Natural Products*. Andrushko V, Andrushko N. (Eds). Wiley, Hoboken. Vol. 2, Chapter 53, 1571-1599.
- [16] Batista Jr JM, da Silva BV. (2014) Determination of the absolute configuration of natural product molecules using vibrational circular dichroism. In *Studies in Natural Products Chemistry*. Vol. 41, Atta-ur-Rahman (Ed). Elsevier, Amsterdam, 311–451.
- [17] Kong LY, Wang P. (2013) Determination of the absolute configuration of natural products. *Chinese Journal of Natural Medicines*, **11**, 193-198.
- [18] Egidi F, Bloino J, Cappelli C, Barone V. (2013) Development of a virtual spectrometer for chiroptical spectroscopies: the case of nicotine. *Chirality*, **25**, 701-708.
- [19] Barone V, Baiardi A, Bloino J. (2014) New developments of a multifrequency virtual spectrometer: stereo-electronic, dynamical, and environmental effects on chiroptical spectra. *Chirality*, **26**, 588-600.
- [20] Sherer EC, Lee CH, Shpungin J, Cuff JF, Da C, Ball R, Bach R, Crespo A, Gong X, Welch CJ. (2014) Systematic approach to conformational sampling for assigning absolute configuration using vibrational circular dichroism. *Journal of Medicinal Chemistry*, **57**, 477-494.
- [21] Casero C, Machín F, Méndez-Álvarez S, Demo M, Ravelo AG, Pérez-Hernández N, Joseph-Nathan P, Estevez-Braun A. (2015) Structure and antimicrobial activity of phloroglucinol derivatives from *Achyrocline satureioides*. *Journal of Natural Products*, **78**, 93-102.
- [22] Taniguchi T, Monde K. (2012) Exciton chirality method in vibrational circular dichroism, *Journal of the American Chemical Society*, **134**, 3695-3698.
- [23] Harada N, Nakanishi K, Berova N. (2012) Electronic CD exciton chirality method: Principles and applications. In *Comprehensive Chiroptical Spectroscopy. Applications in Stereochemical Analysis of Synthetic Compounds, Natural Products, and Biomolecules*. Berova N, Polavarapu PL, Nakanishi K, Woody RW. (Eds). Wiley, Hoboken. Vol. 2, Chapter 4, 115-166.
- [24] Lightner DA, Gurst JE. (2000) Exciton Coupling and Exciton Chirality. In *Organic Conformational Analysis and Stereochemistry from Circular Dichroism Spectroscopy*. Wiley-VCH, New York. Chapter 14, 423-456.
- [25] Buendia-Trujillo IA, Torres-Valencia JM, Joseph-Nathan P, Burgueño-Tapia E. (2014) The absolute configuration of angular 3'-acyloxypyranocoumarins by vibrational circular dichroism exciton chirality. *Tetrahedron: Asymmetry*, **25**, 1418-1423.
- [26] Asai T, Taniguchi T, Yamamoto T, Monde K, Oshima Y. (2013) Structures of spiroinducumides A and B, unprecedented carbon skeletal spiroactones, and determination of the absolute configuration by a vibrational circular dichroism exciton approach. *Organic Letters*, **15**, 4320-4323.
- [27] Massa A, Rizzo P, Monaco G, Zanasi R. (2013) Absolute configuration assignment made easier by the VCD of coupled oscillating carbonyls: the case of (–)-propanedioic acids, 2-(2,3)-dihydro-3-oxo-1H-isoindol-1-yl)-1,3-dimethyl ester. *Tetrahedron Letters*, **54**, 6242-6246.
- [28] Komori K, Taniguchi T, Mizutani S, Monde K, Kuramochi K, Tsubaki K. (2014) Short synthesis of berkeleyamide D and determination of the absolute configuration by the vibrational circular dichroism exciton chirality method. *Organic Letters*, **16**, 1386-1389.
- [29] Sánchez-Castellanos M, Bucio MA, Hernández-Barragán A, Joseph-Nathan P, Cuevas G, Quijano L. (2015) Vibrational circular dichroism (VCD), VCD exciton coupling, and X-ray determination of the absolute configuration of an α,β -unsaturated germacranolide. *Chirality*, **27**, 247-252.
- [30] Buendia-Trujillo IA, Torres-Valencia JM, Joseph-Nathan P, Burgueño-Tapia E. (2015) Absolute configuration assignment of 3',4'-di-O-acylkhellactones using vibrational circular dichroism exciton chirality. *Natural Product Communications*, **10**, 1027-1032.
- [31] Wu T, You X. (2012) Exciton coupling analysis and enolization monitoring by vibrational circular dichroism spectra of camphor diketones. *The Journal of Physical Chemistry, A*, **116**, 8959-8964.
- [32] Konno K, Shiina I, Yui H. (2013) Fine structures in vibrational circular dichroism spectra of chiral molecules with rotatable hydroxyl groups and their application in the analysis of local intermolecular interactions. *Journal of Molecular Structure*, **1035**, 260-266.
- [33] Talavera-Alemán A, Rodríguez-García G, López Y, García-Gutiérrez HA, Torres-Valencia JM, del Río RE, Cerda-García-Rojas CM, Joseph-Nathan P, Gómez-Hurtado MA. (2015) Systematic evaluation of thymol derivatives possessing stereogenic centers. *Phytochemistry Reviews*, DOI:10.1007/s11101-015-9412-6.
- [34] Bustos-Brito C, Sánchez-Castellanos M, Esquivel B, Calderón JS, Calzada F, Yopez-Mulia L, Hernández-Barragán A, Joseph-Nathan P, Cuevas G, Quijano L. (2014) Structure, absolute configuration, and antidiarrheal activity of a thymol derivative from *Ageratina cylindrica*. *Journal of Natural Products*, **77**, 358-363.
- [35] Junior FMS, Covington CL, De Amorim MB, Velozo LSM, Kaplan MAC, Polavarapu PL. (2014) Absolute configuration of a rare sesquiterpene: (+)-3-ishwarone. *Journal of Natural Products*, **77**, 1881-1886.
- [36] Ortiz-León A, Torres-Valencia JM, Manríquez-Torres JJ, Alvarado-Rodríguez JG, Hernández-Balderas U, Cerda-García-Rojas CM, Joseph-Nathan P. (2014) Diastereoselective addition of diazomethane to zaluzanin A. *Natural Product Communications*, **9**, 753-756.
- [37] Brushan R, Brückner H. (2004) Marfey's reagent for chiral amino acid analysis: A review. *Amino acids*, **27**, 231-147.
- [38] Cai S, Du L, Gereá AL, King JB, You J, Cichewicz RH. (2013) Spiro fused diterpene-indole alkaloids from a creek-bottom-derived *Aspergillus terreus*. *Organic Letters*, **15**, 4186-4189.
- [39] Avilés E, Rodríguez AD, Vicente J. (2013) Two rare-class tricyclic diterpenes with antitubercular activity from the Caribbean sponge *Svenzea flava*. Application of vibrational circular dichroism spectroscopy for determining absolute configuration. *Journal of Organic Chemistry*, **78**, 11294-11301.
- [40] Manríquez-Torres JJ, Torres-Valencia JM, Velázquez-Jiménez R, Valdez-Calderón A, Alvarado-Rodríguez JG, Cerda-García-Rojas CM, Joseph-Nathan P. (2013) A macrocyclic dimeric diterpene with a C2 symmetry axis. *Organic Letters*, **15**, 4658-4661.
- [41] Muñoz MA, Chamy C, Bucio MA, Hernández-Barragán A, Joseph-Nathan P. (2014) Absolute configuration of scopadulane diterpenes from *Calceolaria* species. *Tetrahedron Letters*, **55**, 4274-4277.
- [42] Williams RB, Du L, Norman VL, Goering MG, O'Neil-Johnson M, Woodbury S, Albrecht MA, Powell DR, Cichewicz RH, Eldridge GR, Starks CM. (2014) Diterpenes from the endangered goldenrod *Solidago shortii*. *Journal of Natural Products*, **77**, 1438-1444.
- [43] García-Sánchez E, Ramírez-López CB, Talavera-Alemán A, León-Hernández A, Martínez-Muñoz RE, Martínez-Pacheco MM, Gómez-Hurtado MA, Cerda-García-Rojas CM, Joseph-Nathan P, del Río RE. (2014) Absolute configuration of (13R)- and (13S)-labdane diterpenes coexisting in *Ageratina jocotepecana*. *Journal of Natural Products*, **77**, 1005-1012.
- [44] Yun YS, Fukaya H, Nakane T, Takano A, Takahashi S, Takahashi Y, Inoue H. (2014) A new bis-*seco*-abietane diterpenoid from *Hyptis crenata* Pohl ex Benth. *Organic Letters*, **16**, 6188-6191.
- [45] Hernández-Hernández JD, García-Gutiérrez HA, Román-Marín LU, Torres-Blanco YI, Cerda-García-Rojas CM, Joseph-Nathan P. (2014) Absolute configuration of cembrane diterpenoids from *Burcera multijuga*. *Natural Product Communications*, **9**, 1249-1252.

- [46] Yang Y, Liu G, Lei T, Mang C. (2013) Density functional study of chiral structures and spectral properties of the phenolic compound planchol E from pinecones of *Pinus yunnanensis*. *Dali Xueyuan Xuebao*, **12**, 40-43.
- [47] Derewacz DK, McNeese CR, Scalmani G, Covington CL, Shanmugam G, Marnett LJ, Polavarapu PL, Bachmann BO. (2014) Structure and stereochemical determination of hypogaeamicins from a cave-derived actinomycete. *Journal of Natural Products*, **77**, 1759-1763.
- [48] Wink C, Andernach L, Opatz T, Waldvogel SR. (2014) Total synthesis of (±)-oxalicumone C and chiral resolution and elucidation of its absolute configuration. *European Journal of Organic Chemistry*, **2014**, 7788-7792.
- [49] Kurtán T, Antus S, Pescitelli G. (2012) Electronic cd of benzene and other aromatic chromophores for determination of absolute configuration. In *Comprehensive Chiroptical Spectroscopy. Applications in Stereochemical Analysis of Synthetic Compounds, Natural Products, and Biomolecules*; Berova N, Polavarapu PL, Nakanishi K, Woody RW. (Eds). Wiley, Hoboken. Vol. **2**, Chapter 3, 73-114.
- [50] Asai T, Otsuki S, Taniguchi T, Monde K, Yamashita K, Sakurai H, Ozeki T, Oshima Y. Structures and absolute configurations of short-branched fatty acid dimers from an endophytic fungus of *Aloe arborescens*. *Tetrahedron Letters*, **54**, 3402-3405.
- [51] Andernach L, Opatz T. (2014) Assignment of the absolute configuration and total synthesis of (+)-caripyrin. *European Journal of Organic Chemistry*, **2014**, 4780-4784.
- [52] Fukaya H, Hitotsuyanagi Y, Aoyagi Y, Shu Z, Komatsu K, Takeya K. (2013) Absolute structures of stemona-lactam S and tuberostemospirone, alkaloids from *Stemona tuberosa*. *Chemical & Pharmaceutical Bulletin*, **61**, 1085-1089.
- [53] Hitotsuyanagi Y, Shigemori G, Fukaya H, Hikita M, Zhu S, Komatsu K, Takeya K. (2013) Stemona-amines C-E, new alkaloids from *Stemona tuberosa*. *Tetrahedron Letters*, **54**, 6995-6998.
- [54] Qiu S, De Gussem E, Tehrani KA, Sergeyev S, Bultinck P, Herrebout W. (2013) Stereochemistry of the tadalafil diastereoisomers: A critical assessment of vibrational circular dichroism, electronic circular dichroism, and optical rotatory dispersion. *Journal of Medicinal Chemistry*, **56**, 8903-8914.
- [55] Pelletier PJ, Caventou JB. (1820) Examen chimique de plusieurs végétaux de la famille des colchiciées, et du principe actif qu'ils renferment. [Cévadille (*Veratrum sabadilla*); hellébore blanc (*Veratrum album*); colchique commun (*Colchicum autumnale*)]. *Annales de chimie et physique*, **114**, 69-83.
- [56] Virgili A, Quesada-Moreno MM, Avilés-Moreno JR, López-González JJ, García MA, Claramunt RM, Torres MR, Jimeno ML, Reviriego F, Alkorta I, Elguero J. (2014) A spectroscopic study of colchicine in the solid state and in solution by multinuclear magnetic resonance and vibrational circular dichroism. *Helvetica Chimica Acta*, **97**, 471-490.
- [57] Islam N, Niaz S, Manzoor T, Pandith AH. (2014) Theoretical investigations into spectral and non-linear optical properties of brucine and strychnine using density functional theory. *Spectrochimica acta. Part A, Molecular and biomolecular spectroscopy*, **131**, 461-70.
- [58] Marenich AV, Cramer CJ, Truhlar DG. (2009) Universal solvation model based on solute electron density and on continuum model of the solvent defined by the bulk dielectric constant and atomic surface tensions. *The Journal of Physical Chemistry B*, **113**, 6378-6396.
- [59] Mondal S, Naubron JV, Campolo D, Giorgi M, Bertrand MP, Nechab M. (2013) Cooperative use of VCD and XRD for the determination of tetrahydrobenzoisquinolines absolute configuration: A reliable proof of memory of chirality and retention of configuration in enediyne rearrangements. *Chirality*, **25**, 832-839.
- [60] Muñoz MA, Arriagada S, Joseph-Nathan P. (2014) Chiral resolution and absolute configuration of 3*α*,6*β*-dicinnamoyloxypitropane and 3*α*,6-di(1-ethyl-1*H*-pyrrol-2-ylcarbonyloxy)pitropane, constituents of *Erythroxylum* species. *Natural Product Communications*, **9**, 27-30.
- [61] Mang CY, Zhao Y, Li HF, Lan H, Yan Y, Yang MH. (2014) Structural characteristics and spectral behaviours of the preferred conformation of a cyclic pentapeptide, cycloaspeptide G from *Cordyceps*-colonising fungus *Isaria farinosa*. *Molecular Physics*, **116**, 104-112.
- [62] Du L, Robles AJ, King JB, Mooberry SL, Cichewicz RH. (2014) Cytotoxic dimeric epipolythiodiketopiperazines from the ascomycetous fungus *Preussia typharum*. *Journal of Natural Products*, **77**, 1459-1466.
- [63] Andernach L, Sandjo LP, Liermann JC, Buckel I, Thines E, Opatz T. (2013) Assignment of configuration in a series of dioxolanone-type secondary metabolites from *Guignardia bidwellii* - A comparison of VCD and ECD spectroscopy. *European Journal of Organic Chemistry*, **2013**, 5946-5951.
- [64] Lin X, Ji S, Qiao X, Hu H, Chen N, Dong Y, Huang Y, Guo D, Tu P, Ye M. (2013) Density functional theory calculations in stereochemical determination of terpecurcumins J-W, cytotoxic terpene-conjugated curcuminoids from *Curcuma longa* L. *Journal of Organic Chemistry*, **78**, 11835-11848.
- [65] Yamakado R, Mikami K, Takagi K, Azumaya I, Sugimoto S, Matsuoka S, Suzuki M, Katagiri K, Uchiyama M, Muranaka A. (2013) Helicity induction in three π -conjugated chromophores by planar chirality of calixamide. *Chemistry - A European Journal*, **19**, 11853-11857.
- [66] Ortega HE, Shen YY, TenDyke K, Rios N, Cubilla-Rios L. (2014) Polyhydroxylated macrolide isolated from the endophytic fungus *Pestalotiopsis mangiferae*. *Tetrahedron Letters*, **55**, 2642-2645.
- [67] Poopari MR, Dezhahang Z, Xu Y. (2013) A comparative VCD study of methyl mandelate in methanol, dimethyl sulfoxide, and chloroform: explicit and implicit solvation models. *Physical Chemistry Chemical Physics*, **15**, 1655-1665.
- [68] Mazzeo G, Santoro E, Andolfi A, Cimmino A, Troselj P, Petrovic AG, Superchi S, Evidente A, Berova N. (2013) Absolute configurations of fungal and plant metabolites by chiroptical methods. ORD, ECD, and VCD studies on phyllostin, scytolide, and oxysporone. *Journal of Natural Products*, **76**, 588-599.
- [69] Buffeteau T, Cavagnat D, Bisson J, Marchal A, Kapche GD, Battistini I, Da Costa G, Badoc A, Monti JP, Mérillon JM, Waffo-Tégou P. (2014) Unambiguous determination of the absolute configuration of dimeric stilbene glucosides from the rhizomes of *Gnetum africanum*. *Journal of Natural Products*, **77**, 1981-1985.
- [70] de Gussem ED, Herrebout W, Specklin S, Meyer C, Cossy J, Bultinck P. (2014) Strength by joining methods: combining synthesis with NMR, IR, and vibrational circular dichroism spectroscopy for the determination of the relative configuration in hemicalide. *Chemistry - A European Journal*, **20**, 17385-17394.
- [71] Batista Jr. JM, Blanch EW, Bolzani VS. (2015) Recent advances in the use of vibrational chiroptical spectroscopic methods for stereochemical characterization of natural products. *Natural Product Reports*, **32**, 1280-1302.

Study of Anti-Tuberculosis Activity Behaviour of Natural Kaurane and Trachylobane Diterpenes Compared with Structural Properties Obtained by Theoretical Calculations Ana C. F. Soares, Mirela M. W. Cabral, Carlos H. G. Martins, Alexsandro E. Ferreira, Pedro A. S. Bergamo, Leonida K. Omosa, Jacob O. Midiwo, Renato L. T. Parreira and Vladimir C. G. Heleno	763
Canthin-6-one Isolated from <i>Brucea javanica</i> Root Blocks Cancer Cells in the G₂/M phase and Synergizes with Cisplatin Norazwana Samat, Mei Fong Ng, Hui Mei Lee, Sui Kiong Ling, Pei Jean Tan and Vyomesh Patel	771
Chemical Constituents of the Root of <i>Angelica tenuissima</i> and their Anti-allergic Inflammatory Activity Hyun Gyu Choi, In-Gyu Je, Geum Jin Kim, Joo-Won Nam, Sang Hee Shim, Sang-Hyun Kim and Hyukjae Choi	779
Exploiting Substrate Diversity of NRPS Led to the Generation of New Sansanmycin Analogs Shan-Shan Wang, Ning-Ning Zhang, Ning He, Wen-Qiang Guo, Xuan Lei, Qiang Cai, Bin Hong and Yun-Ying Xie	781
Antioxidant Activity and Chemical Composition of Essential Oils of some Aromatic and Medicinal Plants from Albania Entela Hodaj-Çeliku, Olga Tsiftoglou, Lulëzim Shuka, Sokol Abazi, Dimitra Hadjipavlou-Litina and Diamanto Lazari	785
 <u>Accounts/Reviews</u>	
γ-Butyrolactones from <i>Aspergillus</i> Species: Structures, Biosynthesis, and Biological Activities Sabrin R. M. Ibrahim, Gamal A. Mohamed and Amgad I.M. Khedr	791

Natural Product Communications

2017

Volume 12, Number 5

Contents

Gerald Blunden Award (2016)

- | | |
|---|-----|
| Vibrational Circular Dichroism: Recent Advances for the Assignment of the Absolute Configuration of Natural Products
Eleuterio Burgueño-Tapia and Pedro Joseph-Nathan | 641 |
|---|-----|

Original Paper

- | | |
|--|-----|
| Antioxidant Activity of Carvone and Derivatives against Superoxide Ion
Sofia Pombal, Yaiza Hernández, David Díez, Eily Mondolis, Aldahir Mero, Juan Morán-Pinzón, Estela I. Guerrero and Jesús M. Rodilla | 653 |
| Studies in Cyclization of Aromatic Epoxy-acyclicpolyrenes: Lewis Superacids and Titanocene Chloride
Alexis Castillo, José Fco Quilez del Moral and Alejandro F. Barrero | 657 |
| 2α-Acetoxy-15-acetylartemisiifolin, a new Anti-trypanosomal Sesquiterpene Lactone from <i>Mikania guaco</i>
Ahmed A. Hussein, Luz Romero, Jose Luis López-Pérez and Mahabir P. Gupta | 659 |
| Structure-Dependent Cytotoxic Effects of Eremophilanolide Sesquiterpenes
M. Teresa Agulló-Ortuño, Carmen E. Díaz, Azucena González-Coloma and Matías Reina | 663 |
| Highly Functionalized Ring B Labdane Synthesis as Key Intermediate in the Route to Forskolin
Ignacio E. Tobal, Lourdes Castañeda, Alejandro M. Roncero, Rosalina F. Moro, David Díez and Isidro S. Marcos | 667 |
| Lathyrane Diterpenes from the Latex of <i>Euphorbia laurifolia</i>
María Jesús Durán-Peña, María Eugenia Flores-Giubi, José Manuel Botubol-Ares, Felipe Escobar-Montaño, Antonio J. Macías-Sánchez, Luis F. Echeverri, Isidro G. Collado and Rosario Hernández-Galán | 671 |
| Pentacyclic Triterpenoids from <i>Maytenus cuzcoina</i>
Carolina P. Reyes, Ignacio A. Jiménez and Isabel L. Bazzocchi | 675 |
| New Fluvirucinins C₁ and C₂ Produced by a Marine Derived Actinomycete
Margarida Costa, Paz Zúñiga, Ana M ^a Peñalver, Margrét Thorsteinsdottir, Marta Pérez, Librada M Cañedo and Carmen Cuevas | 679 |
| Optimization of the Number of Considered Conformers for the Absolute Configuration Determination of Catechin and Epicatechin Peracetates by VCD
Eleuterio Burgueño-Tapia, Mariano Sánchez-Castellanos and Pedro Joseph-Nathan | 683 |
| Profiling of Phenolic Natural Products in the Seagrass <i>Zostera noltei</i> by UPLC-MS
Lauren Manck, Ester Quintana, Rocio Suárez, Fernando G. Brun, Ignacio Hernández, María J. Ortega and Eva Zubía | 687 |
| The Dimerization of Precocene I
Braulio M. Fraga and Inmaculada Cabrera | 691 |
| Cytotoxic Terphenyl Neolignans from Fungus <i>Terana coerulea</i>: New Natural Corticins D and E, and Revised Structure for Corticin A
Maitane Maisterra, M ^a Ángeles Castro, Luz M ^a Muñoz-Centeno, Ricardo C. Calhelha, Isabel C.F.R. Ferreira and Pablo A. García | 695 |
| Synthesis and Biological Evaluation of Imines Structurally Related to Resveratrol as Dual Inhibitors of VEGF Protein Secretion and <i>hTERT</i> Gene Expression
Rosa Martí-Centelles, Juan Murga, Eva Falomir, Miguel Carda and J. Alberto Marco | 699 |
| Nematicidal Activity of the Essential Oil of Three Varieties of <i>Tagetes minuta</i> from Argentina
Yamile Massuh, Angel Cruz-Estrada, Azucena González-Coloma, Marta S. Ojeda, Julio A. Zygodlo and Maria Fe Andrés | 705 |
| Trypanocidal Effects of Essential Oils from Selected Medicinal Plants. Synergy among the Main Components
Nuria I. Guardo, Paula Sainz, Azucena González-Coloma, Jesús Burillo and Rafael A. Martínez-Díaz | 709 |

Accounts/Reviews

- | | |
|--|-----|
| Ring Rearrangement Metathesis in 7-Oxabicyclo[2.2.1]heptene (7-Oxanorbornene) Derivatives. Some Applications in Natural Product Chemistry
Silvia Roscales and Joaquín Plumet | 713 |
| Occurrence and Chemical Synthesis of Apocarotenoids from Mucorales: A Review
Alejandro F. Barrero, M. Mar Herrador del Pino, Jesús F. Arteaga and José A. González-Delgado | 733 |
| Phytochemicals and Biological Activities of Laurel Tree (<i>Laurus nobilis</i>)
Alfonso Alejo-Armijo, Joaquín Altarejos and Sofia Salido | 743 |

Original Paper

- | | |
|---|-----|
| Cannabidiolic Acid-Mediated Interference with AP-1 Transcriptional Activity in MDA-MB-231 Breast Cancer Cells
Masayo Suzuki, Shuso Takeda, Hiroyuki Okazaki, Kazuhito Watanabe, Masufumi Takiguchi and Hironori Aramaki | 759 |
|---|-----|

Continued Overleaf