


NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering all
Aspects of Natural Products Research



**This Issue is Dedicated to
the Memory of
Professor Ivano Morelli**

Volume 1. Issue 12. 2006
ISSN 1555-9475 (online)
www.naturalproduct.us

EDITOR-IN-CHIEF**DR. PAWAN K AGRAWAL**

*Natural Product Inc.
7963, Anderson Park Lane,
Westerville, Ohio, 43081 USA
agrawal@naturalproduct.us*

EDITORS**PROFESSOR GERALD BLUNDEN**

*The School of Pharmacy & Biomedical Sciences,
University of Portsmouth,
Portsmouth, PO1 2DT U.K.
gands@surefish.co.uk*

PROFESSOR ALESSANDRA BRACA

*Dipartimento di Chimica Bioorganica e Biofarmacia,
Università di Pisa,
via Bonanno 33, 56126 Pisa, Italy
Email: braca@farm.unipi.it*

PROFESSOR DEAN GUO

*State Key Laboratory of Natural and Biomimetic Drugs,
School of Pharmaceutical Sciences,
Peking University,
Beijing 100083, China
gda5958@163.com*

PROFESSOR ERNST HASLINGER

*Institute of Pharmaceutical Chemistry,
University of Graz,
A-8010 Graz, Austria
Ernst.Haslinger@uni-graz.at*

PROFESSOR J. ALBERTO MARCO

*Departamento de Química Orgánica,
Universidade de Valencia,
E-46100 Burjassot, Valencia, Spain
alberto.marco@uv.es*

PROFESSOR YOSHIHIRO MIMAKI

*School of Pharmacy,
Tokyo University of Pharmacy and Life Sciences,
Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan
mimakiy@ps.toyaku.ac.jp*

PROFESSOR M. G. REINECKE

*Department of Chemistry,
Texas Christian University,
Forts Worth, TX 76129, USA
m.reinecke@tcu.edu*

PROFESSOR YASUHIRO TEZUKA

*Institute of Natural Medicine
Toyama Medical and Pharmaceutical University,
2630-Sugitani, Toyama 930-0194, Japan
tezuka@ms.toyama-mpu.ac.jp*

ADVISORY BOARD

Prof. Oyvind Andersen
Bergen, Norway

Prof. Bruno Botta
Roma, Italy

Prof. Carlos Cerda-Garcia-Rojas
Mexico city, Mexico

Prof. Ioanna Chinou
Athens, Greece

Prof. Josep Coll
Barcelona, Spain

Prof. Geoffrey Cordell
Chicago, IL, USA

Prof. Samuel Danishefsky
New York, NY, USA

Dr. Biswanath Das
Hyderabad, India

Prof. Daneel Ferreira
Mississippi, MS, USA

Prof. A.A. Leslie Gunatillika
Tucson, AZ, USA

Prof. Stephen Hanessian
Montreal, Canada

Prof. Michael Heinrich
London, UK

Prof. Kurt Hostettmann
Lausanne, Switzerland

Prof. Martin A. Iglesias Arteaga
Mexico, D. F., Mexico

Prof. Norbert De Kimpe
Gent, Belgium

Prof. Jerzy Jaroszewski
Copenhagen, Denmark

Prof. Teodoro Kaufman
Rosario, Argentina

Prof. Marie Lacaille-Dubois
Dijon, France

Prof. Hartmut Laatsch
Göttingen, Germany

Prof. Hyeon-Kyu Lee
Daejeon, Korea

Prof. Shoei-Sheng Lee
Taipei, Taiwan

Prof. Chun-Nan Lin
Kaohsiung, China

Prof. Francisco Macias
Cadiz, Spain

Prof. Anita Marsaioli
Campinas, Brazil

Prof. Rachel Mata
Mexico D. F., Mexico

Prof. Imre Mathe
Szeged, Hungary

Prof. Joseph Michael
Johannesburg, South Africa

Prof. Virinder Parmar
Delhi, India

Prof. Luc Pieters
Antwerp, Belgium

Prof. Peter Proksch
Düsseldorf, Germany

Prof. Stephen Pyne
Wollongong, Australia

Prof. William Reynolds
Toronto, Canada

Prof. Raffaele Riccio
Salerno, Italy

Prof. Ricardo Riguera
Santiago de Compostela, Spain

Prof. Satyajit Sarker
Coleraine, UK

Prof. Monique Simmonds
Richmond, UK

Prof. Valentin Stonik
Vladivostok, Russia

Prof. Hermann Stuppner
Innsbruck, Austria

Prof. Apichart Suksamrarn
Bangkok, Thailand

Prof. Hiromitsu Takayama
Chiba, Japan

Prof. Paul Wender
Stanford, USA

INFORMATION FOR AUTHORS

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site <http://www.naturalproduct.us>.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the Natural Product Inc. (NPI). A photocopy license is available from the NPI for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

To Subscribe: Natural Product Communications (ISSN# 1555-9475) is a journal published monthly. 2006 subscription price: US\$1,095 (Web edition); US\$1,395 (Print); US\$1,795 (Print + single site online). Orders should be addressed to Subscription Department, Natural Product Communications, 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

Editorial

Special Issue in Memory of Professor Ivano Morelli

The December issue of *Natural Product Communications* is dedicated to Professor Ivano Morelli, an inspiration to scientists, young and old. The more enjoyable element of publishing this memorial issue comes from the recognition and tribute made to an exceptional individual by the many former students and colleagues who are authors of the papers in this issue. Indeed, Professor Morelli has played a key role in the development of phytochemical research, not only within the Italy, but also in the international arena.

Many thanks go to the authors, who have contributed some of their finest work to a timetable that has been quite challenging, and to the manuscript reviewers for their help in making this issue a success.

Finally, the editors would like to pay their tribute to the scholarly, scientific, and mentorship roles of Professor Ivano Morelli. Truly, his pursuit of scientific knowledge concerning organic chemistry and his dedication to his students and colleagues make him an exceptional individual worthy of our admiration, emulation, and the homage being paid to him in these pages of *Natural Product Communications*.

Pawan K. Agrawal
Editor-in-Chief

Obituary

Ivano Morelli (1940-2005)



Prof. Ivano Morelli was born March 11, 1940, in Pisa, Italy. He graduated from Pisa University in 1965 and received an Assistant Professor position in Organic Chemistry in 1966. From 1980 until his death he was Full Professor of Phytochemistry at the Faculty of Pharmacy, Pisa University.

During his career he was appointed as Director of Bioorganic and Biopharmacy Department since 1986 when it was called Institute of Organic Chemistry; in the last years he was also member of the Academic Senate of Pisa University. He served the Italian Society of Phytochemistry as President in the years 1987-1989 and 2004-2005. He was also the Director of the School of Science and Technology of Medicinal Plants of Pisa University from 1985 until 2001 (the year of its disactivation) and Director of several PhD courses during the last 10 years.

His chemical work focused on the study of mosses and reaction mechanisms of epoxides and dibromocyclohexanes during the years 1966-1978; chemical studies of Mediterranean plants belonging to Rosaceae, Apiaceae, and Asteraceae families (1978-1990); studies of plants belonging to developing countries of Latin America, Africa, and Asia (1990-2005); analysis of essential oils of Apiaceae, Asteraceae, and Lamiaceae from 1990 until 2005.

We remember him as a very active scientist and an eclectic man: a man of few words, but open-hearted, honest, concrete, and fair. The wellbeing of his collaborators, students, and friends was for him one of the most important thoughts. The door of his office was never closed and he knew how to listen the requests of everyone. He liked to give suggestions, but at the same time, he knew how to ask suggestions, in a mutual opinions exchange. His open-mindedness, curiosity, diplomacy, and passion brought him far, to Asia, Europe, and America, taking his scientific cooperation and opening his laboratory to many young researchers from different countries. He gives the opportunity to the young scientists to work without any restriction, giving hospitality to any people and to their opinions, since he was researcher of all the “human activities”. His contribution to the knowledge of the plants belonging to Italian and international flora is great but his love for the teaching, the research, the University, and the co-workers was great.

An inexorable leukemia took him far from his dear the 18 April 2005. He is survived by his wife, Iolena, and two sons, Marco and Ilaria. His love for the phytochemistry was stronger than the fear of his disease; he continued to be present among us even if he was far from the Department, during his hard time in the hospital. He had never stopped to fight: his brave heart was a daily teaching. The death obliged him to rest but never the death prevented him from teaching us his last lesson of life.

Alessandra Braca
Dipartimento di Chimica Bioorganica e Biofarmacia,
Universita di Pisa,
via Bonanno 33, 56126 Pisa,
Italy

Natural Product Communications

2006

Volume 1, NUMBER 12

Contents

<u>Original paper</u>	<u>Page</u>
Triterpenes from <i>Maytenus macrocarpa</i> and Evaluation of Their Anti-HIV activity Sonia Piacente, Lourdes Campaner Dos Santos, Naheed Mahmood and Cosimo Pizza	1073
New Oxidized 4-Oxo Fatty Acids from <i>Hygrophorus discoxanthus</i> Gianluca Gilardoni, Marco Clericuzio, Alberto Marchetti, Paola Vita Finzi, Giuseppe Zanoni and Giovanni Vidari	1079
Kenyaloside, a Novel <i>O,O,O</i>-Triglycosylated Naphthalene Derivative from the Exudate of Kenyan <i>Aloe</i> Species Giovanna Speranza, Daniela Monti, Sergio Crippa, Paola Cairolì, Carlo F. Morelli and Paolo Manitto	1085
New Flavonoid Glycosides from <i>Chrozophora senegalensis</i> and Their Antioxidant Activity Antonio Vassallo, Giuseppina Cioffi, Francesco De Simone, Alessandra Braca, Rokia Sanogo, Angelo Vanella, Alessandra Russo and Nunziatina De Tommasi	1089
<i>N1,N2,N3</i>-Trisisopentenyl Guanidine and <i>N1,N2</i>-Diisopentenyl guanidine, Two Cytotoxic Alkaloids from <i>Alchornea cordifolia</i> (Schumacher & Thonn.) Müll. Arg. (Euphorbiaceae) Root Barks Hélène Mavar-Manga, David Chapon, Sara Hoet, Sébastien Block, Marie-Claire. De Pauw-Gillet and Joëlle Quetin-Leclercq	1097
Indole Monoterpenes with Antichemotactic Activity from <i>Psychotria myriantha</i> Cláudia A. Simões-Pires, Fabianne M. Farias, Andrew Marston, Emerson F. Queiroz, Célia G. Chaves, Amélia T. Henriques and Kurt Hostettmann	1101
HPLC Based Activity Profiling for Inhibitors of Human Neutrophil Elastase in <i>Isatis tinctoria</i> Leaf Extracts M. Hamburger, H. G. Rüster and M. F. Melzig	1107
Variation in Artemisinin and Flavonoids Content in Different extracts of <i>Artemisia annua</i> L. Anna Rita Bilia, Caterina Gabriele, Maria Camilla Bergonzi, Pedro Melillo de Malgalhaes and Franco Francesco Vincieri	1111
Antifungal Evaluation of <i>Hypericum triquetrifolium</i> Polar Extracts Against <i>Fusarium</i> spp Daniele Fraternali, Alessandra Bertoli, Laura Giamperi, Anahi Bucchini, Donata Ricci, Francesco Menichini, Elena Trinciarelli and Luisa Pistelli	1117
Antioxidant Activity Analysis for the Selection of <i>Rosmarinus officinalis</i> L. Juan Antonio Garbarino, Nicolás Troncoso, Pia Delpiano, Loreto Carvajal and Alessandra Russo	1123
<i>Hypericum perforatum</i> L., <i>H. maculatum</i> Crantz., <i>H. calycinum</i> L. and <i>H. pulchrum</i> L.: Phytochemical and Morphological Studies Gelsomina Fico, Sara Vitalini, Noemi Colombo and Franca Tomè	1129
Chemical Composition and Antimicrobial Activities of Essential Oil of <i>Stachys glutinosa</i> L. from Sardinia Pintore Giorgio, Chessa Mario, Manconi Paola, Zanetti Stefania, Deriu Antonella and Tirillini Bruno	1133
Molecular Identification of <i>Panax ginseng</i> C.A. Meyer in Ginseng Commercial Products Paola Del Serrone, Lucilla Attorri, Bruno Gallinella, Francesca Romana Gallo, Elena Federici and Giovanna Palazzino	

Continued overleaf

Lipoxygenase Inhibitory Activity of Boropinic Acid, Active Principle from *Boronia pinnata*

Massimo Curini, Francesco Epifano, Salvatore Genovese, Luigi Menghini, Donata Ricci,
Daniele Fraternale, Laura Giamperi, Anahi Bucchini and Emanuele Bellacchio

1141

A Convenient Synthesis of 5'-Iodoresiniferatoxin (I-RTX)

Abdellah Ech-Chahad, Lahboub Bouyazza and Giovanni Appendino

1147

Review /Account**Acaricides of Natural Origin. Part 2. Review of the Literature (2002-2006)**

Guido Flamini

1151

Chemistry and Biological Activity of Triterpene Saponins from *Medicago* Species

Aldo Tava and Pinarosa Avato

1159

Non-nitrogenous Plant-derived Constituents with Antiplasmodial Activity

Anna Rita Bilia

1181

LIST OF AUTHORS

Antonella, D 1133
Appendino, G 1147
Attorri, L 1137
Avato, P 1159

Bellacchio, E 1141
Bergonzi, MC 1111
Bertoli, A 1117
Bilia, AR 1111,1181
Block, S 1097
Bouyazza, L 1147
Braca, A 1089
Bruno, T 1133
Bucchini, A 1117
Bucchini, A 1141

Cairolì, P 1085
Carvajal, L 1123
Chapon, D 1097
Chaves, CG 1101
Cioffi, G 1089
Claire, M 1097
Clericuzio, M 1079
Colombo, N 1129
Crippa, S 1085
Curini, M 1141

de Malgalhaes, PM 1111
Delpiano, P 1123
Dos Santos, LC 1073

Ech-Chahad, A 1147
Epifano, F 1141
Farias, FM 1101
Federici, E 1137
Fico, G 1129
Finzi, PV 1079
Flamini, G 1151
Fraternale, D 1117
Fraternale, D 1141

Gabriele, C 1111
Gallinella, B 1137
Gallo, FR 1137
Garbarino, JA 1123
Genovese, S 1141
Giamperi, L 1117
Giamperi, L 1141
Gilardoni, G 1079
Giorgio, P 1133

Hamburger, M 1107
Henriques, AT 1101
Hoet, S 1097
Hostettmann, K 1101

Mahmood, N 1073
Manitto, P 1085
Marchetti, A 1079
Mario, C 1133
Marston, A 1101
Mavar-Manga, H 1097
Melzig, MF 1107
Menghini, L 1141
Menichini, F 1117
Monti, D 1085
Morelli, CF 1085

Palazzino, G 1137
Paola, M 1133
Pauw-Gillet, MCD ... 1097
Piacente, S 1073
Pistelli, L 1117
Pizza, C 1073

Queiroz, EF 1101
Quetin-Leclercq, J 1097

Ricci, D 1117,1141
Russo, A 1089,1123
Rüster, GU 1107

Sanogo, R 1089
Serrone, PD 1137
Simões-Pires, C 1101
Simone, FD 1089
Speranza, G 1085
Stefania, Z 1133

Tava, A 1159
Tomè, F 1129
Tommasi, ND 1089
Trinciarelli, E 1117
Troncoso, N 1123

Vanella, A 1089
Vassallo, A 1089
Vidari, G 1079
Vincieri, FF 1111
Vitalini, S 1129

Zanoni, G 1079

Triterpenes from *Maytenus macrocarpa* and Evaluation of Their Anti-HIV Activity

Sonia Piacente^a, Lourdes Campaner Dos Santos^b, Naheed Mahmood^c and Cosimo Pizza^{a,*}

^aDipartimento di Scienze Farmaceutiche, University of Salerno, via Ponte Don Melillo, 84084 Fisciano, Salerno, Italy

^bInstituto de Química, UNESP, CP335, CEP 14801-970 Araraquara, SP, Brazil

^cRetroscreen Ltd, Academic Virology, London Hospital, Whitechapel, 64 Turner Street, London E1 2AD, U.K.

pizza@unisa.it

Received: September 8th, 2006; Accepted: November 2nd, 2006

Dedicated to the memory of Professor Ivano Morelli.

A set of pentacyclic triterpenes has been isolated from the bark of *Maytenus macrocarpa* (Celastraceae). It includes two new olean-12-ene derivatives, namely 3 β , 22 α -dihydroxy-olean-12-en-30-oic acid (**1**) and 22 α -hydroxy-olean-12-en-3-oxo-30-oic acid (**2**), and the new urs-12-ene derivative 3-(*E*)-coumaroyluvaol (**7**), along with 10 known compounds possessing olean-12-ene, urs-12-ene, lupane and friedelane skeletons. The structures have been elucidated by extensive spectroscopic methods including 1D- and 2D-NMR experiments, as well as ESI-MS analysis. All isolated compounds were tested for anti-HIV activity in C8166 cells infected with HIV-1_{MN}. The most active compound was 22 α -hydroxy-12-en-3-oxo-29-oic acid (triterpenonic acid A, **4**), with an EC₅₀ value of 1 μ g/mL and a selectivity index of 35.

Keywords: *Maytenus macrocarpa*, Celastraceae, pentacyclic triterpenes, anti-HIV activity.

Species belonging to the genus *Maytenus* (Celastraceae) have been used as a traditional medicine in the Amazonian region against cancer, rheumatism, and inflammation [1-2]. *M. macrocarpa* (R & P) Briquet is endemic to the Amazonian region of Peru and an alcoholic infusion of its bark is used, generally in “aguardiente”, for the treatment of rheumatism, influenza, gastrointestinal diseases, and as an antitumor agent for skin cancer [3]. In previous papers, dammarane [3] and friedelane triterpenes [4] from the stem bark exudates, sesquiterpene polyol esters from the leaves [5], and the nortriterpenes macrocarpins A-D from the roots [6] of *M. macrocarpa* have been reported. Furthermore a dihydro- β -agarofuran sesquiterpene isolated from the roots of *M. macrocarpa* has been reported to act as a modulator of daunomycin resistance in a multidrug-resistant *Leishmania tropica* line. [7].

Here we report the isolation and characterization of three new triterpenes from the bark of *M. macrocarpa*, namely 3 β , 22 α -dihydroxy-12-en-30-oic acid (**1**), 22 α -hydroxy-12-en-3-oxo-30-oic acid (**2**) and 3-(*E*)-*p*-coumaroyluvaol (**7**), along with the known olean-12-ene derivatives maytenfolic acid (**3**), triptotriterpenonic acid A (**4**), 22-*epi*-maytenfolic acid (**5**), 22-*epi*-triptotriterpenonic acid A (**6**), the urs-12-ene derivative 3-(*E*)-caffeoyluvaol (**8**), the lupane-type triterpenes 3-(*E*)-*p*-coumaroylbetulin (**9**), 3-(*Z*)-*p*-coumaroylbetulin (**10**), 3-(*E*)-caffeoylbetulin (**11**), nepeticin (**12**), and friedelane orthosphenic acid (**13**). On the basis of the anti-HIV activity reported for triterpenes closely related to compounds **1-13** isolated from *M. macrocarpa* [8-10], the inhibitory activity of these compounds against HIV-1 replication in acutely infected C8166 cells has been evaluated.




Figure 1: Compounds **1-13** isolated from the bark of *Maytenus macrocarpa*.

Column chromatography of a CHCl_3 extract of the bark of *M. macrocarpa* on silica gel, yielded compounds **1-13**.

Compound **1** was obtained as an amorphous white solid, which showed in the ESI-MS in positive ion mode the quasi-molecular ion $[\text{M}+\text{H}]^+$ at m/z 473, corresponding to the molecular formula $\text{C}_{30}\text{H}_{48}\text{O}_4$. The ^{13}C NMR spectrum showed signals for 30 carbons. The ^1H NMR spectrum showed seven tertiary methyl singlets at δ 0.82, 0.94, 1.00, 1.03, 1.05, 1.26, and 1.33. The presence of two secondary alcoholic functions was evident from the signals at δ 3.19 (1H, dd, $J = 3.5$ and 11.5 Hz) and δ 3.62 (1H, dd, $J = 3.0$ and 11.0 Hz), which correlated in the

HSQC spectrum with the carbon resonances at δ 79.7 and 78.8, respectively. An additional feature of the ^1H NMR spectrum was a signal at δ 5.24 (1H, t, $J = 3.5$ Hz) typical of H-12 of a Δ -12 oleanene. Analysis of ^1H and ^{13}C NMR spectroscopic data and comparison with those of 3 β ,22 α -dihydroxy-12-en-29-oic acid (**3**), known as maytenfolic acid [11], suggested that the difference between the two compounds should be confined to ring E [12] [δ 39.5 (C-17), 49.9 (C-18), 44.3 (C-19), 46.2 (C-20), 40.7 (C-21), 78.8 (C-22) 34.1 (C-29), 181.1 (C-30) in **1**; δ 39.9 (C-17), 48.3 (C-18), 42.3 (C-19), 45.2 (C-20), 38.9 (C-21), 76.9 (C-22), 180.6 (C-29), 21.6 (C-30) in **3**]. Comparison of the ROESY spectra of **1** and **3**

allowed us to clarify the difference between the two compounds. While, in the case of **3**, the ROESY spectrum showed correlations between the signal at δ 1.22 (Me-30) and the signals at δ 2.36 (H-18), 3.52 (H-22), and 1.01 (Me-28), these effects were absent in the ROESY spectrum of **1**, where the signal at δ 1.26 (Me-29) showed a correlation with that at δ 1.33 (Me-27). This evidence led to the conclusion that **1** was 3 β ,22 α -dihydroxy-olean-12-en-30-oic acid, which was named macrocarpoic acid A.

Compound **2** showed, in the ESI-MS in positive ion mode, the quasi-molecular ion $[M+H]^+$ at m/z 471, corresponding to the molecular formula $C_{30}H_{46}O_4$. The compound exhibited, in its 1H NMR spectrum, seven singlet methyl groups (δ 0.95, 1.09, 1.11 x 2, 1.13, 1.26, 1.33), one olefinic proton (δ 5.25), and one methine proton (δ 3.63) linked to an oxygen bearing carbon. In the ^{13}C NMR spectrum the occurrence of signals for 30 carbons, including peaks at δ 123.4, 143.5, 181.0 and 219.0, suggested that **2** was an oxoolean-12-enoic acid [13]. By comparison of NMR data of **2** with those of **1** it was speculated that the position of the carbonyl group (δ 219.0) was at C-3 on the basis of the absence in **2** of the signals at δ 3.19 in the 1H NMR spectrum and 79.7 in the ^{13}C NMR spectrum, observed for **1**. It was confirmed by the long range correlations observed in the HMBC spectrum between the proton signals at δ 1.09 (Me-23) and 1.11 (Me-24) and the carbon resonance at δ 219.0 (C-3). Thus **2** was assigned as 22 α -hydroxy-olean-12-en-3-oxo-30-oic acid, and named macrocarpoic acid B.

The molecular formula $C_{39}H_{56}O_4$ was assigned to compound **7** on the basis of the ESI-MS spectrum, which showed the quasi-molecular ion peak $[M+H]^+$ at m/z 589. The 1H NMR spectrum showed seven methyl groups at δ 0.88 (d, J = 6.5 Hz), 0.96 (s), 0.97 (d, J = 6.5 Hz), 1.01 (s), 1.08 (s), 1.09 (s), and 1.17 (s), two signals ascribable to a primary alcoholic function at δ 3.08 and 3.59 (each 1H, d, J = 11.0 Hz), and a signal for a methine proton linked to an oxygen-bearing carbon at δ 4.58 (dd, J = 3.5 and 11.0 Hz). Further features were signals due to three olefinic protons at δ 5.21 (t, J = 3.5), 6.32 (d, J = 15.9 Hz), and 7.64 (d, J = 15.9 Hz), and signals at δ 6.83 (2H, d, J = 8.0 Hz) and 7.50 (2H, d, J = 8.0 Hz), typical of a 1,4-disubstituted aromatic ring. The ^{13}C NMR spectroscopic data of **7** were similar to those of urs-12-en-3 β ,28-diol, known as uvaol [14], except for the chemical shift of C-3 and the occurrence of an

acyl moiety identified as (*E*)-*p*-coumaroyl. In the HMBC spectrum, the proton at δ 4.58 (H-3) correlated with the carbon resonance at δ 167.2 (C=O), indicating that the (*E*)-coumaroyl group was located at C-3. Thus compound **7** was identified as 3-(*E*)-coumaroyluvaol, and named macrocarpol A.

The known compounds maytenfolic acid (**3**) [11], triptotriterpenonic acid A (**4**) [15], 22-*epi*-maytenfolic acid (**5**) [16], 22-*epi*-triptotriterpenonic acid A (**6**) [16], 3-(*E*)-caffeoyluvaol (**8**) [14], 3-(*E*)-*p*-coumaroylbetulin (**9**) [17], 3-(*Z*)-*p*-coumaroylbetulin (**10**) [18], 3-(*E*)-caffeoylbetulin (**11**) [18], nepeticin (**12**) [12], and orthosphenic acid (**13**) [12] were identified by comparison of their spectroscopic data with those reported in the literature.

The co-occurrence in *M. macrocarpa* of triterpenes belonging to different classes is an unusual finding. For maytenfolic acid (**3**), previously isolated from *M. diversifolia* [11], antileukemic activity has been reported [11], while compounds **9-11**, betulin esters, have been reported to exert antitumor promoting activity [19]. Compounds **7-8** are esters of uvaol, for which antiproliferative and antileukemic activity have been reported [20].

Table 1: Anti-HIV activity of compounds **1-13**.

Compounds	EC ₅₀ ^a	TC ₅₀ ^b
1	10	50
2	10	50
3	10	80
4	1	35
5	100	200
6	50	100
7	10	50
8	10	100
9	12	62
10	10	25
11	5	50
12	10	50
13	10	80

EC₅₀^a = concentration (μ g/mL) that reduced by 50% the production of gp120 in infected C8166 cells. TC₅₀^b = concentration (μ g/mL) that caused 50% cytotoxicity to uninfected C8166 cells.

On the basis of the anti-HIV activity reported for the lupane derivative betulinic acid [8-9], and, to a minor extent, for salaspermic acid, a friedelane derivative closely related to orthosphenic acid [10], the anti-HIV activity of compounds **1-13** in C8166 cells infected with HIV-1_{MN} was tested. The most active compound was **4**, with an EC₅₀ value of 1 μ g/mL and a selectivity index above 30 (Table 1). The investigation of the mechanism of action of **4** revealed that it was more effective when added either

prior to or at the time of virus infection. Further experiments confirmed that it inhibited the binding of gp120 to sCD4 in a dose dependent manner. At a concentration corresponding to its EC₅₀, compound **4** inhibited the interaction between gp120 and CD4 by 55%.

The activity exerted by betulin esters (**9-11**) is very much lower than that reported for betulinic acid and its derivatives by Hashimoto *et al.* [8]. Also orthosphenic acid (**13**) showed an anti-HIV activity lower than that reported for salaspermic acid [10], from which it differs only by the occurrence of a further α -OH group at C-2.

Experimental

General procedures: Optical rotations were measured on a Jasco DIP 1000 polarimeter. UV spectra were obtained on a Beckman DU 670 spectrometer. IR measurements were obtained on a Bruker IFS-48 spectrometer. Melting points were determined using a Bausch & Lomb apparatus. Accurate molecular weights were measured by a Voyager DE mass spectrometer. Samples were analyzed by matrix assisted laser desorption ionization (MALDI) mass spectrometry. A mixture of analyte solution and α -cyano-4-hydroxycinnamic acid (Sigma) was applied to the metallic sample plate and dried. Mass calibration was performed with the ions from ACTH (fragment 18-39) at 2465.1989 Da and Angiotensin III at 931.5154 Da as internal standards. ESI-MS analyses were performed using a ThermoFinnigan LCQ Deca XP Max ion trap mass spectrometer equipped with Xcalibur software. NMR experiments were performed on a Bruker DRX-600 spectrometer at 300 K. All the 2D-NMR spectra were acquired in CD₃OD. Standard pulse sequence and phase cycling were used for DQF-COSY, HSQC, HMBC and ROESY spectra. TLC was performed on silica gel F254 (Merck) plates, and reagent grade chemicals (Carlo Erba) were used throughout.

Plant material: *Maytenus macrocarpa* was collected in the Ucayali Region (Pucallpa), Peru. A voucher specimen is deposited at the Department of Pharmaceutical Sciences, Salerno, Italy.

Extraction and isolation: The dried and powdered bark of *M. macrocarpa* (310 g) was defatted with light petroleum and then extracted by maceration at room temperature with CHCl₃ until exhaustion. The CHCl₃ extract was concentrated under reduced

pressure to a syrupy consistency (2.5 g). The crude extract was chromatographed on a silica gel column using CHCl₃ and increasing amounts of MeOH (up to 20%). After monitoring by TLC [Si gel plates, CHCl₃-MeOH (9:1)], the fractions were combined to give A (350 mg), B (150 mg), C (200 mg) and D (280 mg). Fractions A-D were submitted to HPLC on a μ -Bondapack C-18 column (30 cm x 7.8 mm i.d., flow rate 2.0 mL/min) using MeOH: H₂O in the ratio 85:15 for A-C and 3:1 for D (isocratic conditions). Pure **11** (12 mg, Rt = 12 min), **9** (8 mg, Rt = 15 min) and **10** (6 mg, Rt = 19 min) were obtained from A; **12** (9 mg, Rt = 24 min) from B; **7** (5 mg, Rt = 20 min), **8** (8 mg, Rt = 19 min) from C; **1** (7 mg, Rt = 8 min), **2** (5 mg, Rt = 12 min), **3** (12 mg, Rt = 10 min), **4** (10 mg, Rt = 15 min), **5** (7 mg, Rt = 11 min), **6** (9 mg, Rt = 16.5 min), and **13** (18 mg, Rt = 7 min) from D.

Antiviral assays: The anti-HIV activity and toxicity of compounds **1-13** were assessed in C8166 cells infected with HIV-1_{MN}. Cells were cultured in RPMI 1640 with 10% fetal calf serum. Forty-thousand cells per microtiter plate well were mixed with 5-fold dilutions of compounds prior to addition of 10 CCID₅₀ units of virus and incubated for 5-6 days. Formation of syncytia was examined from 2 days post-infection. The inhibition of HIV-infection was determined by examining syncytia, by estimating antigen gp120 by ELISA, and by measuring cell viability for virus-infected cells and uninfected cell controls using the XTT-formazan method.

Virus infectivity assay: The total progeny virus was titrated in microtiter plates using double dilutions of freshly collected supernatants and C8166 cells. The end point was determined by examining syncytia formation and by the XTT-formazan method. Virus titer (CCID₅₀) is expressed as the reciprocal of the dilution that gave a 50% end point. To measure the effects of compounds on virus infectivity, HIV-1_{MN} (10⁴-10⁵ CCID₅₀) was incubated with test compound at 37°C for 1h, the virus was serially diluted, and the infectivity end-point determined.

Gp120-sCD4 interaction assay: Gp120-sCD4 interaction was measured by ELISA; sCD4 was bound to microtiter plate wells at a concentration of 0.05 μ g/well. Various dilutions of compounds were mixed with equal volumes of recombinant gp120 (0.04 μ g/mL) and added to CD4 coated wells. After incubation at 37°C for 3-5 h, the binding of gp120 was detected using human anti-HIV serum and anti-human Ig conjugated to horseradish peroxidase.

Using WIACALC (Pharmacia LKB), the percent inhibition was calculated from linear logarithmic plots using three concentrations of gp120 alone as standard.

Macrocarpoic acid A (1)

MP: 272-274°C.

$[\alpha]_D$: +48.2° (c 0.1, CHCl₃).

Rf: 0.6 (CHCl₃-MeOH, 9:1).

IR (KBr): 3480, 2970-2880, 1690, 1450, 1360, 1230, 1070 cm⁻¹.

¹H NMR (600 MHz, MeOH): 0.82 (3H, s, Me-23), 0.94 (3H, s, Me-28), 1.00 (3H, s, Me-25), 1.03 (3H, s, Me-24), 1.05 (3H, s, Me-26), 1.26 (3H, s, Me-29), 1.33 (3H, s, Me-27), 3.19 (1H, dd, *J* = 3.5 and 11.5 Hz), 3.62 (1H, dd, *J* = 3.0 and 11.0 Hz), 5.24 (1H, t, *J* = 3.5).

¹³C NMR (150 MHz MeOH): 15.7 (CH₃, C-25), 16.1 (CH₃, C-24), 17.3 (CH₃, C-26), 19.0 (CH₂, C-6), 24.4 (CH₂, C-11), 25.8 (CH₃, C-28), 26.3 (CH₃, C-27), 26.5 (CH₂, C-15), 28.2 (CH₂, C-2), 28.3 (CH₃, C-23), 29.0 (CH₂, C-16), 33.4 (CH₂, C-7), 34.1 (CH₃, C-29), 37.8 (C, C-10), 38.9 (CH₂, C-1), 39.5 (C, C-17), 40.1 (C, C-4), 40.7 (CH₂, C-21), 41.2 (C, C-8), 43.5 (C, C-14), 44.3 (CH₂, C-19), 46.2 (C, C-20), 48.9 (CH, C-9), 49.9 (CH, C-18), 56.5 (CH, C-5), 78.8 (CH, C-22), 79.6 (CH, C-3), 123.5 (CH, C-12), 144.5 (C, C-13), 181.1 (C, C-30).

ESI-MS: 473 [M + H⁺], 495 [M + Na⁺].

HRMS-MALDI: *m/z* [M + Na⁺] calcd for C₃₀H₄₈NaO₄ 495.3450, found 495.3458.

Macrocarpoic acid B (2)

MP: 264-266°C.

$[\alpha]_D$: +76.2° (c 0.1, CHCl₃).

Rf: 0.7 (CHCl₃-MeOH, 9:1).

IR (KBr): 3450, 2980-2840, 1730, 1710, 1450, 1380, 1230, 1120 cm⁻¹.

¹H NMR (600 MHz, MeOH): 0.95 (3H, s, Me-28), 1.09 (3H, s, Me-24), 1.11 (6H, s, Me-23, Me-26), 1.13 (3H, s, Me-25), 1.26 (3H, s, Me-29), 1.33 (3H, s, Me-27), 3.63 (1H, dd, *J* = 3.0 and 11.0 Hz), 5.25 (1H, t, *J* = 3.5).

¹³C NMR (150 MHz MeOH): 15.5 (CH₃, C-25), 17.2 (CH₃, C-26), 20.3 (CH₂, C-6), 21.7 (CH₃, C-24), 24.0 (CH₂, C-11), 25.8 (CH₃, C-28), 26.2 (CH₃, C-27), 26.3 (CH₂, C-15), 26.6 (CH₃, C-23), 26.7 (CH₂, C-7),

29.0 (CH₂, C-16), 34.3 (CH₃, C-29), 35.0 (CH₂, C-2), 37.8 (C, C-10), 39.5 (C, C-17), 40.1 (CH₂, C-1), 40.6 (CH₂, C-21), 40.5 (C, C-8), 43.4 (C, C-14), 44.2 (CH₂, C-19), 46.0 (C, C-20), 48.0 (C, C-4), 47.5 (CH, C-9), 49.8 (CH, C-18), 56.0 (CH, C-5), 78.7 (CH, C-22), 123.4 (CH, C-12), 143.5 (C, C-13), 181.0 (C, C-30), 219.0 (C, C-3).

ESI-MS: 471 [M + H⁺], 493 [M + Na⁺].

HRMS-MALDI: *m/z* [M + Na⁺] calcd for C₃₀H₄₆NaO₄ 493.3294, found 493.3301.

Macrocarpol A (7)

MP: 310-312°C.

$[\alpha]_D$: +42.0° (c 0.1, CHCl₃).

Rf: 0.7 (CHCl₃-MeOH, 9:1).

IR (KBr): 3440, 2930, 1680, 1600, 1188 cm⁻¹.

¹H NMR (600 MHz, MeOH): 0.88 (3H, s, Me-29), 0.96 (3H, s, Me-23), 0.97 (3H, s, Me-30), 1.01 (3H, s, Me-24), 1.08 (3H, s, Me-25), 1.09 (3H, s, Me-26), 1.17 (3H, s, Me-27), 3.08 (1H, d, *J* = 11.0 Hz, H-28a), 3.59 (1H, d, *J* = 11.0 Hz, H-28b), 4.58 (1H, dd, *J* = 3.5, 12.0 Hz), 5.21 (1H, t, *J* = 3.5, H-12), 6.32 (1H, d, *J* = 15.9), 6.83 (1H, d, *J* = 8.0, H-3', H-5'), 7.50 (1H, d, *J* = 8.0, H-2', H-6'), 7.64 (1H, d, *J* = 15.9).

¹³C NMR (150 MHz MeOH): 16.3 (CH₃, C-25), 17.3 (CH₃, C-26), 17.5 (CH₃, C-24), 17.8 (CH₃, C-29), 18.8 (CH₂, C-6), 21.6 (C, C-30), 23.9 (CH₃, C-27), 24.1 (CH₂, C-16), 24.2 (CH₂, C-11), 25.9 (CH₂, C-2), 26.7 (CH₂, C-15), 28.3 (CH₃, C-23), 32.2 (CH₂, C-21), 34.3 (CH₂, C-7), 36.5 (CH, C-22), 37.7 (C, C-17), 38.0 (C, C-10), 38.7 (C, C-4), 39.8 (CH₂, C-1), 40.7 (CH, C-20), 41.3 (CH, C-8), 41.4 (CH, C-19), 43.3 (C, C-14), 49.4 (CH, C-9), 55.3 (CH, C-18), 56.7 (CH, C-5), 70.5 (CH₃, C-28), 115.8 (CH, C-3', C-5'), 116.4 (CH, C-8'), 125.1 (CH, C-12), 127.3 (C, C-1'), 129.5 (CH, C-2', C-6'), 140.8 (C, C-13), 143.8 (CH, C-7'), 157.4 (C, C-4') 167.2 (C, C-9').

ESI-MS: 589 [M + H⁺], 611 [M + Na⁺].

HRMS-MALDI: *m/z* [M + Na⁺] calcd for C₃₉H₅₆NaO₄ 611.4076, found 611.4082.

Acknowledgments - The authors thanks Dr Juan de Dios Zuniga Quiroz of Agro Selva Zuniga, Parque Caceres Dorregaray 86-C Pueblo Libre, Lima 21, Peru, for providing the plant material.

References

- [1] Piacente S, De Tommasi N, Pizza C. (1999) Laevisines A and B: two new sesquiterpene-pyridine alkaloids from *Maytenus laevis*. *Journal of Natural Products*, **62**, 161-163.

- [2] Muhammad I, El Sayed KA, Mossa JS, Al-Said MS, El-Feraly FS, Clark AM, Hufford CD, Oh S, Mayer AMS. (2000) Bioactive 12-oleanene triterpene and secotriterpene acids from *Maytenus undata*. *Journal of Natural Products*, **63**, 605-610.
- [3] Chavez H, Estevez-Braun AE, Ravelo AG, Gonzalez AG. (1997) First examples of dammarane triterpenes isolated from Celastraceae. *Tetrahedron*, **53**, 6465-6472.
- [4] Chavez H, Estevez-Braun AE, Ravelo AG, Gonzalez AG. (1998) Friedelane triterpenoids from *Maytenus macrocarpa*. *Journal of Natural Products*, **61**, 82-85.
- [5] Chavez H, Callo N, Estevez-Braun AE, Ravelo AG, Gonzalez AG. (1999) Sesquiterpene polyolesters from the leaves of *Maytenus macrocarpa*. *Journal of Natural Products*, **62**, 1576-1577.
- [6] Chavez H, Rodriguez G, Estevez-Braun AE, Ravelo AG, Estevez-Reyes R, Gonzalez AG, Fdez-Puente JL, Garcia-Gravalos D. (2000) Macrocarpins A-D, new cytotoxic nor-triterpenes from the leaves of *Maytenus macrocarpa*. *Biorganic & Medicinal Chemistry Letters*, **10**, 759-762.
- [7] Perez-Victoria JM, Tincusi BM, Jimenez IA, Bazzocchi IL, Gupta MP, Castanys S, Gamarro F, Ravelo AG. (1999) New natural sesquiterpenes as modulators of daunomycin resistance in a multidrug-resistant *Leishmania tropica* line. *Maytenus macrocarpa*. *Journal of Medicinal Chemistry*, **42**, 4388-4393.
- [8] Hashimoto F, Kashiwada Y, Cosentino LM, Chen CH, Garrett PE, Lee KH. (1997) Anti-AIDS agents-XXVII. Synthesis and anti-HIV activity of betulinic acid and dihydrobetulinic acid derivatives. *Biorganic & Medicinal Chemistry*, **5**, 2133-2143.
- [9] Sun IC, Chen CH, Kashiwada Y, Wu JH, Wang HK, Lee KH. (2002) Anti-AIDS agents 49. Synthesis, anti-HIV, and anti-fusion activities of IC9564 analogues based on betulinic acid. *Journal of Medicinal Chemistry*, **45**, 4271-4275.
- [10] Chen K, Shi Q, Kashiwada Y, Zhand DC, Hu CQ, Jin JQ, Nozaki H, Kilkuskie R, Tramontano E, Cheng YC, McPhail DR, McPhail AT, Lee KH. (1992) Anti-Aids agents, 6. Salaspermic acid, an anti-HIV principle from *Tripterygium wilfordii*, and the structure-activity correlation with its related compounds. *Journal of Natural Products*, **55**, 340-346.
- [11] Nozaki H, Suzuki H, Hirayama T, Kasai R, Wu RY, Lee KH. (1986) Antitumour triterpenes of *Maytenus diversifolia*. *Phytochemistry*, **28**, 479-485.
- [12] Mahato SB, Kundu AP. (1994) ¹³C NMR spectra of pentacyclic triterpenoids – A compilation and some salient features. *Phytochemistry*, **37**, 1517-1575.
- [13] Kaneda N, Pezzuto JM, Kinghorn AD, Farnsworth NR. (1992) Plant anticancer agents, L. Cytotoxic triterpenes from *Sandoricum koetjape* stems. *Journal of Natural Products*, **55**, 654-659.
- [14] Nakagawa H, Takaishi Y, Fujimoto Y, Duque C, Garzon C, Sato M, Okamoto M, Oshikawa T, Ahmed SU. (2004) Chemical constituents from the Colombian medicinal plant *Maytenus laevis*. *Journal of Natural Products*, **67**, 1919-1924.
- [15] Kutney JP, Hewitt GM, Lee G, Piotrowska K, Roberts M, Rettig SJ. (1992) Studies with tissue cultures of the Chinese herbal plant *Tripterygium wilfordii*. Isolation of metabolites of interest in rheumatoid arthritis, immunosuppression, and male contraceptive activity. *Canadian Journal of Chemistry*, **70**, 1455-1480.
- [16] Silva GDF, Duarte LP, Vieira Filho SA, Doriguetto AC, Mascarenhas YP, Ellena J, Castellano EE, Cota AB. (2002) Epikatoncic acid from *Austroplenckia populnea*: structure elucidation by 2D NMR spectroscopy and X-ray crystallography. *Magnetic Resonance in Chemistry*, **40**, 366-370.
- [17] Rashid MA, Gray AI, Waterman PG. (1992) Coumarins from *Phebalium tuberosum* ssp. *megaphyllum* and *Phebalium filifolium*. *Journal of Natural Products*, **55**, 851-858.
- [18] Fuchino H, Satho T, Tanaka N. (1995) Chemical evaluation of *Betula* species in Japan. I. Constituents of *Betula ermanii* *Chemical and Pharmaceutical Bulletin*, **43**, 1937-1942.
- [19] Gao H, Wu L, Kuroyanagi M, Harada K, Kawahara N, Nakane T, Umehara K, Hirasawa A, Nakamura Y. (2003) Antitumor-promoting constituents from *Chaenomeles sinensis* Koehne and their activities in JB6 mouse epidermal cells. *Chemical and Pharmaceutical Bulletin*, **51**, 1318-1321.
- [20] Es Saady D, Delage C, Simon A, Chulia AJ. (1995) Antiproliferative effects of uvaol. *Fitoterapia*, **66**, 366-369.

New Oxidized 4-Oxo Fatty Acids from *Hygrophorus discoxanthus*

Gianluca Gilardoni, Marco Clericuzio, Alberto Marchetti, Paola Vita Finzi, Giuseppe Zannoni and Giovanni Vidari*

Dipartimento di Chimica Organica, University of Pavia, Via Taramelli 10, 27100 Pavia, Italy

vidari@unipv.it.

Received: July 24th, 2006; Accepted: August 28th, 2006

Dedicated to the memory of Professor Ivano Morelli.

The results are reported from the first investigation of the secondary metabolites of the basidiomycete *Hygrophorus discoxanthus* (Fr.) Rea. Five new oxidized 4-oxo fatty acids (C₁₆, C₁₈) were isolated from the fruiting bodies and their structures established on the basis of their spectroscopic data and an ozonolysis experiment. Preliminary data indicate a moderate fungicidal activity, suggesting a possible function of these acids as chemical deterrents against mushroom parasites and predators.

Keywords: *Hygrophorus discoxanthus*, Basidiomycetes, 4-oxo-fatty acids, fungicidal activities.

In a search for new prototype (bioactive) agents from higher mushrooms (Basidiomycetes) [1], we were attracted by the species *Hygrophorus discoxanthus* (Fr.) Rea (fam. Hygrophoraceae) [2]. This is a mycorrhizal fungal species, growing solitary, scattered to gregarious in hardwood forests, particularly in the presence of *Fagus* trees, and fruiting in the fall. It is easily recognized by a whitish, viscid cap, with an ochreous-brown border, hence the name, and by the widely spaced, cream colored decurrent gills, turning rust-colored on rubbing. Our own field observations revealed that the fruiting bodies of *H. discoxanthus* are hardly ever attacked by either insects or parasitic fungi.

Fungicidal 4-oxo-2-alkenoic fatty acids were recently isolated from *H. eburneus* (Bull.: Fr.) Fr. [3], and related cyclopentenone and cyclopentenedione derivatives were found in the extracts of various *Hygrophorus* species [4,5]. In addition to the common fungal sterol ergosterol and derivatives, the aroma components of various *Hygrophorus* species were investigated by GC-MS [6]; a ceramide was reported from a Chinese *Hygrophorus* species [7], malodorous indole derivatives were isolated from

H. paupertinus A. H. Smith & Hesler [8], while muscaflavine and hygrophoric acid were identified as pigments of some *Hygrophorus* fruiting bodies [9]. No investigation of the secondary metabolites of *H. discoxanthus* has yet appeared in the literature. Along with the ecological observations, this prompted a study of the chemical constituents of this mushroom.

To prevent undesired enzymatic reactions, the fresh fruiting bodies were frozen after collection and extracted with EtOAc at -20°C. The crude extract was subsequently partitioned between *n*-hexane and MeCN, and the residue from the more polar layer was separated by chromatography on multiple reverse-phase C-18 columns to give acids **1-5**. Remarkably, these compounds exhibit brown-ochreous spots on C-18 TLC-plates sprayed with a sulfovanillin solution, followed by heating, and are thus well differentiable from the fungal ubiquitous oleic and linoleic acids, and methyl linoleate, of similar chromatographic polarity, which are detected as purple spots with the same reagent. In addition, TLC-spots of compounds **1** and **2** respond to UV light (fluorescence quenching at 254 nm).

Acids **1-5** (C_{16} or C_{18}) can be divided between those (**1-2**) presenting an δ,ϵ -unsaturated γ -oxocrotonate partial structure and those (**3-5**) containing a chetol system (Figure 1). Additionally, some compounds possess either an internal *Z*-configured double bond or a terminal one. Compounds **3-5** are optically active. Acid **1** was obtained as a whitish sticky solid. The UV spectrum showed an intense absorption band at $\lambda_{\max} = 234$ nm ($\text{Log } \epsilon = 4.34$) attributable to a $\pi \rightarrow \pi^*$ transition of a conjugated keto group, which was corroborated by an intense absorption peak at about 1666 cm^{-1} in the IR spectrum. On the other hand, an IR broad band extending from 3600 to 2800 cm^{-1} , along with a strong band at 1693 cm^{-1} revealed the presence of an unsaturated carboxylic acid. These attributions were firmly confirmed by the signals at $\delta 170.2$ and $\delta 188.1$ in the ^{13}C NMR spectrum of **1**, belonging to an unsaturated carboxylic group and an unsaturated carbonyl group, respectively. The negative ion ESI mass spectrum showed an ion at m/z 291 $[\text{M-H}]^-$ which, in accordance with data obtained from the NMR spectra, corresponded to the molecular formula $C_{18}\text{H}_{28}\text{O}_3$.




Figure 1: Acids **1-5** isolated from *Hygrophorus discoxanthus*.

The upfield portion of the ^1H NMR spectrum of compound **1** was typical of a long chain unsaturated fatty acid, as indicated by the distorted triplet at $\delta 0.88$, integrating for 3H, attributable to the $\omega 1$ methyl group, a broad signal at $\delta 1.10\text{--}1.45$, integrating for 12H, assignable to the $\omega 2\text{--}\omega 7$ methylene protons, and a distorted quartet at $\delta 2.05$ typical of an allylic methylene group (C-11). COSY and HMBC (Figure 2) correlations proved that this group was linked to a 1,2-disubstituted double bond, whose carbon signals were found at $\delta 127.1$ and 131.6 , respectively. The corresponding protons

resonated as well separated doublets of triplets at $\delta 5.30$ and $\delta 5.42$, respectively, and showed a mutual vicinal coupling constant of 10.3 Hz, indicative of a *Z*-configured double bond. Comparison of these data with the literature [10] showed that the structure of compound **1** corresponds, from C-9 to C-18, to that of oleic acid. The remaining eight carbons were assembled as a δ,ϵ -unsaturated γ -oxocrotonate unit, attached to C-9 by a C_2 linker, on the basis of the following NMR information. The proton doublets at $\delta 6.75$ and $\delta 7.48$ (1H each, $J_{\text{AB}} = 15.7$ Hz), which showed HSQC correlations with the carbon signals at $\delta 129.8$ and $\delta 139.7$, respectively, and HMBC correlations (Figure 2) with the signals at $\delta 170.2$ and $\delta 188.1$, indicated an *E*-configured double bond positioned between the carboxylic and the carbonyl group. The carbon signal of the ketone displayed additional HMBC cross peaks with two other olefinic methine resonances at $\delta 6.39$ and $\delta 7.06$ (1H each, vicinal coupling $J_{\text{AB}} = 15.9$ Hz) constituting an *E*-configured double bond, which was joined to C-9 through a CH_2CH_2 group. These two methylenes gave rise to two, well-resolved distorted quartets at $\delta 2.27$ ($\text{H}_2\text{-8}$) and $\delta 2.38$ ($\text{H}_2\text{-7}$), respectively, which showed two and three bond HMBC correlations with both C-6 and C-9 (Figure 2).




Figure 2: Selected HMBC correlations of compound **1**.

The spectral data of compound **2** were closely related to **1**, the most significant difference being the lack of evidence for an internal non-conjugated double bond. In fact, the UV absorption band at $\lambda_{\max} = 235$ nm, along with the IR peaks at 1690 and 1664 cm^{-1} , and the almost superimposable patterns of the ^1H - and ^{13}C NMR signals for the C(1)-C(6) moiety clearly proved that acid **2** contains the same *E,E*-configured δ,ϵ -unsaturated γ -oxocrotonate unit as compound **1**. From the mass spectral data, the length of the fatty acid chain in compound **2** could be determined as C_{16} , while the terminal double bond was identified by the signals from the three spin system at $\delta 5.83$ (1H, *ddt*, $J = 17.0, 10.3, 6.7$ Hz), $\delta 4.95$ (1H, *dtd*, $J = 10.3, 1.8, 1.5$ Hz), and $\delta 5.02$ (1H, *dtd*, $J = 17.0, 1.8, 1.5$ Hz).

The molecular formula $C_{18}\text{H}_{32}\text{O}_4$ of compound **3** was deduced from the ion at m/z 311 $[\text{M-H}]^-$ in the negative ion ESI spectrum, combined with the

protons and carbons counted from the NMR spectra. Remarkably, the ^1H NMR spectrum of **3** did not contain the characteristic signals of the cross conjugated dienone system of **1** and **2**; instead, three overlapping multiplets, each integrating for 2H, were found between δ 2.60–2.82, and were attributed to three different methylene groups adjacent to either saturated carbonyl or carboxylic groups. This assignment was confirmed by the resonances at δ 177.4 and 209.7 in the ^{13}C NMR spectrum, attributed to the carbons of a carboxylic acid and a saturated ketone, respectively. In addition, a broad multiplet at δ 4.10 (1H), which was correlated to a carbon at δ 67.7 in the HSQC spectrum, was firmly assigned to a secondary alcohol. The presence of an internal, non-conjugated, disubstituted olefin was demonstrated by an end absorption band at $\lambda_{\text{max}} = 218$ nm in the UV spectrum, along with the ^{13}C NMR signals of two methines at δ 129.3 and 130.1, which were correlated to an NMR signal at δ 5.25–5.45 in the HSQC spectrum. The AB coupling constant of 10.5 Hz of these two protons proved the Z-configuration of the double bond. A homonuclear COSY experiment, and two and three bonds HMBC correlations (Figure 3) allowed establishment of the 1,4-relationship of the carboxylic group with the ketone, and the 1,3-relationship of the hydroxyl and carbonyl groups.

NMR data alone left the position of the internal double bond undetermined. Therefore, compound **3** was exposed to ozone and, after work-up, the crude reaction mixture was directly subjected to GC analysis. Comparison with an authentic sample revealed heptanal to be formed by ozonolysis of olefin **3**. From all results, the structure of compound **3** was established as (Z)-6-hydroxy-4-oxo-octadec-11-enoic acid.




Figure 3: Selected HMBC correlations of compound **3**.

The NMR data of compounds **4** and **5** were closely related to **3** as regards to the 6-hydroxy-4-oxo-carboxylic acid [C(1)–C(6)] unit. In contrast, other than compound **3**, the acids **4** and **5** did not show the signals of an internal double bond. Instead, in the ^1H NMR spectrum of compound **4**, the pattern of signals

from a three spin system at δ 4.95, 4.99, and 5.83, almost identical to that of acid **2** (see above) were due to a terminal double bond. On the other hand, compound **5** contains a fully saturated fatty acid-like chain, as indicated, in the ^1H NMR spectrum, by the characteristic distorted triplet ($J = 6.8$ Hz) at δ 0.88, assigned to the terminal methyl group, and by a broad peak at δ 1.20–1.60, assigned to the methylenes in the chain. From the mass spectral data, the length of the chain in both compounds **4** and **5** could be determined as C_{16} , thus permitting assignment of the structure of 6-hydroxy-4-oxo-hexadec-15-enoic acid to **4**, and of 6-hydroxy-4-oxo-hexadecanoic acid to **5**.

The absolute configuration of carbinols **3–5** has yet to be determined. Compounds **1–5** have never been isolated from a natural source; acid **5** was obtained previously as a racemate by synthesis [11].

A preliminary qualitative test indicated that acids **1** and **2** are moderately fungicidal against the phytopathogenic fungus *Cladosporium cucumerinum* Ell. et Arth..

The structures **1–5** are closely related to other oxidized C_{16} – C_{22} fatty acids and their derivatives recently isolated from a few *Hygrophorus* species [3–5], for which hypothetical biogenetic relationships have been proposed [3,5]. A rare feature of all these structures is the oxidation to a ketone of the C-4 of the parent fatty acid; a few compounds show an additional site-specific oxidation at C-6, which the optically active alcohols **3–5** indicate to occur under enzyme control. Indeed, 6-hydroxy-4-oxo-carboxylic acids like **3–5** are, to our knowledge, unprecedented in nature. They can be considered advanced biogenetic precursors of hygrophorones F^{12} and G^{12} [4a]. Examining the literature data, it was concluded that each *Hygrophorus* species is characterized by its own pattern of oxidized C_{16} – C_{22} fatty acid derivatives, which may thus be considered a significant chemotaxonomic marker. Moreover, due to the fungicidal and bactericidal properties [3–5], these metabolites likely function as “chemical deterrents”, protecting *Hygrophorus* fruiting bodies against the attack of parasites and predators.

Experimental

General experimental procedures: Optical rotations were determined on a Perkin-Elmer 241 polarimeter; IR spectra were recorded on an FT-IR Perkin Elmer Paragon 1000 PC spectrometer as neat films on NaCl

discs. UV spectra were obtained in spectrometer grade CHCl_3 from a Jasco V-550 spectrophotometer. ^1H and ^{13}C NMR spectra were determined in CDCl_3 on a Bruker CXP 300 spectrometer operating at 300 MHz (^1H) and 75 MHz (^{13}C), respectively. ^1H and ^{13}C chemical shifts (δ , ppm) are relative to residual CHCl_3 signals [δ_{H} 7.26; δ_{C} (central line of t) 77.1, respectively]. 2D NMR spectra (COSY, HSQC, HMBC) were recorded by using standard pulse sequences. The abbreviations s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, and br = broad are used throughout; coupling constants (J) are reported in Hz. ESIMS experiments were carried out using a Finnigan LCQ Advantage MS 1.4 spectrometer, equipped with the Xcalibur 1.4 software. High-resolution ESI mass spectra were determined on a Bruker Apex II FT-ICR mass spectrometer. TLC was performed on sheets precoated with either silica gel F₂₅₄ (Polygram) or with RP-18 F₂₅₄ (Merck, Germany). Compounds were visualized under UV light (254 and 366 nm) and by spraying with a 0.5% solution of vanillin in H_2SO_4 -EtOH (4:1), followed by heating. Preparative column chromatography was carried out on LiChroprep RP-18 (25-40 μm , Merck). Reagent grade solvents, redistilled just before use, were employed for extraction; HPLC grade solvents were used for chromatographic separations. GC analysis was performed with a Perkin Elmer Autosystem gaschromatograph.

Fungal material: Fresh fruiting bodies of *Hygrophorus discoxanthus* (Batsch.: Fr.) Fr. were collected on 16 October 2005 in a mixed conifer and beech wood near Brallo, in the province of Pavia, Italy, at an altitude of 1050 m. The mushroom was identified by one of the authors (M.C.) and a frozen voucher specimen has been deposited at the Dipartimento di Chimica Organica, University of Pavia, Italy.

Extraction and isolation: Fruiting bodies (750 g) were frozen at -20°C , minced, and extracted at -20°C with EtOAc (3 x 1.5 L), followed by MeOH- H_2O (4:1, 1 L), and H_2O (1 L) at 0°C . The light yellow EtOAc solution was concentrated to dryness *in vacuo* at $<30^\circ\text{C}$ to produce an oily residue (2.1 g), which was partitioned between MeCN (0.5 L) and *n*-hexane (0.5 L). Evaporation of the two layers gave crude residues of 1.02 g and 1.08 g, respectively. Acids **1-5** were contained in the MeCN extract (TLC: R_f = 0.55-0.70; RP-18 F₂₅₄, solvent system: MeCN- H_2O , 7:1 v/v), which was subjected to column

chromatography on a LiChroprep RP-18 column (100 g). Elution was performed with a gradient of MeCN- H_2O , starting from a mixture 1:1, v/v, and increasing MeCN regularly every 100 mL, until a final mixture of MeCN- H_2O , 10:1, v/v. The column was then washed with MeCN (100 mL), followed by Me_2CO (100 mL). Thirty-four fractions (A1-A34), of 35 mL each, were collected. Fraction A9 gave acid **4** (12 mg, $1.6 \cdot 10^{-3}$ % of fresh fruiting bodies), fraction A13 gave acid **5** (25 mg, $3.2 \cdot 10^{-3}$ %), and fraction A14 afforded acid **3** (34 mg, $4.6 \cdot 10^{-3}$ %). Linoleic acid (36 mg, $4.8 \cdot 10^{-3}$ %), oleic acid (107 mg, $14 \cdot 10^{-3}$ %), and methyl linoleate (37 mg, $4.9 \cdot 10^{-3}$ %) were obtained by evaporation of fractions A23, A25, and A27, respectively. Fraction A15 (76 mg) was further separated on a LiChroprep RP-18 column (20 g) eluted with a gradient of MeCN- H_2O , starting from a mixture 1:1, v/v, and increasing MeCN regularly every 50 mL, until a final mixture of MeCN- H_2O , 10:1, v/v. Thirteen fractions (B1-B13), of 40 mL each, were collected. Acid **2** (10 mg, $1.4 \cdot 10^{-3}$ %) was isolated by evaporation of fraction B7. Fractions A19 and A20 were pooled together and the residue (110 mg) was further separated on a LiChroprep RP-18 column (20 g) eluted with a gradient of MeCN- H_2O , starting from a mixture 1:1, v/v, and increasing MeCN regularly every 50 mL, until a final mixture of MeCN- H_2O , 10:1, v/v; 14 fractions (C1-C14), each of 35 mL, were collected. Fraction C5 (61 mg) afforded compound **1** (10 mg, $1.4 \cdot 10^{-3}$ %) on successive separation on a LiChroprep RP-18 column (15 g) eluted with MeOH- H_2O , 4:1, v/v.

(2E, 5E, 9Z)-4-Oxo-octadeca-2,5,9-trienoic acid (**1**)

Whitish sticky solid.

R_f: 0.45 (RP18, MeCN- H_2O , 7:1).

IR (film): 3600-2800, 3090, 3050, 2920, 2852, 1693, 1666, 1613, 1278, 1216, 1000, 975, 950 cm^{-1} .

UV/Vis λ_{max} (CHCl_3) nm (log ϵ): 234 (4.34).

^1H NMR: 0.88 (3H, t, J = 6.8 Hz, Me), 1.10-1.45 (12H, brs H_2 -12- H_2 -17), 2.05 (2H, q, J = 6.8 Hz, H_2 -11), 2.27 (2H, distorted q, J = 7.0 Hz, H_2 -8), 2.38 (2H, distorted q, J = 7.0 Hz, H_2 -7), 5.30 (1H, dd, J = 10.3, 6.5 Hz, H-9), 5.42 (1H, dd, J = 10.3, 6.5 Hz, H-10), 6.39 (1H, d, J = 15.9 Hz, H-5), 6.75 (1H, d, J = 15.7 Hz, H-3), 7.06 (1H, dt, J = 15.9, 6.7 Hz, H-6), 7.48 (1H, d, J = 15.7 Hz, H-2).

^{13}C NMR: 13.9 (CH_3 , C-18), 22.5 (CH_2 , C-17), 25.5 (CH_2 , C-8), 27.2 (CH_2 , C-11), 29.2, 29.3, 29.4, 29.5 (4 CH_2 , C-12, C-13, C-14, C-15), 31.6 (CH_2 , C-16), 32.8 (CH_2 , C-7), 127.1 (CH, C-9), 129.4 (CH, C-5),

129.8 (CH, C-3), 131.6 (CH, C-10), 139.7 (CH, C-2), 150.8 (CH, C-6), 170.2 (C, C-1), 188.1 (C, C-4).

Negative ion ESI-FT-ICR-MS: m/z [M – H][–] calcd for C₁₈H₂₇O₃ 291.1960, found 291.1962.

(2E, 5E)-4-Oxo-hexadeca-2,5,15-trienoic acid (2)

Whitish sticky solid.

Rf: 0.5 (RP18, MeCN-H₂O, 7:1).

IR (film): 3600-3200, 3050, 2923, 2851, 1690, 1664, 1625, 1279, 1215, 1000, 915 cm^{–1}.

UV/Vis λ_{max} (CHCl₃) nm (log ε): 235 (4.19).

¹H NMR: 1.30-1.65 (12H, brs H₂-8–H₂-13), 2.06 (2H, q, J = 6.8 Hz, H₂-14), 2.31 (2H, q, J = 7.2 Hz, H₂-7), 4.95 (1H, dtd, J = 10.3, 1.8, 1.5 Hz, H-16E), 5.02 (1H, dtd, J = 17.0, 1.8, 1.5 Hz, H-16Z), 5.83 (1H, ddt, J = 17.0, 10.3, 6.7 Hz, H-15), 6.39 (1H, d, J = 15.9 Hz, H-5), 6.75 (1H, d, J = 15.7 Hz, H-3), 7.06 (1H, dt, J = 15.9, 6.7 Hz, H-6), 7.48 (1H, d, J = 15.7 Hz, H-2).

¹³C NMR: 27.8, 28.7, 28.9, 29.2, 29.3, 29.4 (6 x CH₂, C-8, C-9, C-10, C-11, C-12, C-13), 32.8 (CH₂, C-7), 33.7 (CH₂, C-14), 114.0 (CH₂, C-16), 129.2 (CH, C-5), 129.7 (CH, C-3), 139.0 (CH, C-15), 139.7 (CH, C-2), 151.7 (CH, C-6), 169.5 (C, C-1), 188.2 (C, C-4).

Negative ion ESI-FT-ICR-MS: m/z [M – H][–] calcd for C₁₆H₂₃O₃: 263.1647; found: 263.1649.

(Z)-6-Hydroxy-4-oxo-octadec-11-enoic acid (3)

Whitish sticky solid.

[α]_D²⁵: –340° (c = 10 mg/mL, CHCl₃).

Rf: 0.6 (RP18, MeCN-H₂O, 7:1).

IR (film): 3600-3200, 3010, 2928, 2856, 1713, 1406, 1260, 1201, 1100 cm^{–1}.

¹H NMR: 0.88 (3H, t, J = 6.8 Hz, Me), 1.20-1.65 (14H, brs, H₂-14–H₂-17, H₂-7–H₂-9), 2.05 (4H, q, J = 7.0 Hz, H₂-10, H₂-13), 2.60-2.82 (6H, m, H₂-2, H₂-3, H₂-5), 4.10 (1H, brm, H-6), 5.25-5.45 (2H, m, H-11, H-12).

¹³C NMR: 14.0 (CH₃, C-18), 22.5 (CH₂, C-17), 25.0, 27.0, 27.1, 28.9, 29.5, 29.6 (6 x CH₂, C-8, C-9, C-10, C-13, C-14, C-15), 27.4 (CH₂, C-2), 31.7 (CH₂, C-16), 36.3 (CH₂, C-7), 37.5 (CH₂, C-3), 49.1 (CH₂, C-5), 67.7 (CH, C-6), 129.3, 130.1 (2 x CH, C-11, C-12), 177.4 (C, C-1), 209.7 (C, C-4).

Negative ion ESI-FT-ICR-MS: m/z [M – H][–] calcd for C₁₈H₃₁O₄ 311.2222, found 311.2225.

6-Hydroxy-4-oxo-hexadec-15-enoic acid (4)

Whitish sticky solid.

[α]_D²⁵: –109° (c = 11 mg/mL, CHCl₃).

Rf: 0.7 (RP18, MeCN-H₂O, 7:1).

IR (film): 3600-3200, 3010, 2917, 2850, 1702, 1412, 1250, 1080, 1000, 913 cm^{–1}.

¹H NMR: 1.25-1.65 (14H, brs, H₂-7–H₂-13), 2.04 (2H, q, J = 6.7 Hz, H₂-14), 2.60-2.80 (6H, m, H₂-2, H₂-3, H₂-5), 4.10 (1H, brm, H-6), 4.95 (1H, dtd, J = 10.3, 1.8, 1.5 Hz, H-16E), 4.99 (1H, dtd, J = 17.0, 1.8, 1.5 Hz, H-16Z), 5.83 (1H, ddt, J = 17.0, 10.3, 6.7 Hz, H-15).

¹³C NMR: 27.4 (CH₂, C-2), 25.3, 28.8, 29.0, 29.2, 29.3, 29.4 (6 x CH₂, C-8, C-9, C-10, C-11, C-12, C-13), 33.7 (CH₂, C-14), 36.3 (CH₂, C-7), 37.6 (CH₂, C-3), 49.1 (CH₂, C-5), 67.8 (CH, C-6), 114.0 (CH₂, C-16), 139.1 (CH, C-15), 177.1 (C, C-1), 209.7 (C, C-4).

Negative ion ESI-FT-ICR-MS: m/z [M – H][–] calcd for C₁₆H₂₇O₄ 283.1909, found 283.1911.

6-Hydroxy-4-oxo-hexadecanoic acid (5)

Whitish sticky solid.

[α]_D²⁵: –95° (c = 10 mg/mL, CHCl₃).

Rf: 0.65 (RP18, MeCN-H₂O, 7:1).

IR (film): 3600-3200, 2920, 2855, 1710, 1415, 1255 cm^{–1}.

¹H NMR: 0.88 (3H, t, J = 6.8 Hz, Me), 1.20-1.60 (18H, brs, H₂-7–H₂-15), 2.60-2.80 (6H, m, H₂-2, H₂-3, H₂-5), 4.10 (1H, brm, H-6).

¹³C NMR: 14.0 (CH₃, C-16), 27.6 (CH₂, C-2), 22.8-29.6 (8 x CH₂, C-8, C-9, C-10, C-11, C-12, C-13, C-14, C-15), 36.3 (CH₂, C-7), 37.6 (CH₂, C-3), 49.1 (CH₂, C-5), 67.7 (CH, C-6), 177.2 (C, C-1), 209.5 (C, C-4).

Negative ion ESI-FT-ICR-MS: m/z [M – H][–] calcd for C₁₆H₂₉O₄ 285.2066, found 285.2064.

Ozonolysis of acid 3: A saturated solution of O₃ in CH₂Cl₂-MeOH, 4:1 v/v, was added to compound **3** (3 mg) dissolved in CH₂Cl₂, (0.5 mL) at –78°C. The reaction was quenched after 3 h by adding excess Me₂S and the mixture was left at –20°C overnight. A sample was directly analyzed by GC under the following conditions: column HP-5 (25 m×0.25 mm, 0.33 μm film thickness), injection temperature 250°C, detector (FID) temperature 280°C, carrier gas nitrogen, flow rate 1.27 mL/min, constant flow mode, split splitless injection, ratio 1:35, column temperature program: 40°C for 5 min, then raised to 100°C at a rate of 2°C/min, then raised to 280°C at a rate of 10°C/min, then isothermal at 280°C for 5 min. Enrichment of the peak eluted at 9.69 min with an authentic sample of heptanal, confirmed its identity.

Fungicidal activity: A simple test, adapted from the literature [4a, 12], was carried out to reveal the

possible fungicidal activity of compounds **1-5**. Five solutions of compounds **1-5** in MeOH, each containing approximately 20 µg of substance, were spotted on F₂₅₄ Merck silica gel plastic sheets, which were sprayed with a conidial suspension of *Cladosporium cucumerinum* Ell. et Arth spores in a glucose mineral medium (Czapek broth). The plates were then incubated at 25°C in the dark in a wet chamber (> 95% humidity) for 5 days, when they were overgrown with a dark gray colored mycelium. White spots (inhibition zones), signaling fungicidal activity, were found, in particular in correspondence

with compounds **1** and **2**; they were about eight times smaller than the inhibition area of the reference compound pseudomycin A (20 µg).

Acknowledgments - The authors thank Prof. Mariella Mella and Prof. Giorgio Mellerio for NMR and MS spectra measurements, respectively. Fungicidal tests were carried out by Dr Solveig Tosi. Financial support by the Italian MIUR (Grants COFIN and FIRB) and the University of Pavia (Grant FAR) is acknowledged.

References

- [1] This paper is Part 51 of the series "Fungal Metabolites". Part 50: Clericuzio M, Tabasso S, Bianco MA, Pratesi G, Beretta G, Tinelli S, Zunino F, Vidari G. (2006) Cucurbitane triterpenes from fruiting bodies and cultivated mycelia of *Leucopaxillus gentianus*. *Journal of Natural Products*, submitted.
- [2] Bon M (1990) Flore Mycologique d'Europe. 1. Les Hygrophores Hygrophoraceae Lotsy. In *Documents Mycologiques* mémoire hors série 1. CRDP, Amiens, 1-99.
- [3] Teichert A, Lübken T, Schmidt J, Porzel A, Arnold N, Wessjohann L. (2004) Unusual bioactive 4-oxo-2-alkenoic fatty acids from *Hygrophorus eburneus*. *Zeitschrift für Naturforschung*, **60B**, 25-32.
- [4] (a) Lübken T, Schmidt J, Porzel A, Arnold N, Wessjohann L. (2004) Hygrophorones A-G: fungicidal cyclopentenones from *Hygrophorus* species (Basidiomycetes). *Phytochemistry*, **65**, 1061-1071; (b) Lübken T, Arnold N, Wessjohann L, Böttcher C, Schmidt J. (2006) Analysis of fungal cyclopentenone derivatives from *Hygrophorus* spp. by liquid chromatography/electrospray-tandem mass spectrometry. *Journal of Mass Spectrometry*, **41**, 361-371.
- [5] Gilardoni G, Clericuzio M, Vidari G. Chrysotrienes A and B from *Hygrophorus chrysodon*, unpublished results.
- [6] Breheret S, Talou T, Rapior S, Bessiere JM. (1997) Monoterpenes in the aromas of fresh wild mushrooms (Basidiomycetes). *Journal of Agricultural and Food Chemistry*, **45**, 831-836.
- [7] Qu Y, Zhang H, Liu J. (2004) Isolation and structure of a new ceramide from the basidiomycete *Hygrophorus eburneus*. *Zeitschrift für Naturforschung*, **59B**, 241-244.
- [8] Wood WF, Smith J, Wayman K, Largent DL. (2003) Indole and 3-chloroindole: the source of the disagreeable odor of *Hygrophorus paupertinus*. *Mycologia*, **95**, 807-808.
- [9] Gill M, Steglich W. (1987) Pigments of fungi (macromycetes). In *Progress in the Chemistry of Organic Natural Products*. Vol **51**, Herz W, Grisebach H, Kirby GW, Tamm Ch. (Eds). Springer Verlag, Wien, New York. 1-317.
- [10] (a) Williamson RT, Carney JR, Gerwick WH. (2000) Application of the BIRD sandwich for the rapid and accurate determination of ¹H-¹H NMR coupling constants in higher order spin systems. *Journal of Natural Products*, **63**, 876-878; (b) Stamatov SD, Stawinski J. (2000) A simple and efficient method for direct acylation of acetals with long alkyl-chain carboxylic acid anhydrides. *Tetrahedron*, **56**, 9697-9703; (c) Vieville C, Mouloungui Z, Gaset A. (1995) Synthesis and analysis of the C1-C18 alkyl oleates. *Chemistry and Physics of Lipids*, **75**, 101-108; (d) Rossi R, Carpita A, Quirici MG, Verancini CA. (1982) Insect pheromone components. Use of carbon-13 NMR spectroscopy for assigning the configuration of carbon-carbon double bonds of monoenic or dienic pheromone components and for quantitative determination of Z/E mixtures. *Tetrahedron*, **38**, 639-644.
- [11] Baraldi PG, Barco A, Benetti S, Manfredini S, Simoni D. (1987) Ring cleavage of 3,5-disubstituted 2-isoxazolines by molybdenum hexacarbonyl and water to β-hydroxy ketones. *Synthesis*, **3**, 276-278.
- [12] Gottstein D, Gross D, Lehmann H. (1982) Mikrobiotest mit *Cladosporium cucumerinum* Ell. et Arth. zum Nachweis fungitoxischer Verbindungen auf Dünnschichtplatten. *Archiv für Phytopathologie und Pflanzenschutz*, **20**, 111-116.

Kenyaloside, a Novel *O,O,O*-Triglycosylated Naphthalene Derivative from the Exudate of Kenyan *Aloe* Species^{¶*}

Giovanna Speranza^{a,*}, Daniela Monti^b, Sergio Crippa^a, Paola Cairolì^a, Carlo F. Morelli^a and Paolo Manitto^a

^aDipartimento di Chimica Organica e Industriale, Università degli Studi di Milano, via Venezian 21, 20133 Milano, Italy

^bIstituto di Chimica del Riconoscimento Molecolare, C.N.R., via Mario Bianco 9, 20131 Milano, Italy

giovanna.speranza@unimi.it

Received: July 11th, 2006; Accepted: September 2nd, 2006

Dedicated to the memory of Professor Ivano Morelli.

A new naphthalene *O,O,O*-triglycoside, kenyaloside (**1**), was isolated from the dried exudate of Kenyan *Aloe* species, a bittering and laxative agent. Its structure was established by combined spectral and chemical methods as 1-(β -D-glucopyranosyloxy)-8-(α -L-rhamnopyranosyloxy)-3-(β -D-xylopyranosyloxymethyl)naphthalene.

Keywords: aloes, *Aloe ferox*, naphthalene *O,O,O*-triglycoside, kenyaloside.

As part of a systematic chemical investigation into *Aloe* exudates (bitter aloes) [1], the structural elucidation of a new water-soluble constituent of the exudate of Kenyan *Aloe* species is reported here. This exudate, flowing from the cut leaves of *Aloe ferox* Miller and of its hybrids with *A. spicata* and *A. africana* growing in Kenya [2, 3], when dried, is used as a bittering agent and as a purgative, similarly to Cape aloes [4, 5]. The drug has been reported to contain a number of polyketide metabolites (such as *O*- and/or *C*-glucosides) belonging to the families of 6-phenyl-2-pyrones, 5-methyl-7-hydroxychromones, and 1,8-dihydroxyanthrones (see Ref. 3 for a complete list of such compounds).

The structure of the new product, named kenyaloside (**1**), was determined by spectral and chemical methods. To our knowledge, it represents the first example of a naphthalene glycoside both occurring in *Aloe* species and bearing three different *O*-glycosyl residues [4, 5].

The aqueous extract of the dried exudate of Kenyan *Aloe* species, after partitioning with ethyl acetate, was lyophilized to afford a residue that was chromatographed successively on silica gel and Sephadex LH-20 columns. Kenyaloside (**1**) was obtained in *ca.* 0.1% yield (based on the starting drug). Its molecular formula, C₃₀H₄₀O₁₇, was derived from ESI-HRMS (found: *m/z* 695.21326, calcd for [M+Na⁺] *m/z*: 695.21577). The presence of three *O*-glycosyl residues was suggested by inspection of chemical shifts and coupling constants in the ¹H and ¹³C NMR spectra of **1** (Table 1); in addition, the NOESY spectrum revealed two significant associations between the anomeric proton at δ 5.77 and the upfield aromatic proton, and between another anomeric proton (at δ 4.31) and both the aromatic proton at δ 7.48 and an Ar-CH₂ group (AB system: δ 4.73, 4.94, *J* = 12.4 Hz). ¹H and ¹³C signals due to four aromatic C-H groups, together with the values of ¹H-¹H coupling constants and mutual NOEs, were indicative of a 1,2,3,8-tetrasubstituted naphthalene nucleus.

[¶]Part 19 in the series "Studies on Aloe". For Part 18, see Ref. 1

Table 1: NMR data of compounds **1** and **2** in CD₃OD at 400 MHz (¹H) and 100 MHz (¹³C).^{a, b}

C/H position	kenyaloside (1)			Compound 2		
	δ_H (J, Hz)	δ_C	Selected ¹ H- ¹ H NOEs	δ_H (J, Hz)	δ_C	Selected ¹ H- ¹ H NOEs
1		152.69			152.56	
2		123.06			122.96	
3		134.10			134.12	
4	7.48 s	119.77	7.52; 4.73, 4.94; 4.31	7.47 s	119.73	7.51; 4.71, 4.93
4a		137.03			137.05	
5	7.52 d (8.0)	122.94	7.48; 7.45	7.51 dd (8.3, 1.1)	122.83	7.47; 7.45
6	7.45 dd (7.6, 8.0)	127.97	7.52; 7.33	7.45 dd (7.8, 8.3)	127.97	7.51; 7.32
7	7.33 d (7.6)	109.92	7.45; 5.77	7.32 dd (7.8, 1.1)	109.91	7.45; 5.75
8		153.68			153.66	
8a		114.74			114.71	
COCH ₃		206.70			206.59	
COCH ₃	1.95 s	30.84		1.89	31.62	
CH ₂ O	4.73 d (12.4)	68.80	4.31, 7.48	4.71 d (12.4)	68.78	4.27, 7.47
	4.94 d (12.4)			4.93 d (12.4)		
1'	4.39 d (8.0)	102.46				
2'	3.23 dd (8.0, 8.8)	73.67				
3'	3.33 m ^c	76.85 ^c				
4'	3.33 m ^c	70.67 ^d				
5'	3.33 m ^c	77.10 ^c				
6'	3.67 dd (5.2, 12.0)	61.64				
	3.89 dd (1.6, 12.0)					
1''	5.77 d (1.8)	100.85	7.33	5.75 d (1.9)	100.83	7.32
2''	4.21 dd (1.8, 3.4)	70.56		4.20 dd (1.9, 3.5)	70.66	
3''	3.85 dd (3.4, 9.2)	71.58		3.84 dd (3.5, 9.3)	71.55	
4''	3.57 t (9.2)	72.42		3.56 t (9.3)	72.40	
5''	3.71 m	70.72 ^d		3.70 m	70.71	
CH ₃ (5'')	1.31 d (6.4)	17.02		1.29 d (6.1)	17.02	
1'''	4.31 d (7.2)	103.09	4.73, 4.94, 7.48	4.27 d (7.3)	103.30	4.71, 4.93
2'''	3.29 dd (7.2, 8.8)	73.82		3.23 dd (7.3, 9.0)	73.97	
3'''	3.50 t (8.8)	75.11		3.35 t (9.0)	76.85	
4'''	3.71 m	77.65		3.50 m	70.23	
5'''	3.33 m ^c	63.58		3.19 dd (10.1, 11.5)	65.94	
	4.06 dd (5.2, 12.0)			3.88 dd (5.4, 11.5)		

^a Spectra recorded at 40°C; ^b all assignments were based on extensive 1D and 2D NMR measurements (COSY, TOCSY, NOESY, APT, HMQC and HMBC); ^{c, d} signals with the same superscript are interchangeable; ^e covered by the CH₃OH signal.

heating at 150°C. Silica gel 60, 63-200 µm and 40-63 µm (Merck) was used for column and flash chromatography, respectively.

Plant material: The commercial exudate of Kenyan *Aloe* species used in this investigation was purchased from Sessa Carlo spa (Sesto S. Giovanni, Italy). A voucher specimen is kept at the Dipartimento di Chimica Organica e Industriale, Università di Milano

Extraction and isolation: The dried exudate of Kenyan *Aloe* species (250 g) was finely powdered and extracted with water (750 mL) with vigorous mechanical stirring for 24 h at room temperature. After filtration of the insoluble material, the aqueous solution was partitioned with ethyl acetate (2 x 1 L) and lyophilized to give a brown residue (120 g). Of this residue, 40 g was adsorbed onto sea sand and fractionated by flash chromatography (silica gel, 1.5 Kg) eluting with EtOAc containing increasing amounts of MeOH. Separation was monitored by TLC (eluent *A*) and fractions containing **1** (R_f 0.38) were combined, concentrated (3.5 g) and further purified by flash chromatography (silica gel, 500 g) eluting with EtOAc-EtOH-H₂O, 100:20:10. Fractions

were combined on the basis of TLC analysis (eluent *A*) and evaporated to dryness. The residue (*ca.* 400 mg) was chromatographed over a Sephadex LH-20 column eluted with MeOH-H₂O (1:1) to give kenyaloside (**1**) (200 mg, 0.08% overall yield) as an amorphous powder, pure by TLC (eluent *A*).

Kenyaloside [1-(β-D-glucopyranosyloxy)-8-(α-L-rhamnopyranosyloxy)-3-(β-D-xylopyranosyloxy)methyl)naphthalene (**1**)]

[α]_D: - 84.4° (*c* 0.25, MeOH).

R_f: 0.38 (AcOEt-EtOH-H₂O, 100:20:13).

IR (KBr): 1695, 1652, 1615 cm⁻¹.

UV/Vis λ_{max} (MeOH) nm (log ε): 226 (4.72), 260 (4.36), 290sh (4.30), 338 (3.94) [for dimethyl dianellidin (**3**) [6]: 223 (4.68), 253 (4.04), 331 (3.61)].

¹H NMR (400 MHz, CD₃OD): Table 1.

¹³C NMR (100 MHz, CD₃OD): Table 1.

ESI-HRMS: *m/z* [M + Na⁺] calcd for C₃₀H₄₀NaO₁₇ 695.21577, found 695.21326.

ESI MS: *m/z* 695 [M + Na⁺], 549 [M-146+Na⁺].

Enzymatic hydrolyses: β-Glucosidase (almond emulsin, Sigma, 30 mg) was added to a solution of

kenyaloside (**1**, 50 mg) in H₂O (25 mL), and the mixture was incubated at 37° for 3 h under nitrogen. After adding MeOH, the solution was filtered and concentrated under reduced pressure. Column chromatography of the aqueous residue (eluent: EtOAc-EtOH-H₂O, 100:20:13) gave two fractions. The less polar fraction, after further purification by column chromatography eluting with AcOEt-EtOH (from 5:1 to 1:1), furnished the diglycoside **2** (30 mg, 79%) as a pale yellow powder. The more polar fraction was submitted to column chromatography (eluent AcOEt-EtOH, from 3:1 to 1:1) to give glucose (9 mg, 67%), identified by TLC comparison with an authentic sample (eluent *B*, R_f 0.31).

Compound **2** (20 mg), dissolved in 50 mM phosphate buffer pH 6 (5 mL), was incubated with α -rhamnosidase from *Fusarium oxysporum* CCF 906 (0.2 U) [11] at 35°C for 24 h. After concentration, MeOH was added, the precipitate removed by filtration and the solvent evaporated under reduced pressure. Repeated column chromatographic purification (eluent AcOEt-EtOH, from 5:1 to 1:1 and from 3:1 to 1:1) furnished rhamnose (5 mg, R_f 0.62, eluent *B*, co-TLC with an authentic sample) and a yellow product (10 mg, R_f 0.73, eluent *A*). This was hydrolyzed without further purification (0.1 N HCl, dioxane-H₂O, 1:1, 5 mL; 70°C, 5 h, under nitrogen) to give xylose (R_f 0.50, eluent *B*, co-TLC

with an authentic sample), which was purified (1.5 mg) by column chromatography (eluent AcOEt-EtOH, from 3:1 to 1:1).

8-(α -L-Rhamnopyranosyloxy)-3-(β -D-xylopyranosyloxymethyl)naphthalen-ol (**2**)

$[\alpha]_D$: - 36.2° (*c* 0.07, MeOH).

R_f: 0.54(AcOEt-EtOH-H₂O, 100:20:13).

IR (KBr): 1635 cm⁻¹.

UV/Vis λ_{\max} (MeOH) nm (log ϵ): 225 (4.49), 258sh (4.09), 297 (3.99), 334 (3.80).

¹H NMR (400 MHz, CD₃OD): Table 1.

¹³C NMR (100 MHz, CD₃OD): Table 1.

ESI MS: *m/z* 533 [M + Na⁺], 387 [M-146+Na⁺].

Determination of the absolute configuration of the isolated sugars: The isolated monosaccharides were converted into methyl glycopyranosides followed by treatment with excess *p*-bromobenzoyl chloride, as in ref. 9. Comparison of the CD spectra of the resulting per-*p*-bromobenzoates with those of the analogous derivatives prepared from authentic samples allowed the D-configuration for glucose and xylose and the L-configuration for rhamnose to be established.

Acknowledgments – Thanks are due to Dr Lavinia Duri for technical assistance and to MIUR for financial support.

References

- [1] Speranza G, Morelli CF, Tubaro A, Altinier G, Duri L, Manitto P. (2005) Aloeresin I, an anti-inflammatory 5-methylchromone from Cape aloe. *Planta Medica*, **71**, 79-81.
- [2] Trease GE, Evans WC. (1983) *Pharmacognosy*, Baillière Tindall, London, 404-408.
- [3] Duri L, Morelli CF, Crippa S, Speranza G. (2004) 6-Phenylpyrones and 5-methylchromones from Kenya aloe. *Fitoterapia*, **75**, 520-522.
- [4] Reynolds T. (2004) Aloe chemistry. In *Aloes. The genus Aloe*. Reynolds T (Ed). CRC Press, Boca Raton, USA. 39-74.
- [5] Dagne E, Bisrat D, Viljoen A, Van Wyk B-E. (2000) Chemistry of *Aloe* species. *Current Organic Chemistry*, **4**, 1055-1078.
- [6] Batterham T, Cooke RG, Duewell H, Sparrow LG. (1961) Colouring matters of Australian plants. VIII. Naphthalene derivatives from *Dianella* species. *Australian Journal of Chemistry*, **14**, 637-642.
- [7] Speranza G, Corti S, Manitto P. (1994) Isolation and chemical characterization of a new constituent of Cape aloe having the 1,1-diphenylethane skeleton. *Journal of Agricultural and Food Chemistry*, **42**, 2002-2006.
- [8] (a) Buckingham J. (2005) *Dictionary of Natural Products on CD-ROM*. Chapman & Hall/CRC, England; (b) Editorial (1997) *Planta Medica*, **63**, 195.
- [9] Golik J, Liu H-W, Dinovi M, Furukawa J, Nakanishi K. (1983) Characterization of methyl glycosides at the pico- to nano-gram level. *Carbohydrate Research*, **118**, 135-146.
- [10] Nakanishi K, Kuroyanagi M, Nambu H, Oltz EM, Takeda R, Verdine GL, Zask A. (1984) Recent application of circular dichroism to structural problems, especially oligosaccharide structures. *Pure and Applied Chemistry*, **56**, 1031-1048.
- [11] Monti D, Pišvejcová A, Křen V, Lama M, Riva S. (2004) Generation of an α -L-rhamnosidase library and its application for the selective derhamnosylation of natural products. *Biotechnology and Bioengineering*, **87**, 763-771.
- [12] Pham TN, Hinchley SL, Rankin DWH, Liptaj T, Uhrinpp D. (2004) Determination of sugar structures in solution from residual dipolar coupling constants: methodology and application to methyl β -D-xylopyranoside. *Journal of the American Chemical Society*, **126**, 13100-13110.

New Flavonoid Glycosides from *Chrozophora senegalensis* and Their Antioxidant Activity

Antonio Vassallo^a, Giuseppina Cioffi^a, Francesco De Simone^a, Alessandra Braca^b, Rokia Sanogo^c, Angelo Vanella^d, Alessandra Russo^d and Nunziatina De Tommasi^{a*}

^aDipartimento di Scienze Farmaceutiche, Università di Salerno, Via Ponte Don Melillo, 84084 Fisciano, Salerno, Italy

^bDipartimento di Chimica Bioorganica e Biofarmacia, Università di Pisa, Via Bonanno 33, 56126 Pisa, Italy

^cDepartement Medicine Traditionelle (DMT), INRSP, B.P. 1746, Bamako, Mali

^dDipartimento di Chimica Biologica, Chimica Medica e Biologia Molecolare, Università di Catania, v.le A. Doria 6, 95125 Catania, Italy

detommasi@unisa.it

Received: June 27th, 2006; Accepted: September 27th, 2006

Dedicated to the memory of Professor Ivano Morelli.

Bioassay-directed fractionation of an antioxidant methanol extract of the leaves of *Chrozophora senegalensis* using DPPH assay led to the isolation of three new flavonoid glycosides, quercetin 3-*O*-(6"-caffeoyl)- β -D-glucopyranoside-3'-*O*- β -D-glucopyranoside (**1**), quercetin 3-methyl ether-7-*O*- α -L-rhamnopyranosyl-(1 \rightarrow 6)-(2"-*p*-coumaroyl)- β -D-glucopyranoside (**2**), acacetin 7-*O*-(6"-*p*-coumaroyl)- β -D-glucopyranoside (**3**), along with five known flavonoids, one phenolic derivative, and three megastigmane glycosides. Their structures were established on the basis of detailed spectral analysis. All isolated compounds were tested for their antioxidant activity on DPPH stable radical, superoxide anion, metal chelating activity, and DNA cleavage induced by the photolysis of H₂O₂. Quercetin 3-*O*-(6"-caffeoyl)- β -D-glucopyranoside-3'-*O*- β -D-glucopyranoside (**1**), quercetin 3'-methyl ether-3-*O*- α -L-rhamnopyranoside (**4**), and 4"-methyl ether amenthoflavone (**9**) exhibited the highest antioxidant capacity being also able to modulate hydroxyl radical formation more efficiently than other compounds acting as direct hydroxyl radical scavengers and chelating iron.

Keywords: *Chrozophora senegalensis*, Euphorbiaceae, flavonoids, antioxidant activity.

In recent years, a global trend toward the use of natural phytochemicals present in herbs and functional foods as antioxidants was further increased after that it had been reported that some commonly used synthetic antioxidant compounds, such as butylated hydroxytoluene (BHT) and butylated hydroxyanisole, have long-term toxicological effects, including carcinogenicity [1]. Of particular interest as possible sources of natural antioxidants are medicinal plants traditionally used to treat conditions related to oxidative stress, such as rheumatism and inflammation. In this regard, many phytochemicals with diversified biological properties have shown

promise for the prevention and/or treatment of all diseases in which oxidative stress plays a key role [2]. *Chrozophora senegalensis* (Lam) A Juss. ex Spreng, syn. *Croton senegalensis* (Euphorbiaceae family) is a small tree widely distributed in Mali where it grows wild and is used in folk medicine for the treatment of diarrhea, rheumatism, teniasis, stomachache, rachitis, and venereal diseases. The leaf and root decoctions are also drunk for hairloss [3, 4]. To confirm the use of *C. senegalensis* in Malian traditional medicine, the extracts of the leaves were evaluated for *in vitro* antioxidant activity. A bioassay-guided fractionation procedure showed that

the methanol extract was the active one, while all the other residues were inactive (data not shown). Subsequent fractionation and analysis of the methanol extract led to the isolation and structural characterization of three new flavonoids (**1-3**), together with some known compounds, including five flavonoids (**4-7** and **9**), one phenolic derivative (**8**), and three megastigmane glycosides (**10-12**).

Compound **1** was isolated as a yellow amorphous powder. Its molecular formula was established as $C_{36}H_{36}O_{20}$ by means of ESI-MS ($[M-H]^-$ peak at m/z 787), ^{13}C , ^{13}C -DEPT NMR, and elemental analysis. Analysis of 600 MHz NMR spectra suggested a flavonoid skeleton for compound **1**. The 1H -NMR spectrum (Table 1) indicated a 5,7-dihydroxylated pattern for ring A (two *meta*-coupled doublets at δ 6.16 and 6.33, $J = 1.5$ Hz) and a 3',4'-dihydroxylation pattern for ring B (ABX system signals at δ 6.80, d, $J = 8.5$ Hz; 7.58, dd, $J = 8.5$, 2.5 Hz; 7.67, d, $J = 2.5$ Hz), allowing the aglycon to be recognized as quercetin [5]. The 1H -NMR spectrum of **1** also showed signals ascribable to sugar moieties and a caffeoyl residue (Table 1). Two anomeric protons arising from the sugar moieties appeared at δ 5.26 and 4.88 each (1H, d, $J = 7.5$ Hz), which correlated respectively with signals at δ 103.4 and 104.7 ppm in the HSQC spectrum. All the 1H - and ^{13}C -NMR signals of **1** were assigned using 1D-TOCSY, DQF-COSY, HSQC, and HMBC experiments. Complete assignments of proton and carbon chemical shifts of the sugar portion were accomplished by DQF-COSY and 1D-TOCSY experiments and allowed the identification of the sugars as two terminal β -D-glucopyranosyl units. The configurations of the sugar units were assigned after hydrolysis of **1** with 1 N HCl. The hydrolysate was trimethylsilylated, and GC retention times compared with those of authentic sugar samples prepared in the same manner. The lower field shifts of H_2-6''' (δ 4.32 and 4.23) of one glucosyl unit suggested the substitution site of the caffeoyl moiety. Unequivocal information could be obtained by 2D-NMR spectra; the HMBC experiment indicated correlations between δ 5.26 (H-1''') and 135.6 (C-3), δ 4.88 (H-1'') and 149.0 (C-3'), δ 4.32 and 4.23 (H_2-6''') and 170.0 (COO). Thus, the structure of **1** was determined as quercetin 3-*O*-(6''-caffeoyl)- β -D-glucopyranoside-3'-*O*- β -D-glucopyranoside.

The molecular formula $C_{37}H_{38}O_{18}$ for compound **2** was determined by ESI-MS ($[M-H]^-$ at m/z 769), ^{13}C ,

Table 1: 1H and ^{13}C NMR data of compound **1** (CD_3OD , 600 MHz)^a.

position	δ_H	δ_C
2		159.0
3		135.6
4		179.0
5		163.5
6	6.16 d (1.5)	100.0
7		166.3
8	6.33 d (1.5)	94.2
9		159.0
10		105.8
1'		123.1
2'	7.67 d (2.5)	117.2
3'		149.0
4'		146.4
5'	6.80 d (8.5)	116.0
6'	7.58 dd (2.5, 8.5)	123.5
3'-O-Glc 1''	4.88 d (7.5)	104.7
2''	3.58 dd (7.5, 9.0)	74.8
3''	3.52 t (9.0)	77.3
4''	3.42 t (9.0)	71.2
5''	3.53 m	78.4
6''a	3.95 dd (5.0, 12.0)	62.4
3-O-Glc 1'''	5.26 d (7.5)	103.4
2'''	3.56 dd (7.5, 9.0)	73.6
3'''	3.49 t (9.0)	77.7
4'''	3.40 t (9.0)	71.8
5'''	3.59 m	75.6
6'''a	4.32 dd (5.0, 12.0)	64.2
trans-caffeoyl 1		128.4
2	7.00 d (1.5)	115.4
3		147.5
4		150.1
5	6.81 d (8.8)	116.2
6	6.82 dd (1.5, 8.8)	123.1
α	6.07 d (16.0)	114.6
β	7.39 d (16.0)	147.4
COO		170.0

^a Coupling pattern and coupling constants (J in Hertz) are in parentheses.

^{13}C -DEPT NMR analyses and was supported also by elemental analysis. Its 1H - and ^{13}C -NMR spectra (see Table 2) indicated that it was a quercetin 3-methyl ether derivative [5]. Its 1H -NMR spectrum further displayed signals for two sugar residues that were easily clarified with the help of 1D-TOCSY and DQF-COSY experiments, leading to the identification of one β -D-glucopyranosyl and one α -L-rhamnopyranosyl residue. The configuration of sugar units was determined as reported for compound **1**. The presence of one *p*-coumaroyl moiety was shown in the 1H -NMR spectrum by the signals at δ 7.45 and 6.73 each (2H, d, $J = 8.5$ Hz) and δ 7.41 and 6.38 each (1H, d, $J = 16.0$ Hz). The HSQC spectrum showed glycosidation shifts for C-6'' (δ 67.5) and acylation shift for H-2'' (δ 4.74) and C-2'' (δ 74.5) of the β -D-glucopyranosyl unit. An unambiguous determination of the sequence and linkage sites was obtained from an HMBC experiment, showing cross peak correlations between δ 5.06 (H-1'') and 164.5




Figure 1: Structures of compounds 1-3.

Table 2: ¹H- and ¹³C-NMR data of compounds 2-3 (CD₃OD, 600 MHz)^a.

position	2	3
	δ _H	δ _C
2		157.9
3		139.9
4		180.0
5		164.3
6	6.54 d (2.0)	101.2
7		164.5
8	6.73 d (2.0)	95.8
9		158.8
10		107.1
1'		123.6
2'	7.71 d (1.5)	116.2
3'		145.0
4'		149.6
5'	6.90 d (8.0)	117.3
6'	7.65 dd (1.5, 8.0)	123.3
OMe	3.90 s	56.1
7-O-Glc 1''	5.06 d (7.5)	100.0
2''	4.74 dd (7.5, 9.0)	74.5
3''	3.47 t (9.0)	77.0
4''	3.45 t (9.0)	71.0
5''	3.30 m	77.7
6''a	4.00 dd (5.0, 12.0)	67.5
6''b	3.60 dd (3.0, 12.0)	
Rha 1'''	4.80 d (1.5)	101.9
2'''	3.94 dd (1.5, 3.4)	72.2
3'''	3.88 dd (3.4, 9.5)	71.8
4'''	3.55 t (9.0)	74.5
5'''	4.20 m	69.6
6'''	1.12 d (6.5)	17.6
p-coumaroyl 1		124.9
2,6	7.45 d (8.5)	130.2
3,5	6.73 d (8.5)	116.5
4		161.0
α	6.38 d (16.0)	118.0
β	7.41 d (16.0)	146.8
COO		168.7

^a Coupling pattern and coupling constants (*J* in Hertz) are in parentheses.

(C-7), δ 4.74 (H-2'') and 168.7 (COO), and δ 4.80 (H-1''') and 67.5 (C-6''). Therefore, the structure quercetin 3-methyl ether-7-*O*-α-L-rhamnopyranosyl-(1→6)-(2''-*p*-coumaroyl)-β-D-glucopyranoside was assigned to compound 2.

Compound 3 was obtained as a yellow amorphous powder and its ESI-MS showed an [M-H]⁻ ion peak at *m/z* 591. The molecular formula C₃₁H₂₈O₁₂ was

confirmed by elemental analysis. In the ¹H-NMR spectrum (Table 2) two singlets at δ 6.70 and 3.92, two doublets at δ 6.77 and 6.56 each (1H, d, *J* = 2.0 Hz), and two *o*-coupled protons at δ 7.94 and 7.06 each (2H, d, *J* = 8.5 Hz) were present permitting the identification of the aglycon as apigenin 4'-methyl ether or acacetin [5]. Additionally for 3, resonances of one anomeric proton and one *p*-coumaroyl residue were observed in the ¹H-NMR spectrum at δ 5.05

(1H, d, $J = 7.5$ Hz), 7.45 and 6.75 each (2H, d, $J = 8.5$ Hz) and δ 7.43 and 6.38 each (1H, d, $J = 16.0$ Hz), respectively. 1D-TOCSY, DQF-COSY, and HSQC NMR experiments showed the presence of one β -D-glucopyranosyl unit characterized by an acylation shift at H₂-6 (δ 4.64 and 4.25). The configuration of the glucose unit was determined as reported for compound **1**. HMBC correlations confirmed the substitution sites of each residue allowing compound **3** to be identified as acacetin 7-*O*-(6"-*p*-coumaroyl)- β -D-glucopyranoside.

Compounds **4-12** were identified by 1D- and 2D-NMR spectroscopy and ESI-MS analysis and by comparison of their data with those reported in the literature [9-14] as quercetin 3'-methyl ether-3-*O*- α -L-rhamnopyranoside (**4**), quercetin 3'-methyl ether-3-*O*- α -L-rhamnopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside (**5**), apigenin 7-*O*-(6"-*p*-coumaroyl)- β -D-glucopyranoside (**6**), quercetin 3-methyl ether-7-*O*- α -L-rhamnopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside (**7**), 4-hydroxyphenyl-*O*- α -L-rhamnopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside (**8**), 4'''-methyl ether amenthoflavone (**9**), roseoside (**10**), icaraside B5 (**11**), and ampelopsionoside (**12**).

Table 3: Scavenger effect on DPPH stable radical and superoxide anion of methanol fractions and compounds **1-12** isolated from *C. senegalensis*.

Fracts or Compds	DPPH Test ^a IC ₅₀ (μg/ml) ± ^b SD	Effect on O ₂ ⁻ ^a IC ₅₀ (μg/ml) ± ^b SD
A	178 ± 6.7	0.61 ± 0.04
B	14.22 ± 1.1	2.6 ± 0.35
C	7.01 ± 0.6	0.37 ± 0.03
D	6.47 ± 1.5	0.19 ± 0.05
E	4.56 ± 0.8	0.36 ± 0.02
F	25.65 ± 3.6	0.47 ± 0.03
1	9.75 ± 0.9	0.085 ± 0.002
2	1.08 ± 0.4	0.025 ± 0.003
3	61.59 ± 2.5	2.5 ± 0.4
4	6.69 ± 0.7	0.20 ± 0.01
5	52 ± 0.5	0.85 ± 0.06
6	110 ± 24	1.35 ± 0.09
7	94.33 ± 0.7	0.42 ± 0.05
8	-	-
9	4.31 ± 1.1	2.76 ± 0.01
10	527 ± 0.4	50 ± 0.4
11	32 ± 0.5	0.5 ± 0.01
12	25 ± 0.9	0.015 ± 0.03
^c Trolox	96 ± 1.7	-
^d SOD	-	89 ± 1.5

^aconcentration that inhibited radicals by 50%.

^bn = 6.

^cTrolox (50 μM) and ^dsuperoxide dismutase (SOD) (80 mU/mL) were used as standard; the results are expressed as % of inhibition.

The preliminary *in vitro* biological analysis indicated that compounds **1-7** and **9-12** were able to quench DPPH radicals and exhibited a direct scavenging activity on superoxide anion; this radical was in fact produced by the reduction of β -mercaptoethanol,

excluding the Fenton-type reaction and the xanthine/xanthine oxidase system (Table 3). Quercetin 3-*O*-(6"-caffeoyl)- β -D-glucopyranoside-3'-*O*- β -D-glucopyranoside (**1**), quercetin 3'-methyl ether-3-*O*- α -L-rhamno-pyranoside (**4**), and 4'''-methyl ether amenthoflavone (**9**) exhibited the highest antioxidant capacity. On the other hand, the potent biological activity of quercetin is largely reported in literature [15].

Although both O₂⁻ and H₂O₂ are potentially cytotoxic, most of the oxidative damage in biological systems is caused by the OH radical, which is generated by the reaction between O₂⁻ and H₂O₂ in the presence of transition metal ions [2]. In fact, the OH radical can react with a number of target molecules including proteins, membrane lipids, and DNA.

Table 4: Effect of methanol fractions and compounds **1-12** isolated from *C. senegalensis* (100 μg/mL) on DNA cleavage induced by the photolysis of H₂O₂ and metal chelating activity.

	UD of supercoiled DNA (% of native DNA)	Ferrozine assay ^a IC ₅₀ (μg/mL) ± ^b SD
scDNA	100	
A	11 ± 2.4*	-
B	62.7 ± 3.7*	32 ± 4.5
C	78 ± 4.5*	47.6 ± 3.6
D	76.7 ± 2.6*	16.83 ± 2.5
E	95 ± 4.7*	19.74 ± 3.2
F	65 ± 4.6*	28.41 ± 0.9
1	36 ± 1.2*	13.65 ± 2.8
2	7 ± 1.6*	92 ± 1.9
3	9 ± 2.4*	25 ± 2.5
4	70 ± 2.7*	6.19 ± 0.19
5	10 ± 0.8*	44.64 ± 3.6
6	5 ± 0.9*	-
7	37 ± 0.6*	630 ± 67
8	3.4 ± 0.4*	625 ± 50
9	73 ± 4.7*	18.31 ± 2.4
10	2.6 ± 0.6*	-
11	15.3 ± 1.1*	222 ± 32
12	11.3 ± 3.1*	-
DMSO	75.3 ± 3.1*	-
DTPA	-	77.5 ± 2.3

The hydroxyl radicals generated by the photolysis of H₂O₂ inhibited the supercoiled DNA (SCDNA). Each value represents the mean ± SD of three experiments. *Significant vs. supercoiled DNA ($p < 0.001$).

^aconcentration that inhibited the ferrozine-Fe²⁺ formation by 50%.
^bn = 6.

DTPA (5 μM) and DMSO (1mM) were used as standard; ^cthe result is expressed as % of inhibition.

Based on the data obtained from this study, compounds **1**, **4**, and **9** might also be able to modulate hydroxyl radical formation more efficiently than other compounds acting as direct scavengers and chelating iron. In fact, these natural compounds exhibited a more efficient protection against DNA strand scission induced by OH radicals generated by UV-photolysis of H₂O₂ (Table 4), and showed metal

chelating activity capturing ferrous ions before ferrozine, with an IC_{50} value (concentration that inhibited the ferrozine- Fe^{2+} by 50%) of 13.65, 6.19 and 18.31 $\mu\text{g/mL}$, respectively (Table 4).

These data also suggest that the biological effect of *C. senegalensis* observed from ethnopharmacological studies is due in part to the anti-oxidant action of its active components.

Experimental

General: Optical rotations were measured on a Perkin-Elmer 241 polarimeter equipped with a sodium lamp (589 nm) and a 10 cm microcell. Elemental analysis was obtained from a Carlo Erba 1106 elemental analyzer. UV spectra were recorded on a Perkin-Elmer-Lambda 12 spectrophotometer. A Bruker DRX-600 NMR spectrometer using the UXMNMR software package was used for NMR experiments. ESIMS (negative mode) were obtained using a Finnigan LC-Q Advantage Termoquest spectrometer, equipped with Xcalibur software. TLC was performed on precoated Kieselgel 60 F_{254} plates (Merck, Darmstadt, Germany); compounds were detected by spraying with $Ce(SO_4)_2/H_2SO_4$ (Sigma-Aldrich, St. Louis, Mo, USA) and NTS (Naturstoffe reagent)-PEG (Polyethylene glycol 4000) solutions. Column chromatography was performed over Sephadex LH-20 (Pharmacia); reversed-phase (RP) HPLC separations were conducted on a Waters 515 pumping system equipped with a Waters R401 refractive index detector and Waters U6K injector, using a C_{18} μ -Bondapak column (30 cm x 7.8 mm) and a mobile phase consisting of MeOH- H_2O mixtures at a flow rate of 2 mL/min. GC analyses were performed using a Dani GC 1000 instrument. A Hitachi U-2000 spectrophotometer (Hitachi, Tokyo, Japan) was used for all antioxidant assays.

Plant material and chemicals: The leaves of *Chrozophora senegalensis* were collected in Bandiagara, Mali, in 1999 and identified by Prof. N'Golo Diarra of the Departement Medicine Traditionnelle (DMT), Bamako, Mali where a voucher specimen (DMT n. 0074) is deposited. pBR322 plasmid DNA, 1,1-diphenyl-2-picryl-hydrazyl radical (DPPH), diethylenetriaminepentaacetic acid (DTPA) and 3-(2-pyridyl)-5,6-bis (4-phenyl-sulfonic acid)-1,2,4-triazine (ferrozine) were obtained from Sigma Aldrich Co (St. Louis, USA); β -nicotinamide-adenine dinucleotide (NADH) was obtained from Boehringer Mannheim GmbH (Germany). All other chemicals

were purchased from GIBCO BRL Life Technologies (Grand Island, NY, USA).

Extraction and isolation: The air-dried powdered leaves of *C. senegalensis* (600 g) were defatted with *n*-hexane and extracted successively by exhaustive maceration (3 x 1 L, for 48 h) with $CHCl_3$, $CHCl_3$ -MeOH 9:1, and MeOH. The extracts were concentrated under reduced pressure to afford 13.4, 14.0, 13.8, and 62.4 g of dried residues, respectively. A portion of the MeOH extract (27.0 g) was partitioned between *n*-BuOH and H_2O to give a *n*-BuOH soluble portion (9.0 g); 5.0 g of this residue were chromatographed over a Sephadex LH-20 column (100 cm x 5 cm) with MeOH as the eluent. A total of 115 fractions were collected (10 mL each). These were combined according to TLC analysis [silica 60 F_{254} gel-coated glass sheets with *n*-BuOH-AcOH- H_2O (60:15:25) and $CHCl_3$ -MeOH- H_2O (40:9:1)] to give nine pooled fractions (A-I). Fractions G, H, and I yielded compounds **3** (19.2 mg), **4** (40 mg), and **9** (30 mg), respectively. Fraction A (90 mg) was purified by RP-HPLC using MeOH- H_2O (45:55) to give compounds **10** (6 mg, t_R = 10 min) and **12** (5 mg, t_R = 20 min). Fraction B (36 mg) was purified by RP-HPLC using MeOH- H_2O (1:1) to give compounds **2** (8 mg, t_R = 10 min) and **11** (12 mg, t_R = 20 min). Fraction C (50.5 mg) was purified by RP-HPLC using MeOH- H_2O (45:55) to give compounds **5** (28 mg, t_R = 10 min) and **7** (10.8 mg, t_R = 20 min). Fraction D (100 mg) was purified by RP-HPLC using MeOH- H_2O (45:55) to give compounds **1** (14.5 mg, t_R = 10 min) and **6** (6.5 mg, t_R = 20 min), while fraction E (70 mg) was purified by RP-HPLC using MeOH- H_2O (55:45) to yield compound **3** (11 mg, t_R = 10 min). Finally, fraction F (85 mg) was chromatographed on a RP-HPLC using MeOH- H_2O (1:1) as the eluent to afford compounds **8** (5 mg, t_R = 28 min) and **4** (6.3 mg, t_R = 46 min).

Quercetin 3-*O*-(6''-caffeoyl)- β -D-glucopyranoside-3'-*O*- β -D-glucopyranoside (1)

Yellow amorphous powder.

$[\alpha]_D$: -27° (c 0.1, MeOH).

UV/Vis λ_{max} (MeOH) nm (log ϵ): 267 (3.99), 344 (4.32)

1H NMR (600 MHz, CD_3OD): Table 1.

^{13}C NMR (600 MHz, CD_3OD): Table 1.

ESIMS: m/z 787 $[M - H]^-$.

Anal. Calcd for $C_{36}H_{36}O_{20}$: C, 54.83; H, 4.60. Found C, 54.79; H 4.62.

Quercetin 3-methyl ether-7-O- α -L-rhamnopyranosyl-(1 \rightarrow 6)-(2''-p-coumaroyl)- β -D-glucopyranoside (2)

Yellow amorphous powder.

$[\alpha]_D$: +18° (c 0.1, MeOH).

UV/Vis λ_{\max} (MeOH) nm (log ϵ): 265 (3.92), 356 (4.05).

^1H NMR (600 MHz, CD_3OD): Table 2.

^{13}C NMR (600 MHz, CD_3OD): Table 2.

ESIMS: m/z 769 $[\text{M} - \text{H}]^-$.

Anal. Calcd for $\text{C}_{37}\text{H}_{38}\text{O}_{18}$: C, 57.66; H, 4.97. Found C, 57.68; H 5.00.

Acacetin 7-O-(6''-p-coumaroyl)- β -D-glucopyranoside (3)

Yellow amorphous powder.

$[\alpha]_D$: +11° (c 0.1, MeOH).

UV/Vis λ_{\max} (MeOH) nm (log ϵ): 269 (3.99), 321 (3.76).

^1H NMR (600 MHz, CD_3OD): Table 2.

^{13}C NMR (600 MHz, CD_3OD): Table 2.

ESIMS: m/z 591 $[\text{M} - \text{H}]^-$.

Anal. Calcd for $\text{C}_{31}\text{H}_{28}\text{O}_{12}$: C, 62.84; H, 4.76. Found C, 62.80; H 4.80.

Acid hydrolysis of compounds 1-3: A solution of each compound (**1-3**, 2.0 mg each) in 1 N HCl (1 mL) was stirred at 80°C in a stoppered reaction vial for 4 h. After cooling, the solution was evaporated under a stream of N_2 . Each residue was dissolved in 1-(trimethylsilyl)imidazole and pyridine (0.2 mL), and the solution was stirred at 60°C for 5 min. After drying the solution, the residue was partitioned between water and CHCl_3 . The CHCl_3 layer was analyzed by GC using an L-CP-Chirasil-Val column (0.32 mm x 25 m). Temperatures of the injector and detector were 200°C for both. A temperature gradient system was used for the oven, starting at 100°C for 1 min and increasing up to 180°C at a rate of 5°C/min. Peaks of the hydrolysate were detected by comparison with retention times of authentic samples of L-rhamnose and D-glucose (Sigma Aldrich) after treatment with 1-(trimethylsilyl)imidazole in pyridine.

Antioxidant activity in cell-free systems

Quenching of DPPH: The free radical-scavenging capacity of extracts, fractions and pure compounds was tested by their ability to bleach the stable 1,1-diphenyl-2-picrylhydrazyl radical (DPPH) [16]. The reaction mixture contained 86 μM DPPH and

different concentrations of the natural compounds in 1 mL of ethanol. After 10 min at room temperature the absorbance at $\lambda = 517$ nm was recorded. Trolox (50 μM), a water-soluble derivative of vitamin E, was used as a standard.

Scavenger effect on superoxide anion: Superoxide anion was generated *in vitro* as described by Paoletti *et al.* [17]. The assay mixture contained in a total volume of 1 mL, 100 mM triethanolamine-diethanolamine buffer, pH 7.4, 3 mM NADH, 25 mM/12.5 mM EDTA/ MnCl_2 , 10 mM β -mercaptoethanol; some samples contained the natural compounds at different concentrations. After 20 min incubation at 25°C, the decrease in absorbance was measured at $\lambda = 340$ nm. Superoxide dismutase (SOD) (80 mU/mL) was used as a standard.

DNA cleavage induced by hydrogen peroxide UV-photolysis: The experiments were performed, as previously reported [18], in a volume of 20 μL containing 33 μM in bp (base pair) of pBR322 plasmid DNA in 5 mM phosphate saline buffer (pH 7.4), and the natural compounds at different concentrations. Immediately prior to irradiating the samples with UV light, H_2O_2 was added to a final concentration of 2.5 mM. The reaction volumes were held in caps of polyethylene microcentrifuge tubes, placed directly on the surface of a transilluminator (8000 $\mu\text{W cm}^{-1}$) at 300 nm. The samples were irradiated for 5 min at room temperature. After irradiation 4.5 μL of a mixture, containing 0.25% bromophenol blue, 0.25% xylen cyanol FF, and 30% glycerol, were added to the irradiated solution. The samples were then analyzed by electrophoresis on a 1% agarose horizontal slab gel in Tris-borate buffer (45 mM Tris-borate, 1 mM EDTA). Untreated pBR322 plasmid was included as a control in each run of gel electrophoresis, conducted at 1.5 V/cm for 15 hours. Gel was stained in ethidium bromide (1 $\mu\text{g/mL}$; 30 min) and photographed on Polaroid-Type 667 positive land film. The intensity of each scDNA band was quantified by means of densitometry. Dimethylsulfoxide (DMSO) (1 mM) was used as a standard.

Metal chelating activity: The chelating of ferrous ions by fractions and pure compounds was estimated by the ferrozine assay [19]. Briefly, natural compounds were added to a solution of 0.15 mM FeSO_4 . The reaction was initiated by the addition of 0.5 mM ferrozine and the mixture was shaken

vigorously and left standing at room temperature for ten minutes. After the mixture had reached equilibrium, the absorbance of the solution was then measured spectrophotometrically at 562 nm. DTPA (5 μ M) was used as a standard.

Supplementary data: NMR spectral data for quercetin 3'-methyl ether-3-*O*- α -L-rhamno-pyranoside (4), quercetin 3'-methyl ether-3-*O*- α -L-rhamnopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside (5), apigenin 7-*O*-(6"-*p*-coumaroyl)- β -D-glucopyranoside (6), quercetin 3-methyl ether-7-*O*- α -L-rhamno-

pyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside (7), 4-hydroxyphenyl-*O*- α -L-rhamnopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside (8), 4'''-methyl ether amenthoflavone (9), roseoside (10), icariside B5 (11), and ampelopsionoside (12).

Acknowledgments - This work was supported by the "Bioactive Compounds from Medicinal and Food Plants of Developing Countries" project of the Italian Ministry for University and Research (Ministero dell'Università e della Ricerca, MIUR).

References

- [1] Devi RS, Narayan S, Mohan KV, Sabitha KE, Devi CS. (2003) Effect of a polyherbal formulation, Ambrex, on butylated hydroxy toluene (BHT) induced toxicity in rats. *Indian Journal of Experimental Biology*, **41**, 1294-1299.
- [2] Halliwell B, Gutteridge JMC. (1999) Free radicals in biology and medicine. In: *Studies of Generalized Light Emission (Luminescence/Fluorescence)*. 3rd Ed. Oxford: University Press, 387-388.
- [3] Kerharo J, Adam JC. (1974) *Pharmacopée Sénégalaise Traditionnelle: Plantes Médicinales et Toxicologiques*. Vigot et Frères Ed, Paris, 1011.
- [4] Neuwinger HD. (2000) *African Traditional Medicine. A Dictionary of Plant Use and Application*. Medpharm Scientific Publisher.
- [5] Agrawal PK. (1989) *Carbon-13 NMR of Flavonoids*. Elsevier Science, Amsterdam, 294-364.
- [6] Wolbis M. (1989) Flavonol glycosides from *Sedum album*. *Phytochemistry*, **28**, 2187-2189.
- [7] Markham KR, Ternai B, Stanley R, Geiger H, Mabry TJ. (1978) Carbon-13 NMR studies of flavonoids. III. Naturally occurring flavonoid glycosides and their acylated derivatives. *Tetrahedron*, **34**, 1389-1397.
- [8] Itokawa H, Suto K, Takeya K. (1981) Studies on a novel *p*-coumaroyl glucoside of apigenin and on other flavonoids isolated from patchouli (Labiatae). *Chemical & Pharmaceutical Bulletin*, **29**, 254-256.
- [9] Nawwar MAM, El-Mousallamy AMD, Barakat, HH, Buddrus J, Linscheid M. (1989) Flavonoid lactates from leaves of *Marrubium vulgare*. *Phytochemistry*, **28**, 3201-3206.
- [10] De Tommasi N, Pizza C, Aquino R, Cumandà J, Mahmood N. (1997) Flavonol and chalcone ester glycosides from *Bidens leucantha*. *Journal of Natural Products*, **60**, 270-273.
- [11] Markham KR, Sheppard C, Geiger H. (1987) Carbon-13 NMR of flavonoids. Part IV. Carbon-13 NMR studies of some naturally occurring amentoflavone and hinokiflavone biflavonoids. *Phytochemistry*, **26**, 3335-3337.
- [12] Otsuka H, Takeda Y, Yamasaki K, Takeda Y. (1992) Structural elucidation of dendranthemosides A and B: two new β -ionone glucosides from *Dendranthera shiwogiku*. *Planta Medica*, **58**, 373-375.
- [13] Miyase T, Ueno A, Takizawa N, Kobayashi H, Oguchi H. (1988) Studies on the glycosides of *Epimedium grandiflorum* Morr. var. *thunbergianum* (Miq.) Nakai. III. *Chemical & Pharmaceutical Bulletin*, **36**, 2475-2484.
- [14] Inada A, Nakamura Y, Konishi M, Murata H, Kitamura F, Toya H, Nakanishi T. (1991) A new ionone glucoside and a new phenylpropanoid rhamnoside from stems of *Ampelopsis brevipedunculata* (maxim.) Trautv. *Chemical & Pharmaceutical Bulletin*, **39**, 2437-2439.
- [15] Di Carlo G, Mascolo N, Izzo AA, Capasso F. (1999) Flavonoids: old and new aspects of a class of natural therapeutic drugs. *Life Sciences*, **65**, 337-353.
- [16] Bonina F, Saija A, Tomaino A, Lo Cascio R, Rapisarda P, Dederen JC. (1998) *In vitro* antioxidant activity and *in vivo* photoprotective effect of a red orange extract. *International Journal of Cosmetic Sciences*, **20**, 331-342.
- [17] Paoletti F, Aldinucci D, Mocalli A, Caparrini A. (1986) A sensitive spectrophotometric method for the determination of superoxide dismutase activity in tissue extracts. *Analytical Biochemistry*, **154**, 536-541.
- [18] Russo A, Cardile V, Lombardo L, Vanella L, Vanella A, Garbarino JA. (2005) Antioxidant activity and antiproliferative action of methanolic extract of *Geum quellyon* Sweet roots in human tumor cell lines. *Journal of Ethnopharmacology*, **100**, 323-332.
- [19] Dinis TC, Madeira VM, Almeida LM. (1994) Action of phenolic derivatives (acetoaminophen, salicylate and 5-aminosalicylate) as inhibitors of membrane lipid peroxidation and as peroxy radical scavengers. *Archives of Biochemistry & Biophysics*, **315**, 161-169.

***N1,N2,N3*-Trisopentenyl Guanidine and *N1,N2*-Diisopentenyl Guanidine, Two Cytotoxic Alkaloids from *Alchornea cordifolia* (Schumach.& Thonn.) Müll. Arg. (Euphorbiaceae) Root Barks**

Hélène Mavar-Manga^a, David Chapon^b, Sara Hoet^a, Sébastien Block^a,
Marie-Claire De Pauw-Gillet^c and Joëlle Quetin-Leclercq^{a*}

^aLaboratoire de pharmacognosie, Unité CHAM 72.30, Ecole de Pharmacie, UCL, Av. E. Mounier, 72, 1200 Bruxelles, Belgium

^bLaboratoire de chimie structurale, Unité CSTR, Bâtiment Lavoisier, Place Louis Pasteur, 1, Bte 4, 1348 Louvain-la-Neuve, Belgium

^cLaboratoire d'histologie et de cytologie, département des sciences précliniques, Bat. 6, Allée de la Chimie 3, 4000 Liège 1, Belgium

leclercq@cham.ucl.ac.be

Received: June 24th, 2006; Accepted: September 5th, 2006

Dedicated to the memory of Professor Ivano Morelli.

This paper describes the purification of two guanidine alkaloids: *N1, N2*-diisopentenyl guanidine (DIPG) **1** and *N1,N2,N3*-triisopentenyl guanidine (TIPG) **2** from *Alchornea cordifolia* root bark and reports their cytotoxic properties on cancer (HeLa, Mel-5, J774) and non cancer (WI 38) cells. TIPG showed the highest cytotoxicity with IC₅₀ values from 0.7 to 14.3 µg/mL (2.6 to 54.3 µM) on the four cell lines while DIPG was much less active: IC₅₀ 45.8 and 97.6 µg/mL (234.8 and 500.5 µM) on Mel-5 and HeLa and > 512.8 µM on J774 and WI 38. The results indicate that the cytotoxicity notably decreased with the loss of one isopentenyl substituent.


Keywords: *Alchornea cordifolia*, guanidine alkaloids, triisopentenyl guanidine (TIPG), diisopentenyl guanidine (DIPG), cytotoxicity.

A. cordifolia roots are widely used externally and internally to treat different illnesses [1-5] throughout tropical Africa. There have been several phytochemical studies on leaves and stem bark, but only a few old ones have dealt with the roots. The only compounds isolated were gentisic and anthranilic acids, tannins and a small quantity of alkaloids, one of which could be yohimbine, tannins were also found [3, 6].

In 1995, the antitumor activity of methanolic leaf extracts of *A. cordifolia* as well as fractions of different polarity were evaluated at a single high dose of 100 µg/mL against a panel of 60 human tumor cell lines and showed no significant activity [7].

This work reports the isolation and cytotoxic activity of *N1,N2*-diisopentenyl guanidine (DIPG) **1** and *N1,N2,N3*-triisopentenyl guanidine (TIPG) **2** from the root bark of *A. cordifolia*. *N1,N2,N3*-triisopentenyl guanidine was previously reported to be present in *A. cordifolia* leaves [8] but the reference cited to support the presence of this compound [9] does not mention it. While the crude extract (F) showed no cytotoxic activity on the cell lines tested, the partition of F between hexane and methanol-water gave a polar cytotoxic fraction (F_{MeOH}) while the non-polar fraction (F_{Hex}) showed a lower or no cytotoxicity (Table 1). The active F_{MeOH} fraction contained two major spots that were positive with Dragendorff and anisaldehyde/H₂SO₄ reagents and were purified by

HSCCC. Structures were determined by comparing the ESI-MS, ^{13}C and ^1H NMR spectra with literature data [10-12]. NMR chemical shifts of **1** and **2** are similar because of the electron delocalisation over the guanidine structure. Thus each isopentenyl substituent is nearly identical in NMR spectroscopy.



N1,N2-Diisopentenyl guanidine (DIPG) **1** *N1,N2,N3*-Triisopentenyl guanidine (TIPG) **2**

Table 1: *In vitro* cytotoxicity of *A. cordifolia* root bark extracts and isolated alkaloids (IC_{50} in $\mu\text{g/mL} \pm \text{SEM}$).

Fractions/ compounds	cancer cell lines			non cancer cell line
	HeLa ^a	Mel 5 ^a	J774 ^b	WI 38 ^a
F	>100	91.2 \pm 6.1	ND	>100
F _{Hex}	49.2 \pm 3.9	80.7 \pm 19.5	ND	>100
F _{MeOH}	22.2 \pm 2.2	20.2 \pm 3.5	ND	41.4 \pm 1.9
TIPG	11.1 \pm 2.1	8.5 \pm 1.3	0.7 \pm 0.5	14.3 \pm 1.6
DIPG	97.6 \pm 4.6	45.8 \pm 2.6	> 100	>100
Campto.	0.1 \pm 0.2	0.7 \pm 1.0	ND	2.4 \pm 1.9
Colchicine	ND	ND	0.08 \pm 0.02	ND

Campto. = camptothecin; ^a MTT assay; ^b Alamar BlueTM assay; ND = not determined.

Cytotoxicity was assessed on three human cell lines: two cancer (HeLa, Mel5) and one non cancer (WI 38) and one murine cancer cell line (J774) to detect an eventual selectivity. Both compounds showed a dose dependent cytotoxicity on the tested cell lines. In the first set of tests using MTT, we observed that for both isolated compounds Mel-5 cells proved to be the most sensitive while they were less toxic for WI-38. DIPG was less active and showed even no toxicity for WI-38 (Table 1). We also analysed the cytotoxicity on J774 using the Alamar BlueTM test and found that TIPG was very active (IC_{50} = 0.7 $\mu\text{g/mL}$, 2.6 μM) while DIPG could be considered as not toxic (IC_{50} >100 $\mu\text{g/mL}$, >512.8 μM). This stresses the importance of the three isopentenyl residues for the cytotoxic activity. Nevertheless, both compounds are much less effective than controls, which are highly cytotoxic compounds.

Up to now *N1,N2,N3*-TIPG was only reported in *Alchornea* species: *A. javanensis* [10, 13] and *A. glandulosa* [12]. This last team reported that a crude MeOH leaf extract of *A. glandulosa* and fractions containing **2** exhibited an antiproliferative activity on cancer cells and antimicrobial activities on *Bacillus subtilis* and *Candida tropicalis*. Nevertheless they did not test the activity of the purified compound.

N1,N2-DIPG **2** was isolated previously from *Pterogyne nitens* leaves, under the name of pterogynidine [11, 14]. It was reported to inhibit the growth of a mutant yeast strain lacking a DNA repair mechanism but was not evaluated for its cytotoxicity [11]. This plant also contains an isomer *N1,N1*-DIPG (pterogynine) which was not tested on yeast [14].

Another monosubstituted guanidine alkaloid (galegine) was isolated from different species of *Galega*, *Verbesina* and *Schoenus* [15, 16]. Although it was considered as toxic by its direct effect on pulmonary vascular permeability on some animals [17, 18], there is no *in vitro* toxicity study reported on human cell lines. It has also to be noted that aliphatic guanidine alkaloids have been shown to possess different biological properties: agmatine as hypotensive [19] or aplysillamides A and B as antimicrobials [20], an activity which may also be shared by the guanidines from *A. cordifolia*. In fact, *A. cordifolia* root bark extracts possess antimicrobial properties which may at least be due to alkaloids but also probably to tannins [21, 22].

In conclusion, we isolated, for the first time, two isopentenyl guanidine derivatives from *A. cordifolia* root bark. TIPG was shown to possess cytotoxic activity on different cell lines while DIPG was much less active. This emphasizes the importance of the three isopentenyl substituents but TIPG is at least 10 times less cytotoxic than camptothecin or colchicine. Furthermore, the presence of these compounds could partially explain some uses of *A. cordifolia* in African folk medicine.

Experimental

General: ^{13}C NMR (125.7 MHz) and ^1H NMR (500 MHz) experiments were carried out in CD_3OD with a Bruker Avance 500 spectrometer. Chemical shifts are reported in part per million (ppm). UV spectra were recorded in MeOH with an UVIKON 933 (Kontron Instrument). A Perkin Elmer spectrometer was used for IR spectra. Mass spectra were obtained by direct injection in ESI positive mode using a LCQ Advantage (Thermo Finnigan) mass spectrometer.

Plant material: The fresh leaves and root barks of *A. cordifolia* were collected in Kinshasa and identified at INERA (Institut National pour l'Etude et la Recherche Agronomique, University of Kinshasa). A voucher specimen is deposited at the Belgian National Botanic Garden (BR) bearing the number SP 848103.

Extraction and isolation: Dried root bark powder (250 g) was moistened with a 500 mL of 10% Na₂CO₃ aqueous solution overnight and then extracted in a Soxhlet successively with EtOAc (1.5 l) and CHCl₃ (1.5 l). These extracts were combined and evaporated under reduced pressure. The residue (F: 0.87%) was partitioned between hexane and MeOH-H₂O (8:2) and 500 mg of the polar fraction (F_{MeOH}) purified by HSCCC (High Speed Counter Current Chromatography) (Kromaton III, SEAB, France) with the solvent system hexane/EtOAc/*n*-BuOH/H₂O/AcOH 1: 1: 2: 5: 0.2 in the head to tail elution mode (the lower phase was the mobile phase) and a flow rate of 2 mL/min to give DIPG **1** (10.5 mg) and TIPG **2** (29.8 mg). T_R of **1**: 450–630 mL, **2**: 906–1200 mL. Due to the presence of AcOH in the solvent system, DIPG and TIPG were obtained as the acetates. Detection of eluate was performed by TLC: Merck silica gel 60 F₂₅₄ plate, toluene/EtOAc/MeOH/HCOOH 58:15:20:7; Dragendorff and anisaldehyde/H₂SO₄ reagents.

NI,N2-diisopentenyl guanidine (1)

IR (NaCl) ν_{max} : 3186, 1625, 1448 cm⁻¹.

UV (MeOH) λ_{max} nm: 203, 273.

¹H NMR (CD₃OD): 5.25 (tm, 1H, CH, *J* = 7.0 Hz), 3.78 (d, 2H, N-CH₂, *J* = 7.0 Hz), 1.90 (s, 3H, CH₃ acetate), 1.77 (s, 3H, CH₃), 1.72 (s, 3H, CH₃).

¹³C NMR-APT (CD₃OD): 180.1 (C=O, acetate), 157.3 (C=N), 139.0 (C=CH), 119.5 (CH=C), 40.5 (-CH₂-NH-), 25.6 (CH₃), 24.0 (CH₃, acetate), 18.0 (CH₃).

ESI-MS-MS: *m/z* (rel. int.) = 196 [M]⁺ (100), 127 (8).

Colourless oil, acetate salt, C₁₁H₂₁N₃.

Yield: 4.2%

NI,N2,N3-triisopentenyl guanidine (2)

IR and UV spectra were identical to *NI,N2-diisopentenyl guanidine*.

¹H NMR (CD₃OD): 5.24 (tm, 1H, CH, *J* = 6.8 Hz), 3.82 (d, 2H, N-CH₂, *J* = 6.8 Hz), 1.89 (s, 3H, CH₃ acetate), 1.77 (s, 3H, CH₃), 1.71 (s, 3H, CH₃).

¹³C NMR-APT (CD₃OD): 180.1 (C=O, acetate), 156.1 (C=N), 138.6 (C=CH), 120.0 (CH=C), 40.7 (CH₂-NH-), 25.8 (CH₃), 24.2 (CH₃, acetate), 18.0 (CH₃).

ESI-MS-MS: *m/z* (rel. int.) = 264 [M]⁺ (100), 310 (10), 195 (8).

References

- [1] Lewis WH, Elvin-Lewis PF. (1977) *Medical Botany: Plants Affecting Man's Health*. New York: John Wiley & sons. 720.

Colorless oil, acetate salt, C₁₆H₂₉N₃.

Yield: 11.92%

Cytotoxicity assay: Two methods were used to assess cytotoxicity depending on availability. They both rely on the same properties of living cells whose enzymes transform substrates introduced in the medium into coloured or fluorescent derivatives. The concentrations of these derivatives have been shown to be proportional to the number of living cells in most cases [23, 24].

MTT (Methylthiazoletetrazolium) assay: Cytotoxicity was determined on HeLa (human cervix carcinoma cells), Mel 5 (human node metastasis derived human melanoma cells clone 32 [25]) and WI38 (human lung fibroblasts) cell lines as described previously [26] with the following modifications: the Dulbecco's Modified Eagle Medium was supplemented with 5% heat-inactivated fetal bovine serum, L-glutamine 0.33%, non-essential amino acids 1%, penicillin/streptomycin 1% and sodium pyruvate 1%. Each extract and alkaloids were tested in a concentration range from 3.1 µg/mL to 50 µg/mL. Camptothecin (Sigma) was used as a positive control.

The results are expressed by IC₅₀ values (concentration of compound causing 50% inhibition of cell growth) calculated from graphs using at least five different concentrations of each alkaloid. All experiments were made at least in triplicate.

Alamar BlueTM assay: The assay was run as described previously by Hoet et al. on J774 cells (murine macrophages). Colchicine (Sigma) was used as a positive control [27].

Fluorescence development was calculated as percentage of the control culture, considered as 100%, and IC₅₀ values (concentration of extract that reduced fluorescence intensity by 50%) were calculated by linear interpolation according to Hills [28].

Acknowledgments - The authors wish to thank V. Derrider for SM measurements and G. Muccioli for the 1D-NMR spectra.

- [2] Bouquet A, Debray M. (1974) *Plantes Médicinales de la Côte d'Ivoire*. Travaux et Documents de l'O.R.S.T.O.M. Vol. 32. Paris: O.R.S.T.O.M. 232.
- [3] Kerharo J, Adam JG. (1974) *La Pharmacopée Sénégalaise Traditionnelle: Plantes Médicinales et Toxiques*. Paris: Vigots Frères. 1011.
- [4] Akedengué B. (1992) Medicinal plants used by the Fang traditional healers in Equatorial Guinea. *Journal of Ethnopharmacology*, **37**, 165-173.
- [5] Neuwinger HD. (2000) *African Traditional Medicine: A Dictionary of Plant Use and Applications*. Stuttgart: Medpharm, Scientific Publishers. 589.
- [6] Alves AC, Prista LN. (1962) Gentisic acid as a possible precursor in the biosynthesis of yohimbine. *Anais da Faculdade de Farmacia do Porto*, **22**, 27-33.
- [7] Muanza DN, Euler KL, Williams L, Newman DJ. (1995) Screening for antitumor and anti-HIV activities of nine medicinal plants from Zaire. *International Journal of Pharmacognosy*, **33**, 98-106.
- [8] Lamikanra A, Ogundaini AO, Ogungbamila FO. (1990) Antibacterial constituents of *Alchornea cordifolia* leaves. *Phytotherapy Research*, **4**, 198-200.
- [9] Ogungbamila FO, Samuelsson G. (1990) Smooth-Muscle Relaxing Flavonoids from *Alchornea cordifolia*. *Acta Pharmaceutica Nordica*, **2**, 421-422.
- [10] Hart NK, Johns SR, Lamberton JA, Willing RI. (1970) Alkaloids of *Alchornea javanensis*: the isolation of hexahydroimidazo [1,2- α] pyrimidines and guanidines. *Australian Journal of Chemistry*, **23**, 1679-1693.
- [11] Bolzani VDS, Gunatilaka AAL, Kingston DGI. (1995) Bioactive guanidine alkaloids from *Pterogyne nitens*. *Journal of Natural Products*, **58**, 1683-1688.
- [12] Conegero LD, Ide RM, Nazari AS, Sarragiotto MH, Dias BP, Nakamura CV, de Carvalho JE, Foglio MA. (2003) Chemical constituents of *Alchornea glandulosa* (Euphorbiaceae). *Quimica Nova*, **26**, 825-827.
- [13] Hart NK, Johns SR, Lamberton JA. (1969) Hexahydroimidazopyrimidines, a new class of alkaloids from *Alchornea javanensis*. *Journal of the Chemical Society [Section] D: Chemical Communications*, 1484-1485.
- [14] Corral RA, Orazi OO, De Petruccelli MF. (1969) Studies on plants: XIV. Guanidine alkaloid. *Experientia*, **25**, 1020-1021.
- [15] Susag L, Mathenge S, Benn M. (2003) The alkaloids of two species of *Afrogaleta*. *Biochemical Systematics and Ecology*, **31**, 645-647.
- [16] Oelrichs PB, Vallely PJ, MacLeod JK, Lewis IAS. (1981) Isolation of galegine from *Verbesina encelioides*. *Journal of Natural Products*, **44**, 754-755.
- [17] Keeler RF, Baker DC, Panter KE. (1992) Concentration of galegine in *Verbesina encelioides* and *Galega officinalis* and the toxic and pathologic effects induced by the plants. *Journal of Environmental Pathology, Toxicology and Oncology*, **11**, 75-81.
- [18] Dakshinamurti K, Bhuvaneshwar C. (1970) Effects of guanidine derivatives and oligomycin on swelling of rat liver mitochondria. *Biochemistry*, **9**, 5070-5076.
- [19] Delle Monache G, Volpe AR, Delle Monache F, Vitali A, Botta B, Espinal R, De Bonnevaux SC, De Luca C, Botta M, Corelli F, Carmignani M. (1999) Novel hypotensive agents from *Verbesina caracasana*. 7. Further hypotensive metabolites from *Verbesina caracasana*. *Bioorganic & Medicinal Chemistry Letters*, **9**, 3249-3254.
- [20] Honma K, Tsuda M, Mikami Y, Kobayashi J. (1995) Aplysillamides A and B, new antimicrobial guanidine alkaloids from the Okinawan marine sponge *Psammaphysilla purea*. *Tetrahedron*, **51**, 3745-3748.
- [21] Ebi GC. (2001) Antimicrobial activities of *Alchornea cordifolia*. *Fitoterapia*, **72**, 69-72.
- [22] Tona L, Kambu K, Mesia K, Cimanga K, Apers S, De Bruyne T, Pieters L, Totte J, Vlietinck AJ. (1999) Biological screening of traditional preparations from some medicinal plants used as antidiarrhoeal in Kinshasa, Congo. *Phytomedicine*, **6**, 59-66.
- [23] Mosmann T. (1983) Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays. *Journal of Immunological Methods*, **65**, 55-63.
- [24] O'Brien J, Wilson I, Orton T, Pogan F. (2000) Investigation of the Alamar Blue (resurazin) fluorescent dye for the assessment of mammalian cell cytotoxicity. *European Journal of Biochemistry*, **267**, 5421-5426.
- [25] Degiovanni G, Lahaye T, Herin M, Hainaut P, Boon T. (1988) Antigenic heterogeneity of a human melanoma tumor detected by autologous CTL clones. *European Journal of Immunology*, **18**, 671-676.
- [26] Block S, Stevigny C, De Pauw-Gillet M-C, De Hoffmann E, Llabres G, Adjakidje V, Quetin-Leclercq J. (2002) Ent-Trachyloban-3 β -ol, a new cytotoxic diterpene from *Croton zambesicus*. *Planta Medica*, **68**, 647-649.
- [27] Hoet S, Opperdoes F, Brun R, Adjakidje V, Quetin-Leclercq J. (2004) *In vitro* antitrypanosomal activity of ethnopharmacologically selected Beninese plants. *Journal of Ethnopharmacology*, **91**, 37-42.
- [28] Hills M, Hudson C, and Smith P. (1986) *Global Monitoring of the Resistance of the Malaria Parasites to Drugs: Statistical Treatment of the Micro-Test Data*. WHO working paper n° 2.8.5 for the Informal Consultation on the Epidemiology of Drug Resistance of Malaria Parasites. Geneva: World Health Organization.

Indole Monoterpenes with Antichemotactic Activity from *Psychotria myriantha*: Chemotaxonomic Significance

Cláudia A. Simões-Pires^a, Fabianne M. Farias^b, Andrew Marston^a, Emerson F. Queiroz^a,
Célia G. Chaves^b, Amélia T. Henriques^b and Kurt Hostettmann^{a*}

^aLaboratoire de Pharmacognosie et Phytochimie, École de Pharmacie Genève-Lausanne, Université de Genève, Université de Lausanne, CH-1211 Genève 4, Switzerland

^bPrograma de Pós-graduação em Ciências Farmacêuticas, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil

kurt.hostettmann@pharm.unige.ch

Received: June 6th, 2006; Accepted: August 8th, 2006

Dedicated to the memory of Professor Ivano Morelli.

The alkaloid extract of the aerial parts of *Psychotria myriantha* (Rubiaceae) displayed antichemotactic activity on polymorphonuclear leukocytes (PMN) assessed by the Boyden chamber assay. On analysis of the crude extract by LC/APCI/MS and LC/UV/DAD, two major constituents could be detected. In order to rapidly identify the active compounds, a microfractionation was conducted during LC/UV/DAD analysis. By this means, both the collected compounds could be assayed separately in the Boyden chamber and were shown to inhibit PMN chemotaxis. Their isolation was performed by semi-preparative HPLC and their structures elucidated by classical spectroscopic methods, including UV, NMR, MS and HRMS. Both compounds showed characteristics of monoterpene indole glucoside alkaloids; one of them was identified as strictosidinic acid and the other was a new natural product, myrianthosine. The antichemotactic activity of the compounds may be related to an antiacute inflammation activity.

Keywords: *Psychotria myriantha*, Rubiaceae, antichemotactic activity, glucoside indole monoterpene alkaloids, myrianthosine, strictosidinic acid.

Psychotria myriantha Mull. Arg. is a shrub (up to 2 meters high) occurring in southern Brazil [1]. Up to now, no phytochemical work has been reported on the species. As part of our continuing work on the Rubiaceae [2-5], the alkaloid extract from *P. myriantha* was tested for antichemotactic activity. From this extract, two active indole monoterpene alkaloid glucosides were isolated.

A polar alkaloid-rich extract of *P. myriantha* aerial parts was prepared as described in the experimental section. In order to identify the constituents, the alkaloid extract was analyzed by LC/UV/MS, with an atmospheric pressure chemical ionization (APCI) interface. Compound **1** gave an ion at m/z 517.1 $[M+H]^+$, while compound **2** gave a molecular ion at m/z 531.0 $[M+H]^+$ (Figure 1). The UV spectrum of

compound **1** showed absorptions of an indole chromophore (226 and 280 nm), while compound **2** showed three absorptions due to an extended chromophore at 240 (sh), 290 and 350 nm (Figure 1) [4]. LC microfractionation was performed on the extract and the inhibition of polymorphonuclear leukocyte (PMN) chemotaxis by an antichemotactic assay in a modified Boyden chamber was assessed in collected fractions. By this means, inhibition properties could be rapidly linked to two of the LC peaks at retention times of 11.0 and 12.5 min. The migration (μm) of PMN control cells and PMN treated with *P. myriantha* extract, **1** and **2** are shown in Table 1 as mean \pm S.D.

For full structure determination, **1** and **2** (Figure 2) were isolated by semi-preparative HPLC. Compound




Figure 1: (A) LC-UV-APCI/MS analysis of the alkaloid extract of the aerial parts of *Psychotria myriantha*. (B) UV and mass spectra of compounds **1** and **2**. (LC/UV/MS conditions: see Experimental section).

1 was identified as strictosidinic acid by comparing its data with those of published values [6]. Compound **2** was isolated as an amorphous solid. The high-resolution electrospray mass spectrum (HRESI/MS) showed a $[M+H]^+$ peak at an exact mass of m/z 531.2344 corresponding to the formula $C_{27}H_{35}N_2O_9$, suggesting the presence of an additional methyl group when compared to compound **1**. A careful analysis of 1H , ^{13}C and 2D NMR data indicated the presence of an open secologanin unit. As observed for compound **1**, the 1H NMR spectrum showed the presence of four signals at δ_H 8.17 (H-9, $J = 7.81$ Hz), 7.20 (H-10, $J = 7.33$ and 7.81 Hz), 7.50 (H-11, $J = 7.33$ and 7.81 Hz), 7.56 (H-12, $J = 7.82$ Hz), attributed to the aromatic protons of the indole moiety. However, the presence of two doublets at δ_H 7.93 (H-6, $J = 4.8$ Hz) and 8.23 (H-5, $J = 4.8$ Hz) indicated the presence of a double bond between the carbons H-6 and H-5. Analysis of the COSY and the HMBC spectra for the remainder of the molecule suggested the presence of a secologanin unit. All the same, careful analysis of these data revealed some differences when compared with **1**. The HMBC spectrum showed correlations between the signal at δ_H 1.00 and δ_C 119 (C-18), 135 (C-19) and 96 (C-21) suggesting the presence of a methyl group in position C-20. These data indicated that the ring of the secologanin was open. The attachment of the open secologanin unit to the β -carboline skeleton was ascertained from the HMBC spectrum. The

correlations between the methine proton at C-3 (δ_H 3.11), and the carbon at C-15 (δ_C 30.5) suggested the attachment of the unit at C-3. These data led to the structure of compound **2**. The HMBC spectrum of compound **2** showed correlations between the anomeric proton at δ_H 4.50 and the carbon at δ_C 95.4, suggesting the presence of a sugar moiety attached to C-21. The 2D NMR experiments (COSY, HMBC and HSQC) allowed the identification of the sugar as β -glucose. Acid and enzymatic hydrolysis confirmed the presence of β -D-glucose. These results pointed to a new natural product named myrianthosine.




Figure 2: Structures of compounds **1** and **2**.

Psychotria, one of the largest genera of angiosperms, is taxonomically complex and has been the object of investigation by several authors. Due to classification uncertainties, this genus has been placed close to *Cephaelis* and *Palicourea* [7,8]. Based on morphological parameters and geographical distribution, different authors suggested the division of *Psychotria* into three subgenera: *Psychotria* (pantropical), *Tetramera* (species from Africa and Madagascar) and *Heteropsychotria* (including the remainder of the neotropical species) [9-11].

Polyindoline alkaloids are the main metabolites found in the *Psychotria* genus, resulting from condensation of *N*(b)-methyltryptamine moieties [12-16]. These alkaloids have been found particularly in species classified into the subgenus *Psychotria*.

On the other hand, the subgenus *Heteropsychotria* has been characterized by the presence of monoterpene indole alkaloids. Isodolichantoxide, correantoxide, correantines A, B and C, 20-epi-correantine B, correantine and 10-hydroxycorreantoxide were isolated from leaves of *P. correae* growing in Panama [17].

From *Heteropsychotria* species found in the forests of Southern Brazil, glycosylated indole alkaloids have also been identified, such as lyaloside, strictosamide and nauclefine from *P. suterella* [5]; umbellatine from *P. umbellata* [18]; brachycerine from *P. brachyceras* [3,4]; and *N*- β -D-glucopyranosyl vincosamide from *P. leiocarpa* [2]. Some *Psychotria* species were formerly included in the genus *Cephaelis*, which is considered by some authors as a synonym of the subgenus *Heteropsychotria* [8,11,16]. However, *Cephaelis* alkaloids are usually tyrosine derivatives, whereas monoterpene indole alkaloids found in *Heteropsychotria* species are tryptophan derivatives [2]. The observation of different alkaloid profiles in different subgenera illustrates the importance of chemical analysis of *Psychotria* species, which may help to establish new groupings within this genus. Alkaloid type segregation among *Psychotria* species may also be related to their geographical distribution. It is remarkable that indole monoterpene alkaloids seem to be a constant feature in neotropical *Psychotria*, as observed for species occurring in Southern Brazil [2-5] as well as in Panama [17,19]. The presence of compounds **1** and **2** in *P. myriantha* corroborates this observation. From the biogenetic point of view, indole monoterpene alkaloids are

synthesized from tryptamine and the iridoid secologanin. Different secologanin derivatives have been reported to condense with tryptamine, such as the 10-oxo-1-*epi*-loganin precursor in brachycerine isolated from *P. brachyceras* [3]. The present work is the first report of an indole monoterpene alkaloid (**2**) incorporating an open ring of a secologanin derivative with an additional methyl group. Despite work done on the intermediates in the biosynthesis of secologanin [20], there is no precedent for the monoterpene moiety in the literature, and consequently a biosynthetic scheme for myrianthosine has not been attempted. The fact that strictosidinic acid (**1**) and myrianthosine (**2**) are genuine alkaloids of *P. myriantha* was confirmed by LC analysis of rapidly prepared ethanolic extracts of fresh leaves without acid/base extraction.

Strictosidinic acid (**1**) has previously been isolated from *Hunteria zeylanica* (Retz.) Gardner ex Thwaites (Apocynaceae) and has shown analgesic and antipyretic activities in mice after oral administration [21]. In the present work, it has been demonstrated that this alkaloid, together with myrianthosine (**2**), are responsible for the activity of the alkaloid extract of *P. myriantha* and were able to inhibit *in vitro* polymorphonuclear leukocytes (PMN) chemotaxis (Table 1). This activity has been demonstrated for some second-line anti-inflammatory drugs, such as D-penicillamine [22].

Table 1: Antichemotactic activity of *P. myriantha* alkaloid extract and isolated compounds.

Sample	PMN migration (μ m) ^a
Control cells	129.2 \pm 1.68
<i>P. myriantha</i> alkaloid extract	12.4 \pm 0.84*
myrianthosine	14.6 \pm 0.96*
strictosidinic acid	12.6 \pm 0.96*
genistein ^b	9.9 \pm 1.00*

^aValues given as mean \pm S.D of 10 measurements

^bPositive control

*Statistically significant - $p < 0.005$ (Student's *t*-test)

The inhibition of PMN chemotaxis reported here may be involved in the mechanism of the previously published antipyretic and analgesic activities of strictosidinic acid (**1**) [21] and the analgesic activity of *P. myriantha* extract [23].

Experimental

General: Optical rotation $[\alpha]_D$ was determined using a Perkin-Elmer 241 polarimeter (MeOH, *c* in g/100 mL). ¹H- and ¹³C NMR were recorded on a Varian Unity Inova 500 spectrometer (500 MHz and 125 MHz, respectively) in DMSO-*d*₆; chemical shifts

in ppm as δ rel. to Me₄Si (int. std.). LC/MS was performed directly after UV-DAD measurements. A Finnigan LCQ ion trap (Finnigan MAT, San Jose, CA, USA) with APCI interface was used with the following conditions: capillary temp. 150°C; vaporizer temp. 370°C; positive mode; sheath gas flow: 60 arb, corona needle current 5 μ A; spectra (150-900 m μ). HRESIMS was performed using a Bruker FTMS 4.7T. TLC: silica gel 60 F₂₅₄ Al sheets (Merck), detection at 254 nm and with vanillin-sulfuric acid reagent. LC/UV-DAD analysis of the alkaloid extract was performed on a Hewlett-Packard (Waldbronn, Germany) Series 1100 photodiode array detector (DAD) liquid chromatograph system. The separation was achieved on a Nucleosil 100-5 C₁₈ AB column (125 x 4.6 mm i.d., 5 μ m; Macherey-Nagel) with MeOH/H₂O (containing Et₃N 2 mM) in the gradient mode (10% of MeOH to 100% in 40 min). The flow rate was 1 mL/min; the UV traces were measured at 210 and 254 nm and UV spectra (DAD) were recorded between 200 and 500 nm. HPLC microfractionation: Fractions were collected, after the LC/UV analysis, every 1 min (1 mL) in Eppendorf tubes by a Gilson collector (FC204). After collection, all fractions were evaporated to dryness on a Speedvac system (RCT 90, Jouan). The content of each fraction was suspended in 1 mL of PMN suspension and then used for the antichemotactic assay. Semi-preparative HPLC was carried out with a Shimadzu LC-8A pump equipped with a Knauer UV detector using a Symmetry-Prep column (7 μ m, 19x150 mm, Waters).

Plant material: *P. myriantha* was collected in Reserva Estadual do Turvo, Derrubadas Rio Grande do Sul, Brazil and identified by M. Sobral. A voucher specimen (M. Sobral *et al.*, 8913) was deposited in the ICN Herbarium (Universidade Federal do Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil).

Extraction and isolation: Dried leaves (823 g) were extracted with EtOH (3.5 l) at room temperature. The extract was concentrated under vacuum at 40 °C and an alkaloid extract was obtained by classical acid/base extraction. In addition to the alkaloid rich CH₂Cl₂ extract, it was noticed that the aqueous fraction was also positive for alkaloids with Dragendorff reagent. In order to extract these alkaloids, a second partition was conducted between the residual aqueous fraction and *n*-BuOH. The butanolic extract (1 g) was purified by semi-preparative HPLC using Symmetry-Prep column

(7 μ m, 19 x 150 mm, Waters), MeOH-H₂O (30:70) with Et₃N 2 mM, flow rate 10 mL/min, UV 254 nm) providing alkaloids **1** (33 mg) and **2** (12 mg).

Acid hydrolysis: Compounds **1** and **2** were submitted to hydrolysis with 20 mL of HCl 0.05 N at 65°C during 24h. Sugars were extracted by partition with *n*-BuOH and were compared by TLC (solvent: AcOEt/Formic acid/H₂O 100:20:30; detected with *p*-anisaldehyde-H₂SO₄ reagent) with reference compounds: glucose (R_f 0.40), arabinose (R_f 0.45), fructose (R_f 0.41), rhamnose (R_f 0.64), and galactose (R_f 0.33), all 1 mg/mL in H₂O. Compounds **1** and **2** gave glucose (R_f 0.40).

Enzymatic hydrolysis: Compounds **1** and **2** were treated with β -D-glucosidase in 1 mL NaOAc buffer (pH 5.0) for three days at 40°C. The aglycones were extracted by partition with *n*-BuOH, and submitted to LC/UV-DAD analysis hydrolysis in order to confirm hydrolysis.

Antichemotactic assay: Chemotaxis was measured in a Boyden chamber by the method previously described [24]. Prior to the chemotactic assay, rat leukocytes were treated with 100 μ g/mL of each sample (alkaloids and extract), at 37°C for 1 h. Plasma collected from rats was incubated at 37°C for 30 min with 65 μ g/mL of lipopolysaccharide (LPS) from *Escherichia coli*, and then diluted in Hanks buffer 1:5 (v/v). Chemotactic migration of leukocytes through an 8.0- μ m cellulose nitrate filter, towards the chemotactic stimulant (LPS treated plasma) was measured after incubation for 1 h at 37°C using the micrometer on the fine-focus knob of a Nikon Alphaphot-2 YS2 microscope. The distance from the upper surface of the filter to the lower surface of focus still containing two cells allowed the evaluation of leukocyte migration in five microscopic fields per filter. The assay was carried out in duplicate and measurements were statistically analyzed by Student's *t*-test, using genistein as positive control.

Strictosidinic acid (1)

Amorphous powder.

$[\alpha]_D^{25}$: +143.14° (*c* 0.1, MeOH).

R_f: 0.30: TLC system: ethyl acetate / acetic acid / formic acid / water (100:11:11:10).

¹H NMR (500 MHz, DMSO-*d*₆): 4.10 (1H, d, *J* = 10.4 Hz, H-3), 2.92-3.47 (2H, m, H-5), 2.73-2.92 (2H, m, H-6), 7.41 (1H, d, *J* = 7.81 Hz, H-9), 6.97 (1H, dd, *J* = 7.33 and 7.81 Hz, H-10), 7.05 (1H, dd,

$J = 7.33$ and 7.81 Hz, H-11), 7.29 (1H, d, $J = 7.81$ Hz, H-12), 1.95 - 2.20 (2H, m, H-14), 2.90 (1H, m, H-15), 7.38 (1H, s, H-17), 5.13 (1H, d, $J = 11.23$ Hz, H-18a), 5.30 (1H, d, $J = 17.09$ Hz, H-18b), 5.76 (1H, m, H-19), 2.60 (1H, m, H-20), 5.62 (1H, d, $J = 9.70$ Hz, H-21), 4.63 (1H, d, $J = 7.81$ Hz, H-1'), 3.10 (1H, m, H-2'), 2.90 (1H, m, H-3'), 3.20 (1H, m, H-4'), 4.12 (1H, m, H-5'), 3.50 (1H, m, H-6' α), 3.70 (1H, m, H-6' β).

^{13}C NMR (125 MHz, DMSO- d_6): 132.3 (C), 49.6 (CH), 40.0 (CH $_2$), 19.2 (CH $_2$), 106.0 (C), 126.1 (C), 117.8 (CH), 118.7 (CH), 121.2 (CH), 111.5 (CH), 135.8 (C), 33.7 (CH $_2$), 31.8 (CH), 113.4 (C), 150.0 (CH), 117.8 (CH $_2$), 135.6 (CH), 44.3 (CH), 95.1 (CH), 170.0 (C), 98.9 (CH), 69.8 (CH), 73.1 (CH), 77.2 (CH), 76.5 (CH), 61.0 (CH $_2$).

APCIMS m/z 517.4 $[\text{M}+\text{H}]^+$. HRESIMS m/z 517.2172 $[\text{M}+\text{H}]^+$, (calculated for $\text{C}_{26}\text{H}_{33}\text{N}_2\text{O}_9$, 517.2186).

Myrianthosine (2)

Amorphous powder.

$[\alpha]^{25}_{\text{D}}$: $+80.8^\circ$ (c 0.1, MeOH).

Rf: 0.27; TLC system: ethyl acetate / acetic acid / formic acid / water (100:11:11:10).

^1H NMR (500 MHz, DMSO- d_6): 3.11 (1H, m, H-3), 8.23 (1H, $J = 4.80$ Hz, H-5), 7.93 (1H, $J = 4.80$ Hz,

H-6), 8.17 (1H, d, $J = 7.81$ Hz, H-9), 7.20 (1H, dd, $J = 7.33$ and 7.81 Hz, H-10), 7.50 (1H, dd, $J = 7.33$ and 7.81 Hz, H-11), 7.56 (1H, d, $J = 7.81$ Hz, H-12), 2.70 (2H, m, H-14), 1.25 (2H, m, H-15), 7.40 (1H, s, H-17), 4.60 (1H, d, $J = 17.1$ Hz, H-18a), 4.80 (1H, d, $J = 10.2$ Hz, H-18b), 5.70 (1H, ddd, $J = 17.10$, 10.20 and 2.00 Hz, H-19), 2.60 (1H, m, H-20), 1.00 (3H, d, $J = 7.8$ Hz, CH $_3$ -20), 5.47 (1H, d, $J = 5.3$ Hz, H-21), 4.50 (1H, d, $J = 7.8$ Hz, H-1'), 3.00 (1H, m, H-2'), 3.10 (1H, m, H-3'), 3.20 (1H, m, H-4'), 4.10 (1H, m, H-5'), 3.41 (1H, m, H-6' α), 3.70 (1H, m, H-6' β).

^{13}C NMR (125 MHz, DMSO- d_6): 134.8 (C), 48.5 (CH), 137.0 (CH), 111.8 (CH), 121.0 (C), 121.5 (C), 126.6 (CH), 118.9 (CH), 127.5 (CH), 112.5 (CH), 140.2 (C), 45.6 (CH $_2$), 30.0 (CH $_2$), 112.0 (C), 151.0 (CH), 118.9 (CH $_2$), 134.5 (CH), 45.5 (CH), 95.4 (CH), 170.0 (C), 98.6 (CH), 73.0 (CH), 69.9 (CH), 77.3 (CH), 76.8 (CH), 61.0 (CH $_2$), 10.4 (CH $_3$).

APCIMS m/z positive: 531.2 $[\text{M}+\text{H}]^+$.

HRESIMS m/z $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{27}\text{H}_{35}\text{N}_2\text{O}_9$: 531.2337 ; found: 531.2344 .

Acknowledgments - The Swiss National Science Foundation (grant n° 200020-100083 to K. Hostettmann) and CNPq (Brazil) are gratefully acknowledged for supporting this work.

References

- [1] Dillenburg CR, Porto ML. (1985) Rubiaceae - Tribo Psychotriaceae. *Boletim do Instituto de Biociências - UFRGS*, **39**, 1-76.
- [2] Henriques AT, Lopes SO, Paranhos JT, Gregianini TS, Von Poser GL, Fett-Neto AG, Schripsema J. (2004) N,β -D-glucopyranosyl vincosamide, a light regulated indole alkaloid from the shoots of *Psychotria leiocarpa*. *Phytochemistry*, **65**, 449-454.
- [3] Kerber VA, Gregianini TS, Paranhos JT, Schwambach J, Farias F, Fett JP, Fett-Neto A, Zuanazzi JAS, Quirion JC, Elisabetsky E, Henriques AT. (2003) Brachycerine, a novel monoterpene indole alkaloid from *Psychotria brachyceras*. *Journal of Natural Products*, **66**, 1038.
- [4] Kerber VA, Gregianini TS, Paranhos JT, Schwambach J, Farias F, Fett JP, Fett-Neto AG, Zuanazzi JAS, Quirion JC, Elisabetsky E, Henriques AT. (2001) Brachycerine, a novel monoterpene indole alkaloid from *Psychotria brachyceras*. *Journal of Natural Products*, **64**, 677-679.
- [5] Van De Santos L, Fett-Neto AG, Kerber VA, Elisabetsky E, Quirion JC, Henriques AT. (2001) Indole monoterpene alkaloids from leaves of *Psychotria suterella* Mull. Arg. (Rubiaceae). *Biochemical Systematics and Ecology*, **29**, 1185-1187.
- [6] Arbain D, Putra DP, Sargent MV. (1993) The alkaloids of *Ophiorrhiza filistipula*. *Australian Journal of Chemistry*, **46**, 977-985.
- [7] Nepokroeff M, Bremer B, Sytsma KJ. (1999) Reorganization of the genus *Psychotria* and tribe Psychotrieae (Rubiaceae) inferred from ITS and rbcL sequence data. *Systematic Botany*, **24**, 5-27.
- [8] Taylor CM. (1996) Overview of the Psychotrieae (Rubiaceae) in the Neotropics. *Opera Botanica Belgica*, **7**, 267-270.
- [9] Pettit E. (1964) Les espèces africaines du genre *Psychotria* L. (Rubiaceae). *Bulletin du Jardin Botanique de Bruxelles*, **34**, 1-229.
- [10] Pettit E. (1966) Les espèces africaines du genre *Psychotria* L. (Rubiaceae). II. *Bulletin du Jardin Botanique de Bruxelles*, **36**, 65-189.
- [11] Steyermark J. (1972) Botany of the Guyana highlands. In *Memoirs of the New York Botanical Garden*. Vol. **23**, Maguire B, Cowan S, Wurdack JJ (Eds). New York Botanical Garden Press, New York, 227-832.
- [12] Jannic V, Gueritte F, Laprevote O, Serani L, Martin MT, Sevenet T, Potier P. (1999) Pyrrolidinoindoline alkaloids from *Psychotria oleoides* and *Psychotria lyciiflora*. *Journal of Natural Products*, **62**, 838-843.
- [13] Lajis NH, Mahmud Z, Toia RF. (1993) The alkaloids of *Psychotria rostrata*. *Planta Medica*, **59**, 383-384.

- [14] Libot F, Miet C, Kunesch N, Poisson JE, Pusset J, Sevenet T. (1987) Plants of New-Caledonia. 110. Rubiaceae of Oceania - Alkaloids from *Psychotria oleoides* of New-Caledonia and from *Calycodendron-Milnei* of Vanuatu (New-Hebrides). *Journal of Natural Products*, **50**, 468-473.
- [15] Roth A, Kuballa B, Cabalion P, Anton R. (1985) Preliminary study of the alkaloids of *Psychotria forsteriana*. *Planta Medica*, **5**, 289-289.
- [16] Solis PN, Ravelo AG, Palenzuela JA, Gupta MP, Gonzalez A, Phillipson JD. (1997) Quinoline alkaloids from *Psychotria glomerulata*. *Phytochemistry*, **44**, 963-969.
- [17] Achenbach H, Lottes M, Waibel R, Karikas GA, Correa MD, Gupta MP. (1995) Constituents of tropical medicinal plants. 66. Alkaloids and other compounds from *Psychotria correae*. *Phytochemistry*, **38**, 1537-1545.
- [18] Lopes S, Von Poser GL, Kerber VA, Farias FM, Konrath EL, Moreno P, Sobral ME, Zuanazzi JAS, Henriques AT. (2004) Taxonomic significance of alkaloids and iridoid glucosides in the tribe Psychotrieae (Rubiaceae). *Biochemical Systematics and Ecology*, **32**, 1187-1195.
- [19] Solis PN, Wright CW, Gupta MP, Phillipson JD. (1993) Alkaloids from *Cephaelis dichroa*. *Phytochemistry*, **33**, 1117-1119.
- [20] Uesato S, Matsuda S, Inouye H. (1984) Mechanism for iridane skeleton formation from acyclic monoterpenes in the biosynthesis of secologanin and vindoline in *Catharanthus roseus* and *Lonicera morrowii*. *Chemical & Pharmaceutical Bulletin*, **32**, 1671-1674.
- [21] Reanmongkol W, Subhadhirasakul S, Kongsang J, Tanchong M, Kitti J. (2000) Analgesic and antipyretic activities of *n*-butanol alkaloids extracted from the stem bark of *Hunteria zeylanica* and its major constituent, strictosidinic acid, in mice. *Pharmaceutical Biology*, **38**, 68-73.
- [22] Turner RA, Johnson JA, Semble EL. (1983) Anti-rheumatic drug effects on neutrophil response to chemotactic factors - A comparison of analytical techniques. *Proceedings of the Society for Experimental Biology and Medicine*, **173**, 200-204.
- [23] Both FL, Misturini J, Henriques AT, Elisabetsky E. (2002) Avaliação da atividade analgésica de extratos alcaloídicos de espécies de *Psychotria*. *Revista Brasileira de Plantas Medicinais*, **5**, 41-45.
- [24] Zigmond SH, Hirsch JG. (1973) Leukocyte locomotion and chemotaxis - New methods for evaluation and demonstration of a cell-derived chemotactic factor. *Journal of Experimental Medicine*, **137**, 387-410.

HPLC Based Activity Profiling for Inhibitors of Human Neutrophil Elastase in *Isatis tinctoria* Leaf Extracts

Matthias Hamburger^{a,b}, Georg-Ulrich Rüster^b and Matthias F. Melzig^c

^aInstitute of Pharmaceutical Biology, University of Basel, Klingelbergstrasse 50, CH-4056 Basel, Switzerland

^bInstitute of Pharmacy, University of Jena, Semmelweisstrasse 10, D-07743 Jena, Germany

^cInstitute of Pharmacy, Free University Berlin, Königin-Luise-Strasse 2 u. 4, D-14195 Berlin, Germany

matthias.hamburger@unibas.ch

Received: June 7th, 2006; Accepted: August 5th, 2006

Dedicated to the memory of Professor Ivano Morelli.

In continuation of our search for anti-inflammatory constituents in lipophilic extracts of *Isatis tinctoria*, an HPLC-based profiling for inhibitors of human neutrophil elastase was performed. Using a linear gradient profile, fractions with significant activity (>50%) appeared in the second half of the chromatogram and corresponded to moderately to highly lipophilic constituents. The active fractions of intermediate polarity were further analyzed by HPLC, TLC and GC. They contained mainly α -linolenic, linoleic, *cis*-11-octadecenoic acid, oleic and palmitic acids.

Keywords: *Isatis tinctoria*, anti-inflammatory, human neutrophil elastase, α -linolenic acid, linoleic acid, activity profiling.

The woad plant (*Isatis tinctoria* L, Brassicaceae) is an old indigo dye and medicinal plant of temperate climate zones. The plant was used for the treatment of wounds, ulcers, snakebites, and various inflammatory afflictions, but fell into oblivion with the disappearance of woad cultivation [1]. In contrast, the related *I. indigotica* is a widely used plant in traditional Chinese Medicine up to now [2, 3].

Our interest in woad was spurred by its purported anti-inflammatory properties for which we obtained first, *in vitro* evidence in a broad-based pharmacological screening involving more than 20 clinically relevant targets [4]. Subsequently, we identified the alkaloid tryptanthrin as a potent inhibitor of COX-2 and 5-LOX catalyzed eicosanoid synthesis [5, 6], and an indolin-2-one derivative as inhibitor of histamine release from mast cells [7]. We found that the major inhibitor of 5-LOX in woad extracts was γ -linoleic acid [8]. Further anti-inflammatory activities of *Isatis* constituents have been reported by other groups, for example, inhibition of the expression of inducible

NO synthase [9] by tryptanthrin, and inhibition of IL-6 release and TNF α by indirubin [10]. Lipophilic *Isatis* extracts displayed activity in various *in vivo* models of inflammation, cutaneous allergy and arthritis [11, 12]. In a clinical pilot study, the same extracts reduced inflammation and water loss in a cutaneous irritation model [13].

A significant inhibition of leucocytic elastase from neutrophils had been observed in our initial *in vitro* screening [4]. This enzyme is one of two main proteinases of neutrophils released in connection with inflammation. It cleaves fibrous elastin, collagens, cartilage proteoglycans, and other matrix proteins. Elevated plasma levels of neutrophil elastase are a characteristic feature of an active inflammation [14, 15]. We, therefore, decided to identify the inhibitors of human neutrophil elastase in woad extracts with the aid of HPLC-based activity profiling using an established bioassay [16]. We had previously used the approach of HPLC profiling for

the identification of a variety of bioactive compounds from *I. tinctoria* [5, 7, 8] and other plants [17].

For HPLC profiling, the SFE extract was dissolved in dichloromethane (10 mg/mL) and separated on an analytical C-18 column. Aliquots of 200 µg extract per injection were repeatedly fractionated. A linear gradient from 10% to 100% acetonitrile over 25 min was followed by isocratic elution for an additional 20 min. In a first profiling step, 15 fractions of 3 min each (Fr. 1-15) were collected. The solvent was removed in an evaporator centrifuge and the dry films were redissolved in 180 µL DMSO for the elastase assay, which was carried out according to a published protocol [16], whereby *p*-nitro-aniline release was quantified by measurements of absorbancy at 405 nm. Representative HPLC chromatograms recorded at 254 and 220 nm and the activity profile of fractions 1 to 15 in the elastase assay are shown in Figure 1.




Figure 1: HPLC chromatogram of *Isatis tinctoria* extract recorded at 220 nm (bottom). Vertical lines indicate time windows for fractions 1 to 15. Inhibitory activity of fractions is shown above (mean of 3 independent experiments in two parallels).

Inhibitory activities > 50% were found in fractions 8 (77%), 9 (66%), 14 (78%) and 15 (85%). In a second step, fractions of 0.5 min were collected in the time window 21 to 30 mins, which corresponded to fractions 8 to 10. The HPLC profile of this time window and fractionation steps (8-1 to 10-6) are shown in Figure 2.

Fractions were analyzed by HPTLC (RP-18, HOAc/H₂O (95:5)). Upon staining with Godin's reagent, fractions 8-5, 9-2, 9-5 and 9-6 each showed one single violet spot in the *R_f* range 0.4 to 0.5. The




Figure 2: HPLC chromatogram recorded at 220 nm of the time window from 21 to 30 min, corresponding to fractions 8-1 to 10-6. Vertical lines indicate time windows for fractionation.

R_f values were comparable with α -linolenic, linoleic and oleic acids. For further analysis, fractions 8-1 to 10-6 were submitted to GC after derivatization with TMSH. The results of the fatty acid analysis of the major fractions 8-5, 9-2, 9-5 and 9-6 are shown in Figure 3. Identification was carried out by an overlay of GC chromatograms with a standard reference mix (C-18 FAME Isomer Mix, Supelco). Fraction 8-5 consisted of α -linolenic acid, fraction 9-2 of linoleic acid, whereas fraction 9-5 contained oleic acid and *cis*-11-octadecenoic acid in equal proportions. Fraction 9-6 consisted of palmitic acid.

Inhibition of human neutrophil elastase by saturated and unsaturated fatty acids has been recently reported. The IC₅₀ of α -linolenic, linoleic, oleic and palmitic acids were in the range of 5 to 15 µM [18]. Hence, inhibitory activity in fractions 8 and 9 can be attributed to these fatty acids. HPTLC and GC analysis of fractions 14 and 15, however, showed that they contained only traces of fatty acids. Elastase inhibition in these fractions seems thus due to highly lipophilic compounds, which remain to be identified.

Experimental

Organic solvents for separations were HPLC grade (Roth, Karlsruhe, Germany). HPLC grade water was obtained from a Milli-Q RG water purification system (Millipore, Schwalbach, Germany).

Leaf material of *I. tinctoria* was harvested in 2001 from first year plants grown on experimental plots of the Agricultural Research Station of Thuringia (TLL), Dornburg, Germany. Fresh leaves were dried on a band drier operating at 60°C.

Extracts were prepared by supercritical fluid extraction (SFE) on a pilot plant extractor, with the following conditions: CO₂ + 2000 ppm EtOH as modifier, 700 bar, 50°C. Tryptanthrin concentration in the SFE extract was 0.23%, as determined by ESI-MS. The extract was stored at -32°C. Fractionations were carried out with a HP 1100 HPLC system consisting of autosampler, high-pressure mixing pump, column oven, diode array detector, and HP workstation (Agilent, Waldbronn, Germany) connected to a Gilson FC 2004 fraction collector (Gilson, Middleton, USA). GC analysis of fatty acids was carried out on a Shimadzu 17A system (Shimadzu, Duisburg, Germany) with an AOC 5000 autosampler. TLC analysis was performed on RP-18 F254 HPTLC glass plates (Merck, Darmstadt, Germany).

HPLC separations were performed on a LiChrospher 100 RP-18 cartridge (5 µm, 125 x 4 mm i.d.; Merck, Darmstadt, Germany) using acetonitrile (A) and water (B) as eluents, with the following gradient profile: 10% A (0-2 min), 10% → 100% A (2-25 min), 100% A (25-45 min). The flow rate was set at 1.0 mL/min, and the column oven was at 25°C. HPLC traces were recorded at 220 and 254 nm, and UV-vis spectra from 190-700 nm.

For activity profiling, *Isatis* extract was dissolved in dichloromethane at a concentration of 10 mg/mL. Aliquots of 30 µL of the solution, corresponding to 300 µg extract, were injected for HPLC separation. The column effluent was fractionated into 5 mL vials. Fifteen fractions of 3 min each were collected. The solvent was removed at 50°C in a centrifugal evaporator (Evaporator centrifuge RC 10.22, Jouan GmbH, Unterhaching, Germany). The separation was repeated 4 times. The dried films were redissolved in MeOH (500 µL), transferred into 1.5 mL Eppendorff tubes, and dried again in a centrifugal evaporator prior to shipment for bioassay. In the second round of profiling, the time window from 21 to 30 min was fractionated at higher resolution. A total of 18 fractions of 0.5 min each (8-1 to 10-6) were collected. The fractions were processed as described above.

GC analysis was carried out after derivatization with trimethylsulfoniumhydroxide (TMSH) in MeOH (Macherey-Nagel, Düren, Germany). A 2 µL volume was injected onto a capillary column, DB 225 ms (length 60 m, 0.25 mm i.d., film thickness 0.25 µm) under the following conditions: split (1:20), injector

temperature 260°C, mobile phase: H₂, (42 cm /s), detection: FID, detector temperature 270°C, temperature program: start 70 °C, hold for 2 min, 70 °C → 180°C (heating rate 10°C / min), 180°C → 220°C (heating rate 2 °C / min), hold time 5 min at 220°C, 220°C → 230°C (heating rate 2°C / min), hold 15 min. The identification was carried out in comparison to a standard reference mix (C-18 FAME Isomer Mix, Supelco, Taufkirchen, Germany).

Plates were developed with AcOH/H₂O (95:5) as mobile phase. Compounds were stained with Godin's reagent (equal volumes of solution A and B, freshly mixed; solution A: 1% vanillin in EtOH + 3% HClO₄, solution B: 10% H₂SO₄ in EtOH) followed by heating at 105°C for 3 min (Thermoplate S, Desaga, Heidelberg, Germany). The compound in fraction 8-5 showed the same chromatographic mobility and staining as the reference α -linolenic acid (violet zone, R_f = 0.5), the compound in fraction 9-2 showed the same behaviour as linoleic acid (violet spot, R_f = 0.45), and the spot in fractions 9-5 and 9-6 showed the same behaviour as oleic acid (pale violet staining, R_f = 0.4).

Inhibition of neutrophil elastase was determined with human leucocyte elastase, according to a published procedure [16]. Briefly, 125 µL substrate solution (10 mM MeO-Suc-Ala-Ala-Pro-Val-pNA in Tris-HCl-buffer, 60 mM, pH 7.5) were mixed with 10 µL test solution (test substances solubilized in DMSO) + 445 µL Tris-HCL-buffer, pH 7.5 and vortexed. After the addition of 20 µL enzyme solution (approximately 1.05 mU) the samples were incubated for 1 h at 37°C. The reaction was stopped by addition of 500 µL soybean trypsin inhibitor solution (2 mg/mL Tris-HCL-buffer, pH 7.5) and placed in an ice bath. After vortexing, the absorbance was read at 405 nm. The assays were performed three times with duplicate samples and DMSO controls. Inhibition rates were calculated in percent to DMSO controls, and IC₅₀ values calculated from the dose-inhibition curves by linear regression. As positive control for the elastase assay, the inhibitor GW311616A [19], with an IC₅₀ of 90 nM, was used.

For HPTLC analysis, MeOH solutions (20 µL) of fractions 8-1 to 10-6 were sprayed onto a RP-18 plate with the aid of an AS 30 TLC applicator (Desaga, Heidelberg, Germany) along with reference solutions of α -linolenic, linoleic and oleic acids (1 mg/mL).

Acknowledgments - We thank Dr A. Vetter and Mrs A. Biertümpfel, TLL Jena and Dornburg, for the woad samples, Mrs M. Janka, FU Berlin, for technical assistance, Dr B. Weinreich, Adalbert-Raps-Forschungszentrum, TU München-Weihenstephan,

for preparation of the SFE extract, Prof. G. Jahreis and P. Moeckel, Institute of Food Sciences, University of Jena, for fatty acid analysis, and T. Mohn, University of Basel, for assistance in preparation of figures.

References

- [1] Hurry JB. (1930) *The Woad Plant and its Dye*. Oxford University Press, London. 249-256.
- [2] Chang HM, But PPH. (1986) *Pharmacology and Applications of Chinese Material Medica*, vol. **I**. World Scientific, Singapore. 94.
- [3] Tang W, Eisenbrand G. (1992) *Chinese Drugs of Plant Origin*. Springer, Berlin. 805-808.
- [4] Hamburger M. (2002) *Isatis tinctoria* – From the rediscovery of an ancient medicinal plant towards a novel anti-inflammatory phytopharmaceutical. *Phytochemistry Reviews*, **1**, 333-344.
- [5] Danz H, Stoyanova S, Wippich P, Brattström A, Hamburger M. (2001) Identification and isolation of the cyclooxygenase-2 inhibitory principle in *Isatis tinctoria*. *Planta Medica*, **67**, 411-416.
- [6] Danz H, Stoyanova S, Thomet OAR, Simon HU, Dannhardt G, Ulbrich H, Hamburger M. (2002) Inhibitory activity of tryptanthrin on prostaglandin and leukotriene synthesis. *Planta Medica*, **68**, 875-880.
- [7] Rüster GU, Hofmann B, Hamburger M. (2004) Inhibitory activity of indolin-2-one derivatives on compound 48/80-induced histamine release from mast cells. *Pharmazie*, **59**, 236-237.
- [8] Oberthür C, Jäggi R, Hamburger M. (2005) HPLC based activity profiling for 5-lipoxygenase inhibitory activity in *Isatis tinctoria* leaf extracts. *Fitoterapia*, **76**, 324-332.
- [9] Ishihara T, Kohno K, Ushio S, Kurimoto M. (2000) Tryptanthrin inhibits nitric oxide and prostaglandin E2 synthesis by murine macrophages. *European Journal of Pharmacology*, **407**, 197-204.
- [10] Kunikata T, Tatefuji T, Aga H, Iwaki K, Ikeda M, Kurimoto M. (2000) Indirubin inhibits inflammatory reactions in delayed-type hypersensitivity. *European Journal of Pharmacology*, **410**, 93-100.
- [11] Recio MC, Cerdá-Nicolás M, Potterat O, Hamburger M, Rios JL. (2006) Anti-inflammatory and anti-allergic activity in vivo of lipophilic *Isatis tinctoria* extracts and tryptanthrin. *Planta Medica*, **72**, 539-546.
- [12] Recio MC, Cerdá-Nicolás M, Hamburger M, Rios JL. (2006) Anti-arthritis activity of a lipophilic woad (*Isatis tinctoria*) extract. *Planta Medica*, **72**, 715-720.
- [13] Heinemann C, Schliemann-Willers C, Oberthür C, Hamburger M, Elsner P. (2004) Prevention of experimentally induced irritant contact dermatitis by extracts of *Isatis tinctoria* compared to pure tryptanthrin and its impact on UVB-induced erythema. *Planta Medica*, **70**, 385-390.
- [14] Fischbach W, Becker W, Mossner J, Ohlemüller H, Koch W, Borner W. (1987) Leukocytic elastase in chronic inflammatory bowel diseases: A marker of inflammatory activity. *Digestion*, **37**, 88-95.
- [15] Bieth JG. (1998) Leukocyte elastase. In *Handbook of Proteolytic Enzymes*. Barrett AJ, Rawlings ND, Woessner FF (Eds). Academic Press, London UK. 54-60.
- [16] Melzig MF, Löser B, Ciesielski S. (2001) Inhibition of neutrophil elastase activity by phenolic compounds from plants. *Pharmazie*, **56**, 967-970.
- [17] Dittmann K, Gerhäuser C, Klimo K, Hamburger M. (2004) HPLC-based activity profiling of *Salvia miltiorrhiza* for MAO A and iNOS inhibitory activities. *Planta Medica*, **70**, 909-913.
- [18] Rennert B, Melzig MF (2002) Free fatty acids inhibit the activity of *Clostridium histolyticum* collagenase and human neutrophil elastase. *Planta Medica*, **68**, 767-769.
- [19] Macdonald SJ, Dowle MD, Harrison LA, Shah P, Johnson MR, Inglis GG, Clarke GD, Smith RA, Humphreys D, Molloy CR, Amour A, Dixon M, Murkitt G, Godward RE, Padfield T, Skarzynski T, Singh OM, Kumar KA, Fleetwood G, Hodgson ST, Hardy GW, Finch H (2001) The discovery of a potent, intracellular, orally bioavailable, long duration inhibitor of human neutrophil elastase - GW311616A a development candidate. *Bioorganic and Medicinal Chemistry Letters*, **11**, 895-898.

Variation in Artemisinin and Flavonoid Content in Different Extracts of *Artemisia annua* L.

Anna Rita Bilia^{a*}, Caterina Gabriele^a, Maria Camilla Bergonzi^a, Pedro Melillo de Malgalhaes^b and Franco Francesco Vincieri^a

^aDepartment of Pharmaceutical Sciences, University of Florence, via Ugo Schiff, 6, Sesto Fiorentino-50019-Florence, Italy

^bDivisão de Agrotecnologia, CPQBA-UNICAMP, C.P. 6171, 13.081.970 Campinas, SP, Brazil

ar.bilia@unifi.it

Received: June 30th, 2006; Accepted: October 24th, 2006

Dedicated to the memory of Professor Ivano Morelli.

Artemisia annua L. is a promising and potent antimalarial drug. This activity has been ascribed to its content of artemisinin, a sesquiterpene lactone that is stage specific and very effective against drug-resistant *Plasmodium* species and which has low toxicity. The *in vitro* antiplasmodial activity of artemisinin is enhanced by the flavonoids of the extract, as recently proposed by the authors. Different extracts (tinctures, infusions and decoctions), obtained from a cultivar selected by the University of Campinas (0.52% artemisinin), were analyzed in order to prove the selectivity of the solvents to obtain high yields of both artemisinin and flavonoids. Tinctures 40 and 60% v/v showed a greater power of extraction in comparison with infusions and decoctions. The best performance was obtained using 60% v/v tincture. The extraction efficiency for artemisinin was 40% and for flavonoids was 29.5%. Among aqueous extracts, the best results were obtained by preparing an infusion with boiling water, left to cool for 15 minutes before filtration. The extraction efficiency for artemisinin was 57.5% and for flavonoids was 8.2%. If leaves are boiled for several minutes the artemisinin concentration is decreased, probably due to the heat instability of this constituent. Also microwave could represent a valid alternative method to extract the phytocomplex, the extraction efficiency for artemisinin was 41.0% and that for flavonoids was 7.7%.

Keywords: *Artemisia annua* L., extracts, artemisinin, flavonoids, HPLC/DAD/MS.

Artemisia annua L. (sweet or annual wormwood) is an annual herb endemic to the northern parts of Chahar and Suiyuan provinces in China, where it is known as ‘qinghao’ and has been used as a remedy for chills and fevers for more than 2000 years [1, 2]. Traditionally, the plant is used to prepare a drink, as indicated in the Compendium of Treatments (Ben Cao Gang Mu), written in 1596 AD by Li Shizhen: “take a handful of qinghao, soak it in a sheng (liter) of water, and squeeze out the juice and drink it all” [3, 4]. Nowadays the (daily) dose of *A. annua* given in the Chinese Pharmacopoeia for the treatment of various fevers, including malaria, is 4.5–9 g of dried plant material, extracted by heating with water [5, 6].

Surprisingly, only a few clinical studies based on the use of either extracts of the plant or traditional preparations are available. The first was carried out in 1992 in China: 144 malaria patients were treated with tablets containing either an ethanolic dried extract of *A. annua* or capsules using the same extract, but formulated in oil [7, 8]. Both treatments were found to be effective in reducing parasitaemia and fever at doses equivalent to 80.8 g (tablets) and 73.6 g (capsules) raw herb, given over a three-day period.

In the most recent literature, after development of high artemisinin-yielding plants (>0.5% per dried weight), clinical trials using either teas or decoctions have also been reported. The principal aim of such

investigations is related to the possibility for populations in endemic areas with either scarce or no access to modern medicines or medical services to cultivate selected cultivars of *A. annua* and prepare either teas or decoctions from the plant material, achieving a positive effect in the treatment of malaria.

An herbal tea prepared from a selected cultivar of *A. annua* with a high content of artemisinin (0.58% w/w dried herb) was evaluated in malaria patients in the Republic of Congo [9]. The dose was according to the recommendations of the Chinese Pharmacopoeia (5 g herbal drug/day for five days) and the extraction process was investigated in order to have the maximum extraction efficiency. It was proved that it is better to make an infusion rather than a short decoction (kept boiling for five minutes) in order to obtain the maximum extraction efficiency of more than 40%. In the study, about 90.9% of patients reported complete disappearance of malaria symptoms within the course of the treatment and it was proposed that the bioavailability of artemisinin from the tea preparations may exceed that from pure artemisinin tablets [9].

Two additional interesting papers regarding clinical studies using traditional preparations appeared almost contemporarily in 2004. In the first investigation [10], the patients received 1 L of preparation from 9 g leaves of a special cultivar of *A. annua* containing 1.39% artemisinin, which resulted in a content of 94.5 mg/L artemisinin, administered in five doses of 200 mL each per day. Even if the given dose of artemisinin was only 19% of the usual daily dose of

artemisinin in adults [11], peak plasma levels were 240 ± 75 ng/mL artemisinin, approximately 40% of the peak concentrations reported after intake of 500 mg artemisinin in the form of capsules. The data indicated that artemisinin was absorbed faster from herbal tea preparations than from oral solid dosage forms, but the bioavailability was similar [10]. The other study that appeared in 2004 used an herbal drug containing 1.4% artemisinin and the infusions (5 or 9 g/herb /L; artemisinin content 47 and 94 mg/L, respectively) were administered divided into four doses of 250 mL each. Even if there were a higher rate of recrudescence, most of the reported malaria symptoms either improved or resolved within three days after initiation of therapy, as expected for either an artemisinin or quinine treatment [12]. All these studies have pointed out that the presence of flavonoids in the phytocomplex can enhance either the bioavailability or the activity of artemisinin. Thus, experimental evidence from *in vitro* studies suggests that some flavonoids may enhance the action of artemisinin against *P. falciparum* [13]. In an attempt to find an optimal extraction method for both artemisinin and flavonoids, we have reported the best recovery with *n*-hexane to obtain complete extraction of artemisinin and most of the flavonoids [14].

The aim of the present study was to analyze the qualitative and quantitative composition of different extracts of the aerial parts of a cultivar of *A. annua* (0.52% artemisinin) selected by the University of Campinas. The investigated extraction methods were several techniques of infusion and decoction

Table 1: Artemisinin and flavonoid contents of tincture preparations (T40: 40% v/v; T60: 60% v/v).

Sample	artemisinin % in lyophilized material	artemisinin % in lyophilized material	extraction efficiency (%)	flavonoids % in lyophilized material	flavonoids % in lyophilized material	extraction efficiency (%)
T40	27	0.75	26	41.1	2.64	15.7
T60	41	1.08	40	83.8	3.52	29.5

Table 2: Artemisinin and flavonoid contents of infusion and decoction preparations.

Sample	artemisinin % in lyophilized material	extraction efficiency (%)	flavonoids % in lyophilized material	extraction efficiency (%)
I1	0.72	30.4	5.38	5.61
I2	0.68	27.4	3.24	5.41
I3	0.80	57.5	5.18	8.17
D1	0.68	30.2	3.28	5.34
D2	0.81	35.9	6.18	9.93
M	0.61	41.0	4.89	7.66

I1: sample extracted with 1 L of boiling water, left to cool, filtered and lyophilized; I2: sample extracted with 1 L of boiling water, covered, left to cool, filtered and lyophilized; I3: sample extracted with 1 L of boiling water, left to cool for 15 min, filtered and lyophilized; D1: sample extracted with 1 L of boiling water, kept boiling for 5 min, left to cool, filtered and lyophilized; D2: sample extracted with 1 L of boiling water, kept boiling for 5 min, immediately filtered and lyophilized; M: sample treated with 1 L of water, kept boiling for 5 min with a microwave oven, left to cool, filtered and lyophilized.

(see experimental part) and two tinctures (40% v/v and 60% v/v), prepared according to the European Pharmacopoeia [14].

The flavonoid structures were determined by combining the HPLC/DAD/MS data with those previously reported [15]. Artemisinin was identified by mass spectrometry as the peak at 17.90 min. *n*-Hexane can selectively and exhaustively extract both artemisinin and flavonoids and, for this reason, was considered as a standard for comparison with other preparations.

Very different yields and contents of artemisinin and flavonoids were found in the tested extracts, as reported in Tables 1 and 2. Concerning the percentage of constituents in the *n*-hexane extract, artemisinin made up 12.8% and the total flavonoids 62.8%. In particular, the respective percentages of different flavonoids were 34.3% for casticin and chrysosplenin, 9.4% for artemetin and 19.1% for eupatin. An additional peak was found in the tinctures and in aqueous preparations and identified as chrysosplenol-D by combining the HPLC/DAD/MS data with those previously reported [16].

The qualitative profile of the two tinctures was similar, while both the content of flavonoids and artemisinin was highest in the 60% v/v tincture, as reported in Table 1. The greater amount of ethanol in T60, compared with T40, increased the efficiency of the extraction. The extraction efficiency for artemisinin was 40% and for flavonoids 29.5%.

However, for all constituents of *A. annua*, the tinctures showed a greater power of extraction in comparison to infusions and decoctions, when the amount of solvent used was greater. In addition, after the freeze-drying process of the tinctures, the obtained dried powder of crude extract showed good technological properties and could be very useful in the formulation of either capsules or extemporaneous preparations.

In the case of infusions and decoctions, the highest contents of artemisinin and flavonoids were obtained from samples I3, D2 and M with 57.5%, 35.9% and 41.0% for artemisinin, respectively, and 8.2%, 9.9% and 7.7% for flavonoids. Artemisinin is known to be heat-unstable and Table 2 shows that tea prepared by adding boiling water to the leaves without further

heating (I3) yields higher artemisinin concentrations than if the leaves are boiled for several minutes (D2), according to literature data [9, 10]. No great differences were evidenced between infusion and decoction methods for flavonoid content, but an increase of extraction efficiency was obtained with filtration of the hot solution (I3 and D2). Also a microwave oven can represent a valid alternative method to extract the constituents of *A. annua*, in particular artemisinin.

Experimental

Chemicals: A sample of a selected high-yield cultivar of *A. annua* was provided by P.M.M. of the Universidade Estadual de Campinas (Brazil). The Brazilian hybrid plant was obtained according to the procedure carried out by MEDIPLANT [17]. The percentages of constituents of the herbal drug were artemisinin 0.52% and total flavonoids 2.6%.

Artemisinin was purchased from Sigma (Sigma-Aldrich S.r.l., Milan, Italy). Indena Research Laboratories (Settala, Milan, Italy) kindly provided the reference rutin trihydrate (batch no. K12408717, standard purity 88.17%, considering the content of residual solvents, moisture and amount of impurities), which was used for the calibration of polymethoxylated flavonoids.

All the solvents used for the extraction and HPLC analysis (MeOH, *n*-hexane, dichloromethane, and acetonitrile) were HPLC grade from Merck (Darmstadt, Germany); 85% formic acid was provided by Carlo Erba (Milan, Italy). Water was purified by a Milli-Q_{plus} system from Millipore (Milford, MA).

Preparation of the *n*-hexane extract: The dried aerial parts of a sweet wormwood sample were cut into small pieces with an Osterizer. Samples of 200 g material was exhaustively extracted at room temperature by maceration with 2 L of *n*-hexane for 72 h. The eluates were subsequently taken to dryness under reduced pressure to obtain the crude extract.

Preparation of the tinctures: The dried aerial parts of a sweet wormwood sample were cut into small pieces with an Osterizer. Samples of 10 g of material were extracted at room temperature by maceration with 100 g of ethanol [either 40 or 60% v/v (samples T40 and T60)].

Preparation of the decoctions: Sample D1: A 9 g sample of dried aerial parts of sweet wormwood was extracted with 1 L of boiling water, kept boiling for 5 min, then left to cool and filtered. Sample D2: A 9 g sample of dried aerial parts of sweet wormwood was extracted with 1 L of boiling water, kept boiling for 5 min and immediately filtered. Sample M: A 9 g sample of pieces of sweet wormwood was treated with 1 L of water, kept boiling for 5 min in a microwave oven, then left to cool and filtered. For analytical purposes, all the filtrates were lyophilized and provided 2.15 g, 2.09 g, and 3.16 g of dried product, respectively.

Preparation of the infusions: Sample 11: A 9 g sample of dried sweet wormwood was extracted with 1 L of boiling water, then left to cool and filtered. Sample 12: A 9 g sample of dried aerial parts of sweet wormwood was extracted with 1 L of boiling water, covered, then left to cool and filtered. Sample 13: A 9 g sample of dried aerial parts of sweet wormwood was extracted with 1 L of boiling water, then left to cool for 15 min and filtered. For analytical purposes, the filtrates were lyophilized, providing 2.04, 1.89 and 3.44 g of dried product, respectively.

Table 3: Mobile phases used for HPLC analysis.

Time(min)	A %	B %	C %	Flow (mL/min)
0.00	50	50	0	1.000
15.00	50	50	0	1.000
20.00	0	0	100	1.000
23.00	0	0	100	1.000
28.00	50	50	0	1.000

HPLC-DAD and HPLC-MS systems: The HPLC analyses were performed using a HP 1100 Liquid Chromatograph (Agilent Technologies, Palo Alto, CA, USA) equipped with a HP 1040 Diode Array Detector (DAD), an automatic injector, an auto sampler, a column oven and managed by a HP 9000 workstation (Agilent Technologies, Palo Alto, CA, USA).

Separations were performed on a reversed phase column Purospher®Star RP-18, namely Hibar®. The HPLC system was interfaced with a HP 1100 MSD API-electrospray (Agilent Technologies, Palo Alto, CA, USA). The interface geometry, with an

orthogonal position of the nebulizer with respect to the capillary inlet, allowed the use of analytical conditions similar to those of HPLC-DAD analysis. Mass spectrometry operating conditions were optimized in order to achieve maximum sensitivity values: gas temperature 350°C at a flow rate of 10 L/min, nebulizer pressure 30 p.s.i., quadrupole temperature 30°C, and capillary voltage 3500 V. Full scan spectra from m/z 100 to 800 in the positive ion mode were obtained (scan time 1 s).

A prepacked column RP (250 x 4.6 mm) with particle size 5 μ m (Merck, Darmstadt, Germany) was employed. The eluents were A: water adjusted to pH 3.2 with formic acid; B: acetonitrile; C: methanol. The mobile phase is reported in Table 3. The system was operated with oven temperature at 26°C. Before HPLC analysis, each sample was filtered through a cartridge-type sample filtration unit with a polytetrafluoroethylene (PTFE) membrane ($d = 13$ mm, porosity 0.45 μ m, (Lida Manufacturing Corp.) and immediately injected.

Chromatograms were recorded between 200 and 450 nm. DAD spectra were stored for all peaks exceeding a threshold of 0.1 mAu.

Calibration curves: A calibration curve, obtained from a methanolic solution of artemisinin (1 mg/mL), was used to quantify artemisinin in *n*-hexane extracts and tinctures, while a methanolic solution of artemisinin (0.5 mg/mL) was employed to determine the artemisinin content of infusions and decoctions. The flavonoid amounts were quantified by a methanolic solution of rutin international standard 0.03 mg/mL.

Sample analysis: Samples of 5 mg of the different extracts were accurately weighed and suspended in methanol (1.0 mL). The suspensions were sonicated for 10 min and filtered through a cartridge-type sample filtration unit before HPLC analysis. The tinctures were injected as prepared.

Acknowledgments - The financial support of MIUR (PRIN 2004) and Ente Cassa di Risparmio di Firenze are gratefully acknowledged.

References

- [1] Klayman DL. (1985) Qinghaosu (Artemisinin): an antimalarial drug from China. *Science*, **228**, 1049-1055.
- [2] Hien TT, White, NJ. (1993) Qinghaosu. *Lancet*, **341**, 603-608.

- [3] Dhingra V, Rao KV, Narasu ML. (2000) Current status of artemisinin and its derivatives as antimalarial drugs. *Life Sciences*, **66**, 279-300.
- [4] Li Y, Wu Y-L. (1996) How Chinese scientists discovered Qinghaosu (artemisinin) and developed its derivatives? What are the future perspectives? *Medicine Tropicale*, **58**, 9-12.
- [5] The People's Republic of China. (1985). In: *Pharmacopoeia*, vol. **I**. The People's Health Publisher, Beijing
- [6] Stöger EA. (1991). *Arzneibuch der Chinesischen Medizin*. Deutscher Apotheker-Verlag, Stuttgart, Germany.
- [7] Yao-De W, Qi-Zhong Z, Jie-Sheng J. (1992) Studies on the antimalarial action of gelatin capsule of *Artemisia annua*. *Chung Kuo Chi Sheng Ching Hsueh Yu Chi Sheng Chung Ping Tsa Chih*, **10**, 290-294.
- [8] Yu H, Zhong S. *Artemisia* species in traditional Chinese medicine and the discovery of artemisinin, In: Wright CW. (ed), *Artemisia*. (2002) London: Taylor and Francis, pp. 149-158.
- [9] Mueller MS, Karhagomba IB, Hirt HM, Wemakor E. (2000) The potential of *Artemisia annua* L. as a locally produced remedy for malaria in the tropics: agricultural, chemical and clinical aspects. *Journal of Ethnopharmacology*, **73**, 487-494.
- [10] Rāth K, Taxis K, Walz G, Gleiter CH, Li S-M, Heide L. (2004) Pharmacokinetic study of artemisinin after oral intake of a traditional preparation of *Artemisia annua* L. *American Journal of Tropical Medicine and Hygiene*, **70**, 128-132.
- [11] McIntosh HM, Olliaro P. (2001) *Artemisinin Derivatives for Treating Severe Malaria (Cochrane Review)*. The Cochrane Library 2. Oxford: Updated Software.
- [12] Mueller MS, Runyambo N, Wagner I, Borrmann S, Dietz K, Heide L. (2004) Randomized controlled trial of a traditional preparation of *Artemisia annua* L. in the treatment of malaria. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **98**, 318-21.
- [13] Elford BC, Roberts MF, Phillipson JD, Wilson RJ. (1987) Potentiation of the antimalarial activity of Qinghaosu by methoxylated flavones. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **81**, 434-436.
- [14] European Pharmacopoeia 5th Ed. (2005), pp. 571-572.
- [15] Bilia AR, Magalhães PM, Bergonzi MC, Vincieri FF. (2006) Simultaneous analysis of artemisinin and flavonoids of several extracts of *Artemisia annua* L. obtained from a commercial sample and a selected cultivar. *Phytomedicine*, **13**, 487-493.
- [16] Bilia AR, Lazari D, Messori L, Taglioli V, Temperini C, Vincieri FF. (2002) Simple and rapid physico-chemical methods to examine action of antimalarial drugs with hemin: Its application to *Artemisia annua* constituents. *Life Sciences*, **70**, 769-778.
- [17] Magalhães PM, Delabays N, Sartoratto A. (1997) *Ciência e Cultura. Journal of Brazilian Association for the Advancement of Science*, 49(5/6), September/December, 413-415.

Antifungal Evaluation of *Hypericum triquetrifolium* Polar Extracts Against *Fusarium* spp.

Daniele Fraternale^a, Alessandra Bertoli^c, Laura Giamperi^a, Anahi Bucchini^a, Donata Ricci^{a,*},
Francesco Menichini^c, Elena Trinciarelli^b and Luisa Pistelli^b

^aIstituto di Botanica e Orto Botanico "Pierina Scaramella", University of Urbino "Carlo Bo", Italy

^bDipartimento di Chimica Bioorganica e Biofarmacia, University of Pisa, Italy

^cDipartimento di Scienze Farmaceutiche, University of Calabria, (Arcavacata di Rende, Cosenza) Italy

d.ricci@uniurb.it

Received: July 28th, 2006; Accepted: November 8th, 2006

Dedicated to the memory of Professor Ivano Morelli.

The chemical composition of different kinds of *Hypericum triquetrifolium* extracts was analyzed by LC-DAD-ESI-MS. Hyperoside, rutin, isoquercitrin and biapigenin were the main constituents. As these natural compounds were reported in the literature for their antifungal activity, the total extracts were tested for their antifungal activity against eight phytopathogenic strains of *Fusarium* species.

Keywords: *Hypericum triquetrifolium*, flavonoids, *Fusarium* ssp., antifungal activity.

The genus *Hypericum* comprises more than 400 species, but only 20 can be found in Italy [1]. The species are herbaceous plants and are widely used in phytotherapy in many countries. Several phytochemical investigations of this genus have led to the isolation of many secondary metabolites, some with antidepressant, antimicrobial and antifungal activities [2-4]. Recently, plant extracts and phytochemicals with either antibacterial or antifungal properties have been investigated actively as alternatives to synthetic pesticides due to their perceived increased level of safety and minimal environmental impact [5]. Plant diseases often reduce quality and quantity of agricultural commodities. In fact, plant pathogens are estimated to cause yield reductions in crops of almost 20% worldwide [6,7]. Infestation by micro-organisms in post-harvest storage can effect the health of humans and livestock, especially if the contaminating organism produces toxic residues either in or on consumable products [8-10]. The fungicides made by synthesis provide the primary means for controlling post-harvest fungal decay of cereals,

fruits and vegetables. [8-10]. On the other hand, the extensive use of these synthetic fungicides causes uncontrolled residues and proliferation of resistance in the pathogen populations [11]. Therefore, studies concerning the possible use of biologically active natural products to control decay and prolong storage life of crops have received more and more attention [12,13].

The volatile compounds extracted from different species of *Hypericum* have been tested for their fungicidal activities on *Candida albicans* and *Saccharomyces cerevisiae* by several authors [14,15]. Some compounds from *Hypericum* species, such as xanthenes isolated from *H. roeperanum*, exhibited antifungal activity against *Candida albicans* [16], while xanthenes, a new γ -pyrone and betulinic acid from *H. brasiliense* showed similar activity against the plant pathogenic fungus *Cladosporium cucumerinum* [17]. Also a phloroglucinol derivative from the aerial parts of *H. calycinum* showed a fungicidal activity on the same phytopathogen [18]. Interesting activity of methanolic extracts of

Table 1: Percentage composition of the reference compounds in the analysed *H. triquetrifolium* extracts.

Extracts ^b	Compounds								
	1	2	3	4	5	6	7	8	9
	0.523 ^c	1.000	1.294	1.287	1.563	1.876	2.148	1.678	0.764
	Percentage composition (%)								
MM	6.93	29.8	26.3	21.7	3.12	2.72	4.90	0.3	-
RMMA	3.32	18.5	22.6	11.4	8.5	12.7	21.2	0.6	-
RMMB	5.53	16.8	70.2	-	-	-	-	-	-
SM	1.11	17.6	10.6	2.85	1.92	-	40.2	3.82	-

^a1 = chlorogenic acid; 2 = rutin; 3 = hyperoside; 4 = isoquercitrin; 5 = quercitrin; 6 = quercetin; 7 = I,II biapigenin; 8 = hypericin; 9 = hyperforin

^bMM = methanolic macerate; RMMA= ethyl acetate extract by the repartition of methanolic macerate; RMMB = butanolic extract by the repartition of methanolic macerate; SM = methanolic extract by Soxhlet extraction. ^cRRF= response factor of the constituent relative to rutin area/conc. (mg/mL) x purity/100

Table 2: % Inhibition of fungal strains of various extracts of *H. triquetrifolium*.

Fungal strain	Nystatin	MM			RMM A		
	100 ppm	3200	6400	10000	3200	6400	10000
I	100	0	20.2	50.0	66.6	70.0	70.0
II	100	0	25.3	50.0	66.6	70.0	70.5
III	100	0	24.3	50.4	71.4	70.0	70.0
IV	100	0	15.5	51.2	64.2	70.0	70.0
V	100	0	21.3	55.5	60.0	71.5	72.0
VI	100	0	24.6	50.5	57.1	65.6	67.0
VII	100	0	22.7	53.4	52.0	62.0	70.0
VIII	100	0	26.4	55.5	54.0	55.4	70.0

Fungal strain: *F. culmorum* (I), *F. graminearum* (II), *F. poae* (III), *F. avenaceum* (IV), *F. equiseti* (V), *F. semitectum* (VI), *F. sporotrichoides* (VII), *F. oxysporum* (VIII).

Extracts: MM = methanolic macerate; RMMB= ethyl acetate extract by the repartition of methanolic macerate.

The values are the average of three determinations.

Table 3: % Inhibition of fungal strains of various extracts of *H. triquetrifolium*.

Fungal strain	Nystatin	RMMB			SM		
	100 ppm	3200	6400	10000	3200	6400	10000
I	100	30.0	43.6	50.3	0	13.3	15.0
II	100	30.0	40.2	52.5	0	13.3	15.0
III	100	35.0	44.5	55.5	31.4	45.0	53.0
IV	100	28.0	34.7	50.3	10.7	32.1	40.0
V	100	25.0	36.2	54.5	0	16.0	20.0
VI	100	35.0	47.2	53.1	25.8	42.8	50.1
VII	100	25.0	33.9	50.0	0	8	13.0
VIII	100	25.0	41.5	54.3	0	8	15.0

Fungal strain: *F. culmorum* (I), *F. graminearum* (II), *F. poae* (III), *F. avenaceum* (IV), *F. equiseti* (V), *F. semitectum* (VI), *F. sporotrichoides* (VII), *F. oxysporum* (VIII); Extracts: RMMB = butanolic extract by the repartition of methanolic macerate; SM = methanolic extract by Soxhlet extraction.

The values are the average of three determinations

H. perforatum from Calabria, Italy against phytopathogenic fungi was also reported by Conforti et al. [19]. Other species of *Hypericum* were investigated for their biological activity, such as *H. triquetrifolium* Turra, native to Eastern Europe and the Mediterranean area. This species has been used for its sedative, anthelmintic, anti-inflammatory and antiseptic effects in folk medicine [5]. Extracts of *H. triquetrifolium* showed antimicrobial activity against *Staphylococcus aureus* and *Mycobacterium smegmatis*. In this work, the fungitoxic property of different polar extracts of *H. triquetrifolium*

collected in Calabria was evaluated for the first time against eight phytopathogenic strains of *Fusarium* species [20]. We tested the methanolic extracts, rich in hyperoside, rutin, isoquercitrin and biapigenin since some studies are reported in the literature on these main constituents of *Hypericum* and other related species [21]. Furthermore, the post-infection production of flavonoids and polyphenols in plant species suggests that these compounds might either function as phytoalexins [22] or have a protector role against fungal infection [23].

Some plants do not produce phytoalexins when challenged by pathogens, but release toxins that are normally stored as toxic glycosides in the vacuoles of their cells, for example phenolic and iridoid glycosides, glucosinolates and saponins [24]. If the integrity of the cells is broken when they are penetrated by fungal hyphae, the glycoside comes in contact with hydrolysing enzymes, present in other compartments of the same cell, releasing the toxic aglycone. Although this aglycone is not present *in situ* in the intact plant, it is not strictly a phytoalexin, because the involved enzymes (glycosidases) were already present in the healthy plant and not *de novo* formed [25]. The genus *Fusarium* contains a number of soil borne species with worldwide distribution, which have been known for a long time as plant pathogens and produce secondary metabolites toxic to plants (phytotoxins) and animals (mycotoxins), such as fusaric acid, trichothecenes, fumosins and enniatins [26]. Fusaric acid, a compound with moderate toxicity to plants and animals produced by many *Fusarium* species, was one of the first fungal metabolites implicated in plant pathogenesis. Furthermore the potential of *Fusarium* to serve as a model system for soil borne fungal pathogens is outlined [26].

A majority of known fungal secondary metabolites are not classified as mycotoxins. They may have toxic effects on insects (insecticides), plants (herbicides) and microorganisms (antibiotics) or they may have pharmacological effects on vertebrates or act synergistically with known mycotoxins on vertebrates. *Fusarium* spp produce a series of toxins, such as trichothecenes and zearalenones (responsible for several diseases in plants and animals) [27]. Cereals are often invaded by *Fusarium* species, before and after harvest, and the risk of trichothecene contamination of cereals is therefore of great concern [27]. Other kinds of *Fusarium* toxins, fusarins and fumonisins, have been

proposed to be involved in equine diseases [28]. Moreover, fumonisins are considered cancerogenic and they have been found to occur naturally [29]. Plant extracts rich in chlorogenic acid and isolated derivatives of chlorogenic acid showed antifungal activity on *Fusarium oxysporum*, as reported by Lattanzio [30] and Naidu [31], while other authors [32] demonstrated that the antifungal activity of caffeine in coffee beans was antagonized by chlorogenic acid. In tables 2 and 3 are reported data obtained by the agar diffusion test regarding the percent inhibition of the growth of *Fusarium* strains induced by our extracts. The *H. triquetrifolium* methanolic macerate (MM) tested in these experiments, which contained the higher amount of chlorogenic acid (6.93%) in comparison with the other extracts, exhibited a very weak antifungal activity on the selected eight *Fusarium* strains.

Hyperoside, rutin, and isoquercitrin were the main constituents of *H. triquetrifolium* in the most polar extracts. Hyperoside commonly occurs in a wide range of plants and has shown bactericidal activity [33]. However, several anomalous reports lead to uncertainty as to the antifungal activities of hyperoside. Previous studies demonstrated that it was inactive in *in vitro* bioassays against *Fusarium* spp. and other fungi at >100 µg/mL, while Dall'Agnol et al. reported that crude extracts of *Hypericum* including hyperoside showed no activity against yeast [34]. Hyperoside was tested for antifungal activity on several kinds of *Fusarium* and was considered more potent than some recently discovered natural antifungal products, including some fungicides on the market [35]. It is reported as an important secondary metabolite involved in the control of fungal pathogens *in vitro*, including *Fusarium* species, although the antifungal activity of these compounds in the plant is limited [21]. Hyperoside may serve as a lead compound for the development of fungicides [21]. Although its action against fungi is unknown, its effectiveness, resource availability at low cost, and probable low toxicity to humans make flavonoids potential prototypes for fungicides. Several studies have been carried out on the antifungal activities of these natural compounds. A recent report on the antifungal activity of flavonoids from *Pelargonium radula* showed that *Fusarium graminearum* was strongly inhibited only by the fraction rich in rutin, while the fraction with isoquercitrin as its main constituent inhibited *Candida tropicalis*, *C. lusitaniae* and *Microsporum gypseum* [36].

The extracts tested in this work (Table 1), containing good amounts of hyperoside, rutin and isoquercitrin [MM, RMMA and RMMB (where isoquercitrin was not detected)], did not exhibit major antifungal activity, with the exception of RMMA, which inhibits the growth of *F. poae* (71.4%) and all the other *Fusarium* strains, with values ranging from 52% to 67% at 3200 ppm. Although this was the most active extract, it was not comparable to nystatin (positive control). Only in RMMA are quercetin and quercitrin present in significant amounts (12.7% and 8.5%, respectively) and consequently the detected activity should be due to the presence of these compounds. The inhibitory action of quercitrin and quercetin, as pure compounds, on the mycelial growth of the crop pathogen, *Verticillium albo-atrum*, has been reported [37]. More recently, Conforti et al. [19] demonstrated an antifungal activity, particularly on the phytopathogenic fungus *Pythium ultimum*, of *Hypericum perforatum* extracts and its component, quercetin.

To the best of our knowledge, no data on the antifungal activity against *Fusarium* spp. are reported in the literature for biapigenin, which was the main constituent of the methanolic extract obtained by Soxhlet extraction, even if it was present also in the methanolic macerate. Although the compounds detected in the analyzed extracts as the main constituents have been reported in the literature for their antifungal activity against several microorganisms, such as *Fusarium* spp., these experiments did not show significant antifungal activity of the extracts that contained them.

Experimental

Plant material: The aerial parts of *Hypericum triquetrifolium* were collected on June 2004 at Isola Capo Rizzuto (Calabria, Italy). A voucher specimen (code CLU) was deposited in the herbarium of the Dipartimento di Botanica, University of Calabria, Italy.

Extraction and Purification of standard compounds: The pulverized dried material (400 g) was extracted at room temperature with methanol (MM 12.7 g). The MeOH residue was suspended in H₂O and then partitioned into EtOAc (RMMA 20.2 g) and *n*-BuOH (RMMB 7.6 g), successively. A portion of the EtOAc extract (10 g) was fractionated by gel-permeation and low pressure chromatography (eluted with MeOH) in order to isolate rutin (2)

130.6 mg), quercetin (**6**) (99.1 mg), quercitrin (**5**) (15.3 mg), isoquercitrin (**4**) (120.6 mg), hyperoside (**3**) (380 mg) and I3,II8 biapigenin (**7**) (33.1 mg), which were identified by comparison of their spectral data ($^1\text{H-NMR}$, $^{13}\text{C-NMR}$ and MS) with those reported in the literature [38-41]. Moreover, a portion of the powdered air-dried vegetable material (118 g) was defatted with light petrol in a Soxhlet apparatus and then extracted with methanol (SM 4.5 g). A preliminary screening by TLC [SiO_2 , BAW (60:15:10); RP-18, M-W (7:3); UV 254 nm and 366nm; NTS-PEG] was carried out on each extract in order to show the presence of the marker compounds (**1-9**) for the *Hypericum* genus. The isolated compounds (**2-7**) and the commercial standards of chlorogenic acid (**1**) (Extrasyntheses, Lot.01021203), hypericin (**8**) (Extrasyntheses, Lot: 02072309), and hyperforin (**9**) (Sigma, Lot. 092K1015), containing small quantities of impurities, were analyzed by HPLC-PDA in the same gradient conditions used for the extract samples in order to verify their purity (>98%) before using them as reference compounds.

Sample preparation and LC-DAD-ESI-MS-analyses: Three samples of each extract of *H. triquetrifolium* were dissolved in methanol (2 mg/mL) and filtered through a cartridge-type sample filtration unit with a polytetrafluoroethylene membrane before HPLC analyses (PTFE, 0.45 μm , 25 mm). All the extracts were analysed by the previously described method [42], slightly modified for our analytical equipment. The HPLC system consisted of a Waters W600E liquid chromatography pump equipped with an analytical Lichrosorb RP-18 column (250 x 4.6 mm i.d., 5 μm , Merck), a Rheodyne injection loop, and a Waters 996 photodiode array detector. The optimum efficiencies of separation were obtained using a linear gradient of a mobile phase of water with 0.1% HCOOH (solvent A), CH_3CN (solvent B) and MeOH (solvent C) at a flow rate of 1.0 mL/min. Gradient elution was carried out starting with a mixture of A-B-C (5:95:0) to (85:15:0) in 10 min, then to (50: 40:10) in 20 min, to (10:75:15) in 10 min, to B-A-C (5: 80:15) in 15 min, and then back to the initial condition in 10 min. Prior to running the gradient, the column was equilibrated for 10 min with solvents A and B (5:95 v/v). The total analytical run time for each sample was 65 min. The spectral data from the PDA detector were collected during the whole run in the range 210-600 nm and the peaks were detected at 270 nm and 590 nm.

Chromatographic procedures were performed at room temperature. An aliquot (20 μL) of each sample was analysed in triplicate. The same chromatographic conditions were used for LC-MS analyses performed using these ESI values: sheath gas flow-rate 62 psi, auxiliary gas flow 10 psi, capillary voltage -16 V and capillary temperature 200°C. Full scan spectra from m/z 200 to 700 u in the negative ion mode were obtained. The injected volume of the *Hypericum* extracts was 20 μL of a 1.5 mg/mL solution (methanol).

Identification of each constituent was achieved by comparison of the peak retention times, and UV and mass spectra of the extract sample with those of authentic samples (**1-9**). The standard solutions for the authentic samples (10 mg) (**1-7**) were prepared in methanol. Hypericin (**8**) (4.4 mg) was dissolved in pyridine (2 mL) before adding methanol (8 mL), while hyperforin (11.7 mg) was dissolved in a mixture MeOH -ascorbic acid 0.1% (10 mL). All standard and extract samples were injected alternatively. The linearity of the responses for the rutin reference standard (**2**) and for the constituents (**1, 3-9**) was determined at six levels of concentration with three injections for each level. Rutin was linear from 2.64 ppm to 264 ppm and all the curves had coefficients of linear correlation $r \geq 0.999$. The reproducibility of the injection integration procedure was determined for the constituents (**1-9**). The solutions were injected ten times and the relative standard deviation (R.S.D.) values were calculated (chlorogenic acid 1.56%, rutin 1.02%, hyperoside 0.78%, isoquercitrin 0.71%, quercitrin 0.50%, quercetin 0.56%, I3,II8-biapigenin 0.32%, hypericin 0.98%, hyperforin 1.54%). The repeatability of the method was evaluated by injection of three *H. triquetrifolium* extract solutions of different concentrations (0.5 mg/mL, 1.0 mg/mL, 1.5 mg/mL), each three times. The contents of constituents (**1-9**) were estimated by the following equation:

$$\text{Contents (\%)} = A_{\text{sample}} / \text{RF}_{\text{std}} \times C_{\text{sample}} \times 1 / \text{RRF} \times 100$$

where A_{sample} is the peak area of the considered constituent in the test solution, RF_{std} is the mean response factor of rutin in the reference solutions, C_{sample} is the concentration of the test solution (mg/mL) and RRF is the response factor of the considered constituent, relative to rutin (Table 1).

Biological screening: Fungal plant pathogens used in these tests were *Fusarium culmorum* (Smith)

Saccardo (I), *F. graminearum* Schwabe (II), *F. poae* (Peck) Wollenweber (III), *F. avenaceum* (Corda: Fries) (IV), *F. equiseti* (Corda) Saccardo (V), *F. semitectum* Berkeley et Ravenel (VI), *F. sporotrichoides* Sherbakoff (VII) and *F. oxysporum* Schl. (VIII), kindly supplied by the DI.PRO.VAL. (Dipartimento di Protezione e Valorizzazione Agro-alimentare, Università degli Studi di Bologna). All of the used microorganisms were maintained in potato dextrose agar (PDA, Sigma) and subcultured every 30 days.

The phytopathogenic fungi were tested by an agar dilution method. The extracts were dissolved in DMSO (Sigma) and added to the culture medium at a temperature of 40°-45°C, then poured into Petri dishes (Ø3 cm). Concentrations of 100, 200, 400, 800, 1600, 3200, 6400 and 10000 ppm were tested. The fungi were inoculated as soon as the medium had solidified. A disc (Ø 0.5 cm) of mycelial material, taken from the edge of seven-day old fungal cultures, was placed at the centre of each

Petri dish. The control consisted of a fungal disc placed in PDA, *Hypericum* extract free, + DMSO 1% v/v [43]. The Petri dishes with the inoculum were placed in the dark under controlled temperature conditions of 22 ± 1°C. The efficacy of treatment was evaluated after seven days by measuring the diameter of the fungal colonies when all the free surface of the medium in the control Petri dishes had been covered. The values were expressed in terms of percent inhibition of growth compared to control = 100. The fungicidal activity of the extracts was determined using the technique of Thompson [44] and Carta and Arras [45]: the mycelial discs were transferred from Petri dishes in which no growth was observed (total inhibition = 100) onto fresh plates of PDA, in order to verify, after three days, either the fungistatic or fungicidal activity of such inhibition. All tests were repeated three times.

Acknowledgments - The authors would like to acknowledge the financial support from Regione Marche, Progetto CIPE 17/2003.

References

- [1] Pignatti S. (1982) *Flora d'Italia*. Vol. I (Ed). Edagricole, Bologna IT. 344, 350.
- [2] Ishiguro K, Nagata S, Yamaki M, Fukumoto H, Koichiro I. (1994) Phloroglucinol derivatives from *Hypericum japonicum*, *Phytochemistry*, **35**, 469-471.
- [3] Decosterd L, Stockli-Evans H, Msonthi DJ, Hostettmann K. (1987) New antifungal chromenyl ketones and their pentacyclic dimers from *Hypericum revolutum* Vahl. *Helvetica Chimica Acta*, **70**, 1694-1702.
- [4] Jayasuriya H, Clark M, McChesney JD, Swanson M, Pezzuto J. (1989) Antimicrobial and cytotoxic activity of Rotlerin-type compounds from *H. drummondii*. *Journal of Natural Products*, **52**, 325-331.
- [5] Rai M, Mares D. (2003) Antimycotic activity of essential oils. In *Plant-derived Antimicrobials Current Trends and Future Prospects*. (Ed). Food Products Press, London UK. 307-320.
- [6] Seitz LM, Sauer DB, Mohr HE, Aldis DF. (1982) Fungal growth and dry matter loss during bin storage of high-moisture com. *Cereal Chemistry*, **59**, 9-14.
- [7] Alderman SC, Coats DD, Crowe FJ. (1996) Impact of ergot on Kentucky bluegrass for seed in northeastern Oregon. *Plant Disease*, **80**, 853-855.
- [8] Dieckman MA, Green ML. (1992) Mycotoxins and reproduction in domestic livestock. *Journal of Animal Science*, **70**, 1615-1627.
- [9] Nelson PE, Desjardins AE, Plattner RD. (1993) Fumonisin, mycotoxins produced by *Fusarium* species: biology, chemistry and significance. *Annual Review of Phytopathology*, **31**, 233-252.
- [10] Cheeke PR. (1995) Endogenous toxins and mycotoxins in forage grasses and their effects on livestock. *Journal of Animal Science*, **73**, 909-918.
- [11] Pujol I, Guarro J, Gene J, Sala J. (1997) *In vitro* antifungal susceptibility of clinical and environmental *Fusarium* spp. strains. *Journal of Antimicrobial Chemotherapy*, **39**, 163-167.
- [12] Costa TR, Fernandes OFL, Santos SC, Oliveira CMA, Liao LM, Ferri PH, Paulo JR, Ferreira HD, Sales BHN, Silva MRR. (2000) Antifungal activity of volatile constituents of *Eugenia dysenterica* leaf oil. *Journal of Ethnopharmacology*, **72**, 111-117.
- [13] Alvarez-Castellanos DP, Bishop CD, Pascual-Villalobos MJ. (2001) Antifungal activity of the essential oil of flowerheads of garland chrysanthemum (*Chrysanthemum coronarium*) against agricultural pathogens. *Phytochemistry*, **57**, 99-102.
- [14] Jang-Hee C, Jin-Cheol K, Moo-Key K, Hoi-Seon L. (2002) Fungicidal activities of 67 herb-derived oils against six phytopathogenic fungi. *Agriculture Chemistry and Biotechnology*, **45**, 202-207.
- [15] Schwob I, Bessiere JM, Dherbomez M, Viano J. (2002) Composition and antimicrobial activity of the essential oil of *Hypericum coris*. *Fitoterapia*, **73**, 511-513.
- [16] Rath G, Poterat O, Mavi S, Hostettmann K. (1996) Xanthenes from *Hypericum roeperanum*. *Phytochemistry*, **43**, 513-520.

- [17] Rocha L, Marston A, Kaplan MAC, Stoeckly E, Thuell U, Testa B, Hostettmann K. (1994) An antifungal γ -pyrone and xanthenes with monoamine oxidase inhibitory activity from *Hypericum brasiliense*. *Phytochemistry*, **36**, 1381-1385.
- [18] Decosterd LA, Hoffmann E, Kyburz R, Bray D, Hostettmann K. (1991) A new phoroglucinol derivative from *Hypericum calycinum* with antifungal and *in vitro* antimalarial activity. *Planta Medica*, **57**, 548-551.
- [19] Conforti F, Statti GA, Tundis R, Bianchi A, Agrimonti C, Sacchetti G, Andreotti E, Poli F, Menichini F. (2005) Comparative chemical composition and variability of biological activity of methanolic extracts from *Hypericum perforatum* L. *Natural Product Research*, **19**, 295-303.
- [20] Sakar MK, Tamer AÜ. (1990) Antimicrobial activity of different extracts from some *Hypericum* species, *Fitoterapia*, **61**, 464-466.
- [21] Li S, Zhang Z, Cain A, Wang B, Long M, Taylor J. (2005) Antifungal activity of camptothecin, trifolin, and hyperoside isolated from *Camptotheca acuminata*. *Journal of Agricultural and Food Chemistry*, **53**, 32-37.
- [22] McClure JW. (1975) in *The Flavonoids*, Harborne JB, Mabry TJ, Mabry H. eds, pp. 970-1055 Chapman & Hall, London
- [23] Smith IM. (1971) In: *Fungal Pathogenicity and the Plants Response*, (Ed). Cutting CV, Academic Press, New York USA.
- [24] Osburn AE, (1996) Performed antimicrobial compounds and plant defense against fungal attack. *The Plant Cell*, **8**, 1821-1831.
- [25] Grayer RJ, Kokubun T. (2001) Plant-fungal interactions: the search for phytoalexins and other antifungal compounds from higher plants, *Phytochemistry*, **56**, 253-263.
- [26] Bacon CW, Porter JK, Norred WP, Leslie JF (1996) Production of fusaric acid by *Fusarium* species. *Applied and Environmental Microbiology*, **62**, 4039-4043.
- [27] Betina V. (1993) In: *Chromatography of Mycotoxins: Techniques and Applications*. (Ed). Elsevier, London UK; "Liquid chromatography of mycotoxins" pp. 253-316.
- [28] Ghosal S, Biswas K, Chattopadhyay BK, (1978) Differences in the chemical constituents of *Mangifera indica*, infected with *Aspergillus niger* and *Fusarium moniliforme*. *Phytochemistry*, **17**, 689-694
- [29] Azcona-Oliveira J, Abouzied MM, Plattner RD, Pestka JJ. (1992) Production of monoclonal antibodies to the mycotoxins fumonisins B1, B2 and B3. *Journal of Agricultural and Food Chemistry*, **40**, 531-534
- [30] Lattanzio V, De Cicco V, Di Venere D, Lima G, Salerno M. (1994) Antifungal activity of phenolics against fungi commonly encountered during storage. *Italian Journal of Food Science*, **6**, 23-30.
- [31] Naidu GP. (1988) Antifungal activity in *Codiaeum variegatum* leaf extract. *Current Science*, **57**, 502-504.
- [32] Morozumi S, Wauke T, Kudoh Y, Hitokoto H. (1989) Antifungal effects of commercial foods and spices, and their components. *Bioactive Molecules*, **10**, 155-160.
- [33] Van der Watt E, Pretorius JC. (2001) Purification and identification of active antibacterial components in *Carpobrotus edulis* L. *Journal of Ethnopharmacology*, **76**, 87-91.
- [34] Dall'Agnol R, Ferraz A, Bernardi A, Albring D, Lamb L, Hass M, Schapoval EES. (2003) Antimicrobial activity of some *Hypericum* species. *Phytomedicine*, **10**, 511-516.
- [35] Sridhar SR, Rajavel RR, Masilamani S, Narasimhan S. (2003) Antifungal activity of some essential oils. *Journal of Agricultural and Food Chemistry*, **51**, 7596-7599.
- [36] Pepeljnjak S, Kalodera Z, Zovko M. (2005) Antimicrobial activity of flavonoids from *Pelargonium radula* (Cav) L'Herit. *Acta Pharmaceutica*, **55**, 431-435.
- [37] Picman AK, Schneider EF, Picman J. (1995) Effect of flavonoids on mycelial growth of *Verticillium albo-atrum*. *Biochemical Systematics and Ecology*, **23**, 663-683.
- [38] Agrawal PK, Markham KR. (1989) *Carbon-13 NMR of Flavonoids*. Agrawal PK (Ed). Elsevier, London UK. 513-530.
- [39] Markham KR, Ternai B. (1976) Carbon-13 NMR of flavonoids. II. Flavonoids other than flavone and flavonol aglycons. *Tetrahedron*, **32**, 2607-2612.
- [40] Seto T, Yasuda I, Akiyama K. (1992) Purgative activity and principals of the fruits of *Rosa multiflora* and *R. wichuraiana*. *Chemical and Pharmaceutical Bulletin*, **40**, 2080-2082.
- [41] Berghöfer R, Hölzl J. (1987) Biflavonoids in *Hypericum perforatum*. Part I. Isolation of I3,II8-biapigenin. *Planta Medica*, 216-217.
- [42] Brolis M, Gabetta B, Fuzzati N, Pace R, Panzeri F, Peterlongo FJ. (1998) Identification by high-performance liquid chromatography–diode array detection–mass spectrometry and quantification by high-performance liquid chromatography–UV absorbance detection of active constituents of *Hypericum perforatum*. *Journal of Chromatography A*, **825**, 9-16.
- [43] Nostro A, Cannatelli MA, Mugolino AD, Procopio F, Alonzo V. (2002) *Helichrysum italicum* extract interferes with the production of enterotoxins by *Staphylococcus aureus*. *Letters in Applied Microbiology*, **35**, 181-184.
- [44] Thompson DP. (1989) Fungitoxic activity of essential oil components on food storage fungi. *Mycologia*, **81**, 151-153.
- [45] Carta C, Arras G. (1987) Azione inibitrice in vitro di olii essenziali nei confronti di alcuni patogeni di piante ornamentali. *La difesa delle piante*, **10**, 195-202.

Antioxidant Activity Analysis for the Selection of *Rosmarinus officinalis* L.

Juan Antonio Garbarino^a, Nicolás Troncoso^b, Pia Delpiano^b, Loreto Carvajal^b and Alessandra Russo^c

^aDepartment of Chemistry, University T.F. Santa Maria, Casilla 110-V, Valparaíso, Chile

^bFundación Chile, Área Agroindustrias, Av. Parque Antonio Rabat Sur 6165 Vitacura, Santiago, Chile

^cDepartment of Biological Chemistry, Medical Chemistry and Molecular Biology, University of Catania, v.le A. Doria 6, 95125, Catania, Italy

alrusso@unict.it; ales0303@libero.it

Received: June 23rd; Accepted: August 25th, 2006

Dedicated to the memory of Professor Ivano Morelli.

Rosmarinus officinalis L. presents a high genetic variability, which is reflected in the chemical composition of the different individuals, and consequently in its biological activity, including antioxidant capacity. The aim of the present research was to correlate the chemical composition of methanolic extracts of the dried leaves of eight rosemary accessions with their antioxidant activity for the selection of plants to optimize the use of rosemary. The eight samples examined, starting from a collection of more than 160 individuals selected by BOTANE Ltd, were cultivated at Illapel, north central Chile, using the same cultivation techniques. The free radical-scavenging capacity was tested by the ability of extracts to bleach the stable 1,1-diphenyl-2-picryl-hydrazyl radical (DPPH) and to inhibit superoxide anion ($O_2^{\cdot-}$) and hydroxyl radical ($\cdot OH$) production. The metal chelating activity was estimated by the ferrozine assay. All extracts (1-8) contained high concentrations of carnosic acid, and to a minor extent rosmarinic acid, and exhibited antioxidant activity. However, extracts 7 and 8, containing 31.7 and 26.1% of carnosic acid, respectively, have shown a higher biological effect, confirming that the antioxidant activity of *R. officinalis* leaves is primarily related to this phenolic diterpene and suggesting that the measure of antioxidant activity could be considered a good method in the selection of this plant for its optimization. Interestingly, our experimental evidence also suggests that air pollution negatively influences the carnosic acid content. In fact, samples 3 and 4, with a low carnosic acid content, originated from a highly polluted metropolitan area of Santiago city.

Keywords: *Rosmarinus officinalis* L., leaf extract, carnosic acid, free radicals, antioxidant activity.

There is abundant evidence that reactive oxygen and nitrogen species (ROS and RNS) are implicated in several physiological processes, such as in host defence against invading pathogens and signal transduction. An overproduction of such reactive species, however plays a major role in several pathophysiological conditions. The ROS and RNS formed may cause cellular and subcellular damage by peroxidation of membrane lipids, by denaturing cellular proteins and by the breaking of DNA strands, disrupting cellular functions [1]. Lipid oxidation may also reduce the flavor and nutritive value of fats, oils and lipid-containing products. Unsaturated fatty acids

are sensitive to oxidation because of their chemical structure, and protein cross-linking, denaturation, polypeptide chain scission, enzyme inactivation and amino acid destruction in the presence of oxidizing lipids have been reported [1]. In this regard, many nutritive and non-nutritive phytochemicals, containing principally polyphenolic compounds and with diverse biological properties, have shown promising responses for the prevention and/or intervention of all diseases in which oxidative stress plays a key role [2]. A large number of polyphenolic compounds with antioxidant activity have been identified in the Labiatae plant

Rosmarinus officinalis L., including phenolic diterpenes such as carnosic acid, carnosol, rosmanol, epirosmanol, 7-methylepirosmanol, and methyl carnosate. In addition, several flavonoids, such as genkwanin, hispidulin 7-*O*-glucoside, cirsimaritin, luteolin, and isoscutellarein 7-*O*-glucoside, are found in Labiatae plants; the phenolic compounds rosmarinic and caffeic acids are also present [3-8]. *R. officinalis* (rosemary) is a typical Mediterranean species, but now is cultivated all over the world. Usually the plant is clonally propagated because of the poor germinability of its seeds and the genetic diversity of the seedlings [9]. *R. officinalis* presents, in fact, a high genetic variability, which is reflected in the chemical composition of the different individuals, and probably in its biological activity.

This plant is used as a spice and folk medicine around the world, as well as in cosmetics. The leaves are used in the preparation of alcoholic beverages, herbal soft drinks and in food preservation. In medicine, the extract is receiving increasing attention due to its antimicrobial, anti-inflammatory and antioxidative constituents [9].

The antioxidant properties of rosemary have been well documented, and there are several reports that have established carnosic acid as the major phenolic diterpenoid present in rosemary leaves with antioxidant activity [10]. Recently this phenolic compound has attracted wide interest as a potential therapeutic agent against several diseases, and research was started to investigate new biological activities. Studies showed that it has chemopreventive, anti-neoplastic [11-13] and radioprotective-antimutagenic [14] effects.

Therefore, the aim of the present research was to correlate the chemical composition of the methanolic extracts of leaves from eight different rosemary accessions, cultivated in Chile using the same cultivation techniques, with their antioxidant activity for the selection of *R. officinalis*, with the view to optimize its use.

Table 1: Origin of plant materials.

Accessions	Country
1	Santiago (Illapel), Chile
2	Santiago (Las Condes), Chile
3	Santiago (Renca), Chile
4	Santiago (Renca), Chile
5	Goodwood, Canada
6	Goodwood, Canada
7	Goodwood, Canada
8	Goodwood, Canada

The plant accessions were collected throughout Chile and other countries (Table 1), and were clonally propagated in order to maintain genetic uniformity. The fresh rosemary samples were dried, and double extraction for 159 hours was conducted for total extraction of carnosic and rosmarinic acids, which was confirmed by chromatographic analysis (data not shown). The yields of extraction of the samples are given in Table 2.

Table 2: Yield of extraction.

	Dry leaves g	First Extraction g	Second Extraction g	Total extract g	Yield %
1	58.0	5.4	4.8	10.2	17.6
2	50.3	5.3	3.1	8.4	16.7
3	52.1	5.6	5.0	10.6	20.3
4	51.7	4.9	3.3	8.2	15.9
5	51.3	3.7	3.5	7.2	14.0
6	50.1	5.0	4.0	9.0	18.0
7	55.4	6.6	6.1	12.7	22.9
8	65.9	6.1	5.9	12.0	18.2

Table 3: Content of carnosic acid and rosmarinic acid of methanol extracts of leaves from different accessions of *Rosmarinus officinalis*.

Extracts	Carnosic acid (%)	Rosmarinic acid (%)
1	13.8±2.6	1.14±0.11
2	19.3±2.2	0.46±0.13
3	14.3±2.6	0.79±0.08
4	10.8±3.2	1.17±0.11
5	11.7±3.0	0.84±0.08
6	12.8±2.8	1.32±0.15
7	31.7±4.9	0.41±0.14
8	26.1±3.1	0.43±0.08

n=6

The biological effects exhibited by these rosemary samples, under our experimental conditions, could be related to an overall effect of the phenolic compounds present in the extracts, but carnosic acid, as previously reported [10], seems to play a key role in the antioxidant activity. All extracts (1-8) containing high concentrations of carnosic acid (Table 3), and, to a minor extent, rosmarinic acid, exhibited antioxidant properties (Tables 4, 5). However, extracts 7 and 8 containing 31.7 and 26.1% of carnosic acid, respectively (Table 3), have shown a higher antioxidant capacity. In fact, the results, summarized in Table 4, showed that all extracts exhibited DPPH free radical scavenging activity, but samples 7 and 8 exhibited higher capacity with IC₅₀ values (concentration that inhibited radicals by 50%) of 9.2 and 8.6%, respectively. As DPPH is a synthetic radical, we also investigated the superoxide anion scavenging capacity of these extracts using the method of Paoletti [15], which excludes the Fenton-type reaction and the xanthine/xanthine oxidase system. Also in this assay, samples 7 and 8 showed a major superoxide scavenging effect (Table 4).

Table 4: Scavenger effect of leaf methanol extracts of different accessions of *Rosmarinus officinalis* on DPPH stable radical and superoxide anion.

	DPPH Test	Superoxide radicals
	^a IC ₅₀ (µg/mL)	
1	14.8±1.1	18.3±0.9
2	16.1±1.1	22.9±0.8
3	16.8±0.9	24.3±0.7
4	17.9±1.2	24.0±0.4
5	19.4±1.5	26.9±1.1
6	15.8±0.7	20.8±1.1
7	9.2±1.2	13.0±1.1
8	8.6±0.5	12.0±0.8
^b Trolox	95±1.4	-
^c SOD	-	87±3.4

^aconcentration that inhibited radicals by 50%. Values represent the mean ± SD of three experiments, performed in duplicate.

^bTrolox (50 µM) and ^csuperoxide dismutase (SOD) (80 mU/mL) were used as a standard; the results are expressed as % of inhibition.

Although both O₂^{•-} and H₂O₂ are potentially cytotoxic, most of the oxidative damage in biological systems is caused by the ·OH radical, which is generated by the reaction between O₂^{•-} and H₂O₂ in the presence of transition metal ions [1]. Based on the data obtained from this study, rosemary extract might also be able to modulate hydroxyl radical formation, acting as a direct scavenger and chelating ion. In fact, all extracts examined exhibited protection against DNA strand scission induced by ·OH radicals, generated by UV-photolysis of H₂O₂ (Table 5), and showed metal chelating activity capturing ferrous ions before ferrozine (Table 5). Also in these assays, samples 7 and 8 exhibited a higher effect (Table 5).

Table 5: Effect of methanol extracts of leaves from different accessions of *Rosmarinus officinalis* (100 µg/mL) on DNA cleavage induced by the photolysis of H₂O₂ and metal chelating activity.

	UD of supercoiled DNA	Ferrozine assay
	^a % of native DNA	^b IC ₅₀ (µg/mL)
scDNA	100	
1	46±1.4*	83.2±1.4
2	51±1.7*	89.4±1.8
3	53±2.1*	94.5±0.9
4	50±1.9*	96.7±1.2
5	55±1.8*	106.3±1.5
6	45±0.9*	86.3±0.7
7	83±1.6*	62.5±1.2
8	81±1.5*	57.5±0.8
^c DTPA	-	77±2.7

^aThe hydroxyl radicals generated by the photolysis of H₂O₂ reduced the supercoiled DNA (SCDNA).

^bconcentration that inhibited the ferrozine-Fe²⁺ formation by 50%.

^cDTPA (5 µM) was used as a standard; the results are expressed as % of inhibition.

Values represent the mean ± SD of three experiments, performed in duplicate. *significant vs. supercoiled DNA (*p*<0.001).

Interestingly, our results, similar to other works reporting that environmental elements affect carnosic acid concentrations [16], also reveal a correlation between carnosic acid concentration and

air pollution. In fact, samples 3 and 4, with low carnosic acid contents, 14.3 and 10.8%, respectively, correlating with a low antioxidant activity, originated from a heavily polluted metropolitan area of Santiago city, which ranks as one of the most polluted cities in the world. The main ambient pollutants of concern include carbon monoxide (CO), nitrogen oxides (NO), sulphur dioxide (SO₂), tropospheric ozone (O₃), and particulate matter (PM) [16-18].

In summary, our results suggest that the measure of antioxidant activity could be considered a good method for the selection of this plant for its optimization. Interestingly, our experimental evidence also suggests that air pollution negatively influences the carnosic acid content, justifying further studies to explain the mechanisms involved in this effect.

Experimental

Materials: Rosmarinic acid and carnosic acid, used as standards, were obtained from Addipharma and Sigma, respectively. The water used was Milli-Q quality, methanol and acetonitrile (MeCN) were from Merck and HPLC grade. Inorganic reagents from Merck were PA quality.

1,1-diphenyl-2-picryl-hydrazyl radical (DPPH), diethylenetriaminepentaacetic acid (DTPA), 3-(2-pyridyl)-5,6-bis (4-phenyl-sulfonic acid)-1,2,4-triazine (ferrozine) and pBR322 plasmid DNA were obtained from Sigma Aldrich Co (St. Louis, USA); β-nicotinamide-adenine dinucleotide (NADH) was obtained from Boehringer Mannheim GmbH (Germany). All other chemicals were purchased from GIBCO BRL Life Technologies (Grand Island, NY, USA).

Plant materials: Plant accessions were kindly borrowed from the special collection of more than 160 individuals used by BOTANE Ltd. to establish its crops and were collected throughout Chile and other countries (Table 1). Accessions were clonally propagated in order to maintain genetic uniformity. Each accession was propagated and established in a greenhouse and then transplanted to square-shaped blocks. Accessions were planted in November 2001 at Illapel, in north central Chile, about 250 Km north of Santiago (31°43'S; 71°07'W; 391 m.a.s.l.). Sampling was performed in a random pattern in the square shaped blocks and on freshly grown material.

Sample preparation: The fresh rosemary samples were dried at 40°C in a forced air circulation oven (Memmert ULM500). Leaves were manually separated and ground in a vertical hammer mill (Peruzzo Milly model 35.010) at 12,000 rpm and 0.8 mm mesh. Sample humidity was determined employing a Sartorius MA30 infrared system. Samples were mixed with 500 mL of methanol and stirred for 15 h at 20 °C in the dark. After stirring and filtering under vacuum, the filtrate was evaporated to dryness in a Rotavapor. The samples were extracted again for 144 hours, as previously described, and the filtrate was evaporated to dryness in a Rotavapor. Double extraction for 159 hours was conducted for total extraction of carnosic and rosmarinic acids, which was confirmed by chromatographic analysis. The extraction yields from the rosemary samples are given in Table 2.

Chromatographic conditions: A binary MeCN-H₂O acidified gradient was used for elution, as previously reported [19]. Two different procedures were developed. Method I, for simultaneous resolution of the three compounds of interest (CA, C, RA), the solvents A and B were MeCN-H₂O-H₃PO₄ (65.1%:34.9%:0.02%) and MeCN-H₂O-H₃PO₄ (22%:78%:0.25%), respectively. At a flow of 1.5 mL/min, the eluent consisted of 100% B during the initial 2 min, then the percentage of solvent A was increased to 100% at 2.1 min and remained at this level for the next 6 min. At 8.1 min the percentage of solvent B was again increased to 100%, where it remained for the last 2 min of the run time. With this method, the retention times were for RA *t_r* = 2.7 min, for C *t_r* = 5.7 min, and for CA *t_r* = 6.6 min. This method requires sample extraction with methanol: water (2:1) in order to extract all lipo-soluble and hydro-soluble antioxidants.

Method II: This chromatographic procedure is isocratic with solvent A as eluent for 6 min. The retention times were for C *t_r* = 1.8 min, and for CA *t_r* = 2.5 min. With this procedure, hydro-soluble compounds are not resolved, so sample extraction was simply performed with methanol. The detection wavelengths selected to quantify carnosic acid and rosmarinic acid were 230 and 330 nm, respectively, in order to avoid mobile phase absorption.

Antioxidant activity

Quenching of DPPH: Since the DPPH test can accommodate a large number of samples in a short period and is sensitive enough to detect natural

compounds at low concentrations, it was used in the present study for a primary screening of the methanolic extracts of *R. officinalis* free radical-scavenging activity. The assay provides information on the reactivity of test compounds with a stable free radical. Because of its odd electron, DPPH gives a strong absorption band at 517 nm in visible spectroscopy (deep violet color). As this electron becomes paired off in the presence of a free radical scavenger, the absorption vanishes, and the resulting decolorization is stoichiometric with respect to the number of electrons taken up. The reaction mixture contained 86 µM DPPH, and different concentrations of the extracts (5-100 µg/mL) in 1 mL of ethanol. After 10 min at room temperature, the absorbance at $\lambda = 517$ nm was recorded [20]. Trolox (50 µM), a water-soluble derivative of vitamin E, was used as a standard. A Hitachi U-2000 spectrophotometer (Hitachi, Tokyo, Japan) was used.

Scavenger effect on superoxide anion: Superoxide anion was generated *in vitro* during the autoxidation of β -mercapto-ethanol, as described by Paoletti *et al.* [15]. The assay mixture contained, in a total volume of 1 mL, 100 mM triethanolamine-diethanolamine buffer, pH 7.4, 3 mM NADH, 25 mM/12.5 mM EDTA/MnCl₂, 10 mM β -mercapto-ethanol; some samples contained methanolic extracts of the samples examined of *R. officinalis*, at different concentrations (5-100 µg/mL). After 20 min incubation at 25°C, the decrease in absorbance was measured at $\lambda = 340$ nm. Superoxide dismutase (SOD) (80 mU/mL) was used as a standard. A Hitachi U-2000 spectrophotometer (Hitachi, Tokyo, Japan) was used.

DNA cleavage induced by hydrogen peroxide UV-photolysis: The experiments were performed as previously reported [21], in a volume of 20 µL containing 33 µM in bp of pBR322 plasmid DNA in 5 mM phosphate saline buffer (pH 7.4), and the extracts. Immediately prior to irradiating the samples with UV light, H₂O₂ was added to a final concentration of 2.5 mM. The reaction volumes were held in caps of polyethylene microcentrifuge tubes, placed directly on the surface of a transilluminator (8000 µW cm⁻¹) at 300 nm. The samples were irradiated for 5 min at room temperature. After irradiation, 4.5 µL of a mixture containing 0.25% bromophenol blue, 0.25% xylen cyanol FF, and 30% glycerol were added to the irradiated solution. The samples were then analyzed by electrophoresis on a 1% agarose horizontal slab gel in Tris-borate buffer

(45 mM Tris-borate, 1 mM EDTA). Untreated pBR322 plasmid was included as a control in each run of gel electrophoresis, conducted at 1.5 V/cm for 15 hours. Gel was stained in ethidium bromide (1 µg/mL; 30 min), and photographed on Polaroid-Type 667 positive land film. The intensity of each scDNA band was quantified by means of densitometry.

Metal chelating activity: The chelating of ferrous ions by the methanolic extracts from the 8 samples examined of *R. officinalis* were estimated by the ferrozine assay [22]. Briefly, the extracts (5-200 µg/mL) were added to a solution of 0.15 mM FeSO₄. The reaction was initiated by the addition of 0.5 mM ferrozine and the mixture was shaken

vigorously and left standing at room temperature for ten minutes. After the mixture had reached equilibrium, the absorbance of the solution was then measured spectrophotometrically at 562 nm. DTPA (5 µM) was used as a standard. A Hitachi U-2000 spectrophotometer (Hitachi, Tokyo, Japan) was used.

Statistical analysis: Statistical analyses were performed using the statistical software package SYSTAT, version 9 (Systat Inc., Evanston IL, USA).

Acknowledgements - The authors gratefully thank CORFO for its financial support through the FDI AT-11 grant and Dr C. Wright for proof reading the manuscript.

References

- [1] Halliwell B. (1994) Free radicals, antioxidants and human disease: curiosity, cause of consequence? *Lancet*, **344**, 721-724.
- [2] Dryden GW, Song M, McClain C. (2006) Polyphenols and gastrointestinal diseases. *Current Opinion in Gastroenterology*, **22**, 165-170.
- [3] Aruoma OI, Halliwell B, Aeschbach R, Loligers J. (1992) Antioxidant and pro-oxidant properties of active rosemary constituents: carnosol and carnosic acid. *Xenobiotica*, **22**, 257-268.
- [4] Munne-Bosch S, Schwarz K, Alegre L. (1999) Enhanced formation of R-tocopherol and highly oxidized abietane diterpenes in water stressed rosemary plants. *Plant Physiology*, **121**, 1047-1052.
- [5] Cuvelier ME, Richard H, Berset C. (1996) Antioxidative activity and phenolic composition of pilot plant extracts of sage and rosemary. *Journal of American Oil Chemistry Society*, **73**, 645-652.
- [6] Del Bano M J, Lorente J, Castillo J, Benavente-García O, Mariñ MP, Del Rio JA, Ortuno, A, Ibarra I. (2004) Flavonoid distribution during the development of leaves, flowers, stems and roots of *Rosmarinus officinalis*. Postulation of a biosynthetic pathway. *Journal of Agricultural Food and Chemistry*, **52**, 4987-4992.
- [7] Lamaison JL, Petitjean-Freytet C, Carnat A. (1991) Medicinal Lamiaceae with antioxidant properties, a potential source of rosmarinic acid. *Pharmaceutica Acta Helvetica*, **66**, 185-188.
- [8] Cuvelier ME, Bondety V, Berset C. (2000) Behaviour of phenolic antioxidants in a partitioned medium: structure-activity relationship. *Journal of American Oil Chemistry Society*, **77**, 819-823.
- [9] Flamini G, Cioni PL, Morelli I, Macchia M, Ceccarini L. (2002) Main agronomic-productive characteristics of two ecotypes of *Rosmarinus officinalis* L. and chemical composition of their essential oils. *Journal of Agricultural and Food Chemistry*, **50**, 3512-3517.
- [10] Backle M, Leupold G, Parlar H. (2003) Rapid quantitative enrichment of carnosic acid from rosemary (*Rosmarinus officinalis* L.) by isoelectric focused adsorptive bubble chromatography. *Journal of Agricultural and Food Chemistry*, **51**, 1297-1301.
- [11] Offord EA, Gautier JC, Avanti O, Scaletta C, Runge F, Kramer K, Applegate LA. (2002) Photoprotective potential of lycopene, β-carotene, vitamin E, vitamin C and carnosic acid in UVA-irradiated human skin fibroblasts. *Free Radical Biological and Medicine*, **32**, 1293-1303.
- [12] Sharabani H, Izumchenko E, Wang Q, Kreinin R, Steiner M, Barvish Z, Kafka M, Sharoni Y, Levy J, Uskokovic M, Studzinski GP, Danilenko M. (2006) Cooperative antitumor effects of vitamin D3 derivatives and rosemary preparations in a mouse model of myeloid leukemia. *International Journal of Cancer*, **118**, 3012-3021.
- [13] Huang MT, Ho CT, Wang ZY, Ferraro T, Lou YR, Stauber K., Ma W, Georgiadis C, Laskin JD, Conney AH. (1994) Inhibition of skin tumorigenesis by rosemary and its constituents carnosol and ursolic acid. *Cancer Research*, **54**, 701-708.
- [14] Del Bano MJ, Castillo J, Benavente-Garcia O, Lorente J, Martin-Gil R, Acevedo C, Alcaraz M. (2006) Radioprotective-antimutagenic effects of rosemary phenolics against chromosomal damage induced in human lymphocytes by gamma-rays. *Journal of Agricultural and Food Chemistry*, **54**, 2064-2068.
- [15] Paoletti F, Aldinucci D, Mocalli A, Caparrini A. (1986) A sensitive spectrophotometric method for the determination of superoxide dismutase activity in tissue extracts. *Analytical Biochemistry*, **154**, 536-541.
- [16] Wellwood CR, Cole RA. (2004) Relevance of carnosic acid concentrations to the selection of rosemary, *Rosmarinus officinalis* (L.), accessions for optimization of antioxidant yield. *Journal of Agricultural and Food Chemistry*, **52**, 6101-6107.

- [17] Gil L, Martinez V, Riquelme R, Ancic P, Gonzalez G, Rodriguez L, Adonis M. (2003) Occupational and environmental levels of mutagenic PAHs and respirable particulate matter associated with diesel exhaust in Santiago, Chile. *Journal of Occupational Environmental Medicine*, **45**, 984-992.
- [18] Bell ML, Davis DL, Gouveia N, Borja-Aburto VH, Cifuentes LA. (2006) The avoidable health effects of air pollution in three Latin American cities: Santiago, Sao Paulo, and Mexico City. *Environmental Research*, **100**, 431-440.
- [19] Troncoso N, Sierra H, Carvajal L, Delpiano P, Gunther G. (2005) Fast high performance liquid chromatography and ultraviolet visible quantification of principal phenolic antioxidants in fresh rosemary *Journal of Chromatography A*, **1100**, 20-25.
- [20] Bonina F, Saija A, Tomaino A, Lo Cascio R, Rapisarda P, Dederen JC. (1998) *In vitro* antioxidant activity and *in vivo* photoprotective effect of a red orange extract. *International Journal of Cosmetic Sciences*, **20**, 331-342.
- [21] Russo A, Cardile V, Lombardo L, Vanella L, Vanella A, Garbarino JA. (2005) Antioxidant activity and antiproliferative action of methanolic extract of *Geum quellyon* Sweet roots in human tumor cell lines. *Journal of Ethnopharmacology*, **100**, 323-332.
- [22] Dinis TC, Madeira VM, Almeida LM. (1994) Action of phenolic derivatives (acetoaminophen, salicylate and 5-aminosalicylate) as inhibitors of membrane lipid peroxidation and as peroxy radical scavengers. *Archives of Biochemistry and Biophysics*, **315**, 161-169.

Hypericum perforatum L., *H. maculatum* Crantz., *H. calycinum* L. and *H. pulchrum* L.: Phytochemical and Morphological Studies

Gelsomina Fico*, Sara Vitalini, Noemi Colombo and Franca Tomè

Dipartimento di Biologia, Università degli Studi di Milano,
via Celoria 26, 20133 Milano, Italy

gelsomina.fico@unimi.it

Received: June 22nd, 2006; Accepted: July 12th, 2006

Dedicated to the memory of Professor Ivano Morelli.

Four species of *Hypericum* growing in Italy were characterized morphologically and chemically: *Hypericum perforatum* L., *H. maculatum* Crantz., *H. calycinum* L. and *H. pulchrum* L. The composition of secondary metabolites (phloroglucinols, naphthodianthrones, flavonoids) in the aerial parts of plants collected in different habitats was analysed. The four species show different compositions of phloroglucinols and naphthodianthrones, but there was no qualitative difference in flavonoid content of the species analysed. Study of main-constituent variation during the ontogenetic cycle showed that hypericin decreases and hyperforin increases during the reproductive phase. In St. John's Wort, hypericin and hyperforin are thought to be localised in black nodules. Our investigation shows no clear correlation between either the presence or absence of nodules and hypericin or hyperforin content.

Keywords: flavonoids, *Hypericum*, naphthodianthrones, ontogenetic cycle, phloroglucinols.

The genus *Hypericum* (Guttiferae) comprises herbs and shrubs, distributed all over the world, with long, opposite leaves and flowers usually organised in a terminal inflorescence.

Many ancient writers wrote about the medical properties of this genus and in particular of St. John's Wort, noting its use as a vulnerary and as a balm for wounds, burns, ulcers, and bites [1-2]. In recent years *Hypericum perforatum* has received increasing attention for the treatment of mild and moderate depression [3-4].

The great interest on *H. perforatum* and its potential for human health have encouraged us to investigate the productivity of some *Hypericum* species growing in Italy. In this work four species were studied: *H. perforatum* L., *H. maculatum* Crantz., *H. calycinum* L. and *H. pulchrum* L. All these species are herbaceous plants. *H. perforatum* is characterized by a two – winged stem and black nodules over the

whole plant; *H. maculatum* is different only for the four – edged stem. *H. pulchrum* is characterized by sessile leaves, small flowers, hirsute sepals with black nodules, and petals, stems and sepals with translucent glands. *H. calycinum* has typical inflorescences, but the black nodules and translucent glands are absent [5]. We have characterized each species chemically and morphologically. The chemical study has been concerned with the composition of flavonoids, phloroglucinols and naphthodianthrones; the morphological analysis has regarded the presence and distribution of secretory structures.

In this study we have analysed the secondary metabolites with interesting and demonstrated biological activity (**a**, chlorogenic acid; **b**, rutin; **c**, hyperoside; **d**, isoquercitrin; **e**, quercitrin; **f**, quercetin; **g**, hypericin; **h**, hyperforin) [3-4,6-8]. The MeOH extracts were analysed by RP-HPLC. The identification of peaks was effected on the basis of

the comparison of retention times and the use of a spectral library based on pure compounds previously described.

The qualitative analysis has shown four different profiles. The first profile, belonging to *H. perforatum* is characterized by the presence of all compounds under study: **a** ($\lambda = 270$ nm; Rt = 13.5), **b** ($\lambda = 270$ nm, Rt = 25.2), **c** ($\lambda = 270$ nm, Rt = 25.8), **d** ($\lambda = 270$ nm, Rt = 26.3), **e** ($\lambda = 270$ nm, Rt = 30.2), **f** ($\lambda = 270$ nm, Rt = 37.4), **g** ($\lambda = 590$ nm, Rt = 42.9), **h** ($\lambda = 270$ nm, Rt = 51.3). The second profile, typical of *H. maculatum* is characterized by the absence of hyperforin. The third profile is characterized by the absence of hypericin (*H. calycinum*). The fourth profile, where hypericin and hyperforin are absent, characterises *H. pulchrum*.

For quantitative analysis we produced eight calibration curves, as described in “experimental”. For all compounds, a linear relationship between peak area and concentration was observed, with a correlation coefficient always better than $r = 0.997$. Analysis of the four species under study was performed during the flowering phase (Table 1). In all species, the content of **a** was highly variable from

0.16 to 4.98%. Rutin (**b**) and hyperoside (**c**) (not always detectable separately) were the more abundant flavonoids, except in *H. calycinum* where quercitrin (**e**) and quercetin (**f**) were more important. *H. perforatum* was characterized by the presence of hypericin (**g**) (0.13-0.18 %) and hyperforin (**h**) (up to 10%). A considerable amount of **g** was also detected in *H. maculatum* and of **h** in *H. calycinum* (0.54%). The analysis of samples of *H. perforatum* collected at different altitudes showed that the contents of chlorogenic acid, flavonoids and hypericin seem not to be affected by the altitude; on the contrary very significant decreases were found in the **h** content in the sites at higher altitudes (from 10% to 3%) (Table 1).

Furthermore in *H. perforatum*, the analysis of metabolite content was performed during the reproductive phase (pre-flowering I, flowering II and fruiting phase III) with the following results. **a**: the content was variable in the different populations studied; **b**, **c**, **d**, **e**: there was a gradual decrease from the I to III phase; **f**: the content of this compound reached the maximum level during the II phase; **g**: the content decreased with values in a range of less than 10%; **h**: the content was low in the I phase and

Table 1: Secondary metabolites content during the flowering phase (% dry wt.).

Samples (Altitude)	a	b	c	d	e	f	g	h
<i>H. perforatum</i>								
4 (176 m)	3.45		6.65	2.27	2.97	0.14	0.13	10.05
8 (180 m)	4.98		7.24	7.10	1.50	0.47	0.13	10.76
9 (186 m)	4.70		6.12	3.20	4.10	0.10	0.13	10.10
5 (200 m)	4.62	9.35	2.08	1.30	5.44	0.10	0.15	10.65
12 (200 m)	0.16		7.24	5.89	3.04	0.45	0.14	10.20
7 (470 m)	2.27		8.24	3.70	2.18	0.23	0.13	7.50
11 (500m)	1.27	7.98	2.98	2.59	0.86	0.58	0.18	8.25
6 (800 m)	0.99	5.16	2.88	2.29	0.94	0.23	0.16	5.10
10 (900 m)	3.26	11.78	2.09	2.35	1.50	0.10	0.17	5.40
1 (1090 m)	0.63	4.08	1.89	1.71	1.72	0.57	0.14	3.54
3 (1400 m)	1.73	8.42	1.92	1.36	0.36	1.55	0.15	3.09
2 (1600 m)	0.47	11.13	2.07	1.56	0.47	0.70	0.13	3.74
<i>H. maculatum</i>								
13 (560 m)	0.64		10.72	3.16	0.19	2.13	0.12	-
<i>H. calycinum</i>								
14 (180 m)	-	0.30	0.63	0.37	2.28	1.54	-	0.54
<i>H. pulchrum</i>								
15 (560 m)	0.24	4.30	0.61	2.55	3.52	0.30	-	-

(a) chlorogenic acid, (b) rutin, (c) hyperoside, (d) isoquercitrin, (e) quercitrin, (f) quercetin, (g) hypericin and (h) hyperforin.

Table 2: Secondary metabolite content (% dry wt.) during the reproductive phase of *H. perforatum* collected in Comabbio.

	a	b	c	d	e	f	g	h
Phase I	3.69	12.95	3.56	3.48	2.09	0.12	0.13	4.80
Phase II	2.27		8.24	3.70	2.18	0.23	0.13	7.50
Phase III	0.56	3.75	1.26	1.07	1.51	0.15	0.11	8.00

Table 3: Localities and identification numbers of samples.

Samples	Locality	Herbarium No.
<i>H. perforatum</i>		
	Valle d'Aosta	
1	Anthey-St. André (AO - 1090 m)	Hy-101
2	Colle de Joux (AO - 1600 m)	Hy-102
3	Crest (AO - 1400 m)	Hy-103
	Lombardia	
4	Canegrate (MI - 176 m)	Hy-104
5	Collebeato (BS - 200 m)	Hy-105
6	Colle Brianza (LC - 800 m)	Hy-106
7	Comabbio (VA - 470 m)	Hy-107
8	Parabiago Canale Villoresi (MI - 180 m)	Hy-108
9	Parabiago Santa Maria (MI - 186 m)	Hy-109
10	Pezzaze (BS - 900 m)	Hy-110
	Friuli Venezia Giulia	
11	Costa (UD - 500 m)	Hy-111
12	Monte Spaccato (TS - 200 m)	Hy-112
<i>H. maculatum</i>		
	Friuli Venezia Giulia	
13	Ampezzo Carnico (UD - 560 m)	Hm-101
<i>H. calycinum</i>		
	Lombardia	
14	Brescia (BS - 180 m)	Hc-101
<i>H. pulchrum</i>		
	Piemonte	
15	Mondovì (CN - 560 m)	Hp-101

reached the maximum in the III phase, with an increase of more than 50%. Table 2 shows an example (sample collected in Comabbio).

Morphological characteristics

***H. perforatum*:** Leaves: presence, distribution and density of black nodules (b.n.) is variable among populations: sometimes only on the upper side of the lamina, and sometimes on both sides. Along with b.n., it is possible to find translucent glands, which confer the typical aspect at the leaves. Stems: b.n. are always present; there are only differences in the density of these structures. Petals and sepals: b.n. are always present on the borders. In some populations there are also secretory canals on all surfaces. Ovary: the surface is rich in translucent glands. B.n. are only present on the placenta. Stamen: one b.n. is always present between the thecae of anthers.

***H. maculatum*:** This species shows, in particular, red glands on the stems.

***H. calycinum*:** B.n. are completely absent.

***H. pulchrum*:** B.n. are only on the sepals. From the above results it appears that *H. maculatum* and *H.*

calycinum may be considered a good source of phloroglucinols and naphthodianthrone, respectively. It is worthwhile to notice that the production of **g** in *H. maculatum* is comparable, in quantity, to that of the well-known *H. perforatum*. It is also important to note the significant influence of altitude on the productivity of **h** in *H. perforatum*. Finally, even if many authors report that **g** and **h** are localised in b.n. [9-10], our investigation shows no clear correlation between the presence/absence of nodules and either hypericin or hyperforin content.

Experimental

Plant material: Fifteen populations of *Hypericum* belonging to the species *H. perforatum*, *H. maculatum*, *H. calycinum* and *H. pulchrum* were collected in different localities of Valle d'Aosta, Piemonte, Lombardia and Friuli Venezia Giulia, in Northern Italy during the summer of 2000 and determined according to Pignatti [5]. Voucher specimens are deposited in the Dipartimento di Biologia, Università di Milano. Table 3 shows localities, altitude and identification numbers of samples.

Extraction and separation: Dried powdered aerial parts (1 g), taken 20-25 cm from the apex, as described in the Italian F.U., were extracted in a Soxhlet apparatus with 200 mL of MeOH for six hours. From the extract solution, 4 mL was diluted to 10 mL and submitted to RP-HPLC on a Merck LiChrospher 100 RP-18 column (5 μ m, 250 x 4 mm, flow rate 1 mL min⁻¹) with ternary gradient elution [A: H₂O (acidified at 0.3% with H₃PO₄); B: ACN; C: MeOH; gradient: 0 min 100% A; 10 min 85% A, 15% B; 30 min 70% A, 20% B; 40 min 10% A, 75% B; 55 min 5% A, 80% B; minimum re-equilibration time between two injections: 10 min]. The detection range was 270-590 nm.

Chlorogenic acid (**a**), rutin (**b**), hyperoside (**c**), isoquercitrin (**d**), quercitrin (**e**), quercetin (**f**), hypericin (**g**) and hyperforin (**h**) were obtained commercially, **a-g** from Extrasynthese, Genay, France, and **h** from PhytoLab GmbH e Co. KG, Labor Addipharma, Hamburg, Germany. These

compounds were used to produce a spectral library in order to identify chromatographic peaks. The concentration of pure compounds was 0.4 mg mL⁻¹ and the injection volume was 15 μ L. The analytical chromatographic analyses were performed with a Merck-Hitachi L 6200 system with a Hewlett Packard 1040 photo diode array detector, controlled by HP-Chemstation (Hewlett Packard) software. Calibration curves for **a-h** were realized with solutions of known concentrations (0.4, 0.2, 0.1, 0.05, 0.025 mg/mL).

Morphological analyses: The morphological analyses were performed using a stereomicroscope, model MZ 6, Leica Microsystems S.p.A, Milano, Italy.

Acknowledgements - This research was carried out with the financial support of MIUR, Italy.

References

- [1] Matthioli A. (1668) *I discorsi di M. Pietro*. Edizione veneziana di Vincenzo Valglisi.
- [2] Fournier P. (1947) *Le livre des plantes médicinales et vénéneuses de France*. Paul Lechevalier Editeur, Paris.
- [3] Chatterjee SS, Noldner M, Koch E, Erdelmeier C. (1998) Antidepressant activity of *Hypericum perforatum* and hyperforin – the neglected possibility. *Pharmacopsychiatry*, **31**, (Suppl.), 7-15.
- [4] Francis AJP. (2005) Antidepressant action of St. John's Wort, *Hypericum perforatum*: a test of the circadian hypotheses. *Phytomedicine*, **12**, 167-172.
- [5] Pignatti S. (1982) *Flora d'Italia*. vol. 2, Edagricole, Bologna.
- [6] Ozturk Y. (1997) Testing the antidepressant effects of *Hypericum* species on animal models. *Pharmacopsychiatry*, **30** (Suppl. 2), 125-128.
- [7] Raffa RB. (1998) Screen of receptor and uptake-site activity of hypericin component of St. John's Wort reveals sigma-receptor binding. *Life Sciences*, **62**, 265-270.
- [8] Hosseinzadeh H, Karimi GR, Rakhshanizadeh M. (2005) Anticonvulsant effect of *Hypericum perforatum*: role of nitric oxide. *Journal of Ethnopharmacology*, **98**, 207-208.
- [9] Briskin DP, Leroy A, Gawienowski M. (2000) Influence of nitrogen on the production of hypericins by St. John's Wort. *Plant Physiology and Biochemistry*, **38**, 413-420.
- [10] Ciccarelli D, Andreucci AC, Pagni AM. (2001) The "black nodules" of *Hypericum perforatum* L. subsp. *perforatum*: Morphological, anatomical, and histochemical studies during the course of ontogenesis. *Israel Journal of Plant Sciences*, **49**, 33-40.

Chemical Composition and Antimicrobial Activities of Essential Oil of *Stachys glutinosa* L. from Sardinia

Pintore Giorgio^{a*}, Chessa Mario^a, Manconi Paola^a, Zanetti Stefania^b, Deriu Antonella^b and Tirillini Bruno^c

^aDipartimento Farmaco Chimico Tossicologico, Università di Sassari, Via Muroni 23/A, 07100 Sassari, Italy

^bDipartimento di Scienze Biomediche Sez. di Microbiologia sperimentale e clinica, Università di Sassari, Italy

^cIstituto di Botanica, Università di Urbino, via Bramante 28, I-61029 Urbino, Italy

pintore@uniss.it

Received: July 13th, 2006; Accepted: September 12th, 2006

Dedicated to the memory of Professor Ivano Morelli.

The oil composition of *Stachys glutinosa* L. from two different areas of Sardinia was analyzed by GC/MS. The oil from Gallura plants was characterized by the four main constituents: terpinen-4-ol (12.7%), α -terpinyl acetate (10.6%), *trans*-cadina-1(6),4-diene (8.5%), and α -terpineol (8.4%) whilst α -cedrene (19.2%), α -terpineol (18.5%), terpinen-4-ol (12.6%), and α -terpinyl acetate (8.6%) were the main compounds in the oil from Ulassai plants. The oils showed good bacteriostatic activities against *Vibrio cholerae* (MIC 0.6%), all the *Candida* tested (1.25%) and *Rodotorula rubra* (2.5%). There were also bactericidal activities against *Candida glabrata* (1.25 %) and *Rodotorula rubra* (2.5%).

Keywords: *Stachys glutinosa* L., Lamiaceae, Sardinia, essential oil, terpinen-4-ol, α -cedrene, antimicrobial.

Stachys glutinosa L. is a fruticose dwarf shrub, widespread in Sardinia, Corsica and the Capraia Islands [1], and is very common on different substrata from sea level to the mountains. The plant is covered by weak thorny stems that emanate an unpleasant smell [2]. The plant is used for medicinal purposes (mostly as an antispasmodic and antiseptic) in folk medicine [3]. A few studies refer to the composition of the essential oil of *S. glutinosa* from Corsica [4,5] and one about the principal compounds of essential oils from Sardinian plants [6].

The aim of this research was to determine the composition of the essential oil of wild *S. glutinosa* in different pedological soils and on the antimicrobial activities of the oils against soil-borne pathogens, mycotoxic species, phytopathogens and opportunistic human pathogens.

Table 1 shows the composition of the essential oils obtained from *S. glutinosa* L. plants harvested in the Gallura and Ulassai areas. Compounds are listed in order of their elution from an HP-5 column. In the oil from the Gallura plants, fifty-nine compounds were identified representing 95.7% of the oil, while in the oil from the Ulassai plants forty-seven compounds were identified representing 97.1% of the oil.

The oil from the Gallura plants was characterized by the four main constituents: terpinen-4-ol (12.7%), α -terpinyl acetate (10.6%), *trans*-cadina-1(6),4-diene (8.5%), and α -terpineol (8.4%). α -Cedrene (19.2%), α -terpineol (18.5%), terpinen-4-ol (12.6%), and α -terpinyl acetate (8.6%) were the main compounds in the oil from the Ulassai plants.

Table 1: Percentage composition of the essential oils of *S. glutinosa*.

Compounds	KI	Gallura (%)	Ulassai (%)
α -pinene	937	0.4	3.1
β -pinene	978	0.1	0.8
myrcene	996	n.d	0.4
α -phellandrene	1008	0.2	0.2
α -terpinene	1019	0.4	2.3
β -phellandrene	1029	3.5	4.7
γ -terpinene	1060	3.3	4.1
terpinolene	1089	0.3	1.1
<i>trans</i> -sabinene hydrate	1099	n.d	1.2
linalool	1106	1.6	n.d.
<i>cis</i> -p-menth-2-en-1-ol	1126	n.d.	0.5
<i>trans</i> -pinocarveol	1143	n.d.	0.1
<i>neo</i> -isopulegol	1147	n.d.	0.3
isopulegol	1148	0.1	n.d.
terpinen-4-ol	1177	12.7	12.6
thuj-3-en-10-al	1188	n.d.	0.1
α -terpineol	1189	8.4	18.5
Myrtenol	1198	n.d.	0.3
<i>trans</i> -piperitol	1211	n.d.	0.3
linalyl acetate	1260	4	0.5
terpinen-4-ol acetate	1301	0.4	n.d.
Carvacrol	1315	0.1	n.d.
<i>trans</i> -sabinyl acetate	1323	n.d.	1.1
δ -elemene	1329	1	n.d.
α -terpinyl acetate	1348	10.6	8.6
α -copaene	1366	0.2	n.d.
β -bourbonene	1373	0.9	0.8
β -cubebene	1381	0.1	0.1
β -elemene	1383	0.5	n.d.
geranyl acetate	1386	2.9	n.d.
(<i>Z</i>)- β -damascone	1399	n.d.	0.3
(<i>Z</i>)-caryophyllene	1404	2.9	n.d.
α -cedrene	1414	n.d.	19.2
β -cedrene	1416	0.6	0.3
β -copaene	1426	0.1	n.d.
β -gurjunene	1432	0.1	n.d.
(<i>Z</i>)- β -farnesene	1441	0.2	0.5
α -himachalene	1447	0.8	n.d.
Alloaromadendrene	1451	n.d.	0.7
<i>cis</i> -muurola-3,5-diene	1452	0.1	n.d.
<i>trans</i> -muurola-3,5-diene	1454	0.1	n.d.
(<i>E</i>)- β -farnesene	1456	0.8	n.d.
<i>cis</i> -muurola-4(14),5-diene	1465	0.1	n.d.
<i>trans</i> -cadina-1(6),4-diene	1470	8.5	0.9
γ -curcumene	1474	0.3	n.d.
γ -himachalene	1479	0.1	n.d.
germacrene D	1487	n.d.	0.4
<i>cis</i> - β -guaiene	1493	n.d.	0.1
Bicyclogermacrene	1495	2.9	n.d.
α -muurolene	1497	0.5	n.d.
<i>trans</i> -muurola-4(14),5-diene	1499	0.1	n.d.
<i>trans</i> - β -guaiene	1501	1.1	0.2
germacrene A	1508	1.5	0.2
δ -amorphene	1514	3.2	0.9
δ -cadinene	1522	0.1	n.d.
<i>trans</i> -calamenene	1529	0.2	0.2
Elemol	1542	1.8	n.d.
geranyl butanoate	1566	n.d.	0.3
Spathulenol	1570	2.2	n.d.
caryophyllene oxide	1573	n.d.	6.8
Globulol	1579	0.7	0.3
Viridiflorol	1589	0.9	0.2
Guaiol	1602	0.4	n.d.
10- <i>epi</i> - γ -eudesmol	1621	0.3	n.d.
10- <i>epi</i> - α -eudesmol	1625	n.d.	0.1
caryophylla-4(14),8(15)-dien-5-ol *	1630	n.d.	0.6
γ -eudesmol	1631	1.1	n.d.
α -muurolol	1637	n.d.	0.2
<i>epi</i> - α -cadinol	1638	0.7	1.2
<i>epi</i> - α -muurolol	1642	0.4	n.d.
Cubenol	1645	2.6	0.6

Table 1 (contd.)

Valerianol	1658	0.3	n.d.
14-hydroxy-9- <i>epi</i> -(<i>E</i>)-caryophyllene	1667	n.d.	0.4
helifolenol A	1674	0.4	n.d.
<i>epi</i> - α -bisabolol	1677	5.7	n.d.
eudesma-4(15),7-dien-1- β -ol	1683	0.9	0.4
<i>epi</i> -laurenene	1890	0.7	n.d.
Isopimara-9(11),15-diene	1894	n.d.	0.4
Sclarene	1986	0.4	n.d.

* = correct isomer not identified

n.d. = not detected

The sesquiterpene hydrocarbons contributed the highest percentage (27.0%) of the oil from the Gallura plants. This fraction was dominated by *trans*-cadina-1(6),4-diene (8.5%). The oxygenated monoterpene fraction represented the 22.9% of the total oil, terpinen-4-ol (12.7%) being the most abundant compound. The oxygenated sesquiterpenoids and esters were also relatively high representing 18.4% and 17.9% of the total oil respectively, whereas in the oil from the Ulassai plants the oxygenated monoterpenes constituted the highest portion (33.9%). This fraction was also dominated by α -terpineol (18.5%). The sesquiterpene hydrocarbons represented 24.5% of the total oil, α -cedrene (19.2%) being the major compound. The monoterpene hydrocarbons, oxygenated sesquiterpenes and esters were also relatively high representing 16.7%, 10.8% and 10.2%, respectively, of the total oil.

If we consider the principal components, α -cedrene distinguishes the Ulassai oil from the Gallura oil. *Epi*- α -bisabolol, on the contrary, distinguishes the Gallura oil from the Ulassai oil, with 5.7% in the former and none detected in the latter. Other minor components present only in Gallura oil were (*Z*)-caryophyllene (2.9%), bicyclogermacrene (2.9%), elemol (1.8%), γ -eudesmol (1.1%), and δ -elemene (1%), whereas, caryophyllene oxide (6.8%), *trans*-sabinene hydrate (1.2%) and *trans*-sabinyl acetate (1.1%), were the minor compounds present only in the Ulassai oil. Previous studies on *S. glutinosa* harvested in Corsica and Sardinia have proposed the three chemotypes A, B and C, characterized by the presence of α -terpineol-*trans*-caryophyllene, β -phellandrene, and terpinen-4-ol [5]. According to these proposed division, both the oils might belong to chemotype C.

Antimicrobial activity: The two tested oils of *S. glutinosa* presented similar antimicrobial activities the mean values of Minimal Inhibitory Concentration (MICs) and Minimal Bactericidal Concentration

(MBC) as summarized in Table 2. The oil of *S. glutinosa* exhibited good bacteriostatic effects against yeast clinical strains, particularly *Candida glabrata*, with the same value of MIC and MBC (1.25%) (*Rodotorula rubra* (MIC and MBC = 2.5%) and against *V. cholerae* 01 (MIC = 0.6%). The oils exhibited moderate bacteriostatic and bactericidal activities in general, but had a good bacteriostatic activities against *Vibrio cholerae* (0.6%). Other interesting bacteriostatic activities were against all the *Candida* tested (1.25%) and *Rodotorula rubra* (1.25%). Bactericidal activities at the tested concentration were against *Candida glabrata* (1.25%) and *Rodotorula rubra* (1.25%). Significantly, all the clinical and environmental strains to have multi-drug resistance, for example: *A. hydrophila* to ampicillin (192 µg/mL), ceftazidime (125 µg/mL), and gentamicin (125 µg/mL), *S. epidermidis* to ampicillin/sulbactam (64 µg/mL), norfloxacin (125 µg/mL), and gentamicin (125 µg/mL); *V. cholerae* to ampicillin (256 µg/mL), ceftazidime (125 µg/mL), cefotaxime (256 µg/mL), doxycycline (32 µg/mL), and amoxycillin-clavulanate (256 µg/mL). All the bacteria are susceptible to streptomycin (range 0.010 – 0.06 mg/mL) and the yeasts are susceptible to bifonazole (0.02 – 0.05 mg/mL). Other studies [7] of the antimicrobial activity of *Stachys* essential oil was of different species and subspecies.

Table 2: Antimicrobial activity of the essential oil of *Stachys glutinosa*.

Microorganism	MIC(%)	MCB(%)
<i>Aeromonas sobria</i>	1.25	2.5
<i>Candida albicans</i>	1.25	>2.5
<i>Candida glabrata</i>	1.25	1.25
<i>Candida krusei</i>	1.25	1.25
<i>Candida parapsilosis</i>	1.25	>2.5
<i>Enterococcus faecalis</i>	>2.5	>2.5
<i>Escherichia coli</i>	>2.5	>2.5
<i>Klebsiella pneumoniae</i>	>2.5	>2.5
<i>Aeromonas hydrophila</i>	1.25	>2.5
<i>Rodotorula rubra</i>	1.25	1.25
<i>Staphylococcus aureus</i> (ATCC)	>2.5	>2.5
<i>Staphylococcus aureus</i>	>2.5	>2.5
<i>Staphylococcus epidermidis</i>	2.5	2.5
<i>Streptococcus group D</i>	2.5	2.5
<i>Vibrio alginolyticus</i>	>2.5	>2.5
<i>Vibrio cholerae</i>	0.6	2.5

Experimental

Collection of Plant Material: Plants of *S. glutinosa* L. growing in Ulassai (central Sardinia) on limestone, and Gallura (northern Sardinia) on granite were collected during flowering, in June 2004. Voucher specimens were deposited in the Herbarium SASSA [Dipartimento di Scienze del Farmaco, Università di Sassari], under Acquisition No. 1099

Isolation of the Essential Oil: Fresh plant material was subjected to hydrodistillation using a Clevenger-type apparatus for 2 h yielding 0.25% of yellowish oil. The oil was dried over anhydrous sodium sulfate and stored in sealed vials under refrigeration prior to analysis.

Gas Chromatography: The GC analyses were carried out using a Hewlett Packard 5890 Series II dual FID instrument equipped with HP-WAX and HP-5 capillary columns (30 m x 0.25 mm, 0.25 µm film thickness), working with the following temperature programmed: 10 min at 60°C, and subsequently at 5°C/min up to 220°C; injector and detector temperatures, 250°C; carrier gas, helium (1 mL/min); split ratio, 1 : 20.

Gas Chromatography-Mass Spectrometry: GC-MS analyses were carried out using a Hewlett Packard 6890-5973 GC-MS system operating in the EI mode at 70 eV, using two different columns, a HP Innowax (30 m x 0.25 mm, film thickness 0.50 µm) capillary column and a DB-5 (30 m x 0.25 mm, film thickness 0.25 µm) capillary column. The temperature programmed for HP Innowax was 60-260°C at a rate of 3°C/min, held for 10 min, and for the HP 5 it was 60-300°C at a rate of 3°C/min. Injector and transfer line temperatures were 220°C and 280°C, respectively. Helium was used as the carrier gas, flow rate 1 mL/min. Split ratio, 1 : 10.

Identification of the Components: The identification of the components was made for both the columns, by comparison of their retention time with respect to n-paraffin (C6-C22) internal standards. The mass spectra and retention indices (RI) were compared with those of commercial (NIST 98 and WILEY) and home-made library mass spectra built up from pure compounds and MS literature data [8,9,10,11,12,13].

Area percentages were obtained electronically from the GC-FID response without the use of an internal standard or correction factors.

Microorganisms: A total of 17 strains of bacteria and yeast were investigated, ten were isolated from patients, three from environmental sources and four ATCC (American Type Culture Collection) strains were used as quality control strains. The isolates were identified to the species level by standard procedures, and some clinical and environmental ones were tested for virulence phenotype. Antimicrobial susceptibility to 14 different antibiotics (ampicillin, amoxycillin

clavulanate, amikacin, piperacillin, cefotaxime, ceftazidime, ceftriaxone, ciprofloxacin, ofloxacin, gentamicin, doxycycline, imipenem, meropenem, trimethoprim-sulphamethoxazole) was determined by the Kirby Bauer method, according to the general qualitative assay described by Barry (1986) [14]. Protease activity was tested on Nutrient agar containing 1.5% skim milk, production of protease was shown by the formation of a clear zone caused by casein degradation; hemolysis test was assayed by culturing each strain on agar plates containing rabbit erythrocytes 5%; Hep-2 cells (human laryngeal carcinoma) were used for the adhesion assay. The bacteriostatic and bactericide activities were determined by measuring the Minimal Inhibitory Concentration (MICs) and the Minimal Bactericidal Concentration (MBC) of *Stachys glutinosa* oil performed in microtiter plates using a bacterial inoculum (taken from Luria Berani broth after overnight culture) with a turbidity equivalent to 0.5 MacFarland standard. The essential oils were suspended in the medium with a 0.5% Tween 80 as

emulsifier and tested at different concentrations; values of MICs and MBCs are expressed as percent vol/vol of total oil and culture medium used as diluent. Bacterial strains were as follows: *Aeromonas hydrophyla* (isolated from patients), *Aeromonas sobria* (isolated from patients), *Candida albicans* (isolated from patients), *Candida glabrata* (isolated from patients), *Candida krusei* (isolated from patients), *Candida parapsilosis* (isolated from patients), *Enterococcus faecalis* (ATCC 24212), *Escherichia coli* (ATCC 35218), *Klebsiella pneumoniae* (ATCC 700603), *Rodotorula rubra* (isolated from patients), *Staphylococcus aureus* (isolated from patients), *Staphylococcus aureus* (ATCC 43300), *Staphylococcus epidermidis* (isolated from patients), *Streptococcus group D* (isolated from patients), *Vibrio alginolyticus* (soil-borne pathogens), and *Vibrio cholerae* (soil-borne pathogens). All micro-organism species were tested in triplicate.

Acknowledgments - The work was financially supported by INTERREG III and ex 60% MURST.

References

- [1] Pignatti S. (1982) In *Flora d'Italia*, Edagricole:Bologna, 462-469.
- [2] Camarda I. (1980) – Le piante endemiche della Sardegna. 70. *Stachys glutinosa* L. *Bollettino Società Sarda di Scienze Naturali*, **19**, 261-267.
- [3] Atzei A. (2003) In *Le Piante nella Tradizione Popolare della Sardegna*, Carlo Delfino Editore
- [4] Pelissier Y, Marion C, Quastana C, Milhau M, Malan A, Bessiere JM. (1996) Composés volatils de lamiaceae originaires de Corse. *Rivista Italiana EPPOS*, **7** (Spec. Num.), 526-535.
- [5] Mariotti JP, Costa J, Bianchini A, Bernardini AF, Casanova J. (1997) Composition and Variability of the Essential Oil of *Stachys glutinosa* L. from Corsica (France). *Flavour and Fragrance Journal*, **12**, 205-209.
- [6] Mariotti JP, Tomi F, Bernardini AF, Costa J, Casanova J. (1996) Composition chimique d'huile essentielles de *Stachys glutinosa* de Corse et de Sardigne. *Rivista Italiana EPPOS* 1996, **7** (Spec. Num.), 536-540.
- [7] Skaltsa HD, Demetzos C, Lazari D, Sokovic M. (2003) (Essential oil analysis and antimicrobial activity of eight *Stachys* species from Greece. *Phytochemistry*, **64**, 743-752.
- [8] Adams RP. (2001) In *Identification of Essential Oil Components by Gas Chromatography/Quadrupole Mass Spectroscopy*, Allured Publishing Corporation, Carol Stream, IL, USA.
- [9] Davies NW. (1990) Gas chromatographic retention indices of monoterpenes and sesquiterpenes on methyl silicone and Carbowax 20M phases. *Journal of Chromatography A*, **503**, 1-24.
- [10] Heller SR, Milne GWA. (1983) In *EPA/NIH Mass Spectral Data Base*, U. S. Government Printing Office: Washington, DC.
- [11] Jennings WG, Shibamoto T. (1980) In *Qualitative Analysis of Flavour and Fragrance Volatiles by Glass Capillary Gas Chromatography*, Academic Press, New York.
- [12] McLafferty FW, Staufner DB. (1989) In *The Wiley NBS Registry of Mass Spectral Data*, John Wiley and Sons, New York.
- [13] Stenhagen E, Abrahamsson S, McLafferty FW. (1974) In *Registry of Mass Spectral Data*, John Wiley and Sons, New York.
- [14] Barry AL. (1986) In *Procedure for Testing Antimicrobial Agents in Agar Media: Theoretical Considerations*, ed. V. Lorian, Baltimore, 1986.

Molecular Identification of *Panax ginseng* C.A. Meyer in Ginseng Commercial Products⁺

Paola Del Serrone^a, Lucilla Attorri^b, Bruno Gallinella^b, Francesca Romana Gallo^b,
Elena Federici^b and Giovanna Palazzino^{b,*}

^aConsiglio per la Ricerca e la Sperimentazione in Agricoltura, Via Salaria 31, I-00016 Monterotondo, Rome, Italy

^bIstituto Superiore di Sanità, Viale Regina Elena 299, I-00161 Rome, Italy

*palazzin@iss.it

Received: May 13th, 2006; Accepted: May 27th, 2006

Dedicated to the memory of Professor Ivano Morelli.

Molecular techniques (PCR and RFLP) were used to verify the presence of *Panax ginseng* C.A. Meyer in commercial products containing ginseng. DNA, extracted from four vegetable forms present in marketed products, was amplified with 18df/28ccr primers. The RFLP of the DNA amplified products, obtained using *Inf* I, *Sau* 3A1 and *Taq* I endonucleases, allowed the identification of *P. ginseng* and its differentiation from *P. quinquefolium*. *P. ginseng* was detected in 9 out 16 samples tested which, according to the declaration on the labels, contained the drug. Negative results were obtained for products containing the dried extract of the drug. A comparison of the results acquired using the molecular techniques with those using HPLC is also reported.

Keywords: *Panax ginseng*, *P. quinquefolium*, Molecular identification technique, Ginseng commercial products.

Molecular biology constitutes a new frontier for phytochemical analysis, allowing the improvement of previous knowledge, as well as the acquisition of new data. Recently, molecular techniques have been successfully utilised in order to validate plant drugs, overcoming the limitations of traditional analyses [1-4]. In the present study, PCR and RFLP were used in order to authenticate *Panax ginseng* C.A. Meyer (Korean ginseng) in different vegetable forms of ginseng commercial products and to differentiate it from other *Panax* species and from some of their adulterants. A comparison with HPLC identification results was also made.

There is an ongoing question over the labelling of herbal products as “Ginseng”. Currently the word “Ginseng” is used to sell a variety of herbs associated with certain claimed therapeutic properties (Table 1). This can be confusing as they neither contain the

same constituents nor display the same biochemical properties.

The quality of ginseng commercial products influences their effectiveness and safety of use and depends on the employed raw materials. The most active constituents of *P. ginseng* are steroidal saponins, called ginsenosides. So far 22 ginsenosides have been isolated and characterised, based on triterpene aglycone moieties with dammarane and oleanane structures and on the sugar unit sequences. In the monograph entitled “Ginseng” [5] the European Pharmacopoeia (Ph. Eur.) reports only the whole or cut dried root of *P. ginseng*, that must contain ginsenoside Rf and not less than 0.40% of combined ginsenosides Rg1 and Rb1, calculated with reference to the dried drug. This is, therefore, the only true ginseng (Korean ginseng).

Since the 1990s differentiation and research on the various species of *Panax* have been reported and the

⁺Part 1 in the Series: “Molecular identification of herbal drugs”

use of TLC, GLC and HPLC, as well as chemical techniques have allowed efficient separation and isolation of ginsenosides [6-9]. However, as reported in the Ph. Eur. [5], TLC and HPLC are the official tools to detect the presence of ginsenosides in root commercial samples. These methods need time (two working days), and a large quantity of plant material, as well as reference standards, that are often difficult to obtain.

Table 1: Main recoverable species in ginseng commercial products other than the European Pharmacopoeia species.

Botanical Name	Common names
<i>Panax ginseng</i> C.A. Meyer (Araliaceae) Ph. Eur. species	Korean ginseng Asian ginseng Chinese ginseng Ren shen Panax schinseng Jiln ginseng
<i>Panax quinquefolium</i> Linn	American ginseng
<i>Panax notoginseng</i> Burkill	San-chi ginseng
<i>Panax pseudoginseng</i> N. Wallich	Himalayan ginseng, Tien-chi ginseng
<i>Panax japonicus</i> or <i>P. japonicum</i> C.A. Meyer	Japanese ginseng, Ginseng bamboo
<i>Panax trifolium</i> Linn	Dwarf ginseng
<i>Panax zingiberensis</i> C.Y. Wu & Feng	Ginger ginseng
<i>Panax stipuleanatus</i> Tsai & Feng	Pingbiann ginseng
<i>Panax vietnamensis</i> Ha Thi Dung & I.V. Grushvitskii	Vietnamese ginseng
Other genera and families	
<i>Eleutherococcus senticosus</i> Maxim (Araliaceae)	Siberian ginseng, Wujia
<i>Echinopanax horridus</i> Decne & Planch. (Araliaceae)	Alaskan ginseng, Devil's club
<i>Aralia nudicaulis</i> Blume (Araliaceae)	Wild ginseng, Salsaparilla
<i>Rumex hymenosepalus</i> J. Torrey (Polygonaceae)	Red Desert ginseng
<i>Pfaffia paniculata</i> Kuntze (Amaranthaceae)	Brazilian ginseng, Suma
<i>Pseudostellaria heterophylla</i> Pax (Caryophyllaceae)	Sometimes used as a ginseng substitute
<i>Caulophyllum thalictroides</i> Regel (Berberidaceae)	Yellow or Blue ginseng, Blue cohosh
<i>Triosteum perfoliatum</i> Linn. (Caprifoliaceae)	Fever root, sometimes called Ginseng
<i>Codonopsis tangshen</i> Oliver (Campanulaceae)	Sometimes used as a ginseng substitute
<i>Lepidium meyenii</i> Walp. (Cruciferae)	Maca, Peruvian ginseng Andean ginseng
<i>Withania somnifera</i> Dun. (Solanaceae)	Indian ginseng Ashwaganda

The objective of the present study was to develop and standardise a reliable and easy molecular method for authentication of *P. ginseng* in the different forms of commercial ginseng products and to compare these results with HPLC identification made on the same samples [10].

Useful amounts of DNA were extracted from all the considered samples (1-19). PCR amplification was made using 18df/28ccr primers. The DNA amplified products, digested with the endonucleases *Hinf*I, *Taq*I and *Sau* 3A1, gave fragments of 170 bp; 230 bp and

280 bp; and 120 bp and 580 bp, respectively, characteristic for *P. ginseng*. Similarly, fragments of 60 bp and 100 bp; and 106 bp, 170 bp and 260 bp, were obtained for *P. quinquefolium* (American ginseng) with the use of *Hinf*I and *Taq*I, respectively (Figure 1).




Figure 1: Agarose gel electrophoresis of *Hinf*I, *Sau* 3A1 and *Taq*I restriction fragments obtained from amplified amplicons using the 18df/28ccr primer pair, specific for conserved region 18S-28S of *Panax* species.

For a total of 486 determinations (19 samples, 3 repetitions, 3 extraction protocols, following PCR and restriction with 3 endonucleases), the molecular analysis confirmed the presence of *Panax* species in 12 out of 19 samples tested, as shown in Table 2. Three of the positive samples, named PQ, PQT, PQP (6, 11, 12), were commercial ginseng products containing *P. quinquefolium* in the form of dried body root, dried root tails and dried root prongs, respectively. Among the samples labelled PG (1-5, 7-10, 13-19), the presence of *P. ginseng* was confirmed, as reported on the label, in 9 out of 16 samples, but not in three dried body root products (4, 5, 18) or in four dried extract samples (15-17, 19). The presence of adulterants such as *Mirabilis jalapa* L. and *Phytolacca acinosa* Roxb could be excluded in the considered samples since no specific *Sau* 3A1 digestion fragments [11] for either plant were visualised in agarose gel.

Table 2: DNA identification (PCR and RFLP) of *Panax ginseng* and HPLC detection of ginsenoside Rf in ginseng commercial products.

Sample		DNA identification				HPLC
		Body root	Root tails	Root prongs	Dried extracts	Ginsenoside Rf
1	PGB	+				+
2	PGR	+				+
3	PGB02	+				+
4	PGBH1	–				–
5	PGBH2	–				–
6	PQ	+				–
7	PGBR		+			+
8	PGT99		+			–
9	PGTO3		+			+
10	PGTO4		+			+
11	PQT		+			–
12	PQP			+		–
13	PGRB	+				–
14	PGBDSPR	+				+
15	PGPHRB				–	–
16	PGNGLC				–	–
17	PGRKPS				–	+
18	PGNST	–				–
19	PGext				–	+

PG: samples labelled as *P. ginseng*.PQ: samples labelled as *P. quinquefolium*.

1-12: raw materials; 13-18: commercial preparations as capsules or tablets;

19: laboratory hydromethanoholic extract.

HPLC analysis [10], conducted as reported in the Ph. Eur. [5], revealed the presence of all ginsenosides used as reference standards. As required by the Ph. Eur. monograph [5], ginsenosides Rg1 and Rb1 were identified in all tested samples and Rf, characteristic of *P. ginseng*, was detected in only 9 out of the 16 PG samples, which claimed to be based on *P. ginseng*. In particular, ginsenoside Rf was absent from the three PQ samples (6, 11, 12), as expected for *P. quinquefolium* products, and from seven PG samples (4, 5, 8, 13, 15, 16, 18).

The PCR and RFLP results were in accordance with the HPLC data (presence of ginsenoside Rf, characteristic of *P. ginseng*) for the majority of the tested commercial products and in accordance with the species, *P. ginseng*, declared on their labels. PCR and RFLP/HPLC afforded negative results for five products, in the form of dried body root (4, 5, 18) and of dried extract (15, 16), which excluded the presence of *P. ginseng*, although this was declared on the label. On the other hand, discordance between the molecular results and HPLC data was noted when the preparation was based on dried extracts (17, 19).

Molecular methods have been used to unequivocally allow the authentication to species level of the genus *Panax* and the results were not affected by the nature of the drug. Compared with other methods that detect genome-wise polymorphism simultaneously, such RAPD [12], AP-PCR and AFLP [13], the method applied in this research, based on PCR followed by

RFLP, is more reliable for large scale screening of commercial products, is rapid (one working day), and the results are easily readable. However, the procedure failed when the commercial products were dried extracts. A sample of reference *P. ginseng* was processed as described in the European Pharmacopoeia monograph [5], by boiling the root powder in 50% (v/v) aqueous methanol for 1h, to obtain a dried extract (19). This prepared extract resulted in a negative result in the molecular analytical procedure, as expected, because DNA molecules are not soluble in the hydromethanolic solvent, whereas the extract gave a positive for Rf on HPLC examination.

Experimental

The analysed commercial ginseng products as raw materials in the form of body root, root tails and root prongs, and as capsules and tablets containing also dried extract were obtained from national health care stores. To protect the Manufacturers' identities the sample sources were labelled as reported in Table 2.

An AB GeneAmp PCR System 9700 thermal cycler was used for the PCR analysis. *P. ginseng* and *P. quinquefolium* dried roots, used as references for the molecular analysis, were kindly provided by the Department of Plant Biology of the University of La Sapienza, Rome, Italy. Chromatography was performed on a Waters chromatographic system equipped with a Waters 600 MS multisolvent delivery system and a Waters 717 Auto sampler. A Waters 996 Photodiode Array Detector was used to monitor the eluates at 203 nm. The chromatographic data were analysed using a Waters Millennium Software version 3.2. Chromatography was performed at room temperature (25° C). Ginsenosides Rg1, Rb1, Rb2, and Rc–Rf, purchased from Extrasynthese, France, were used as reference standards.

Molecular analysis: DNA was extracted from 50-100 mg of each sample using an Invitrogen Easy-DNA Kit [14] and two other molecular protocols [15, 16] in order to compare their effectiveness in the extraction of useful amounts of DNA for molecular analysis. Body roots were previously treated with liquid nitrogen, while the other samples were used directly in the DNA procedures following the Manufacturers' instructions. Sometimes, it was necessary to precipitate with isopropyl alcohol, rinse with 70% ethanol, resuspend in 10 mM TE (Tris-HCl, pH 8.0, 1 mM EDTA) and precipitate a second

time in the presence of 0.3 M sodium acetate and 2 volumes of ethanol. The final pellet, after a second rinse in 70% ethanol, was resuspended in sterile distilled water (50 μ L). The PCR amplification was performed on all ginseng DNA samples using oligonucleotide primers 18df/28ccr. This primer pair amplifies the conserved region, 18S-28S, including ITS1 and ITS2, highly variable regions for *Panax* species [17]. Amplification reactions were performed with reaction mixtures and with reaction conditions previously reported [11], using DNA *Taq* polymerase W1 (Invitrogen, Italy). For the RFLP analysis, amplified 18S rDNA fragments (11 μ L aliquots) were separately digested in a final volume of 20 μ L at 37°C for 3h and 65°C for 16h with 1.5-2 Units for each of the following endonucleases: *Hinf*I, *Sau* 3A1, *Taq* I (New England BioLabs, UK). Restriction fragments were analysed by electrophoresis in 1% agarose gels buffered in 0.5 X TBE [TBE buffer: 90

mM Tris (hydroxymethyl)-aminomethane, 90 mM boric acid, 3 mM ethylene-diaminetetraacetate Na salt, pH 8.3] and visualised by UV light after staining with ethidium bromide. The size marker was 50 bp ladder (Invitrogen, Italy).

Chemical analysis: Analysis of the principal active constituents was performed on all available samples by the HPLC method described in the “Ginseng” monograph of the Ph. Eur. [5], using a 5 μ m (25 cm x 4.6 mm) Kromasil KR100-5NH2 E6170 column, a mobile phase filtered on an Alltech nylon membrane 47 mm, 0.45 μ m, and degassed by a Waters in line degasser, at a flow rate of 1 mL/min.

Acknowledgments - The authors thank Professor Marcello Nicoletti (University “La Sapienza”, Rome, Italy) for his suggestions to improve the paper.

References

- [1] Linacre A, Thorpe J. (1998) Detection and identification of cannabis by DNA. *Forensic Science International*, **91**, 71-76.
- [2] Lau DT, Shaw PC, Wang J, But PPH. (2001) Authentication of medicinal *Dendrobium* species by the internal transcribed spacer of ribosomal DNA. *Planta Medica*, **67**, 456-460.
- [3] Miller Coyle H, Palmbach T, Juliano N, Ladd C, Lee HC. (2003) An overview of DNA methods for the identification and individualization of marijuana. *Croatian Medical Journal*, **44**, 315-321.
- [4] Johnson EL, Saunders JA, Mischke S, Helling CS, Emche SD. (2003) Identification of *Erythroxylum* taxa by AFLP DNA analysis. *Phytochemistry*, **64**, 187-197.
- [5] European Pharmacopoeia (2005) Ginseng. 5th Edition: monograph 1523.
- [6] Wang X, Sakuma T, Asafu-adjaye E, Shiu GK. (1999) Determination of ginsenosides in plant extracts from *Panax ginseng* and *Panax quinquefolius* L. by LC/MS/MS. *Analytical Chemistry*, **71**, 1579-1584.
- [7] Chan TW, But PPH, Cheng SW, Kwok IM, Lau FW, Xu HX. (2000) Differentiation and authentication of *Panax ginseng*, *Panax quinquefolius*, and ginseng products by using HPLC/MS. *Analytical Chemistry*, **72**, 1281-1287.
- [8] Zhang Haijiang, Wu Yongjiang, Cheng Yiyu (2003) Analysis of 'SHENMAI' injection by HPLC/MS/MS. *Journal of Pharmaceutical and Biomedical Analysis*, **31**, 175-183.
- [9] Yoon SR, Nah JJ, Kim SK, Kim SC, Nam KY, Jung DW, Nah SY. (1998) Determination of ginsenosides Rf and Rg2 from *Panax ginseng* using enzyme immunoassay. *Chemical and Pharmaceutical Bulletin*, **46**, 1144-1147.
- [10] Palazzino G, Gallinella B, Turchetto L, Gallo FR, Federici E, Iurilli R, Zanitti L. (2004) Ginseng: quali-quantitative analysis in Italian commercial products. *FITOMED 2004*. 1st Intersociety Congress on Medicinal Plants, 16-19 September 2004, Trieste, Italy.
- [11] Ngan F, Shaw T, But P, Wang J. (1999) Molecular authentication of *Panax* species. *Phytochemistry*, **50**, 787-791.
- [12] Shaw PC, But PPH. (1995) Authentication of *Panax* species and their adulterants by random-primed polymerase chain reaction. *Planta Medica*, **61**, 466-469.
- [13] Ha WY, Shaw PC, Liu J, Yau, CFF, Wang J. (2002) Authentication of *Panax ginseng* and *Panax quinquefolius* using amplified fragment length polymorphism (AFLP) and directed amplification of minisatellite region DNA (DAMD). *Journal of Agricultural and Food Chemistry*, **50**, 1871-1875.
- [14] Invitrogen (2003) Easy-DNA kit instruction for genomic DNA isolation. *Invitrogen Life Technologies*, Manual 25-0056 Version F.
- [15] Del Serrone P, Minucci C, Barba M, Conti M, Boccardo G. (1995) Ottimizzazione della diagnosi di fitoplasmi in vite. *Petria*, **5**, 161-170.
- [16] Kuske CR, Banton KI, Adorada DL, Stark PC, Hill KK, Jackson PJ. (1998) Small-scale DNA sample preparation method for field PCR detection of microbial cells and spores in Soil. *Applied and Environmental Microbiology*, **64**, 2463-2472.
- [17] Wen J, Zimmer EA. (1996) Phylogeny and biogeography of *Panax ginseng* L. (the *Ginseng* genus, Araliaceae): Interferences from ITS sequences of nuclear ribosomal DNA. *Molecular Phylogenetics and Evolution*, **6**, 167-177.

Lipoxygenase Inhibitory Activity of Boropinic Acid, Active Principle of *Boronia pinnata*

Massimo Curini^{a*}, Francesco Epifano^b, Salvatore Genovese^a, Luigi Menghini^b, Donata Ricci^c, Daniele Fraternali^c, Laura Giamperi^c, Anahi Bucchini^c and Emanuele Bellacchio^d

^aDipartimento di Chimica e Tecnologia del Farmaco, Sezione di Chimica Organica, Via del Liceo, 06123 Perugia, Italy

^bDipartimento di Scienze del Farmaco, Via dei Vestini 31, 66013 Chieti Scalo, Italy

^cIstituto di Botanica e Orto Botanico, Via Bramante 28, 61029 Urbino, Italy

^dCSS Hospital, IRCCS, San Giovanni Rotondo and CSS Mendel Institute, Viale Regina Margherita 261, 00198 Roma, Italy

curmax@unipg.it

Received: March 21st, 2006; Accepted: April 24th, 2006

Dedicated to the memory of Professor Ivano Morelli.


Boropinic acid and other natural prenyloxy-cinnamic and benzoic acids were easily synthesized in high yield by a two-step sequence from the corresponding *p*-hydroxy aromatic acids and were assayed for radical scavenging activity using the DPPH test and for inhibition of enzymatic lipid peroxidation mediated by soybean 5-lipoxygenase. Compared to other acids and to known antioxidant compounds like BHT, Trolox and ascorbic acid, boropinic acid was far more active in the lipoxygenase test ($IC_{50} = 7.6$ ng/mL, $p < 0.05$). The recorded inhibition value suggested that boropinic acid acted as an enzyme inhibitor rather than a mere radical or peroxide scavenger. This hypothesis was confirmed by studying the interaction between boropinic acid and soybean 5-lipoxygenase by molecular modelling techniques.

Keywords: anti-inflammatory activity, antioxidant activity, boropinic acid, lipoxygenase, prenyloxy acids.

Secondary metabolites of phenylpropanoic acid biosynthetic origin containing sesquiterpenyl, monoterpenyl and isopentenyl chains attached to a phenol group represent quite a rare group of natural products. Some of these compounds, including coumarins [1], anthraquinones [2], xanthenes [3], flavonoids [4] and carboxylic acids, have been recently studied chemically and pharmacologically. Among the latter, cinnamic and benzoic acids have been shown recently to have valuable biological properties [5].


To the best of our knowledge, only five prenyloxy-phenylpropanoic acids have been reported from natural sources: 3-(4'-geranyloxyphenyl)-2-*trans* propenoic acid (1), 3-(4'-geranyloxy-3'-methoxyphenyl)-2-*trans* propenoic acid (2), isolated from

Acronychia baueri Schott [6], boropinic acid (3), extracted from *Boronia pinnata* Sm. [7], valencic acid (4), isolated from *Citrus sinensis* L. and *Aegle marmelos* [8], and 4-isopentenyl-3-methoxy benzoic acid (5), isolated as a methyl ester from the liverwort *Trichocolea lanata* (Ehrh.) Dumm. [9]. The aim of this study was to synthesize these natural prenyloxy-carboxylic acids and to test their antioxidant activity.



- 1 $R^1 = -CH=CH-COOH$, $R^2 = -H$, $R^3 = \text{geranyl}$
2 $R^1 = -CH=CH-COOH$, $R^2 = -OCH_3$, $R^3 = \text{geranyl}$
3 $R^1 = -CH=CH-COOH$, $R^2 = -OCH_3$, $R^3 = \text{isopentenyl}$
4 $R^1 = -COOH$, $R^2 = -H$, $R^3 = \text{isopentenyl}$
5 $R^1 = -COOH$, $R^2 = -OCH_3$, $R^3 = \text{isopentenyl}$

The synthesis of compounds **1**, **3**, **4** and **5** was accomplished following an environmentally friendly route similar to that reported for the synthesis of compound **2** [5]. Compound **1** was obtained in 97% overall yield starting from commercially available *p*-coumaric acid that was first converted into its methyl ester by refluxing in MeOH catalyzed by concentrated H₂SO₄, then alkylated with geranyl bromide and hydrolyzed in a basic medium (Scheme 1).



Scheme 1: Reagents and conditions: a) MeOH, conc. H₂SO₄ (cat.), reflux, 12 h; b) geranyl bromide, K₂CO₃, acetone, reflux, 2h; c) NaOH 2N, 70°C, 1h

Compounds **3** and **5** were obtained, using the same reaction conditions as above, in 96% and 98% yield from ferulic acid and vanillic acid, respectively, while compound **4** was synthesized in 99% yield by a one-pot alkylation-basic hydrolysis from commercially available methyl *p*-hydroxy benzoate and employing, in all cases, 4-bromo-2-methyl-2-butene as alkylating agent.

Table 1: DPPH radical scavenging activity of prenyloxy-carboxylic acid.

Compound	IC ₅₀ , μmol/mL ^a
1	0.065 ± 0.0060
2	0.011 ± 0.0011
3	0.011 ± 0.0008
4	0.011 ± 0.0008
5	0.052 ± 0.0048
Ascorbic acid	6.24 × 10 ⁻⁴ ± 3.97 × 10 ⁻⁵
BHT	5.24 × 10 ⁻⁴ ± 4.87 × 10 ⁻⁵
Trolox	0.30 × 10 ⁻⁴ ± 0.39 × 10 ⁻⁵

^a p < 0.05 using Student's *t* test

Compounds **1-5** were first assayed to evaluate their radical scavenging activity by the DPPH test [10], using Trolox (6-hydroxy-2,5,7,8-tetramethyl-chroman-2-carboxylic acid), BHT (butyl hydroxy toluene) and ascorbic acid as positive controls. Results are reported in Table 1.

As shown in Table 1, no acid exhibited an appreciable radical scavenging activity compared to the controls, particularly those having a free phenolic

hydroxyl group like Trolox and BHT. So, the lack of any significant radical scavenging ability of compounds **1-5** may be due to the alkylation of the phenol moiety with either a geranyl or isopentenyl group.

We then evaluated the inhibition of polyunsaturated fatty acid (PUFA) peroxidation catalyzed by soybean 5-lipoxygenase (5-LOX) [11]. Assessing the inhibitory effect of a chemical on this enzyme is noteworthy, as lipoxygenases are nowadays recognized as playing a major role in cancer cell growth, metastasis, invasiveness, cell survival and induction of tumor necrosis factor (TNF) [12,13]. More particularly, it has been observed that the inhibition of the 5-LOX pathway has a chemopreventative effect in lung carcinogenesis, prevents the biological activation of different types of carcinogens, decreases cell proliferation, and induces apoptosis [14-16]. We used soybean 5-LOX in our study as, despite differences in the number of aminoacids between plant and mammalian LOXs, it has been reported that these proteins are similar in topology, with high similarities in the respective active sites and mechanism of catalysis [17].

Table 2: Inhibition of 5-LOX-mediated PUFA peroxidation by prenyloxy-carboxylic acid.

Compound	IC ₅₀ , μmol/mL ^a
1	0.006 ± 0.0005
2	0.262 ± 0.0220
3	2.89 × 10 ⁻⁵ ± 2.62 × 10 ⁻⁶
4	> 100
5	> 100
Ascorbic acid	0.105 ± 0.0072
BHT	0.023 ± 0.0052
Trolox	0.047 ± 0.0048

^a p < 0.05 using Student's *t* test

Results on the inhibition of lipoxygenase mediated lipid peroxidation are reported in Table 2. Trolox, BHT and ascorbic acid were used as positive controls. As reported in Table 2, the pattern of antioxidative activity of **1-5** is similar to that recorded for the radical scavenging activity, with the notable exception of boropinic acid (**3**). This cinnamic acid derivative is far more active, not only in respect to compounds **1**, **2**, **4** and **5**, but also when compared to the positive controls. The value recorded for boropinic acid suggested that it did not act as a mere reducing agent like Trolox, BHT and ascorbic acid, but more likely as an effective 5-LOX inhibitor. To rationalize tentatively the inhibitory mechanism observed for boropinic acid and the lack of efficacy




Figure 1: Comparative positions of boropinic acid (green sticks) and DHB (white sticks) in their complexes with LOX after protein backbone alignment (for simplicity, bond orders, hydrogen atoms and the protein are not shown).




Figure 2: Modelled lipoxigenase / boropinic acid complex. The protein backbone is represented as green ribbons. Boropinic acid is represented as stick and dots. Amino acid residues with atoms within 5 Å from boropinic acid atoms are represented as sticks (amino acids with hydrophobic side chains are in blue and the remaining amino acids are in magenta). For simplicity, hydrogen atoms of the ligand and the displayed residues are not shown. It can be noted that the prenyloxy group can be inserted in a cluster of hydrophobic amino acids. In this position, boropinic acid is potentially involved in interactions with two polar groups similar to the case of the DHB ligand (observed in the PDB structure 1n8q). The groups of the protein implied in the polar interactions are the carboxy-terminal group, Ile 857, and the amide side chain of Gln 514 (both groups are highlighted by the orange mesh).

of the other four natural carboxylic acids we have inferred a possible lipoxigenase/ligand docking by comparative modelling.

Among ligands reported to have an inhibitory effect on soybean lipoxigenase [17], 3,4-dihydroxybenzoic acid (DHB) showed the most significant structural similarity with boropinic acid and we have adopted its geometry of binding to the same cavity in the

lipoxigenase protein as a model for the docking of boropinic acid. We have initially assigned to the phenyl ring of boropinic acid the same position and orientation known for the phenyl ring of DHB in the lipoxigenase/DHB complex (PDB structure 1n8q). In order to minimize collisions with the protein atoms, boropinic acid was subjected to structure optimization. The final docking of boropinic acid (**3**) required minor adjustments with respect to its initial position of insertion in the protein cavity. In Figure 1 are shown the mutual positions of boropinic acid and the DHB, as viewable if the two respective theoretical and experimental complexes with lipoxigenase were superimposed by aligning the protein backbones. The interactions of **3** and DHB should involve at least some common amino acid residues in the protein cavity. A peculiar feature of the modelled lipoxigenase/boropinic acid complex (Figure 2) is the possibility for the apolar atoms of the isopentenyl group to be oriented and enter in van der Waals contact with a cluster of hydrophobic amino acids. Since this additional interaction might contribute to the enhancement of the complex stability, it seems that the loss of activity of lipoxigenase in the presence of boropinic acid could be the result of enzyme inhibition as a consequence of stable ligand docking in the active site, while similar experiments performed for all the other acids showed that the geranyloxy-cinnamic ones are too big to fit the active site while benzoic acids are too small to get more stable interaction with the enzyme.

In summary, the results described herein provide a new high yielding and environmentally friendly synthetic route to prenyloxy aromatic acids and indicate boropinic acid as a lead compound of a novel class of selective LOXs inhibitors.

Experimental

General experimental: For the synthesis of compounds **1-5** the same general procedure as that reported previously was followed [5].

3-(4'-Geranyloxyphenyl)-2-*trans* propenoic acid (**1**)

Yield: 97%.

MP: 156-157°C.

IR (KBr): 3550, 1690 cm⁻¹.

¹H NMR: [6]

¹³C NMR (100 MHz CDCl₃): 16.1 (CH₃), 17.5 (CH₃), 25.6 (CH₃), 26.2 (CH₂), 39.4 (CH₂), 64.9 (CH₂), 115.3 (CH), 117.6 (CH), 119.8 (CH), 123.8 (CH),

128.3 (C), 129.3 (CH), 131.4 (C), 141.6 (C), 144.2 (CH), 157.7 (C), and 168.9 (C).

Anal. Calcd for $C_{19}H_{24}O_3$: C, 75.97; H, 8.05; O, 15.98. Found C, 75.96; H, 8.07; O, 15.99.

3-(4'-Geranyloxy-3'-methoxyphenyl)-2-*trans* propenoic acid (2)

Yield: 96%.

Analytical data are in full agreement with those reported in the literature [5]

Boropinic acid (3)

Yield: 96%.

Analytical data are in full agreement with those reported in the literature [7]

Valencic acid (4)

Yield: 99%.

MP: 131-132°C.

IR: [21]

1H NMR: [19]

^{13}C NMR (100 MHz $CDCl_3$): 18.7 (CH_3), 26.2 (CH_3), 66.5 (CH_2), 116.0 (CH), 120.9 (CH), 122.1 (C), 132.2 (CH), 139.1 (C), 162.7 (C), 170.5 (C).

Anal. Calcd for $C_{12}H_{14}O_3$: C, 69.89; H, 6.84; O, 23.27. Found C, 69.88; H, 6.82; O, 23.26.

4-Isopentenyl-3-methoxy benzoic acid (5)

Yield: 98%.

MP: 141-142°C.

IR (KBr): 3600, 1695 cm^{-1} .

1H NMR (400 MHz $CDCl_3$): 1.70 (3H, s, CH_3), 1.74 (3H, s, CH_3), 3.83 (3H, s, OCH_3), 4.55-4.61 (2H, m, OCH_2), 5.72-5.77 (1H, m, olefinic proton), and 7.01-7.72 (3H, m, aromatic protons)

^{13}C NMR (100 MHz $CDCl_3$): 18.0 (CH_3), 27.0 (CH_3), 55.9 (CH_3), 66.2 (CH_2), 115.4 (CH), 116.4 (CH), 119.9 (CH), 124.6 (CH), 125.8 (C), 138.0 (C), 147.3 (C), 150.7 (C), and 168.8 (C).

Anal. Calcd for $C_{13}H_{16}O_4$: C, 66.09; H, 6.83; O, 27.09. Found C, 66.08; H, 6.82; O, 27.07.

DPPH test: Radical scavenging activity was determined by a spectrophotometric method based on the reduction of an ethanol solution of 1,1-diphenyl-2-picrylhydrazyl (DPPH). Tests were carried out in triplicate. Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid), BHT (butylated hydroxytoluene) and ascorbic acid were used as positive controls and purchased from SIGMA.

Inhibition of lipid peroxide formation test: Lipid peroxidation inhibitory activity was evaluated using the 5-lipoxygenase test in conjunction with positive controls. The activity of the enzyme was assayed spectrophotometrically according to the method of Holman, as modified by Sud'ina *et al.* [11]. The assay mixture (1 mL) contained: 10 mM linoleic acid, the sample (or the same quantity of solvent as reference) and 50 mM sodium phosphate, pH 6.8. This mixture was maintained at 20°C for 20 min. Subsequently, 0.18 $\mu g\ mL^{-1}$ commercial 5-lipoxygenase was added to the mixture and the formation of hydroperoxides from linoleic acid was observed spectrophotometrically at 235 nm at 20 °C.

Docking experiments: Molecular Dynamics (MD) of lipoxygenase (PDB structure 1n8q) was performed with the program NAMD (v2.5) [18] after having removed all heteroatoms from the structure (ligands and crystallization water molecules). The simulation was carried out in explicit solvent under periodic boundary conditions using the CHARMM 27 parameter set. The system was minimized for 500 steps (1 fs per step) and the MD simulation was carried out for 500000 steps (500 ps) at a temperature of 310 K. Molecular representations were made with the program PyMol.

Acknowledgments - Authors from Perugia and Chieti would like to acknowledge financial support from MIUR National Project "Sviluppo di processi sintetici ecocompatibili nella sintesi organica" COFIN 2004. Authors from Urbino would like to acknowledge financial support from Regione Marche, Progetto CIPE 17/2003.

References

- [1] Curini M, Cravotto G, Epifano F, Giannone G. (2006) Chemistry and biological activity of natural and synthetic prenyloxycoumarins. *Current Medicinal Chemistry*, **13**, 199-222.
- [2] Pattanaprateeb P, Ruangrunsi N, Cordell GA. (2005) Cytotoxic constituents from *Cratoxylum arborescens*. *Planta Medica*, **71**, 181-183.
- [3] Bilia AR, Yusuf AW, Braca A, Keita A, Morelli I. (2000) New prenylated anthraquinones and xanthenes from *Vismia guianensis*. *Journal of Natural Products*, **63**, 16-21.

- [4] Magalhães AF, Tozzi A, Magalhães EG, Blanco IS, Soriano MC. (2004) Flavonoids from *Lonchocarpus muehlbergianus*. *Annales de Academia Brasileira de Ciencias*, **76**, 651-661.
- [5] Curini M, Epifano F, Genovese G. (2005) Synthesis of a novel prodrug of 3-(4'-geranyloxy-3'-methoxyphenyl)-2-*trans* propenoic acid. *Bioorganic and Medicinal Chemistry Letters*, **15**, 5049-5052, and references cited herein.
- [6] Prager RH, Thregold HM. (1966) Some neutral constituents of *Acronychia baueri*. *Australian Journal of Chemistry*, **19**, 451-452.
- [7] Ito C, Itoigawa M, Otsuka T, Tokuda H, Nishino H, Furukawa H. (2000) Constituents of *Boronia pinnata*. *Journal of Natural Products*, **63**, 1344-1348.
- [8] Ali MS, Pervez MK. (2004) Marmenol a 7-geranyloxycoumarin from the leaves of *Aegle marmelos* Corr. *Natural Product Research*, **18**, 141-146.
- [9] Perry NB, Foster LM, Lorimer SD, May BC, Weavers RT. (1996) Isoprenyl phenyl ethers from liverworts of the genus *Trichocolea*: cytotoxic activity, structural corrections and synthesis. *Journal of Natural Products*, **59**, 729-733.
- [10] Santosh Kumar S, Priyadarsini KI, Sains KB. (2002) Free radical scavenging activity of vanillin and *o*-vanillin using 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical. *Redox Reports*, **7**, 35-40.
- [11] Sud'ina GF, Mirzoeva OK, Pushkareva MA, Korshunova GA, Sumbatyan NV, Vafolomeev SD. (1993). Caffeic acid phenethyl ester as a lipoxygenase inhibitor with antioxidant properties. *Federation of European Biochemical Societies*, **329**, 21-24.
- [12] Chan MM. (1995) Inhibition of tumor necrosis factor by curcumin, a phytochemical. *Biochemical Pharmacology*, **49**, 1551-1556.
- [13] Hallahan DE, Virudachalam S, Kufe, DW, Weichselbaum RR. (1966) Ketoconazole attenuates radiation induction of tumor necrosis factor. *International Journal of Radiation in Oncology Biology and Physics*, **29**, 777-780.
- [14] Rioux N, Castonguay A. (1998) Inhibitors of lipoxygenase: a new class of cancer chemopreventive agents. *Carcinogenesis*, **19**, 1393-1400.
- [15] Moody TV, Leyton J, Martinez A, Hong S, Malkinson A, Mulshine JL. (1998) Lipoxygenase inhibitors prevent lung carcinogenesis and inhibit non-small cell lung cancer growth. *Experimental Lung Research*, **24**, 617-628.
- [16] Ghosh J, Myers CE. (1998) Inhibition of tumor arachidonate 5-lipoxygenase triggers massive apoptosis in human prostate cancer cell. *Proceedings of the Natural Academy of Science USA*, **95**, 13182-13187.
- [17] Skrzypczak AK, Jankun E, Zhou K, Jankun J. (2003) Inhibition of lipoxygenase by (-) epigallocatechin gallate: X-ray analysis at 2.1 Å reveals degradation of EGCG and shows soybean LOX-3 complex with EGC instead. *International Journal of Molecular Medicine*, **12**, 415-422.
- [18] Phillips JC, Braun R, Wang W, Gumbart J, Tajkhorshid E, Villa E, Chipot C, Skeel RD, Kale L, Schulten K. (2005) Scalable molecular dynamics with NAMD. *Journal of Computational Chemistry*, **26**, 1781-1802.
- [19] Takemura Y, Kawaguchi H, Maki S, Juichi M, Omura M, Ito C, Furukawa H. (1996) Constituents of domestic Citrus plants. Part XXVI. Studies on the constituents of Yalaha. Structures of a new acridone alkaloid and two new coumarins. *Chemical & Pharmaceutical Bulletin*, **44**, 804-809.

A Convenient Synthesis of 5'-Iodoresiniferatoxin (I-RTX)

Abdellah Ech-Chahad,^{a,b} Lahboub Bouyazza^a and Giovanni Appendino^{b,*}^aUniversité Hassan 1er, Faculté de Sciences et Techniques, BP 577, 2600 Settat, Morocco^bDipartimento di Scienze Chimiche, Alimentari, Farmaceutiche e Farmacologiche
Università del Piemonte Orientale, Via Bovio 6, 28100 Novara, Italy

appendino@pharm.unipmn.it

Received: June 16th, 2006; Accepted: July 5th, 2006


Dedicated to the memory of Professor Ivano Morelli.

Starting from resiniferonol orthophenylacetate (ROPA, **2**) and commercial 5-iodovanillin (**5a**), a convenient synthesis of the ultrapotent vanilloid antagonist 5'-iodoresiniferatoxin (**1a**) was achieved, overcoming the problems involved in the direct iodination of either resiniferatoxin (**1b**) or homovanillic acid (**3a**).

Keywords: Vanilloid antagonists, 5'-iodoresiniferatoxin, resiniferatoxin, TRPV1, resiniferonol orthophenylacetate.

The study of ion channels strongly depends on the availability of compounds that can either activate or inhibit their function with high selectivity and potency [1]. While there is no shortage of ligands for sodium-, potassium-, and calcium-channels [1], most channels of the TRP type still await de-orphanization in terms of small molecule activators and/or inhibitors [2]. A remarkable exception is TRPV1, the capsaicin receptor, for which a large number of ligands (vanilloids) are available [3]. Most TRPV1 activators are either natural products or compounds derived from (or inspired by) natural products. Conversely, vanilloid antagonists are mainly synthetic compounds that have emerged from the random screening of chemical libraries. Nevertheless, none of them approaches the potency of 5'-iodoresiniferatoxin (I-RTX, **1a**), a natural product-derived ligand. I-RTX inhibits TRPV1 activation with a one-digit nanomolar $K_{(i)}$, [4]. While impressive, this value probably even underestimates the actual potency of I-RTX, whose intracellular penetration is slow compared to the time frame of most assays for vanilloid activity [5]. I-RTX is not only important as a molecular probe, but also as a potential drug, and has been investigated, with impressive results, in animal models of antitussive [6] and analgesic [7] activity.

I-RTX was serendipitously discovered by Wahl while attempting to prepare a radioactive derivative of the ultrapotent vanilloid agonist resiniferatoxin (RTX, **1b**) [4]. The molecular bases for the reversal of activity induced by aromatic iodination *ortho* to the phenolic hydroxyl are unknown, but a similar observation was made with capsaicinoids for iodination at the carbons *ortho*- and *meta*- to the phenolic hydroxyl [8]. Remarkably, iodination of RTX *meta* to the phenolic hydroxyl generated instead a partial agonist (**1c**) [9].




Despite the relevance of I-RTX for pharmacological research, a convenient synthesis of this compound has not yet been reported. The original synthesis by Wahl [4] was improved by a Merck group [10], and is based on the iodination of RTX with the sodium iodide/chloramine T system. Since RTX is labile in these conditions, the reaction requires careful control, and must be quenched at incomplete conversion. [10]. After HPLC purification, I-RTX was eventually

obtained in *ca* 22% yield. Given the low yield of the reaction, its problematic scale-up, and the exorbitant price of RTX [11], this synthesis is unsuitable to produce the amounts of I-RTX needed to profile its bioactivity in *in vivo* experiments.

We reasoned that resiniferol orthophenylacetate (ROPA, **2**), the terpenoid core of RTX, would be a more convenient starting point for the synthesis of I-RTX, both in terms of availability of the starting material and purification of the final product. Thus, while RTX is a highly offensive compound, ROPA can be manipulated under normal laboratory conditions, and can be obtained relatively easily from the partially hydrolyzed latex of *Euphorbia resinifera* Berg., a household plant [12]. Conversely, the isolation of RTX from the native latex is difficult and hazardous due to its obnoxious properties and to the occurrence of irritant and tumor-promoting ingenol and deoxyphorbol esters that share the polarity and chromatographic behavior of RTX [12]. Finally, carrying out the iodination at the stage of a simple vanillyl derivative will also solve the problem of the instability of the terpenoid core of ROPA in the iodinating conditions. A similar strategy has been reported for the preparation of 6'-iodoresiniferatoxin (**1c**) [9], but, surprisingly, no attempt has been made so far to extend this strategy to its more important 5'-isomer.


The iodination of homovanillic acid (**3a**) was first investigated (Scheme 1). This compound and its esters have been reported to be poor substrates for aromatic iodination [9], and also, in our hands, complex mixtures were obtained with a variety of iodinating conditions. However, methyl homovanillate (**3b**) could be iodinated, albeit in poor yield, with the *N*-iodosuccinimide (NIS) - trifluoroacetic acid (TFA) protocol [13]. The compound obtained (**4a**) contained *ca.* 5-10% of an impurity, tentatively identified as the product of α , 5'-bis-iodination (**4b**) on the basis of MS and ^1H NMR spectroscopic evidence. Thus, a peak corresponding to the incorporation of two iodine atoms was observed in the MS, while the ^1H NMR spectrum showed two additional *meta*-coupled aromatic protons. This by-product could not be removed by either chromatography or crystallization. After hydrolysis and Mitsunobu esterification [14] with ROPA, I-RTX (**1a**) was obtained, still contaminated, however, with the corresponding bis-iodinated impurity. Since preparative HPLC could not afford a completely pure material, this approach, though simple, was abandoned, and an alternative

strategy based on the homologation of 5-iodovanillic acid was explored (Scheme 2).



Scheme 1: Iodination of methyl homovanillate (**3b**).

While homovanillic acid (**3a**) is expensive, 5-iodovanillin (**5a**) is cheap and commercially available in high purity [15]. After acetylation and oxidation, an acetylated carboxylic acid precursor for the one-carbon Arndt-Eisert homologation was obtained (**5c**). Reaction with oxalyl chloride and next with trimethylsilyldiazomethane [16] afforded the stable diazoketone **6**. The Wolf rearrangement of **6** in water with silver oxide gave mainly the corresponding acyloin **7**, while the reaction was successful after switching to the methanol-silver benzoate system. [17]. The resulting acetylated methyl ester was next hydrolyzed (LiOH, THF-water), affording 5'-iodohomovanillic acid (**4a**) as a crystalline compound in 43% yield overall from **5a** (Scheme 2).



Scheme 2: Synthesis of 5'-iodohomovanillic acid (**4a**) from commercial 5-iodovanillin (**5a**).

The final Mitsunobu coupling of **4a** and ROPA (**2**) could be carried out with crude ROPA (*ca.* 80%, HPLC) and the DIAD-TPP redox couple. After solvent removal, the residue was dissolved in toluene and cooled to remove the hydrazodicarboxylate-triphenylphosphine oxide crystalline adduct [18]. Further purification by gravity column chromatography on neutral alumina affording I-RTX (**1a**) as a colorless foam in a reproducible yield of 52% and a HPLC purity of *ca* 95%.

In conclusion, a convenient synthesis of I-RTX, an ultrapotent vanilloid antagonist, has been reported,

overcoming the problems posed by the iodination of RTX and homovanillic acid and filling an important gap in vanilloid research.

Experimental

Acetyl-5-iodovanillic acid (5c): To a solution of 5-iodovanillin acetate (**5b**, 1g, 3.12 mmol, prepared from commercial 5-iodovanillin (**5a**) by reaction with Ac₂O-pyridine) in acetone (10 mL), freshly prepared Jones reagent [19] was added (3 mL). After stirring overnight at room temperature, the reaction was worked up by concentration and filtration over Celite. The filtrate was then extracted with diethylether, and the organic phase was washed with brine. After drying and removal of the solvent, the residue was crystallized from diethylether to afford 800 mg (76%) of **5c** as a white powder.

MP: 199°C.

Rf: 0.37 (light petroleum -EtOAc 7:3).

IR (KBr): 3072, 1764, 1683, 1572, 1409, 1294, 1193, 1168, 1037 cm⁻¹.

¹H NMR (300 MHz, CDCl₃): 2.39 (3H, s, Ac), 3.89 (3H, s, OMe), 7.64 (1H, d, *J* = 1.5 Hz, Ar-H), 8.16 (1H, d, *J* = 1.5 Hz, Ar-H).

¹³C NMR (75 MHz, CDCl₃): 20.9 (CH₃), 56.5 (CH₃), 91.7 (C), 113.8 (CH), 129.0 (CH), 132.8 (C), 145.4 (C), 151.6 (C), 167.5 (C), 170.3 (C).

CI-EIMS: *m/z* [M+ H]⁺ 321 [C₁₀H₉IO₅ + H]⁺

α-Diazo-5-iodoacetovanillone acetate (6): To a cooled solution of **5c** (700 mg, 2.1 mmol) in dry CH₂Cl₂ (4 mL), oxalyl chloride (0.73 mL, 8.4 mmol, 4 mol. equiv.) and cat. DMF (0.20 mL) were added. After stirring for 1 h at 0°C and 90 min at room temperature, the reaction was worked up by evaporation, and the residue dissolved in THF-acetonitrile (1:1, 10 mL). After cooling to 0°C, TMSCHN₂ (2M in diethylether, 1.84 mL, 3.94 mmol, 1.9 mol. equiv.) was added. The brownish-colored reaction was stirred at 0°C for 30 h, and then quenched by the addition of 0.5 N acetic acid. The reaction was then worked up by the addition of satd NaHCO₃, and the organic phase was separated, washed with brine and evaporated. The residue was purified by gravity column chromatography on silica gel (25 g, light petroleum -EtOAc 8:2 as eluant) to afford 471 mg (66%) **6** as a yellowish powder.

MP: 143°C.

Rf: 0.32 (light petroleum -EtOAc 7:3).

IR (KBr): 3113, 3027, 2412, 2117, 1768, 1566, 1407, 1282, 1196, 1027, 903 cm⁻¹.

¹H NMR (300 MHz, CDCl₃): 2.23 (3H, s, Ac), 3.73 (3H, s, OMe), 5.71 (s, 1H), 7.29 (1H, d, *J* = 1.5 Hz, Ar-H), 7.49 (1H, d, *J* = 1.5 Hz, Ar-H).

¹³C NMR (75 MHz, CDCl₃): 20.9 (CH₃), 56.5 (CH₃), 91.7 (C), 110.8 (CH), 128.7 (CH), 136.6 (C), 144.3 (C), 151.9 (C), 167.6 (C), 184.0 (C), 225.1 (C).

CI-EIMS: *m/z* [M+ H]⁺ 361 [C₁₁H₉IN₂O₄ + H]⁺

Methyl 5'-iodohomovanillate (8): To a refluxing solution of **6** (350 mg, 0.97 mmol) in methanol (4 mL), freshly prepared silver benzoate [17] (140 mg) and triethylamine (2 mL) were added. After refluxing for 1 h, the reaction was worked up by filtration over silica gel and evaporation, and the residue was crystallized from diethylether to afford 260 mg (73%) **8** as an amorphous brownish powder.

Rf: 0.45 (light petroleum -EtOAc 7:3).

IR (KBr): 3644, 1767, 1737, 1463, 1415, 1279, 1189, 1042, 1010, 901 cm⁻¹.

¹H NMR (300 MHz, CDCl₃): 2.35 (3H, s, Ac), 3.54 (2H, s), 3.69 (3H, OMe), 3.80 (3H, s, OMe), 6.86 (1H, d, *J* = 1.5 Hz, Ar-H), 7.29 (1H, d, *J* = 1.5 Hz, Ar-H).

¹³C NMR (75 MHz, CDCl₃): 20.9 (CH₃), 40.5 (CH₂), 52.4 (CH₃), 56.2 (CH₃), 91.7 (C), 113.8 (CH), 130.9 (CH), 134.2 (C), 141.9 (C), 151.4 (C), 168.1 (C), 171.3 (C).

CI-EIMS: *m/z* [M+ H]⁺ 365 [C₁₂H₁₃IO₅ + H]⁺

5'-Iodohomovanillic acid (4a): To a solution of **8** (240 mg, 0.66 mmol) in water-THF 2:1 (3 mL), LiOH (194 mg, 4.6 mmol, 7 mol. equiv.) was added. After stirring at room temperature overnight, the reaction was diluted with water, extracted with EtOAc, sequentially washed with 2 N H₂SO₄ and brine, and then evaporated. The residue was purified by crystallization from CH₂Cl₂, affording 190 mg of a white powder.

MP: 178°C.

Rf: 0.15 (light petroleum -EtOAc 6:4).

IR (KBr): 3412, 1710, 1506, 1273, 1222, 1161, 1025, 884, 824 cm⁻¹.

¹H NMR (300 MHz, CDCl₃): 3.54 (2H, s), 3.88 (3H, s, OMe), 6.10 (1H, -OH, s), 6.76 (1H, d, *J* = 1.5 Hz, Ar-H), 7.21 (1H, d, *J* = 1.5 Hz, Ar-H).

¹³C NMR (75 MHz, acetone-d₆): 44.7 (CH₂), 55.5 (CH₃), 88.9 (C), 114.2 (CH), 124.9 (CH), 129.6 (C), 148.0 (C), 148.1 (C), 172.0 (C).

CI-EIMS: *m/z* [M+ H]⁺ 309 [C₉H₉IO₄ + H]⁺

5'-IodoRTX (1a): To a cooled (0°C) stirred solution of ROPA (**2**, 220 mg, 0.47 mmol) and 5'-iodohomovanillic acid (**4a**, 145 mg, 0.47 mmol, 1

mol. equiv.) in dry THF (2 mL), triphenylphosphine (TPP, 147 mg, 0.56 mmol, 1.2 mol. equiv.) and diisopropylazodicarboxylate (DIAD, 0.101 mL, 0.56 mmol, 1.2 mol. equiv.) were added. After stirring at room temperature for 2 h, the reaction was worked up by evaporation, and the residue was dissolved in toluene (*ca.* 5 mL) and cooled to 4°C overnight. After filtration of the copious white precipitate, the filtrate was purified by gravity column chromatography on alumina (25 mL, light petroleum-EtOAc 8:2 as eluant) to afford 184 mg (52%) of **1a**, having physical and spectroscopic (¹H NMR) properties identical to those reported in ref. 10.

¹³C NMR (75 MHz, CDCl₃): 10.4 (CH₃), 18.9 (CH₃), 19.9 (CH₃), 33.1 (CH), 35.8 (CH₂), 39.2 (CH), 39.4 (CH₂), 40.5 (CH₂), 41.1 (CH₂), 55.4 (CH₃), 56.4 (CH), 72.0 (C), 73.5 (C), 80.6 (CH), 81.2 (C), 84.5 (C), 110.8 (CH₂), 112.0 (CH), 117.9 (C), 126.7 (CH), 127.3 (C), 127.4 (CH), 128.9 (CH), 131.2 (CH), 131.4 (CH), 134.0 (C), 135.0 (C), 136.7 (C), 145.1 (C), 146.1 (C), 146.5 (C), 158.5 (CH), 171.1 (C), 208.4 (C).

Acknowledgments - We are grateful to Università del Piemonte Orientale for financial support to A.E.-C, and to Dr Sara Bacchiega for preliminary work on the iodination of vanillic derivatives.

References

- [1] Ashcroft FM. (2006) From molecule to malady. *Nature*, **440**, 440-447.
- [2] Clapham DE. (2003) TRP channels as cellular sensors. *Nature*, **426**, 517-524.
- [3] Appendino G, Muñoz E, Fiebich BL. (2003) TRPV1 (vanilloid receptor, capsaicin receptor) agonists and antagonists. *Expert Opinion Therapeutic Patents*, **13**, 1825-1837.
- [4] Wahl P, Foged C, Tullin S, Thomsen C. (2001) Iodo-resiniferatoxin, a new potent vanilloid receptor antagonist. *Molecular Pharmacology*, **59**, 9-15.
- [5] The K_(i) value for I-RTX after two hour incubation with cells transfected with TRPV1 has been estimated to be 4.2 pM (Lazar J, Braun DC, Toth A, Wang Y, Pearce LV, Pavlyukovets VA, Blumberg PM, Garfield SH, Wincovitch S, Choi HK, Lee J. (2006) Kinetics of penetration influence the apparent potency of vanilloids on TRPV1. *Molecular Pharmacology*, **69**, 1166-1173).
- [6] Trevisani M, Milan A, Gatti Z, Zanasi A, Harrison S, Fontana G, Morice AH, Geppetti P. (2004) Antitussive activity of iodo-resiniferatoxin in guinea pigs. *Thorax*, **59**, 769-772.
- [7] Rigoni M, Trevisani M, Mazzieri D, Nadaletto R, Tognetto M, Creminon C, Davis JB, Campi B, Amatesi S, Geppetti P, Harrison S. (2003) Neurogenic responses mediated by vanilloid receptor-1 (TRPV1) are blocked by the high affinity antagonist, iodo-resiniferatoxin. *British Journal of Pharmacology*, **138**, 977-985.
- [8] Appendino G, Daddario N, Minassi A, Schiano Morello A, De Petrocellis L, Di Marzo V. (2005) The taming of capsaicin. Reversal of vanilloid activity of *N*-acylvanillamines by aromatic iodination. *Journal of Medicinal Chemistry*, **48**, 4663-4669.
- [9] McDonnell ME, Zhang SP, Dubin AE, Dax SL. (2002) Synthesis and *in vitro* evaluation of a novel iodinated resiniferatoxin derivative that is an agonist at the human vanilloid VR1 receptor. *Bioorganic & Medicinal Chemistry Letters*, **12**, 1189-1192.
- [10] Seabrook GR, Sutton KG, Jarolimek W, Hollingworth JH, Teague S, Webb J, Clark N, Boyce S, Kerby J, Lai Z, Chou M, Middleton R, Kaczorowski G, Jones AB. (2002) Functional properties of the high-affinity TRPV1 (VR1) vanilloid receptor antagonist (4-hydroxy-5-iodo-3-methoxyphenylacetate ester) iodo-resiniferatoxin. *Journal of Pharmacology and Experimental Therapeutics*, **303**, 1052-1060.
- [11] The price of RTX in the 2006 Sigma Catalogue is 280,60 Euro/1mg
- [12] Fattorusso E, Lanzotti V, Tagliatela-Scafati O, Tron GC, Appendino G. (2002) Bisnorsesquiterpenoids from *Euphorbia resinifera* Berg. and an expeditious procedure to obtain resiniferatoxin from its fresh latex. *European Journal of Organic Chemistry*, 71-78.
- [13] Castanet A-S, Colobert F, Broutin P-E. (2002) Mild and regioselective iodination of electron-rich aromatics with *N*-iodosuccinimide and catalytic trifluoroacetic acid. *Tetrahedron Letters*, **43**, 5047-5048
- [14] Appendino G, Minassi A, Daddario N, Bianchi F, Tron GC. (2002) Chemoselective esterification of phenolic acids and alcohols. *Organic Letters*, **4**, 3839-3841.
- [15] The price of homovanillic acid in the 2006 Aldrich Catalogue is 12,10 Euro/100mg, while 5-iodovanillin is over 50 times cheaper (12,00 Euro/5 g).
- [16] Aoyama T, Shioiri T. (1980) Trimethylsilyldiazomethane, a new, stable, and safe reagent for the classical Arndt-Eisert synthesis. *Tetrahedron Letters*, **21**, 4461-4462.
- [17] Newman MS, Beal PF. (1950) An improved Wolff rearrangement in homogeneous medium. *Journal of American Chemical Society*, **72**, 5163-5165.
- [18] Anderson NG, Lust DA, Colapret KA, Simpson JH, Malley ME, Gougoutas JZ. (1996) Sulfonation with inversion by Mitsunobu reaction: An improvement on the original conditions. *Journal of Organic Chemistry*, **61**, 7955-7962.
- [19] Eisenbraun EJ. (1973) Cyclooctanone. *Organic Synthesis, Collective Volume 5*, 310-312.

Acaricides of Natural Origin. Part 2. Review of the Literature (2002-2006)[†]

Guido Flamini

Dipartimento di Chimica Bioorganica e Biofarmacia, Via Bonanno 33, 56126 Pisa, Italy

flamini@farm.unipi.it

Received: June 14th, 2006; Accepted: November 7th, 2006

Dedicated to the memory of Professor Ivano Morelli.

Acari are responsible for millions of dollars worth of damage each year as a result of infestations of animals, plants and man. They directly affect our health and prosperity as animal and plant parasites, vectors of disease, and producers of allergens. The indiscriminate use of pesticides has quickly induced resistance in many parasites. At present, the control of acarid parasitic diseases in agriculture, human and veterinary medicine is mainly based on the use of drugs; and for this reason the lack of effective drugs often prevents the control of some parasitic diseases, making them more serious and important. The use of commercial drugs involves many problems, besides the drug-resistance shown by the most important parasites. Environmental damage and the toxicity of many synthetic drugs, represent the main problems that strongly limit drug use. In addition, drug residues in plant and animal food products are important reasons for further economic losses for farmers and must be regarded as potentially hazardous to man and the environment. Plant-derived compounds are generally more easily degradable and could show a smaller negative environmental impact with respect to synthetic drugs. For these reasons, the evaluation of the antiacarid activity of plant extracts is increasingly being investigated in order to obtain new leads, as demonstrated by recent studies that have evaluated and confirmed the effectiveness of many plant compounds on bacteria, fungi, protozoa, helminths and arthropods. This review will be limited to the class Arachnida, sub-class Acaridi, particularly to their control in agriculture, veterinary and human medicine using natural methods.

Keywords: Acaricides, natural, human and veterinary medicine.

Mites and ticks, collectively known as the Acari, are of interest to humans for a variety of reasons. They affect our health and well being directly as plant, animal and human parasites, vectors of disease, and producers of allergens. The Class Arachnida, to which the order Acari belongs, together with the Classes Insecta, Crustacea and others, constitute the Phylum Arthropoda. All the classes contain species useful to man, but also many pests that are responsible for millions of dollars worth of economic losses each year as a result of infestations of animals, man and plants. This review will be limited to the order Acari, particularly to their control with natural methods in agriculture, veterinary and human medicine. This report follows and upgrades the previous one [1].

The indiscriminate use of inorganic pesticides destroyed many harmless species, including natural enemies of these mites and ticks [2]. After organochlorine and organophosphate pesticides were introduced, resistance was quickly acquired by many arthropod parasites, including acari; fortunately many useful predatory mites became resistant too. The emergence of resistance to parasiticides is one of the most serious challenges faced by man. Perhaps it is the simplicity of treating parasite attacks with either very effective drugs or pesticides on a routine basis, and the proven cost-effective gains in productivity that accrue in the short term, that has led to the predominance of synthetic pesticides [3]. Broadly speaking, resistance is the ability of the parasites to survive doses of drugs that would normally kill them at the same stage of development. The resistance is inherited and selected because the survivors of the

[†]For Part 1 see Ref. 1

pesticide treatment pass the genes for resistance on to their offspring. Drug susceptibility is a resource that needs to be preserved, using appropriate techniques of parasite management. The application of synthetic chemical substances is still the common method to either control or eradicate parasites of plant and animals, but many acaricides have non-specific properties, affecting other organisms (crops, non-vertebrates and vertebrates). Plants are the richest source of organic compounds on Earth, many of which are endowed with pesticide properties.

In veterinary medicine, the control of ectoparasites is of great importance due to their effects on livestock profitability and the health status of animals. Infestations of livestock can cause intense irritation, leading to poor condition, weight loss, reduced milk yield, and hide and fleece damage. Furthermore, many species of acari are responsible for transmission of diseases either to the host animals themselves or act as vectors of a number of diseases to humans [4].

Apiculture

During the 90s, several cases of resistance of *Varroa* mites (Acari: Varroidae) to common acaricides employed in beekeeping were reported from different countries [5–8]. In Italy, the consequences of the resistance led to disastrous colony losses. Available statistics show that in certain districts, losses often exceeded 70% and, in some locations, even reached 90% [9]. *Varroa* mites suck the body fluids from adults and brood, preferring the latter, especially the drone brood. The problem of developing suitable treatments was difficult in the case of the *Varroa* mites because most substances effective against the parasites have unacceptable side effects on bees. Since the creation of the EU *Varroa* experts' group, several lines of research in alternative control measures have been explored: apicultural techniques for reducing the number of mites, increasing bee resistance, and searching for acaricidal products that are generally recognized as safe for humans, such as some natural derivatives [10]. Many natural compounds have been evaluated for their effectiveness against *Varroa* mites [1]. During the five-year period employed in this review, some papers have described the use of simple carboxylic acids, such as formic [11–16] and oxalic [17–19] acids. Their efficacy has been known since 1980 [20]. Furthermore, a paper has appeared in the literature [21] about two different formulations of thymol, a well-known varroacidal agent. The main

goals of this study were to determine their effectiveness against *V. destructor* in an apiary in Sardinia (Italy), taking into account natural mite mortality in control hives and, simultaneously, to determine the persistence of both formulations and residues in honey and wax. Both thymol formulations, after the treatments, reduced significantly the levels of mite infestations of adult bees and sealed brood, but their efficacy, expressed as percentage of mortality, was lower for both products than the efficacy previously obtained with the same products under other experimental conditions. The residues were relatively higher in wax than in honey, because of the lipophilicity of thymol.

Other monoterpenes seem to be valuable as control agents for this mite, both in laboratory assays and in field treatments. The most active ones were linalyl acetate, myrtenyl acetate, perillyl acetate and thymyl acetate. In the field trials, all four monoterpenoid-treated groups were statistically significantly different from the control group in reduction of *V. destructor* infestation, yielding a 51–64% reduction of the mite when compared with the control group [22]. In an Argentinean study, the repellent and acaricidal effects of some essential oils from the most typical wild plant species from the northern part of the country were evaluated against *V. destructor*, using a complete exposure test. The lowest LD₅₀ values for mites were registered for *Acantholippia seriphioides* (1.27 µL per cage) and *Schinus molle* (2.65 µL per cage) after 24 hours and for *Wedelia glauca* (0.59 µL per cage) and *A. seriphioides* (1.09 µL per cage) after 72 hours of treatment [23].

Veterinary and Human Medicine

(a) Ticks: Particularly in the tropics, but also in many other countries, arthropod-borne diseases are among the major limiting factors to the efficient production of livestock and poultry. These diseases cause weakening, lameness, blindness, wasting, congenital defects, abortions, sterility, and death of the infested animals. Some exotic arthropod-borne diseases of livestock are zoonotic and affect humans as well as animals. Some of the most devastating of all animal diseases caused by arthropod-borne blood protozoa, include babesiosis of cattle, sheep, goats, horses, and swine; theileriosis, the East Coast fever syndrome, and Mediterranean fever; the trypanosomiasis causing illness in cattle, sheep, goats, camels, pigs, dogs, and many wild game

species; as well as several arthropod-borne protozoa that cause diseases of birds. The most prominent groups of arthropods that transmit etiological agents pathogenic to livestock are those that are hematophagous, such as ticks. The tick-borne diseases they transmit are among the most significant animal health deterrents to efficient livestock production. Ticks are obligate ectoparasites of vertebrates and they parasitize all vertebrate groups, except fishes.

The family Ixodidae comprises approximately 80% of all tick species, with the most economically important ixodid ticks that attack livestock in tropical regions belonging to the genera *Amblyomma*, *Boophilus*, *Rhipicephalus* and *Hyalomma*. *Ixodes scapularis* L., the black-legged tick, is the primary vector of disease-causing agents in humans in North America, especially Lyme disease, human granulocytic ehrlichiosis, and human babesiosis. Among the various strategies that have been considered for their control, one is the use of tick pheromones to facilitate targeted delivery of acaricides in the ticks natural habitat. Ticks use many different pheromones to regulate their behavior, especially for sexual activity. Some cause assembly, a type of behavior known as arrestment, defined as the cessation of kinetic activity. The chemical composition of the arrestment pheromone of *I. scapularis* has been identified so that it can be used as an aid in tick control [24]. The isolates that caused the arrestment were characterized as guanine and xanthine. The strongest responses were found when hematin was also present. These substances were used to formulate a preparation containing a mixture of pheromones and acaricidal substances, such as DEET (*N,N*-diethyl-*m*-toluamide) and permethrin. In this way, the efficacy of the acaricides was enhanced.

In further searching for alternative methods of reducing Lyme disease, the activity against *I. scapularis* nymphs was determined of 15 natural products isolated from the essential oil components extracted from the heartwood of *Chamaecyparis nootkatensis*. Nootkatone was the most effective eremophilane sesquiterpene, with an LC_{50} value of 0.029 mg/mL. Residual LC_{50} values for nootkatone did not differ significantly at 4 weeks post-treatment from the observations made after the initial 24 hours treatment. Among nymphal tick repellents, the most active compound was valencene-13-ol, with a repellent dose (RD_{50}) of 0.03 mg/mL at 4 hours, compared to 3.8 mg/mL for DEET. The ability of

these natural products to kill and repel ticks at relatively low concentrations may represent a future alternative to the use of synthetic pesticides [25].

Boophilus species are one-host ticks, which occur in all tropical and sub-tropical regions of the world, where they feed preferably on cattle. They are the main vectors of *Babesia* species, *B. bovis* and *B. bigemina*, causing babesiosis in cattle. *Boophilus* ticks, together with many other tick species, also transmit *Anaplasma marginale*, the rickettsia that causes anaplasmosis of cattle on all continents. The naturally occurring avermectins and milbemycins are fermentation products of actinomycetes in the genus *Streptomyces*. They are 16-membered, macrocyclic lactones, which have structural similarities to antibacterial macrolides and antifungal polyenes, but lack their antifungal and antibacterial activities and do not inhibit either protein or chitin synthesis [26]. Milbemycins, first described from a culture of *S. hygroscopicus*, are structurally similar to the avermectins, but lack the disaccharide substituent at C13 [27]. Mishima et al. [28] first reported the acaricidal activity of milbemycins. Moxidectin, a synthetically modified milbemycin derived from the fermentation product nemadectin [29], is used for insect and helminth control in animal health applications. Since 2001, only one paper about the use of moxidectin against this tick has been published [30]. It works like other macrocyclic lactones opening chloride channels in the nerve cells, causing paralysis. The macrocyclic lactone, moxidectin has a broad-spectrum activity against important internal and external parasites of cattle, including ticks [31–33]. The efficacy of a 1% injectable formulation of moxidectin at the dose of 0.20 mg/kg body weight by subcutaneous injection was greater than 95%. Furthermore, there was no evidence of either any local or systemic adverse reaction in treated animals and all cattle remained healthy throughout the trial period. For rapidly screening many compounds, a larval immersion microassay that offers superior sensitivity and flexibility to accommodate multiple formulations has been developed using the tick *Amblyomma americanum* (L.). This assay proved suitable for the identification and characterization of active molecules from natural product libraries, and it can be a useful tool to prioritize molecules for further *in vivo* testing in animal models [34].

(b) House dust mites: The term "house dust mites" is applied to a large number of mites found in association with dust in dwellings. Unlike some other

kinds of mites, house dust mites are not parasites of living plants, animals, or humans. House dust mites primarily live on dead skin cells regularly shed by humans and their animal pets. Skin cells and squames, commonly called dandruff, are often concentrated in parlor and sitting rooms, mattresses, frequently used furniture and associated carpeted areas, and may harbor large numbers of these microscopic mites. For most people, house dust mites are not harmful. The medical significance of house dust mites arises because their microscopic molted skins and feces, being major constituents of house dust, induces allergic reactions in some individuals. For those individuals, inhaling the house dust allergen triggers either rhinitis or bronchial asthma. Expert panel reports and position statements from the European Union, the US National Heart, Lung and Blood Institute (NHLBI), and the American Academy of Allergy, Asthma and Immunology (AAAAI) have recommended dust mite allergen avoidance as an integral part of asthma management [35–38]. House dust mites belong to different genera and species; the main ones are *Dermatophagoides pteronyssinus*, *D. farinae* and *Euroglyphus maynei* (Acari: Pyroglyphidae). However, there is great variation in the acarid fauna of the different regions of the world. *Dermatophagoides pteronyssinus* (literally "skin-eating mites") is considered as the true house dust mite and has a cosmopolitan distribution. Together with *D. farinae* (=flour, also infests stored food), it accounts for 80–90% of the total mite population generally found in houses. No pesticides are currently labeled for house dust mites. However, some commercial products are available for treatment of house dust mites and their allergens. The active ingredients are benzyl benzoate and tannic acid.

Recently, a new daphnane diterpenoid, rediocide F, was isolated together with the known redioides A, C and E, from the *n*-hexane extract of *Trigonostemon reidioides* roots by bioassay-guided fractionation using acaricidal activity on *Dermatophagoides pteronyssinus*. The structure of rediocide F was established as the demethyl analog of rediocide C. All the compounds exhibited potent activity against *D. pteronyssinus* with LC₅₀ values of 2.53, 0.78, 5.59 and 0.92 µg/cm², respectively [39]. Three *Uvaria* species, namely *U. klaineana*, *U. mocoli* and *U. versicolor* were tested *in vitro* against *D. pteronyssinus*. The most active extracts were the crude methanol and *n*-hexane extracts of *U. versicolor* stems, with EC₅₀ values of 0.095 g/m² and 0.12 g/m², respectively. The successive bioassay-

guided fractionation of the *n*-hexane extract led to the isolation of benzyl benzoate, which exhibited an EC₅₀ value of 0.045 g/m². A new flavanone, versuvanone, and the known oxoaporphine liriodenine were also isolated from this species and showed EC₅₀ values > 1.5 g/m². A weak acaricidal activity (0.85 g/m²) was observed for the dichloromethane extract of *U. klaineana*, due again to the presence of benzyl benzoate. *U. mocoli* extracts were inactive [40]. Other researchers tested the acaricidal activity of materials derived from the rhizome of *Cnidium officinale* Makino against *T. putrescentiae* adults using direct contact application and fumigation methods. The biologically active constituent was identified as butylidenephthalide by spectroscopic analyses. On the basis of 24-hours LD₅₀ values, the acaricidal activity of butylidenephthalide (5.80 µg/cm²) was more pronounced than that of the standard drugs benzyl benzoate (9.75 µg/cm²) and DEET (16.26 µg/cm²). Butylidenephthalide caused lethargy in the treated mites, leading to death without knockdown, whereas benzyl benzoate and DEET caused death associated with uncoordinated behavior. In a fumigation test with *T. putrescentiae* adults, butylidenephthalide was much more effective in closed containers than in open ones, indicating that the effects of this compound was largely due to action in the vapor phase [41]. Among essential oils, those obtained from *Pinus* species revealed promising activity against *T. putrescentiae*. The oils obtained from *P. pinea*, *P. halepensis*, *P. pinaster* and *P. nigra* have been evaluated for their acaricidal activity by aerial diffusion. Among them, the oil from *P. pinea* showed the best activity (100% deaths, while those from *P. halepensis* and *P. pinaster* were partially effective only at the higher dose); and the dose of 8 µL showed a percentage of dead mites statistically higher than that of the lower dose. The main constituents of the essential oil of *P. pinea* branches were α-pinene, β-caryophyllene, myrcene, 1,8-cineole, and limonene. Of these, α-pinene, β-caryophyllene, and myrcene were ineffective, whereas 1,8-cineole and limonene showed 100% acaricidal activity at 8 µL. Only 1,8-cineole maintained 100% acaricidal activity, also at the lower concentration of 6 µL [57].

Agriculture

Plant-feeding mites play important roles as agricultural pests of timber, fruits, vegetables, forage crops, and ornamentals. In many instances, lack of information about the correct identity of the mites, as

well as inadequate knowledge regarding their biology and ecology, have hampered our ability to combat effectively these mite pests. Their small size and cryptic appearance make mites difficult to detect, and thus infestations are often overlooked. Once established in a new area, certain biological characteristics allow rapid escalation to pest status. Miticidal compounds, as in veterinary and human medicine, cannot be toxic to the plant host and no harmful residues must be found in foods. Furthermore, in agriculture, an additional feature is requested: they must be devoid of undesirable effects on useful non-target organisms, like pollinators and predator arthropods. The main species are *Tetranychus* sps, *Oligonychus* sps (Acari: Tetranychidae), *Phyllocoptruta oleivora*, and *Tegolophus australis* (Acari: Eriophyidae). Among these, the two-spotted spider mite, *Tetranychus urticae*, a polyphagous pest, is probably one of the most dangerous for crops and ornamentals, particularly in glasshouses. Its high reproductive capacity enables it to cause serious damage in a short period. Furthermore, this parasite has developed resistance to many synthetic acaricides (see i.e. [42–45]), apart from the fact that many of these substances are toxic to useful non-target arthropods [46–48].

Among promising plant species, *Chenopodium ambrosioides* var. *ambrosioides* has been evaluated. An emulsifiable concentrate (UDA-245), obtained from the essential oil was compared with commercially available pesticides for their effectiveness to control the adult stage and egg hatch of the twospotted spider mite, *Tetranychus urticae* and the European red mite, *Panonychus ulmi*. A 0.5% UDA-245 was more effective than 0.7% neem oil on adult twospotted spider mites. In the case of the European red mite, UDA-245 was as effective as 0.006% (AI) abamectin. Furthermore, UDA-245 at 0.5% significantly reduced egg hatch of the twospotted spider mite, 5 and 9 days after treatment, and of the European red mite 6 days after treatment. Egg hatch was significantly lower using 0.006% abamectin, 0.7% neem oil, and 1.0% insecticidal soap than UDA-245. Residual tests indicated that UDA-245 may be persistent in the environment only for a few hours. Only 23% mortality was noted when mites were introduced on bean leaves 1 hour after treatment with 2% UDA-245. At the recommended dose of 0.5%, UDA-245 was not considered to be phytotoxic to most plants tested, i.e., lettuce, roses, and tomatoes. Results suggest that a greenhouse

integrated pest management program using UDA-245 could effectively and selectively control mite infestations by treating "hot spots", with negligible effect on biological control agents when treating before introduction or when natural enemies are absent [49].

Other natural derivatives effective against *Tetranychus urticae* can be obtained from leaves of wild tomato species, *Lycopersicon hirsutum*, *L. pennellii* and *L. pimpinellifolium*. Crude chloroform, ethanol and *n*-hexane extracts of the leaves were tested for their antibiosis and for their repellency. The antibiosis was assayed as a 6-hours no-choice test. The method for repellency utilized a ring bioassay. Chloroform leaf extracts of *L. hirsutum* exhibited the greatest antibiotic activity, and the *n*-hexane extracts exhibited the greatest repellency. Among the major chemical compounds of the extracts, α -curcumene, α -zingiberene, β -caryophyllene, 2-undecanone, and 2-tridecanone were detected. Lethality of the extracts was mainly associated with the presence of high concentrations of 2-tridecanone, while repellency of extracts was mainly associated with the presence of β -caryophyllene [50]. Another useful plant genus that can control *Tetranychus urticae* is *Taxus*. *T. cuspidata* and *T. media* var. *Hicksii* contain paclitaxel, among other taxoids, on the surface of the needles. These compounds were extracted by 5 seconds dipping of the needles in water at 96°C, 60°C and 40°C. The extracts with the higher concentration of paclitaxel were more harmful to the mites, increasing their mortality, prolonging development and lowering the average fecundity [51]. The citrus red mite, *Panonychus citri*, feeds on leaves, fruit and green twigs of all *Citrus* species. The infestation can result in heavy leaf drop, twig die-back and even death of large limbs. Various natural derivatives have been evaluated against this pest. A recent investigation has suggested that *Panonychus citri* is the most common causative allergen in citrus-cultivating farmers with either asthma or allergic rhinitis. Citrus red mite is a common sensitizing allergen among children living around citrus orchards [52].

It has been observed that spraying a *Mikania micrantha* alcohol extract on *Panonychus citri* could significantly decrease the survival rates of its eggs, larvae and nymphs. Furthermore, both the fecundity and the longevity of female *P. citri* fed on the leaves treated with *M. micrantha* alcohol extracts were significantly reduced. In a field experiment, the

efficacy of *M. micrantha* alcohol extracts was compared with water, alcohol, and pyridaben, a widely used acaricide in commercial control of red mite in sweet orange orchards. The survival rates of *P. citri* eggs, larvae and nymphs in the treatments were lower than the control, and were better than the pyridaben treatment [53].

Another study evaluated the acaricidal activity against this mite of a ginkgolic acid, 6-[(Z)-10-heptadecenyl]-2-hydroxybenzoic acid, isolated from the external seed coat of *Ginkgo biloba* [54]. Laboratory bioassays showed that this compound possessed powerful contact toxicity, similar to that of pyridaben and significantly superior to that of omethoate. Furthermore, it showed a quick-acting acaricidal activity, and was much faster-acting than either pyridaben or omethoate.

It has been observed that in the *Ageratum conyzoides* intercropped *Citrus* orchards, the populations of *Panonychus citri* were reduced. This could be explained by increases in the population of the predatory mite *Amblyseius newsami*, an effective natural enemy of citrus red mite. In fact the study showed that *A. conyzoides* produced and released volatile allelochemicals into the air in the intercropped citrus orchard, and these volatiles influenced the olfactory responses of *A. newsami* and *P. citri*. It has been observed that *A. conyzoides* fresh leaves, its essential oil, and major constituents, demethoxy-ageratochromene, β -caryophyllene, α -bisabolene, and (*E*)- β -farnesene, attracted *A. newsami* and slightly repelled *P. citri* [55]. To evaluate if some natural miticides were less toxic to useful arthropods than to phytophagous mites, some trials were conducted in apple orchards [56]. The European red mite, *Panonychus ulmi*, was the dominant and more harmful phytophagous species, followed by the apple rust mite, *Aculus schlechtendali*. Two predacious mites, *Typhlodromus pyri*, and *Zetzellia mali*, were often found in the orchards. Abamectin had favorable

selectivity, being more toxic to the two phytophagous mites than to the useful ones.

Conclusions

The control of parasitic diseases is mainly based on the use of effective drugs, both in agriculture and human and veterinary medicine; for this reason the lack of effective drugs often prevents the control of some parasitic diseases, making them more serious and important. At present, however, the use of commercial drugs involves many problems that strongly limit their use: foremost, the drug-resistance problem shown by the most important parasites, the environmental damage and the toxicity of many synthetic drugs.

Since plant-derived compounds are generally more easily degradable and could show reduced environmental damage with respect to synthetic drugs, at present the evaluation of the antiparasitic activity of plant extracts is being increasingly investigated, as demonstrated by recent studies that have evaluated and confirmed the effectiveness of many plant compounds on bacteria, fungi, protozoa, helminths and arthropods.

Perhaps human and veterinary medicine are the most suitable fields for a real application of natural drugs. Treatment of these pathologies is mostly topical, and particular drug-formulations are not required. Furthermore, generally only a few treatments are necessary to kill all the parasites. In agriculture, in spite of the studies performed to date, these substances are perhaps still far from their most effective use: their main useful feature, that is their biodegradability, is also their weakness. Often, many products are not able to persist in the environment for a period of time sufficient for pest control. Further studies are necessary to prepare better formulations that allow us to solve this problem. Other important future research topics should concentrate on the evaluation of the toxicity of these compounds, an unknown feature for many natural compounds.

References

- [1] Flamini G. (2003) Acaricides of natural origin, personal experiences and review of literature (1990-2001). In *Studies in Natural Product Chemistry — Bioactive Natural Products, Part I*. Atta-ur-Rahman (Ed). Elsevier, Amsterdam, 381–451.
- [2] Massee AM. (1958) The effect on the balance of arthropod populations in orchards arising from unrestricted use of chemicals. Proceedings of the 10th International Congress of Entomology “*Contrasting Approaches to Orchard Entomology*”, pp. 155-194.
- [3] Zajac AM, Sangster NC, Geary TG. (2000) Why veterinarians should care more about parasitology?. *Parasitology Today*, **16**, 504–506.
- [4] Taylor MA. (2001) Recent developments in ectoparasiticides. *Veterinary Journal*, **161**, 253–268.

- [5] Dujin T, Jovanovic V, Suvakov D, Milkovic Z. (1991) Effects of extended utilization of amytrase-based preparations on the formation of resistant strains of *Varroa jacobsoni*. *Veterinarski Glasnik*, **45**, 851–855.
- [6] Lodesani M, Colombo M, Spreafico M. (1995) Ineffectiveness of Apistan® treatment against the mite *Varroa jacobsoni* Oud. in several districts of Lombardy (Italy). *Apidologie*, **26**, 67–72.
- [7] Trouiller J. (1998) Monitoring of *Varroa jacobsoni* resistance to pyrethroids in Europe. *Apidologie*, **28**, 537–546.
- [8] Elzen PJ, Baxter JR, Spivak M, Wilson WT. (1999) Amitraz resistance in *Varroa*: new discovery in North America. *American Bee Journal*, **139**, 362.
- [9] Milani N. (2000) The resistance to chemotherapy in parasites and pathogens of the honeybee. *Proceedings of the International Conference MOMEDITO*, Prague, 117–131.
- [10] Colin ME, Ciavarella F, Otero-Colina G, Belzunces LP. (1994) A method for characterizing the biological activity of essential oils against *Varroa jacobsoni*. In *New perspectives on Varroa*. Matheson A (Ed). International Bee Research Association, Cardiff, 109–114.
- [11] Satta A, Floris I, Eguaras M, Cabras P, Garau V, Melis M. (2005) Formic acid-based treatments for control of *Varroa destructor* in a Mediterranean area. *Journal of Economic Entomology*, **98**, 267–273.
- [12] Bahreini R, Tahmasebi G, Nowzari J, Talebi M. (2004) A study of the efficacy of formic acid in controlling *Varroa destructor* and its correlation with temperature in Iran. *Journal of Apicultural Research*, **43**, 158–161.
- [13] Elzen PJ, Westervelt D, Lucas R. (2004) Formic acid treatment for control of *Varroa destructor* (Mesostigmata: Varroidae) and safety to *Apis mellifera* (Hymenoptera: Apidae) under southern United States conditions. *Journal of Economic Entomology*, **97**, 1509–1512.
- [14] Ostermann DJ, Currie RW. (2004) Effect of formic acid formulations on honey bee (Hymenoptera: Apidae) colonies and influence of colony and ambient conditions on formic acid concentration in the hive. *Journal of Economic Entomology*, **97**, 1500–1508.
- [15] Elzen PJ. (2003) Suitability of formic acid to control *Varroa destructor* and safety to *Apis mellifera* in the southwestern U.S. *Southwestern Entomologist*, **28**, 261–266.
- [16] Eguaras M, Palacio MA, Faverin C, Basualdo M, Del Hoyo ML, Velis G, Bedascarrasbure EL. (2003) Efficacy of formic acid in gel for *Varroa* control in *Apis mellifera* L.: importance of the dispenser position inside the hive. *Veterinary Parasitology*, **111**, 241–245.
- [17] Rademacher E, Harz M. (2005) Oxalic acid for the control of varroosis in honey bee colonies - a review. *Apidologie*, **37**, 98–120.
- [18] Hatjina F, Haristos L. (2005) Indirect effects of oxalic acid administered by trickling method on honey bee brood. *Journal of Apicultural Research*, **44**, 172–174.
- [19] Gregorc A, Planinc I. (2002) The control of *Varroa destructor* using oxalic acid. *Veterinary Journal*, **163**, 306–310.
- [20] Ritter W, Ruttner F. (1980) Neue Wege in der Behandlung der Varroatose. *Allgemeine Deutsche Imkerzeitung*, **14**, 151–155.
- [21] Floris I, Satta A, Cabras P, Garau V, Angioni A. (2004) Comparison between two thymol formulations in the control of *Varroa destructor*: effectiveness, persistence, and residues. *Journal of Economic Entomology*, **97**, 187–191.
- [22] Fassbinder C, Grodnitzky J, Coats JR. (2002) Monoterpenoids as possible control agents for *Varroa destructor*. *Journal of Apicultural Research*, **41**, 83–88.
- [23] Ruffinengo S, Eguaras M, Floris I, Faverin C, Bailac P, Ponzi M. (2005) LD₅₀ and repellent effects of essential oils from Argentinian wild plant species on *Varroa destructor*. *Journal of Economic Entomology*, **98**, 651–655.
- [24] Sonenshine DE, Adams T, Sandra AA, McLaughlin J, Webster FX. (2003) Chemical composition of some components of the arrestment pheromone of the black-legged tick, *Ixodes scapularis* (Acari: Ixodidae) and their use in tick control. *Journal of Medical Entomology*, **40**, 849–859.
- [25] Panella NA, Dolan MC, Karchesy JJ, Xiong Y, Peralta-Cruz J, Khasawneh M, Montenieri JA, Maupin GO. (2005) Use of novel compounds for pest control: insecticidal and acaricidal activity of essential oil components from heartwood of Alaska yellow cedar. *Journal of Medical Entomology*, **42**, 352–358.
- [26] Fisher MH, Mrozik H. (1989) Chemistry. In *Ivermectin and Abamectin*. Campbell WC (Ed). Springer, New York, 1–23.
- [27] Takiguchi Y, Mishna H, Okuda M, Tenao M. (1980) Milbemycins, a new family of macrolide antibiotics: fermentation, isolation and physico-chemical properties. *Journal of Antibiotics*, **33**, 1120–1127.
- [28] Mishima H, Kurabayashi M, Tamura C, Sato S, Kuwano H, Aoki ASA. (1975) Structures of milbemycin beta1, beta2, and beta3. *Tetrahedron Letters*, **16**, 711–714.
- [29] Carter GT, Nietsche JA, Hertz MR. (1988) LLF28249 antibiotic complex: a new family of antiparasitic macrocyclic lactones: isolation, characterization and structures of LL-F28249a,b,g,l. *Journal of Antibiotics*, **41**, 519–529.
- [30] Aguilar-Tipacamu G, Rodriguez-Vivas RI. (2003) Effect of moxidectin against natural infestation of the cattle tick *Boophilus microplus* (Acarina: Ixodidae) in the Mexican tropics. *Veterinary Parasitology*, **111**, 211–216.
- [31] Sibson GJ. (1994) The effects of moxidectin against natural infestation of cattle tick *Boophilus microplus*. *Australian Veterinary Journal*, **71**, 22–23.
- [32] Remington B, Kieran P, Coob R, Boder D. (1997) The application of moxidectin formulations for control of the cattle tick (*Boophilus microplus*) under Queensland field conditions. *Australian Veterinary Journal*, **75**, 588–591.

- [33] Gugliemone A, Mangold AJ, Muñoz CM, Scherling N, Garcia RF, Anziani OS, Loppolo M. (2000) Moxidectin pour-on for control of natural populations of the cattle tick *Boophilus microplus* (Acarina: Ixodidae). *Veterinary Parasitology*, **87**, 237–241.
- [34] White W, Hunter P, Plummer PR, Kemper CJ, Miller RJ, Davey RB, Kemp DH, Hughes S, Smimth CK, Gutierrez JA. (2004) An *in vitro* larval immersion microassay for identifying and characterizing candidate acaricides. *Journal of Medical Entomology*, **41**, 1034–1042.
- [35] Bahir A, Goldberg A, Mekori YA, Confino-Cohen R, Morag H, Rosen Y, Monakir D, Rigler S, Cohen AH, Horev Z, Noviski N, Mandelberg A. (1997) Continuous avoidance measures with or without acaricide in dust mite-allergic asthmatic children. *Annals of Allergy, Asthma and Immunology*, **78**, 506–512.
- [36] van der Heide S, Kauffman HF, Dubois AEJ, de Monchy JGR. (1997) Allergen-avoidance measures in homes of house-dust-mite-allergic asthmatic patients: effects of acaricides and mattress encasings. *Allergy*, **52**, 921–927.
- [37] Tovey E, Marks G. (1999) Methods and effectiveness of environmental control. *Journal of Allergy and Clinical Immunology*, **103**, 179–191.
- [38] Arlian LG, Platts-Mills TAE. (2001) The biology of dust mites and the remediation of mite allergens in allergic disease. *Journal of Allergy and Clinical Immunology*, **107**, s406–s413.
- [39] Soonthornchareonnon N, Sakayarojkul M, Isaka M, Mahakittikun V, Chuakul W, Wongsinkongman P. (2005) Acaricidal daphnane diterpenoids from *Trigonostemon reidioides* (Kurz) Craib roots. *Chemical and Pharmaceutical Bulletin*, **53**, 241–243.
- [40] Akendengue B, Ngou-Milama E, Bourobou-Bourobou H, Essouma J, Roblot F, Gleye C, Laurens A, Hocquemiller R, Loiseau P, Bories C. (2003) Acaricidal activity of *Uvaria versicolor* and *Uvaria klaineana* (Annonaceae). *Phytotherapy Research*, **17**, 364–367.
- [41] Kwon JH, Ahn YJ. (2003) Acaricidal activity of *Cnidium officinale* rhizome-derived butyridenephthalide against *Tyrophagus putrescentiae* (Acari: Acaridae). *Journal of Chemical Ecology*, **59**, 119–123.
- [42] Flexner JL, Westigard PH, Hilton R, Croft BA. (1995) Experimental evaluation of resistance management for twospotted spider mite (Acari: Tetranychidae) on southern Oregon Pear: 1981-1993. *Journal of Economic Entomology*, **88**, 1517–1524.
- [43] Herron GA, Rophail J. (1998) Tebufenpyrad (Pyranica (R)) resistance detected in two-spotted spider mite *Tetranychus urticae* Koch (Acari: Tetranychidae) from apples in Western Australia. *Experimental and Applied Acarology*, **22**, 633–641.
- [44] Herron GA, Learmonth SE, Rophail J, Barchia I. (1997) Clofentezine and fenbutatin oxide resistance in the two-spotted spider mite, *Tetranychus urticae* Koch (Acari: Tetranychidae) from deciduous fruit tree orchards in Western Australia. *Experimental and Applied Acarology*, **21**, 163–169.
- [45] Ho CC. (2000) Spider-mite problems and control in Taiwan. *Experimental and Applied Acarology*, **24**, 453–462.
- [46] Fischer-Colbrie P, El-Borolossy M. (1988) Nebenwirkungen von Winter- und Austriebbehandlungen auf Raubmilbenpopulationen an Apfelbaumen. *Pflanzenschutzberichte*, **49**, 125–131.
- [47] Stolz M. (1990) Testing side effects of various pesticides on the predatory mite *Phytoseiulus persimilis* Athias-Henriot (Acarina: Phytoseiidae) in laboratory. *Pflanzenschutzberichte*, **51**, 127–138.
- [48] Stolz M. (1994) Efficacy of different concentrations of seven pesticides on *Phytoseiulus persimilis* A.H. (Acarina: Phytoseiidae) and on *Tetranychus urticae* K. (Acarina: Tetranychidae) in laboratory and semifield test. *Bulletin OILB SROP*, **17**, 49–54.
- [49] Chiasson H, Bostanian NJ, Vincent C. (2004) Acaricidal properties of a *Chenopodium*-based botanical. *Journal of Economic Entomology*, **97**, 1373–1377.
- [50] Antonious GF, Snyder J. (2006) Natural products: repellency and toxicity of wild tomato leaf extracts to the two-spotted spider mite, *Tetranychus urticae* Koch. *Journal of Environmental Science and Health, Part B*, **41**, 43–55.
- [51] Furmanowa M, Kropczynska D, Zobel A, Glowinski K, Oledzka H, Jozefowicz J, Sahajdak A, Jozetczyk A. (2002) Influence of water extracts from the surface of two Yew (*Taxus*) species on mites (*Tetranychus urticae*). *Journal of Applied Toxicology*, **22**, 107–109.
- [52] Lee MH, Cho SH, Park HS, Bahn JW, Lee BJ, Son JW, Kim YK, Koh YY, Min KU, Kim YY. (2000) Citrus red mite (*Panonychus citri*) is a common sensitizing allergen among children living around citrus orchards. *Annals of Allergy, Asthma and Immunology*, **85**, 200–204.
- [53] Cen Y, Pang X, Xu C, Deng Q. (2005) Control effects of *Mikania micrantha* alcohol extract on citrus red mite *Panonychus citri*. *Ying yong sheng tai xue bao* (=The Journal of Applied Ecology), **16**, 754–757.
- [54] Pan W, Luo P, Fu R, Gao P, Long Z, Xu F, Xiao H, Liu S. (2006) Acaricidal activity against *Panonychus citri* of a ginkgolic acid from the external seed coat of *Ginkgo biloba*. *Journal of Chemical Ecology*, **62**, 283–287.
- [55] Kong C, Hu F, Xu X, Zhang M, Liang W. (2005) Volatile allelochemicals in the *Ageratum conyzoides* intercropped citrus orchard and their effects on mites *Amblyseius newsami* and *Panonychus citri*. *Journal of Chemical Ecology*, **59**, 2193–2203.
- [56] Hardman JM, Franklin JL, Moreau DL, Bostanian NJ. (2003) An index for selective toxicity of miticides to phytophagous mites and their predators based on orchard trials. *Pest Management Science*, **59**, 1321–1332.
- [57] Macchioni F, Cioni PL, Flamini G, Morelli I, Perrucci S, Franceschi A, Macchioni G, Ceccarini L. (2002) Acaricidal activity of pine essential oils and their main components against *Tyrophagus putrescentiae*, a stored food mite. *Journal of Agricultural and Food Chemistry*, **50**, 4586–4588.

Chemical and Biological Activity of Triterpene Saponins from *Medicago* Species

Aldo Tava^{*a} and Pinarosa Avato^b

^aC.R.A. Istituto Sperimentale per le Colture Foraggere, viale Piacenza 29, 26900 Lodi, Italy

^bDipartimento Farmaco-Chimico, Università di Bari, via Orabona 4, 70125 Bari, Italy

aldo.tava@entecra.it

Received: June 1st, 2006; Accepted: September 7th, 2006

Dedicated to the memory of Professor Ivano Morelli.

Naturally occurring saponins are a large group of triterpene and steroid glycosides characterized by several biological and pharmacological properties. The *Medicago* genus represents a valuable source of saponins which have been extensively investigated. This review summarizes the chemical features of saponins from *Medicago* species and their biological activity, with particular attention to their antimicrobial, insecticidal, allelopathic and cytotoxic effects. Influence of saponins on animal metabolism is also reported.

Keywords: *Medicago*, *M. sativa*, saponins, chemical structure, biological activity.

Saponins are a large group of plant metabolites including triterpenoids, steroids and steroidal alkaloids glycosylated with one or more sugar chains [1, 2]. They are commonly distinguished by their surfactant and hemolytic activities. Naturally occurring saponins display a broad spectrum of biological and pharmacological properties such as fungicidal, molluscicidal, antibacterial, antiviral and antitumor activities [2-6]. Due to their chemical, physical and physiological characteristics, commercial products containing plant saponins are available and used in the pharmaceutical, cosmetic and food industries [7-8]. Some saponins are the starting material for the semisynthesis of drugs and some are used as emulsifiers and foaming agents in food. Plant extracts rich in saponins have been used as folk detergents and are ingredients of cosmetic preparations such as lipsticks, shampoos and toothpaste.

Saponins are produced by many plant species and their distribution in the plant kingdom seems to be correlated with the structural type. That is, steroidal saponins have been found almost exclusively in the Monocotyledons, while triterpenoid saponins mainly

occur in the Dicotyledons and are practically absent in the Gymnosperms [6, 9]. The *Leguminosae* have been extensively investigated for their saponin content and within this family of plants, the *Medicago* genus represents a particularly rich source of bioactive saponins [10-15].

The genus *Medicago* includes 83 different species, the most known represented by *M. sativa* L. (syn. *M. media* Pers.), or alfalfa, a highly valued forage crop [16, 17]. The chemical structure of saponins from several species within the genus has been determined [18-46]. Generally they are complex mixtures of high-molecular weight triterpene glycosides with medicagenic acid, hederagenin, zanhic acid, bayogenin and soyasapogenols A and B as the dominant aglycones. Recently the 2 β ,3 β -dihydroxy-23-oxo-olean-12-en-28-oic acid has been identified as a new aglycone moiety in the two species *M. arborea* [45] and *M. hybrida* [46]. A summary of the structural types of saponins isolated from the various species of *Medicago* is reported in Figure 1.

Sugars or sugar chains are generally linked at the C-3 position of the aglycone (monodesmosides) and




Figure 1: Chemical structure of sapogenins detected in *Medicago* species.

additionally at the C-28 position, giving the corresponding bidesmosides. A tridesmoside saponin (extra sugar at the C-23 position) has only been identified in *M. sativa* [36] and *M. truncatula* [42].

The most abundant monosaccharide units found in the *Medicago* saponins are: arabinose, rhamnose, xylose, glucose, and glucuronic acid. Saponins in the *Medicago* species are produced in all the plant organs: leaves, flowers, roots, seeds and sprouts [10-49]. Their content in the plant material changes as a function of several factors, such as plant organs, genotype, cutting, year and stage of growth, and environmental effects, as reported for *M. sativa*, the most studied species of the genus [50-55]. The chemotaxonomic significance of saponins has also been investigated as their composition can discriminate among *Medicago* species [56-59]. The occurrence of saponins in the *Medicago* genus is long known [60], and their composition has been studied in several species. Structure elucidation of complex saponin mixtures differentiates their aglycone composition [61, 62]. In particular, investigation of several annual and perennial wild and cultivated *Medicago* species, showed variability in the aglycone composition of the saponins from each species.

Medicagenic acid was detected in some of them and soyasapogenol B was often present in the form of soyasaponin I, a common saponin of the *Leguminosae* family [10, 63].

Chemical analysis of saponins is not simple due to their 'soapy' properties due to sugars in the molecules. Their presence can be evaluated by biological tests involving their toxic haemolytic [64], fungicidal [65] and insecticidal [66] properties. Chemical methods also have been used, such as TLC [10], HPLC [67], GC and GC/MS [52, 68, 69], the last technique being used to analyze and quantify only the aglycone moieties. Capillary electrophoresis [70] and LC/MS methods [40, 42, 43] have also been employed for the identification and quantification of saponins in the plant extracts.

Structure investigation of *Medicago* saponins is usually performed by preliminary identification of the sapogenins and sugars released after acid hydrolysis from pure saponins obtained by direct and reverse-phase chromatographic separation of the raw saponin mixtures. Detailed information on the saponin structure, however, could be obtained only by a combination of analytical methods, including

MS [71] and NMR analyses [72, 73], performed on pure compounds. The MS spectra allowed subsequent fragmentation of the sugar chains to give the corresponding aglycones, NMR analyses (^1H , ^{13}C and 2D experiments) allowed the determination of all the carbon atoms and the sugar linkage in the molecules, while the absolute configuration of monosaccharides was generally obtained by GC analyses with a chiral capillary column. Detailed investigations on saponin chemical structures have until now been reported for *M. arabica*, *M. arborea*, *M. hybrida*, *M. lupulina*, *M. polymorpha*, *M. sativa* and *M. truncatula* [19-46]. Characterized saponins from these species of *Medicago* are listed in Tables 1-7.

Table 1: Saponins identified in *M. arabica* leaves [41].

Aglycone	3 OH substituted	28 COOH substituted
2 β -Hydroxy oleanolic acid	α -L-Ara(1 \rightarrow 2)- β -D-Glc	β -D-Glc
Hederagenin	α -L-Ara	-
Hederagenin	α -L-Ara	β -D-Glc
Hederagenin	β -D-Glc(1 \rightarrow 2)- α -L-Ara	-
Hederagenin	β -D-Glc(1 \rightarrow 2)- α -L-Ara	β -D-Glc
Hederagenin	α -L-Ara(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- α -L-Ara	β -D-Glc
Bayogenin	α -L-Ara	-
Bayogenin	α -L-Ara	β -D-Glc

M. arabica leaves are characterized by the presence of short sugar chain saponins, including mono and bidesmosides of 2- β -hydroxyoleanolic acid, hederagenin and bayogenin (Table 1). *M. arborea* leaves produce saponins containing up to seven sugars, identified as mono and bidesmosides of medicagenic and zanhic acid (Table 2). Saponins from *M. hybrida* roots are characterized by the presence of short sugar chain bidesmosides of hederagenin and medicagenic acid (Table 3). *M. lupulina* leaves contain mono and disaccharide saponins of hederagenin and medicagenic acid (Table 4), while saponins from the leaves of *M. polymorpha* predominantly consist of short sugar chain bidesmosides of hederagenin and caulophyllogenin (Table 5). Saponins from the roots and the aerial parts of *M. sativa* are a complex mixture of both short and long sugar chains of mono and bidesmosidic compounds with hederagenin, medicagenic acid, zanhic acid and soyasapogenols as the most representative aglycones. In this species a tridesmoside saponin containing eight monosaccharide units and a β -maltoside derivative, 3-*O*-[α -D-glucopyranosyl(1 \rightarrow 4)- β -D-glucopyranosyl] medicagenic acid, were also identified in the aerial parts (Table 6). *M. truncatula* saponins from both roots and aerial parts (Table 7) are long sugar chain bidesmosides of medicagenic and zanhic acid.

Table 2: Saponins identified in *M. arborea* leaves [45].

Aglycone	3 OH substituted	28 COOH substituted
2 β -Hydroxy oleanolic acid	α -L-Rha(1 \rightarrow 2)- α -L-Ara(1 \rightarrow 2)- β -D-Glc	-
Bayogenin	β -D-GlcA	β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara
2 β ,3 β -Dihydroxy-23-oxo-olean-28-oic acid	β -D-GlcA	β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara
Medicagenic acid	β -D-Glc	α -L-Rha(1 \rightarrow 2)- α -L-Ara
Medicagenic acid	β -D-GlcA	α -L-Rha(1 \rightarrow 2)- α -L-Ara
Medicagenic acid	β -D-Glc	β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara
Medicagenic acid	β -D-GlcA	β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara
Medicagenic acid	β -D-Glc(1 \rightarrow 2)- β -D-Glc	β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara
Medicagenic acid	β -D-GlcA	β -D-Api(1 \rightarrow 3)-[β -D-Xyl(1 \rightarrow 4)]- α -L-Rha(1 \rightarrow 2)- α -L-Ara
Zanhic acid	β -D-Glc	α -L-Ara(1 \rightarrow 3)- α -L-Rha(1 \rightarrow 2)- α -L-Ara
Zanhic acid	β -D-GlcA	β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara
Zanhic acid	β -D-Glc	α -L-Ara(1 \rightarrow 3)-[β -D-Xyl(1 \rightarrow 4)]- α -L-Rha(1 \rightarrow 2)- α -L-Ara
Zanhic acid	β -D-GlcA	β -D-Api(1 \rightarrow 3)-[β -D-Xyl(1 \rightarrow 4)]- α -L-Rha(1 \rightarrow 2)- α -L-Ara
Zanhic acid	β -D-GlcA	α -L-Ara(1 \rightarrow 3)-[β -D-Xyl(1 \rightarrow 4)]- α -L-Rha(1 \rightarrow 2)- α -L-Ara
Zanhic acid	β -D-Glc(1 \rightarrow 2)- β -D-Glc	α -L-Ara(1 \rightarrow 3)-[β -D-Xyl(1 \rightarrow 4)]- α -L-Rha(1 \rightarrow 2)- α -L-Ara
Zanhic acid	α -L-Ara(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- β -D-Glc	β -D-Api(1 \rightarrow 3)-[β -D-Xyl(1 \rightarrow 4)]- α -L-Rha(1 \rightarrow 2)- α -L-Ara
Soyasapogenol A	α -L-Rha(1 \rightarrow 2)- β -D-Gal(1 \rightarrow 2)- β -D-GlcA	α -L-Rha
Soyasapogenol B	α -L-Rha(1 \rightarrow 2)- β -D-Gal(1 \rightarrow 2)- β -D-GlcA	-

Table 3: Saponins identified in *M. hybrida* roots [46].

Aglycone	3 OH substituted	28 COOH substituted
Oleanolic acid	β -D-Gal(1 \rightarrow 2)- β -D-GlcA	β -D-Glc
Oleanolic acid	β -D-Gal(1 \rightarrow 2)- β -D-GlcA	α -L-Rha(1 \rightarrow 4)- β -D-Glc
Hederagenin	β -D-Glc	-
Hederagenin	β -D-GlcAMe	-
Hederagenin	β -D-Glc(1 \rightarrow 2)- α -L-Ara	-
Hederagenin	β -D-GlcA	β -D-Glc
Hederagenin	β -D-GlcAMe	β -D-Glc
Hederagenin	α -L-Rha(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- β -D-Glc	β -D-Glc
Bayogenin	β -D-Glc	β -D-Glc
2 β ,3 β -Dihydroxy-23-oxo-olean-12-en-28-oic acid	β -D-GlcA	β -D-Glc
Medicagenic acid	β -D-Glc	-
Medicagenic acid	β -D-Glc	β -D-Glc
Medicagenic acid	β -D-GlcA	β -D-Glc
Medicagenic acid	β -D-Glc(1 \rightarrow 2)- β -D-Glc	β -D-Glc

Table 4: Saponins identified in *M. lupulina* [28].

Aglycone	3 OH substituted	28 COOH substituted
Hederagenin	β -D-Glc	-
Medicagenic acid	β -D-Glc	-
Medicagenic acid	β -D-Glc	β -D-Glc
Soyasapogenol B	α -L-Rha(1 \rightarrow 2)- β -D-Gal(1 \rightarrow 2)- β -D-GlcA	-

Table 5: Saponins identified in *M. polymorpha*.

Aglycone	3 OH substituted	28 COOH substituted	Ref.
Oleanolic acid	α -L-Rha(1 \rightarrow 2)- α -L-Ara	β -D-Glc-(1 \rightarrow 6)- β -D-Glc	[38]
Hederagenin	α -L-Rha(1 \rightarrow 2)- α -L-Ara	-	[38]
Hederagenin	α -L-Ara	β -D-Glc-(1 \rightarrow 6)- β -D-Glc	[38]
Hederagenin	α -L-Rha(1 \rightarrow 2)- α -L-Ara	β -D-Glc	[38]
Hederagenin	α -L-Rha(1 \rightarrow 2)- α -L-Ara	β -D-Glc-(1 \rightarrow 6)- β -D-Glc	[38]
Caulophyllogenin	α -L-Rha(1 \rightarrow 2)- α -L-Ara	β -D-Glc	[38]
Caulophyllogenin	α -L-Rha(1 \rightarrow 2)- α -L-Ara	β -D-Glc-(1 \rightarrow 6)- β -D-Glc	[38]
Soyasapogenol B	α -L-Rha(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- β -D-GlcA	-	[33]

Branched sugar chain saponins were identified in this species, as in *M. arborea* and *M. sativa*. Methyl ester derivative of saponins were also found in *M. hybrida* and *M. sativa*, but these were recognized as artifacts obtained during the extraction with methanol [74].

The nature of the saccharide units, their position on the molecule and the similarity of the sugar chains on saponins from the different species, have suggested high enzymatic selectivity for the sugar position. Hederagenin often contains an α -L-arabinopyranose unit as the first sugar in its 3-*O* position. Alternatively a β -D-glucopyranose or the corresponding uronic derivative are present as in *M. hybrida* root saponins. The second monosaccharide unit linked at the C-2 position of α -L-arabinopyranose can be α -L-rhamnopyranose, as in *M. polymorpha*, or β -D-glucopyranose as in *M. arabica* and *M. sativa*.

By contrast, in all the studied species of *Medicago*, saponins of medicagenic and zanhic acids are always characterized by the presence of β -D-glucopyranose or β -D-glucuronopyranose units as the first sugar in the 3-*O* position. The second monosaccharide β -D-glucopyranose, linked predominantly at the C-2 position, as in *M. arborea*, *M. hybrida* and *M. sativa*, or at the C-3 position, as in *M. truncatula* suggesting the presence of a specific glucosyltransferase in this species. Different sugar linkage positions were also detected in the 3-*O* disaccharide chain of *M. sativa*, in which the 1 \rightarrow 3 and 1 \rightarrow 4 linkage between the first and the second monosaccharide were found. Trisaccharides are predominantly 3-*O*- β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranosyl-(1 \rightarrow 2)- β -D-glucopyranosyl derivatives.

The C-28 glycosylated saponins showed the presence of the β -D-glucopyranose unit esterified at the carboxylic group, the disaccharide chain 28-*O*- β -D-glucopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside (only found in *M. polymorpha*), and chains with more than two sugars, always characterized by α -L-arabinopyranose, directly linked at the C-28, and an α -L-rhamnopyranose in the central position, linked (1 \rightarrow 4) with a β -D-xylopyranose. Branching points are formed by α -L-arabinopyranose or β -D-apiofuranose linked (1 \rightarrow 3) at the β -D-xylopyranose unit. These features are typical of saponins extracted from *M. arborea*, *M. sativa* and *M. truncatula* and suggest high enzymatic selectivity for the sugar position independent of the involved genin.

Table 6. Saponins identified in *M. sativa* leaves and roots.

Aglycone	3 OH substituted	28 COOH substituted	Ref.
Hederagenin	β -D-Glc(1 \rightarrow 2)- α -L-Ara	-	[13]
Hederagenin	β -D-Glc(1 \rightarrow 2)- α -L-Ara	β -D-Glc	[37]
Hederagenin	β -D-Glc(1 \rightarrow 2)- α -L-Ara	β -D-Glc	[27]
Hederagenin	β -D-Glc(1 \rightarrow 3)- β -D-Xyl	β -D-Glc	[37]
Hederagenin	α -L-Ara(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- α -L-Ara	-	[23]
Hederagenin	α -L-Ara(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- α -L-Ara	β -D-Glc	[24]
Bayogenin	β -D-Gal(1 \rightarrow 2)- β -D-GlcA	β -D-Glc	[39]
Medicagenic acid	-	β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara	[34]
Medicagenic acid	β -D-Glc	-	[19]
Medicagenic acid	β -D-GlcA	-	[31]
Medicagenic acid	β -D-Glc	β -D-Glc	[22]
Medicagenic acid	β -D-Glc	α -L-Rha(1 \rightarrow 2)- α -L-Ara	[30]
Medicagenic acid	β -D-GlcA	α -L-Rha(1 \rightarrow 2)- α -L-Ara	[36]
Medicagenic acid	β -D-Glc	β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara	[25]
Medicagenic acid	β -D-GlcA	β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara	[31]
Medicagenic acid	β -D-GlcA Me ester	β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara	[39]
Medicagenic acid	α -D-Glc(1 \rightarrow 4)- β -D-Glc	-	[29]
Medicagenic acid	β -D-Glc(1 \rightarrow 3)- β -D-Glc	β -D-Glc	[39]
Medicagenic acid	β -D-Gal(1 \rightarrow 2)- β -D-Glc	β -D-Glc	[27]
Medicagenic acid	β -D-Glc(1 \rightarrow 2)- β -D-Glc	β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara	[27]
Medicagenic acid	α -L-Rha(1 \rightarrow 6)- β -D-GlcA(1 \rightarrow 2)- β -D-Glc	-	[20]
Medicagenic acid	β -D-Glc(1 \rightarrow 6)- β -D-Glc(1 \rightarrow 3)- β -D-Glc	-	[21]
Medicagenic acid	α -L-Rha(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- β -D-Glc	-	[39]
Medicagenic acid	α -L-Rha(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- β -D-Glc	β -D-Glc	[27]
Medicagenic acid	β -D-Glc(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- β -D-Glc	β -D-Glc	[39]
Medicagenic acid	β -D-Glc(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- β -D-Glc	β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara	[39]
Medicagenic acid	β -D-Glc(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- β -D-Glc	β -D-Api(1 \rightarrow 3)-[β -D-Xyl(1 \rightarrow 4)]- α -L-Rha(1 \rightarrow 2)- α -L-Ara	[39]
Medicagenic acid	β -D-Glc(1 \rightarrow 2)-[α -L-Rha(1 \rightarrow 3)]- β -D-Glc(1 \rightarrow 2)- β -D-Glc	β -D-Glc	[32]
Medicagenic acid	Glc-malonyl	-	[40]
Medicagenic acid	Glc-malonyl	Glc	[40]
Zanhic acid	β -D-Glc(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- β -D-Glc	β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara	[39]
Zanhic acid	β -D-Glc(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- β -D-Glc	β -D-Api(1 \rightarrow 3)-[β -D-Xyl(1 \rightarrow 4)]- α -L-Rha(1 \rightarrow 2)- α -L-Ara	[39]
Zanhic acid	β -D-Glc(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- β -D-Glc	β -D-Api(1 \rightarrow 3)- β -D-Xyl(1 \rightarrow 4)- α -L-Rha	[36]
Zanhic acid	β -D-Glc(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- β -D-Glc	23 COOH substituted: α -L-Ara β -D-Api(1 \rightarrow 3)- β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara 23 COOH substituted: α -L-Ara	[36]
Soyasapogenol A	α -L-Rha(1 \rightarrow 2)- β -D-Gal(1 \rightarrow 2)- β -D-GlcA	α -L-Rha	[39]
Soyasapogenol B	β -D-Glc(1 \rightarrow 2)- β -D-GlcA	-	[26]
Soyasapogenol B	α -L-Rha(1 \rightarrow 2)- β -D-Glu(1 \rightarrow 2)- β -D-GlcA	-	[26]
Soyasapogenol B	α -L-Rha(1 \rightarrow 2)- β -D-Gal(1 \rightarrow 2)- β -D-GlcA	-	[26]
Soyasapogenol E	α -L-Rha(1 \rightarrow 2)- β -D-Gal(1 \rightarrow 2)- β -D-GlcA	-	[26]
Soyasapogenol E	α -L-Rha(1 \rightarrow 2)- β -D-Gal(1 \rightarrow 2)- β -D-GlcA	22- <i>O</i> -maltol	[35]

A very interesting structural feature of these substances, is the presence of an aldehyde group at the C-23 position in 2 β ,3 β -dihydroxy-23-oxo-olean-12-en-28-oic acid (Figure 1), a new aglycone of saponins from *M. arborea* and *M. hybrida*. This metabolite might in fact represent an interesting

biosynthetic intermediate in the oxidative steps that lead from a methyl group to the corresponding carboxylic acid [1, 75]. That is, if we consider the following genins found in the genus *Medicago*: 2 β -hydroxyoleanolic acid, bayogenin, 2 β ,3 β -dihydroxy-23-oxo-olean-12-en-28-oic acid and medicagenic

Table 7: Saponins identified in *M. truncatula* leaves and roots.

Aglycone	3 OH substituted	28 COOH substituted	Ref.
Hederagenin	GlcA	-	[40]
Hederagenin	Glc-Ara	Glc	[40]
Medicagenic acid	Glc	-	[40]
Medicagenic acid	Glc-malonyl	-	[40]
Medicagenic acid	Glc-Glc	-	[40]
Medicagenic acid	Glc	Glc	[40]
Medicagenic acid	Glc-malonyl	Glc	[40]
Medicagenic acid	β -GlcA	β -Glc	[44]
Medicagenic acid	β -Glc	β -Xyl(1 \rightarrow 4)- α -Rha(1 \rightarrow 2)- α -Ara	[42, 44]
Medicagenic acid	β -GlcA	β -Xyl(1 \rightarrow 4)- α -Rha(1 \rightarrow 2)- α -Ara	[42, 44]
Medicagenic acid	β -Glc(1 \rightarrow 3)- β -Glc	α -Rha(1 \rightarrow 2)- α -Ara	[44]
Medicagenic acid	β -Glc(1 \rightarrow 3)- β -Glc	β -Xyl(1 \rightarrow 4)- α -Rha(1 \rightarrow 2)- α -Ara	[44]
Medicagenic acid	β -Glc(1 \rightarrow 3)- β -Glc	α -Ara(1 \rightarrow 3)-[β -Xyl(1 \rightarrow 4)]- α -Rha(1 \rightarrow 2)- α -Ara	[44]
Medicagenic acid	β -Glc(1 \rightarrow 3)- β -Glc	β -Api-(1 \rightarrow 3)-[β -Xyl(1 \rightarrow 4)]- α -Rha(1 \rightarrow 2)- α -Ara	[44]
Zanhic acid	β -Glc	β -Xyl(1 \rightarrow 4)- α -Rha(1 \rightarrow 2)- α -Ara	[44]
Zanhic acid	β -Glc(1 \rightarrow 3)- β -Glc	α -Rha(1 \rightarrow 2)- α -Ara	[44]
Zanhic acid	β -Glc(1 \rightarrow 3)- β -Glc	α -Rha[4-Ac](1 \rightarrow 2)- α -Ara	[44]
Zanhic acid	β -Glc(1 \rightarrow 3)- β -Glc	β -Xyl(1 \rightarrow 4)- α -Rha(1 \rightarrow 2)- α -Ara	[44]
Zanhic acid	β -Glc(1 \rightarrow 3)- β -Glc	α -Ara(1 \rightarrow 3)- α -Rha(1 \rightarrow 2)- α -Ara	[44]
Zanhic acid	β -Glc(1 \rightarrow 3)- β -Glc	β -Api(1 \rightarrow 3)- α -Rha(1 \rightarrow 2)- α -Ara	[44]
Zanhic acid	β -Glc(1 \rightarrow 3)- β -Glc	α -Ara(1 \rightarrow 3)-[β -Xyl(1 \rightarrow 4)]- α -Rha(1 \rightarrow 2)- α -Ara	[44]
Zanhic acid	β -Glc(1 \rightarrow 3)- β -Glc	β -Api-(1 \rightarrow 3)-[β -Xyl(1 \rightarrow 4)]- α -Rha(1 \rightarrow 2)- α -Ara	[44]
Zanhic acid	Glc-Glc-Glc	Xyl-Rha-Ara, 23 COOH substituted: Ara	[42]
Zanhic acid	Glc-Glc-Glc	Api-Xyl-Rha-Ara, 23 COOH substituted: Ara	[42]
Soyasapogenol B	α -Rha(1 \rightarrow 2)- β -Gal(1 \rightarrow 2)- β -GlcA	-	[40, 43]
Soyasapogenol B	α -Rha(1 \rightarrow 2)- β -Xyl(1 \rightarrow 2)- β -GlcA	-	[43]
Soyasapogenol E	α -Rha(1 \rightarrow 2)- β -Gal(1 \rightarrow 2)- β -GlcA	-	[40, 43]

acid, all the oxidative products at C-23 can be observed. The above genins all possess the same stereochemistry (2 β ,3 β) in the hydroxylated triterpene carbons with the different functional groups at the C-23 position. The presence of an aldehyde group in 2 β ,3 β -dihydroxy-23-oxo-olean-12-en-28-oic acid, identified for the first time in *Medicago* spp, indicates a possible biosynthetic pathway for the sapogenins of this genus. Accordingly, medicagenic acid may originate from bayogenin by subsequent oxidative enzymatic steps involving the formation of 2 β ,3 β -dihydroxy-23-oxo-olean-12-en-28-oic acid while bayogenin may originate by a selective oxidative demethylation at C-23 from 2 β -hydroxyoleanolic acid. In a similar way, the two 16 α -hydroxy triterpenes found in this genus, caulophyllogenin and zanhic acid (Figure 1) probably originate by enzymatic oxidation of hederagenin and medicagenic acid, respectively. The biosynthesis of these compounds in the genus *Medicago* has never been extensively investigated, and only a few papers have been published [76-78].

Saponin extracts as well as purified saponins from selected species of *Medicago* have different biological properties [10-14]. Their antimicrobial, insecticidal, allelopathic and cytotoxic effects are described below. The influence of saponins on animal metabolism is also reviewed.

Antimicrobial activity

Saponins are likely to be implicated in plant defense mechanisms against microbial or fungal infections. In some plants wounding of tissues in response to a pathogenic attack causes the hydrolysis of saponins to derivatives with strong antibiotic activity [79, 80]. On the other hand, resistance to infestation by certain fungi in plants such as oat is associated with the specific presence of saponins (e.g. avenacins). Nevertheless, antifungal efficacy of saponins has been demonstrated *in vitro* for a number of plant species [2, 7, 79, 81] but little data is available on their antibacterial activity [2].

A compilation of microorganisms used to assess antifungal and antibacterial activity of saponins from *Medicago* spp. is reported in Table 8. Data derive from incubation of *Medicago* dry meals, saponin extracts and purified saponins from different species and plant organs.

Antifungal efficacy of *Medicago* has been primarily studied with the model fungus *Trichoderma viride* [10, 29, 31, 63, 65, 82-85] which appeared particularly sensitive to the presence of saponins in the growth medium. A bioassay to determine the content of saponins in the plant was developed [58] based on saponin toxicity towards this fungus. Growth of *T. viride* was in fact found inversely correlated with the amount of *Medicago* saponins in the incubation medium thus representing a useful index to evaluate the total percentage of these metabolites.

As described (Table 8), saponins from *Medicago* have been assayed *in vitro* against phytopathogenic species and their activity well established not only against specific pathogens of *Medicago*, but also against some fungi generally pathogenic to cereals [10, 14, 28, 31, 65, 85-95]. A higher antifungal activity was found for the saponins from the roots than from the aerial parts of *M. sativa* [92, 93, 95]. Furthermore, assays with purified saponins from the same species [93] indicated that the growth of the two pathogens *Botrytis tulipae* and *Phloma narcissi* was mostly affected by the following compounds: medicagenic acid; 3-*O*- β -D-glucopyranosyl-medicagenic acid; 3-*O*- β -D-glucopyranosyl-28-*O*-[β -D-xylopyranosyl(1 \rightarrow 4)- α -L-rhamopyranosyl(1 \rightarrow 2)- α -L-arabinopyranoside] medicagenic acid and 3-*O*- β -D-glucuronopyranosyl-28-*O*-[β -D-xylopyranosyl(1 \rightarrow 4)- α -L-rhamopyranosyl(1 \rightarrow 2)- α -L-arabinopyranoside] medicagenic acid. The screening of saponin extracts from several *Medicago* spp. has shown that *M. arabica* possesses antifungal efficacy several times higher than that of *M. sativa* [95] and the most sensitive pathogens were *Rhizoctonia solani*, *B. tulipae*, *P. narcissi*, *Fusarium oxysporium* ssp. *tulipae* and *Pestalotia* ssp.

Besides their phytopathogenic potential the antimicrobial activity of saponins from *Medicago* against human pathogens has also been investigated [14, 96-104]. Preliminary studies have concerned the effect against some yeasts and dermatophytes of a gluco derivative of medicagenic acid named G2 and later identified as 3-*O*- β -D-glucopyranosyl-

medicagenic acid. The compound was found to be particularly effective against *Cryptococcus neoformans* with an MFC of 4 μ g/mL [2, 7, 79, 80]. More recent investigations [104] on dermatophytes have shown that *Trichophyton interdigitale* and *Microsporium gypseum* were susceptible to *Medicago* saponins especially to glycosides of medicagenic acid, such as 3-*O*- β -glucopyranoside (MIC < 62.5 μ g/mL) the most bioactive phytochemical.

The study of the antifungal activity of saponins from *M. sativa*, *M. arborea* and *M. arabica* against a selection of medically important yeasts (*Candida albicans*, *C. tropicalis*, *Saccharomyces cerevisiae*, *Cryptococcus laurentii* and *Blastomyces capitatus*) [103] has shown that *S. cerevisiae* was the most susceptible, being highly inhibited when treated with the sapogenin mixtures from the aerial parts of the three different species of *Medicago* (MICs of 125, 62.5 and 175 μ g/mL for *M. sativa*, *M. arabica* and *M. arborea*, respectively). A very low MIC value (42.5 μ g/mL) was observed when the same strain was treated with medicagenic acid, which represents the dominant aglycone found in *M. sativa* (50%) and *M. arborea* (30%) aerial organs. Medicagenic acid also inhibited the two mycetes *C. tropicalis* and *B. capitatus*, with an MIC of 125 μ g/mL.

Although strongly antifungal, saponins are reported to have only weak or no growth inhibitorial effects against bacteria [80]. To the best of our knowledge only one investigation has been carried out to evaluate the antibacterial activity of saponins from *Medicago* species and they were found not very active (MICs > 500 μ g/mL) against Gram negative bacteria [103]. Nevertheless, they displayed some efficacy against selected Gram positive bacteria [103]. In particular, sapogenins obtained on acid hydrolysis of saponins from *M. arabica* aerial parts and roots were the most effective, showing good growth inhibitorial activity towards three different strains of *S. aureus*, two strains of *E. faecalis*, and against *B. subtilis* and *B. cereus* (Table 8). *In vitro* antibacterial assays with purified aglycones from *Medicago* saponins showed that medicagenic acid had significant activity against *S. aureus* (MIC 52.5 μ g/mL) and two strains of *E. faecalis* (MICs 50 and 32.5 μ g/mL)

The *in vitro* effects of some saponins from *M. sativa* on rhizosphere bacteria suspension, showed that 3-*O*- β -D-glucopyranosylmedicagenic acid sodium salt

Table 8: Overview of antimicrobial studies with saponins from *Medicago* spp.

Saponin source	Microorganisms	Ref.
a) Phytopathogenic fungi		
<i>M. sativa</i> leaves Total saponins	<i>Fusarium oxysporum</i> , <i>F. solani</i> , <i>Phytophthora drechsleri</i> <i>Phoma</i> sp., <i>Rhizoctonia solani</i> , <i>Verticillium albo-atrum</i>	[65]
<i>M. sativa</i> tops <i>M. sativa</i> roots Total saponins	<i>Alternaria solani</i> <i>Pythium myriotylum</i> <i>P. butleri</i> , <i>P. sp. PRL2142</i> , <i>Sclerotium rolfsii</i>	[86, 87]
<i>M. sativa</i> roots Total extract Total saponins Saponin sugars Sapogenins	<i>Rhizoctonia solani</i>	[88]
<i>M. sativa</i> roots 3- <i>O</i> - β -D-Glc Medicagenic acid	<i>Aspergillus niger</i> , <i>Fusarium oxysporum</i> sp. <i>Lycopersici</i> , <i>Phytophthora cinnamomi</i> , <i>Rhizopus mucco</i> , <i>Sclerotium rolfsii</i>	[82]
<i>M. sativa</i> roots 3- <i>O</i> -[α -D-Glc(1 \rightarrow 4)- β -D-Glc] Medicagenic acid (Medicagenic acid β -maltoside)	<i>Aspergillus niger</i> <i>Fusarium oxysporum</i> sp. <i>Lycopersici</i> , <i>Pythium aphanidermatum</i> , <i>Rhizoctonia solani</i> , <i>Sclerotium rolfsii</i>	[29]
<i>M. sativa</i> roots Compound G2 (3- <i>O</i> - β -D-Glc Medicagenic acid)	<i>Aspergillus niger</i> <i>Fusarium oxysporum</i> , <i>Geotrichum candidum</i> <i>Pythium aphanidermatum</i> , <i>Phytophthora cinnamomi</i> <i>Rhizoctonia solani</i> , <i>Rhizopus mucco</i> , <i>Sclerotium rolfsii</i>	[14]
Meal from aerial parts of: <i>M. arabica</i> , <i>M. doliata</i> , <i>M. heyneana</i> , <i>M. murex</i> , <i>M. sativa</i> Total saponins from aerial parts of: <i>M. arabica</i> , <i>M. heyneana</i> , <i>M. murex</i> , <i>M. polymorpha</i> , <i>M. sativa</i>	<i>Cephalosporium gramineum</i>	[89, 91, 94]
<i>M. sativa</i> roots Total saponins Total prosapogenins 3- <i>O</i> - β -D-Glc Medicagenic acid Medicagenic acid		
Meal from aerial parts of: <i>M. arabica</i> , <i>M. doliata</i> , <i>M. heyneana</i> , <i>M. murex</i> , <i>M. sativa</i>	<i>Gaeumannomyces graminis</i> v. <i>tritici</i>	[90, 91]
<i>M. sativa</i> roots Total saponins Total prosapogenins 3- <i>O</i> - β -D-Glc Medicagenic acid Medicagenic acid		
<i>M. sativa</i> aerial parts <i>M. sativa</i> roots Total saponins	<i>Alternaria zinniae</i> , <i>Botrytis cinerea</i> , <i>B. tulipae</i> , <i>Phoma narcissi</i> , <i>P. poolensis</i> , <i>Rhizoctonia solani</i>	[92]
<i>M. sativa</i> leaves and roots 3- <i>O</i> -[α -L-Ara(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- α -L-Ara]-28- <i>O</i> - β -D-Glc Hederagenin 3- <i>O</i> - β -D-Glc Medicagenic acid 3- <i>O</i> - β -D-Glc-28- <i>O</i> - β -D-Glc Medicagenic acid 3- <i>O</i> - β -D-Glc-28- <i>O</i> -[β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara] Medicagenic acid 3- <i>O</i> - β -D-GlcAc-28- <i>O</i> -[β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara] Medicagenic acid 3- <i>O</i> -[β -D-Glc(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- β -D-Glc]-23- α -L-Ara-28- <i>O</i> - [β -D-Api(1 \rightarrow 3)- β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara] Zanhic acid Soyasaponin I Hederagenin, Medicagenic acid, Soyasapogenol B	<i>Botrytis tulipae</i> <i>Phoma narcissi</i>	[93]

Table 8 (Contd.)

<i>M. arabica</i> shoots Total saponins	<i>Alternaria tenui</i> , <i>Botrytis cinerea</i> , <i>B. tulipae</i> , <i>Fusarium oxysporium</i> sp. <i>Callistephi</i> , <i>F. oxysporium</i> sp. <i>Narcissi</i> , <i>F. oxysporium</i> sp. <i>Tulipae</i> , <i>Pestalotia ssp.</i> , <i>Phoma narcissi</i> , <i>P. poolensis</i> , <i>Pythium ultimum</i> , <i>Rhizoctonia solani</i> , <i>Stangospora curtisii</i>	[95]
<i>M. sativa</i> Compound G2 (3- <i>O</i> - β -D-Glc Medicagenic acid)	b) Human pathogenic fungi <i>Candida albicans</i> , <i>C. guilliermondii</i> , <i>C. krusei</i> , <i>C. parapsilopsis</i> , <i>C. pseudotropicalis</i> , <i>C. tropicalis</i> , <i>Cryptococcus neoformans</i> , <i>Epidermophyton floccosum</i> , <i>Geotrichum candidum</i> , <i>Microsporium canis</i> , <i>Rhodotorula glutinis</i> , <i>Torulopsis candida</i> , <i>T. glabrata</i> , <i>Trycophyton mentagrophytes</i> , <i>T. mentagrophytes</i> var. <i>granulare</i> , <i>Trichopyton rubrum</i> , <i>T. tonsurans</i>	[14, 96-100]
3- <i>O</i> - β -D-Glc Medicagenic acid	<i>Scopulariopsis brevicaulis</i> , <i>Trycophyton mentagrophytes</i>	[101, 102]
<i>M. arabica</i> tops Total saponins, Sapogenins, Bayogenin	<i>Blastomyces capitatus</i> , <i>Candida albicans</i> , <i>C. tropicalis</i> , <i>Cryptococcus laurentii</i> , <i>Saccharomyces cerevisiae</i>	[103]
<i>M. arabica</i> roots Total saponins, Sapogenins		
<i>M. arborea</i> tops Total saponins, Prosapogenins, Sapogenins		
<i>M. sativa</i> tops Total saponins, Prosapogenins, Sapogenins Medicagenic acid, Hederagenin		
<i>M. sativa</i> roots Total saponins, Sapogenins		
<i>Medicago</i> sp. 3- <i>O</i> - α -L-Ara-Hederagenin 3- <i>O</i> -[α -L-Ara(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- α -L-Ara]-Hederagenin 3- <i>O</i> - β -D-Glc-Medicagenic acid 3- <i>O</i> - β -D-Glc-28- <i>O</i> - β -D-Glc-Medicagenic acid 3- <i>O</i> - β -D-GlcAc-28- <i>O</i> - β -D-Glc-Medicagenic acid 3- <i>O</i> -[β -D-Glc(1 \rightarrow 2)- β -D-Glc]-28- <i>O</i> - β -D-Glu-Medicagenic acid 3- <i>O</i> - β -D-Glc-28- <i>O</i> -[β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara]-Medicagenic acid 3- <i>O</i> - β -D-GlcAc-28- <i>O</i> -[β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara]-Medicagenic acid 3- <i>O</i> -[β -D-Glc(1 \rightarrow 2)- β -D-Glc]-28- <i>O</i> -[β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara]-Medicagenic acid 3- <i>O</i> -[β -D-Glc(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- β -D-Glc]-28- <i>O</i> -[β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara]-Zanhic acid Soyasaponin I Hederagenin, Medicagenic acid	<i>Microsporium gypseum</i> , <i>Trichophyton interdigitale</i>	[104]
<i>M. arabica</i> tops Total saponins, Sapogenins, Bayogenin	c) Bacteria <i>Acinebacter baumanii</i> , <i>Bacillus subtilis</i> , <i>B. cereus</i> , <i>Enterococcus faecalis</i> , <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Staphylococcus aureus</i>	[103]
<i>M. arabica</i> roots Total saponins, Sapogenins, Hederagenin		
<i>M. sativa</i> roots Total saponins, Sapogenins		
<i>M. arborea</i> tops Total saponins, Prosapogenins, Sapogenins		
<i>M. sativa</i> tops Total saponins, Prosapogenins, Sapogenins, Medicagenic acid		
<i>M. sativa</i> 3- <i>O</i> - β -D-Glc-28- <i>O</i> - β -D-Glc Medicagenic acid Soyasaponin I 3- <i>O</i> - β -D-Glc Medicagenic acid Na ⁺ salt Medicagenic acid Na ⁺ salt	d) Soil Bacteria <i>Agrobacterium tumefaciens</i> , <i>Bacillus thuringiensis</i> , <i>Curtobacterium flacumafaciens</i> , <i>Pseudomonas fluorescens</i>	[105]

Table 8 (Contd.)

e) Others	
Total saponins from aerial parts of:	
<i>M. aculeata</i> , <i>M. arabica</i> , <i>M. blanchiana</i> , <i>M. carstiensis</i>	[10, 29,
<i>M. ciliaris</i> , <i>M. coerulea</i> , <i>M. coronata</i> , <i>M. disciformis</i>	31, 51,
<i>M. doliata</i> , <i>M. falcata</i> , <i>M. glutinosa</i> , <i>M. granadensis</i>	63, 65,
<i>M. hemicycla</i> , <i>M. heyneana</i> , <i>M. hybrida</i> , <i>M. intertexta</i>	82-85]
<i>M. laciniata</i> , <i>M. lupulina</i> , <i>M. minima</i> , <i>M. murex</i>	
<i>M. muricoleptis</i> , <i>M. noeana</i> , <i>M. orbicularis</i> , <i>M. polyceratia</i>	
<i>M. polymorpha</i> , <i>M. praecox</i> , <i>M. radiata</i> , <i>M. rigidula</i>	
<i>M. rotata</i> , <i>M. rugosa</i> , <i>M. sativa</i> , <i>M. sauvagei</i> , <i>M. scutellata</i>	
<i>M. soleirolia</i> , <i>M. tornata</i> , <i>M. turbinata</i> , <i>M. truncatula</i>	
<i>M. sativa</i> roots	
Total saponins	
3- <i>O</i> -[α -L-Ara(1 \rightarrow 2)- β -D-Glc(1 \rightarrow 2)- α -L-Ara] Hederagenin	
3- <i>O</i> - β -D-Glc Medicagenic acid	
3- <i>O</i> - β -D-GlcAc Medicagenic acid	
3- <i>O</i> -[α -D-Glc(1 \rightarrow 4)- β -D-Glc] Medicagenic acid (Medicagenic acid β -maltoside)	
3- <i>O</i> - β -D-Glc-28- <i>O</i> - β -D-Glc-Medicagenic acid	
3- <i>O</i> - β -D-Glc-28- <i>O</i> -[β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara] Medicagenic acid	
3- <i>O</i> - β -D-GlcAc-28- <i>O</i> -[β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara] Medicagenic acid	
3- <i>O</i> -[β -D-Glc(1 \rightarrow 2)- β -D-Glc]-28- <i>O</i> -[β -D-Xyl(1 \rightarrow 4)- α -L-Rha(1 \rightarrow 2)- α -L-Ara] Medicagenic acid	
<i>M. lupulina</i> roots	
3- <i>O</i> - β -D-Glc Medicagenic acid	
3- <i>O</i> - β -D-Glc-28- <i>O</i> - β -D-Glc Medicagenic acid	
Medicagenic acid	

could negatively affect them and, could negatively affect them and, in most cases, its activity corresponded to that of the corresponding aglycone, medicagenic acid disodium salt [105].

Investigations on the structure-activity relationships of *Medicago* saponins have led to contrasting results [11, 86, 106]. The number, kind and sequence of the sugar residues in the molecules have been differently correlated with their antimicrobial effects. A detailed study on the activity of different saponins from alfalfa roots against *T. viride* indicated that the monodesmoside derivatives of medicagenic acid were more active than the related bidesmosides, even though no straight correlation between the number of sugars in the molecule and its bioactivity could be established [31]. Moreover, the antifungal activity of medicagenic acid and its derivatives were reported as dependent on the presence of functional groups, such as carboxy and hydroxy in the molecule. In some studies, reduction of bioactivity was related to the presence of a sugar moiety at the 28-*O*-position of the saponin [11, 86, 106].

Bioassays with saponins from *Medicago* sp. against human pathogenic fungi and bacteria indicated that

the sugar moieties are not required for antimicrobial activity [103]. This study proved that saponins were more active than the related prosapogenins and saponins.

Insecticidal activity

Toxicity of saponins to insects is known, and it has been suggested that they might also provide plant protection from insect predation [107]. To support this hypothesis, the herbivore-induced response of alfalfa was recently examined through assays with *Spodoptera littoralis* larvae, and it was observed that the levels of total saponins increases in the young foliage of damaged plants [108].

Several works on the insecticidal and antifeedant properties of saponins against several classes of insects have been published [109, 110]. A list of saponin source and related insects and pests on which they have been tested, is reported in Table 9. Saponins from alfalfa roots and shoots were reported to be active against the peach aphid (*Myzus persicae*) [109], and found to be toxic to the larvae of the grass grub (*Costelytra zealandica*) [111]. Several species of locusts have shown increasing mortality when fed

on alfalfa; their larvae developed more slowly and the emerging adults were smaller than when they were fed saponin-free herbage [109]. Alfalfa root saponins, rich in medicagenic acid, are toxic to the flour beetle (*Tribolium castaneum*) and their toxicity increased when some of the sugars were removed by hydrolysis [109]. On the contrary, it has been described that several alfalfa pests, such as alfalfa weevil (*Hypera postica*), spotted aphid (*Tereosaphid maculata*), clover root curculio (*Stoma hispidulus*), and seed chalcid (*Bruchophagus rodii*) are hardly affected by a saponin-rich diet, suggesting that they have evolved strategies to overcome the toxicity of the saponins of the plant on which they prey [109].

Crude mixtures and purified saponins from alfalfa leaves were tested against potato leafhopper (*Empoasca fabae* Harris) and pea aphid (*Acyrtosiphon pisum* Harris). Larvae were fed with a diet containing 0.01-5.0% saponins for a few days. An increase of mortality was observed for all the tested organisms, in particular for those fed on saponins containing medicagenic acid [112].

Saponins extracted from the leaves of 41 alfalfa varieties, with a different content of saponins and sapogenins, were assayed *in vitro* against larvae of the yellow mealworm (*Tenebrio molitor* L.). Results showed a good correlation between larvae mortality and saponin concentration so this biological assay was proposed to detect alfalfa saponins in plant material [66, 113].

Alfalfa saponin mixtures also have been tested against the summer fruit tortrix moth (*Adoxophyes orana* F.v.R.), the European grape moth (*Lobesia botrana* Den. & Schiff.) and the European corn borer (*Ostrinia nubilalis* Hb.). The increasing amount of saponins added to the diet (from 1 to 1000 ppm) increased larval mortality from 11.3% at 1 ppm to 46.1% at 1000 ppm. The contact effect accounted for a maximum of 22.7% mortality. No appreciable differences were detected in the insecticidal activity exerted by crude saponins derived from alfalfa leaves and roots [12].

Saponins isolated from the aerial parts of alfalfa were tested against the Colorado potato beetle (*Leptinotarsa decemlineata* Say). Larvae were fed on potato leaves sprayed with 0.5 and 1% saponin solutions; no repellent effects were observed for any of the tested compounds, but insect feeding proved to be less intense on saponin-treated leaves. The larvae

fed on saponin treated leaves had the lowest body weight gain, suggesting the antifeedant activity of the compounds. The insect mortality from eating saponin-treated leaves was 100% at both tested concentrations [114]. Other experiments showed that the larvae of Colorado potato beetles reared on potato leaves treated with a 0.5% solution of total saponins from *M. sativa* roots and tops, died after 4-6 days because of fasting. Lower saponin doses (from 0.1 to 0.001%) reduced the insect feeding less causing an inhibition in growth and an extension of the larval stage. Mortality was reached at a level of 76.7-100%. No evident differences have been found in saponin activity from the tops or the roots of alfalfa [115].

The Colorado potato beetle was also used to differentiate insecticidal activity of saponins from *M. arabica*, *M. hybrida* and *M. murex* roots and tops. Total saponins were included in the insect diet as a solution applied on potato leaves on which larvae were reared. All saponins reduced larval feeding, growth rate and mortality in a dose dependant manner. All the saponins showed a high insecticidal activity at the concentration of 0.5%. Saponins from *M. murex* roots and from *M. arabica* and *M. hybrida* aerial parts were found to be the most active, probably due to the differences in their saponin composition [116].

Crude alfalfa root saponins, their prosapogenins produced by alkaline hydrolysis, and medicagenic acid sodium salt, were tested in field trials against spider mite (*Tetranychus urticae* Koch.) and hop aphid (*Phoron humuli* Schrank). Plants were sprayed with a 0.1 and 0.2% solution of saponin products. Prosapogenins were the most active against both phytophages, while crude saponins and medicagenic acid sodium salt were less active [117].

Dried alfalfa leaf and root tissues incorporated in an artificial diet to give the final saponin concentration of 0.1, 0.5 or 1.6% mg/g fresh weight, a cholesterol-precipitable saponin fraction from the plant leaves and a total saponin mixture from the roots were used to evaluate their toxic potential against the polyphagous insect european corn borer. The growth and development of larvae were significantly inhibited after feeding. Root saponins were somewhat more harmful than saponins from the tops [118].

Total saponins from *M. sativa* roots and leaves and individual saponins and sapogenins were tested on a polyphagous pest, the army-worm *Spodoptera*

Table 9: List of pests and insects used to evaluate the insecticidal activity of saponins from the *Medicago* spp.

Saponin source	Insect	Ref.
<i>M. sativa</i> tops	Grass grub (<i>Costelytra zealandica</i>)	[111]
Total saponins	Pea aphid (<i>Acyrtosiphon pisum</i> Harris)	[112]
	Potato leafhopper (<i>Empoasca fabae</i> Harris)	
	Yellow mealworm (<i>Tenebrio molitor</i>)	[66, 113]
	Colorado potato beetle (<i>Leptinotarsa decemlineata</i> Say)	[114]
<i>M. sativa</i> tops	Alfalfa weevie (<i>Hypera postica</i>)	[109]
<i>M. sativa</i> roots	Clover root curculio (<i>Stona hispidulus</i>)	
Total saponins	Flour beetle (<i>Tribolium castraneum</i>)	
	Peach aphid (<i>Myrus persicae</i>)	
	Seed chalcid (<i>Bruchophagus rodoli</i>)	
	Spotted aphid (<i>Thereoaphid maculata</i>)	
	Several species of locusts	
	European corner borer (<i>Ostrinia nubilalis</i> Hb.)	[12]
	European grape moth (<i>Lobesia botrana</i> Den. & Schiff.)	
	Summer fruit tortrix moth (<i>Adoxophyes orana</i> F.v.R.)	
	Colorado potato beetle (<i>Leptinotarsa decemlineata</i> Say)	[115]
<i>M. arabica</i> tops	Colorado potato beetle (<i>Leptinotarsa decemlineata</i> Say)	[116]
<i>M. arabica</i> roots		
Total saponins		
<i>M. hybrida</i> tops		
<i>M. hybrida</i> roots		
Total saponins		
<i>M. murex</i> tops		
<i>M. murex</i> roots		
Total saponins		
<i>M. sativa</i> roots	Hop aphid (<i>Phoron humuli</i> Schrank)	[117]
Total saponins	Spider mite (<i>Tetranychus urticae</i> Koch.)	
Prosapogenins		
Medicagenic acid Na ⁺ salt		
<i>M. sativa</i> tops	European corner borer (<i>Ostrinia nubilalis</i> Hb.)	[118]
Total saponins		
Cholesterol-precipitable saponins		
<i>M. sativa</i> roots		
Total saponins		
<i>M. sativa</i>	Army-warm (<i>Spodoptera littoralis</i> Boisd.)	[119]
3- <i>O</i> -β-D-Glc-28- <i>O</i> -[α-L-Ara(1→2)-β-D-Glc(1→2)-α-L-Ara]		
Hederagenin		
3- <i>O</i> -β-D-Glc Medicagenic acid		
3- <i>O</i> -β-D-Glc-28- <i>O</i> -β-D-Glc Medicagenic acid		
3- <i>O</i> -β-D-Glc-28- <i>O</i> -[β-D-Xyl(1→4)-α-L-Rha(1→2)-α-L-Ara]		
Medicagenic acid		
3- <i>O</i> -β-D-GlcAc-28- <i>O</i> -[β-D-Xyl(1→4)-α-L-Rha(1→2)-α-L-Ara] Medicagenic acid		
Soyasaponin I,		
Hederagenin, Medicagenic acid, Soyasapogenol A,		
Soyasapogenol B, Soyasapogenol E,		
3- <i>O</i> -β-D-Glc Medicagenic acid Na ⁺ salt		
3- <i>O</i> -β-D-Glc-28- <i>O</i> -β-D-Glc Medicagenic acid Na ⁺ salt		
Soyasaponin I Na ⁺ salt, Medicagenic acid Na ⁺ salt		

littoralis. Total saponins (1, 10 and 100 ppm) and a series of pure saponins (10 ppm) and sapogenins (20 ppm) were given in the food and their effects examined during larval development as well as in the resulting pupae and adults. At 1 ppm, root saponins caused a nearly 70% mortality and the emerged

females exhibited about 60% fertility reduction. Total saponins from the aerial parts were less active, although the increase of mortality and the reduction of fecundity were significant. All the pure saponins lowered the food consumption and reduced the larval growth rate although to a different extent. Aglycones

influenced the larval development in a similar way, medicagenic acid was found to be the most active, hederagenin and soyasapogenols A and B exhibited only moderate activity, while soyasapogenol E was inactive. Medicagenic acid sodium salt and its 3-*O*- β -D-glucopyranosyl and medicagenic acid 3-*O*- β -D-glucopyranosyl-28-*O*- β -D-glucopyranoside derivatives were the most active substances. Additionally, all the tested α -L-arabinopyranosyl glycosides were inactive, while the corresponding aglycones or glycosides were active. Based on those results it has been suggested that glycosylated saponins are bioactive only when they are hydrolyzed by insect gut glycosidases and release an active aglycone; complex glycosides containing arabinopyranosyl units apparently resist the action of the enzymes [119].

The spraying of winter wheat with different saponin concentrations at various phenological phases had no negative effects on growth parameters, grain yield and quality of wheat flour. Results indicated that saponins (0.01-0.1%) can be applied on a wheat crop as fungicides or insecticides [120].

Allelopathic effects

Alfalfa, as other forage legumes, has the reputation as an important rotation crop to improve nitrogen availability in the soil. In some cases the increase of nitrogen by *Medicago* does not correspond to an increase of grain yield in the succeeding rotated crop, suggesting that some factors might interfere with the utilization of nitrogen. This effect has been experimentally correlated with the presence, in alfalfa plant material, of saponins which display allelopathic activity [11, 13, 121-124].

The role of alfalfa saponins as allelopathic agents was first reported in 1955 by Mishutin and Naumova [125] who observed that growth of cotton was influenced by the use of alfalfa as a rotation crop. Detrimental effects on cotton-seed germination was also shown in *in vitro* assays with alfalfa saponins. Later investigations have shown that saponins from various species of *Medicago* act as allelochemicals, some with a defined specificity towards different plants [126-130]. The allelopathic potential of medicagenic acid glycosides has been noted. Depending on their concentration they may function as plant growth inhibitors (high concentrations) or

stimulators (low concentrations) [121, 129, 130]. They also inhibit the growth of several weeds and cereals [11, 13, 121, 128-130]: elongation of roots and shoots of *Bromus secalimus* and *Echinochloa crus-galli* was inhibited by 10 ppm saponin (19-11 and 28-17%, respectively), while growth of wheat roots was 50% reduced at the concentration of 100 ppm compared to the control [129, 130]. Other saponins such as soyasapogenol B and hederagenin glycosides were in general found less active as growth inhibitors than medicagenic acid derivatives [121, 129].

The different allelopathic potential of *Medicago* species was related to their different content of saponins. Thus, for example, saponins (medicagenic and soyasapogenol glycosides) isolated from the seeds of *M. lupulina* were able to inhibit the growth of the cereals oat, barley, wheat and rye, whereas saponins from the seeds of *M. sativa* (containing only soyasapogenol glycosides) had no effects on wheat and rye development, but only on that of barley and oat [121, 129]. The use of plant material of various physiological ages indicated that alfalfa at immature stages is more phytotoxic since it likely contains higher amounts of allelochemicals [126].

Soil texture also was found to influence the inhibitory activity of alfalfa saponins [121, 127]. In a detailed study it has been in fact shown that finely powdered alfalfa roots in sandy soil causes a more pronounced detrimental effect on wheat growth than incorporation in heavy clay soils.

An autotoxic effect of alfalfa also has been reported [131]. Despite their allelopathic activity, however, saponins produced by the plant species seem not to be involved in the autotoxic effects which instead have been attributed to water-soluble phenolic components [132-135]. In particular the isoflavonoids medicarpin and its methoxy analogue, 4-methoxymedicarpin, and chlorogenic acid purified from alfalfa leaves were assayed in *in vitro* experiments and found to contribute to the plant autoallelopathy which results in a yield decrease, low seed germination and poor growth when alfalfa is sown in soils where the same species was previously cropped.

The physiological mechanism of action of saponins as allelopathic agents is not clear. Inhibition of seed germination has been correlated with a decrease in

oxygen diffusion through the seed coat [121], while seedling growth retardation has not been well studied. Comparison of the allelopathic effects of structurally different saponins from *Medicago* species revealed some structure-activity relationships. As found for antimicrobial activity, monodesmosides were in general more active than the related bi- and tridesmosides while medicagenic acid glycosides having glucose at the C-3 position were more active than similar compounds substituted with glucuronic acid, and zanhic acid glycosides were more effective than the 3-*O*-glucuronides of medicagenic acid [11, 121, 129].

Effects on animals

The significance of natural saponins in animal nutrition has been widely investigated [136, 137]. Studies of the effects of the saponins from *Medicago* species have been carried out only for *M. sativa*, due to the importance of this species as forage and as an industrial source of leaf protein concentrate used in animal diets. An excellent review on this topic is available [138]. Saponins may have significant effects on all the phases of animal metabolism from ingestion to excretion. Alfalfa saponins influence rumen fermentation and affect microbial protein synthesis in the rumen, the site of nutrient digestion. Moreover they suppress fermentation in rumen cultures [139], and *in vivo* investigations [140] have confirmed a general decrease of fermentation associated with a symptomatic decrease of volatile fatty acids and cellulose digestion. A significant reduction of protozoa in rumen of sheep receiving alfalfa saponins was also reported [140]. Moreover, endogenous bacteria appeared morphologically modified when treated with alfalfa saponins [139]. All these effects on animal nutrition have been related to the ability of saponins, or their aglycones, to interact with cell membrane sterols and other metabolites [138, 141]. Saponins are in fact able to complex cholesterol, and their anti-nutritional effects were lowered by addition of cholesterol to the diet. Retardation of growth by alfalfa dietary saponins has been observed in livestock and laboratory animals, probably due to the bitter and astringent sensory characteristics of the processed grain products. One mechanism that might account for the growth depressing effects of saponins is the lowering of feed intake because of unpalatability.

No clear information is available on the lethal dose or minimum inhibition concentration of alfalfa saponins towards livestock. Animal species differ in their susceptibility to saponins, however. Poultry are more sensitive than other farm animals. A variety of alfalfa with 1.47% of saponins caused an average reduction of 11% in weight gain of chicks compared to a low-saponin variety containing 0.59% of the active compounds [142, 143]. No effects were reported when calves were fed with alfalfa hay containing up to 2.62% saponin [144]. Though accurate estimates of detrimental saponin levels are lacking, high- and low-saponin germplasm has been defined in the literature (and set as a goal in breeding programs), mostly based on responses of monogastric animals, or biological assays (e.g. *Trichoderma viride* test and the hemolytic test). Conventionally an average concentration of about 2.0% and 0.8% were considered to be high and low, respectively [143, 145].

Determination of saponins by semi-quantitative methods based on biological assays may give erroneous results. For instance, glycosides of zanhic acid are weakly detectable by biological tests, although they are classified as toxic/moderately toxic compounds, with an LD₅₀ value of 562 mg/kg body weight calculated for hamsters [146]. Sensory test trials on human volunteers, using saponins isolated from alfalfa aerial parts, showed that zanhic acid tridesmoside is the most bitter, astringent and throat-irritating compound of all the tested saponins [36]. This compound is also reported to have the highest intestinal membrane depolarizing activities compared to other alfalfa saponins [147]. It also has been described as causing breathing problems and nervous system perturbations to hamsters, followed by death after 24h. Bloat syndromes were observed at necropsy [146].

Rats fed alfalfa saponins at levels of 1% in the diet for up to 26 weeks showed no toxic effects; a potentially beneficial reduction of serum cholesterol and triglycerides was observed instead [148]. No adverse reactions have been detected in the non-human primate, *Macaca fascicularis*, following consumption of a mixture of alfalfa top saponins for up to 78 weeks. The metabolites decreased cholesterolemia without changing the level of high density lipoprotein-cholesterol; hence, they reduced the total cholesterol/high density lipoprotein-

cholesterol ratio. Furthermore, saponins decreased intestinal adsorption of cholesterol, and increased excretion of neutral steroids and bile acids [149]. As these compounds interact with cholesterol and directly interfere with its absorption, a possible application in some human pathologies can be hypothesized, although toxicity of alfalfa saponins for human consumption needs detailed investigation.

Furthermore, *in vitro* studies indicated that saponins from *M. sativa* roots and aerial parts have some effects on pancreatic lipase activity. Results showed that they stimulated lipolytic activity and did not influence the proteolytic and amylolytic activities of Neopancreatum, a mixture of porcine pancreatic enzymes such as trypsin, chymotrypsin, lipase and amylase. An increase of the stimulatory effects of saponins was observed when sodium cholate was added to the medium [150].

Cytotoxic and tumor-promoter inhibitory activities

Although the cytotoxicity of triterpenoid saponins is known [151], saponins from *Medicago* species have never been extensively investigated, although saponins from *M. sativa* leaves showed dose-dependant growth inhibition *in vitro* of human leukemic cell line K562 [12]. No significant effects on clonogenic survival were observed when purified saponins from *M. sativa* roots, leaves and seeds were tested *in vitro* against MCF7 human breast carcinoma cells and HeLa human cervical carcinoma cells, although MCF7 was more sensitive to the treatment. Inhibition of tumoral cell growth was instead observed when saponins were used in association with cis-platin. The growth of MCF7 cells was 18-33% (saponin concentration 25 µg/mL; cis-platin 4 µg/mL), compared to 40% survival when only cis-platin was used. Root and seed saponins were found to be more active than saponins from leaves. All the tested saponins enhanced the cis-platin induced toxicity, although HeLa cells were significantly less affected [152]. As reported [153], saponins seem to act as promoters probably affecting cell membrane permeability cis-platin diffusion in the cells.

The lack of information on the cytotoxicity of saponins from *Medicago* spp. does not allow additional indications of their activity, but their antitumoral, chemopreventive and antimutagenic

properties can be extrapolated from those of bioactive saponins from other plants but found in *Medicago* spp. For example, saponins from soybean, including soyasaponin I (soyasapogenol B 3-*O*- α -L-rhamopyranosyl(1 \rightarrow 2)- β -D-glucopyranosyl(1 \rightarrow 2)- β -D-glucuronopyranosyl) found in almost all the studied *Medicago* species, are known for their chemopreventive properties [154-156]. Orally consumed soybean saponins are not adsorbed in the small intestine and appear to reach the colon [157] where they exert their beneficial effects. They are able to suppress the growth of human colon carcinoma cells *in vitro* [158, 159], and to inhibit the chemically induced colonic aberrant crypt formation in CF1 mice [160]. Soyasaponin I from *W. brachybotrys* has also been shown to strongly inhibit mouse skin tumor promotion [161].

Antimutagenic and antiproliferative [162-164] activity has also been observed for some hederagenin monodesmosides from *Hedera helix*, including 3-*O*- α -L-arabinopyranosyl hederagenin and 3-*O*- α -L-rhamopyranosyl(1 \rightarrow 2)- α -L-arabinopyranosyl hederagenin in *M. arabica* and *M. polymorpha*, respectively.

The saponin 3-*O*- α -L-rhamopyranosyl(1 \rightarrow 2)- β -D-glucopyranosyl (1 \rightarrow 2) - β -D- glucuronopyranosyl soyasapogenol E from *Wistaria brachybotrys* (Leguminosae), named wistaria saponin D, showed antitumor promoting activity [165]. The same sapogenin has also been found in *M. sativa* and *M. truncatula*.

In this context, the mutagenic activity of some saponins and sapogenins from *M. sativa* have been evaluated. Soyasaponin I, in a concentration up to 500 µg, medicagenic acid (up to 200 µg) and its 3-*O*-glucopyranosyl derivative (up to 200 µg), were tested according to the Ames assay against *Salmonella typhimorium* strains TA97, TA98, TA100 and TA102. Results showed that saponins did not increase the number of his⁺ revertants in any of the strains, neither in the absence nor in the presence of metabolic activation (S9 fraction from rat liver) [166].

Conclusion

Saponins from the *Medicago* genus are a complex group of pentacyclic triterpene glycosides which

display antimicrobial, insecticidal, allelopathic and cytotoxic properties, together with antinutritional effects. Particularly studied *M. sativa*, the most important species within the genus from an agronomic point of view.

The biological activities of *Medicago* saponins are related to their chemical structure in that monodesmosides are more active than the

corresponding bidesmosides, and the aglycone and the nature and position of the sugar in the molecule might be important factors in determining their efficacy.

Based on their bioactivity, plant saponins are already used commercially. Data summarized here might suggest further applications of saponins from *Medicago*.

References

- [1] Dewick PM. (2001) *Medicinal Natural Products – A Biosynthetic approach*. Wiley, New-York.
- [2] Hostettmann K, Marston A. (1995) *Chemistry and Pharmacology of Natural Products. Saponins*. Phillipson JD, Baxter H (Eds). Cambridge University Press, UK.
- [3] Milgate J, Roberts DCK. (1995) The nutritional and biological significance of saponins. *Nutritional Research*, **8**, 1223-1249.
- [4] Lacaille-Dubois MA, Wagner H. (2000) Bioactive saponins from plants: an update. In *Studies in Natural Products Chemistry*. Vol. **21**, Atta-ur-Rahman (Ed). Elsevier, London, UK. 633-687.
- [5] Francis G, Keem Z, Makkar HPS, Becker K. (2002) The biological action of saponins in animal systems. A review. *British Journal of Nutrition*, **88**, 587-605.
- [6] Sprag SG, Light ME, van Staden J. (2004) Biological activities and distribution of plant saponins. *Journal of Ethnopharmacology*, **94**, 219-243.
- [7] Tanaka O, Tamura Y, Masuda H, Mizutani K. (1996) Application of saponins in food and cosmetics: saponins of *Mohave yucca* and *Sapindus mukurossi*. In *Advances in Experimental Medicine and Biology. Saponins Used in Food and Agriculture*. Vol. **405**. Waller GR, Yamasaki K. (Eds). Plenum Press, New-York. 1-11.
- [8] San Martin R, Briones R. (2000) Quality control of commercial quillaja (*Quillaja saponaria* Molina) extracts by reverse phase HPLC. *Journal of the Science of Food and Agriculture*, **88**, 587-605.
- [9] Bruneton J. (1999) *Pharmacognosy – Phytochemistry, Medicinal Plants*. Lavoisier Publishing, Paris, 671-719.
- [10] Jurzysta M, Waller GR. (1996) Antifungal and haemolytic activity of aerial parts of alfalfa (*Medicago*) species in relation to saponin composition. In *Advances in Experimental Medicine and Biology. Saponins Used in Traditional and Modern Medicine*. Vol. **404**. Waller GR, Yamasaki K. (Eds). Plenum Press, New-York. 565-574.
- [11] Oleszek W. (1996) Alfalfa saponins: structure, biological activity and chemotaxonomy. In *Advances in Experimental Medicine and Biology. Saponins used in Food and Agriculture*. Vol. **405**. Waller GR, Yamasaki K. (Eds). Plenum Press, New-York. 155-170.
- [12] Tava A, Odoardi M. (1996) Saponins from *Medicago* spp.: chemical characterization and biological activity against insects. In *Advances in Experimental Medicine and Biology. Saponins Used in Food and Agriculture*. Vol. **405**. Waller GR, Yamasaki K. (Eds). Plenum Press, New-York. 97-109.
- [13] Timbekova AE, Isaev MI, Abubakirov NK. (1996) Chemistry and biological activity of triterpenoid glycosides from *Medicago sativa*. In *Advances in experimental medicine and biology. Saponins Used in Food and Agriculture*. Vol. **405**, Waller GR, Yamasaki K (Eds). Plenum Press, New York USA. 171-182.
- [14] Zehavi U, Polacheck I. (1996) Saponins as antimycotic agents: glycosides of medicagenic acid. In *Advances in Experimental Medicine and Biology. Saponins Used in Traditional and Modern Medicine*. Vol. **404**. Waller GR, Yamasaki K. (Eds). Plenum Press, New-York. 535-546.
- [15] Oleszek W. (2000) Alfalfa saponins: Chemistry and Application. In: *Phytochemicals as Bioactive Agents*. Technomic Publ. Comp., Inc., USA, 167-188.
- [16] Lesins KA, Lesins I. (1979) *Genus Medicago (Leguminosae). A taxogenetic Study*. Junk W. (Ed), Publisher, London.
- [17] Heyn CC. (1963) *The Annual Species of Medicago*. Magnes Press, Hebrew University, Jerusalem.
- [18] Potter GC, Kummerow FA. (1954) Chemical similarity and biological activity of the saponins isolated from alfalfa and soybeans. *Science*, **120**, 224-225.
- [19] Morris RJ, Dye WB, Gisler DS. (1961) Isolation, purification and structural activity of an alfalfa root saponins. *Journal of Organic Chemistry*, **26**, 1241-1243.
- [20] Morris RJ, Hussey EW. (1965) A natural glycoside of medicagenic acid. An alfalfa blossom saponin. *Journal of Organic Chemistry*, **30**, 166-168.

- [21] Gestetner B. (1971) Structure of a saponin from lucerne (*Medicago sativa*). *Phytochemistry*, **10**, 2221-2223.
- [22] Timbekova AE, Abubakirov NK. (1984) Triterpene glycosides from alfalfa. I. Medicoside G. A novel bidesmoside from *Medicago sativa*. *Khimiya Prirodnikh Soedinenii*, 451-458.
- [23] Timbekova AE, Abubakirov NK. (1985) Triterpene glycosides from alfalfa. II. Medicoside C. *Khimiya Prirodnikh Soedinenii*, 805-808.
- [24] Timbekova AE, Abubakirov NK. (1986) Triterpene glycosides from alfalfa. VII. Medicoside I. *Khimiya Prirodnikh Soedinenii*, 607-610.
- [25] Timbekova AE, Abubakirov NK. (1986) Triterpene glycosides of alfalfa. IV. Medicoside J. *Khimiya Prirodnikh Soedinenii*, 610-613.
- [26] Kitagawa I, Taniyama T, Murakami T, Yoshihara M, Yoshikawa M. (1988) Saponin and sapogenol. XLVI. On the constituents of aerial part of american alfalfa, *Medicago sativa* L. The structure of dehydrosoyasaponin I. *Yakugaku Zasshi*, **108**, 547-551.
- [27] Massiot G, Lavaud C, Le Men-Olivier L, van Binst G, Miller SPF, Fales HM. (1988) Structural elucidation of alfalfa root saponins by mass spectrometry and nuclear magnetic resonance analysis. *Journal of Chemical Society Perkin Transaction I*, 3071-3079.
- [28] Oleszek W, Price KR, Fenwick GR. (1988) Triterpene saponins from the roots of *Medicago lupulina* L. (black medic trefoil). *Journal of Science of Food and Agriculture*, **43**, 289-297.
- [29] Levy M, Zehavi U, Naim M, Polacheck I. (1989) Isolation, structure determination and antifungal activity of a new alfalfa root saponin. *Carbohydrate Research*, **193**, 115-123.
- [30] Timbekova AE, Larin MF, Yagudaev MR, Abubakirov NK. (1989) Triterpene glycosides of alfalfa. V. Medicoside H. *Khimiya Prirodnikh Soedinenii*, 673-677.
- [31] Oleszek W, Price KR, Colquhoun IJ, Jurzysta M, Ploszynski M, Fenwick GR. (1990) Isolation and identification of alfalfa (*Medicago sativa* L.) root saponins: their activity in relation to a fungal bioassay. *Journal of Agricultural and Food Chemistry*, **38**, 1810-1817.
- [32] Timbekova AE, Verechagin AL, Semenov AA, Abubakirov NK. (1990) Triterpene glycosides of alfalfa. VI. Medicoside L. *Khimiya Prirodnikh Soedinenii*, 221-227.
- [33] Mahato SB. (1991) Triterpenoid saponins from *Medicago hyspida*. *Phytochemistry*, **30**, 3389-3393.
- [34] Massiot G, Lavaud C, Besson V, Le Men-Olivier L, van Binst G. (1991) Saponins from aerial parts of alfalfa (*Medicago sativa*). *Journal of Agricultural and Food Chemistry*, **39**, 78-82.
- [35] Massiot G, Lavaud C, Benkhaled M, Le Men-Olivier L. (1992) Soyasaponin VI, a new maltol conjugate from alfalfa and soyabean. *Journal of Natural Products*, **55**, 1339-1341.
- [36] Oleszek W, Jurzysta M, Ploszynski M, Colquhoun IJ, Price KR, Fenwick GR. (1992) Zanic acid tridesmoside and other dominant saponins from alfalfa (*Medicago sativa* L.) aerial parts. *Journal of Agricultural and Food Chemistry*, **40**, 191-196.
- [37] Timbekova AE, Shashkov AS, Abubakirov NK. (1993) Triterpene glycosides from alfalfa. VII. Medicosides E and F. *Khimiya Prirodnikh Soedinenii*, 701-705.
- [38] Kinjo J, Uemura H, Nakamura M, Nohara T. (1994) Two new triterpenoidal glycosides from *Medicago polymorpha* L. *Chemical and Pharmacological Bulletin*, **42**, 1339-1341.
- [39] Bialy Z, Jurzysta M, Oleszek W, Piacente S, Pizza C. (1999) Saponins in alfalfa (*Medicago sativa* L.) root and their structural elucidation. *Journal of Agricultural and Food Chemistry*, **47**, 3185-3192.
- [40] Huhman DV, Sumner LW. (2002) Metabolic profiling of saponins in *Medicago sativa* and *Medicago truncatula* using HPLC coupled to an electrospray ion-trap mass spectrometer. *Phytochemistry*, **59**, 347-360.
- [41] Zbigniew B, Jurzysta M, Mella M, Tava A. (2004) Triterpene saponins from aerial parts of *Medicago arabica* L. *Journal of Agricultural and Food Chemistry*, **52**, 1095-1099.
- [42] Huhman DV, Berhow MA, Sumner LW. (2005) Quantification of saponins in aerial and subterranean tissues of *Medicago truncatula*. *Journal of Agricultural and Food Chemistry*, **53**, 1914-1920.
- [43] Kapusta I, Bogdan J, Stochmal A, Oleszek W. (2005) Determination of saponins in aerial parts of barrel medic (*Medicago truncatula*) by liquid chromatography-electrospray ionization/mass spectrometry. *Journal of Agricultural and Food Chemistry*, **53**, 7654-7660.
- [44] Kapusta I, Stochmal A, Perrone A, Piacente S, Pizza C, Oleszek W. (2005) Triterpene saponins from barrel medic (*Medicago truncatula*) aerial parts. *Journal of Agricultural and Food Chemistry*, **53**, 2164-2170.
- [45] Tava A, Mella M, Avato P, Argentieri MP, Bialy Z, Jurzysta M. (2005) Triterpene saponins from leaves of *Medicago arborea* L. *Journal of Agricultural and Food Chemistry*, **53**, 9954-9965.
- [46] Bialy Z, Jurzysta M, Mella M, Tava A. (2006) Triterpene Saponins from the Roots of *Medicago hybrida*. *Journal of Agricultural and Food Chemistry*, **54**, 2520-2526.
- [47] Pedersen MW. (1975) Relative quantity and biological activity of saponins in germinated seeds, roots, and foliage of alfalfa. *Crop Science*, **15**, 541-543.

- [48] Gorski PM, Miersch J, Ploszynski M. (1991) Production and biological activity of saponins and canaverine in alfalfa seedlings. *Journal of Chemical Ecology*, **17**, 1135-1143.
- [49] Oleszek W. (1998) Composition and quantitation of saponins in alfalfa (*Medicago sativa* L.) seedlings. *Journal of Agricultural and Food Chemistry*, **46**, 960-962.
- [50] Berrang B, Davis KH Jr, Wall ME, Hanson CH, Pedersen ME. (1974) Saponins of two alfalfa cultivars. *Phytochemistry*, **13**, 2253-2260.
- [51] Quazi HM (1975) Effect of cultivar and season on the concentration of saponins in lucerne (*Medicago sativa* L.). *New Zealand Journal of Agricultural Research*, **18**, 227-232.
- [52] Tava A, Oleszek W, Jurzysta M, Berardo N, Odoardi M. (1993) Alfalfa saponins and sapogenins: isolation and quantification in two different cultivars. *Phytochemical Analysis*, **4**, 269-274.
- [53] Tava A, Pecetti L. (1998) Hemolytic activity and saponin content in lucerne (*Medicago sativa* complex) genotypes. *Journal of Genetics and Breeding*, **52**, 33-37.
- [54] Tava A, Odoardi M, Oleszek W. (1999) Seasonal changes of saponin content in five alfalfa (*Medicago sativa*) cultivars. *Agricoltura Mediterranea*, **129**, 111-116.
- [55] Pecetti L, Tava A, Romani M, De Benedetto MG, Corsi P. (2006) Variety and environment effects on the dynamics of saponins in lucerne (*Medicago sativa* L.). *European Journal of Agronomy*, **25**, 187-192.
- [56] Jurzysta M, Burda S, Oleszek W, Ploszynski M. (1988) The chemotaxonomic significance of larcytrin and medicagenic acid in the tribe Trigonellae. *Canadian Journal of Botany*, **66**, 363-367.
- [57] Jurzysta M, Small E, Nozzolillo C. (1988) The evolution of hemolytic saponin content in wild and cultivated alfalfa (*Medicago sativa*, Fabaceae). *Economic Botany*, **44**, 226-235.
- [58] Small E, Jurzysta M, Nozzolillo C. (1990) Hemolysis, a sinapomorphic discriminator for an expanded genus *Medicago* (Leguminosae). *Taxon*, **37**, 354-363.
- [59] Jurzysta M, Burda S, Oleszek W, Ploszynski M. (1992) Chemical composition of seed saponins as a guide to the classification of *Medicago* species. *Canadian Journal of Botany*, **70**, 1384-1387.
- [60] Walter ED, Van Atta GR, Thompson CR, Maclay WD. (1954) Alfalfa saponins. *Journal of the American Chemical Society*, **76**, 2271-2273.
- [61] Jurzysta M. (1973) Isolation and chemical characterization of saponins from lucerne seeds (*Medicago media* Pers.) *Acta Societatis Botanicorum Poloniae*, **42**, 201-207.
- [62] Gorsky PM, Jurzysta M, Burda S, Oleszek W, Ploszynsky M. (1984) Studies on *Medicago lupulina* saponins. 2. Isolation, chemical characterization and biological activity of saponins from *Medicago lupulina* tops. *Acta Societatis Botanicorum Poloniae*, **53**, 527-533.
- [63] Jurzysta M, Nowacki E. (1979) Saponins in the genus *Medicago*. *Acta Agrobotanica*, **32**, 13-17.
- [64] Jurzysta M. (1979) Haemolytic micromethod for rapid estimation of toxic alfalfa saponins. *Acta Agrobotanica*, **32**, 5-11.
- [65] Zimmer DE, Pederson MW, McGuire CF. (1967) A bioassay for alfalfa saponins using the fungus *Trichoderma viride* Pers. *Crop Science*, **7**, 223-224.
- [66] Pracros P. (1988) Mesure de l'activité des saponins de la lucerne par les larves du ver de la farine: *Tenebrio molitor* L. (Coléoptère, *Tenebrionidae*). I. Comparaison avec les résultats de divers tests biologiques. *Agronomie*, **8**, 257-263.
- [67] Nowacka J, Oleszek W. (1994) Determination of alfalfa (*Medicago sativa* L.) saponins by high performance liquid chromatography. *Journal of Agricultural and Food Chemistry*, **42**, 727-729.
- [68] Jurzysta M, Jurzysta A. (1978) Gas-liquid chromatography of trimethylsilyl ethers of soyasapogenols and medicagenic acid. *Journal of Chromatography*, **148**, 517-520.
- [69] Rao D, Boris G. (1987) Simple gas chromatographic method for the determination of medicagenic acid in alfalfa (*Medicago sativa*). *Journal of Chromatography*, **410**, 169-175.
- [70] Tava A, Chiari M, Oleszek W. (2000) Separation of alfalfa (*Medicago sativa* L.) saponins as their borate complexes by capillary electrophoresis. In *Saponins in Food, Feedstuffs and Medicinal Plants*. Chapter 5, Oleszek W, Marston A (Eds). Kluwer Academic Publishers, Dordrecht The Netherlands. 43-56.
- [71] Lee MK, Ling YC, Jurzysta M, Waller GR. (1996) Saponins from alfalfa, clover, and mungbeans analysed by electrospray ionization-mass spectrometry as compared with positive and negative FAB-mass spectrometry. In *Advances in Experimental Medicine and Biology: Saponins Used in Food and Agriculture*. Vol 405, Waller GR, Yamasaki K (Eds). Plenum Press, New York USA. 353-364.
- [72] Agrawal PK. (1996) A systematic NMR approach for the determination of the molecular structure of steroidal saponins. In *Advances in Experimental Medicine and Biology: Saponins Used in Food and Agriculture*. Vol 405, Waller GR, Yamasaki K (Eds). Plenum Press, New York USA. 299-315.
- [73] Mella M, Jurzysta M, Ricci M, Tava A. (2002) NMR investigation of saponins and sapogenins from *Medicago* species. *Proceedings of XXXII National Congress on Magnetic Resonance*. Pavia, Italy P-3.

- [74] Tava A, Mella M, Bialy Z, Jurzysta M. (2003) Stability of saponins in alcoholic solutions: ester formation as artifacts. *Journal of Agricultural and Food Chemistry*, **51**, 1797-1800.
- [75] Torsell, KBG. In *Natural Product Chemistry - A Mechanistic, Biosynthetic and Ecological Approach*. Swedish Pharmaceutica Society, 1997.
- [76] Morris RJ, Tankersley DL. (1963) The synthesis of the β -D-glucoside of medicagenic acid, an alfalfa root saponin. *The Journal of Organic Chemistry*, **28**, 240-242.
- [77] Nowacki E, Jurzysta M, Dietrych-Szostak D. (1976) Zur biosynthese der medicagensäure in keimenden luzernesamen. *Biochemie und Physiologie der Pflanzen*, **156**, 183-186.
- [78] Suzuki H, Achnine L, Xu R, Matsuda SPT, Dixon RA. (2002) A genomic approach to the early stages of triterpene saponin biosynthesis in *Medicago truncatula*. *The Plant Journal*, **32**, 1033-1048.
- [79] Osbourn A. (1996) Saponins and plant defence - a soap story. *Trends in Plant Science*, **1**, 4-9.
- [80] Osbourn A. (2003) Saponins in cereals. *Phytochemistry*, **62**, 1-4.
- [81] Wina E, Muetzel S, Becker K. (2005) The impact of saponin-containing plant materials on ruminant production-A review. *Journal of Agricultural and Food Chemistry*, **53**, 8093-8105.
- [82] Levy M, Zehavi U, Naim M, Polacheck I. (1986) An improved procedure for the isolation of medicagenic acid 3-O- β -D-Glucopyranoside from alfalfa roots and its antifungal activity on plant pathogens. *Journal of Agricultural and Food Chemistry*, **34**, 960-963.
- [83] Nonaka M. (1986) Variable sensitivity of *Trichoderma viride* to *Medicago sativa* saponins. *Phytochemistry*, **25**, 73-75.
- [84] Gruiz K. (1996) Fungitoxic activity of saponins: practical use and fundamental principles. In *Advances in Experimental Medicine and biology. Saponins Used in traditional and modern medicine*. Vol. **404**. Waller GR, Yamasaki K. (Eds). Plenum Press, New-York. 527-534.
- [85] Jurzysta M, Bialy Z. (1999) Antifungal and haemolytic activity of roots of alfalfa (*Medicago* spp.) in relation to saponin composition. In *Modern fungicides and antifungal compounds II*. Lyr H, Russel PE, Sisler HD (Eds). Intercept: Andover, UK. 445-451.
- [86] Gestetner B, Assa Y, Henis Y, Birk Y, Bondi A. (1971) Lucerne saponins IV.- Relationship between their chemical constitution, and haemolytic and antifungal activities. *Journal of Agricultural and Food Chemistry*, **22**, 168-172.
- [87] Assa Y, Gestetner B, Chet I, Henis Y. (1972) Fungistatic activity of Lucerne saponins and digitonin as related to sterols. *Life Sciences*, **11**, Part II, 637-647.
- [88] Levanon D, Henis Y, Okon Y, Dovrat A. (1982) Alfalfa saponins and microbial transformations of nitrogen in peat. *Soil Biology & Biochemistry*, **14**, 501-504.
- [89] Martyniuk S, Jurzysta M, Bialy Z, Wróblewska B. (1995) Alfalfa root saponins affect the growth and sporulation of *Cephalosporium gramineum*. In *Environmental Biotic Factors in Integrated Plant Disease Control*. Manka M (Ed). The Polish Phytopathological Society, Poznan. 395-398.
- [90] Martyniuk S, Wróblewska B, Jurzysta M, Bialy Z. (1995) Saponins as inhibitors of cereal pathogens: *Gaeumannomyces graminis* v. *tritici* and *Cephalosporium gramineum*. In *Modern Fungicides and Antifungal Compounds*. Lyr H, Russel PE, Sisler HD (Eds.) Intercept Ltd., UK. 193-197.
- [91] Martyniuk S, Jurzysta M, Wróblewska B. (1999) Influence of powdered aerial parts of various *Medicago* species on the growth of *Gaeumannomyces graminis* and *Cephalosporium gramineum*. *Bulletin of the Polish Academy of Sciences, Biological Sciences*, **47**, 2-4.
- [92] Saniewska A, Jurzysta M, Bialy Z. (2001) Differential antifungal activity of alfalfa (*Medicago sativa* L.) saponins originated from roots and aerial parts of some ornamental plant pathogens. *Acta Agrobotanica*, **54**, 31-43.
- [93] Saniewska A, Jurzysta M, Bialy Z. (2003) The effect of alfalfa (*Medicago sativa*) saponins on *Botrytis tulipae* and *Phoma narcissi* growth. *Phytopathologica Polonica*, **27**, 15-27.
- [94] Martyniuk S, Bialy Z, Jurzysta M. (2004) Antifungal activity of aerial parts and saponins of *Medicago* ssp. against *Cephalosporium gramineum*. *Allelopathy Journal*, **13**, 75-82.
- [95] Saniewska A, Jarecka Z, Bialy Z, Jurzysta M. (2005) Antifungal activity of saponins from *Medicago arabica* L. shoots against some pathogens. *Allelopathy Journal*, **16**, 105-112.
- [96] Polacheck I, Zehavi U, Naim M, Levy M, Evron R. (1986) Activity of compound G2 isolated from alfalfa roots against medically important yeasts. *Antimicrobial Agents and Chemotherapy*, **30**, 290-294.
- [97] Polacheck I, Zehavi U, Naim M, Levy M, Evron R. (1986) The susceptibility of *Cryptococcus neoformans* to an antimycotic agent (G2) from alfalfa. *Zentralblatt für Bakteriologie, Mikrobiologie und Hygiene. Series A*, **261**, 481-486.
- [98] Evron R, Polacheck I, Guizie M, Levy M, Zehavi U. (1988) Activities of compound G2 isolated from alfalfa roots against dermatophytes. *Antimicrobial Agents and Chemotherapy*, **32**, 1586-1587.
- [99] Evron R, Guizie M, Zehavi U, Polacheck I. (1990) Activity of compound G2 isolated from alfalfa roots in experimental dermatophyte infection. *Antimicrobial Agents and Chemotherapy*, **34**, 1600-1601.

- [100] Polacheck I, Zehavi U, Naim M, Levy M, Evron R. (1986) The susceptibility of *Cryptococcus neoformans* to an antimycotic agent (G2) from alfalfa. *Zentralblatt für Bakteriologie, Mikrobiologie und Hygiene. Series A*, **261**, 481-486.
- [101] Spiewak R, Szostak W, Jurzysta M, Bialy Z. (2000) Inhibiting action of medicagon acid 3-glucoside on the growth of *Scropulariopsis brevicaulis* in vitro. *Abstract Book*, 10th International Mycological Symposium of PDS – Mycology 2000, Poland, pg 113.
- [102] Spiewak R, Szostak W, Jurzysta M, Bialy Z. (2000) Inhibiting action of medicagon acid 3-glucoside on the growth of *Trichophyton mentagrophytes* in vitro. *Abstract Book*, 10th International Mycological Symposium of PDS – Mycology 2000, Poland, pg 114.
- [103] Avato P, Bucci R, Tava A, Vitali C, Rosato A, Bialy Z, Jurzysta M. (2006) Antimicrobial activity of saponins from *Medicago* sp.: structure –activity relationship. *Phytotherapy Research*, **20**, 454-457.
- [104] Houghton PJ, Patel N, Cheung CY, Jurzysta M, Bialy Z. (2005) Antifungal activity of saponins from *Medicago* species. *Abstract Book*, 53rd GA Annual Congress, Florence, p. 192.
- [105] Hoagland R. (2001) Effects of alfalfa saponins on in vitro physiological activity of soil and rhizosphere bacteria. *Journal of Crop Protection*, **4**, 349-361.
- [106] Oleszek W. (2000) Saponins. In *Natural Food Antimicrobial Systems*. Naidu AS (Ed). CRC Press, London, 1-30.
- [107] Applebaum SW, Marco S, Birk Y. (1969) Saponins as a possible factor of resistance of legume seed to attack of insects. *Journal of Agricultural and Food Chemistry*, **17**, 618-621.
- [108] Agrelli J, Oleszek W, Stochmal A, Olsen M, Anderson P. (2003) Herbivore-induced responses in alfalfa (*Medicago sativa*). *Journal of Chemical Ecology*, **29**, 303-320.
- [109] Applebaum SW, Birk Y. (1979) In *Herbivores. Their Interaction with Secondary Plant Metabolites*. Rosenthal GA, Janzen DH (Eds). Academic Press, New York. 539-566.
- [110] Oakenfull D, Sidhu GS. (1989) Saponins. In *Toxicant of Plant Origin. Volume II. Glycosides*. Cheeke PR (Ed), CRC Press, Boca Raton, Florida (USA), 97-141.
- [111] Sutherland ORW, Hutchins RFN, Greenfield WJ. (1982) Effect of lucerne saponins and lotus condensed tannins of survival of grass grub, *Costelytra zealandica*. *New Zealand Journal of Zoology*, **9**, 511-516.
- [112] Horber E, Leath KT, Berrang B, Macarian V, Hanson CH. (1974) Biological activities of saponin components from DuPuits and Lahontan alfalfa. *Entomology Experimental & Application*, **17**, 410-424.
- [113] Pracros P. (1988) Mesure de l'activité des saponines de la luzerne par les larves du ver de farine: *Tenebrio molitor* L. (Coléoptère, *Tenebrionidae*). II – Recherche des fractions de saponines responsables des effets antinutritionnelles observés. *Agronomie*, **8**, 793-799.
- [114] Waligora D, Krzymanska J. (1994) The influence of secondary plant substances: glucosinolates, alkaloids and saponins on the feeding of Colorado potato beetle (*Leptinotarsa decemlineata* Say). *Proc. XXXIV Session IOR*, vol. **II**, Poznan, pp. 9-12.
- [115] Szczepanik M, Krystkowiak K, Jurzysta M, Bialy Z. (2001) Biological activity of saponins from alfalfa tops and roots against colorado potato beetle larvae. *Acta Agrobotanica*, **54**, 35-45.
- [116] Szczepanik M, Bialy Z, Jurzysta M. (2004) The insecticidal activity of saponins from various *Medicago* spp. against Colorado potato beetle, *Leptinotarsa decemlineata* Say. *Allelopathy Journal*, **14**, 177-186.
- [117] Puszkar L, Jastrzebski A, Jurzysta M, Bialy Z. (1994) Alfalfa saponins as a chance in the integrated hop protection. *Proc. XXXIV Session IOR*, vol. **II**, Poznan, pp. 255-259.
- [118] Nozzolillo C, Arnason JT, Campos C, Donskov N, Jurzysta M. (1997) Alfalfa leaf saponins and insect resistance. *Journal of Chemical Ecology*, **21**, 995-1002.
- [119] Adel MM, Sehnal f, Jurzysta M. (2000) Effects of alfalfa saponins on the moth *Spodoptera littoralis*. *Journal of Chemical Ecology*, **26**, 1065-1078.
- [120] Podolska G, Bialy Z, Jurzysta M, Waller GR. (2003) Effect of application of alfalfa root saponins aqueous solution on the plant structure, yield and quality of winter wheat. *Allelopathy Journal*, **11**, 171-184.
- [121] Waller GR. (1989) Biochemical frontiers of allelopathy. *Biologia Plantarum*, **31**, 418-447.
- [122] Oleszek W, Jurzysta M, Gorski PM. (1992) Alfalfa saponins the allelopathic agents. In *Allelopathy: Basic and Applied Aspects*. Rizvi SJH, Rizvi V. (Eds). Chapman & Hall, London. 151-167.
- [123] Oleszek W. (1993) Allelopathic potential of alfalfa (*Medicago sativa*) saponins: their relation to antifungal and hemolytic activities. *Journal of Chemical Ecology*, **19**, 1063-1074.
- [124] Oleszek W, Hoagland RE, Zablotowicz RM. (1999) Ecological significance of plant saponins. In: *Principles and Practices in Plant Ecology. Allelochemical Interactions*. Inderjit Dakshini KMM, Foy CL (Eds.). CRC Press LLC, Washington DC, 451-465.
- [125] Mishustin BN, Naumova AN. (1955) Secretion of toxic substances by alfalfa and their effect on cotton and soil microflora. *Akademia Nauk USSR Izvestija, Seriya Biologicheskaya*, **6**, 3-9.
- [126] Guenzi WD, Kehr WR, McCalla TM. (1964) Water-soluble phytotoxic substances in alfalfa forage: variation with variety, cutting, year, and stage of growth. *Agronomy Journal*, **56**, 499-500.

- [127] Oleszek W, Jurzysta M. (1987) The allelopathic potential of alfalfa root medicagenic acid glycosides and their fate in soil environments. *Plant and Soil*, **98**, 67-80.
- [128] Wyman-Simpson CL, Waller GR, Jurzysta M, McPherson JK, Young CC. (1991) Biological activity and chemical isolation of root saponins of six cultivars of alfalfa (*Medicago sativa* L.). *Plant and Soil*, **135**, 83-94.
- [129] Waller GR, Jurzysta M, Thorne RLZ. (1993) Allelopathic activity of root saponins from alfalfa (*Medicago sativa* L.) on weeds and wheat. *Botany Bulletin Academia Sinica*, **34**, 1-11.
- [130] Waller GR, Jurzysta M, Thorne RLZ. (1995) Root saponins from alfalfa (*Medicago sativa* L.) and their allelopathic activity on weeds and wheat. *Allelopathy Journal*, **2**, 21-30.
- [131] Hall MH, Hederlong PR. (1989) Alfalfa autotoxic fraction characterization and initial separation. *Crop Science*, **29**, 425-428.
- [132] Miller RW, Kleiman R, Powell RG. (1988) Germination and growth inhibitors of alfalfa. *Journal of Natural Products*, **51**, 328-330.
- [133] Dornbos DL Jr, Spencer GF, Miller RW. (1990) Medicago delays alfalfa seed germination and seedlings growth. *Crop Science*, **30**, 162-166.
- [134] Chung I-M, Seigler D, Miller DA, Kyung S-H. (2000) Autotoxic compounds from fresh alfalfa leaf extracts: identification and biological activity. *Journal of Chemical Ecology*, **26**, 315-327.
- [135] Xuan TD, Tsuzuki E, Terao H, Matsuo M, Khanh TD. (2003) Correlation between growth inhibitory exhibition and suspected allelochemicals (phenolic compounds) in the extract of alfalfa (*Medicago sativa* L.). *Plant Production Science*, **6**, 165-171.
- [136] Price KR, Johnson IT, Fenwick GR. (1987) The chemistry and biological significance of saponins in food and feedstuffs. *CRC Critical Reviews in Food Science and Nutrition*, **26**, 27-135.
- [137] Cheeke PR. (1996) Biological effects of feed and forage saponins and their impacts on animal production. In *Advances in experimental medicine and biology. Saponins Used in Food and Agriculture*. Vol **405**, Waller GR, Yamasaki K (Eds). Plenum Press, New York USA. 377-385.
- [138] Sen S, Makkar HPS, Becker K. (1998) Alfalfa saponins and their implication in animal nutrition. *Journal of Agricultural and Food Chemistry*, **46**, 131-140.
- [139] Lu CD, Jorgensen NA. (1987) Alfalfa saponins affect site and extent of nutrient digestion in ruminants. *Journal of Nutrition*, **117**, 919-927.
- [140] Lu CD, Tsai LS, Schaefer DM, Jorgensen NA. (1987) Alteration of fermentation in continuous culture of mixed rumen bacteria. *Journal of Dairy Science*, **70**, 799-805.
- [141] Wallace RJ, McEwan NR, McIntosh FM, Teferedegne B, Newbold CJ. (2002) Natural products as manipulators of rumen fermentation. *Asian-Australian Journal of Animal Science*, **15**, 1458-1468.
- [142] Pedersen MW, Zimmer DE, McAllister DR, Anderson JO, Wilding MD, Taylor GA, McGuire CF. (1967) Comparative studies of several alfalfa varieties using chemical and biochemical assay. *Crop Science*, **7**, 349-352.
- [143] Pedersen MW, Wang L. (1971) Modification of saponin content of alfalfa through selection. *Crop Science*, **11**, 833-835.
- [144] Howard RE. (1988) Antiquality factors and nonnutritive chemical components. In *Alfalfa and alfalfa Improvements*. Hanson AA, Barnes DK, Hill RR Jr. (Eds). ASA Inc. CSSA Inc. SSSA Inc. Publishers, Madison, USA. 493-514.
- [145] Majak W, Fesser AC, Goplen BP, Pedersen MW. (1980) Relationship between ruminant bloat and composition of alfalfa herbage. II. Saponins. *Canadian Journal of Animal Science*, **60**, 699-708.
- [146] Oleszek W, Nowacka-Zaborska JM, Minta M, Zmudzki J. (1995) Effect of alfalfa saponin-zanhic acid tridesmoside on hamsters (*Mesocricetus auratus*). In *Current trends in Fruit and Vegetable Phytochemistry*. Garcia-Viguera C, Castaner M, Gil MI, Ferreras F, Tomas-Barberan FA (Eds). Consejo Superior Investigaciones Cientificas, Madrid, Spain. 293-297.
- [147] Oleszek W, Nowacka J, Gee JM, Wortley GM, Johnson IT. (1994) Effects of some purified alfalfa (*Medicago sativa*) saponins on transmural potential difference in mammalian small intestine. *Journal of Science of Food and Agriculture*, **65**, 35-39.
- [148] Malinow MR, McLaughlin P, Papworth L, Stafford C, Kohler GO, Livingston AL, Cheeke PR. (1977) Effects of alfalfa saponins on intestinal cholesterol adsorption in rats. *The American Journal of Clinical Nutrition*, **30**, 2061-2067.
- [149] Malinow MR, Connor WE, McLaughlin P, Stafford C, Lin DS, Livingston AL, Kohler GO, McNulty WP. (1981) Cholesterol and bile acid balance in *Macaca fascicularis*: effects of alfalfa saponins. *The Journal of Clinical Investigation*, **67**, 156-162.
- [150] Sroka Z, Jurzysta M, Tylcz J, Rzakowska-Bodalska H. (1997) Stimulation of pancreatic lipase activity by saponins isolated from *M. sativa* L. *Zeitschrift fur Naturforschung*, **52c**, 235-239.
- [151] Bader G, Plohm B, Hiller K, Franz F. (1996) Cytotoxicity of triterpenoid saponins. Part 1: activities against tumor cells *in vitro* and haemolytic index. *Pharmazie*, **51**, 414-417.
- [152] Singuaroli I. (2002) Attività citotossica di saponine da *M. sativa* in linee tumorali umane. Doctoral Thesis. Università degli Studi di Pavia, Facoltà di Farmacia, Dipartimento di Farmacologia Sperimentale ed Applicata.
- [153] Gaidi G, Correia M, Chauffert B, Beltramo JL, Wagner H, Lacaille-Dubois MA. (2002) Saponin-mediated potentiation of cisplatin accumulation and cytotoxicity in human colon cancer cells. *Planta Medica*, **68**, 70-72.

- [154] Messina MJ, Persky V, Setchell KDR, Barnes S. (1994) Soy intake and cancer risk: a review of the *in vitro* and *in vivo* data. *Nutrition and Cancer*, **21**, 113-131.
- [155] Rao AV, Sung MK. (1995) Saponins as anticarcinogens. *Journal of Nutrition*, **125**, 117s-124s.
- [156] Fournier DB, Erdman JW Jr, Gordon GB. (1998) Soy, its components, and cancer prevention: a review of the *in vitro*, animal, and human data. *Cancer Epidemiology Biomarkers Prevention*, **7**, 1055-1065.
- [157] Gestneter B, Birk Y, Tencer Y. (1968) Soybean saponins. Fate of ingested soybean saponins and the physiological aspect of their hemolytic activity. *Journal of Agricultural and Food Chemistry*, **16**, 1031-1035.
- [158] Sung MK, Kendall CWC, Koo MM, Rao AV. (1995) Effect of soybean saponins and gypsophylla saponin on growth and viability of colon carcinoma cells in culture. *Nutrition and Cancer*, **23**, 259-270.
- [159] Oh YJ, Sung MK. (2001) Soybean saponins inhibit cell proliferation by suppressing PKC activation and induce differentiation of HT-21 human colon adenocarcinoma cells. *Nutrition and Cancer*, **39**, 132-138.
- [160] Koratkar R, Rao AV. (1997) Effect of soya bean saponins on azoxymethane-induced preneoplastic lesions in the colon of mice. *Nutrition and Cancer*, **27**, 206-209.
- [161] Konoshima T, Kokumai M, Kozuka M, Tokuda H, Nishino H, Iwashima A. (1992) Anti-tumor-promoting activities of afromosin and soyasaponin I isolated from *Wisteria brachybotrys*. *Journal of Natural Products*, **55**, 1776-1778.
- [162] Elias R, De Meo M, Vidal-Ollivier E, Laget M, Balansard G, Dumenil G. (1990) Antimutagenic activity of some saponins isolated from *Calendula officinalis* L., *C. arvensis* L. and *Hedera helix* L. *Mutagenesis*, **5**, 327-331.
- [163] Danloy S, Quetin-Leclercq J, Coucke P, De Pauw-Gillet MC, Elias R, Balansard G, Angenot L, Bassleer R. (1994) Effects of α -hederin, a saponin extracted from *Hedera helix*, on a cell cultured *in vitro*. *Planta Medica*, **60**, 45-49.
- [164] Park HJ, Kwon SH, Lee JH, Lee KH, Miyamoto KI, Lee KT. (2001) Kalopanaxsaponin A is a basic saponin structure for the anti-tumor activity of hederagenin monodesmosides. *Planta Medica*, **67**, 118-121.
- [165] Konoshima T, Kozuka M, Haruma M, Ito K. (1991) Constituents of leguminous plants, XIII. New triterpenoid saponins from *Wisteria brachybotrys*. *Journal of Natural Products*, **54**, 830-836.
- [166] Czegezot H, Rahden-Staron I, Oleszek W, Jurzysta M. (1994) Isolation and studies of the mutagenic activity of saponins in the Ames test. *Acta Poloniae Pharmaceutica – Drug Research*, **51**, 133-136.

Non-nitrogenous Plant-derived Constituents with Antiplasmodial Activity

Anna Rita Bilia

Department of Pharmaceutical Sciences, University of Florence, via Ugo Schiff, 6,
Sesto Fiorentino-50019-Florence, Italy

ar.bilia@unifi.it

Received: August 2nd, 2006; Accepted: September 27th, 2006


Dedicated to the memory of Professor Ivano Morelli.

The paper is a compilation of the studies reported in the literature concerning non-nitrogenous natural constituents that have shown antiplasmodial activity and aims to provide a basis for further *in vivo* studies as well as for clinical trials to develop new antimalarial agents. Due to the increasingly unsatisfactory outcomes for *N*-heterocyclic drugs, coupled with the rising incidence of the deadly *falciparum* malaria, the advent of non-nitrogenous lead compounds is timely, signaling a new era of antimalarial chemotherapy. Currently a few non-nitrogenous molecules are used in therapy, but many promising molecules of plant origin are under study, such as peroxide sesquiterpenes, quinoid triterpenes, quassinoids, gallic acid derivatives, lignans, flavonoids and biflavonoids, xanthenes, naphthoquinones and phenylanthraquinones. Many of these constituents are isolated from plants used traditionally to treat malaria and fever. Ethnopharmacology can still be considered as a rich source of lead molecules.

Keywords: Plant-derived non-nitrogenous, malaria, *in vitro* and *in vivo* studies, terpenoids, polyphenols.

Malaria is one of the oldest life-threatening parasitic diseases diffused in the tropical regions of the world. It causes more than 300 million acute illnesses and at least 1-2.7 million deaths annually (mainly children under the age of five in sub-Saharan Africa). The majority of malaria deaths are due to cerebral malaria and other complications as a result of malaria-related anemia, and the cost in human life, incapacity for work, programs of control and medical treatment are enormous [1,2]. There are four types of human malaria: *Plasmodium vivax*, *P. falciparum*, *P. malariae*, and *P. ovale*, the first two of which are the most common, and *P. falciparum* is the most deadly type of malaria infection. The malaria situation is aggravated by the appearance of strains of *P. falciparum* resistant to antimalarial drugs as well as by the resistance of vector *Anopheles* mosquitoes to DDT and other insecticides. These are the principal factors that contribute to the difficulty of malaria control and it is unrealistic to think about eradication of this disease by means of destruction of the vector or use of vaccination. Studies in a number


of African countries have shown that the emergence of chloroquine-resistant malaria parasites is associated with a two-fold increase in malaria deaths, but in one study in Mlomp, Senegal it was shown that malaria mortality in children under the age of four increased 11-fold within six years of the emergence of chloroquine-resistance [3]. Thus, chloroquine (**1**) represents one of the most effective anti-malarial drugs, but if used as monotherapy its effectiveness is rapidly lost.



Chloroquine is an analog of quinine (**2**), a natural constituent, which is not only considered as the most important lead molecule for the synthesis of the majority of the existing antimalarial drugs but is also currently used in therapy, especially in severe and

complicated cases of malaria caused by chloroquine-resistant strains of *P. falciparum*.

Quinine was isolated in 1820 from *Cinchona* sp. bark, because the antimalarial properties of these plants had been known for several centuries. Jesuit missionaries in Peru around 1630 discovered that the bark of the cinchona tree allayed fever and a few years later exported the bark to Europe, where it was included in pharmacopoeias to treat fever.




From the 1930s, chemically related molecules such as chloroquine (**1**), mefloquine, amodiaquine, mepaquine and pamaquine were developed [1,2]. These molecules are all characterized by the presence of nitrogen heterocycle moieties, and apart from the diminished effectiveness due to resistance by *P. falciparum*, they generally present some disadvantages and risks for the patient or user. Chloroquine is limited in its geographical use, only working in the Middle East, Mexico and Central America. Mefloquine is expensive, 100 times more so than chloroquine, and has resulted in seizures and psychiatric disorders. Halofantrine is equally expensive, unsuitable for prophylaxis, and has led to cases of cardiotoxicity. Even quinine is never totally effective, and its toxic side effects deter its usage [1,2]. Against this disheartening backdrop of the increasingly unsatisfactory performance of the *N*-heterocyclic drugs coupled with the rising incidence of the deadly *falciparum* malaria, the advent of non-nitrogenous lead compounds was not only timely, but also fortuitous and signalled a new era of antimalarial chemotherapy.

Plant-derived non-nitrogenous antimalarials have made, and continue to make, an immense contribution to malaria chemotherapy. In particular, artemisinin (**3**), isolated from the Chinese plant *Artemisia annua* L., has recently been used successfully against malaria resistant to chloroquine and the aim of this review is to consider the potential of plants to provide new antimalarial treatments.

In the last decades many plant extracts, especially those from species with a reputation for use in


traditional medicines, have been evaluated in the laboratory for their *in vitro* antiplasmodial activities and some have also been tested *in vivo*, usually in mice infected with *P. berghei* or *P. yoelii* [4,5]. In some cases, the constituent(s) responsible for the observed activities have been isolated by bioassay-guided fractionation and their structures elucidated, many of them being non-nitrogenous derivatives.



This paper reports on the antiplasmodial evaluation of these constituents according to their structures: terpenoids, polyphenols and other constituents.

Terpenoids

In the last three decades all the classes of terpenoids have been investigated to evaluate their antimalarial potency.




Among monoterpenoids, a simple molecule, an iridoid related aglycone, compound **4**, isolated from the roots of *Scrophularia lepidota* Boiss. (Scrophulariaceae), showed a low anti-plasmodial activity (the 50% inhibitory concentration (IC₅₀) was 240 μ M). The result was nevertheless interesting because **4** showed a weak FabI enzyme inhibitory activity (IC₅₀=590 μ M): FabI is a key enzyme of *Plasmodium falciparum* fatty acid biosynthesis and it can be used as a novel biological target to be used in the search for novel antiplasmodial constituents [6].


Bioassay-guided fractionation of the dichloromethane extract of the fruits of *Renealmia cinnamata* (Zingiberaceae), whose fruits are widely used in Cameroon to treat fevers, led to the isolation of six sesquiterpenoids of which two known ones, **5** and **6**, were the most active; their IC₅₀ values were 6.8 and 7.4 μ M, respectively, using 3D7 chloroquine-sensitive *P. falciparum* strains [7].

Among sesquiterpenes some lactone derivatives deserve to be mentioned because of the interesting activity [8-11]. A bioassay-guided fractionation of *Neurolaena lobata* (L.) R. Br. (Asteraceae), an

important medicinal plant in Central America and the Caribbean region, where it is used for a variety of diseases including malaria, resulted in the isolation of seven sesquiterpene lactones that showed IC_{50} values ranging from 0.62 to 19.27 μM against the NF54 strain (chloroquine-sensitive) and the clone A1A9 (chloroquine-resistant) of *P. falciparum* *in vitro* [8]. The most active components were neurolenin A (**7**) with IC_{50} of 0.92 μM and neurolin B (**8**) with IC_{50} of 0.62 μM . It was found that the structural requirements for high antiplasmodial activity *in vitro* is an α,β -unsaturated keto function. Additionally, a free hydroxy function at C-8 increased the antiplasmodial activity [8, 12].



The sesquiterpene dilactone 16,17-dihydro-brachycalixolide (**9**) was isolated from *Vernonia brachycalyx* (Asteraceae), a herb growing in East Africa and used by the Maasai, the Kipsigis and other East African tribes as a treatment for parasitic diseases [9]. This compound showed an IC_{50} of 26.9 μM using the 3D7 chloroquine-sensitive *P. falciparum* strain. The IC_{50} values for other tested strains, K39, V1/S and Dd2, were in a similar range, 8.3, 5.9 and 32 μM . This compound also strongly inhibited the proliferation of human lymphocytes at the same concentrations [13].




Another known sesquiterpene lactone, brevilin A was isolated from *Centipeda minima*, a plant used by the Chinese people to treat colds, nasal allergies, asthma, malaria and amoebiasis [10]. Brevilin A showed an IC_{50} of 9.42 μM against the W2 chloroquine-resistant strain [10].

Recently, the antimalarial activity of lactucin and lactupicrin isolated from *Cichorium intybus* L. (Asteraceae) was also determined against the HB3 clone of strain *Honduras-1* of *Plasmodium* which is

chloroquine sensitive and pyrimethamine resistant. The complete inhibitory activity (IC_{100}) for lactucin was 38.5 μM and the value for lactucopicrin was 126 μM [11].


Four sesquiterpene lactones of the pseudo-guaianolide type, the typical constituents of *Arnica montana* L. (Asteraceae), i.e. helenalin (**10**), dihydrohelenalin and their acetates, have shown activities against asexual blood forms of *Plasmodium falciparum* *in vitro* cultures (NF54, clone A1A9) [14]. The IC_{50} values of the four compounds were in the range from 0.23 to 7.41 μM and the most active constituent was helenalin (**10**), whose potency was comparable to that found for artemisinin (IC_{50} 0.14 μM).



Because of the cytotoxic effects of sesquiterpene lactones on various types of cells, the cytotoxic/antiplasmodial ratio was also evaluated as a measure of therapeutic efficiency. Using the cytotoxicity data obtained for helenalin and artemisinin against the human carcinoma cell lines GLC4 and COLO 320, ten times higher toxicity was found for helenalin, which makes its therapeutic usefulness questionable [14].


Several sesquiterpene lactones isolated from *Eupatorium semialatum* Benth. (Asteraceae), a plant used in the traditional medicine of Central America for malaria and dysentery, were assayed *in vitro* for their activities against *Plasmodium falciparum* (K1 strain) using the pLDH-assay [15]. All the compounds were tested and exhibited a moderate activity (IC_{50} 8.9-31.7 μM) if compared to chloroquine (IC_{50} 0.18 μM). Nevertheless these results concerning their *in vitro* activity could justify the traditional use of the plant against malaria [15].

Several diterpenoids with different structures were also reported for their antiplasmodial activity. Among the abietane-type derivatives, 3-*O*-benzoyl-hosloppone (**11**) was isolated from the roots of *Hoslundia opposita* (Lamiaceae) used in East and West Africa to treat malaria [16]. The IC_{50} against the multidrug resistant strain K₁ of *Plasmodium falciparum* was 0.95 μM and the activity of this molecule was attributed to the presence of an α,β -unsaturated carbonyl moiety [16,17].



Several studies have recently been carried out on labdane and isopimarane diterpenoids, but most of them have shown a modest *in vitro* activity against chloroquine-sensitive *P. falciparum* strains [18-20]. Among the tested constituents only 8(9),15-isopimaradien-3 β -ol, isolated from *Platycladus orientalis* (L.) Franco (Cupressaceae), gave interesting IC₅₀ values (7.1 μ g/mL, 24.6 μ M) in the inhibition of the growth of 3D7 *P. falciparum* strain [18].

Clerodane diterpenoids with a mild antiplasmodial activity have recently been isolated from two species of Flacourtiaceae, *Laetia procera* (Poepp.) Eichler a typical species of French Guiana [21] and *Casearia grewiifolia* Vent., growing widely in the northern and northeastern parts of Thailand and used traditionally as a tonic and a febrifuge [22].



The compounds isolated from *C. grewiifolia* were tested against K1 multidrug resistant strains using artemisinin as positive control, while the derivatives isolated from *L. procera* were tested against F32 Tanzania (a chloroquine-sensitive strain) and FcB1-Columbia (a chloroquine-resistant strain), using chloroquine as a positive control. The most active clerodane diterpenoids were compounds **12-14** isolated from *L. procera*, showing activities against *P. falciparum* with an IC₅₀ as low as 0.5 μ M on both FcB1 and F32 strains. The IC₅₀ values were 0.62 and


0.54 μ M, respectively, in the two strains for compound **12**, 0.57 and 0.59 μ M for **13**, and 0.58 and 0.66 μ M for **14** [21]. It was also observed in this study that the hydrolysis of the diacetal moiety lowered their biological activity [21].

In 1997 Bringmann and coworkers [23] first reported on the moderate activity of betulinic acid against *P. falciparum* *in vitro*, with an IC₅₀ of 23.0 μ M. This widespread constituent was isolated after a bioassay-guided fractionation from *Triphyophyllum peltatum* (Dioncophyllaceae) and *Ancistrocladus heyneanus* (Ancistrocladaceae). In 1999 Steele et al. [24] confirmed the *in vitro* activity of betulinic acid, but found that it was ineffective in *in vivo* experiments. Betulinic acid was also isolated from an ethanol extract of the root bark of the Tanzanian tree *Uapaca nitida* Mull-Arg. (Euphorbiaceae) used in Tanzania to treat malaria. It showed *in vitro* antiplasmodial IC₅₀ values similar to those obtained in the study of Bringmann et al. in 1997 [24]; the IC₅₀ values against chloroquine resistant (K1) and sensitive (T9-96) *P. falciparum* strains were 43.0 μ M and 63.6 μ M, respectively. The *in vitro* activities of several related triterpenes were also evaluated. Betulin was found to be inactive at 1164 μ M for both K1 and T9-96. Ursolic acid exhibited IC₅₀ values similar to betulinic acid, 80.0 μ M and 61.4 μ M, respectively. Oleanolic acid exhibited higher IC₅₀ values, 194.7 μ M and 154.8 μ M against K1 and T9-96, respectively. Thus, among the triterpenes, betulinic acid showed the highest activity and for this reason was further tested for *in vivo* activity in a murine malaria model (*P. berghei*). However, the top dosage of 250 mg/kg/day was ineffective in reducing parasitaemia and exhibited some toxicity, and thus not advisable for clinical use [24].


An investigation of *Gardenia saxatilis* Geddes (family Rubiaceae), a plant with folkloric use against malaria and distributed in the northeastern part of Thailand, led to the isolation of several triterpenoids which were assayed for antiplasmodial activity using the K1 multidrug resistant strain [25].

Four compounds, namely messagenic acid A (**15**) and messagenic acid B (**16**), the 27-*O-p*-(*Z*)- and 27-*O-p*-(*E*)-coumarate esters of betulinic acid, and a mixture of uncarinic acid E (27-*O-p*-(*E*)-coumaroyloxy-oleanolic acid) (**17**) and 27-*O-p*-(*E*)-coumaroyloxy-ursolic acid (**18**) showed moderate activity with IC₅₀ values of 2.43, 6.14 and 4.69 μ M, respectively. The results indicated that *p*-coumarate moieties at the



27-position contributed to antiplasmodial activity. As both the *p*-(*Z*)-coumarate ester **15** and the isomeric *p*-(*E*)-coumarate ester **16** were active in the assay, it was noteworthy that the difference in geometry of the double bond in the ester moieties did not significantly effect antiplasmodial activity of the triterpenes, while the introduction of a methoxyl group to the 3-position of *p*-(*E*)-coumarate moiety gave a ferulate moiety which resulted in a loss of activity [25].



A very recent investigation [26] tested several ceanothane- and lupane-type triterpenes isolated from the root bark of *Ziziphus cambodiana* Pierre (Rhamnaceae) were antiplasmodial activity. 3-*O*-Vanillylceanothic acid (**19**), 2-*O*-*E*-*p*-coumaroyl alphitollic acid (**20**) and zizyberenalic acid (**21**) exhibited significant *in vitro* antiplasmodial activity against the parasite *Plasmodium falciparum* (K1 multidrug resistant strain), with IC₅₀ values of 5.81, 1.45 and 6.61 μM, respectively.




A comparison of the structures of the tested compounds indicated that the *p*-coumaroyl moiety in **20** and the vanillyl group of compound **19** were crucial for high antiplasmodial potential. Introduction of a double bond in ring A of the ceanothane-type triterpene **21** greatly increased the inhibitory activity in the antiplasmodial assay [26].




Another extensive investigation on the antimalarial effects of triterpenoids isolated from several species of the genus *Cimicifuga* was carried out by Takahara and coworkers [27]. Fifty-nine compounds belonging to five different structural groups were investigated. Almost all the compounds tested showed activity in the 1–56 μM concentration range against *Plasmodium falciparum* FCR-3 strain.

Twenty-five compounds had an IC₅₀ 1–3 μM and nineteen of them had a common 16, 23:23, 26:24, 25-triepoxy group in the side-chain moieties. The most active compound was (26*S*)-*O*-methylactein (**22**) [27].

Studies on a species of the Celastraceae family, *Celastrus paniculatus* Willd. from Thailand, known locally as Kra-Thong-Lai and sold in the form of pressed pills for the treatment of malaria, led to the isolation of a moderately active antiplasmodial constituent, a quinonoid triterpene, pristimerin (**23**). The IC₅₀ value against K1 strain was 0.42 μM [28].



These findings were also confirmed by an investigation on another species of the Celastraceae family, *Salacia krauss*, a small shrub growing in Mozambique and KwaZulu-Natal Province, South Africa and traditionally used to treat bilharzia and dysentery. Thus, a bioassay-guided fractionation of the roots resulted in the isolation of six quinone methides including pristimerin [29]. Each of these compounds was tested *in vitro* against two strains of *P. falciparum*, a chloroquine-resistant strain (K1) and a chloroquine-sensitive reference strain (NF54). The highest activities were found for isoiguesterol (**24**) with an IC_{50} of 22.9 ng/mL (51.1 nM) against K1 and IC_{50} of 54.1 ng/mL (127 nM) against NF54.



Another constituent, 17-(methoxycarbonyl)-28-norisoiguesterin (**25**), displayed an IC_{50} of 27.6 ng/mL (60.9 nM) against K1 and an IC_{50} of 37.1 ng/mL (81.9 nM) against NF54. In addition, all the six isolated quinone methides were found to be cytotoxic against the human adenocarcinoma cell line HT-29 in the range of 1300 ng/mL up to 6060 ng/mL. They displayed, however, a 10-100-fold higher activity against plasmodia than against HT-29 cells, thus indicating some selectivity. Furthermore, compound **25** was also tested *in vivo* against *P. berghei* in mice. However, parenteral administration at 10 mg/kg body weight lead to the death of mice after 1 day, whereas 5 mg/kg and 1mg/kg parenteral as well as 30 mg/kg per oral neither cured mice nor reduced parasitaemia of *Plasmodium berghei* significantly [29].


Recently four tanshinones, i.e. 20-norditerpenes with an abietane-type skeleton containing a quinone moiety in the C-ring, were isolated from *Perovskia abrotanoides* Kar. (Lamiaceae) and moderately inhibit growth of cultured malaria parasites (3D7 strain of *Plasmodium falciparum*); the IC_{50} values ranged from 12.5 to 26.9 μ M [30].

Artemisinin and other peroxides

Among the terpenoid derivatives, artemisinin (qinghaosu) (**3**) is one of the most well-known antiparasmodial drugs, it has few adverse side effects, making this by far the most useful natural product discovered to date to treat chloroquine-resistant malaria.


Artemisinin is an unusual sesquiterpene trioxane lactone containing an endoperoxide bridge which is essential for its activity. It (**3**) was isolated in 1972 by Chinese scientists from *Artemisia annua* (Asteraceae), a Chinese herb that has been used for over 2,000 years as a remedy for chills and fever. It was quickly observed that this molecule is a rapidly acting antimalarial drug effective against chloroquine and other drug-resistant parasites, and was as good as quinine (but less toxic) for the treatment of cerebral malaria. It is very active *in vitro*, with IC_{50} values between 1-100 nM depending on the *Plasmodium* strain [31,32]. As artemisinin is a non-polar compound, derivatives including ethers (artemether, arteether) and esters (sodium artesunate, sodium artelinate) were prepared to improve its formulation characteristics. These derivatives are now increasingly used as an alternative to quinine [33].

A number of other naturally occurring peroxides, not only from *Artemisia* sp. but also from other members of the Asteraceae (*Achillea millefolium*, *Anthemis nobilis*, *Heterothalamus psiadioides*), have also been tested [34]. It was found that although all of them showed some activity, none was as active as artemisinin (**3**). A weakly active peroxide (1*S*)-1-hydroxy- α -bisabolol oxide A acetate was isolated from *Artemisia abrotanum*, a plant widely cultivated in Europe for its aromatic properties. This compound showed interesting antiparasmodial *in vitro* activity, the IC_{50} being 17.9 μ M [35].




The functional group associated with the activity, namely the endoperoxide, is also present in the structure of another natural antimalarial, yingzhaosu A (**26**) first isolated in 1979 from another Chinese plant, *Artabotrys uncinatus* (Lam.) Merr. (Annonaceae). This constituent is a typical 1,2-dioxane and it occurs as a decomposition product from the stored roots of a sparsely growing vine [36].


Although the evidence of its antimalarial activity is largely anecdotal, **26** is reported to be active against *P. berghei*. However, owing to the limitations imposed by a poor supply of yingzhaosu A, the total synthesis of **26** was proposed in 1991 starting from R-(-)-carvone [37]. Total synthesis proved to be long and tedious, but in 1994 efforts led to semisynthetic first generation derivatives as potential drug candidates. Structurally related but simplified analogues containing the 2,3-dioxabicyclo [3.3.1] nonanes were synthesized. The analogue arteflene (Ro 42-1611, **27**) is a highly active, synthetic antimalarial endoperoxide [38] which can be considered a new lead molecule because of its lower rate of recrudescence, longer lasting therapeutic effects, and a longer half-life than that of artemisinin (**3**) and its commercial derivatives [39]. Later on, a series of endoperoxides containing a sulfide or a sulfone group were synthesized and some members of this class of sulfone endoperoxides have a good *in vivo* therapeutic index (efficacy/toxicity) [40].




Several other endoperoxides have been isolated from plant sources, but most of them did not show high potency against *P. falciparum* strains. The exception to this is ascaridole (**28**), isolated from *Chenopodium ambrosioides* (Chenopodiaceae) and reported to be a potent inhibitor of plasmodial growth; at a concentration of 0.05 μM , development of plasmodium was arrested after 3 days [41]. Zingiberene 3,6- β -endoperoxide and zingiberene 3,6- α -endoperoxide isolated from two Brazilian species, *Eupatorium rufescens* and *Senecio selloi*, were reported to be active with an IC_{50} value of 49 μM against FCH-5 *Plasmodium* strains [42]. 10,12-Peroxy calamene (**29**), a sesquiterpene with an endoperoxide group similar in structure to artemisinin, was isolated from *Cyperus rotundus*, a Tanzanian plant used traditionally to treat malaria; it showed an IC_{50} value of 2.33 μM against the K1 strain [43].



Two epimers, nardoperoxide (**30**) and isonardoperoxide (**31**) were isolated from *Nardostachys chinensis* (Valerianaceae) and tested for antimalarial activity [44-46]. Their EC_{50} values against *P. falciparum* were 1.5 μM and 0.6 μM , respectively, values comparable with that of quinine (0.11 μM). In addition, studies of cytotoxicity against FM3A and KB cells showed that the selectivity (cytotoxicity/antimalarial activity) of these compounds was comparable to that of quinine. Therefore, these compounds could be considered as promising leads for a new class of antimalarial drugs.



Finally, a diterpene peroxide (**32**) isolated from the spice cardamom, *Amomum krevanh* Pierre (Zingiberaceae) showed an antiplasmodial activity about one-tenth that of artemisinin, having an IC_{50} of 0.17 μM [47].




Quassinoids

Quassinoids are terpenoid bitter principles of the *Simaroubaceae* family, including the genera *Ailanthus*, *Brucea*, *Eurycoma* and *Simarouba*. Originally, these bitter substances were termed *quassin*, after a man by the name of "Quassi" who treated fever with the bark of these plants [48, 49]. Chemically they are degraded triterpenes and are categorized into five groups according to their basic skeleton.

Many quassinoids display a wide range of biological activities *in vitro* and/or *in vivo*, and their activity is related to both the position and nature of the ester group and, on the other hand, to the substitution of the A nucleus [48]. Constituents with antiplasmodial activity are mainly represented by the C-20 skeleton. An α,β -unsaturated ketone in ring A and an oxymethylene bridge in ring C are generally considered necessary for antimalarial activity [49].

At the end of the 1940s it was demonstrated that the majority of *Simaroubaceae* have activity on malaria in birds [50] and at the beginning of the 1980s a strong antimalarial activity *in vitro* of many derivatives was demonstrated [51]. Although several quassinoids are cytotoxic, results do indicate that cytotoxicity and antimalarial activity are not correlated, suggesting that the antimalarial activity is not merely cytotoxicity, but that selectivity is present [52]. Therefore, more investigations should be carried out in order to obtain specific information regarding the mechanism of action of these compounds.


The first molecules tested *in vivo* using *P. berghei*-infected mice were bruceine B (**33**) and brusatol (**34**). They showed some activity, but they were found to be toxic at higher levels than were necessary for antimalarial activity [53].



Active quassinoids have also been isolated from the fruits of *Simarouba amara* of the Republic of Panama [54], *Ailanthus altissima* [55], *Simana cedron* [56], the Brazilian plant *Simaba guianensis* [57], *Eurycoma longifolia* [58], the Central African *Hannoa chlorantha* and *Hannoa klaineana* [59], the Guinanan *Picrolemma pseudocoffea* [60], and stems of the Indonesian plant *Quassia indica* [61]. All the tested quassinoids showed good activity against chloroquine-resistant and chloroquine-sensitive strains of *P. falciparum* and against *P. vinckei petteri* or *P. berghei* in mice. Studies on the structure-activity relationships of the quassinoids [48,62] indicated that the type and presence of an ester group at C-15 was vital for the antiplasmodial activity. Ring A substitution also affected the activity, with a diosphenol moiety in ring A giving the highest activity. The glycosides were found to be generally less active than the corresponding aglycones [48,62].

The most active quassinoids reported in the literature are gutolactone (**35**) and simalikalactone D (**36**) isolated from the bark of *Simaba guianensis* collected near Manaus, Brazil. They were tested against two *Plasmodium falciparum* strains: the W-2 Indochina, a chloroquine-resistant strain, and the D-6 Sierra

Leone, a mefloquine-resistant strain. Most notably, the activity was the same for the two different strains, since compound **35** showed IC₅₀ values of about 9 nM and compound **36** displayed an IC₅₀ of about 3.4 nM. Both compounds presented *in vitro* antimalarial activity similar to or better than that of known antimalarials used as standards (chloroquine, mefloquine, artemisinin, quinine) [57].




Limonoids


Limonoids are bitter constituents which have a polyoxygenated triterpenoid skeleton biosynthetically connected to the quassinoids. Limonoids are most often found in the family Meliaceae and less frequently in the families Rutaceae and Cneoraceae. Of the over 300 limonoids known today, about one-third are accounted for by neem (*Azadirachta indica*) and Chinaberry (*Melia azedarach*) [63, 64]. The first limonoid found active against *Plasmodium* was gedunin (**37**) with an IC₅₀ of 0.040 μM (0.02 μg/mL) isolated from *Melia azedarach* [65]. However, this compound was not active *in vivo* against *Plasmodium berghei* in mice. [66].

On the other hand, recently it has been found that the combination of gedunin with chloroquine has an additive effect [67]. In addition, a recent *in vivo* reinvestigation [68] of the antimalarial activity of gedunin (**37**) in CD-1 mice infected with *Plasmodium berghei* led to some interesting results. When orally administered at 50 mg kg⁻¹ day⁻¹ for four days, gedunin (**37**) was able to suppress the parasitaemia level by 44%. However, no clear dose-response effects were observed in the 0-100 mg kg⁻¹ day⁻¹ dose range. Preliminary pharmacokinetics in Sprague-Dawley rats showed poor absorption, but a binary treatment of 50 mg kg⁻¹ day⁻¹ gedunin with 25 mg kg⁻¹ day⁻¹ dillapiol, a cytochrome P450 inhibitor, increased parasitaemia clearance in mice to 75%. A clear dose-response curve was observed in the 0-50 mg kg⁻¹ day⁻¹ gedunin dose range when administration was combined with 25 mg kg⁻¹ day⁻¹ dillapiol. In addition, 7-methoxygedunin, a semi-synthetic derivative which is more stable to degradation than gedunin, suppressed the level in

mice by 67% at 50 mg kg⁻¹ day⁻¹. When administered at this dose in combination with 25 mg kg⁻¹ day⁻¹ dillapiol, clearance increased to 80%. These results demonstrate the potential efficacy of gedunin and the value of combination therapy [68].



Studies on the leaves of *Azadirachta indica* collected in India resulted in the isolation of four limonoids active against the chloroquine-resistant K1 strain of *P. falciparum* [69]. Further investigations on *A. indica* have been carried out by Jones and coworkers [70] and Dhar and coworkers [71]. Jones and his co-workers looked at azadirachtin (**38**) and a series of 17 semisynthetic derivatives and their effects *in vitro* on male gamete production from malarial microgametocytes.




Azadirachtin (**38**) and three of the semisynthetic derivatives were found to inhibit the formation of mobile male gametes *in vitro*. This study indicated that the presence of a hemiacetal group at C-11 was vital to the activity. Dhar and coworkers [71] investigated the seeds of *A. indica* and found that the extract was active against all the erythrocytic stages of *P. falciparum*. In addition, the neem extracts also revealed a gametocytocidal effect with inhibition of the asexual stages of the parasite. All stages of maturation of the gametocytes were affected, unlike artemisinin and primaquine which only affect the immature stages [71].

Polyphenols


Over the last three decades studies on polyphenol plant constituents have shown antiplasmodial activity by almost all the classes of polyphenols. Simple galloyl derivatives isolated from *Swintonia forworthi* Elmer (Anacardiaceae), a large tree of the Philippines, showed activity against two strains of *P. falciparum* (W-2, a chloroquine-resistant one and

D-6, a chloroquine-sensitive one). Methyl gallate showed an IC₅₀ of 19 μM for the D6 and an IC₅₀ of 10.9 μM for the W2 strain. Methyl 3-*O*-galloylgallate showed an IC₅₀ of 28.8 μM and for W2 a value of 13.7 μM against D6. Methyl gallate demonstrated a selectivity index of >5 towards the D6 strain and >8 towards the W2 strain when compared with cytotoxicity towards BC1, Lu1, CoI2, KB-V1, and LNCaP cancer cells, while methyl digallate demonstrated a selectivity index of >4 against both strains [72].




Ellagic acid (**39**) and 3,4,5-trimethoxyphenyl-(6'-*O*-galloyl)-*O*-β-D-glucopyranoside, isolated from *Tristanopsis calobuxus* Bronghiart & Gris, *T. yatensis* J.W. Dawson and *T. glauca* Bronghiart & Gris (Myrtaceae) inhibited the growth of chloroquine-sensitive and resistant clones. Their IC₅₀ values were 0.5 and 3.2 μM, respectively [74].

Gossypol (**40**), the most abundant component of cottonseed (cotton=*Gossypium* sp., Malvaceae), is known for a variety of biological activities, including antispermatogenic, anticancer, antiparasitic and antiviral activity. It also demonstrated a weak antimalarial activity against both chloroquine-sensitive and chloroquine-resistant strains of *P. falciparum*, with IC₅₀ values in the order of 10 μM. [75].




Three prenylated stilbenes, isolated from the edible fruits of *Artocarpus integer* (Moraceae), popular among the people in Thailand, exhibited moderate activity. Their EC₅₀ values against the K1 multidrug

resistant strain were 5.66 μM , 26.3 μM and 32.0 μM , respectively, with the novel compound **41**, being the most active [76]. Two other stilbenes, longistylin A and C, isolated from the roots and leaves of *Cajanus cajan* (L.) Millsp. (Fabaceae) showed a moderately high activity *in vitro* against the chloroquine-sensitive *Plasmodium falciparum* strain 3D7 [77].



Bioassay-guided fractionation of the leaves from *Andira inermis* led to the isolation of numerous polyphenol constituents including isoflavones, dihydroflavonols and three novel 2-arylbenzofuran-3-carbaldehydes, and inermal A–C. Andinermal A (**42**) exhibited the strongest antiplasmodial activity *in vitro* with IC_{50} values of 6.69 μM against the poW strain (chloroquine-sensitive) and 11.3 μM against the Dd2 strain (chloroquine-resistant). Andinermal C (**43**) was slightly less active and the values were 17.8 μM (poW) and 19.0 μM (Dd2), respectively [78].




Two 5-methylcoumarin epoxides and several 4-phenylcoumarins have been found to be active against *P. falciparum* strains *in vitro*. The first compounds were isolated from the roots of *Vernonia brachycalyx* Hoffm. (Asteraceae), an herb used by the Maasai, the Kipsigis and other East African tribes as a treatment for parasitic diseases [79]. Their structures were 2'-epicycloisobrachycoumarinone epoxide (**44**) and cycloisobrachycoumarinone epoxide (**45**) epoxide, both of which showed antiplasmodial activity against chloroquine-sensitive (3D7) and chloroquine-resistant (Dd2) strains of *P. falciparum* *in vitro*. IC_{50} values for the strain 3D7 were 160 and 111 μM , respectively, while the IC_{50} values for the strain Dd2 were 54 μM for both compounds [79].

A second group of coumarins active against *Plasmodium* strains was isolated from the stem bark of *Exostema mexicanum* (Rubiaceae), used in Latin American folk medicine as a quinine substitute for malaria treatment. The most lipophilic compound, 4',5,7,8-tetramethoxy-4-phenylcoumarin (*O*-methyl-

exostemin), revealed the strongest antiplasmodial activity with IC_{50} values of 10.5 μM (poW strain *P. falciparum*) and 4.68 μM (Dd2 strain) [80].

Four coumarins, theraphins A–D, isolated from *Kayea assamica* King & Prain (Clusiaceae), an evergreen tree used as a remedy for treating fevers in India, were tested against a panel of human cancer cell lines to assay their cytotoxicity, and tested for antimalarial activity against the D6 (chloroquine-sensitive) and W2 (chloroquine-resistant) clones of *Plasmodium falciparum*. The constituents were characterized by a 1-hydroxypropyl moiety linked to C-4, a 1-oxobutyl moiety linked to C-8 and an isoprenyl chain linked to C-6. They showed modest antiplasmodial activities, with IC_{50} values in the range 9.7–11.1 μM against the D6 clone, and IC_{50} values in the range 5.1–10.4 μM against the W2 clone. However, their Selectivity Indices ($\text{SI} = \text{KB } \text{IC}_{50} / \text{P. falciparum } \text{IC}_{50}$) were less than 1.0, although the values for theraphin D (i.e. 11(S)-(-)-8,8-dimethyl-5-hydroxy-4-(1-hydroxypropyl)-10-(1-oxobutyl)-2*H*,8*H*-benzo(1,2-*b*:3,4-*b'*)di-pyran-2-one) were 4.70 and 5.02 for the D6 and W2 clones, respectively. These observations indicated that the coumarin derivatives possess little potential as antimalarial drugs, although appropriate structure modifications of some of them might improve the SI level leading to derivatives of greater antimalarial potential [81].


Among neolignan derivatives, polysyphorin (**46**), isolated from *Rhaphidophora decursiva* (Araceae), a vine growing in Vietnam, showed antiplasmodial activity. From the same plant was also isolated a new active benzoperoxide, raphidecurperoxin (**47**) [82]. Compounds **46** and **47** were tested against the oral epidermoid cancer line KB and cultures of the chloroquine-sensitive clone D6 and chloroquine-resistance clone W2 of *P. falciparum*. The IC_{50} of the neolignan was 0.92 μM (D6 strain) and 0.84 μM (W2 strain) with selectivity indices of 5 and 6, respectively, compared to the KB cell line.




Therefore, compound **46** appears promising and further evaluation in *in vivo* antimalarial models should be pursued. It was also interesting to note that

47, which contains a peroxide ester, a moiety similar to the endoperoxide bridge of artemisinin, showed a moderate antimalarial activity; its IC_{50} against the D6 strain was 1.76 μM and against W2 was 1.37 μM . However, due to its poor SI values (0.7 and 1, respectively), this compound is not considered of great interest as an antimalarial agent [82].

Recently the neolignan nitidanin (**48**) has been isolated from *Grewia bilamellata* Gagnep. (Tiliaceae). It displayed weak antimalarial activity in cultures of *P. falciparum* clones D6 and W2 (IC_{50} 21.2 and 18.4 μM , respectively). The same derivative tested against the human oral epidermoid carcinoma cell line (KB) showed a minimal cytotoxicity (ED_{50} >99.0 μM) and thus its selectivity index (SI) expressed as ED_{50} (KB)/ IC_{50} (*P. falciparum*) was high against both *Plasmodium* clones (>4.6 and 5.4, respectively). Thus, this molecule could represent a model structure because several neolignans with antimalarial activity were previously reported, but were highly cytotoxic [83].




Several other lignans are reported to have antiplasmodial activity. Two of them, termilignan and anolignan B were isolated from *Terminalia bellerica* (Combretaceae), a species extensively used in the Indian system of traditional medicine for the treatment of fever, cough, diarrhea, dysentery and skin conditions [84]. These compounds were tested against the chloroquine-susceptible strain 3D7 of *Plasmodium falciparum* and showed IC_{50} values of 9.6 and 20.5 μM , respectively [84].




Another antiplasmodial lignan was isolated from a palm, *Euterpe precatorea* Mart. (Aracaceae): it was the 8-5'linked lignan dehydrodiconiferyl dibenzoate showing a similar antiplasmodial activity. The IC_{50} value was 12 μM when the compound was tested against the chloroquine-sensitive 3D7 *Plasmodium falciparum*. [85]

Phytochemical investigation of the aerial parts of *Bonamia spectabilis* (Choisy) Hall. (Convolvulaceae) led to the isolation of some active tetrahydrofuran-type sesquilignans (**49-52**). The derivatives were tested for their antiplasmodial activity against a chloroquine-sensitive strain (PoW) and a chloroquine-resistant clone (Dd2) of *Plasmodium falciparum*. Bonaspectin C 4''-O-glucoside (**49**), its aglycone (**49a**), and bonaspectin D 4''-O-glucoside (**50**) revealed the highest antiplasmodial activities (IC_{50} values: 1.3, 2.0, 6.5 μM [PoW]; 1.7, 4.6, 3.7 μM [Dd2], respectively).




The sesqueneolignans **51** and **52** revealed antiplasmodial activity with IC_{50} values of 9.9, 3.0 μM (PoW) and 10.9, 8.5 μM (Dd2), respectively.



There was no significant difference of activity between the chloroquine-sensitive strain PoW and the chloroquine-resistant clone Dd2, however the phenylpropanoid dimers showed lower antiparasitic activities than the related trimers [86].

Several flavonoids, including biflavonoids, have been recognised for their antiplasmodial activity. Among them two flavanones, exiguaflavanone A (**53**) and exiguaflavanone B (**54**), were isolated from *Artemisia indica* from Thailand [87]. The assay was carried out with *P. falciparum* (K1, multidrug-resistant strain) and the constituents exhibited an IC_{50} of 10.8 μM and 16.0 μM , respectively [87].




Two flavones, 5,7,4'-trimethoxyflavone and 5,7,3',4'-tetramethoxyflavone isolated from *Kaempferia parviflora* (Zingiberaceae), another plant from Thailand, [88] showed a weak antiplasmodial activity (IC₅₀ values were 11.9 and 12.5 μ M, respectively).

Three flavonol glycosides (all kaempferol derivatives) isolated from *Hydrangea macrophylla* Seringe var. *thunbergii* Makino (Hydrangeaceae), a Japanese plant, were tested for the antimalarial properties and the cytotoxic activity against KB3-1 cells. The compounds exhibited characteristic anti-malarial activity: in particular, approximately 60% of proliferation of the parasite was inhibited even at the concentration of 0.5 ng/mL. On the other hand, these flavonol glycosides have little influence on the growth of KB 3-1 representing the host cell [89].

Other flavonoids, namely (R)-4'-methoxydalbergione, obtusafuran, 7,4'-dihydroxy-3'-methoxyisoflavone, and isoliquiritigenin, isolated from the heartwood of *Dalbergia louvelii*, inhibit the growth of *P. falciparum* *in vitro*. Their IC₅₀ values ranged from 5.8 to 8.7 μ M [90].

Five rotenoids, a chalcone and an isoflavone isolated from the stem bark of *Milletia usaramensis* subsp. *usaramensis* (Fabaceae), a plant of Kenya were tested against chloroquine-resistant (W2) and chloroquine-sensitive (D6) strains of *P. falciparum*. The chalcone 4'-O-geranylisoliquiritigenin was the most potent compound (IC₅₀ values were 8.7 and 10.6 μ M, respectively). Among the rotenoids, those containing a prenyl or a 2,2-dimethylpyrano substituent were most potent (IC₅₀ values were between 19.4 and 70.1 μ M) [91].




Another simple chalcone, licochalcone A (**55**) isolated from *Glycyrrhiza* species in different

amounts, is reported as having *in vitro* and *in vivo* antimalarial activity [92, 93].

In *in vivo* tests against *P. yoelii* in mice, oral doses of 1000 mg/kg resulted in the complete eradication of the malaria parasite and no toxicity was noted [92]. *In vitro* the IC₅₀ was 1.78 μ M (0.6 μ g/mL) on the chloroquine-sensitive (3D7) and chloroquine-resistant (Dd2) strain of *P. falciparum*. An intraperitoneal injection of 15 mg/kg/ of licochalcone A twice daily for three days led to the survival of mice infected with *P. yoelii* and clearance of parasites. Oral administration (50 mg/kg) of a suspension of this flavonoid to mice infected with *P. yoelii* revealed that after four days, that the animals were no longer infected, thus it is efficient in controlling the infection [93]. A further investigation on licochalcone A showed that it is a potent membrane-active agent that transforms normal erythrocytes into echinocytes in parallel with the inhibition of growth of *Plasmodium falciparum* cultures. Thus, the *in vitro* antiplasmodial effect apparently is an indirect effect on the host cell. This effect could also be transiently observed *in vivo* after intravenous administration of the compound, but the cells returned quickly to the normal shape, presumably as the result of redistribution of licochalcone A in lipophilic compartments of the body or removal of damaged erythrocytes [94]. Recently it has also been demonstrated that licochalcone A can inhibit the bc(1) complex (ubiquinol-cytochrome c reductase) as well as complex II (succinate ubiquinone reductase, SQR) of *Plasmodium falciparum* mitochondria at very low concentrations. Because the property of the *P. falciparum* bc(1) complex is different from that of the mammalian host, chalcones could be promising candidates for a new antimalarial drug [95].

Xanthohumol (**56**), an isomer of licochalcone A, and seven derivatives isolated from *Humulus lupulus* L (Cannabinaceae) were tested for their *in vitro* antiplasmodial activity against the chloroquine-sensitive strain poW and the multiresistant clone Dd2. Of the eight compounds tested, four possessed activity with IC₅₀ values <25 μ M against at least one of the two strains of *Plasmodium falciparum*. The main hop constituent, the chalcone xanthohumol, was the most active with IC₅₀ values of 8.2 μ M (poW) and 24.0 μ M (Dd2). Three of these compounds were additionally active in the haemin-degradation assay [96].



Other prenylated chalcones isolated from *Crotalaria orixensis* L. (Fabaceae) have been tested for *in vitro* antiplasmodial activity against NF-54 chloroquine sensitive strains. The most active compound was 3',5'-diprenyl-4,2',4'-trihydroxy chalcone, which inhibited the parasites 100% at 5.09 μ M [97]. Within the same study it was shown that substitution at the 4' and 4-hydroxyl groups decreases the activity. The presence of prenyl moieties can affect the activity positively especially with free 4,4'-dihydroxy systems [97].

Structure-activity relationship studies of antimalarial chalcones were carried out using a series of forty oxygenated derivatives obtained by synthesis [98]. Good antimalarial activity was found among alkoxyated chalcones with polar A rings, in particular those substituted with electron-withdrawing groups or replaced by quinoline rings. The size characteristics of ring B (large, alkoxyated) and the electronic properties of ring A (electron deficient) are considered as important for antimalarial activity [98].


Two studies reported the antiplasmodial activity of prenylated flavonoids isolated from *Erythrina abyssinica* L. (Leguminose), a species widely used in Africa to treat infectious diseases. Flavonoids were tested against two *Plasmodium* strains: the chloroquine-sensitive D6 and the chloroquine-resistant W2 clones. Chalcones, flavanones and isoflavones with prenyl moieties showed weak activity against both strains with IC_{50} values ranging from 4.9 to 27.7 μ M [99,100]. From another *Erythrina* species, *Erythrina saculeuxii*, several flavanones, isoflavones, and isoflavanones with isoprenyl moieties were isolated. These compounds displayed a similar antiplasmodial activity against the chloroquine-sensitive D6 and the chloroquine-resistant W2 *Plasmodium* strains. Their IC_{50} values ranged from 4.9 to 28.0 μ M [101].

Three new prenylated flavonoids, namely the two flavanones 5,7,3'-trihydroxy-4',5'-(2'',2'''-dimethylpyran)-8,2'-di(3-methyl-2-butenyl)-(2*S*)-flavanone and 5,7,3'-trihydroxy-4'-methoxy-8,2'-di(3-methyl-2-

butenyl)-(2*S*)-flavanone and the flavan 7,3',4'-trihydroxy-6-methoxy-8,2'-di(3-methyl-2-butenyl)-(2*S*)-flavan, were isolated from the roots of *Dendrobium lanceolatum* (Dunn) Schindl. (Fabaceae) and assayed against the parasite *Plasmodium falciparum* (K1, multidrug-resistant strain) and several cancer cell lines. They exhibited antimalarial activity with IC_{50} values of 5.3, 7.1, and 6.9 μ M, respectively. However the flavanones were also cytotoxic, in particular the first flavanones showed strong cytotoxicity against the cancer cell lines KB, BC, and NCI-H187 with IC_{50} values of 2.4, 3.3, and 1.2 μ M, respectively, while the latter showed moderate cytotoxicity against the NCI-H187 cell line with an IC_{50} value of 17.5 μ M [102].

The dihydrochalcone, 2',4,6'-trihydroxy-4'-methoxydihydrochalcone (asebogenin) isolated from *Piper hispidum* Sw (Piperaceae), a species used by the indigenous population of Central America to treat malaria or fever, exhibited an IC_{50} of 56 μ M for poW strains and 35 μ M for Dd2 strains [103].


Two common flavone glycosides, luteolin 7-*O*- β -D-glucopyranoside (**57**) and chrysoeriol 7-*O*- β -D-glucopyranoside (**58**), isolated from *Phlomis brunneogaleata* Hub.-Mor. (Lamiaceae), were determined to be the major anti-malarial principles of this plant. Their IC_{50} values were 5 and 13 μ M, respectively, using a K1 strain (chloroquine- and pyrimethamine-resistant). The same compounds, tested with skeletal myoblast L6 cells in order to evaluate their cytotoxicity, did not show any activity at the maximum tested dose of 90 μ g/mL (about 200 μ M). In addition, compound **57** showed a promising FabI-inhibiting effect (the IC_{50} was about 22.2 μ M) [104].



Bioassay-guided fractionation of a *Satureja parvifolia* (Philippi) Epling. (Lamiaceae) MeOH extract led to the isolation, among others, of eriodictyol and luteolin as its active components against *Plasmodium falciparum* K1 strain. The IC_{50} value of luteolin was 22.3 μ M while that of eriodictyol was 59.7 μ M. Besides their moderate antiplasmodial activity, flavonoids showed a very low toxicity on the mammalian KB cell line and


eriodictyol was the most selective compound as a result of its rather low cytotoxicity (IC_{50} 604.2 μ M) [105].

A very recent investigation reported on the inhibition by several flavonoids of different enzymes of *Plasmodium falciparum* fatty acid biosynthesis: α -ketoacyl-ACP-reductase (FabG), α -hydroxacyl-ACP-dehydratase (FabZ), and enoyl-ACP-reductase (FabI). About forty related structures were investigated and several compounds were found to have very good activity against all three enzymes. The flavones and flavonols exhibiting a simple substitution pattern (that is, no hydroxy groups on ring B and one or two hydroxy groups on rings A/C) show moderate inhibition effects toward FabG (10-100 μ M), FabZ (20-30 μ M), and FabI (10 μ M) while flavonoids having more than one hydroxyl substitution on ring B exhibited strong activity toward all three enzymes (IC_{50} 0.5-8 μ M). The methylation of any of the hydroxy groups in flavonols generally abolishes almost all activity against all three enzymes. Among the flavanones tested, only 5,7-dimethoxy-8-methylflavanone showed some inhibitory activity against FabZ (40 μ M). The isoflavonoids tested showed moderate and selective activity only against FabZ with IC_{50} values in the range of 7-30 μ M. The most active compounds were C-3 galloyl acid esters of catechins, which are strong inhibitors of all three enzymes (IC_{50} 0.2-1.1 μ M). Catechins and epicatechins, carrying a free hydroxy group at C-3, neither inhibit the enzymes nor have antiparasmodial activity. This study suggests that flavonoids and analogues are promising antimalarial agents, thus adding new targets to the broad spectrum of biological activities demonstrated by these compounds [106].




Within the same investigation it was shown for the first time that C-3 galloyl acid esters of catechins had *in vitro* activity against chloroquine-sensitive (NF54) and -resistant (K1) *P. falciparum* strains in the low to submicromolar range. The most active compound was (-)-catechin gallate (**59**) with EC_{50} values of 3.2 and 0.4 μ M, respectively [106].

The same finding was also confirmed by another recent investigation [107]. Within this study two *P. falciparum* strains were investigated, namely 3D7, a chloroquine-sensitive one, and F9CR-1/FVO, a chloroquine-resistant one. Remarkably, pronounced plasmodicidal effects on both tested parasite strains were measured for (-)-epigallocatechin gallate (**60**, IC_{50} 30 μ M for 3D7 and 20 μ M for F9CR-1/FVO), and (-)-epicatechin gallate (**61**, IC_{50} 7 μ M for 3D7 and 5 μ M for F9CR-1/FVO).




Furthermore a synergism was observed between artemisinin and these two derivatives on the 3D7 drug-sensitive parasite strain using sublethal doses of artemisinin, ranging from 1 to 10 nM, both of them in the presence (and in the absence) of 15 μ M (-) epigallocatechin gallate (**60**) or of 5 μ M (-) epicatechin gallate (**61**) [107].



Several biflavonoids have shown antiparasmodial activity. Among biflavanones, 7,7'-di-O-methyltetrahydromentoflavone (**62**) isolated from *Rhus retinorrhoea* (Anacardiaceae), a tree growing in the southern parts of Saudi Arabia, showed weak antiparasmodial activity but no cytotoxicity [108].



The compound exhibited weak antimalarial activity against *Plasmodium falciparum* (W2 clone) with an IC_{50} of 1.6 μ M, and activity against *P. falciparum* (D6 clone) with an IC_{50} of 4.6 μ M. [108].



Two biflavanone isomers of **62**, namely sikokianin B (**63**) and sikokianin C (**64**) with moderate activity (IC_{50} values of about 1 μM) against a chloroquine-resistant strain (K1) and a drug-sensitive strain (FCR3) of *Plasmodium falciparum* were also isolated from *Wikstroemia indica* (Linne) C.A. Meyer (Thymelaeaceae). Their activity for the K1 strain was nearly the same as chloroquine but they were less than 2% as active as artemisinin [109].



- 66** $R_1 = H, R_2 = CH_3, R_3 = H$
67 $R_1 = CH_3, R_2 = CH_3, R_3 = H$
68 $R_1 = H, R_2 = CH_3, R_3 = CH_3$
69 $R_1 = CH_3, R_2 = CH_3, R_3 = CH_3$


Recently, the antiplasmodial activity of eight other natural biflavones was evaluated [110]. Lanaroflavone (**65**) showed the highest antiplasmodial activity (IC_{50} of 0.48 μM) when studied *in vitro* on a K1 chloroquine-resistant strain of *Plasmodium falciparum*. Other biflavones of the amentoflavone type, namely bilobetin (**66**), ginkgetin (**67**), isoginkgoetin (**68**) and sciadopitysin (**69**) showed medium activity (IC_{50} values were 6.7, 2.0, 3.5, and 1.4 μM , respectively). Lanaroflavone also exhibited a high selectivity index value ($SI=159$), indicating selective antiplasmodial activity and no significant cytotoxicity [110].

A new biflavanoid, ent-naringeninyl-(I-3 α ,II-8)-4'-O-methylnaringenin, isolated from the root bark of *Garcinia livingstonei* collected in Tanzania, showed moderate activity against *P. falciparum* (chloroquine-sensitive Ghana strain); the IC_{50} was 6.7 μM . Within the same assay the biflavonoids (+)-volkensiflavone and (+)-morelloflavone were also tested and displayed IC_{50} values of 6.0 and 48.0 μM , respectively [111].

Several phenylanthraquinones showed considerable activity with only a little cytotoxicity as well, whereas the individual anthraquinone and phenyl moieties were completely inactive. Knipholone (**70**) and three of its natural derivatives, along with seven structurally-related but simplified compounds, have been examined for their antiplasmodial activity against asexual erythrocytic stages of two strains of *Plasmodium falciparum in vitro* (K1/chloroquine-resistant and NF54/chloroquine-sensitive). All the phenylanthraquinones showed considerable activity with IC_{50} values 0.38-2.37 μM for the K1 strain and 0.42-2.64 for the NF 54 strain. Knipholone (**70**) and its natural derivatives can therefore be considered a new group of potential antimalarials [112].


From another *Bulbine* species, *B. frutescens* (L.) Wild (Asphodelaceae), three novel phenylanthraquinones were isolated, namely 4'-O-demethylknipholone-4'-O-beta-D-glucopyranoside (**70a** a glycoside derivative of knipholone), and gaboroquinones A and B. These were tested against the chloroquine- and pyrimethamine-resistant K1 strain and against the strain NF54 of *P. falciparum* which is sensitive to all known drugs.

The glycoside **70a** displayed the best activity (IC_{50} 0.7 μM for both strains) and did not exhibit any cytotoxic effects on mammalian cells, at least at concentration below 0.15 mM, the highest concentrations tested [113].





Morinda lucida is widely used in West Africa to treat malaria and other tropical diseases. Anthraquinones isolated from this plant have been tested against chloroquine-susceptible (3D7) and chloroquine-resistant (Dd2) strains. Their activity was moderate in both strains, with IC_{50} values between 21.4 and 87.8 μM . Structure-activity relationships studies showed that an aldehyde group at C-2 and a phenolic hydroxy group at C-3 enhance activity of these anthraquinones against *Plasmodium* strains [114,115]

Xanthenes from *Garcinia dulcis* and *G. cowa* (Clusiaceae) have been investigated for antiparasmodial activity [116,117]. *G. cowa* is widely distributed in Thailand where it is used as an antipyretic, while *G. dulcis* is mostly known for its disinfective activity [116,117]. Among the five xanthenes isolated from *G. dulcis*, the most active against chloroquine-sensitive strains of *P. falciparum* (T9/94 line) is garciniaxanthone (**71**) with an IC_{50} of 2.06 μ M. The presence of isoprenyl moieties at C-2, C-7 or C-8 enhanced the antiparasmodial activity [116,117].




Recently, twenty-two xanthenes isolated from *Calophyllum caledonicum* and *Garcinia vieillardii*, (Clusiaceae) were tested against chloroquine-resistant strains of *Plasmodium falciparum* (FcB1/colombia) [118]. The most potent xanthenes were found to be **72**, **73** and **74** (IC_{50} of c.a. 1.0 μ g/mL) which are 1,3,7 trioxygenated and prenylated at the positions 2 and 8. The relationship between antimalarial activity and molecular structure of xanthenes has been explored. Firstly, the position of the hydroxyl groups appears to be important, as indicated by the observed differences in activity. Indeed, oxygenation at the positions 1, 3 and 7, seems to improve antimalarial activity. Secondly, substitution with a 1,1-dimethylallyl chain or the presence of an additional pyran ring appear to be activity-enhancing factors, as well as substitution with two isopentenyl chains or combination of one isopentenyl chain and a pyranic ring. Moreover, hydroxylation of the prenyl side chain is not required for activity [118].



In addition, the *in vivo* antimalarial activity of some hydroxyxanthenes was recently demonstrated for the first time [119].

Another study reported on a series of oxygenated xanthenes which were synthesized and evaluated *in vivo*, using four-day suppressive assays against *Plasmodium berghei* ANKA in BALB/c mice. When given at a dose of 20 mg/kg/day for four days, most of the compounds produced significant chemosuppression of parasitaemia. The most active compound was 1,3,6,8-tetrahydroxyxanthone, which reduced the percentage of erythrocytes infected by 70.5%, followed by norlichexanthone (44.3%) and its isomer, 1,3,8-trihydroxy-6-methylxanthone (37.0%). While di-C-allyl-dihydroxyxanthone showed lower but still notable activity (33.4%), 1,3-dihydroxyxanthone was much less active (15.1%). This is the first demonstration of the antimalarial activity of some hydroxyxanthenes *in vivo* [119]. In a different investigation, four xanthenes isolated from the roots of *Andrographis paniculata* Nees (Acanthaceae), namely 1,8-di-hydroxy-3,7-dimethoxy-xanthone, 4,8-dihydroxy-2,7-dimethoxy-xanthone, 1,2-dihydroxy-6,8-dimethoxyxanthone and 3,7,8-trimethoxy-1-hydroxy xanthone, were assayed *in vitro* using a chloroquine-sensitive strain FSG. 1,2-Dihydroxy-6,8-dimethoxy-xanthone was the most active (IC_{50} of 4 μ g/mL), and it was tested *in vivo* in mice with a *Plasmodium berghei* infection using the Peters' 4-day test. A substantial reduction (62%) of parasitaemia was observed in mice with a 30 mg/kg dose. *In vitro* cytotoxicity against mammalian cells revealed that 1,2-dihydroxy-6,8-dimethoxy-xanthone is non-cytotoxic with an $IC_{50} > 32 \mu$ g/mL [120].




Finally, in 2006 several papers reported on the antimalarial activity of some natural xanthenes. A new prenylated xanthone, 5-O-methylcelecoxanthone (**75**), together with six related constituents from the roots of *Cratoxylum cochinchinense* (Lour.) Blume (Clusiaceae) have been tested for antiparasmodial and cytotoxic activity. Four derivatives including the new one showed cytotoxic activity against the human lung cancer cell line (NCI-H187) with IC_{50} values ranging from 1.4 μ M to 0.011 mM. In the same concentration


ranges they showed antimalarial activity against *Plasmodium falciparum* with IC_{50} values from 5.62 μ M and 0.015 mM [121].

Five other previously known prenylated xanthenes isolated from the root bark of *Garcinia livingstonei* collected in Tanzania, were tested against a chloroquine-sensitive Ghana strain of *P. falciparum*. The dimeric xanthone garcilivin A (**76**) showed the highest antiparasitic activity (IC_{50} 6.7 μ M) but it was cytotoxic in the same range of concentration (IC_{50} 2.0 μ M against MRC-5 cells). Its diastereoisomer garcilivin C and the monomeric xanthenes showed IC_{50} values ranging from 10 to 68 μ M against *Plasmodium* with remarkable selectivity against MRC-5 cells (IC_{50} > 32 μ M) [111].

Three polyprenylated structurally related xanthenes (gaboxanthone, symphonin and globuliferin) isolated from *Symphonia globulifera* L (Guttiferae), a tree whose bark is used in the Northwestern province of Cameroon to treat malaria, were tested for their anti-plasmodial activity against the W2 strain of *P. falciparum*, which is resistant to chloroquine and other antimalarials.



They all exhibited good to moderate activity relative to chloroquine, and symphonin (**77**) had the best potency (IC_{50} was 1.29 μ M). From the structure–activity relationship, it appeared that the cyclization of one of the isopentenyl groups (positions 2 and 4) to give a pyran ring increases the potency of xanthenes. The best result was obtained when the dimethylpyran ring is attached to positions 3 and 4 of the xanthone nucleus as in symphonin (**77**) [122]. A benzophenone, guttiferone A was also isolated and found to be moderately active (IC_{50} 3.17 μ M) [122].




A further study on prenylated xanthenes was carried out on a new prenylated xanthenedione, 1,2-dihydro-3,6,8-trihydroxy-1,1,7-tri(3-methylbut-2-enyl) xanthen-2,9-dione and five known xanthenes isolated from


the stem bark of *Allanblackia monticola* Staner L.C. The compounds were tested on two strains of *Plasmodium falciparum*, F32 (chloroquine sensitive) and FcM29 (chloroquine resistant). The IC_{50} values obtained ranged from 1.4 to 21 μ M. Their cytotoxicity was estimated on human melanoma cells (A375) and the cytotoxicity/antiplasmodial ratio was found to be between 40 and 70 [123].

Other Constituents

Lapachol (**78**), a simple hydroxynaphthoquinone, is known for many pharmacological properties including antimalarial activity. It is present in many members of the Bignoniaceae family and it has been used as a template for the synthesis of the antimalarial drug atovaquone (**79**) [124].




In a very recent paper a naphthoquinone–anthraquinone coupled pigment named newbouldiaquinone A (**80**) together with other naphthoquinones isolated from *Newbouldia laevis* Seem. (Bignoniaceae), a tropical African species widely used for the treatment of several diseases including malaria, were tested *in vitro* against *P. falciparum* NF54 and R strains [125].




The most active compounds were newbouldiaquinone A, lapachol, α -lapachone and β -lapachone which showed a moderate suppression of parasitic growth [125].

Several papers report the isolation of active naphthoquinones from Bignoniaceae. Five furanonaphthoquinones isolated from *Tabebuia ochracea* ssp. *neochrysantha* (Bignoniaceae), a plant used traditionally in the Amazon to treat malaria, were tested against *P. falciparum* and *P. berghei* *in vitro*. The most active constituent was represented by a mixture of two compounds that could not be

separated: 5- and 8-hydroxy-2-(1'-hydroxyethyl)naphtho[2,3-*b*]furan-4,9-dione. The IC₅₀ values obtained with this mixture were 0.17 μ M (against *P. berghei*) and 0.67 μ M (against FcB2 chloroquine-resistant strain of *P. falciparum*). For the former parasite, the IC₅₀ value for chloroquine was 0.05 μ M, while for *P. falciparum* the IC₅₀ value was 0.11 μ M. These results indicate that the furanonaphthoquinones isolated from *T. ochracea* are potential antimalarial compounds [126]. Four naphthoquinoids isolated from *Kigelia pinnata* (Bignoniaceae) root bark were assessed *in vitro* against chloroquine-sensitive (T9-96) and chloroquine-resistant (K1) *Plasmodium falciparum* strains and for cytotoxicity using KB cells. The most active one, 2-(1-hydroxyethyl)naphtho[2,3-*b*]furan-4,9-dione (**81**), has good activity against both strains; IC₅₀ values were 627 nM for the K1 strain and 718 nM for the T9-96 strain [127].



Several novel structurally related, prenylated naphthoquinones (sterekunthals A and B, pyranokunthones A and B) and one novel prenylated anthraquinone (anthrakunthone) isolated from the root bark of *Stereospermum kunthianum* Cham (Bignoniaceae), a plant used in Uganda to treat fever, have been tested against the chloroquine-sensitive strain poW and the chloroquine-resistant clone Dd2. The quinones showed different degrees of activity against the two strains of *P. falciparum* and sterekunthal A (**82**) was the most effective one [IC₅₀ values: 3.85 μ M (PoW); 1.18 μ M (Dd2)].





It was also shown that the 4-hydroxy group is an important structural feature for the antiparasmodial activity of these compounds, as sterekunthal B is distinctly less active than pinnatal [128]. The IC₅₀ values were comparable to those of related naphthoquinones isolated from *Kigelia pinnata* DC [127]. On the other hand, these compounds also exhibited marked toxicity against endothelial ECV-304 cells and hence their antiparasmodial effect seems to be due to general cytotoxicity [129].

A number of isofuranonaphthoquinones isolated from *Bulbine capitata* Poelln. (Asphodelaceae) showed only weak antiparasmodial activity both against the 3D7 (chloroquine-sensitive) and the K1 (chloroquine-resistant) strains. The plant is used in Botswana for its claimed antibiotic and antipyretic properties. The IC₅₀ values for both strains were between 23 and 92 μ M, suggesting that these compounds are unlikely to have a significant *in vivo* activity when used alone [129].

Plumbagin (5-hydroxy-2-methyl-1,4-naphthoquinone), isolated from *Nepenthes thorelii*, a species related to *N. ampullaria* and used to treat malaria in Malaysia, was active against *P. falciparum*, with an IC₅₀ value of 0.27 μ M. The quinone structure is believed to be essential for the activity whereas the presence of a heteroatom such as oxygen or chlorine in synthetic derivatives at position 3 of the naphthoquinone nucleus causes weakening or loss of activity [130].

Another interesting group of constituents tested for antimalarial activity are the anthranoids. A highly active derivative of this class is vismione H (**83**), isolated from *Vismia guineensis* (Clusiaceae). The IC₅₀ against the sexual erythrocytic stages of *P. falciparum* (NF 54, clone A1A9) was 0.23 μ M [131].



From another species of *Vismia*, *V. orientalis* Engl., a plant used in traditional medicine in Tanzania, vismione D (**84**) was isolated and exhibited antiparasmodial activity against *Plasmodium falciparum* strain K1 (IC₅₀ 2.4 μ M). However, it was also found slightly cytotoxic against human L6 cells (IC₅₀ 10 μ M) [132].

Concluding remarks

The prevalence of malaria in tropical zones worldwide, together with the lack of a vaccine and the appearance of strains of malaria parasite resistant to commercially available anti-malarial drugs based

on quinoline derivatives, makes the search for new effective anti-malarial drugs a global demand.

From the examination of the literature of the last decades it appears that a large number of plants used as antimalarial in the traditional medicine or related species have been investigated. Bioassay-guided fractionation of the extracts was generally used to find the active constituents and a large number of non-nitrogenous molecules have been found to possess a moderate to high *in vitro* antiplasmodial activity. However, only a few compounds have also been tested for *in vivo* antimalarial activities. Based on the literature compilation reported here the following three main conclusions can be drawn.

Firstly, only a few molecules result possessing a moderate to high activity and therefore should be considered for further investigations. They including peroxide sesquiterpenes, quinoid triterpenes, quassinoids, gallic acid derivatives, lignans, flavonoids and biflavonoids, xanthones, naphthoquinones and phenylanthraquinones.

Secondly, cytotoxicity of many of these derivatives has been evaluated in order to obtain the selectivity index, and results indicate that cytotoxicity and antimalarial activity are generally not correlated. It would be highly advantageous to consider these molecules as potential new antimalarial drugs.

Thirdly, although some of the investigated compounds are not particularly active, they are nevertheless interesting because they might strengthen chloroquine activity or restore chloroquine sensitivity in resistant strains of *P. falciparum*. Partially effective treatments might be beneficial in that the course of the disease is shortened, perhaps reducing anaemia and lowering the risk of death or serious illness from other anaemia-related diseases. Other possible benefits could be the alleviation of symptoms such as pain and fever and immunomodulation leading to increased immunity.

Another important aspect, not yet developed, is the search for molecules with little or no antiplasmodial activity which can synergistically act with known antimalarial drugs against *Plasmodium*. Thus, it is known that several flavonoids of *A. annua* can promote and enhance the antiplasmodic activity of artemisinin [133, 134], and recently it has been demonstrated that epigallocatechin gallate, epicatechin gallate and green tea extract not only have moderate antiplasmodial activity but also produce synergism in the presence of sublethal doses of artemisinin [107]. Also these molecules could have an important role in fighting malaria.

Acknowledgments - The financial support of MIUR (PRIN 2004) and Ente Cassa di Risparmio di Firenze is gratefully acknowledged.

References

- [1] Winstanley PA. (2000) Chemotherapy for *falciparum* malaria: the armoury, the problems and the prospects. *Parasitology Today*, **16**, 146-153.
- [2] Trigg PI, Kondrachine AV. (1998) Commentary: malaria control in the 1990s. *Bulletin World Health Organization*, **76**, 11-16.
- [3] Trape J-F, Pison S, Spiegel A, Enel C, Rogier C. (2002) Combating malaria in Africa. *Trends in Parasitology*, **18**, 224-230.
- [4] del Rayo Camacho Corona M, Croft SL, Phillipson JD. (2000) Natural Products as a source of antiprotozoal drugs. *Current Opinion in Anti-infective Investigational Drugs*, **2**, 47-62.
- [5] Schwikard S, van Heerden FR. (2002) Antimalarial activity of plant metabolites. *Natural Product Reports*, **19**, 675-692.
- [6] Tasdemir D, Güner ND, Perozzo R, Brun R, Dönmez AA, Çalis I, Rüedi P. (2005) Anti-protozoal and plasmodial FabI enzyme inhibiting metabolites of *Scrophularia lepidota* roots. *Phytochemistry*, **66**, 355-362.
- [7] Tchuendem MHK., Mbah JA, Tsopmo A, Ayafor JF, Sterner O, Okunje CC, Iwu MW, Schuster BM. (1999) Anti-plasmodial sesquiterpenoids from the African *Reneilimia cinnamata*. *Phytochemistry*, **52**, 1095-1099.
- [8] François G, Passreiter CM, Woerdenbag HJ, Van Looveren M. (1996) Antiplasmodial activities and cytotoxic effects of aqueous extracts and sesquiterpene lactones from *Neurolena lobata*. *Planta Medica*, **62**, 126-129.
- [9] Oketch-Rabah HA, Lemmich E, Dossaji SF, Theander TG, Olsen CE, Cornett C, Kharazmi A, Christensen SB. (1997) Two new antiprotozoal 5-methylcoumarins from *Vernonia brachycalyx*. *Journal of Natural Products*, **60**, 458-461.
- [10] Yu HW, Wright CW, Cai Y, Phillipson JD, Kirby GC, Warhurst DC. (1994) Antiprotozoal activities of *Centipeda minima*. *Phytotherapy Research*, **8**, 436-438.
- [11] Bischoff TA, Kelley CJ, Karchesy Y, Laurantos M, Nguyen-Dinh P, Arefi AG. (2004) Antimalarial activity of Lactucin and Lactucopicrin: sesquiterpene lactones isolated from *Cichorium intybus* L. *Journal of Ethnopharmacology*, **95**, 455-457.
- [12] Passreiter CM, Medillana Aldana BE. (1998) Variability of sesquiterpene lactones in *Neurolena lobata* of different origin. *Planta Medica*, **64**, 427-430.

- [13] Oketch-Rabah HA, Christensen SB, Frydenvang K, Dossaji SF, Theander TG, Cornett C, Watkins WM, Kharazmi A, Lemmich E. (1998) Antiprotozoal properties of 16,17-dihydrobrachycaloxolide from *Vernonia brachycalyx*. *Planta Medica*, **64**, 559-562.
- [14] François G, Passreiter CM. (2004) Pseudoguaianolide sesquiterpene lactones with high activities against the human malaria parasite *Plasmodium falciparum*. *Phytotherapy Research*, **18**, 184-186.
- [15] Lang G, Passreiter CM, Wright CW, Filipowicz NH, Addae-Kyereme J, Medinilla BE, Castillo JJ. (2002) Antiplasmodial activities of sesquiterpene lactones from *Eupatorium semialatum* *Zeitschrift für Naturforschung*, **57c**, 282-286.
- [16] Achenbach H, Waibel R, Nkunya MHH, Weenen H. (1992) Antimalarial compounds from *Hoslundia opposita*. *Phytochemistry*, **31**, 3781-3784.
- [17] Weenen H, Nkunya MH, Bray DH, Mwasumbi LB, Kinabo LS, Kilimali VAEB. (1990) Antimalarial activity of Tanzanian medicinal plants. *Planta Medica*, **56**, 368-370.
- [18] Asili J, Lambert M, Ziegler HL, Stärk D, Sairafianpour M, Witt M, Asghari G, Ibrahimi IS, Jaroszewski JW. (2004) Labdanes and isopimaranes from *Platycladus orientalis* and their effects on erythrocyte membrane and on *Plasmodium falciparum* growth in the erythrocyte host cells. *Journal of Natural Products*, **67**, 631-637.
- [19] Duker-Eshun G, Jaroszewski JW, Asomaning WA, Oppong-Boachie F, Olsen CE, Christensen SB. (2002) Antiplasmodial activity of labdanes from *Aframomum latifolium* and *Aframomum scepstrum*. *Planta Medica*, **68**, 642-644.
- [20] del Rayo Camacho M, Phillipson JD, Croft SL, Kirby GC, Warhurst DC, Solis PN. (2001) Assessment of the antiprotozoal activity of *Galphimia glauca* and the isolation of new nor-secofriedelanes and nor-friedelanes. *Phytochemistry*, **56**, 203-210.
- [21] Jullian V, Bonduelle C, Valentin A, Acebey L, Duigou A-G, Prévost M-F, Sauvain M. (2005) New clerodane diterpenoids from *Laetia procera* (Poepp.) Eichler (Flacourtiaceae), with antiplasmodial and antileishmanial activities. *Bioorganic & Medicinal Chemistry Letters*, **15**, 5065-5070.
- [22] Kanokmedhakul S, Kanokmedhakul K, Kanarsa T, Buayairaksa M. (2005) New Bioactive Clerodane Diterpenoids from the Bark of *Casearia grewiifolia*. *Journal of Natural Products*, **68**, 183-188.
- [23] Bringmann G, Saeb W, Aké Assi L, François G, Narayanan ASS, Peters K, Peters E-M. (1997) Betulinic acid: isolation from *Triphyophyllum peltatum* and *Ancistrocladus heyneanus*, antimalarial activity and crystal structure of benzyl ester. *Planta Medica*, **63**, 255-257.
- [24] Steele JCP, Warhurst DC, Kirby GC, Simmonds MSJ. (1999) *In vitro* and *in vivo* evaluation of betulinic acid as an antimalarial. *Phytotherapy Research*, **13**, 115-119.
- [25] Suksamrarn A, Tanachatchairatana T, Kanokmedhakul S. (2003) Antiplasmodial triterpenes from twigs of *Gardenia saxatilis* *Journal of Ethnopharmacology*, **88**, 275-277.
- [26] Suksamrarn S, Phanseeta P, Kunchanawatta K, Distaporn T, Ruktansing S, Suksamrarn A. (2006) Ceanothane- and Lupane-Type Triterpenes with Antiplasmodial and Antimycobacterial Activities from *Ziziphus cambodian*. *Chemical and Pharmaceutical Bulletin*, **54**, 535-537.
- [27] Takahara M, Kusano A, Shibano M, Kusano G, Koizumi K, Suzuki R, Kim H-S, Wataya Y. (1998) Antimalarial activity and nucleoside transport inhibitory activity of the triterpenic constituents of *Cimicifuga* spp. *Biological and Pharmaceutical Bulletin*, **21**, 823-828.
- [28] Pavanand K, Webster HK, Yongvanitchit K, Kun-anake A, Dechatiwongse T, Nutakul W, Bansiddhi J. (1989) Schizontocidal activity of *Celastrus paniculatus* Willd. against *Plasmodium falciparum* *in vitro*. *Phytotherapy Research*, **3**, 136-139.
- [29] Figueiredo JN, Răz B, Séquin U. (1998) Novel quinone methides from *Salacia kraussii* with *in vitro* antimalarial activity. *Journal of Natural Products*, **61**, 718-723.
- [30] Sairafianpour M, Christensen J, Staerk D, Budnik BA, Kharazmi A, Bagherzadeh K, Jaroszewski JW. (2001) Leishmanicidal, antiplasmodial, and cytotoxic activity of novel diterpenoid 1,2-quinones from *Perovskia abrotanoides*: new source of tanshinones. *Journal of Natural Products*, **64**, 1398-1403.
- [31] van Agtmael MA, Eggelte TA, van Bostel CJ. (1999). Artemisinin drugs in the treatment of malaria: from medicinal herb to registered medication. *Trends in Pharmacological Sciences*, **20**, 199-204.
- [32] Klayman DL. (1985) Qinghaosu (artemisinin): an antimalarial drug from China. *Science*, **31**, 1049-1055.
- [33] Wilairatana P, Looareesuwan S. (2002) The clinical use of artemisinin and its derivatives in the treatment of malaria. Wright CW (Ed). Taylor and Francis, London, UK, 289-307.
- [34] Rücker G, Walter RD, Manns D, Mayer R. (1991) Antimalarial activity of some natural peroxides. *Planta Medica*, **57**, 295-296.
- [35] Cubucku B, Bray DH, Warhurst DC, Mericli AH, Ozhatay N, Sariyar G. (1990) *In vitro* antimalarial activity of crude extracts and compounds from *Artemisia abrotanum* L. *Phytotherapy Research*, **4**, 203-204.
- [36] Liang X-T, Yu DQ, Wu WL, Deng HC. (1979) The structure of yingzhaosu A. *Acta Chimica Sinica*, **37**, 215-230.
- [37] Xu XX, Zhu J, Huang DZ, Zhou WS. (1991) Total synthesis of (+)-Yingzhaosu A. *Tetrahedron Letters*, **32**, 5785-5788.
- [38] Hofheinz W, Burgin H, Gocke E, Jaquet C, Masciadri R, Schmid G, Stohler H, Urwyler H, (1994) Ro42-1611 (arteferene), a new effective antimalarial: chemical structure and biological activity. *Tropical Medicinal Parasitology*, **45**, 261-265.

- [39] Jaquet C, Stohler HR, Chollet J, Peters W. (1994) Antimalarial activity of the bicyclic peroxide Ro 42-1611 (arteflene) in experimental models. *Tropical Medicine and Parasitology*, **45**, 266–271.
- [40] Borstnik K, Paika I, Shapiro TA, Posner GH. (2002) Chemotherapeutic peroxides: artemisinin, yingzhaosu A and related compounds. *International Journal for Parasitology*, **32**, 1661–1667.
- [41] Pollack Y, Segal R, Golenser J. (1990) The effect of ascaridole on the *in vitro* development of *Plasmodium falciparum*. *Parasitology Research*, **76**, 570–572.
- [42] Rücker G, Walter RD, Manns D, Mayer R. (1991) Antimalarial activity of some natural peroxides *Planta Medica*, **57**, 295–296.
- [43] Thebtaranonth C, Thebtaranonth Y, Wanauppathamkul S, Yuthavong Y. (1995) Antimalarial sesquiterpenes from tubers of *Cyperus rotundus*: structure of 10,12-peroxycalamenene, a sesquiterpene endoperoxide *Phytochemistry*, **40**, 125–128.
- [44] Schulte KE, Rücker G, Glauch G. (1967) On some components of underground parts from *Nardostachys chinensis* Batalin. *Planta Medica*, **15**, 274–281.
- [45] Takaya Y, Kurumada K-I, Takeuji Y, Kim H-S, Shibata S, Ikemoto N, Wataya Y, Oshima Y. (1998) Novel antimalarial guaiane-type sesquiterpenoids from *Nardostachys chinensis* roots *Tetrahedron Letters*, **39**, 1361–1364.
- [46] Takaya Y, Takeuji Y, Akasaka M, Nakagawasai O, Tadano T, Kisara K, Kim H-S, Wataya Y, Niwa M, Oshima Y. (2000) Novel guaiane endoperoxides, Nardoguaianone A–D, from *Nardostachys chinensis* roots and their antinociceptive and antimalarial activities. *Tetrahedron*, **56**, 7673–7678.
- [47] Kamchonwongpaisan S, Nilanont C, Tarnchompoo B, Thebtaranonth C, Thebtaranonth Y, Yuthavong Y, Kongsaree P, Clardy J. (1995) An antimalarial peroxide from *Amomum krervanh* Pierre. *Tetrahedron Letters*, **36**, 1821–1824.
- [48] Phillipson JD, O'Neill MJ. (1986) Novel antimalarial drugs from plants. *Parasitology Today*, **2**, 355.
- [49] Phillipson JD, Wright CW, Kirby GC, Whurst DC. (1995) Phytochemistry of some plants used in traditional medicine for the treatment of protozoal diseases. In *Phytochemistry of the Plants Used in Traditional Medicine*. Hostettmann K, Marston A, Maillard M, Hamburger M. (Eds.) Oxford University Press, London UK. 95–136.
- [50] Spencer CF, Koniuszy FR, Rogers EF, Shavel J, Easton NR, Kaczka EA, Kuehl FA, Phillips RF, Walti A, Folkers K. (1947) Survey of plants for antimalarial activity. *Lloydia*, **10**, 145–148.
- [51] Trager W, Polonsky J. (1981) Antimalarial activity of quassinoids against chloroquine-resistant *P. falciparum* *in vitro*. *American Journal of Tropical Medicine and Hygiene*, **30**, 531–537.
- [52] Anderson MM, O'Neill MJ, Phillipson JD, Warhurst DC. (1991) *In vitro* cytotoxicity of a series of quassinoids from *Brucea javanica* fruits against KB cells. *Planta Medica*, **57**, 62–64.
- [53] O'Neill MJ, Bray DH, Boardman P, Chan KL, Phillipson JD. (1987) Plants as sources of antimalarial drugs, Part 4: Activity of *Brucea javanica* fruits against chloroquine-resistant *Plasmodium falciparum* *in vitro* and against *Plasmodium berghei* *in vivo*. *Journal of Natural Products*, **50**, 41–48.
- [54] O'Neill MJ, Bray DH, Boardman P, Wright CW, Phillipson JD, Warhurst DC, Gupta MP, Correya M, Solis P. (1988) Plants as sources of antimalarial drugs, Part 6: Activities of *Simarouba amara* fruits. *Journal of Ethnopharmacology*, **22**, 183–190.
- [55] Okunade AL, Bikoff RE, Casper SJ, Oksman A, Goldberg DE, Lewis WH. (2003) Antiplasmodial activity of extracts and quassinoids isolated from seedlings of *Ailanthus altissima* (Simaroubaceae). *Phytotherapy Research*, **17**, 675–677.
- [56] Moretti C, Deharo E, Sauvain M, Jardel C, David PT, Gasquet M. (1994) Antimalarial activity of cedronin. *Journal of Ethnopharmacology*, **43**, 57–61.
- [57] Cabral JA, McChesney JD, Milhous WK. (1993) A new antimalarial quassinoid from *Simaba guianensis*. *Journal of Natural Products*, **56**, 1954–1961.
- [58] Kuo P-C, Damu AG, Lee K-H, Wu T-S. (2004) Cytotoxic and antimalarial constituents from the roots of *Eurycoma longifolia* *Bioorganic & Medicinal Chemistry*, **12**, 537–544.
- [59] François G, Diakanamwa C, Timperman G, Bringmann G, Steenackers T, Atassi G, Van Looveren M, Holenz J, Tassin J-P, Aké Assi L, Vanhaelen-Fastre R, Vanhaelen M. (1998) Antimalarial and cytotoxic potential of four quassinoids from *Hannoa chlorantha* and *Hannoa klaineana*, and their structure-activity relationships. *International Journal of Parasitology*, **28**, 635–640.
- [60] Fandeur T, Moretti C, Polonsky J. (1985) *In vitro* and *in vivo* assessment of the antimalarial activity of sergeolide. *Planta Medica*, **51**, 20–23.
- [61] Kitagawa H, Mahmud T, Yokota K, Nakagawa S, Mayumi T, Kobayashi M, Shibuya H. (1996) Indonesian medicinal plants. XVII. Characterization of quassinoids from the stems of *Quassia indica*. *Chemical and Pharmaceutical Bulletin*, **44**, 2009–2014.
- [62] Lee K-H, Tani S, Imakura Y. (1987) Antimalarial agents, 4. Synthesis of a brusatol analog and biological activity of brusatol-related compounds. *Journal of Natural Products*, **50**, 847–851.
- [63] Omar S, Zhang J, MacKinnon S, Leaman D, Durst T, Philogene BJR, Arnason JT, Sanchez-Vindas PE, Poveda L, Tamez PA, Pezzuto JM. (2003) Traditionally-used antimalarials from the Meliaceae. *Current Topics in Medicinal Chemistry*, **3**, 133–139.
- [64] Roy A, Saraf S. (2006) Limonoids: Overview of Significant Bioactive Triterpenes Distributed in Plants Kingdom. *Biological & Pharmaceutical Bulletin*, **29**, 191–201.

- [65] Khalid SA, Farouk A, Geary TG, Jensen JB. (1986) Potential antimalarial candidates from African plants: An *in vitro* approach using *Plasmodium falciparum*. *Journal of Ethnopharmacology*, **15**, 201-209.
- [66] Bray DH, Warhurst DC, Connolly JD, O'Neill MJ, Phillipson JD. (1990) Plants as sources of antimalarial drugs VI. Activity of some Melillaceae plants and their constituents limonoids. *Phytotherapy Research*, **4**, 29-35.
- [67] Bickii J, Njifutue N, Foyere JA, Basco LK, Ringwald P. (2000) *In vitro* antimalarial activity of limonoids from *Khaya grandiflora* CDC (Meliaceae). *Journal of Ethnopharmacology*, **69**, 27-33.
- [68] Omar S, Godard K, Ingham A, Hussain H, Wongpanich V, Pezzuto J, Durst T, Eklu C, Gbeassor M, Sanchez-Vidas P, Poveda L, Philogene BJR. (2003) Antimalarial activities of gedunin and 7-methoxygedunin and synergistic activity with dillapiol. *Annals of Applied Biology*, **143**, 135-141.
- [69] Joshi SP, Rojatkhar SR, Nagasampagi BA. (1998) Antimalarial activity of neem (*Azadirachta indica*) *Journal of Medicinal and Aromatic Plant Science*, **20**, 1000-1003.
- [70] Jones IW, Denholm AA, Ley SV, Lovell H, Wood A, Sinden RE. (1994) Sexual development of malaria parasites is inhibited *in vitro* by the neem extract azadirachtin, and its semi-synthetic analogues. *FEMS Microbiological Letters*, **120**, 267-273.
- [71] Dhar R, Zhang K, Talwar GP, Garg S, Kumar N. (1998) Inhibition of the growth and development of asexual and sexual stages of drug-sensitive and resistant strains of the human malaria parasite *Plasmodium falciparum* by Neem (*Azadirachta indica*) fractions *Journal of Ethnopharmacology*, **61**, 31-39.
- [72] Horgen FD, Madulid DA, Angerhofer CK, Pezzuto JM, Soejarto DD, Farnsworth NR. (1997) Isolation of gallic acid esters as antiplasmodial constituents of *Swintonia foxyworthii* (Anacardiaceae). *Phytomedicine*, **4**, 353-356.
- [73] Subeki S, Matsuura H, Takahashi K, Yamasaki M, Yamato O, Maeda Y, Katakura K, Kobayashi S, Trimurningsih T, Chairul C, Yoshihara T. (2005) Anti-babesial and anti-plasmodial compounds from *Phyllanthus niruri*. *Journal of Natural Products*, **68**, 537-539.
- [74] Verotta L, Dell'Agli M, Giolito A, Guerrini M, Cabalion P., Bosisio E. (2001) *In vitro* antiplasmodial activity of extracts of *Tristanopsis* species and identification of the active constituents ellagic acid and 3,4,5-trimethoxyphenyl-(6'-O-galloyl)-O-beta-D-glucopyranoside. *Journal of Natural Products*, **64**, 603-607.
- [75] Deck LM, Royer RE, Chamblee BB, Hernandez VM, Malone RR, Torres JE, Hunsaker LA, Piper RC, Makler MT, Vander Jagt DL. (1998) Selective inhibitors of human lactate dehydrogenases and lactate dehydrogenase from the malarial parasite *Plasmodium falciparum*. *Journal of Medicinal chemistry*, **41**, 3879-3887.
- [76] Boonlaksiri C, Oonant W, Kongsaree P, Kittakoo P, Tanticharoen M, Thebtaranonth Y. (2000) An antimalarial stilbene from *Artocarpus integer*. *Phytochemistry*, **54**, 415-417.
- [77] Duker-Eshun G, Jaroszewski JW, Asomaning WA, Oppong-Boachie F, Brøgger CS. (2004) Antiplasmodial constituents of *Cajanus cajan*. *Phytotherapy Research: PTR*, **18**, 128-130.
- [78] Kraft C, Jenett-Siems K, Siems K, Solis PN, Gupta MP, Bienzle U, Eich E. (2001) Andinermals A-C, antiplasmodial constituents from *Andira inermis*. *Phytochemistry*, **58**, 769-774.
- [79] Oketch-Rabah HA, Lemmich E, Dossaji SF, Theander TG, Olsen CE, Cornett C, Kharazmi A, Christensen SB. (1997) Two new antiprotozoal 5-methylcoumarins from *Vernonia brachycalyx*. *Journal of Natural Products*, **60**, 458-61.
- [80] Koeler I, Jenett-Siems K, Gonzalez JC, Hernandez MA, Ibarra RA, Berendsohm WG, Bienzle U., Eich E. (2001) *In vitro* antiplasmodial activity of 4-phenylcoumarins from *Exostema mexicanum*. *Planta Medica*, **67**, 89-91.
- [81] Lee K-H, Chai H-B, Tamez PA, Pezzuto JM, Cordell GA, Win KK, Tin-Wa M. (2003) Biologically active alkylated coumarins from *Kayea assamica*. *Phytochemistry*, **64**, 535-541.
- [82] Zhang H-J, Tamez PA, Hoang VD, Tan GT, Hung NV, Xuan LT, Huong LM, Cuong NM, Thao DT, Soejarto DD, Fong HHS, Pezzuto JM. (2001) Antimalarial compounds from *Rhaphidophora decursiva*. *Journal of Natural Products*, **64**, 772-777.
- [83] Ma C, Zhang HJ, Tan GT, Van Hung H, Cuong NM, Soejarto DD, Fong HHS. (2006) Antimalarial Compounds from *Grewia bilamellata*. *Journal of Natural Products*, **69**, 346-350.
- [84] Valsaraj R, Pushpangadan P, Smitt UW, Adsersen A, Christensen SB, Sittie A, Nyman U, Nielsen C, Olsen CE. (1997) New Anti-HIV-1, Antimalarial, and Antifungal Compounds from *Terminalia belleric*. *Journal of Natural Products*, **60**, 739-743.
- [85] Jensen JF, Kvist LP, Christensen SB. (2002) An Antiplasmodial Lignan from *Euterpe precatoria*. *Journal of Natural Products*, **65**, 1915-1917.
- [86] Kraft C, Jenett-Siems K, Köhler I, Tofern-Reblina B, Siems K, Bienzle U, Eich E. (2002) Antiplasmodial activity of sesquignans and sesquieolignans from *Bonamia spectabilis*. *Phytochemistry*, **60**, 167-173.
- [87] Chanphen R, Thebtaranonth Y, Wanauppathamkul S, Yuthavong Y. (1998) Antimalarial Principles from *Artemisia indica*. *Journal of Natural Products*, **61**, 1146-1150.
- [88] Chavi Y, Khanchara P, Supawadee D, Varima W, Prasat K. (2004) Bioactive flavonoids from *Kaempferia parviflora*. *Fitoterapia*, **75**, 89-92.
- [89] Murakami N, Mostaqul HM, Tamura S, Itagaki S, Horii T, Kobayashi M. (2001) New anti-malarial flavonol glycoside from *Hydrangeae Dulcis Folium*. *Bioorganic & Medicinal Chemistry Letters*, **11**, 2445-2447.

- [90] Beldjoudi N, Mambu L, Labaïed M, Grellier P, Ramanitrahasimbola D, Rasoanaivo P, Martin MT, Frappier F. (2003) Flavonoids from *Dalbergia louvelii* and their antiplasmodial activity. *Journal of Natural Products*, **66**, 1447-1450.
- [91] Yenesew A, Derese S, Midiwo JO, Oketch-Rabah HA, Lisgarten J, Palmer R, Heydenreich M, Peter MG, Akala H, Wangui J. (2003) Anti-plasmodial activities and X-ray crystal structures of rotenoids from *Millettia usaramensis* subspecies *usaramensis*. *Phytochemistry*, **64**, 773-779.
- [92] Chen M, Theander TG, Christensen SB, Hvidt L, Zhai L, Kharazmi A. (1994) Licochalcone A, a new antimalarial agent inhibits *in vitro* growth of the human malaria parasite *Plasmodium falciparum* and protects mice from *P. yoelii* infection. *Antimicrobial Agents and Chemotherapy*, **38**, 1470-75.
- [93] Kharazmi A, Chen M, Theander T, Christensen SB. (1997) Discovery of oxygenated chalcones as novel antimalarial agents. *Annals of Tropical Medicine and Parasitology*, **91**, S91-95.
- [94] Ziegler HL, Hansen HS, Staerk D, Christensen SB, Hägerstrand H, Jaroszewski JW. (2004) The antiparasitic compound licochalcone a is a potent echinocytogenic agent that modifies the erythrocyte membrane in the concentration range where antiplasmodial activity is observed. *Antimicrobial Agents and Chemotherapy*, **48**, 4067-4071.
- [95] Mi-Ichi F, Miyadera H, Kobayashi T, Takamiya S, Waki S, Iwata S, Shibata S, Kita K. (2005) Parasite mitochondria as a target of chemotherapy: inhibitory effect of Licochalcone A on the *Plasmodium falciparum* respiratory chain. *Annals of the New York Academy of Sciences*, **1056**, 46-54.
- [96] Frölich S, Schubert C, Bienzle U, Jenett-Siems K. (2005) *In vitro* antiplasmodial activity of prenylated chalcone derivatives of hops (*Humulus lupulus*) and their interaction with haemin. *Journal of Antimicrobial Chemotherapy*, **55**, 883-887.
- [97] Shweta N, Tanvir K, Srinivasa Rao M, Srivastava K, Puri SK. (2005) Prenylated chalcones isolated from *Crotalaria* genus inhibits *in vitro* growth of the human malaria parasite *Plasmodium falciparum*. *Bioorganic & Medicinal Chemistry Letters*, **15**, 2453-2455.
- [98] Liu M, Wilairat P, Croft SL, Tand A L-C, Go M-L. (2003) Structure-Activity Relationships of antileishmanial and antimalarial chalcones. *Bioorganic & Medicinal Chemistry*, **11**, 2729-2738.
- [99] Yenesew A, Derese S, Irungu B, Midiwo JO, Waters NC, Liyala P, Akala H, Heydenreich M, Peter MG. (2003) Flavonoids and isoflavonoids with antiplasmodial activities from the root bark of *Erythrina abyssinica*. *Planta Medica*, **69**, 658-663.
- [100] Yenesew A, Induli M, Derese S, Midiwo JO, Heydenreich M, Peter MG, Akala H, Wangui J, Liyala P, Waters NC. (2004) Anti-plasmodial flavonoids from the stem bark of *Erythrina abyssinica*. *Phytochemistry*, **65**, 3029-3032.
- [101] Andayi AW, Yenesew A, Derese S, Midiwo JO, Gitu PM, Jondiko OJ, Akala H, Liyala P, Wangui JW. (2006) Antiplasmodial flavonoids from *Erythrina saculeuxii*. *Planta Medica*, **72**, 187-189.
- [102] Kanokmedhakul S, Kanokmedhakul K, Nambuddee K, Kongsaree P. (2004) New bioactive prenylflavonoids and dibenzocycloheptene derivative from roots of *Dendrobium lanceolatum*. *Journal of Natural Products*, **67**, 968-972.
- [103] Jenett-Siems K, Mockenhaupt FP, Bienzle U, Gupta MP, Eich E. (1999) *In vitro* antiplasmodial activity of Central American medicinal plants. *Tropical Medicine and International Health*, **4**, 611-615.
- [104] Kirmizibekmez H, Calis I, Perozzo R, Brun R, Dönmez AA, Linden A, Rüedi P, Tasdemir D. (2004) Inhibiting activities of the secondary metabolites of *Phlomis brunneogaleata* against parasitic protozoa and plasmodial enoyl-ACP Reductase, a crucial enzyme in fatty acid biosynthesis. *Planta Medica*, **70**, 711-717.
- [105] van Baren C, Anao I, Leo Di Lira P, Debenedetti S, Houghton P, Croft S, Martino V. (2006) Triterpenic acids and flavonoids from *Satureja parvifolia*. Evaluation of their antiprotozoal activity. *Zeitschrift fuer Naturforschung C, Journal of Biosciences*, **61**, 189-192.
- [106] Tasdemir D, Lack G, Brun R, Rüedi P, Scapozza L, Perozzo R. (2006) Inhibition of *Plasmodium falciparum* fatty acid biosynthesis: evaluation of FabG, FabZ, and FabI as drug targets for flavonoids. *Journal of Medicinal Chemistry*, **49**, 3345-3353.
- [107] Sannella A, Messori L, Casini A, Vincieri FF, Bilia AR, Maiori G, Severini C. (2006) Antimalarial Properties of Green Tea: A First Report. Personal Communication.
- [108] Ahmed MS, Galal AM, Ross SA, Ferreira D, Elsohly MA, Ibrahim A-RS, Mossa JS, El-Feraly FS. (2001) A weakly antimalarial biflavanone from *Rhus retinorrhoea*. *Phytochemistry*, **58**, 599-602.
- [109] Nunome S, Ishiyama A, Kobayashi M, Otoguro K, Kiyohara H, Yamada H, Omura S. (2004) *In vitro* antimalarial activity of biflavonoids from *Wikstroemia indica*. *Planta Medica*, **70**, 72-76.
- [110] Weniger B, Vonthron-Sénécheau C, Kaiser B, Brun R, Anton R. (2006) Comparative antiplasmodial, leishmanicidal and antitrypanosomal activities of several biflavonoids. *Phytomedicine*, **13**, 176-180.
- [111] Mbwambo ZH, Kapingu MC, Moshi MJ, Machumi F, Apers S, Cos P, Ferreira D, Marais JPI, Vanden BD, Maes A. (2006) Antiparasitic activity of some xanthenes and biflavonoids from the root bark of *Garcinia livingstonei*. *Journal of Natural Products*, **69**, 369-372.
- [112] Bringmann G, Menche D, Bezabih M, Abegaz BM, Kaminsky R. (1999) Antiplasmodial activity of knipholone and related natural phenylanthraquinones. *Planta Medica*, **65**, 757-758.
- [113] Abegaz BM, Bezabih M, Msuta T, Brun R, Menche D, Mühlbacher J, Bringmann G. (2002) Gaboroquinones A and B and 4'-O-demethylknipholone-4'-O-beta-D-glucopyranoside, Phenylanthraquinones from the Roots of *Bulbine frutescens*. *Journal of Natural Products*, **65**, 1117-1121.

- [114] Koumaglo K, Gbeassor M, Nikabu O, de Souza C, Werner W. (1990) Effects of three compounds extracted from *Morinda lucida* on *Plasmodium falciparum*. *Planta Medica*, **58**, 533-34.
- [115] Sittie AA, Lemmich E, Olsen CE, Hviid L, Kharazmi A, Nkrumah FK, Christenson SB. (1999) Structure-activity studies: *in vitro* antileishmanial and antimalarial activities of anthraquinones from *Morinda lucida*. *Planta Medica*, **65**, 259-263.
- [116] Likhitwitayawuid K, Chanmahasathien W, Ruangrunsi N, Krungkrai J. (1989) Xanthenes with antimalarial activity from *Garciniadulcis*. *Planta Medica*, **64**, 281-282.
- [117] Likhitwitayawuid K, Phadungcharoen T, Krungkrai J. (1998) Antimalarial xanthenes from *Garcinia cowa*. *Planta Medica*, **64**, 70-72.
- [118] Haya A-E, Hélesbeuxa J-J, Duvala O, LabaRedb M, Grellieb P, Richommea P. (2004) Antimalarial xanthenes from *Calophyllum caledonicum* and *Garcinia vieillardii*. *Life Sciences*, **75**, 3077-3085.
- [119] Fotie J, Nkengfack AE, Rukunga G, Tolo F, Peter MG, Heydenreich M, Fomum ZT. (2003) *In vivo* antimalarial activity of some oxygenated xanthenes. *Annals of Tropical Medicine and Parasitology*, **97**, 683-688.
- [120] Dua VK, Ojha VP, Roy R, Joshi BC, Valecha N, Devi CU, Bhatnagar MC, Sharma VP, Subbarao SK. (2004) Anti-malarial activity of some xanthenes isolated from the roots of *Andrographis paniculata*. *Journal of Ethnopharmacology*, **95**, 247-251.
- [121] Laphookhieo S, Syers JK, Kiattansakul R, Chantapromma K. (2006) Cytotoxic and antimalarial prenylated xanthenes from *Cratoxylum cochinchinense*. *Chemical and Pharmaceutical Bulletin*, **54**, 745-747.
- [122] Ngouela S, Lenta BN, Nougoué DT, Ngoupayo J, Boyom FF, Tsamo E, Gut J, Rosenthal PJ, Connolly JD. (2006) Anti-plasmodial and antioxidant activities of constituents of the seed shells of *Symphonia globulifera* Linn f. *Phytochemistry*, **67**, 302-306.
- [123] Azebaze AGB, Meyer M, Valentin A, Nguemfo EL, Fomum ZT, Nkengfack AE. (2006) Prenylated xanthone derivatives with antiplasmodial activity from *Allanblackia monticola* STANER L.C. *Chemical & Pharmaceutical Bulletin*, **54**, 111-113.
- [124] Makinde JM, Amusan OO, Adesogan EK. (1988) The antimalarial activity of *Spathodea campanulata* stem bark extract on *Plasmodium berghei berghei* in mice. *Planta Medica*, **54**, 122-125.
- [125] Eyong KO, Folefoc GN, Kuete V, Beng VP, Krohn K, Hussain H, Nkengfack AE, Saefel M, Sarite SR, Hoerauf A. (2006) Newbouldiaquinone A: A naphthoquinone-anthraquinone ether coupled pigment, as a potential antimicrobial and antimalarial agent from *Newbouldia laevis*. *Phytochemistry*, **67**, 605-609.
- [126] Perez HA, Diaz F, Medina JD. (1997) Chemical Investigation and *in vitro* Antimalarial Activity of *Tabebuia ochracea* ssp. *neochrysantha*. *International Journal of Pharmacognosy*, **35**, 227-231.
- [127] Weiss CR, Moideen SV, Croft SL, Houghton PJ. (2000) Activity of extracts and isolated naphthoquinones from *Kigelia pinnata* against *Plasmodium falciparum*. *Journal of Natural Products*, **63**, 1306-1309.
- [128] Onegi B, Kraft C, Köhler I, Freund M, Jenett-Siems K, Siems K, Beyer G, Melzigd MF, Bienzlee U, Eich E. (2002) Antiplasmodial activity of naphthoquinones and one anthraquinone from *Stereospermum kunthianum*. *Phytochemistry*, **60**, 39-44.
- [129] Bezabih M, Abegaz BM, Dufall K, Croft K, Skinner-Adams T, Davis TM. (2001) Antiplasmodial and antioxidant isofuranonaphthoquinones from the roots of *Bulbine capitata*. *Planta Medica*, **67**, 340-344.
- [130] Likhitwitayawuid K, Kaewamatawong R, Ruangrunsi N, Krungkrai J. (1998) Antimalarial naphthoquinones from *Nepenthes torelia*. *Planta Medica*, **64**, 237-241.
- [131] François G, Steenackers T, Assi LA, Steglich W, Lamottke K, Holenz J, Bringmann G. (1999) Vismione H and structurally related anthranoid compounds of natural and synthetic origin as promising drugs against the human malaria parasite *Plasmodium falciparum*: structure-activity relationships. *Parasitology Research*, **85**, 582-588.
- [132] Mbwambo ZH, Apers S, Moshi MJ, Kapingu MC, Van Miert S, Claeys M, Brun R, Cos P, Pieters L, Vlietinck A. (2004) Anthranoid compounds with antiprotozoal activity from *Vismia orientalis*. *Planta Medica*, **70**, 706-710.
- [133] Elford BC, Roberts MF, Phillipson JD, Wilson RJ. (1987) Potentiation of the antimalarial activity of Qinghaosu by methoxylated flavones. *Transaction of the Royal Society of Tropical Medicine and Hygiene*, **81**, 434-436.
- [134] Bilia AR, Lazari D, Messori L, Taglioli V, Temperini C, Vincieri FF. (2002) Simple and rapid physico-chemical methods to examine action of antimalarial drugs with hemin: Its application to *Artemisia annua* constituents. *Life Sciences*, **70**, 769-778.

Natural Product Communications

List of Referees 2006

**The editors of *Natural Product Communications* wish
to thank the following scientists for kindly reviewing the articles submitted
to the journal**

Ahiahonu P, Canada
Agrawal A, USA
Ahmed AA, Egypt
Ahmed VU, Pakistan
Alagona G, Italy
Alali FQ, Jordan
Antus S, Hungary
Appendino G, Italy
Arnold N, Germany
Attard E, Malta
Avilov SA, Russia
Ayyad S, Egypt

Bailleul F, France
Baldovini N, France
Bankova V, Bulgaria
Banting L, UK
Barbosa Filho JM, Brazil
Barrero AF, Spain
Baser KHC, Turkey
Bhakuni RS, India
Bhutani KK, India
Biagi M, Italy
Bilia AR, Italy
Blanco EM, Spain
Bohm B, Canada
Braz-Filho R, Brazil
Bruno M, Italy

Calabria L, USA
Campos MG, Portugal
Capasso F, Italy
Carvalho MG, Brazil
Casanova J, France
Cerecetto H, Uruguay
Charrouf Z, Morocco
Chattopadhyay S, India
Chauhan PMS, India
Chopra M, UK
Chowdhury PK, India

Cimino G, Italy
Connolly JD, UK
Conserva L, Brazil
Cortés D, Spain
Crawford D, USA
Curini M, Italy

Del Rio J, Spain
Dembitsky VM, Israel
Demirci F, Turkey

El Sayed KA, USA
El-Seedi H, Sweden
Emerenciano V, Brazil
Ernst E, UK

Fales HM, USA
Fernández-Barrero A, Spain
Figueiredo AC, Portugal

Galal AM, Egypt
Ganapaty S, India
Garcia-Viguera C, Spain
Gaspar A, South Africa
Gibbons S, UK
Glasl S, Austria
Glivandete MP, Brazil
González-Coloma A, Spain
Gorham J, UK
Gorinstein S, Israel
Guyot M, France

Hamburger M, Switzerland
Hill RA, UK
Hisham A, Oman
Houghton P, UK

Ibrahim R, Canada
Iinuma M, Japan
Iwasa K, Japan

Jirovetz L, Austria
Joël B, France

Kalinin V, Russia
Karchesy J, USA
Kaushik N, India
Khare N, India
Khripach VA, Belarus
Kingston DGI, USA
Kondo R, Japan
Kouno I, Japan
Krasutsky PA, USA
Krishna NR, USA
Kutschy P, Slovak Republic

Lanzotti V, Italy
Lee HK, Korea
Lee HS, Korea
Lemos TLG, Brazil
Likhitwitayawuid K, Thailand
Lockwood B, UK
Loiseau P, France
Luc A, Belgium
Luo YR, USA

Mabry TJ, USA
Mahmoud AA, Saudi Arabia
Majinda RRT, Botswana
Makarieva TN, Russia
Marcello Nicoletti M, Italy
Marston A, Switzerland
Marzouk MSA, Egypt
Massiot G, France
May E, UK
Melzig MF, Germany
Miraldi E, Italy
Montaut S, Canada
Moraes SM, Brazil
Morris GA, UK
Mulder P, UK

List of Referees

Natural Product Communications Vol. **1** (1-12) 2006

Narender T, India
Nemeth E, Hungary
Nonato MG, Philippines

Oleszek W, Poland
Orabi KY, Kuwait
Orihara Y, Japan

Paré P, USA
Parente JP, Brazil
Perry NB, New Zealand
Pessoa ODL, Brazil
Pichette A, Canada
Pino Alea J, Cuba
Pintore G, Italy
Pizza C, Italy
Politi M, UK
Prakash O, USA
Priyadarsini KI, India

Quetin-Leclercq J, Belgium

Raestrelli L, Italy
Reichling J, Germany
Rojas J, Venezuela
Rondon M, Venezuela

Ross SA, USA
Rosselli S, Italy
Rossi-Bergmann B, Brazil
Russo A, Italy

Sahu NP, India
Sandor A, Hungary
Saunders K, UK
Sautreau AV, UK
Seifert K, Germany
Sener B, Turkey
Setzer WN, USA
Shen YC, Taiwan
Shin J, Korea
Singh B, India
Singh SB, USA
Smith JA, Australia
Sosa S, Italy
Spada A, Italy
Speranza G, Italy
Spring O, Germany
Stuessy T, Austria
Subbaraju GV, India
Suchy M, Canada
Tanaka T, Japan

Tanaka K, Japan
Ikeda T, Japan
Tane P, Cameroon
Teixeira VL, Brazil
Thurston DT, UK
Tinto WF, West Indies
Tomás-Barberán FA, Spain
Tripathi AK, India
Tykwinski R, Canada

Valant-Vetschera K, Austria
Vidari G, Italy

Wani MC, USA
Waterman PG, Australia
Wawer I, Poland
Wiesman Z, Israel
Williamson EM, UK
Wolfender J, Switzerland
Wollenweber E, Germany
Wong J, UK
Wu, CL, Taiwan

Yahara S, Japan

Zacchino S, Argentina

Natural Product Communications

2006

Volume 1

Natural Product Communications 1 (1-12) 1-1204 (2006)

ISSN 1934-578X (print)

ISSN 1555-9475 (online)

EDITOR-IN-CHIEF

DR. PAWAN K AGRAWAL

Natural Product Inc.
7963, Anderson Park Lane,
Westerville, Ohio, 43081 USA
agrawal@naturalproduct.us

EDITORS

PROFESSOR GERALD BLUNDEN

The School of Pharmacy & Biomedical Sciences,
University of Portsmouth,
Portsmouth, PO1 2DT U.K.
gands@surefish.co.uk

PROFESSOR ALESSANDRA BRACA

Dipartimento di Chimica Bioorganica e Biofarmacia,
Universita di Pisa,
via Bonanno 33, 56126 Pisa, Italy
Email: braca@farm.unipi.it

PROFESSOR DEAN GUO

State Key Laboratory of Natural and Biomimetic Drugs,
School of Pharmaceutical Sciences,
Peking University,
Beijing 100083, China
gda5958@163.com

PROFESSOR ERNST HASLINGER

Institute of Pharmaceutical Chemistry,
University of Graz,
A-8010 Graz, Austria
Ernst.Haslinger@uni-graz.at

PROFESSOR J. ALBERTO MARCO

Departamento de Química Orgánica,
Universidad de Valencia,
E-46100 Burjassot, Valencia, Spain
alberto.marco@uv.es

PROFESSOR YOSHIHIRO MIMAKI

School of Pharmacy,
Tokyo University of Pharmacy and Life Sciences,
Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan
mimakiy@ps.toyaku.ac.jp

PROFESSOR M. G. REINECKE

Department of Chemistry,
Texas Christian University,
Forts Worth, TX 76129, USA
m.reinecke@tcu.edu

PROFESSOR YASUHIRO TEZUKA

Institute of Natural medicine
Toyama Medical and Pharmaceutical University,
2630-Sugitani, Toyama 930-0194, Japan
tezuka@ms.toyama-mpu.ac.jp

ADVISORY BOARD

Prof. Oyvind Andersen
Bergen, Norway

Prof. Bruno Botta
Roma, Italy

Prof. Carlos Cerda-Garcia-Rojas
Mexico city, Mexico

Prof. Ioanna Chinou
Athens, Greece

Prof. Josep Coll
Barcelona, Spain

Prof. Geoffrey Cordell
Chicago, IL, USA

Prof. Samuel Danishefsky
New York, NY, USA

Dr. Biswanath Das
Hyderabad, India

Prof. Daneel Ferreira
Mississippi, MS, USA

Prof. A.A. Leslie Gunatillika
Tucson, AZ, USA

Prof. Stephen Hanessian
Montreal, Canada

Prof. Michael Heinrich
London, UK

Prof. Kurt Hostettmann
Lausanne, Switzerland

Prof. Martin A. Iglesias Arteaga
Mexico, D. F., Mexico

Prof. Norbert De Kimpe
Gent, Belgium

Prof. Jerzy Jaroszewski
Copenhagen, Denmark

Prof. Teodoro Kaufman
Rosario, Argentina

Prof. Marie Lacaille-Dubois
Dijon, France

Prof. Hartmut Laatsch
Göttingen, Germany

Prof. Hyeon-Kyu Lee
Daejeon, Korea

Prof. Shuei-Sheng Lee
Taipei, Taiwan

Prof. Chun-Nan Lin
Kaohsiung, China

Prof. Francisco Macias
Cadiz, Spain

Prof. Anita Marsaioli
Campinas, Brazil

Prof. Rachel Mata
Mexico D. F., Mexico

Prof. Imre Mathe
Szeged, Hungary

Prof. Joseph Michael
Johannesburg, South Africa

Prof. Virinder Parmar
Delhi, India

Prof. Luc Pieters
Antwerp, Belgium

Prof. Peter Proksch
Düsseldorf, Germany

Prof. Stephen Pyne
Wollongong, Australia

Prof. William Reynolds
Toronto, Canada

Prof. Raffaele Riccio
Salerno, Italy

Prof. Ricardo Riguera
Santiago de Compostela, Spain

Prof. Satyajit Sarker
Coleraine, UK

Prof. Monique Simmonds
Richmond, UK

Prof. Valentin Stonik
Vladivostok, Russia

Prof. Hermann Stuppner
Innsbruck, Austria

Prof. Apichart Suksamrarn
Bangkok, Thailand

Prof. Hiromitsu Takayama
Chiba, Japan

Prof. Paul Wender
Stanford, USA

INFORMATION FOR AUTHORS

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site <http://www.naturalproduct.us>.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the publisher. A photocopy license is available from the publisher for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

To Subscribe: Natural Product Communications (ISSN# 1555-9475) is a journal published monthly. 2006 subscription price: US\$1,095 (Web edition); US\$1,395 (Print); US\$1,795 (Print + single site online). Orders should be addressed to Subscription Department, Natural Product Communications, 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

Natural Product Communications

Contents of Volume 1 2006

Number 1

- 1 Alkamides from *Piper nigrum* L. and Their Inhibitory Activity against Human Liver Microsomal Cytochrome P450 2D6 (CYP2D6)
Subehan, Tepy Usia, Shigetoshi Kadota and Yasuhiro Tezuka
- 9 5,7-Dihydroxy-5,6,7,8-tetrahydro-1*H*-azocin-2-one from a Marine-derived *Streptomyces* sp.
Serge Fotso, Shao Jie Wu, Song Qin and Hartmut Laatsch
- 15 Jasomontanone, a Novel Bicyclic Sesquiterpene from the Leaves of *Jasonia Montana*
Ahmed A. Mahmoud
- 21 Lignans from the Stem of *Cinnamomum camphora*
TJ Hsieh, CH Chen, WL Lo and CY Chen
- 27 Clarification of the Saponin Composition of *Ranunculus ficaria* Tubers
Andrew Marston, Martine Cabo, Christian Lubrano, Jean-Renaud Robin, Claude Fromageot and Kurt Hostettmann
- 33 Hederacine A and Hederacine B from *Glechoma hederaceae* Inhibit the Growth of Colorectal Cancer Cells *in vitro*
Yashodharan Kumarasamy, Lutfun Nahar, Paul Kong-Thu-lin, Marcel Jaspars and Satyajit D. Sarker
- 37 Anticoagulant effect and Constituents of *Baccharis illinita*
Moacir Geraldo Pizzolatti, Luiz Gonzaga Verdi, Inês Maria C. Brighente, Terezinha de Jesus C. Neiva, Jan Schripsema and Raimundo Braz Filho
- 43 3,5-Trimethoxybenzene and 2,4,6-Trimethoxystyrene are the Major Components in the Leaf Oil of *Eugenia confusa* from Abaco Island, Bahamas
William N. Setzer, Joseph A. Noletto and Michael A. Vincent
- 47 Seasonal Variation in Bromophenol Content of *Polysiphonia lanosa*
Nagwa A. Shoeib, Michael C. Bibby, Gerald Blunden, Peter A. Linley, David J. Swaine and Colin W. Wright
- 51 First Synthesis of (±)-Monotesone B and New Syntheses of (±)-Lonchocarpol A and (±)-Bavachin
Ágnes Kenéz and Sándor Antus
- 57 Automatic Structure Elucidation through Data Base Search and 2D NMR Spectral Analysis
Jean-Marc Nuzillard and Vicente de Paulo Emerenciano
- 65 Pharmacological Properties of Crocetin and Crocin (Digentiobiosyl Ester of Crocetin) from Saffron
Liang Xi and Zhiyu Qian
- 77 Herbal remedies: promises with risk
Francesca Borrelli, Raffaele Capasso and Francesco Capasso

Number 2

- 81 C-Prenylflavonoids from *Derris heyneana*
Seru Ganapaty, Pannakal S. Thomas, Jangam S. Josaphine, Ni Ni Than and Hartmut Laatsch
- 87 Growth Inhibitory Polyacetylenes from Galls of *Hedera rhombea* Bean
Sayumi Yamazoe, Koji Hasegawa, Kiyotake Suenaga and Hideyuki Shigemori
- 95 A New Colchicinoid from *Colchicum tauri*, an Unexplored Meadow Saffron Native to Jordan
Feras Q. Alali, Amani S. Ma'aya'h, Ahmad Alkofahi, Amjad Qandil, Chen Li, Jason Burgess, Mansukh C. Wani and Nicholas H. Oberlies
- 101 Constituents of *Erythrina lysistemon*: Their Brine Shrimp Lethality, Antimicrobial and Radical Scavenging Activities
Benard F. Juma and Runner R. T. Majinda
- 109 Two New Dibenzylobutyrolactone Type Lignans from the Stems of *Kadsura heteroclita*
Wei Wang, Xiaochi Ma, Peng Liu, Rongxia Liu, Shuhong Guan and Dean Guo
- 113 Chemical Composition and Antibacterial Activity of the Essential Oil of *Lasiocephalus longipenicillatus* (*Senecio longipenicillatus*)
María Rondón, María Araque, Antonio Morales, María Gualtieri, Janne Rojas, Katalin Veres and Imre Máthé

Cumulative Index
Natural Product Communications Vol. 1 (1-12) 2006

- 117 Antimicrobial and Antioxidant Activities of the Essential Oil of Resin of *Protium heptaphyllum*
P. N. Bandeira, A. M. Fonseca, S. M. O. Costa, M. U. D. S. Lins, O. D. L. Pessoa, F. J. Q. Monte, N. A. P. Nogueira and T. L. G. Lemos
- 121 Distribution and Chemotaxonomic Significance of *N*-Methylprolines in Selected Plant Families
Gerald Blunden, Asmita V. Patel, Nigel Armstrong and Maricela Adrian Romero
- 131 Novel Synthesis of Prenylated Phenols and Their Antioxidant Properties
Soumyaditya Mula, Birija S. Patro, Govind P. Kalena and Subrata Chattopadhyay
- 139 Biogenic Iodine and Iodine-Containing Metabolites
Valery M Dembitsky

Number 3

- 177 Two New Lanostanoid Triterpenes from the Fruit Body of *Ganoderma lucidum*-the Major Component of SunRecome®
Shu-hong Guan, Min Yang, Xuan Liu, Jia-meng Xia, Xiao-ming Wang, Hui Jin and De-an Guo
- 183 Isolation and Identification of *neo*-Clerodane Diterpenes from *Ajuga nipponensis* Makino
Josep Coll and Yudelso A. Tandrón
- 191 Two New C₁₈-Diterpenoid Alkaloids from *Aconitum Piepunense*
Le Cai, Dong-Lin Chen and Feng-Peng Wang
- 195 The Composition and Antimicrobial Activity of the Essential Oil of *Teucrium stocksianum* subsp. *stocksianum* Leaf from Oman
Abdulkhader Hisham, Nirmal Pathare and Salim Al-Saidi
- 201 Analysis of the Leaf Essential Oil of *Stauranthus perforatus* from Monteverde, Costa Rica
Jennifer M. Schmidt and William N. Setzer
- 205 Comparative Study of the Chemical Composition of the Essential Oil of *Lippia oreganoides* Collected in Two Different Seasons in Venezuela
Janne Rojas, Antonio Morales, Sara Pasquale, Alejandro Márquez, María Rondón, Katalin Veres and Imre Máthé
- 209 Chemical Constituents and Effect of Topical Application of *Oleum Hyperici* on Skin Sensitivity to Simulated Sun Exposure
Elisabetta Miraldi, Marco Biagi and Daniela Giachetti
- 215 Bioproduction of Diosgenin in Callus Cultures of *Balanites aegyptiaca*: Effect of Growth Regulators, Explants and Somatic Embryogenesis
Bishnu P. Chapagain, Vinod Saharan, Dan Pelah, Ram C. Yadav and Zeev Wiesman
- 223 AFLP Analysis for Genetic Diversity in *Capsicum annuum* and Related Species
Sanjog T. Thul, Ajit K. Shasany, Mahendra P. Darokar and Suman P. S. Khanuja
- 229 A Theoretical Elucidation of the Radical-Scavenging Power of Cyanidin
Hong-Fang Ji, Hong-Yu Zhang and Liang Shen
- 237 The Stereoselective Synthesis of the Nonnatural Enantiomers of Communiols A-C. A Stereochemical Correction
Juan Murga, Eva Falomir, Miguel Carda and J. Alberto Marco
- 247 Structures and Biological Activities of Plant Glycosides: Cholestane Glycosides from *Ornithogalum saundersiae*, *O. thyrsoides* and *Galtonia candicans*, and Their Cytotoxic and Antitumor Activities
Yoshihiro Mimaki
- 255 Camptothecins: Some Recent Chemical Studies
Biswanath Das, Maddeboina Krishnaiah, Katta Venkateswarlu and Ratna Das

Number 4

- 265 Two New Sarasinosides from the Sponge *Melophlus sarasinorum*
Elena A. Santalova, Vladimir A. Denisenko, Pavel S. Dmitrenok, Dmitrii V. Berdyshev and Valentin A. Stonik
- 273 Guyonianin A and B, Two Polyester Diterpenes from Algerian *Euphorbia guyoniana*
Ahmed A. Ahmed, Noureddine Gherraf, Ashraf A. El-Bassuony, Salah Rhouati, Mahmoud H. Gad, Shinji Ohta and Toshifumi Hirata
- 283 Guaianolides from the Aerial Parts of *Centaurea hololeuca*
Sergio Rosselli, Antonella Maria Maggio, Rosa Angela Raccuglia, Monique S. J. Simmonds, Nelly A. Arnold and Maurizio Bruno
- 287 Radical Scavenging-Flavonoids from *Erythrina abyssinica*
Francis Machumi, Gomotsang Bojase-Moleta, Renameditswe Mapitse, Ishmael Masesane and Runner R. T. Majinda

Cumulative Index
Natural Product Communications Vol. 1 (1-12) 2006

- 293 Hydroperoxysterols from the Brazilian Brown Seaweeds *Dictyopteris justii* and *Spatoglossum schroederi* (Dictyotales): A Defensive Strategy Against Herbivory
V. L. Teixeira, J. P. Barbosa, F. D. Rocha, M. A. C. Kaplan, P. J. Houghton and R. C. Pereira
- 299 Triterpenoids and Sterols from the Stem of *Alnus formosana* Burk
Chien-Kuang Chen, Shiou-Ling Tuh, Chung-Hsiung Chen, Chen-Meng Kuo, and Shoei-Sheng Lee
- 303 Chemical Composition and Bioactivity of the Leaf Oil of *Calyptanthus pallens* (Poir.) Griseb. from Abaco Island, Bahamas
Anita Bansal, Amelia K. Boehme, Lauren C. Eiter, Jennifer M. Schmidt, William N. Setzer and Michael A. Vincent
- 307 An Alternative Approach for the Detection of Ethephon (2-Chlorethylphosphonic acid) Residues in Apples
Dietmar Kröpfel, Klaus Schweiger, Franz Siegfried Wagner and Elke Prettnner
- 313 3,3-Diisopentenyl-*N*-methyl-2,4-quinoldione from *Esenbeckia almarillia*: The Antitumor Activity of this Alkaloid and its Derivatives
Fátima M. Nunes, Bartholomeu A. Barros-Filho, Maria C. F. de Oliveira, Marcos C. de Mattos, Manoel Andrade-Neto, Francisco G. Barbosa, Jair Mafezoli, Raquel C. Montenegro, Cláudia Pessoa, Manoel O. de Moraes, Letícia V. Costa-Lotufo, Fabio C. S. Galetti, Celio L. Silva, and Ana O. De Souza
- 319 Clerodane Diterpenes from *Croton* Species: Distribution and a Compilation of their ¹³C NMR Spectral Data
Sebastião F. Palmeira Júnior, Lucia M. Conserva and José Maria Barbosa Filho

Number 5

- 345 Ganoderic Acid TR, a new Lanostanoid with 5 α -Reductase Inhibitory Activity from the Fruiting Body of *Ganoderma lucidum*
Jie Liu, Kuniyoshi Shimizu and Ryuichiro Kondo
- 351 Novel Metabolites from the Stem Bark of *Brombya* sp. nova (Gap Creek) (Rutaceae)
Clynton W. Halstead, Paul I. Forster and Peter G. Waterman
- 357 Three New C₁₉-Diterpenoid Alkaloids from *Delphinium bonvalotii*
Yong He, Dong-Lin Chen and Feng-Peng Wang
- 363 Polyanxanthone, a Xanthone from the Stem Bark of *Garcinia polyantha*
Justin Komguem, Alain M. Lannang, Jean G. Tangmouo, Gabin N. Louh, Fernande N. Ngounou, David Lontsi, Muhammad I. Choudhary and Beiban L. Sondengam
- 367 Antioxidant Activity of Metabolites from *Coleonema album* (Rutaceae)
Lindy L. Esterhuizen, Riaan Meyer and Ian A. Dubery
- 377 Isolation and Characterization of 2,4,5-Trimethoxy-benzaldehyde and 1-(2,4,5-trimethoxyphenyl)-Ethanone from *Pachypodanthium staudtii*, and their general Toxicity towards Brine Shrimps
Lutfun Nahar and Satyajit D. Sarker
- 381 The Potential Angiotensin-Converting Enzyme Inhibitory Activity of Oleanolic Acid in the Hydroethanolic Extract of *Crataegus monogyna* Jacq.
Everaldo Attard and Henrietta Attard
- 387 Volatile Constituents of *Geranium tuberosum* L. from Iran
Andrea Barra, Hassan Norouzi-Arasi, Sajjad Sedaghat-Sharehjini and Nicolas Baldovini
- 391 *Tephrosia toxicaria* Pers Essential Oil: Chemical Composition and Larvicidal Activity
Walber H. F. Ribeiro, Jackson N. Vasconcelos, Angela M. C. Arriaga, Maria C. F. de Oliveira, Manoel Andrade-Neto, Telma L. G. Lemos, Gilvandete M. P. Santiago, Ronaldo F. Nascimento and Jair Mafezoli
- 395 Chemical Composition and Antibacterial Activity of the Essential Oils of *Blainvillea rhomboidea* (Asteraceae)
Andreza Maria L. Pires, Maria Rose Jane R. Albuquerque, Edson P. Nunes, Vânia M. M. Melo, Edilberto R. Silveira and Otília Deusdênia L. Pessoa
- 399 Influence of Roasting and Seed Collection on Argan Oil Odorant Composition
Zoubida Charrouf, Halima El Hamchi, Silvia Mallia, Giuseppe Licitra and Dominique Guillaume
- 405 Acetylenic Terrestrial Anticancer Agents
Valery M Dembitsky and Dmitri O Levitsky

Number 6

- 431 New Oleanane Glycosides from the Roots of *Gomphrena macrocephala*
Minpei Kuroda, Taku Aoshima, Mitsue Haraguchi, Maria Cláudia Marx Young, Hiroshi Sakagami and Yoshihiro Mimaki

Cumulative Index
Natural Product Communications Vol. 1 (1-12) 2006

- 441 New Clerodane and Halimane Diterpenes from the Leaves and Woody Stems of *Casearia grayi* (Flacourtiaceae/Salicaceae)
Ashik Mosaddik, Paul I. Forster, Ron Booth and Peter G. Waterman
- 449 A Pregnane Derivative and an Anti-plasmodial Labdane Diterpenoid from the Stem Bark of *Turraenthus africanus*
Thomas M. Akam, Pierre Tane, Hippolyte K. Wabo, Joseph N. Yong, Samuel N. Y. Fanzo-Free, Joseph D. Connolly, Cameron Evans and Louis J. Farrugia
- 453 Butanolides from the Stem of *Cinnamomum kotoense*
Chung-Yi Chen
- 457 Flavonoid C-glycosides from *Pterocephalus sanctus* Growing in Egypt
Fahem A. Ahmed and Abdelaaty A. Shahat
- 461 Chemical Constituents from *Alnus formosana* Burk. II. Polar Constituents from the Leaves
Shoei-Sheng Lee, Su-Chang Chen, Chien-Kuang Chen, Chung-Hsiung Chen and Chen-Meng Kuo
- 465 A Phytochemical Investigation of *Nectandra membranacea* from Monteverde, Costa Rica
Xiujun Wu, Bernhard Vogler, William A. Haber and William N. Setzer
- 469 Nematicidal Activity of the Essential Oils from *Pilocarpus microphyllus* (Rutaceae) Samples
Nirla R. Romero, Manoel Andrade-Neto, Francisco J. T. Gonçalves, Raimundo R. G. Nascimento, Franciglauber S. Bezerra, Maria C. F. de Oliveira, Francisco G. Barbosa, Antonio M. E. Bezerra, Jair Mafezoli and José S. P. Sena
- 475 Galanthamine Content of Bulbs and *in vitro* Cultures of *Leucojum aestivum* L.
Mamadou F. Diop, Agata Ptak, Françoise Chrétien, Max Henry, Yves Chapleur and Dominique Laurain-Mattar
- 481 Antibiotic-induced Alterations in the Osmotic Resistance of Erythrocytes is Modulated by β -Carotene and L-Ascorbic Acid
Suaib Luqman, K V Obli Prabhu, Anirban Pal, Dharmendra Saikia, Mahendra P Darokar and Suman P S Khanuja
- 487 A Simple Differential Pulse Polarographic Method for the Determination of Artemisinin in *Artemisia annua* Chhanda
Debnath, Ernst Haslinger and Astrid Ortner
- 495 Chemical Evolution in the Asteraceae. The Oxidation-Reduction Mechanism and Production of Secondary Metabolites
Vicente P. Emerenciano, D. Cabrol-Bass, Marcelo J. P. Ferreira, Sandra A.V. Alvarenga, Antonio J. C. Brant, Marcus T. Scotti and Karina O. Barbosa
- 509 Some Biological Properties of Curcumin: A Review
Badreldin H. Ali, Husnia Marriif, Salwa A. Noureldayem, Amel O. Bakheit and Gerald Blunden

Number 7

- 523 Argolic acid A and Argolic Methyl Ester B, Two New Cyclopentano-monoterpenes Diol from *Nepeta argolica*
Ahmed A. Ahmed, Husaam E. Hassan, Mohamed F. Hegazy, Olga Tzakou, Maria Couladis, Abou El-Hamed H. Mohamed, Mohamed A. Abdella and Paul Paré
- 527 Chemical Constituents of *Gentiana macrophylla* Pall. from Shaanxi
Qianliang Chen, Wenji Sun, Guangzhong Tu, Zhangyan Shi and Yongmin Zhang
- 531 Bioactive Sesquiterpene Lactones from *Eupatorium kiurunense*
Ya-Ching Shen, Kuang-Liang Lo, Yao Haur Kuo and Ashraf Taha Khalil
- 537 A Novel Sesquiterpene Polyol Ester from the *Celastrus rosthornianus* with Anti-tumor Activities
Kuiwu Wang and Yuanjiang Pan
- 541 Leishmanicidal, Antifungal, and Cytotoxic Activity of Triterpenoid Glycosides Isolated from the Sea Cucumber *Neothytone gibbosa*
Rosalba Encarnacion-Dimayuga, Jesús Iván Murillo-Álvarez, Carsten Christophersen, Manuel Chan-Bacab, María Luz García Reiriz and Susana Zacchino
- 549 Prenylated 2-arylbenzofurans from two Species of *Artocarpus*
Yana M. Syah, Euis H. Hakim, Lukman Makmur, Valentina A. Kurdi, Emilio L. Ghisalberti, Norio Aimi and Sjamsul A. Achmad
- 553 Isolation and Structure of Cordifolin, a Novel Insecticidal Oxygenated Chalcone, from the stem of *Tinospora cordifolia* Miers
Najam A. Shakil and Dinesh B. Saxena
- 557 Oestrogenic Isoflavone Content in Natural Strains of Subterranean Clover (*Trifolium subterraneum* L.) from Sardinia
Aldo Tava, Luciano Pecetti, Alessia Bertoli and Efsio Piano
- 563 Antioxidant Effects on Lipid Peroxidation by Superoxide of Echinoisoflavanone and Sophoraisoflavanone D from *Sophora chrysophylla* Seem.
Shizuo Toda and Yoshiaki Shirataki

Cumulative Index
Natural Product Communications Vol. 1 (1-12) 2006

- 567 Chemical compositions of the bark essential oils of *Croton monteверdensis* and *Croton niveus* from Monteverde, Costa Rica
William N. Setzer
- 573 Larvicidal Activity against *Aedes aegypti* L. (Diptera: Culicidae) of Essential Oils of *Lippia* Species from Brazil
Gilvandete M. P. Santiag, Telma L. G. Lemos, Otilia D. L. Pessoa, Ângela M. C. Arriaga, Francisco J. A. Matos, Mary Anne S. Lima, Hécio S. Santos, Maria da Conceição L. Lima, Francisco G. Barbosa, João H. S. Luciano, Edilberto R. Silveira and Gustavo H. A. de Menezes
- 577 Alteration in the Respiratory Pathway of the Freshwater Fish *Channa punctatus* Induced by *Euphorbia royleana* Stem Bark Extract
Sudhanshu Tiwari and Ajay Singh
- 585 The Diterpenoids of the Genus *Marrubium* (Lamiaceae)
Franco Piozzi, Maurizio Bruno, Sergio Rosselli and Antonella Maggio
- 593 Dendrimers as Drug Carriers. A New Approach to Increase the Potential of Bioactive Natural Products
Costas Demetzos

Number 8

- 601 A Sesquiterpene, Clerodane Diterpenes and a Furanone from the Roots of *Casearia multinervosa* (Flacourtiaceae/Salicaceae)
Ashik Mosaddik and Peter G. Waterman
- 609 Diterpenoid Constituents from the Brown Alga *Dictyota menstrualis* (Dictyotaceae, Phaeophyta)
Diana N. Cavalcanti, Claudia M. Rezende, Angelo C. Pinto and Valéria L. Teixeira
- 613 Triterpenoids, Including One with Smooth Muscle Relaxant Activity, from *Rubus idaeus* (Raspberry) Leaves
Janne Rojas Vera, Christopher G. Dacke, Asmita V. Patel and Gerald Blunden
- 619 Anti-trypanosomal Alkaloids from *Xymalos monospora*
Dieudonne Ngamga, Pierre Tane, Donna Rattendi, Cyrus Bacchi, Christopher C. Okunji, Maurice M. Iwu, Brian M. Schuster and Olov Sterner
- 623 Flavonoids from *Rosa damascena* Mill.
Neeraj Kumar, Bikram Singh and Vijay K Kaul
- 627 Chemodiversity of Exudate Flavonoids in *Baccharis concinna* and Three Further South-American *Baccharis* species
Eckhard Wollenweber, Karin M. Valant-Vetschera and G. Wilson Fernandes
- 633 Induction Effects of Apigenin, Luteolin and Vinpocetin on Neutral Endopeptidase (NEP) and Angiotensin-Converting Enzyme Activity (ACE) of SK-N-SH Cells
Shereen Ayoub and Matthias F. Melzig
- 641 Bioactivity of Hirsutanolol, Oregonin and Genkwanin, Isolated from the Seeds of *Alnus glutinosa* (Betulaceae)
Yashodharan Kumarasamy, Philip J. Cox, Marcel Jaspars, Lutfun Nahar and Satyajit D. Sarker
- 645 Chemical Composition and Antimicrobial Activity of the Essential oil of *Saccocalyx satireioides* Coss. et Dur.
Hocine Laouer, Salah Akkal, Claire Debarnot, Bruno Canard, Uwe J. Meierhenrich and Nicolas Baldovini
- 651 *Pycnanthus angolensis* (Welw) Excell: Volatile Oil Constituents and Antimicrobial activity
Anica Simic, Dietmar Kroepfl, Nebojsa Simic and Isiaka A. Ogunwande
- 655 Antimicrobial Activity and Chemical Composition of Essential Oil of *Eupatorium glutinosum* (Lam.)
Hesham R. El-Seedi
- 661 Novel lapachol derivatives and their antioxidant activity
João P. S. Wenceslau, Dávila F. de Souza, Maria C. F. de Oliveira, Telma L. G. Lemos, Antonia L. de Sousa, Maria T. S. Trevisan and Marcos C. de Mattos
- 665 Pregnane Glycosides
Nilendu Panda, Sukdeb Banerjee, Nirup B. Mandal and Niranjan P. Sahu

Number 9

- 697 A New Ionone Glucoside and Terpenoid Constituents from *Achillea biebersteinii* and their Antifungal Activity
Ahmed A. Mahmoud and Shar S. Al-Shihry
- 705 Smooth Muscle Relaxant Triterpenoid Glycosides from *Rubus idaeus* (Raspberry) Leaves
Janne Rojas Vera, Christopher G. Dacke, Gerald Blunden and Asmita V. Patel

Cumulative Index
Natural Product Communications Vol. 1 (1-12) 2006

- 711 8-Oxoadenine, 9-Methyl-8-Oxoadenine, and trihydroxylated sterols from a Far Eastern Thorectidae Sponge
Tatyana N. Makarieva, Alla G. Guzii, Andrei S. Dmitrenok, Pavel S. Dmitrenok, Vladimir B. Krasokhin and Valentin A. Stonik
- 715 Occurrence in Kava Roots of Kava Lactone-Yielding Precursor(s)
Mani Naiker, Sunny Y Prasad, Ranjeeta D. Singh, Joslin A. Singh and Tevita N. Voro
- 721 Thiocarbamates from *Moringa oleifera*
Amit Tewari and Rajendra S. Bhakuni
- 727 Phenolics and iridoids of *Lippia alba*
Thierry Hennebelle, Sevser Sahpaz, Henry Joseph and François Bailleul
- 731 Flavonoid and Flavone C-Glycosides from *Dregea volubilis*
Nilendu Panda, Debayan Mandal, Nirup B. Mandal, Niranjan P. Sahu and Sukdeb Banerjee
- 735 Antioxidant and Anti-proliferative Active Constituents of *Tecoma stans* against Tumor Cell Lines
Mohamed SA Marzouk, Amira M Gamal-Eldeen, Mona A Mohamed and Mortada M El-Sayed
- 745 Polyphenolic Profile and Biological Study of *Salvia fruticosa*
Fatma A. Moharram, Ibrahim I. Mahmoud, Madeha R. Mahmoud and Samah A. Sabry
- 751 Characterization of Seed Oil Components from *Nephelium lappaceum* L.
Pinarosa Avato, Isabella Rosito, Paride Papadia and Francesco P. Fanizzi
- 757 The Chemical Composition of *Phymatidium delicatulum* and *P. tillandsioides* (Orchidaceae) Floral Oils
Mariza G. Reis, Rodrigo B. Singer, Renato Gonçalves and Anita J. Marsaioli
- 763 Volatile Constituents of *Ficus exasperata* leaves
Mubo A. Sonibare, Isiaka A. Ogunwande, Tameka M. Walker, William N. Setzer, Mike O. Soladoye and Emmanuel Essien
- 767 $TiCl_3$ as a New Catalyst for the Imino Diels-Alder Reaction
Narayan V. Mayekar, Sandip K. Nayak and Subrata Chattopadhyay
- 773 Acetylenic Aquatic Anticancer Agents and Related Compounds
Valery M Dembitsky, Dmitri O Levitsky, Tatyana A Glorizova and Vladimir V Poroikov

Number 10

- 813 *Phytochemical Studies on Stemona Plants: Isolation of New Tuberostemonine and Stemofoline Alkaloids.*
Thanapat Sastraruji, Araya Jatisatienr, Kritchaya Issakul, Stephen G. Pyne, Alison T. Ung, Wilford Lie and Morwenna C. Williams
- 819 Alkaloids from the Root of *Flueggea virosa*
Li-She Gan and Jian-Min Yue
- 825 Constituents of Medicinal Plants of Lombok: A New Antibacterial Alkaloid from *Voacanga foetida* (Bl.) Rolfe
Surya Hadi and John B. Bremner
- 831 A General and Efficient Approach to the Proposed Structures of Frog Toxins: The 5-Alkylindolizidines
Brendon S. Gourlay, Ian Little, John H. Ryan and Jason A. Smith
- 839 Progress on Cassaine-Type Diterpenoid Ester Amines and Amides (*Erythrophleum* Alkaloids)
Jing Qu, Shishan Yu, Wenzhao Tang, Yunbao Liu, Yue Liu, and Jing Liu
- 851 Alkaloids and Coumarins from *Ruta* Species
Ayhan Ulubelen and Mehmet Öztürk
- 859 Steroidal Glycoalkaloids: Isolation, Structure, Analysis, and Biosynthesis
Emilio L. Ghisalberti
- 885 N-Demethylation of Alkaloids
Shanti Thavaneswaran, Kristy McCamley and Peter J. Scammells
- 899 Asymmetric Synthesis of Quinine: A Landmark in Organic Synthesis
Vijay Nair, Rajeev S. Menon and Sreekumar Vellalath
- 907 Preparation of Cyclic Peptide Alkaloids Containing Functionalized Tryptophan Residues
Alexander K. L. Yuen and Craig A. Hutton

Number 11

- 921 Chemodiversity of Exudate Flavonoids, as Highlighted by Selected Publications of Eckhard Wollenweber
Karin M. Valant-Vetschera and Brigitte Brem
- 927 Two New Cyclopentanoids from the Endophytic Fungus *Aspergillus sydowii* Associated with the Marine Alga *Acanthophora spicifera*
Franka Teuscher, Wenhan Lin, Victor Wray, RuAngelie Edrada, K. Padmakumar, Peter Proksch and Rainer Ebel
- 935 5-Deoxynevadensin, a Novel Flavone in Sunflower and Aspects of Biosynthesis during Trichom Development
Jens Göpfert, Jürgen Conrad and Otmar Spring
- 941 Flavonol Triglycosides from the Leaves of *Silphium albiflorum*
Malgorzata Wojcińska, Jeffrey Williams, Tom J. Mabry, Ahmed A. Ahmed, Barry D. Davis, Gabor Tóth, Nabil H. El-Sayed, Irena Matlawska and Jennifer Clevinger
- 949 Two Novel Natural Flavonoids from *Primula palinuri*
Munekazu Iinuma, Toshiyuki Tanaka, Masayoshi Oyama and Eckhard Wollenweber
- 953 Bioactive Flavonoids from *Tephrosia purpurea*
Mona E. S. Kassem, Mohamed Sharaf, Manal H. Shabana and Nabil A. M. Saleh
- 957 Carbon-13 NMR Chemical Shift of Methyl Group: a Useful Parameter for Structural Analysis of C-Methylated Flavonoids
Pawan K. Agrawal, Chandan Agrawal and Shravan Agrawal
- 961 Antitumor and Immunostimulatory Activity of Two Chromones and Other Constituents from *Cassia petersiana*
Pierre C. Djemgou, Donatien Gatsing, Marguérite Tchuendem, Bonaventure T. Ngadjui, Pierre Tane, Ahmed A. Ahmed, Amira M. Gamal-Eldeen, Godwin I. Adoga, Toshifumi Hirata and Tom J. Mabry
- 969 Comparative Investigation of O- and C-Glycosylflavones in Leaves of Six *Santalum insulare* (Santalaceae) varieties
Jean-François Butaud, Phila Raharivelomanana, Denis Loquet, Jean-Pierre Bianchini, Robert Faure and Emile M. Gaydou
- 973 Comparative Analysis of Over-the-Counter Tablet Preparations of Isoflavones Extracted from Soy Available in Portugal
Maria G. Campos, António H. Paranhos, Miguel P. Matos, Maria T. Câmara, Margarida M. Cunha, Paula R.O. Pinto, Armando J.D. Silvestre, Francisco M. L. Amado and Carlos P. Neto
- 981 Characterization of a Novel Flavone O-Methyltransferase Gene in Rice
Jian-Min Zhou, Yukiharu Fukushima, Xiao-Feng Wang and Ragai K. Ibrahim
- 985 Biological Activities of Limonoids, Catechins, Procyanidins and Extracts from *Xylocarpus granatum*
Helle Wangenstein, Gia M. Duong, Mahiuddin Alamgir, Mokadez Sarder, Anne B. Samuelsen and Karl E. Malterud
- 991 New Insight into the Ring Contraction of 2'-Benzyloxyflavanones
István Németh, Katalin Gulácsi, Sándor Antus, Sándor Kéki and Miklós Zsuga
- 997 The B Group Vitamins and Mineral Elements in the Selective Removal of Wheat Kernel Layers
Biljana V. Vucelic-Radovic, Vjaceslav M. Nasic, Mirjana A. Demin and Mirjana M. Milovanovic
- 1003 Antibacterial Activity and Irritation Potential of Selected Essential Oil Components – Structure-Activity Relationship
Jürgen Reichling, Ulrike Suschke, Jürgen Schneele and Heinrich Konrad Geiss
- 1013 The *Lasthenia californica* Story: It Started with Flavonoids
Bruce A. Bohm and Nishanta Rajakaruna
- 1023 Plant Sources of Propolis: an Update from a Chemist's Point of View
Vassya Bankova, Milena Popova and Boryana Trusheva
- 1029 Distribution of Surface Flavonoids in Bulgarian Plants
M. Nikolova and S. Ivancheva
- 1037 Phytochemical Quality and Bioactivity of Edible Sprouts
Diego A. Moreno, Santiago Pérez-Balibrea and Cristina García-Viguera
- 1049 Nutritional Relevance of Flavonoids in Disease Prevention
Janet Kyle and Garry Duthie
- 1061 Translational Investigation of Turmeric for Arthritis Treatment: a Review of Lessons Learned
Janet L Funk and Barbara N Timmermann
- 1067 Plant Epicuticular Waxes: Chemistry, Form, Self-Assembly and Function
Kerstin Koch and Wilhelm Barthlott

Number 12

- 1073 Triterpenes from *Maytenus macrocarpa* and Evaluation of Their Anti-HIV activity
Sonia Piacente, Lourdes Campaner Dos Santos, Naheed Mahmood and Cosimo Pizza
- 1079 New Oxidized 4-Oxo Fatty Acids from *Hygrophorus discoxanthus*
Gianluca Gilardoni, Marco Clericuzio, Alberto Marchetti, Paola Vita Finzi, Giuseppe Zanoni and Giovanni Vidari
- 1085 Kenyaloside, a Novel *O,O,O*-Triglycosylated Naphthalene Derivative from the Exudate of Kenyan *aloe* Species
Giovanna Speranza, Daniela Monti, Sergio Crippa, Paola Cairoli, Carlo F. Morelli and Paolo Manitto
- 1089 New Flavonoid Glycosides from *Chrozophora senegalensis* and Their Antioxidant Activity
Antonio Vassallo, Giuseppina Cioffi, Francesco De Simone, Alessandra Braca, Rokia Sanogo, Angelo Vanella, Alessandra Russo and Nunziatina De Tommasi
- 1097 *N1,N2,N3*-Trisisopentenyl Guanidine and *N1,N2*-Diisopentenyl guanidine, Two Cytotoxic Alkaloids from *Alchornea cordifolia* (Schumacher & Thonn.) Müll. Arg. (Euphorbiaceae) Root Barks
Hélène Mavar-Manga, David Chapon, Sara Hoet, Sébastien Block, Marie-Claire. De Pauw-Gillet and Joëlle Quetin-Leclercq
- 1101 Indole Monoterpenes with Antichemotactic Activity from *Psychotria myriantha*
Cláudia A. Simões-Pires, Fabianne M. Farias, Andrew Marston, Emerson F. Queiroz, Célia G. Chaves, Amélia T. Henriques and Kurt Hostettmann
- 1107 HPLC Based Activity Profiling for Inhibitors of Human Neutrophil Elastase in *Isatis tinctoria* Leaf Extracts
M. Hamburger, H. G. Rüster and M. F. Melzig
- 1111 Variation in Artemisinin and Flavonoids Content in Different Extracts of *Artemisia annua* L.
Anna Rita Bilia, Caterina Gabriele, Maria Camilla Bergonzi, Pedro Melillo de Malgalhaes and Franco Francesco Vincieri
- 1117 Antifungal Evaluation of *Hypericum triquetrifolium* Polar Extracts Against *Fusarium* spp
Daniele Fraternale, Alessandra Bertoli, Laura Giamperi, Anahi Bucchini, Donata Ricci, Francesco Menichini, Elena Trinciarelli and Luisa Pistelli
- 1123 Antioxidant Activity Analysis for the Selection of *Rosmarinus officinalis* L.
Juan Antonio Garbarino, Nicolás Troncoso, Pia Delpiano, Loreto Carvajal and Alessandra Russo
- 1129 *Hypericum perforatum* L., *H. maculatum* Crantz., *H. calycinum* L. and *H. pulchrum* L.: Phytochemical and Morphological Studies
Gelsomina Fico, Sara Vitalini, Noemi Colombo and Franca Tomè
- 1133 Chemical Composition and Antimicrobial Activities of Essential Oil of *Stachys glutinosa* L. from Sardinia
Pintore Giorgio, Chessa Mario, Manconi Paola, Zanetti Stefania, Deriu Antonella and Tirillini Bruno
- 1137 Molecular Identification of *Panax ginseng* C.A. Meyer in Ginseng Commercial Products
Paola Del Serrone, Lucilla Attorri, Bruno Gallinella, Francesca Romana Gallo, Elena Federici and Giovanna Plazzino
- 1141 Lipoygenase Inhibitory Activity of Boropinic Acid, Active Principle of *Boronia pinnata*
Massimo Curini, Francesco Epifano, Salvatore Genovese, Luigi Menghini, Donata Ricci, Daniele Fraternale, Laura Giamperi, Anahi Bucchini and Emanuele Bellacchio
- 1147 A Convenient Synthesis of 5'-Iodoresiniferatoxin (I-RTX)
Abdellah Ech-Chahad, Lahboub Bouyazza and Giovanni Appendino
- 1151 Acaricides of Natural Origin. Part 2. Review of the Literature (2002-2006)
Guido Flamini
- 1159 Chemistry and Biological Activity of Saponins from *Medicago* Species
Aldo Tava and Pinarosa Avato
- 1181 Non-nitrogenous Plant-derived Constituents with Antimalarial Activity
Anna Rita Bilia

Natural Product Communications

Author Index of Volume 1 2006

- Abdella, AM 523
Achmad, SA 549
Adoga, GI 961
Agrawal, C 957
Agrawal, PK 957
Agrawal, S 957
Ahmed, AA 273,523,697,941,961
Ahmed, FA 457
Aimi, N 549
Akam, TM 449
Akkal, S 645
Alali, FQ 95
Alamgir, M 985
Albuquerque, MRJR 395
Ali, BH 509
Alkofahi, A 95
Al-Saidi, S 195
Al-Shihry, SS 697
Alvarenga, SAV 495
Amado, FML 973
Andrade-Neto, M 313,391,469
Antonella, D 1133
Antus, S 51,991
Aoshima, T 431
Appendino, G 1147
Araque, M 113
Arasi, HN 387
Armstrong, N 121
Arnold, NA 281
Arriaga, AMC 391,573
Attard, E 381
Attard, H 381
Attorri, L 1137
Avato, P 751,1159
Ayoub, S 633
- Bacchi, C 619
Bailleul, F 727
Bakheit, AO 509
Baldovini, N 387,645
Bandeira, PN 117
Banerjee, S 665,731
Bankova, V 1023
Bansal, A 303
Barbosa, FG 313,469,573
Barbosa, JP 293
Barbosa, KO 495
Barra, A 387
Barros-Filho, BA 313
Barthlott, W 1067
Bellacchio, E 1141
Berdyshev, DV 265
Bergonzi, MC 1111
- Bertoli, A 557,1117
Bezerra, AME 469
Bezerra, FS 469
Bhakuni, RS 721
Biagi, M 209
Bianchini, JP 969
Bibby, MC 47
Bilia, AR 1111,1181
Block, S 1097
Blunden, G 47,121,509,613,705
Boehme, AK 303
Bohm, BA 1013
Bojase-Moleta, G 287
Booth, R 441
Borrelli, F 77
Bouyazza, L 1147
Braca, A 1089
Brant, AJC 495
Brem, B 921
Bremner, JB 825
Brighente, IMC 37
Bruno, M 281,585
Bruno, T 1133
Bucchini, A 1117,1141
Burgess, J 95
Butaud, JF 969
- Cabo, M 27
Cabrol-Bass, D 495
Cai, L 191
Cairolì, P 1085
Câmara, MT 973
Campos, MG 973
Canard, B 645
Capasso, F 77
Capasso, R 77
Carda, M 237
Carvajal, L 1123
Cavalcanti, DN 609
Chan-Bacab, M 541
Chapagain, BP 215
Chapleur, Y 475
Chapon, D 1097
Charrouf, Z 399
Chattopadhyay, S 131,767
Chaves, CG 1101
Chen, CH 20,461
Chen, C-H 299
Chen, C-K 299
Chen, CK 461
Chen, CY 20,453
Chen, D-L 191,357
Chen, Q 527
- Chen, SC 461
Choudhary, MI 363
Chrétien, F 475
Christophersen, C 541
Cioffi, G 1089
Claire, M 1097
Clericuzio, M 1079
Clevinger, J 941
Coll, J 183
Colombo, N 1129
Connolly, JD 449
Conrad, J 935
Conserva, LM 319
Costa, SMO 117
Costa-Lotufu, LV 313
Couladis, M 523
Cox, PJ 641
Crippa, S 1085
Cunha, MM 973
Curini, M 1141
- Dacke, CG 613,705
Darokar, MP 223,481
Das, B 255
Das, R 255
Davis, BD 941
de Malgalhaes, PM 1111
de Mattos, MC 313,661
de Menezes, GHA 573
de Moraes, MO 313
de Oliveira, MCF 313,391,469,661
de Sousa, AL 661
de Souza, AO 313
de Souza, DF 661
Debnath, C 487
Delpiano, P 1123
Dembitsky, VM 139,405,773
Demetzos, C 593
Demin, MA 997
Denisenko, VA 265
Dimayuga, ER 541
Diop, MF 475
Djemgou, PC 961
Dmitrenok, AS 711
Dmitrenok, PS 265, 711
Dos Santos, LC 1073
Dubery, IA 367
Duong, GM 985
Duthie, G 1049
- Ebel, R 927
Ech-Chahad, A 1147

Author Index
Natural Product Communications Vol. **1** (1-12) 2006

- Edrada, RA 927
 Eiter, LC 303
 El Hamchi, H 399
 El-Bassuony, AA 273
 El-Sayed, MM 735
 El-Sayed, NH 941
 El-Seedi, HR 655
 Emerenciano, VP 57,495
 Epifano, F 1141
 Essien, E 763
 Esterhuizen, LL 367
 Evans, C 449
- Falomir, E 237
 Fanizzi, FP 751
 Fanzo-Free, SNY 449
 Farias, FM 1101
 Farrugia, LJ 449
 Faure, R 969
 Federici, E 1137
 Fernandes, GW 627
 Ferreira, MJP 495
 Fico, G 1129
 Filho, JMB 319
 Filho, RB 37
 Finzi, PV 1079
 Flamini, G 1151
 Fonseca, AM 117
 Forster, PI 351,441
 Fotso, S 9
 Fraternal, D 1117,1141
 Fromageot, C 27
 Fukushi, Y 981
 Funk, JL 1061
- Gabriele, C 1111
 Gad, MH 273
 Galetti, FCS 313
 Gallinella, B 1137
 Gallo, FR 1137
 Gamal-Eldeen, AM 735,961
 Gan, LS 819
 Ganapaty, S 81
 Garbarino, JA 1123
 García-Viguera, C 1037
 Gatsing, D 961
 Gaydou, EM 969
 Geiss, HK 1003
 Genovese, S 1141
 Gherraf, N 273
 Ghisalberti, EL 549,859
 Giachetti, D 209
 Giamperi, L 1117,1141
 Gilardoni, G 1079
 Giorgio, P 1133
 Glorizova, TA 773
 Gonçalves, FJT 469
 Gonçalves, R 757
 Göpfert, J 935
 Gourlay, BS 831
 Gualtieri, M 113
 Guan, S 109
 Guan, S-H 177
 Guillaume, D 399
- Gulácsi, K 991
 Guo, D 109,177
 Guzii, AG 711
- Haber, WA 465
 Hadi, S 825
 Hakim, EH 549
 Halstead, CW 351
 Hamburger, M 1107
 Haraguchi, M 431
 Hasegawa, K 87
 Haslinger, E 487
 Hassan, HE 523
 He, Y 357
 Hegazy, MF 523
 Hennebelle, T 727
 Henriques, AT 1101
 Henry, M 475
 Hirata, T 273,961
 Hisham, A 195
 Hoet, S 1097
 Hostettmann, K 27,1101
 Houghton, PJ 293
 Hsieh, TJ 20
 Hutton, CA 907
- Ibrahim, RK 981
 Iinuma, M 949
 Issakul, K 813
 Ivancheva, S 1029
 Iwu, MM 619
- Jaspars, M 33,641
 Jatisatienr, A 813
 Ji, H-F 229
 Jin, H 177
 Josaphine, JS 81
 Joseph, H 727
 Juma, BF 101
 Justin, KJ 363
- Kadota, S 1
 Kalena,GP 131
 Kaplan, MAC 293
 Kassem, MES 953
 Kaul, VK 623
 Kéki, S 991
 Kenéz, Á 51
 Khalil, AT 531
 Khanuja, SPS 223,481
 Koch, K 1067
 Kondo, R 345
 Kong-Thu-lin, P 33
 Krasokhin, VB 711
 Krishnaiah, M 255
 Kröpfel, D 307,651
 Kumar, N 623
 Kumarasamy, Y 33,641
 Kuo, C-M 299
 Kuo, CM 461
 Kuo, YH 531
 Kurdi, VA 549
 Kuroda, M 431
 Kyle, J 1049
- Laatsch, H 9,81
 Lannang, AM 363
 Laouer, H 645
 Laurain-Mattar, D 475
 Lee, SS 299,461
 Lemos, TLG 391,573,661
 Levitsky, DO 405, 773
 Li, C 95
 Lie, W 813
 Licitra, G 399
 Lima, MAS 573
 Lima, MCL 573
 Lin, W 927
 Linley, PA 47
 Lins, MUDS 117
 Little, I 831
 Liu, J 345,839
 Liu, P 109
 Liu, R 109
 Liu, X 177
 Liu, Y 839
 Lo, KL 531
 Lo, WL 20
 Lontsi, D 363
 Loquet, D 969
 Louh, GN 363
 Lubrano, C 27
 Luciano, JHS 573
 Luqman, S 481
- Ma, X 109
 Ma'aya'h, AS 95
 Mabry, TJ 941,961
 Machumi, F 287
 Mafezoli, J 313,391,469
 Maggio, A 585
 Maggio, AM 281
 Mahmood, N 1073
 Mahmoud, AA. 15
 Mahmoud, IM 745
 Mahmoud, MR 745
 Majinda, RRT 101,287
 Makarieva, TN 711
 Makmur, L 549
 Mallia, S 399
 Malterud, KE 985
 Mandal, D 731
 Mandal, NB 665,731
 Manitto, P 1085
 Mapitse, R 287
 Marchetti, A 1079
 Marco, JA 237
 Mario, C 1133
 Márquez, A 205
 Marriif, H 509
 Marsaioli, AJ 757
 Marston, A 27,1101
 Marzouk, MSA 735
 Masesane, I 287
 Máthé, I 113,205
 Matlawska, I 941
 Matos, FJA 573
 Matos, MP 973
 Mavar-Manga, H 1097

Author Index
Natural Product Communications Vol. **1** (1-12) 2006

- Mayekar, NV 767
 McCamley, K 885
 Meierhenrich, UJ 645
 Melo, VMM 395
 Melzig, MF 633,1107
 Menghini, L 1141
 Menichini, F 1117
 Menon, RS 899
 Meyer, R 367
 Milovanovic, MM 997
 Mimaki, Y 247,431
 Miraldi, E 209
 Mohamed, AEH 523
 Mohamed, MA 735
 Moharram, FA 745
 Monte, FJQ 117
 Montenegro, RC 313
 Monti, D 1085
 Morales, A 113,205
 Morelli, CF 1085
 Moreno, DA 1037
 Mosaddik A 441,601
 Mula, S 131
 Murga, J 237
 Murillo-Álvarez, JI 541
- Nahar, L 33,377,641
 Naiker, M 715
 Nair, V 899
 Nascimento, RF 391
 Nascimento, RRG 469
 Nayak, SK 767
 Neiva, TJC 37
 Németh, I 991
 Nesic, VM 997
 Neto, CP 973
 Ngadjui, BT 961
 Ngamga, D 619
 Ngounou, FN 363
 Nikolova, M 1029
 Noletto, JA 43
 Noureldayem, SA 509
 Nunes, EP 395
 Nunes, FM 313
 Nuzillard, JM 57
- Oberlies, NH 95
 Obli Prabu, KV 481
 Ogunwande, IA 651,763
 Ohta, S 273
 Okunji, CC 619
 Ortner, A 487
 Oyama, M 949
 Öztürk, M 851
- Padmakumar, K 927
 Pal, A 481
 Palazzino, G 1137
 Palmeira Júnior, SF 319
 Pan, Y 537
 Panda, N 665,731
 Paola, M 1133
 Papadia, P 751
 Paranhos, AH 973
- Paré, P 523
 Pasquale, S 205
 Patel, AV 121,613,705
 Pathare, N 195
 Patro, BS 131
 Pauw-Gillet, MCD 1097
 Pecetti, L 557
 Pelah, D 215
 Pereira, RC 293
 Pérez-Balibrea, S 103
 Pessoa, C 313
 Pessoa, ODL 117,395,573
 Piacente, S 1073
 Piano, E 557
 Pinto, AC 609
 Pinto, PRO 973
 Piozzi, F 585
 Pires, AML 395
 Pistelli, L 1117
 Pizza, C 1073
 Pizzolatti, MG 37
 Popova, M 1023
 Poroikov, VV 773
 Prasad, SY 715
 Prettner, E 307
 Proksch, P 927
 Ptak, A 475
 Pyne, SG 813
- Qandil, A 95
 Qian, Z 63
 Qin, S 9
 Qu, J 839
 Queiroz, EF 1101
 Quetin-Leclercq, J 1097
- Raccuglia, RA 281
 Raharivelomanana, P 969
 Rajakaruna, N 1013
 Rattendi, D 619
 Reichling, J 1003
 Reiriz, MLG 541
 Reis, MG 757
 Rezende, CM 609
 Rhouati, S 273
 Ribeiro, WHF 391
 Ricci, D 1117,1141
 Robin, JR 27
 Rocha, FD 293
 Rojas, J 113,205
 Romero, NR 469
 Romero, AM 121
 Rondón, M 113,205
 Rosito, I 751
 Rosselli, S 281,585
 Russo, A 1089,1123
 Rüster, GU 1107
 Ryan, JH 831
- Sabry, SA 745
 Saharan, V 215
 Sahpaz, S 727
 Sahu, NP 665,731
- Saikia, D 481
 Sakagami, H 431
 Saleh, NAM 953
 Samuelsen, AB 985
 Sanogo, R 1089
 Santiago, GMP 391,573
 Santos, HS 573
 Sarder, M 985
 Sarker, SD 33,377,641
 Sastraruji, T 813
 Saxena, DB 553
 Scammells, PJ 885
 Schmidt, JM 201,303
 Schneelee, J 1003
 Schripsema, J 37
 Schuster, BM 619
 Schweiger, K 307
 Scotti, MT 495
 Sena, JSP 469
 Serrone, PD 1137
 Setzer, WN 43,201,303,465,567,763
 Shabana, MH 953
 Shahat, AA 457
 Shakil, NA 553
 Sharaf, M 953
 Sharehjini, SS 387
 Shasany, AK 223
 Shen, L 229
 Shen, YC 531
 Shi, Z 527
 Shigemori, H 87
 Shimizu, K 345
 Shirataki, Y 563
 Shoeib, NA 47
 Silva, CL 313
 Silveira, ER 395,573
 Silvestre, AJD 973
 Simic, A 651
 Simic, N 651
 Simmonds, MSJ 281
 Simões-Pires, C 1101
 Simone, FD 1089
 Singer, RB 757
 Singh, A 577
 Singh, B 623
 Singh, JA 715
 Singh, RD 715
 Smith, JA 831
 Soladoye, MO 763
 Sondengam, BL 363
 Sonibare, MA 763
 Speranza, G 1085
 Spring, O 935
 Stefania, Z 1133
 Sterner, O 619
 Stonik, VA 265,711
 Subehan I
 Suenaga, K 87
 Sun, W 527
 Suschke, U 1003
 Swaine, DJ 47
 Syah, YM 549
- Tanaka, T 949

Author Index

Natural Product Communications Vol. **1** (1-12) 2006

- Tandrón, YA 183
Tane, P 449,619,961
Tang, W 839
Tangmouo, JG 363
Tava, A 557,1159
Tchuendem, M 961
Teixeira, VL 293,609
Teuscher, F 927
Tewari, A 21
Tezuka, Y 1
Than, NN 81
Thavaneswaran, S 885
Thomas, PS 81
Thul, ST 223
Timmermann, BN 1061
Tiwari, S 577
Toda, S 563
Tomè, F 1129
Tommasi, ND 1089
Tóth, G 941
Trevisan, MTS 661
Trinciarelli, E 1117
Troncoso, N 1123
Trusheva, B 1023
Tu, G 527
Tuh, S-L 299
Tzakou, O 523

Ulubelen, A 851
Ung, AT 813

Usia, T 1

Valant-Vetschera, KM 627,921
Vanella, A 1089
Vasconcelos, JN 391
Vassallo, A 1089
Vellalath, S 899
Venkateswarlu, K 255
Vera, JR 47,613,705
Verdi, LG 37
Veres, K 113,205
Vidari, G 1079
Vincent, MA 43,303
Vincieri, FF 1111
Vitalini, S 1129
Vogler, B 465
Voro, TN 715
Vucelic-Radovic, BV 997

Wabo, HK 449
Wagner, SF 307
Walker, TM 763
Wang, F-P 191,357
Wang, K 537
Wang, W 109
Wang, XF 981
Wang, X-M 177
Wangenstein, H 985
Wani, MC 95
Waterman, PG 351,441,601

Wenceslau, JPS 661
Wiesman, Z 215
Williams, J 941
Williams, MC 813
Wojcińska, M 941
Wollenweber, E 627,949
Wray, V 927
Wright, CW 47
Wu, SJ 9
Wu, X 465

Xi, L 63
Xia, J-M 177

Yadav, RC 215
Yamazoe, S 87
Yang, M 177
Yong, JN 449
Young, MCM 431
Yu, S 839
Yue, JM 819
Yuen, AKL 907

Zacchino, S 541
Zanoni, G 1079
Zhang, H-Y 229
Zhang, Y 527
Zhou, JM 981
Zsuga, M 991

Natural Product Communications

2006

Key Word Index of Volume 1

- AbyssinoneVII 287
Acaricides 1151
Acanthophora spicifera 927
Acetogenins 773
Acetylenic compounds 405,773
Acetylenic metabolites 405,773
Achillea biebersteinii 697
Aconitum piepunense 191
Activity profiling 1107
Acylated dihydrofuranone 601
Acylated glycosides 461
Acylglycerols 757
Aedes aegypti 391,573
AFLP 223
Ajuga nipponensis 183
Alchornea cordifolia 1097
Alcohols 773
Algae 139,773
Algicolous fungi 927
Alkaloids 95,191,313,357,619,735,773,813,819,825,831,839,851,859,885,899,907,1097,1101
Alkamides 1
Alkylindolizidines 831
Alnus formosana 299,461
Alnus glutinosa 641
Aloe ferox 1085
Aloes 1085
Amaranthaceae 431
Amaryllidaceae 475
Amberlyst 131
Amentoflavone 633
Amides 527,839
Amines 839,885
Angiosperms 1029
Angiotensin-converting enzyme (ACE) activity 633
Angiotensin-converting enzyme inhibitory activity 381
Annona 121
Annonaceae 377
Anthocyanins 229
Antibacterial activity 395,645,825,1003
Antibacterial and antifungal activities 645
Tetrahydrofuran derivatives 237
Antibacterial 113,641
Antibiotics 481
Anticancer 405,509,773
Antichemotactic activity 1101
Anticoagulant 37
Antifeedants activity 183,281
Antifungal activity 51,645,1117
Anti-HIV activity 1073
Anti-inflammatory activity 509,1061,1107,1141
Antileishmanial 541
Antimicrobial activity 81,101,117,195,303,651,655,985,1133
Antioxidant activity 117,131,367,509,641,661,973,985,1089,1123
Anti-proliferative 735
Antitrypanosomal alkaloids 619
Trypanosomiasis 619
Antitumor activity 247,313,961
Antiviral activity 609,645
Apigenin 633
Apples 307
Antiplasmodial activity 449,1181
Araliaceae 87
Argan oil 399
Argania spinosa 399
Artemisia annua L. 1111
Artemisia annua 487
Artemisinin 487, 1111
Arthritis treatment 1061
Argolic acid A 523
Argolic methyl ester B 523
Artocarpus altilis 549
Artocarpus champeden 549
Artoindonesianin A-1 549
Artoindonesianin B-1 549
2-Arylbenzofuran 549
Aryl migration 991
Asclepiadaceae 731
Aspergillus sydowii 927
Asteraceae 15,37,113,281,395,495,655,697,941
Astereae-Asteraceae 627
Aurones 957
Azocine 9
B group vitamins 997
Baccharis confertifolia 627
Baccharis illinita 37
Baccharis 627
Baccharis lycioides 627
Baccharis linearis 627
Bacteria 139,405
Bactericidal activity 825
Balanites aegyptiaca 215
Bark essential oil 567
Basidiomycetes 1079
Bavachin 51
Benign prostatic hyperplasia 345
Berberidaceae 121
 α -Bergamotene 651
Betulaceae 299,461,641
Benzopyranones 351
2'-Benzylxyflavanones 991
Bicyclic sesquiterpene 15
Biebersteiniside 697
Bignoniaceae 735
Bioactive flavonoids 953
Bioactivity 839,1037
Biochemical studies 255
Biogenesis 585
Biological activities 319, 585,665,1159
Bioproduction 215
Biosynthesis of steroidal glycoalkaloids 859
bis-Benzylisoquinoline alkaloid 619
Blainvillea rhomboidea 395
Bond dissociation enthalpy 229
Bone 1061
Bonvalotidine A 357
Bonvalotidine B 357
Bonvalotidine C 357
Borneol 567
Boropinic acid 1141

Key Word Index

Natural Product Communications Vol. 1 (1-12) 2006

- Boronia pinnata* 1141
 Bran layers 997

Brassica 1037
 Breast carcinoma 735
 Brine shrimp lethality 101,641
Brombya platynema 351
Brombya sp. 351
 Brombyins 351
 Bromophenols 47
 Bulgarian plants 1029
 Butanolides 453
 Butin apiofuranosylglucopyranoside 623
 15 β -Butoxy-14,15-dihydronorsecurinine 819
 15 α -Butoxy-14,15-dihydronorsecurinine 819
 Burseraceae 117

 C₁₈-diterpenoid alkaloids 191
 C₁₉-diterpenoid alkaloid 357
 α -Cadinol 201
 Callus 215
Calyptanthus pallens 303
 Camptothecins 255
 Cancer prevention 1049
 Capitate glandular trichome 935
Capsicum germplasm 223
¹³C NMR 319,387,585,957
 Carnosic acid 1123
 β -carotene 481
 Carotenoids 773
 Carvacrol 205
Casearia grayi 441
Casearia multinervosa 601
 Cassaine-type diterpenoid ester amines 839
 Cassaine-type diterpenoid ester amides 839
Cassia petersiana 961
 Catechins 985
 α -Cedrene 1133
 Celastraceae 537,1073
Celastrus rosthornianus 537
 Celogentin 907
Centaurea hololeuca 281
 Centrifugal partition chromatography 27
 C-glycosides 457,731
 C-glycosylflavones 969
 Chalcone glycoside 623
 Chalcone 81,553,949
Channa punctatus 577
 Chemical databases 57
 Chemical evolution 495
 Chemical modifications 255
 Chemical systematics 351
 Chemodiversity 921
 Chemosystematics 281
 Chemotaxonomy 121,299,549,609
 Chiral analysis 645
 Cholestane glycosides 247
 Chromones 961
Chrozophora senegalensis 1089
 Chalcone synthase (CHS) 935
 Cinchona alkaloids 899
 Cineole 567
 1,8-Cineole 763
Cinnamomum camphora 21
Cinnamomum kotoense 453
 2'''-Cinnamoylorigonin 461
 Clerodane diterpenes 183,319,441,601
 C-methylated flavonoids 957
 Colchicaceae 95
 Colchicine 95
 Colchicinoids 95
Colchicum tauri 95

Coleonema album 367
 Colon cancer 33
 Communiols 237
 Complestatin 907
 Corals 139
 Costic acid 465
 Coumarins 351,367,495,851
 C-prenylated flavanones 51
 C-prenylflavonoids 81
Crataegus monogyna 381
 Crocetin 65
 Crocin 65
 Chloroethylphosphonic acid 307
Crocus sativus 65
Croton 319
Croton monteverdensis 567
Croton niveus 567
Curcuma longa 509
 Curcumin 509
 Cyanidin 229
 Cyanobacteria 139,773
 Cyanolipids 751
 Cyclic peptide alkaloids 907
 Cyclopentanoids 927
 Cyclopentanomonoterpene 523
 p-Cymene 205, 763
 CYP2D6 1
 Cytochrome P450 2D6 1
 Cytotoxic activity 21,47,177,247,405,431,531,541,773,961,1097
 Cytotoxic alkaloids 1097
 Cytotoxicity 303,1097

 2D NMR 57
Delphinium bonvalotii 357
 Dendrimer 593
 Density functional theory 229
Derris heyneana 81
 7,8-Dehydromoskachen D 351
 5-Deoxynevadensin 935
 Diarylheptanoids 461
 (+)-Diasamin 21
 Diazonamide 907
 Dibenzylbutyrolactone lignan 109
Dictyopteris justii 293
Dictyota menstrualis 609
 Dictyotaceae 609
 Dietary intakes 1049
 Differential pulse polarography 487
 2,3-Dihydrobenzo[b]furans 991
 Dihydroflavonols, C-methylated 957
 Diisopentenyl guanidine (DIPG) 1097
 Diisopentenyl-N-methyl-quinoldione 313
 Diosgenin 215
 Diterpenoid alkaloids 191,357
 Dipsacaceae 457
 Disease prevention 1049
 Diterpenes 37,183,273,319,441,601,609
 Diterpenoids 449,585, 609,697
 DPPH 101,641
Dregea volubilis 731
 Dregeanin 731
 Drug carriers 593
 Brown Alga 609
 Drug-herb interaction 1

 Echinosoflavanone 563
 Ecological aspects 627
 Edaphic races 1013
 Embryogenesis 215,475
 Enamines 661
 Endophytic fungus 927
 Enzyme induction 633

Key Word Index

Natural Product Communications Vol. 1 (1-12) 2006

- Epicuticular waxes 1067
 Epideoxyloganic acid 523
Erythrina abyssinica 287
Erythrina lysistemon 101
 Erythrocentaurin acid 527
Erythrophleum alkaloids 839
 Erythrocytes 481
Esenbeckia almarillia 313
 Essential oil 43,113,117,195,201,205,303,391,395,465,469,567,573,645,651,655,763,1003,1133
 Ethephon 307
 Ethephon residues in apples 307
Eugenia confusa 43
 Eupakirunsin F 531
 Eupakirunsin G 531
 Eupakirunsin H 531
 Eupakirunsin I 531
Eupatorium glutinosum 655
Eupatorium kiirunense 531
Euphorbia guyoniana 273
Euphorbia royleana 577
 Euphorbiaceae 273,319,567,1089
 Expert systems 57
 Explants 215
 Exudate flavonoids 627,921,949
 Exudate 949
 Fabaceae 953

 Fatty acids 773,1079
 Fabaceae 81
 Fermentation 9
Ficus exasperate 763
 Fish 773
 Flacourtiaceae 441,601
 Flavanone glycosides 623
 Flavanone 51,81,287,957
 Flavone 229,731,935
 Flavone C-glycosides 457,731,745
 Flavone glycosides 727,745,1089
 Flavonoid C-glycosides 457
 Flavonoids 37,81,101,287,367,461,495,563,627,641,727,745,921,935,949,953,957,969,1029,1049,1089,1111,1117,1129
 Flavonol glycosides 457,941,953,1089
 Floral oils 757
Flueggea virosa 819
 Fragranol 387
 Fragranyl acetate 387
 Free radicals 367,1123
 Friedelane triterpenoids 1073
 Frog toxins 831
 Fungi 405
 Fungicidal 1079
 Furanone 601
 Furoquinolines 351,767
Fusarium ssp. 1117

 Galanthamine 475
 Gall 87
Galtonia candicans 247
 Ganoderic acid TR 345
Ganoderma lucidum 177,345
Garcinia polyantha 363
 Gas chromatography-mass spectra 655
 GC 557,751
 GC/MS analyses 307,399,557,763
 GC-O analysis 399
 Genetic diversity 223
 Genkwanin 641
Gentiana macrophylla 527
 Gentianaceae 527
Geranium tuberosum 387
 Germacrene D 113,201

 Ginseng commercial products 1137
 GLC-MS 711
Glechoma hederaceae 33
 Glucoside indole monoterpene alkaloids 1101
 Glucosinolates 1037
 Glycerols 773
 Glycosides 27,247,265,431,457,461,541,623,665,697,721,727,731,735,745,859,941,953,969,1085,1089,1101
 Glycoalkaloids 859
Gomphrena macrocephala 431
 Guaianolides 281
 Guanidine alkaloids 1097
 Guttiferae 363
 Guyonianin A 273
 Guyonianin B 273

 Halimane diterpenes 441
 Heart disease prevention 1049
Hedera rhombea 87
 Hederacine A 33
 Hederacine B 33
 Helianthus annuus 935
 Hepatoprotective 745
 Herbal remedies 77
 HET-CAM assay 1003
 Hirsutanolol 641
 HPLC 183,475
 HPLC/DAD isoflavones 973
 HPLC/DAD/MS 1111
 HSC-2 cells 431
 Human and veterinary medicines 1151
 α -Humulene 113
 Inhibitors of human neutrophil elastase 1107
 Human pulp cells 431
 Humulene sesquiterpenes 601
 Hydrocarbons 757
 Hydroethanolic extract 381
 Hydroperoxysterol 293
 Hydroperoxyvinyl cholesterol 293
 2'-Hydroxy-2-methoxychalcone 949
 1-Hydroxy-2-tridecanone. 469
Hygrophorus discosanthus 1079
Hypericum perforatum 209
Hypericum triquetrifolium 1117
Hypericum 1129
 Hypoglycemic 745

 Ilicic acid 465
 Imino Diels-Alder reaction 767
 Immunoproliferative agents 961
 Immunostimulatory activity 961
 Individual and infraspecific differentiation 627
 Indolizidines 831
 Inflammation 1061
 Infraspecific variability 1029
 Insect growth regulatory activity 553
 Insecticidal 553
in vitro Culture 475
 Iodine 139
 Iodine-containing metabolite 139
 Iodoperoxidase 139
 5'-Iodoresiniferatoxin (I-RTX) 1147
 Ionization potential 229
 Ionone glucoside 697
 Iridoids 527,727
Isatis tinctoria 1107
 3-Isocostic acid 465
 Isoflavanone 563
 Isoflavones 81,557,973
 Isoflavone quantification 557
 Isoquinoline alkaloids 619
 Isorhamnetin 3,7-O-trisides 941

Key Word Index

Natural Product Communications Vol. 1 (1-12) 2006

- Jasomontanone 15
Jasonia Montana 15
 Jordan medicinal plants 95

Kadsura heteroclita 109
 Kauranes 37
 Kava lactones 715
 Kava lactone-yielding precursors 715
Kava 715
 Kenyaloside 1085
 Kernel roasting 399
 Kernel 997
 Kotomolide 453

 Labdanes 449,585
 Labiatae 183
 Lactones 453,715
 Lamiaceae 33,195,523,585,745,1113
 Lanostane triterpenes 177,345
 Lapachol derivatives 661
 Larvicidal activity 391,573
 L-ascorbic acid 481
Lasiocephalus longipenicillatus 113
Lasthenia californica 1013
 Lauraceae 21,453
Laurus 121
 Leaf extract of *Rosmarinus officinalis* 1123
 Legumes 1037
 Leguminosae 287,961
Leucosium aestivum 475
 Lignan 21,109
 Liliaceae 859
 Limonoids 985
 Linalool 567
 Linoleic acid 1107
 Lipid peroxidation 563
 Lipids 773
 15-Lipoxygenase 985
 Lipoxygenase inhibitory activity 1141
Lippia 573
Lippia alba 573,727
Lippia gracillis 573
Lippia microphylla 573
Lippia nodiflora 573
Lippia Oreganoides 205
 Lombine 825
 Lombok 825
 Lonchocarpol A 51
 LSD 57
 Luteolin 633

Madicago sativa 1159
 Macrophage proliferation 961
 Malaria 1181
 MALDI-TOF MS 265
 Marine saponins 541
 Marine Streptomycetes 9
 Marine-derived fungi 927
Marrubium 585
 Mass spectroscopy 475, 665
Maytenus macrocarpa 1073
Medicago 1159
 Medicinal properties 65
 Meliaceae 121,449
Meloidogyne incognita 469
Melophlus sarasinorum 265
 Menispermaceae 553
 9-Methyl-8-oxoadenine 711
 Minerals 997,1037
 Molecular identification technique 1137
 Mollusks 139

 Monimiaceae 619
 Monomethylated adenines 711
 Intracellular cyclic-adenosine monophosphate (cAMP) 633
 Monoterpene alkaloid 735
 Monoterpene diol 523
 Monotesone B 51
 Moraceae 549,763
Moringa oleifera 721
 Moringaceae 721
 Thiocarbamates 721
 MS of steroidal glycoalkaloids 859
 Myrianthosine 1101
 Myristicaceae 651
 Myrtaceae 43,303

 Nanotechnology 593
 Naphthalene *O,O,O*-triglycoside 1085
 Naphthodianthrone 1129
 Bioactive natural products 593
 N-Demethylation 885
Nectandra membranacea 465
 Nematicidal activity 469
Neo-clerodane diterpenes 183
 Neolignene 101
Neothyone gibbosa 541
 Neothyoside A 541
 Neothyoside B 541
 Neothyoside C 541
 Neothyosides 541
Nepeta argolica 523
 Nepetonic acid 523
Nephelium lappaceum L.751
 Neutral endopeptidase (NEP) 633
 N-methylprolines 121
 NMR 183,265,457,537,665,751,859,957
 norlanostanes 265
 norsecurinic acid 819
 norsecurinine derivatives 819
Nothapodytes foetida 255
 Nutritional relevance of flavonoids 1049

 Oleanane triterpenoids 431,613,705,1073
 Olean-12-ene-3 β ,29-diol 613
 Olean-18-ene-1 β ,2 α ,3 α -triol 613
 Oleanane glycosides 27,431
 Oleanolic acid 381
Oleum Hyperici 209
O-Glycosylflavones 969
O-Methyltransferase 981
O-Methyltransferase gene 981
 Ontogenetic cycle 1129
 Opiates 885
 Orchidaceae 757
 Oregonin 641
 Organogenesis 475
Ornithogalum saundersiae 247
Ornithogalum thyrsoides 247
Oryza sativa L 981
 Osmotic fragility 481
 OSW-1 247
 Oxidation number 495
 Oxidative steps 495
 8-Oxoadenine 711
 4-Oxo-fatty acids 1079

Panax quinquefolium 1137
Pachypodanthium staudtii 377
Panax ginseng 1137
 Partial least squares regression 495
 PGR 215
 Pharmacokinetics 65
 Pharmacological activity 851

Key Word Index

Natural Product Communications Vol. 1 (1-12) 2006

- Phenolics 101,131
 Prenylated arylbenzofurans 549
 Prenylated phenols 131
 Phaeophyta 609
 Phenylethanoid 727,735
 Phenylpropanoids 1003
 Phloroglucinols 1129
Phoradendron 121
Phymatidium 757
Phymatidium delicatulum 757
Phymatidium tillandsioides 757
 Phytoestrogens 973
 (*E*)-phytol 763
 Piepunendine A 191
 Piepunendine B 191
 α -Pinene 113, 201,303,567
 β -Pinene 567
Pilocarpus microphyllus 469
Piper nigrum 1
 Pipericyclobutanamide 1
 Piperonyl derivatives 351
 Plant growth inhibitors 87
 Plant sources of propolis 1023
 Plant-derived non-nitrogenous antiplasmodials 1181
 Polarographic method 487
 Polyacetylenes 87, 405,495,773
 Polyanxanthone 363
 Polyethers 773
 Polyester diterpenes 273
 Polyphenols 1181
Polysiphonia lanosa 47
 Post-menopause treatment 973
 Pregnane glycosides 665
 Pregnanes 449,665
 Prenylation 131
 Prenyloxy acids 1141
Primula palinuri 949
 Primulaceae 949
 Principle component analysis 223
 Proanthocyanidins 985
 Procyanidins 985
 Propolis 1023
Protium heptaphyllum 117
Psychotria myriantha 1101
 Pterocarpanes 991
Pterocephalus sanctus.457
 Purines 711
Pycnanthus angolensis 651
 Pyranoquinolines 767
 Pyrroles 831
- Qinjiao 527
 Qinjiaoamide 527
 Quercetin 229
 Quercetin 3,7-*O*-triosides 941
 Quinine 899
 Quinolone alkaloid 313
 Quinoldione derivatives 313
- Radical scavenger 101,229,287
 Rambutan 751
 Ranunculaceae 27,191
Ranunculus ficaria 27
 Raspberry leaves 705
 Raspberry 613
 5 α -Reductase 345
 Resiniferatoxin 1147
 Resiniferonol orthophenylacetate 1147
 Respiratory alteration 577
 Rhamnaceae 121
 Rhamnosides 121
 Rhodomelaceae 47
- Rice 981
 Ring-contraction 991
Rosa damascene 623
 Rosaceae 623
 Rose flowers 623
Rosmarinus officinalis L.1123
 Rubiaceae 1101
Rubus idaeus 613,705
Ruta 851
 Rutaceae 313,351,367,469
- Saccocalyx satureioides* 645
 Saffron 65
 Salicaceae 441,601
Salvia fruticosa 745
 Santalaceae 969
Santalum insulare 969
 Sapindaceae 121,751
 Saponins 541,1159
 Sarasinose A₄ 265
 Sarasinose A₅ 265
 Scavenging 229,367
 Schizandraceae 109
 Scouring 997
 SDE extraction 387
 Seasonal variation 47
 Sea Cucumber 541
 Secondary metabolites 495
 Seed oil 751
 Self-assembly 1067
 Serpentine soils 1013
 (+)-Sesamin 21
 Sesquiterpene lactones 281,531
 Sesquiterpenes 15,465,531,537,601,773
 Silphium albiflorum 941
 Silylation 307
 SISTEMAT 57
 Site of substitution 957
 Sleeping sickness 619
 Smooth muscle relaxant 613,705
 Solanaceae 859
Solanum 859
Sophora chrysophylla 563
 Sophoraisoflavanone D 563
 Soy 973
 Spathulenol 651
Spatoglossum schroederi 293
 Speciation model 1013
 Spectral characteristics 839
Spodoptera litura 553
 Sponge 265,711,139,773
 Sprouts 1037
Stachys glutinosa L. 1133
Stauranthus perforatus 201
 Stemofoline 813
Stemona alkaloids 813
 Stereochemical correction 237
 Stereoselective synthesis 237,899
 Steroidal alkaloids 859
 Steroidal glycoalkaloids 859
 Sterols 293,299,711,773
 Strictosidinic acid 1101
 Structure elucidation 57
 Structure-activity relationship 1003
 Sesquiterpene polyol ester 537
 Sun protection factor 209
 SunRecome 177
 Superoxide anion 563
 Surface flavonoids 1029
 Synthesis 51,131,237,313,661,831,899,907,991
 Systematic significance 921

Key Word Index
Natural Product Communications Vol. **1** (1-12) 2006

- Tamarixetin diglycoside 953
Taxonomic markers 1023
Tecoma stans 735
Tephrosia purpurea 953
Tephrosia toxicaria 391
Terpenes 1003
Terpenoids 495,697,1181
Terpinen-4-ol 651,1133
 α -Terpineol 303
Tertiary *N*-methyl amines 885
Tetrapedia 757
Teucrium stocksianum.195
Thorectidae 711
Thymol 205
Tinospora cordifolia 553
Titanium(III) chloride 767
TMC-95A/B 907
Topical activity 209
trans-Pinocarveol 303
Triacylglycerols 751
Trichome development 935
Tricin 981
2-Tridecanone 469
Trifolium subterraneum 557
Trihydroxyflavone-benzoate 949
Trihydroxyphenyl-methoxyphenyl-propenone 553
Triisopentenyl guanidine (TIPG) 1097
Trimethoxybenzaldehyde 377
5,6,7-Trimethoxydictamnine 351
1,3,5-Trimethoxybenzene 43
2,4,6-Trimethoxystyrene 43
1-(2,4,5-Trimethoxyphenyl)-ethanone 377
Triterpenes 177,345,381,1073
Triterpene glycosides 27,265,541
Triterpene saponins 27,1159
Triterpenoid glycosides 705
Triterpenoids 299,613
Tropane alkaloid 33
TRV1 1147
Tryptophan 907
Tuberostemonine 813
Toxicity 377
Tunicates 139
Turmeric 1061
Turraeanthus africanus 449
Turraeasterodione 449
Ursane triterpenoids 613,705,1073
Urs-12-ene-3 β ,30-diol 613
Vanilloid antagonists 1147
Verbenaceae 205,727
Vinpocetine 633
Vitamins 1037
Voacanga foetida 825
Volatile oil 651
Wax chemistry 1067
Wheat 997
Wollenweber, E 921
Xanthone 363
Xylocarpus granatum 985
Xymalos monospora 619

Lipoxygenase Inhibitory Activity of Boropinic Acid, Active Principle of <i>Boronia pinnata</i> Massimo Curini, Francesco Epifano, Salvatore Genovese, Luigi Menghini, Donata Ricci, Daniele Fraternale, Laura Giamperi, Anahi Bucchini and Emanuele Bellacchio	1141
A Convenient Synthesis of 5'-Iodoresiniferatoxin (I-RTX) Abdellah Ech-Chahad, Lahboub Bouyazza and Giovanni Appendino	1147
<u>Review /Account</u>	
Acaricides of Natural Origin. Part 2. Review of the Literature (2002-2006) Guido Flamini	1151
Chemistry and Biological Activity of Saponins from <i>Medicago</i> Species Aldo Tava and Pinarosa Avato	1159
Non-nitrogenous Plant-derived Constituents with Antimalarial Activity Anna Rita Bilia	1181

Natural Product Communications

2006

Volume 1, NUMBER 12

Contents

<u>Original paper</u>	<u>Page</u>
Triterpenes from <i>Maytenus macrocarpa</i> and Evaluation of Their Anti-HIV activity Sonia Piacente, Lourdes Campaner Dos Santos, Naheed Mahmood and Cosimo Pizza	1073
New Oxidized 4-Oxo Fatty Acids from <i>Hygrophorus discoxanthus</i> Gianluca Gilardoni, Marco Clericuzio, Alberto Marchetti, Paola Vita Finzi, Giuseppe Zanon and Giovanni Vidari	1079
Kenyaloside, a Novel <i>O,O,O</i>-Triglycosylated Naphthalene Derivative from the Exudate of Kenyan <i>Aloe</i> Species Giovanna Speranza, Daniela Monti, Sergio Crippa, Paola Cairolì, Carlo F. Morelli and Paolo Manitto	1085
New Flavonoid Glycosides from <i>Chrozophora senegalensis</i> and Their Antioxidant Activity Antonio Vassallo, Giuseppina Cioffi, Francesco De Simone, Alessandra Braca, Rokia Sanogo, Angelo Vanella, Alessandra Russo and Nunziatina De Tommasi	1089
<i>N1,N2,N3</i>-Trisopentenyl Guanidine and <i>N1,N2</i>-Diisopentenyl guanidine, Two Cytotoxic Alkaloids from <i>Alchornea cordifolia</i> (Schumach. & Thonn.) Müll. Arg. (Euphorbiaceae) Root Barks Hélène Mavar-Manga, David Chapon, Sara Hoet, Sébastien Block, Marie-Claire. De Pauw-Gillet and Joëlle Quetin-Leclercq	1097
Indole Monoterpenes with Antichemotactic Activity from <i>Psychotria myriantha</i> Cláudia A. Simões-Pires, Fabianne M. Farias, Andrew Marston, Emerson F. Queiroz, Célia G. Chaves, Amélia T. Henriques and Kurt Hostettmann	1101
HPLC Based Activity Profiling for Inhibitors of Human Neutrophil Elastase in <i>Isatis tinctoria</i> Leaf Extracts M. Hamburger, H. G. Rüster and M. F. Melzig	1107
Variation in Artemisinin and Flavonoids Content in Different extracts of <i>Artemisia annua</i> L. Anna Rita Bilia, Caterina Gabriele, Maria Camilla Bergonzi, Pedro Melillo de Malgalhaes and Franco Francesco Vincieri	1111
Antifungal Evaluation of <i>Hypericum triquetrifolium</i> Polar Extracts Against <i>Fusarium</i> spp Daniele Fraternale, Alessandra Bertoli, Laura Giamperi, Anahi Bucchini, Donata Ricci, Francesco Menichini, Elena Trinciarelli and Luisa Pistelli	1117
Antioxidant Activity Analysis for the Selection of <i>Rosmarinus officinalis</i> L. Juan Antonio Garbarino, Nicolás Troncoso, Pia Delpiano, Loreto Carvajal and Alessandra Russo	1123
<i>Hypericum perforatum</i> L., <i>H. maculatum</i> Crantz., <i>H. calycinum</i> L. and <i>H. pulchrum</i> L.: Phytochemical and Morphological Studies Gelsomina Fico, Sara Vitalini, Noemi Colombo and Franca Tomè	1129
Chemical Composition and Antimicrobial Activities of Essential Oil of <i>Stachys glutinosa</i> L. from Sardinia Pintore Giorgio, Chessa Mario, Manconi Paola, Zanetti Stefania, Deriu Antonella and Tirillini Bruno	1133
Molecular Identification of <i>Panax ginseng</i> C.A. Meyer in Ginseng Commercial Products Paola Del Serrone, Lucilla Attorri, Bruno Gallinella, Francesca Romana Gallo, Elena Federici and Giovanna Palazzino	

Continued inside back cover